Comparative genomics of sex determination related genes reveals shared evolutionary patterns between bivalves and mammals, but not *Drosophila*

Filippo Nicolini[1,2]*, Sergey Nuzhdin[3], Fabrizio Ghiselli[1], Andrea Luchetti[1], Liliana Milani[1]

[1]*Department of Biological, Geological and Environmental Sciences, University of Bologna, Bologna (BO), Italy.*

[2]*Fano Marine Center, Fano (PU), Italy.*

[3]*Department of Molecular and Computational Biology, University of Southern California, Los Angeles, CA, USA.*

*Author for correspondence: filippo.nicolini6@unibo.it

Running head: Sex determination related genes in bivalves

Abstract

The molecular basis of sex determination (SD), while being extensively studied in model organisms, remains poorly understood in many animal groups. Bivalves, a diverse class of molluscs with a variety of reproductive modes, represent an ideal yet challenging clade for investigating SD and the evolution of sexual systems. However, the absence of a comprehensive framework has limited progress in this field, particularly regarding the study of sex-determination related genes (SRGs). In this study, we performed a genome-wide sequence evolutionary analysis of the Dmrt, Sox, and Fox gene families in more than 40 bivalve species. For the first time, we provide an extensive and phylogenetic-aware dataset of these SRGs, and we find support to the hypothesis that *Dmrt-1L* and *Sox-H* may act as primary sex-determining genes, by showing their high levels of sequence diversity within the bivalve genomic context. To validate our findings, we

studied the same gene families in two well-characterized systems, mammals and *Drosophila*. In the former, we found that the male sex-determining gene *Sry* exhibits a pattern of amino acid sequence diversity similar to that of *Dmrt-1L* and *Sox-H* in bivalves, consistent with its role of master SD regulator. In contrast, no such pattern was observed among genes of the fruit fly SD cascade, which is controlled by a chromosomic mechanism. Overall, our findings highlight similarities in the sequence evolution of some mammal and bivalve SRGs, possibly driven by a comparable architecture of SD cascades. This work underscores once again the importance of employing a comparative approach when investigating understudied and non-model systems.

Keywords

Dmrt genes; Fox genes; molecular evolution; molluscs; reproduction; Sox genes.

INTRODUCTION

In sexually reproducing organisms, the mechanism of sex determination (SD), i.e., the process by which the male or female identity of an organism (or gonadic tissue) is established, is highly diverse, ranging from strictly genetic systems to environmentally-dependent processes (**Haag & Doty, 2005**; **Uller & Helanterä, 2011**; **Bachtrog et al., 2014**; **Beukeboom & Perrin, 2014**). Characterising the molecular basis of SD is crucial for understanding not only the reproductive biology but also the evolutionary pressures shaping these systems (**Wilkins et al., 1995**; **Ellegren & Parsch, 2007**; **Grath & Parsch, 2016**; **Nicolini et al., 2023a**), as sex-determination related genes (SRGs; including primary sex-determining genes [SDGs]) are those responsible for the phenotypic differences between males and females, thanks to their sex-biased expression and interactions (**Ellegren & Parsch, 2007**; **Beukeboom & Perrin, 2014**; **Grath & Parsch, 2016**). One key aspect of SRGs is that they often exhibit accelerated rates of sequence evolution, due to their involvement in sex-related traits and reproduction. This represents the effect of sexual selection and/or adaptation, which act on sex-biased genes and produce highly divergent proteins at the interspecific level (**Civetta & Singh, 1998**; **Ellegren & Parsch, 2007**; **Meisel, 2011**; **Grath & Parsch, 2016**). Rapid sequence evolution is known for *Sex-determining Region Y* (*Sry*) of therians (**Pamilo & O'Neill, 1997**; **Mawaribuchi et al. 2012**), *Doublesex and mab-3 related gene W* (*Dm-W*) of the African clawed frog *Xenopus laevis*, and *Doublesex and mab-3 related gene Y* (*Dmy*) of the medaka fish *Oryzias latipes* (**Mawaribuchi et al. 2012**), all of which are master SDGs, that is, genes whose expression is primarily responsible for the establishment of the sex of the organism. Evolution under episodic diversifying selection has been detected also in *Drosophila* for genes involved in the SD cascade (e.g., *Sex-lethal* [*Sxl*], *transformer* [*tra*], and *doublesex* [*dsx*]), in correspondence with its establishment in the genus common ancestor (**Mullon et al., 2012**; **Baral et al., 2019**); though, rapid sequence evolution does not seem to concern extant amino acid sequences (**Haerty et al., 2007**; **Baral et al., 2019**), as they are globally evolving

under purifying selection, especially in their catalytic domain (**Mullon et al., 2012**; **Baral et al., 2019**). However, concerning the *dsx* genes, higher rates of nucleotide and amino acid sequence evolution can be observed for male-specific regions, if compared to female-specific regions and dimerization domains (**Baral et al., 2019**).

While SD has been extensively studied in model organisms, like mammals, insects, and roundworms, comparatively little is known about the molecular mechanisms in non-model organisms. A remarkable example is represented by bivalve molluscs, which exhibit a wide variety of reproductive modes and sexual systems (**Breton et al., 2018**). Notwithstanding the considerable importance in the human socio-economic landscape (reviewed in **Haszprunar & Wanninger, 2012**; **Gomes-dos-Santos et al., 2020**), the study of SD mechanisms in bivalves has been hampered by the striking divergence among species (**Li et al., 2022**), and thus largely overlooked and limited to few case studies (**Breton et al., 2018**; **Nicolini et al., 2023a**). So far, no master SDG has been unambiguously identified, and the only working hypothesis on the functioning of the SD gene regulatory network is available for the Pacific oyster *Magallana* (formerly *Crassostrea*) *gigas* (**Zhang et al., 2014**). Nonetheless, the field still lacks both a robust functional investigation and an evolutionary framework in which to place the current knowledge (**Nicolini et al., 2023a**). As a matter of fact, major efforts have been dedicated to identify sex-biased genes through differential gene expression (DGE) analyses (e.g., **Ghiselli et al., 2012**; **Milani et al., 2013**; **Teaniniuraitemoana et al., 2014**; **Zhang et al., 2014**; **Capt et al., 2018**; **Afonso et al., 2019**), but very few have leveraged cutting-edge techniques to investigate their actual role in SD and/or gonad differentiation and development (e.g., **Liang et al., 2019**; **Sun et al., 2022**).

Components of the *doublesex* and *mab-3* related transcription factor (Dmrt), *Sry*-related HMG-box (Sox), and forkhead box (Fox) gene families (henceforth referred to as DSFG) are notoriously known as key actors in several developmental processes across Metazoa (**Benayoun et al., 2011**; **Matson & Zarkower, 2012**; **Sarkar & Hochedlinger, 2013**; **Mawaribuchi et al., 2019**),

including SD in certain clades: the aforementioned *Dm-W*, *Dmy*, and *dsx* all belong to the Dmrt

gene family, while *Sry* belongs to the Sox gene family; *Fox-L2*, which takes part in most of the

vertebrate SD processes as a downstream effector of the female pathway, belongs to the Fox gene

families. Members of the DSFGs have been identified as putative SRGs also in bivalves, thanks to

both DGE analyses and *in-situ* hybridization (ISH; e.g., **Naimi et al., 2009**; **Li et al., 2018**; **Liang**

et al., 2019; **Yue et al., 2021**), suggesting that their role in morphological and sexual development

is maintained also in the clade. However, the clear role of DSFGs has yet to be elucidated,

probably as a consequence to the lack of (i) a systematic classification of the gene families and (ii)

a comprehensive understanding of their evolutionary history.

To overcome such limitations, this study aims to perform a thorough investigation of the DSFG

families in bivalves, with the attempt to provide a high-quality resource to be used as a reference

for future studies. Through the analysis of more than 40 annotated bivalve genomes and

transcriptomes, we aim (i) to describe the complete set and evolutionary history of DSFGs in

bivalves by means of phylogenetic inferences, manual curation, and orthology prediction; (ii) to

identify DSFGs potentially involved in bivalve SD by investigating their sequence evolution in a

genome-wide context. Our hypothesis is that, if any of the DSFGs is directly involved in SD (i.e.,

is a SDG), then we should expect it to show a higher rate of sequence evolution, as already found

in previous studies (**Pamilo & O'Neill, 1997**; **Mawaribuchi et al. 2012**) and discussed earlier;

this characteristic, in turn, would be reflected in a high diversity of the extant amino acid

sequences across the bivalve clade. To assess the robustness and reliability of our approach, we

additionally applied our pipeline to two non-bivalve datasets, composed of mammal and

Drosophila species, respectively (hereon referred to as the 'mammal dataset' and the 'fruit fly

dataset'). By choosing two clades for which SD is well characterised, we wanted to compare our

results with those obtained on taxa for which a deeper and more detailed knowledge is available.

In particular, mammals and *Drosophila* provide two different frameworks to study the patterns of

molecular evolution in SDGs: the former is a system where SD is completely genetic (i.e., the

development into a male or into a female is triggered by the up- or downregulation of *Sry* in undifferentiated gonads, respectively), and for which a high rate of sequence evolution is already known for the male SDG *Sry*; the latter is instead a system where SD is chromosomic, thus lacks a master SDG (the sexual fate of the individual is determined by the ratio between autosomes and X chromosomes), and no high rate of sequence evolution has been detected in extant amino acid sequences of SRGs (which are globally evolving under purifying selection). In this sense, the mammal and the fruit fly datasets represent opposing control datasets to be compared to bivalves, as it is expected that a higher rate of sequence evolution concerns only master SDGs (like *Sry* in therians; i.e., the top regulatory part of the SD cascade), but not the downstream genes (i.e., the bottom effectors). Therefore, we test our pipeline on mammals and fruit flies with the following expectations: (i) in the mammalian dataset, *Sry* should be detected as fast-evolving (**Pamilo & O'Neill, 1997**; **Mawaribuchi et al. 2012**); while (ii) in the fruit fly dataset, no gene among those working within the sex-determining cascade should result as evolving at a higher pace (**Haerty et al., 2007**; **Mullon et al., 2012**; **Baral et al., 2019**). In this way, we are able to assess the reliability of results in bivalves, by comparing them to a null hypothesis.

This work offers novel insights into the evolutionary dynamics of SRGs and contributes a valuable genomic resource for understanding SD in bivalves, one of the most ecologically and economically important groups of marine organisms. In particular, here we provide the first extensive phylogenetic-based classification of DSFGs in bivalves, covering many species from the major bivalve orders, along with a comprehensive investigation of their sequence evolution.

MATERIALS AND METHODS

Dataset of bivalve annotated genomes and transcriptomes

Annotated genome assemblies of bivalves, gastropods, and cephalopods were obtained from various publicly available resources (**Supp. Tab. S1**). Isoforms from genome annotations were

removed using the AGAT toolkit (v0.8.0; **Dainat, unpublished**). The nucleotide coding

sequences for *Sinonovacula constricta* (Adapedonta) was not available for download. To avoid

excluding the species from our analyses, the corresponding file was generated in-house by

mapping the annotated protein sequences on the reference genome using miniprot (v0.13-0; **Li,**

2023). In order to provide an extensive identification of SRGs also for underrepresented bivalve

orders (mainly belonging to the Heterodonta clade), 14 additional species represented by

sequenced transcriptomes were included in the analyses, and were obtained from **Piccinini et al.,**

2021, **Iannello et al., 2023** and **Iannello et al., in preparation**. The resulting set of annotated

genomes and transcriptomes (hereafter referred to as the 'comprehensive bivalve dataset') was

checked for completeness using BUSCO with the Metazoa reference dataset (v5.2.2; **Manni et al.,**

2021).

Identification and classification of Dmrt, Sox and Fox genes in bivalves

Members of DSFG families were retrieved from the comprehensive bivalve dataset with

hmmsearch (v3.3.2; http://hmmer.org/). The signature catalytic domains of the DSFG families

were used as queries. Specifically, HMM profiles were built after the Pfam databases for the DM

domain (PF00751), the HMG box (PF00505), and the forkhead domain (PF00250), respectively.

Obtained hits were then annotated using (i) the PANTHER HMM standalone sequence scoring

against the PANTHER library v18.0 and (ii) RPS-BLAST (v2.5.0+) against CDD (pre-compiled

version, downloaded on 09/11/23). In both cases, hits with an E-value of 10^{-5} were retained.

Genes which were correctly annotated by both systems (**Supp. Tab. S2**) were kept for subsequent

analyses. DSFGs from *Homo sapiens*, *Drosophila melanogaster*, and *Caenorhabditis elegans*

(**Supp. Tab. S3**; the reference species) were retrieved from NCBI and were used as reference

genes for annotation (see below). Classification and nomenclature of each family were retrieved

from: **Mawaribuchi et al., 2019** for Dmrt genes; **Phochanukul & Russell, 2010** and **Sarkar &**

Hochedlinger, 2013 for Sox genes; **Mazet et al., 2003** for Fox genes. The alignments of mollusc

and reference DSFGs were guided by the aforementioned Pfam HMM profiles and performed with Clustal Omega (v1.2.3; **Sievers et al., 2011**), then trimmed with trimAl (v1.4.rev15; **Capella-Gutiérrez et al., 2009**) with a gap threshold of 40%. Resulting alignments were manually inspected to remove sequences with incomplete catalytic domains, then aligned and trimmed again as before. Phylogenetic trees were inferred using IQ-TREE (v2.1.4-beta COVID-edition; **Minh et al., 2020**) with automatic model selection (**Kalyaanamoorthy et al., 2017**) and 1,000 bootstrap replicates. The phylogenetic tree of Dmrt genes was midpoint rooted, as no clear outgroup has been found so far (**Wexler et al., 2014**). Phylogenetic trees of Sox and Fox gene families were rooted using two fungi MATA-1 sequences (XP_62685912.1, CCD57795.1) and two Amoebozoa forkhead-like domains (XP_004368148.1, XP_004333268.1), respectively (**Heenan et al., 2016**; **Nakagawa et al., 2016**). The rooting was performed with Gotree (v0.4.5; **Lemoine & Gascuel, 2021**). To identify and annotate bivalve homology groups within each gene family, we utilized the algorithm implemented in Possvm (v1.2; **Grau-Bové & Sebé-Pedrós, 2021**), with DSFGs from the reference species as reference annotation. To better establish the orthology relationships among ambiguous groups of Dmrt and Fox genes, we ran a series of other phylogenetic reconstructions (see **Discussion**), by using the same pipeline as before. In the case of *Fox-Y* genes, we also used Fox gene sequences from the sea urchin *Strongylocentrotus purpuratus*, as given by **Tu et al.** (**2006**). All the phylogenetic trees were plotted using the R package 'ggtree' (**Yu et al., 2017a**).

Sequence diversity of bivalve single-copy orthogroups

As a metrics to measure the sequence diversity of bivalve DSFGs, and test whether those putatively involved in SD showed higher values than other genes, we used the amino acid sequence divergence (AASD). This metric is fast and straightforward to obtain, as it only requires the amino acid alignment and the corresponding best-fit substitution model. To this purpose, we produced amino acid alignments of bivalve single-copy orthologous (SCOs) groups and built the

distribution of their median AASD. To further save computational time and prevent over-represented bivalve clades (such as Ostreida and Mytilia) to bias the analysis, SCOs were obtained after a reduced dataset comprised of 34 bivalve species (hereafter referred to as the 'reduced bivalve dataset'; **Fig. 1**; **Supp. Tab. S1**), which included, for each bivalve genus, only the best genomes and transcriptomes in terms of either BUSCO scores or assembly statistics (**Supp. Tab. S1**). *Archivesica marissinica* and *Saccostrea glomerata* were removed, as their annotated coding sequences contain many stop codons. Genes were clustered in orthogroups using OrthoFinder (v2.5.5; **Emms & Kelly, 2019**) with DIAMOND ultra-sensitive and default parameters. Resulting orthogroups were splitted into SCOs composed by at least 17 species (50% of the bivalve reduced dataset) using DISCO (v1.3.1; **Willson et al., 2022**). Amino acid and nucleotide sequences were then aligned using Clustal Omega as implemented in TranslatorX (v1.1; **Abascal et al., 2010**), and jointly trimmed using trimAl with a gap threshold of 40% and the removal of spurious sequences (-resoerlap 50 -seqoverlap 50). Eventually, SCOs containing (i) internal stop codons, (ii) with less than 17 species left, or (iii) containing DSFGs were removed from downstream analyses. The best amino acid substitution model was inferred for each trimmed alignment using ModelFinder as implemented in IQTREE2 (model search was restricted to the following matrices: Blosum62, cpREV, Dayhoff, DCMut, FLU, HIVb, HIVw, JTT, JTTDCMut, LG, mtART, mtMAM, mtREV, mtZOA, rtREV, VT, WAG) and the corresponding pairwise amino acid distances were computed with the function 'dist.ml' from the 'phangorn' R package (**Schliep 2011**). We decided to use the pairwise amino acid distance instead of the tip-to-tip phylogenetic distance (which accounts for a more comprehensive evolutionary signal) to save computational time. However, to check whether the two metrics were comparable to each other, we randomly selected 200 SCOs and computed the maximum likelihood (ML) trees using IQTREE2, with ModelSelection restricted as before. Then, the tip-to-tip pairwise distances were obtained with the R package 'adephylo' (**Jombart & Dray, 2010**). The same pipeline to obtain SCOs and AASD was also utilized on the DSFG families. The distribution of AASD was then built after the median values of pairwise distances of

each SCO, and genes were categorised accordingly into three groups: Group 1, consisting of genes

from the 1% upper quantile of the distribution; Group 2, consisting of genes between the 1% and

5% upper quantiles; and Group 3, consisting of all the remaining genes. Group 1 and Group 2

genes will be referred to as 'highly divergent genes'.

Mammals and *Drosophila* spp. as test datasets

To validate our approach for the study of bivalve SRG molecular evolution, we run the same

analysis on two additional datasets, consisting of reference genomes of mammals and *Drosophila*

species (**Supp. Tab. S4–S5**, respectively), whose sex-determining mechanisms are well studied

and characterised. As a matter of fact, despite it is well known that SDGs tend to evolve faster

than genes not involved in SD, the hypothesis has never been tested extensively across the entire

phylogenetic diversity of a group: molecular evolution of SDGs and SRGs has mainly been tested

on single species or inside the boundaries of taxonomic genera (**Stothard et al., 2003**; **Haerty et**

al., 2007; **Mank et al., 2007**; **Mullon et al., 2012**; **Papa et al., 2017**; **Ghiselli e al., 2018**). For

both mammals and fruit flies, annotated genomes were downloaded from NCBI using the

command-line tool 'datasets', then processed using the same pipeline and scripts as before (**Fig.**

1).

GO term enrichment

After having obtained the distributions of AASD in the three datasets (Bivalvia, Mammalia, and

Drosophila) we performed a gene ontology (GO) enrichment analysis of highly divergent genes.

To do so, we firstly selected one gene per SCO, giving priority to few chosen species: (i) for

bivalves, we selected genes from *Pecten maximus*, or alternatively from *Crassostrea gigas*,

Hyriopsis bialata, *Tridacna squamosa*, and *Solen grandis*; (ii) for mammals, we selected genes

from *H. sapiens*, or alternatively from *Bubalus bubalis*, *Panthera tigris*, *Camelus dromedarius*,

and *Monodelphis domestica*; (iii) for fruit flies, we selected genes from *D. melanogaster*, or

alternatively from *Drosophila hydei*, *Drosophila pseudoobscura*, and *Drosophila suzukii*.

Afterwards, we annotated the obtained datasets with the corresponding GO terms using the OMA

browser (accessed 18/09/2024; **Altenhoff et al., 2024**). The GO-term enrichment was then

performed with the R package 'topGO' using the Fisher's exact test (**Alexa et al., 2024**).

RESULTS

Assembly of the bivalve, mammal, and fruit fly datasets

The comprehensive bivalve dataset on which we retrieved and analysed DSFGs consists of 29

bivalve genomes, 14 bivalve transcriptomes, and 7 outgroup genomes (5 gastropods and 2

Octopus spp.; **Supp. Tab. S1**). BUSCO statistics for complete single-copy genes spanned from

the 64.9% in *Modiolus modiolus* to the 99.4% of *Perna viridis*, with a median value of 94.7%. We

were able to get at least one representative species for 11 different bivalve orders, covering a good

proportion of the phylogenetic diversity of the clades Pteriomorphia, Palaeoheterodonta, and

Imparidentia, and thus building the most extensive genomic and transcriptomic dataset for bivalve

comparative analyses so far. Unfortunately, no genomes or transcriptomes for Protobranchia,

Archiheterodonta, and Anomalodesmata were available at the time of the project, thus we were

not able to include any of those clades in our analysis. The mammal dataset consists of 32 species

and 1 outgroup (*Gallus gallus*, Aves; **Supp. Tab. S4**), and covers 12 major orders, while the fruit

fly dataset consists of 17 species and 1 outgroup (*Anopheles gambiae*, Culicidae; **Supp. Tab. S5**),

and covers 2 *Drosophila* subgenera (i.e., *Drosophila* and *Sophophora*). BUSCO statistics for

complete single-copy genes were generally higher than those of bivalves, with a median of 98.3%

for mammals and of 99.8% for fruit flies (**Supp. Tab. S4–S5**).

Dmrt, Sox, and Fox complements in bivalves

262 Our pipeline managed to successfully identify and annotate DSFGs in bivalves, as proved by the

263 same analysis in mammals and fruit flies (see following sections). We retrieved four main

orthology groups of Dmrt genes in bivalves (**Fig. 2**; **Supp. Fig. S1**; **Supp. Tab. S6**), three

corresponding to the groups present in the Bilateria common ancestor (*Dmrt-2*, *Dmrt-3*, and *Dmrt-*

4/5; **Mawaribuchi et al., 2019**), and one additional group with no clear ortholog among reference

genes, and thus putatively specific to molluscs (named *Dmrt-1L*, as per **Li et al., 2018b**; **Evensen**

et al., 2022). The *Dmrt-4/5* subgroup shows a group-specific expansion in Palaeoheterodonta and

Heterodonta, while *Dmrt-1L* is completely absent from Heterodonta. The degree of missing data

for Dmrt genes in bivalves is ~35% (**Supp. Tab. S7**). The ubiquitin-binding CUE-like DMA

domain has been annotated in most of the *Dmrt-3* and *Dmrt-4/5* genes, while an additional DM

domain has been annotated in *Dmrt-1L* genes in Mytilida and the gastropod *Pomacea canaliculata*

(**Supp. Tab. S6**). Additionally, we retrieved six main orthology groups of Sox genes, none of

which is restricted to molluscs or bivalves (**Fig. 2**; **Supp. Fig. S2**; **Supp. Tab. S6**). Five Sox

groups (*Sox-B1/2*, *Sox-C*, *Sox-D*, *Sox-E*, and *Sox-F*) are those traditionally considered to be

present in the Bilateria common ancestor (**Phochanukul & Russell, 2010**), while one has been

identified outside mammals only recently (*Sox-H*, or *Sox-30*; **Han et al., 2010**). *Sox-B2* and *Sox-*

B1 have been grouped in the same clade, as in our phylogenetic reconstruction *Sox-B2* is

paraphyletic with respect to *Sox-B1* (**Supp. Fig. S2**), despite being traditionally recognised as a

separate paralogy group in humans, fruit flies, and roundworms. The degree of missing data for

Sox genes in bivalves is ~8% (**Supp. Tab. S7**). The Sox N-terminal signature domain was

annotated for *Sox-E* genes (**Supp. Tab. S6**). Concerning Fox genes, we retrieved 27 main

orthology groups (**Fig. 2**; **Supp. Fig. S3**; **Supp. Tab. S6**), two of which are specific to molluscs

(*Fox-OG13/NA*, *Fox-OG16/NA*). Additionally, other potential mollusc-specific Fox groups have

been identified, but these have been excluded from the final orthology analysis as they are present

in less than half of bivalve species (**Supp. Tab. S6**). The two major Fox gene subgroups, Group I

(monophyletic, specific to Metazoa; includes *Fox-A*, *Fox-B*, *Fox-C*, *Fox-D*, *Fox-E*, *Fox-F*, *Fox-G*,

Fox-H, *Fox-L1*, *Fox-L2*, *Fox-Q2*) and Group II (paraphyletic, specific to Opisthokonta; includes

Fox-O, *Fox-P*, *Fox-J2*, *Fox-J1*, *Fox-K*, *Fox-N2/3*, *Fox-N1/4*; **Larroux et al., 2008**), have been

recovered, including the four Fox genes that were present in the Bilateria common ancestor (*Fox-*

C, *Fox-F*, *Fox-L1*, and *Fox-Q1*; **Shimeld et al., 2010**). Two putative lineage-specific expansions

have been recovered for *Fox-OG28/NA*, one regarding *Mytilus* spp. and one regarding the two

Myida species (**Fig. 2**; **Supp. Fig. S3**). The degree of missing data for Fox genes in bivalves is

~22% (**Supp. Tab. S7**). The forkhead-associated (FHA) domain was annotated for *Fox-K* genes,

the *Fox-P* coiled-coil signature domain was annotated for *Fox-P* genes, while both the forkhead

N- and C-terminal signature domains were annotated for *Fox-A* genes (**Supp. Tab. S6**).

Amino acid sequence divergence of Dmrt, Sox, and Fox genes in bivalves

We produced amino acid alignments of ~11k SCOs, 32 of which belongs to the DSFG families.

Of these, 112 were assigned to Group 1 (1% upper quantile), 447 to Group 2 (5% upper quantile),

and 10,603 to Group 3. Most of the DSFGs (29/32) fell in Group 3 (**Fig. 3**), which means they

have a median AASD comparable to the vast majority of other genes in bivalves (median level of

the genomes). Just *Dmrt-1L*, *Sox-H*, and *Sox-F* showed higher divergences, and have been

accordingly placed in Group 2. Overall, pairwise AASD proved to be a good approximation of the

tip-to-tip distances calculated on the corresponding ML phylogenetic trees ($R = 0.84$, $p < 2.2E–16$,

considering 200 randomly-selected trees; **Fig. 3C**), and it also shows no influence from the

alignment length ($R = 0.11$) or the number of represented species ($R = –0.23$; **Fig. 3D–E**). Genes

from Group 1 and Group 2 are strongly involved in cellular regulatory processes (such as those

related to the metabolism of nucleic acids, proteins, and other macromolecules), but also in

development and response to external stimuli, as shown by the GO-term enrichment analysis

(**Tab. 1**; **Supp. Tab. S8**).

Dmrt, Sox, and Fox genes, and amino acid sequence divergence in the test datasets

Most of the already-recognised DSFG orthology groups in mammals and fruit flies have been

identified. In mammals, we retrieved 7 Dmrt orthology groups with ~3.1% of missing data, 20 Sox

orthology groups with ~8.1% of missing data, and 42 Fox orthology groups with ~4.6% of missing data (**Supp. Fig. S4A, S5–S7**; **Supp. Tab. S9**). Of these, just *Sox-5* was not included in the subsequent AASD analysis, as it did not meet the 50%-species occupancy threshold. OrthoFinder analysed ~650M genes, and the number of SCOs used in the AASD analysis is >16k (**Fig. 4A**). From the distribution of median AASD, 163 genes were assigned to Group 1, 649 to Group 2, and 15,355 to Group 3. Most of the DSFGs (66/68) fell in Group 3 (**Fig. 4B**), while *Sry* and *Fox-D4* showed higher divergences, and have been accordingly placed in Group 1 and 2, respectively. Genes from Group 1 and Group 2 show a strong enrichment in immune-related functions (such as innate and adaptive immune response, defence response to bacteria and viruses, lymphocyte metabolism, etc.), but also in reproductive processes (such as spermatogenesis; **Tab. 1**; **Supp. Tab. S8**). Concerning *Drosophila*, we retrieved 4 Dmrt orthology groups with ~1.7% of missing data, 7 *Sox* orthology groups with ~3.9% of missing data, and 17 Fox genes with ~8.3% of missing data (**Supp. Fig. S4B, S9–S10**; **Supp. Tab. S10**). OrthoFinder analysed ~240M, and the distribution of median AASD was built after >12k SCOs (**Fig. 4C**). 126 genes were assigned to Group 1, 501 to Group 2, and 11,880 to Group 3. All the DSFGs have been used in the AASD analysis, but none of them have been placed in Group 1 or 2, that is, all the DSFGs in *Drosophila* have an AASD comparable to the median level of the genome (**Fig. 4D**). Genes of Group 1 and Group 2 show a GO-term enrichment in meiotic processes, such as chromosome/chromatid organisation, and retrotransposon silencing (**Tab. 1**; **Supp. Tab. S8**).

DISCUSSION

A new manually-curated and phylogenetic-based reference dataset of Dmrt, Sox, and Fox genes in bivalves

The annotation and characterisation process of a gene family may harbour many overlooked challenges in a certain clade of organisms (**Vizueta et al., 2020**). For example, the presence of

highly conserved catalytic domains may hamper the correct identification of the components of a

gene family because of insufficient phylogenetic signal, as it is the case for Hox and ParaHox

genes and their homeobox motif (**Baldwin-Brown et al., 2018**; **Nicolini et al., 2023b**).

Conversely, the components of dynamic gene families characterised by abrupt and sequential

duplication events may be difficult to sort into separate groups. As a matter of fact, varying levels

of sequence heterogeneity and gene copy numbers makes the inference of orthologous groups

hard, as for certain clans of the P450 family (**Dermauw et al., 2020**). Regardless of the causes,

having a solid and wide phylogenetic context, in which to study gene duplications/losses and

orthology relationships, is crucial to overcome these difficulties. In the same way, manual curation

and visual inspection of multiple sequence alignments, phylogenetic trees, and gene structures (in

terms of domain composition, start and stop codons, and other feature representations) is helpful,

despite being time-demanding and possibly less reproducible. In this study, we characterised the

full complement of DSFGs in the vast class of bivalves, by leveraging sequence domain

annotation, phylogenetics, and manual curation of the dataset. Our aim was to obtain the most

reliable gene complements as possible, combined with a vast taxonomic dataset, a solid

phylogenetic inference, an openly available dataset of gene sequences, and a reproducible pipeline

for the annotation of gene identity. By doing so, we want to provide a reliable resource for future

studies of DSFGs, either focused on bivalves or on Metazoa in general.

Concerning the Dmrt gene family, we identified orthologs of the vertebrate *Dmrt-2*, *Dmrt-3*, and

Dmrt-4/5 (*A1/A2;* **Fig. 2**; **Supp. Fig. S1**; **Supp. Tab. S6**), which are also expected to have been

present in the Bilateria common ancestor (**Mawaribuchi et al., 2019**). **Wang et al. (2023)** found

that *Dmrt-4/5* is duplicated in *Mercenaria mercenaria* and *Cyclina sinensis* (Venerida), and in

Dreissena polymorpha (Myida), and we confirm this result by tracing back the duplication event

to the split between Palaeoheterdonta (here represented by Unionida) and Heterodonta (here

represented by Venerida, Myida, Sphaeriida, Adapedonta, Cardiida, and Lucinida; **Fig. 2**).

Furthermore, we confirm *Dmrt-1L* to be present in many bivalve species (mainly belonging to the

Ostreida, Pectinida, Mytilida, and Unionida orders; **Fig. 2**), as well as in gastropods and *Octopus*.

Though, our phylogenetic analysis did not retrieve any unambiguous orthology relationship

among *Dmrt-1L* and either vertebrate *Dmrt-1* or *Drosophila dsx* genes, as instead it was proposed

in previous works (**Li et al., 2018b**; **Evensen et al., 2022**). As a matter of fact, the amino acid

sequence of the *Dmrt-1L* DM domain does not recall that of any other Dmrt gene. Furthermore, it

must be considered that various phylogenetic analyses have recovered both *Dmrt-1* and *dsx* genes

to be restricted to vertebrates and arthropods, respectively (**Wexler et al., 2014**; **Mawaribuchi et**

al., 2019; **Panara et al., 2019**), that is, they do not have any direct ortholog outside their relative

clades. Thus, if *Dmrt-1L*, *dsx*, and *Dmrt-1* were true orthologs, their origin would need to be

placed at least in the Bilateria common ancestor, which seems however to be not the case. All

considered, we thus confirm that *Dmrt-1L* is not orthologous to *Dmrt-1* and *dsx* and is rather a

mollusc-specific gene (**Evensen et al., 2022**). The monophyly of the *Dmrt-1L* group is not

supported by the phylogenetic tree inferred with Dmrt genes from molluscs and the reference

species (**Supp. Fig. S1**); though, it is recovered when analysing just genes from mollusc species

(**Supp Fig. S11**). To this regard, we speculate that in our analysis, the difficulty in obtaining the

monophyly of *Dmrt-1L* genes may have arisen primarily because of the many *C. elegans*-

restricted genes (**Supp. Tab. S3**), which are placed among bivalve genes (**Supp. Fig. S1**), but also

because of the high AASD of *Dmrt-1L* genes (see **High amino acid sequence divergence**

identifies putative sex-determining genes), which hampers a straight-forward phylogenetic

reconstruction. Furthermore, our broad-context analysis allowed us to identify some cases of

incorrect gene identification in bivalves, which have arisen because of erroneous or ambiguous

annotations in previous works, as a result of limited datasets or analyses. For example, (i) the

scallop-specific cluster of Dmrt genes retrieved by **Wang et al. (2023)** rather belongs to the *Dmrt-*

1L group, and (ii) the classification of Dmrt genes in *Crassostrea* species provided by **Zeng et al.**

(2024) needs to be revised following as follows: *Dmrt-1* genes are *Dmrt-4/5*; *Dmrt-2* genes are

Dmrt-3; *Dmrt-3* genes are *Dmrt-1L*; hence, *Crassostrea* species do not have *Dmrt-2* genes.

For what concerns the Sox gene family, bivalves (or molluscs) do not show any major clade-

restricted gene, as only the five Bilateria-specific Sox groups (*Sox-B1/2*, *Sox-C*, *Sox-D*, *Sox-E*, and

Sox-F) and *Sox-H* have been identified (**Fig. 2**; **Supp. Fig. S2**; **Supp. Tab. S6**), in accordance

with previous findings (**Yu et al., 2017b**; **Evensen et al., 2022**; **Wang & Nie , 2024**). *Sox-B1/2* is

clearly made up of two subgroups (i.e., *Sox-B1* and *Sox-B2*), as expected, but their respective

identity could not be unambiguously established, as *Sox-B1/2* genes of reference species do not

form separate clusters (**Supp. Fig. S2**), even when inferring the phylogenetic tree only of

components of the *Sox-B1/2* group from molluscs and reference species (**Supp. Fig. S12**).

Compared to Dmrt and Sox genes, the Fox gene family appears as the most dynamic in terms of

gene presence/absence, as already shown by other works (**Wu et al., 2020**; **Schomburg et al.,**

2022; **Seudre et al., 2022**). Our phylogenetic analysis successfully recovered Group I and Group

II of Fox genes (**Larroux et al., 2008**), which include the four Fox genes that were present in the

Bilateria common ancestor (*Fox-C*, *Fox-F*, *Fox-L1*, and *Fox-Q1*; **Fig. 2**; **Supp. Fig. S3**; **Supp.**

Tab. S6; **Shimeld et al., 2010**). To our knowledge, this is the first broad-taxonomic identification

and classification of Fox genes in bivalves, as up to now they have been systematically

characterised only in *C. gigas* (**Yang et al., 2014**), *Patinopecten yessoensis* (**Wu et al., 2020**), and

Ruditapes philippinarum (**Liu et al., 2024**). Firstly, our analysis confirms the absence in molluscs

of *Fox-I*, *Fox-Q1*, *Fox-R*, *Fox-S* (**Supp. Fig. S3**), which are in fact thought to have emerged with

the diversification of deuterostomes or vertebrates (**Yang et al., 2014**; **Wu et al., 2020**;

Schomburg et al., 2022; **Seudre et al., 2022**). Furthermore, we have found many Fox groups that

appeared as mollusc-specific and/or still-unnamed at a first analysis. However, a more in-depth

investigation revealed a different scenario. *Fox-OG2/NA* appears close to the human *Fox-M* gene

in the phylogenetic tree, but they do not form a monophyletic group (**Supp. Fig. S3**). However, by

comparing *Fox-OG2/NA* sequences and phylogenetic tree with those analysed by **Yang et al.**

(2014), **Wu et al., (2020)**, **Schomburg et al. (2022)**, **Seudre et al. (2022)**, it appears clear that

this group of Fox genes is indeed *Fox-M*. However, our analysis has failed to retrieve a

monophyletic relationship among bivalve and human *Fox-M* genes, even when inferring a tree

with just *Fox-J2*, *Fox-M*, *Fox-O*, and *Fox-P* complements (**Supp. Fig. S13**), which belong to the

same Fox group. Regarding the *Fox-OG39/NA* group, it does not have any homolog in reference

species (**Supp. Fig. S3**) but is found to belong to the *Fox-AB* group by sequence comparison with

previous works (**Yang et al., 2014**; **Wu et al., 2020**; **Seudre et al., 2022**). *Fox-AB* was formerly

described only in the sea urchin *Strongylocentrotus purpuratus* and the lancelet *Branchiostoma*

floridae (**Tu et al., 2006**; **Yu et al., 2008**), but was later identified also in several Spiralia lineages,

including molluscs (e.g., **Yang et al., 2014**; **Wu et al., 2020**; **Seudre et al., 2022**). A similar

situation concerns *Fox-OG15/NA* and *Fox-OG28/NA*, which again could not be named based on

orthology relationships with the reference species genes (**Supp. Fig. S3**), but actually represent

two lineage-specific expansions of the *Fox-Q2* group (named *Fox-Q2b* and *Fox-Q2c*), as already

appointed in previous studies (**Yang et al., 2014**; **Wu et al., 2020**). This observation fits within

the wider context of the *Fox-Q2* group expansion in Bilateria and, particularly, in Spiralia, that led

to remarkable differences in their gene copy numbers across various clades (**Seudre et al., 2022**).

Two additional Fox genes have been previously identified in bivalves and were named *Fox-Y* and

Fox-Z (**Yang et al., 2014**; **Wu et al., 2020**). In our analysis, these Fox genes were identified as

Fox-OG13/NA and *Fox-OG16/NA*, after sequence comparison with Fox genes from *C. gigas* and

P. yessoensis. On one hand, *Fox-Y* was firstly identified in *S. purpuratus* (**Tu et al., 2006**) and

only recently in a few bivalve species (**Yang et al., 2014**; **Wu et al., 2020**). However, when

analysing bivalve and *S. purpuratus* Fox genes, we failed in retrieving such a clear orthology

relationship, as *S. purpuratus* *Fox-Y* does not fall within the phylogenetic range of bivalve *Fox-*

OG13/NA, which contains the supposed *Fox-Y* orthologs (**Supp. Fig. S14**). Also, the forkhead

domains of *Fox-OG13/NA* genes were annotated as 'forkhead domain P' (**Supp. Tab. S6**). On the

other hand, *Fox-Z* was firstly identified in bivalves and in several other protostomes, thanks to a

phylogenetic work including the brachiopod *Lingula unguis*, the annelid *Capitella teleta*, the

scorpion *Centruroides sculpturatus*, and the centipede *Strigamia maritima* (**Wu et al., 2020**).

However, later works have not recovered this Fox gene, even when analysing annelids (**Seudre et**

al., 2022) and panarthropods (**Schomburg et al., 2022**) in a more focused effort. In this case, the

forkhead domains were annotated as either a generic 'forkhead domain' or a 'forkhead domain

Q2' (**Supp. Tab. S6**). All considered, we argue that bivalves possess two additional Fox groups

(here *Fox-OG13/NA* and *Fox-OG16/NA*; **Fig. 2**; **Supp. Fig. S3**; **Supp. Tab. S6**) which are shared

with other mollusc species, as revealed also by other authors. However, given the discordant

results of the phylogenetic hypothesis and domain annotation, we think that a more thorough

investigation on their orthology relationships with Fox genes from other Metazoa is needed, and

thus we chose to not employ their former names *Fox-Y* and *Fox-Z.*

Besides the DSFG groups discussed so far, it must be also considered that many orphan genes

have been identified (**Supp. Fig. S1–S3**; **Supp. Tab. S6**). For example, **Wu et al. (2020)**

identified a duplication event of *Fox-H* genes in *C. gigas*, which has been recovered also in our

analysis for the entire Ostreida clade (*Fox-OG36/NA*; **Supp. Fig. S3**). Similarly, a gene orthology

group putatively specific to Pteriomorphia has been identified among Sox genes (*Sox-OG1/NA*).

Of course, these genes deserve as much attention as their widely distributed paralogs, as they may

constitute true group-specific expansions and may play fundamental roles in some biological

processes. However, they have not been discussed here or included in **Fig. 2** for clarity purposes,

but their sequences are available in on GitHub (see the **Data Availability Statement**).

Overall, our analysis clearly shows the importance of adopting a wide-angle approach when

characterising the members of a gene family, especially for large ones such as the Fox genes

(**Schomburg et al., 2022**). As a matter of fact, the presence of duplication events and orphan

genes needs to be addressed with a broad taxonomic dataset, in order to account for possible mis-

annotations, gene phylogenetic misplacements, and sequence heterogeneity. Additionally, many

reference species need to be included for the gene identification process, to consider distantly

related genes and obtain a solid annotation. Our gene annotation pipeline also resulted to be very

solid, even with non-model organisms and sub-optimal genomic and transcriptomic resources as

those of bivalves. As a matter of fact, by running the same pipeline on two additional datasets

composed of mammal and fruit fly genomes, we were able to obtain high-quality orthology groups

in accordance with previous knowledge on the clades (**Supp. Fig. S4-S10**; **Supp. Tab. S9–S10**),

with little or no need of manual curation. Furthermore, the present work represents the first broad

analysis of DSFGs in both mammals and fruit flies, as so far attention has been mainly dedicated

to single well-studied organisms or little clades (e.g., **Jackson et al., 2010**).

High amino acid sequence divergence identifies putative sex-determining genes

Sex-biased genes tend to evolve more rapidly than unbiased genes at the level of their protein

sequences. Accelerated rates have been observed in both male-biased genes (reviewed in **Parsch**

and Ellegren 2013; **Grath and Parsch 2016**) and female-biased genes (e.g., **Papa et al. 2017**;

Ghiselli et al. 2018), but also in SRGs and SDGs (**O'Neil & Belote, 1992**; **Whitfield et al., 1993**;

de Bono & Hodgkin, 1996). For example, it has been shown that *Dm-W*, *Dmy*, and *Sry* (which are

SDGs in the African clawed frog *X. laevis*, in the medaka fish *O. latipes*, and in eutherians,

respectively) all have higher substitution rates than their paralogues (*Dmrt-1* for *Dm-W* and *Dmy*,

Sox-3 for *Sry*), particularly when considering their DNA-binding domains (**Mawaribuchi et al.**

2012). Similarly, both a burst of positive selection and a relaxation of purifying selection has been

detected in *Drosophila Sxl* in correspondence with its recruitment at the top of the sex-determing

cascade. The same signs of relaxed purifying selection have been found in the downstream targets

of *Sxl*, that is, *transformer* (*tra*) and *dsx*, despite no evidence of positive selection has been

detected (**Mullon et al., 2012**).

Considering these shared features of SRGs and SDGs, we decided to look for signs of accelerated

sequence evolution in DSFGs of bivalves to evaluate if any of them could be *a priori* associated

with SD by employing the tools of molecular evolution. However, we analysed patterns of

sequence evolution not only among putative SRGs and their close paralogs, but also considering

the genomic context in which these genes evolve. Our aim was to check whether SRGs show

higher rates of sequence evolution when compared to other genes not involved in SD and not

belonging to the same gene family. To do so, we obtained the AASD median values of more than

11k SCOs from bivalve genomes (**Fig. 3A**), in order to build a statistical distribution to be used as

a reference: if SRGs/SDGs (in this case, DSFGs) truly evolve faster than other genes, we may

expect them to fall within the 5% (or even 1%) upper quantile of the distribution (**Fig. 3B**), i.e.,

within highly divergent genes (Group 1 and Group 2 genes of the distribution). We chose to use

the AASD as a metric of sequence evolution (instead of the tip-to-tip distances of phylogenetic

trees, which account for more comprehensive evolutionary models) to save computational time.

As a matter of fact, the AASD median values proved to be a good approximation of the tip-to-tip

median distances in 200 randomly selected genes (**Fig. 3C**; R = 0.84, p < 2.2E−6).

Among DSFGs, three fell within the 5% upper quantile, namely *Dmrt-1L*, *Sox-H*, and *Sox-F*.

Interestingly, on the basis of DGE analyses, *Dmrt-1L* and *Sox-H* have been already proposed to be

involved in the male SD pathway of *C. gigas* (inset in **Fig. 3B**; **Zhang et al., 2014**), while *Sox-H*

to be also a candidate gene involved in the development of male germ cells in the Manila clam

Ruditapes philippinarum (**Ghiselli et al., 2012**). Specifically, *Sox-H* would play a major role in *C.*

gigas SD, by interacting with *Dmrt-1L* and determining the onset of the male phenotype

development; at the same time, both *Sox-H* and *Dmrt-1L* would inhibit *Fox-L2*, which instead is

necessary to start the female phenotype development. *Dmrt-1L* and *Sox-H* have been appointed

several other times to be involved in male-gonad development and differentiation, through DGE

(e.g., **Teaniniuraitemoana et al., 2014**; **Capt et al, 2018**; **Afonso et al., 2019**), ISH (e.g., **Naimi**

et al., 2009; **Li et al., 2018**; **Liang et al., 2019**; **Yue et al., 2021**), and RNA interference (**Liang et**

al., 2019; **Sun et al., 2022**). Therefore, the high AASD of *Dmrt-1L* and *Sox-H* is coherent with

previous works, strengthening their role as putative SRGs.

The relationship between high gene AASD and the involvement in SD is particularly strengthened

when looking at the patterns of AASD in the test datasets, which corroborates the solidity of our

analysis: (i) in the mammal dataset—which represents a strictly genetic SD system with a master

and rapidly-evolving SDG—one of the genes from the 5% upper quantile of the distribution is *Sry*

(**Fig. 4A–B**), the male sex-determining gene in eutherians (inset in **Fig. 4B**); (ii) in the fruit fly

dataset—which represents a chromosomic SD system and with no differences in the rates of

sequence evolution among SRGs—none of the DSFG exhibit significantly higher AASD (**Fig.**

4C–D), including the downstream effector *dsx* (inset in **Fig. 4D**). Also, *Sxl* and *tra*, both involved

in the SD pathway of *Drosophila* (inset in **Fig. 4D**) do not belong to the group of highly divergent

genes, as they have a mean amino acid divergence of about 0.09 and 0.9, respectively (Group 3;

Fig. 4D). Therefore, it can be argued that both *Dmrt-1L* and *Sox-H* may not only be SRGs, but

may participate in bivalve SD as primary SDGs, which is reflected in their high AASD, as it is

observed for *Sry* in mammals. As a matter of fact, if they were involved in SD just as intermediate

actors of the signalling cascade, we should have not observed a high AASD, as *Drosophila Sxl*,

tra, and *dsx* seem to suggest. Overall, these patterns of molecular evolution concerning SRGs and

SDGs are also supported by the way SD regulatory networks evolve. It has been proposed that the

sex-determining cascades tend to arise and be established with a bottom-up mechanism (**Wilkins**

et al., 1995; **Mullon et al., 2012**; **Beukeboom & Perrim, 2014**; **Capel, 2017**). This means that

the regulatory relationships among genes at the bottom of the cascade are settled prior to the

regulatory relationships among genes at the top and, consequently, upstream regulators are

progressively recruited to fine-tune diverse SD signals. These evolutionary patterns eventually

produce gene regulatory networks in which the divergence of the upstream triggers is higher than

that of downstream effectors, in terms of both identity and sequence composition (**Beukeboom &**

Perrim, 2014). This mechanism has been proposed for *Drosophila* species (**Mullon et al., 2012**),

C. elegans (**Stothard & Pilgrim, 2003**), and vertebrates, despite in the latter case it has been

questioned several times (reviewed in **Capel, 2017**).

Two main objections can be moved against our approach: (1) the distribution of AASD is not

appropriate for this kind of inference, as it does not represent the true gene evolutionary (or

substitution) rates (which instead are those usually utilized when dealing with SRGs and SDGs);

(2) the three datasets are not comparable one to each other, as they take into consideration very

different animal groups, with different taxonomic rankings and different divergence times (thus,

the patterns of AASD may be the products of other confounding factors not directly related to

SD). Concerning the first objection, we are aware that the AASD does not represent the

evolutionary rate itself, but rather its product. However, the two features are tightly linked, as on

the long term highly divergent proteins tend to be produced by genes with high evolutionary (or

substitution) rates (**Echave et al., 2016**). By performing a GO term enrichment, it emerged that

highly divergent genes of the mammal dataset are mainly involved in the immune response and

male spermatogenesis (**Tab. 1**; **Supp. Tab. S8**), which are two processes notoriously connected

with rapid sequence evolution (i.e., higher evolutionary rates; **Swanson & Vacquier, 2002**;

Murat et al., 2023; **Vinkler et al., 2023**). Similarly, highly divergent genes from the fruit fly

dataset show an enrichment for GO terms associated with meiotic-related functions (such as the

formation of the synaptonemal complex by the products of *c(2)M*, *c(3)G*, *corona*, and *corolla*

genes; **Tab. 1**; **Supp. Tab. S8**), which again are known to evolve rapidly (**Hemmer et al., 2016**).

In other words, the test datasets—which include well-studied and characterised model systems—

allow us to directly link the high AASD (as computed in this work) with high rates of sequence

evolution (as found in previous works). This consideration can thus be extended also to the

bivalve dataset: highly divergent genes in terms of AASD, which include some DSFGs and show

an enrichment for GO-terms associated to macromolecule metabolism and morphological

development (**Tab. 1**; **Supp. Tab. S8**), are also genes with accelerated substitution rates (**Ghiselli**

et al., 2018; **Iannello et al., 2023**).

Concerning the second objection, we chose two test datasets with different characteristics as we

wanted to check the extent of our hypothesis, that is, molecular evolution can be used to look for

putative primary SDGs in taxonomic-wide analyses. The difference in divergence times and

taxonomy ranks for bivalves and therians (Late Cambrian [about 498 million years ago, Mya;

Song et al., 2023] and Early Mesozoic [166–123 Mya; **Álvarez-Carretero et al., 2022**],

respectively) do not seem to influence the sequence diversity of SRGs, as both *Dmrt-1L/Sox-H* for

bivalves and *Sry* for mammals exhibit high AASD with respect to their own distributions,

regardless of their age. *Dmrt-1L* and *Sox-H* (which are mollusc- and Bilateria-specific,

respectively) are undoubtedly older than *Sry,* which emerged in the Theria common ancestor

(**Foster et al., 1992**), but each one of them can be considered a highly divergent gene in bivalves

and mammals, respectively (i.e., genes that are included in the 5% upper quantile of bivalve and

mammal AASD distributions). Conversely, the difference in divergence times and taxonomic

ranks for *Drosophila* (Paleocene/Eocene boundary [about 56 Mya; **Russo et al., 2013**]) may seem

to be influencing the results for the dataset, resulting in a false negative. In other words, it can be

argued that: (i) the genes included in the SD cascade of *Drosophila* (such as *Sxl, tra*, and *dsx*; inset

in **Fig. 4D**) may have a high AASD which, however, has not been detected by our methodological

approach (for example, this may be traced back to the young diversification age of *Drosophila*

species if compared to bivalves); (ii) the species included in the analysis are congeneric, thus the

sequence differentiation of SRGs may exist not at the amino acid level but at the nucleotide one.

To better disentangle this issue and further discuss the fruit fly dataset, we repeated the analysis of

the AASD only on species of the *Crassostrea* genus (*C. gigas*, *Crassostrea angulata*, *Crassostrea*

ariakensis, and *Crassostrea virginica*), which are much younger (Middle Cretaceous [less than

100 Mya; **Qi et al., 2023**]), thus comparable to *Drosophila*. Results showed that, even when

analysing a smaller bivalve dataset, encompassing only 4 species of recent origin, the high AASD

of *Dmrt-1L* persists, that is, *Dmrt-1L* is still grouped together with highly divergent genes (**Supp.**

Fig. S15). The same has not been recovered for *Sox-H*, which fell in genes from Group 3 (the

group corresponding to the 95% interval of the AASD distribution; see **Materials and Methods**)

but still have the second highest AASD median value among DSFGs (**Supp. Fig. S15**).

Of course, we should not expect that highly divergent genes are only those involved in SD but

may also participate in other processes (as discussed earlier and shown by GO term enrichments;

Tab. 1; **Supp. Tab. S8**). Besides the genes of interest for SD (*Dmrt-1L/Sox-H* for bivalves, and

Sry for mammals), other components of the DSFG families have been retrieved with a high

AASD, despite they have never been linked directly to SD so far: *Sox-F* in bivalves (**Fig. 3B**) and

Fox-D4 in mammals (**Fig. 4B**). This implies that our approach cannot be used to unambiguously

identify SDGs alone. Instead, the analysis is meant to be used to detect highly divergent genes

and, subsequently, by comparison with literature and a more thorough and focused functional

investigation, detect putative SDGs among them. In this view, the mammal dataset exemplifies the

importance of putting the results of our pipeline (as those of any other comparative genomics

analysis) into the correct evolutionary and genomic context: among DSFGs of mammals, two

genes exhibit high AASD, one of which is directly related to SD (*Sry*), while the other has a

function connected with neural development (*Fox-D4*; **Klein et al., 2013**). Thus, the high AASD

may arise either because of the involvement in the upper SD pathway or because of other life-

history traits connected with the gene, respectively. Regarding bivalves, *Dmrt-1L* and *Sox-H* show

a sharp connection with SD as a putative primary SDG, either when considering their molecular

evolutionary features or when looking at their gene expression and possible function in gonad

development (**Naimi et al., 2009; Teaniniuraitemoana et al., 2014; Zhang et al., 2014; Capt et

al, 2018; Li et al., 2018; Afonso et al., 2019; Liang et al., 2019; Yue et al., 2021**). It is difficult

to further speculate on the actual involvement in SD of *Dmrt-1L* and *Sox-H* without any additional

information on their biology, also considering that genes evolve not only at the level of their

amino acid sequences, but also at the levels of exon-intron structures, indels, and regulatory

relationships (among others). Nonetheless, molecular evolution proves to be a valuable tool to

investigate genes putatively involved in SD, and to identify major targets onto which dedicate

future research effort.

CONCLUSIONS
619

Genes functioning in reproductive processes, and particularly SDGs, are often among the most

variable in animal genomes, in terms of both sequence composition and regulatory interactions

(**Swanson et al., 2001**; **Bachtrong et al., 2014**). Such high evolutionary rates may be traced back

both to adaptive evolution (either as natural or sexual selection) or to non-adaptive processes

(**Vicoso & Charlesworth, 2006**; **Meisel & Connallon, 2013**; **Parsch & Ellegren, 2013**; **Grath**

& Parsch, 2016) and often results in striking differences in reproductive and sexual systems even

among closely-related species. In the present work we took advantage of this characteristic to

identify SDGs among the DSFG families in bivalves. By comprehensively analysing the

phylogenetic history and AASD in a broad taxonomic dataset, we appointed *Dmrt-1L* and *Sox-H*

as putative SDGs, thus confirming results in previous works that found them to be transcribed in a

male-biased manner and/or strongly involved in male-gonad formation (**Naimi et al., 2009;**

Teaniniuraitemoana et al., 2014; Zhang et al., 2014; Capt et al, 2018; Li et al., 2018; Afonso

et al., 2019; Liang et al., 2019; Yue et al., 2021). Future studies would now need to further

investigate their evolutionary history. For example, considering that SRGs tend to accumulate in

the genomic neighbourhood where primary SDGs are located (**Capel, 2017**), analysing the

genomic location of DSFGs in bivalve genomes may provide enlightening results. Similarly,

revealing the genetic interactions of *Dmrt-1L* and *Sox-H*, through functional and genome editing

assays, would undoubtedly benefit our understanding of their role in the sexual processes of

bivalves.

LIST OF ABBREVIATIONS

AASD: amino acid sequence divergence

DGE: differential gene expression

DM [domain]: *dsx* and *mab-3* [domain]

Dmrt: *doublesex* and *mab-3* related transcription factor

Dmrt-1L: *doublesex and mab-3 related transcription factor 1-like*

Dm-W: *Doublesex and mab-3 related gene W*

Dmy: *Doublesex and mab-3 related gene Y*

DSFGs: Dmrt, Sox, and Fox genes.

dsx: *doublesex*

Fox: forkhead box

ISH: *in-situ* hybridization

Mya: million years ago

SCO: single copy orthogroup

SD: sex determination

SDG: sex-determining gene

Sox: *Sry*-related HMG-box

SRG: sex-determination related gene

Sry: *Sex-determining region Y*

ACKOWLEDGEMENTS

We would like to thank all the people who helped improving this work, by having promoted open discussions and/or exchanged ideas. But above all, we would like to thank Giobbe Forni, Mariangela Iannello and Giovanni Piccinini for always having offered their support whenever it was needed. This work was supported by the Canziani bequest funded to F.G. and A.L., by the European Union - NextGenerationEU under the National Recovery and Resilience Plan (PNRR) –

Mission 4 Education and research – Component 2 From research to business - Investment 1.1,

Notice Prin 2022 - DD N. 104 del 2/2/2022, entitled "Unraveling Sex Determination in Bivalves:

An Integrative Single-Cell Analysis during the Early Embryo Development of *Mytilus*

galloprovincialis", proposal code 2022CYM559 - CUP J53D23006200006 to L.M., and by the

Saltonstall-Kennedy Program by NOAA Fisheries (SK-NOAA) funded to S.N.

REFERENCES

Abascal, F., Zardoya, R., & Telford, M. J. (2010). TranslatorX: multiple alignment of nucleotide

sequences guided by amino acid translations. Nucleic acids research, 38(suppl_2), W7-W13.

Afonso, L. F., Americo, J. A., Soares-Souza, G. B., Torres, A. L. Q., Wajsenzon, I. J. R., &

Rebelo, M. D. F. (2019). Gonad transcriptome of golden mussel Limnoperna fortunei reveals

potential sex differentiation genes. biorxiv, 818757.

Albertin, C. B., Simakov, O., Mitros, T., Wang, Z. Y., Pungor, J. R., Edsinger-Gonzales, E., ... &

Rokhsar, D. S. (2015). The octopus genome and the evolution of cephalopod neural and

morphological novelties. Nature, 524(7564), 220-224.

Alexa, A., Rahnenfuhrer, J. (2024). topGO: Enrichment Analysis for Gene Ontology. R package

version 2.56.0.

Allen, S. L., Delaney, E. K., Kopp, A., & Chenoweth, S. F. (2017). Single-molecule sequencing of

the Drosophila serrata genome. G3: Genes, Genomes, Genetics, 7(3), 781-788.

Altenhoff, A. M., Warwick Vesztrocy, A., Bernard, C., Train, C. M., Nicheperovich, A., Prieto

Baños, S., ... & Glover, N. M. (2024). OMA orthology in 2024: improved prokaryote

coverage, ancestral and extant GO enrichment, a revamped synteny viewer and more in the

OMA Ecosystem. Nucleic Acids Research, 52(D1), D513-D521.

Álvarez-Carretero, S., Tamuri, A. U., Battini, M., Nascimento, F. F., Carlisle, E., Asher, R. J., ...

& Dos Reis, M. (2022). A species-level timeline of mammal evolution integrating

phylogenomic data. Nature, 602(7896), 263-267.

Bachtrog, D., Mank, J. E., Peichel, C. L., Kirkpatrick, M., Otto, S. P., Ashman, T. L., ... & Tree of

Sex Consortium. (2014). Sex determination: why so many ways of doing it?. PLoS biology,

12(7), e1001899.

Bai, C. M., Xin, L. S., Rosani, U., Wu, B., Wang, Q. C., Duan, X. K., ... & Wang, C. M. (2019).

Chromosomal-level assembly of the blood clam, Scapharca (Anadara) broughtonii, using long

sequence reads and Hi-C. GigaScience, 8(7), giz067.

Baldwin-Brown, J. G., Weeks, S. C., & Long, A. D. (2018). A new standard for crustacean

genomes: the highly contiguous, annotated genome assembly of the clam shrimp Eulimnadia

texana reveals HOX gene order and identifies the sex chromosome. Genome biology and

evolution, 10(1), 143-156.

Baral, S., Arumugam, G., Deshmukh, R., & Kunte, K. (2019). Genetic architecture and sex-

specific selection govern modular, male-biased evolution of doublesex. Science advances,

5(5), eaau3753.

Benayoun, B. A., Caburet, S., & Veitia, R. A. (2011). Forkhead transcription factors: key players

in health and disease. Trends in Genetics, 27(6), 224-232.

Beukeboom, L.W., Perrin, N. (2014). The evolution of sex determination. Oxford: Oxford

University Press.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics, 30(15), 2114-2120.

Bredemeyer, K. R., Hillier, L., Harris, A. J., Hughes, G. M., Foley, N. M., Lawless, C., ... &

Murphy, W. J. (2023). Single-haplotype comparative genomics provides insights into lineage-

specific structural variation during cat evolution. Nature genetics, 55(11), 1953-1963.

Breton, S., Capt, C., Guerra, D., & Stewart, D. (2018). Sex-determining mechanisms in bivalves.

Transitions between sexual systems: Understanding the mechanisms of, and pathways

between, dioecy, hermaphroditism and other sexual systems, 165-192.

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using

DIAMOND. Nature methods, 12(1), 59-60.

Capella-Gutiérrez, Salvador, José M. Silla-Martínez, and Toni Gabaldón. "trimAl: a tool for

automated alignment trimming in large-scale phylogenetic analyses." Bioinformatics 25.15

(2009): 1972-1973.

Capt, C., Renaut, S., Ghiselli, F., Milani, L., Johnson, N. A., Sietman, B. E., ... & Breton, S.

(2018). Deciphering the link between doubly uniparental inheritance of mtDNA and sex

determination in bivalves: clues from comparative transcriptomics. Genome biology and

evolution, 10(2), 577-590.

Chakraborty, M., Chang, C. H., Khost, D. E., Vedanayagam, J., Adrion, J. R., Liao, Y., ... &

Emerson, J. J. (2021). Evolution of genome structure in the Drosophila simulans species

complex. Genome research, 31(3), 380-396.

Civetta, A., & Singh, R. S. (1998). Sex-related genes, directional sexual selection, and speciation.

Molecular biology and evolution, 15(7), 901-909.

Corrochano-Fraile, A., Davie, A., Carboni, S., & Bekaert, M. (2022). Evidence of multiple

genome duplication events in Mytilus evolution. BMC genomics, 23(1), 340.

Dainat J. AGAT: Another Gff Analysis Toolkit to handle annotations in any GTF/GFF format.

(Version v0.8.0). Zenodo.

de Bono, M., & Hodgkin, J. (1996). Evolution of sex determination in Caenorhabditis: unusually

high divergence of tra-1 and its functional consequences. Genetics, 144(2), 587-595.

Deng, T., Pang, C., Lu, X., Zhu, P., Duan, A., Tan, Z., ... & Liang, X. (2016). De novo

transcriptome assembly of the Chinese swamp buffalo by RNA sequencing and SSR marker

discovery. PLoS One, 11(1), e0147132.

Dermauw, W., Van Leeuwen, T., & Feyereisen, R. (2020). Diversity and evolution of the P450

family in arthropods. Insect biochemistry and molecular biology, 127, 103490.

Dong, C., Zhang, L., Xia, S., Sosa, D., Arsala, D., & Long, M. (2022). New gene evolution with

subcellular expression patterns detected in PacBio-sequenced genomes of Drosophila genus.

bioRxiv, 2022-11.

Echave, J., Spielman, S. J., & Wilke, C. O. (2016). Causes of evolutionary rate variation among

protein sites. Nature Reviews Genetics, 17(2), 109-121.

Elbers, J. P., Rogers, M. F., Perelman, P. L., Proskuryakova, A. A., Serdyukova, N. A., Johnson,

W. E., ... & Burger, P. A. (2019). Improving Illumina assemblies with Hi□C and long reads:

An example with the North African dromedary. Molecular Ecology Resources, 19(4), 1015-

1026.

Ellegren, H., & Parsch, J. (2007). The evolution of sex-biased genes and sex-biased gene

expression. Nature Reviews Genetics, 8(9), 689-698.

Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome biology, 20, 1-14.

Evensen, K. G., Robinson, W. E., Krick, K., Murray, H. M., & Poynton, H. C. (2022).

Comparative phylotranscriptomics reveals putative sex differentiating genes across eight

diverse bivalve species. Comparative Biochemistry and Physiology Part D: Genomics and

Proteomics, 41, 100952.

Fan, H., Wu, Q., Wei, F., Yang, F., Ng, B. L., & Hu, Y. (2019). Chromosome-level genome

assembly for giant panda provides novel insights into Carnivora chromosome evolution.

Genome biology, 20, 1-12.

Farhat, S., Bonnivard, E., Pales Espinosa, E., Tanguy, A., Boutet, I., Guiglielmoni, N., ... &

Allam, B. (2022). Comparative analysis of the Mercenaria mercenaria genome provides

insights into the diversity of transposable elements and immune molecules in bivalve

mollusks. BMC genomics, 23(1), 192.

Foster, J. W., Brennan, F. E., Hampikian, G. K., Goodfellow, P. N., Sinclair, A. H., Lovell-Badge,

R., ... & Marshall Graves, J. A. (1992). Evolution of sex determination and the Y

chromosome: SRY-related sequences in marsupials. Nature, 359(6395), 531-533.

Foote, A. D., Liu, Y., Thomas, G. W., Vinař, T., Alföldi, J., Deng, J., ... & Gibbs, R. A. (2015).

Convergent evolution of the genomes of marine mammals. Nature genetics, 47(3), 272-275.

Fouret, J., Brunet, F. G., Binet, M., Aurine, N., Enchéry, F., Croze, S., ... & Legras-Lachuer, C.

(2020). Sequencing the genome of Indian flying fox, natural reservoir of nipah virus, using

hybrid assembly and conservative secondary scaffolding. Frontiers in Microbiology, 11, 1807.

Gerdol, M., Moreira, R., Cruz, F., Gómez-Garrido, J., Vlasova, A., Rosani, U., ... & Figueras, A.

(2020). Massive gene presence-absence variation shapes an open pan-genome in the

Mediterranean mussel. Genome biology, 21, 1-21.

Ghiselli, F., Milani, L., Chang, P. L., Hedgecock, D., Davis, J. P., Nuzhdin, S. V., & Passamonti,

M. (2012). De novo assembly of the Manila clam Ruditapes philippinarum transcriptome

provides new insights into expression bias, mitochondrial doubly uniparental inheritance and

sex determination. Molecular biology and evolution, 29(2), 771-786.

Ghiselli, F., Iannello, M., Puccio, G., Chang, P. L., Plazzi, F., Nuzhdin, S. V., & Passamonti, M.

(2018). Comparative transcriptomics in two bivalve species offers different perspectives on

the evolution of sex-biased genes. Genome biology and evolution, 10(6), 1389-1402.

Gomes-dos-Santos, A., Lopes-Lima, M., Castro, L. F. C., & Froufe, E. (2020). Molluscan

genomics: the road so far and the way forward. Hydrobiologia, 847(7), 1705-1726.

Gomes-dos-Santos, A., Lopes-Lima, M., Machado, A. M., Marcos Ramos, A., Usié, A., Bolotov,

I. N., ... & Froufe, E. (2021). The Crown Pearl: a draft genome assembly of the European

freshwater pearl mussel Margaritifera margaritifera (Linnaeus, 1758). DNA research, 28(2),

dsab002.

Gomes-dos-Santos, A., Lopes-Lima, M., Machado, M. A., Teixeira, A., C. Castro, L. F., &

Froufe, E. (2023). PacBio Hi-Fi genome assembly of the Iberian dolphin freshwater mussel

Unio delphinus Spengler, 1793. Scientific Data, 10(1), 340.

Gómez-Chiarri, M., Warren, W. C., Guo, X., & Proestou, D. (2015). Developing tools for the

study of molluscan immunity: the sequencing of the genome of the eastern oyster, Crassostrea

virginica. Fish & shellfish immunology, 46(1), 2-4.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., ... & Regev, A.

(2011). Trinity: reconstructing a full-length transcriptome without a genome from RNA-Seq

data. Nature biotechnology, 29(7), 644.

Grath, S., & Parsch, J. (2016). Sex-biased gene expression. Annual review of genetics, 50(1), 29-

44.

Grau-Bové, X., & Sebé-Pedrós, A. (2021). Orthology clusters from gene trees with Possvm.

Molecular Biology and Evolution, 38(11), 5204-5208.

Haag, E. S., & Doty, A. V. (2005). Sex determination across evolution: connecting the dots. PLoS

biology, 3(1), e21.

Haas, BJ. https://github.com/TransDecoder/TransDecoder.

Habtewold, T., Wagah, M., Tambwe, M. M., Moore, S., Windbichler, N., Christophides, G., ... &

Lawniczak, M. (2023). A chromosomal reference genome sequence for the malaria mosquito,

Anopheles gambiae, Giles, 1902, Ifakara strain. Wellcome Open Research, 8.

Han, F., Wang, Z., Wu, F., Liu, Z., Huang, B., & Wang, D. (2010). Characterization, phylogeny,

alternative splicing and expression of Sox30 gene. BMC Molecular Biology, 11, 1-11.

Hart, S. F., Yonemitsu, M. A., Giersch, R. M., Garrett, F. E., Beal, B. F., Arriagada, G., ... &

Metzger, M. J. (2023). Centuries of genome instability and evolution in soft-shell clam, Mya

arenaria, bivalve transmissible neoplasia. Nature Cancer, 4(11), 1561-1574.

Haszprunar, G., & Wanninger, A. (2012). Molluscs. Current Biology, 22(13), R510-R514.

Haerty, W., Jagadeeshan, S., Kulathinal, R. J., Wong, A., Ravi Ram, K., Sirot, L. K., ... & Singh,

R. S. (2007). Evolution in the fast lane: rapidly evolving sex-related genes in Drosophila.

Genetics, 177(3), 1321-1335.

Heenan, P., Zondag, L., & Wilson, M. J. (2016). Evolution of the Sox gene family within the

chordate phylum. Gene, 575(2), 385-392.

Hemmer, L. W., & Blumenstiel, J. P. (2016). Holding it together: rapid evolution and positive

selection in the synaptonemal complex of Drosophila. BMC evolutionary biology, 16, 1-17.

Hoskins, R. A., Carlson, J. W., Wan, K. H., Park, S., Mendez, I., Galle, S. E., ... & Celniker, S. E.

(2015). The Release 6 reference sequence of the Drosophila melanogaster genome. Genome

research, 25(3), 445-458.

Iannello, M., Forni, G., Piccinini, G., Xu, R., Martelossi, J., Ghiselli, F., & Milani, L. (2023).

Signatures of extreme longevity: a perspective from bivalve molecular evolution. Genome

Biology and Evolution, 15(11), evad159.

Inoue, K., Yoshioka, Y., Tanaka, H., Kinjo, A., Sassa, M., Ueda, I., ... & Itoh, T. (2021).

Genomics and transcriptomics of the green mussel explain the durability of its byssus.

Scientific Reports, 11(1), 5992.

Ip, J. C. H., Xu, T., Sun, J., Li, R., Chen, C., Lan, Y., ... & Qiu, J. W. (2021). Host–endosymbiont

genome integration in a deep-sea chemosymbiotic clam. Molecular Biology and Evolution,

38(2), 502-518.

Jackson, B. C., Carpenter, C., Nebert, D. W., & Vasiliou, V. (2010). Update of human and mouse

forkhead box (FOX) gene families. Human genomics, 4, 1-8.

Jombart, T., & Dray, S. (2010). Adephylo: exploratory analyses for the phylogenetic comparative

method. Bioinformatics, 26(15), 1-21.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., & Jermiin, L. S. (2017).

ModelFinder: fast model selection for accurate phylogenetic estimates. Nature methods,

14(6), 587-589.

Kenny, N. J., McCarthy, S. A., Dudchenko, O., James, K., Betteridge, E., Corton, C., ... &

Williams, S. T. (2020). The gene-rich genome of the scallop Pecten maximus. Gigascience,

9(5), giaa037.

Kim, B. Y., Wang, J. R., Miller, D. E., Barmina, O., Delaney, E., Thompson, A., ... & Petrov, D.

A. (2021). Highly contiguous assemblies of 101 drosophilid genomes. Elife, 10, e66405.

Klein, S. L., Neilson, K. M., Orban, J., Yaklichkin, S., Hoffbauer, J., Mood, K., ... & Moody, S. A.

(2013). Conserved structural domains in FoxD4L1, a neural forkhead box transcription factor,

are required to repress or activate target genes. PloS one, 8(4), e61845.

Knudsen, B., Kohn, A. B., Nahir, B., McFadden, C. S., & Moroz, L. L. (2006). Complete DNA

sequence of the mitochondrial genome of the sea-slug, Aplysia californica: conservation of

the gene order in Euthyneura. Molecular Phylogenetics and Evolution, 38(2), 459-469.

Lan, Y., Sun, J., Chen, C., Sun, Y., Zhou, Y., Yang, Y., ... & Qian, P. Y. (2021). Hologenome

analysis reveals dual symbiosis in the deep-sea hydrothermal vent snail Gigantopelta aegis.

Nature communications, 12(1), 1165.

Larroux, C., Luke, G. N., Koopman, P., Rokhsar, D. S., Shimeld, S. M., & Degnan, B. M. (2008).

Genesis and expansion of metazoan transcription factor gene classes. Molecular biology and

evolution, 25(5), 980-996.

Lemoine, F., & Gascuel, O. (2021). Gotree/Goalign: toolkit and Go API to facilitate the

development of phylogenetic workflows. NAR Genomics and Bioinformatics, 3(3), lqab075.

Li, C., Liu, X., Liu, B., Ma, B., Liu, F., Liu, G., ... & Wang, C. (2018). Draft genome of the

Peruvian scallop Argopecten purpuratus. GigaScience, 7(4), giy031.

Li, R., Zhang, L., Li, W., Zhang, Y., Li, Y., Zhang, M., ... & Bao, Z. (2018). FOXL2 and

DMRT1L are yin and yang genes for determining timing of sex differentiation in the bivalve

mollusk Patinopecten yessoensis. Frontiers in Physiology, 9, 1166.

Li, F., Bian, L., Ge, J., Han, F., Liu, Z., Li, X., ... & Chen, S. (2020). Chromosome□level genome

assembly of the East Asian common octopus (Octopus sinensis) using PacBio sequencing and

Hi□C technology. Molecular ecology resources, 20(6), 1572-1582.

Li, A., Dai, H., Guo, X., Zhang, Z., Zhang, K., Wang, C., ... & Zhang, G. (2021). Genome of the

estuarine oyster provides insights into climate impact and adaptive plasticity.

Communications Biology, 4(1), 1287.

Li, X. Y., Mei, J., Ge, C. T., Liu, X. L., & Gui, J. F. (2022). Sex determination mechanisms and

sex control approaches in aquaculture animals. Science China Life Sciences, 65(6), 1091-

1122.

Li, H. (2023). Protein-to-genome alignment with miniprot. Bioinformatics, 39(1), btad014.

Liang, S., Liu, D., Li, X., Wei, M., Yu, X., Li, Q., ... & Qin, Z. (2019). SOX2 participates in

spermatogenesis of Zhikong scallop Chlamys farreri. Scientific Reports, 9(1), 76.

Liao, Y., Zhang, X., Chakraborty, M., & Emerson, J. J. (2021). Topologically associating domains

and their role in the evolution of genome structure and function in Drosophila. Genome

research, 31(3), 397-410.

Liu, X. L., Zhang, Z. F., Shao, M. Y., Liu, J. G., & Muhammad, F. (2012). Sexually dimorphic

expression of foxl2 during gametogenesis in scallop Chlamys farreri, conserved with

vertebrates. Development Genes and Evolution, 222, 279-286.

Liu, C., Zhang, Y., Ren, Y., Wang, H., Li, S., Jiang, F., ... & Fan, W. (2018). The genome of the

golden apple snail Pomacea canaliculata provides insight into stress tolerance and invasive

adaptation. Gigascience, 7(9), giy101.

Liu, X., Li, C., Chen, M., Liu, B., Yan, X., Ning, J., ... & Wang, C. (2020). Draft genomes of two

Atlantic bay scallop subspecies Argopecten irradians irradians and A. i. concentricus.

Scientific Data, 7(1), 99.

Liu, T., Zhang, Y., Nie, H., Sun, J., & Yan, X. (2024). Characterization and expression patterns of

the Fox gene family under heat and cold stress in Manila clam Ruditapes philippinarum based

on genome-wide identification. Comparative Biochemistry and Physiology Part D: Genomics

and Proteomics, 101313.

Mahajan, S., Wei, K. H. C., Nalley, M. J., Gibilisco, L., & Bachtrog, D. (2018). De novo assembly

of a young Drosophila Y chromosome using single-molecule sequencing and chromatin

conformation capture. PLoS biology, 16(7), e2006348.

Mai, D., Nalley, M. J., & Bachtrog, D. (2020). Patterns of genomic differentiation in the

Drosophila nasuta species complex. Molecular biology and evolution, 37(1), 208-220.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO

update: novel and streamlined workflows along with broader and deeper phylogenetic

coverage for scoring of eukaryotic, prokaryotic, and viral genomes. Molecular biology and

evolution, 38(10), 4647-4654.

Matson, C. K., & Zarkower, D. (2012). Sex and the singular DM domain: insights into sexual

regulation, evolution and plasticity. Nature Reviews Genetics, 13(3), 163-174.

Mawaribuchi, S., Yoshimoto, S., Ohashi, S., Takamatsu, N., & Ito, M. (2012). Molecular

evolution of vertebrate sex-determining genes. Chromosome Research, 20, 139-151.

Mawaribuchi, S., Ito, Y., & Ito, M. (2019). Independent evolution for sex determination and

differentiation in the DMRT family in animals. Biology Open, 8(8), bio041962.

Mazet, F., Yu, J. K., Liberles, D. A., Holland, L. Z., & Shimeld, S. M. (2003). Phylogenetic

relationships of the Fox (Forkhead) gene family in the Bilateria. Gene, 316, 79-89.

McCartney, M. A., Auch, B., Kono, T., Mallez, S., Zhang, Y., Obille, A., ... & Gohl, D. M.

(2022). The genome of the zebra mussel, Dreissena polymorpha: a resource for comparative

genomics, invasion genetics, and biocontrol. G3, 12(2), jkab423.

Mead, D., Fingland, K., Cripps, R., Miguez, R. P., Smith, M., Corton, C., ... & Blaxter, M. (2020).

The genome sequence of the eastern grey squirrel, Sciurus carolinensis Gmelin, 1788.

Wellcome Open Research, 5.

Meisel, R. P. (2011). Towards a more nuanced understanding of the relationship between sex-
biased gene expression and rates of protein-coding sequence evolution. Molecular Biology
and Evolution, 28(6), 1893-1900.

Meisel, R. P., & Connallon, T. (2013). The faster-X effect: integrating theory and data. Trends in
genetics, 29(9), 537-544.

Milani, L., Ghiselli, F., Nuzhdin, S. V., & Passamonti, M. (2013). Nuclear genes with sex bias in
Ruditapes philippinarum (Bivalvia, Veneridae): mitochondrial inheritance and sex
determination in DUI species. Journal of Experimental Zoology Part B: Molecular and
Developmental Evolution, 320(7), 442-454.

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A.,
& Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic
inference in the genomic era. Molecular biology and evolution, 37(5), 1530-1534.

Moreno, J. A., Dudchenko, O., Feigin, C. Y., Mereby, S. A., Chen, Z., Ramos, R., ... & Mallarino,
R. (2024). Emx2 underlies the development and evolution of marsupial gliding membranes.
Nature, 1-9.

Mullon, C., Pomiankowski, A., & Reuter, M. (2012). Molecular evolution of Drosophila Sex-
lethal and related sex determining genes. BMC evolutionary biology, 12, 1-11.

Murat, F., Mbengue, N., Winge, S. B., Trefzer, T., Leushkin, E., Sepp, M., ... & Kaessmann, H.
(2023). The molecular evolution of spermatogenesis across mammals. Nature, 613(7943),
308-316.

Murata, Y., Nikaido, M., Sasaki, T., Cao, Y., Fukumoto, Y., Hasegawa, M., & Okada, N. (2003).
Afrotherian phylogeny as inferred from complete mitochondrial genomes. Molecular
phylogenetics and evolution, 28(2), 253-260.

Naimi, A., Martinez, A. S., Specq, M. L., Diss, B., Mathieu, M., & Sourdaine, P. (2009).
Molecular cloning and gene expression of *Cg-Foxl2* during the development and the adult

gametogenetic cycle in the oyster *Crassostrea gigas*. Comparative Biochemistry and

Physiology Part B: Biochemistry and Molecular Biology, 154(1), 134-142.

Nakagawa, S., Gisselbrecht, S. S., Rogers, J. M., Hartl, D. L., & Bulyk, M. L. (2013). DNA-

binding specificity changes in the evolution of forkhead transcription factors. Proceedings of

the National Academy of Sciences, 110(30), 12349-12354.

Nicolini, F., Ghiselli, F., Luchetti, A., & Milani, L. (2023). Bivalves as emerging model systems

to study the mechanisms and evolution of sex determination: a genomic point of view.

Genome Biology and Evolution, 15(10), evad181.

Nicolini, F., Martelossi, J., Forni, G., Savojardo, C., Mantovani, B., & Luchetti, A. (2023).

Comparative genomics of Hox and ParaHox genes among major lineages of Branchiopoda

with emphasis on tadpole shrimps. Frontiers in Ecology and Evolution, 11, 1046960.

O'Neil, M. T., & Belote, J. M. (1992). Interspecific comparison of the transformer gene of

Drosophila reveals an unusually high degree of evolutionary divergence. Genetics, 131(1),

113-128.

Orkin, J. D., Montague, M. J., Tejada-Martinez, D., De Manuel, M., Del Campo, J., Cheves

Hernandez, S., ... & Melin, A. D. (2021). The genomics of ecological flexibility, large brains,

and long lives in capuchin monkeys revealed with fecalFACS. Proceedings of the National

Academy of Sciences, 118(7), e2010632118.

Paggeot, L. X., DeBiasse, M. B., Escalona, M., Fairbairn, C., Marimuthu, M. P., Nguyen, O., ... &

Dawson, M. N. (2022). Reference genome for the California ribbed mussel, Mytilus

californianus, an ecosystem engineer. Journal of Heredity, 113(6), 681-688.

Pamilo, P., & O'Neill, R. J. (1997). Evolution of the Sry genes. Molecular biology and evolution,

14(1), 49-55.

Panara, V., Budd, G. E., & Janssen, R. (2019). Phylogenetic analysis and embryonic expression of

panarthropod Dmrt genes. Frontiers in zoology, 16, 1-18.

Papa, F., Windbichler, N., Waterhouse, R. M., Cagnetti, A., D'Amato, R., Persampieri, T., ... &

Papathanos, P. A. (2017). Rapid evolution of female-biased genes among four species of

Anopheles malaria mosquitoes. Genome research, 27(9), 1536-1548.

Paris, M., Boyer, R., Jaenichen, R., Wolf, J., Karageorgi, M., Green, J., ... & Prud'homme, B.

(2020). Near-chromosome level genome assembly of the fruit pest Drosophila suzukii using

long-read sequencing. Scientific reports, 10(1), 11227.

Parsch, J., & Ellegren, H. (2013). The evolutionary causes and consequences of sex-biased gene

expression. Nature Reviews Genetics, 14(2), 83-87.

Penaloza, C., Gutierrez, A. P., Eöry, L., Wang, S., Guo, X., Archibald, A. L., ... & Houston, R. D.

(2021). A chromosome-level genome assembly for the Pacific oyster Crassostrea gigas.

GigaScience, 10(3), giab020.

Phochanukul, N., & Russell, S. (2010). No backbone but lots of Sox: Invertebrate Sox genes. The

international journal of biochemistry & cell biology, 42(3), 453-464.

Piccinini, G., Iannello, M., Puccio, G., Plazzi, F., Havird, J. C., & Ghiselli, F. (2021). Mitonuclear

coevolution, but not nuclear compensation, drives evolution of OXPHOS complexes in

bivalves. Molecular Biology and Evolution, 38(6), 2597-2614.

Powell, D., Subramanian, S., Suwansa-Ard, S., Zhao, M., O'Connor, W., Raftos, D., & Elizur, A.

(2018). The genome of the oyster Saccostrea offers insight into the environmental resilience

of bivalves. DNA Research, 25(6), 655-665.

Qi, H., Cong, R., Wang, Y., Li, L., & Zhang, G. (2023). Construction and analysis of the

chromosome-level haplotype-resolved genomes of two Crassostrea oyster congeners:

Crassostrea angulata and Crassostrea gigas. GigaScience, 12, giad077.

Ran, Z., Li, Z., Yan, X., Liao, K., Kong, F., Zhang, L., ... & Xu, J. (2019). Chromosome□level

genome assembly of the razor clam Sinonovacula constricta (Lamarck, 1818). Molecular

ecology resources, 19(6), 1647-1658.

Ranz, J. M., Go, A. C., González, P. M., Clifton, B. D., Gomes, S., Jaberyzadeh, A., ... & Civetta,

A. (2023). Gene expression differentiation in the reproductive tissues of Drosophila willistoni

subspecies and their hybrids. Molecular Ecology, 32(13), 3605-3623.

Renschler, G., Richard, G., Valsecchi, C. I. K., Toscano, S., Arrigoni, L., Ramírez, F., & Akhtar,

A. (2019). Hi-C guided assemblies reveal conserved regulatory topologies on X and

autosomes despite extensive genome shuffling. Genes & development, 33(21-22), 1591-1612.

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., ... & Jarvis, E. D.

(2021). Towards complete and error-free genome assemblies of all vertebrate species. Nature,

592(7856), 737-746.

Rogers, R. L., Grizzard, S. L., Titus☐McQuillan, J. E., Bockrath, K., Patel, S., Wares, J. P., ... &

Moore, C. C. (2021). Gene family amplification facilitates adaptation in freshwater unionid

bivalve Megalonaias nervosa. Molecular Ecology, 30(5), 1155-1173.

Russo, C. A., Mello, B., Frazão, A., & Voloch, C. M. (2013). Phylogenetic analysis and a time

tree for a large drosophilid data set (Diptera: Drosophilidae). Zoological Journal of the

Linnean Society, 169(4), 765-775.

Sanchez-Flores, A., Peñaloza, F., Carpinteyro-Ponce, J., Nazario-Yepiz, N., Abreu-Goodger, C.,

Machado, C. A., & Markow, T. A. (2016). Genome evolution in three species of cactophilic

Drosophila. G3: Genes, Genomes, Genetics, 6(10), 3097-3105.

Sarkar, A., & Hochedlinger, K. (2013). The sox family of transcription factors: versatile regulators

of stem and progenitor cell fate. Cell stem cell, 12(1), 15-30.

Schliep, K. P. (2011). phangorn: phylogenetic analysis in R. Bioinformatics, 27(4), 592-593.

Schomburg, C., Janssen, R., & Prpic, N. M. (2022). Phylogenetic analysis of forkhead

transcription factors in the Panarthropoda. Development genes and evolution, 232(1), 39-48.

Seudre, O., Martín-Zamora, F. M., Rapisarda, V., Luqman, I., Carrillo-Baltodano, A. M., &

Martín-Durán, J. M. (2022). The Fox gene repertoire in the annelid Owenia fusiformis reveals

multiple expansions of the foxQ2 class in Spiralia. Genome Biology and Evolution, 14(10),

evac139.

Shimeld, S. M., Boyle, M. J., Brunet, T., Luke, G. N., & Seaver, E. C. (2010). Clustered Fox

genes in lophotrochozoans and the evolution of the bilaterian Fox gene cluster.

Developmental biology, 340(2), 234-248.

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., ... & Higgins, D. G. (2011).

Fast, scalable generation of high□quality protein multiple sequence alignments using Clustal

Omega. Molecular systems biology, 7(1), 539.

Smith, C. H. (2021). A high-quality reference genome for a parasitic bivalve with doubly

uniparental inheritance (Bivalvia: Unionida). Genome Biology and Evolution, 13(3),

evab029.

Song, H., Wang, Y., Shao, H., Li, Z., Hu, P., Yap-Chiongco, M. K., ... & Kocot, K. M. (2023).

Scaphopoda is the sister taxon to Bivalvia: Evidence of ancient incomplete lineage sorting.

Proceedings of the National Academy of Sciences, 120(40), e2302361120.

Stammnitz, M. R., Gori, K., Kwon, Y. M., Harry, E., Martin, F. J., Billis, K., ... & Murchison, E.

P. (2023). The evolution of two transmissible cancers in Tasmanian devils. Science,

380(6642), 283-293.

Stothard, P., & Pilgrim, D. (2003). Sex□determination gene and pathway evolution in nematodes.

Bioessays, 25(3), 221-231.

Sun, J., Zhang, Y. U., Xu, T., Zhang, Y., Mu, H., Zhang, Y., ... & Qian, P. Y. (2017). Adaptation

to deep-sea chemosynthetic environments as revealed by mussel genomes. Nature ecology &

evolution, 1(5), 0121.

Sun, D., Yu, H., & Li, Q. (2022). Examination of the roles of Foxl2 and Dmrt1 in sex

differentiation and gonadal development of oysters by using RNA interference. Aquaculture,

548, 737732.

Swanson, W. J., & Vacquier, V. D. (2002). The rapid evolution of reproductive proteins. Nature

reviews genetics, 3(2), 137-144.

Teaniniuraitemoana, V., Huvet, A., Levy, P., Klopp, C., Lhuillier, E., Gaertner-Mazouni, N., ... &

Le Moullac, G. (2014). Gonad transcriptome analysis of pearl oyster Pinctada margaritifera:

identification of potential sex differentiation and sex determining genes. BMC genomics, 15,

1-20.

Teng, W., Fu, H., Li, Z., Zhang, Q., Xu, C., Yu, H., ... & Li, Q. (2023). Parallel evolution in

Crassostrea oysters along the latitudinal gradient is associated with variation in multiple genes

involved in adipogenesis. Molecular Ecology, 32(19), 5276-5287.

Tu, Q., Brown, C. T., Davidson, E. H., & Oliveri, P. (2006). Sea urchin Forkhead gene family:

phylogeny and embryonic expression. Developmental biology, 300(1), 49-62.

Tvedte, E. S., Gasser, M., Sparklin, B. C., Michalski, J., Hjelmen, C. E., Johnston, J. S., ... &

Dunning Hotopp, J. C. (2021). Comparison of long-read sequencing technologies in

interrogating bacteria and fly genomes. G3, 11(6), jkab083.

Uller, T., & Helanterä, H. (2011). From the origin of sex-determining factors to the evolution of

sex-determining systems. The Quarterly review of biology, 86(3), 163-180.

Vicoso, B., & Charlesworth, B. (2006). Evolution on the X chromosome: unusual patterns and

processes. Nature Reviews Genetics, 7(8), 645-653.

Vilstrup, J. T., Seguin-Orlando, A., Stiller, M., Ginolhac, A., Raghavan, M., Nielsen, S. C., ... &

Orlando, L. (2013). Mitochondrial phylogenomics of modern and ancient equids. PloS one,

8(2), e55950.

Vinkler, M., Fiddaman, S. R., Těšický, M., O'Connor, E. A., Savage, A. E., Lenz, T. L., ... &

Westerdahl, H. (2023). Understanding the evolution of immune genes in jawed vertebrates.

Journal of evolutionary biology, 36(6), 847-873.

Vizueta, J., Sánchez☐Gracia, A., & Rozas, J. (2020). BITACORA: A comprehensive tool for the

identification and annotation of gene families in genome assemblies.

1065 Wang, G., Dong, S., Guo, P., Cui, X., Duan, S., & Li, J. (2020). Identification of Foxl2 in

1066  freshwater mussel Hyriopsis cumingii and its involvement in sex differentiation. Gene, 754,

1067  144853.

1068 Wang, C., Wallerman, O., Arendt, M. L., Sundström, E., Karlsson, Å., Nordin, J., ... & Lindblad-

1069  Toh, K. (2021). A novel canine reference genome resolves genomic architecture and uncovers

1070  transcript complexity. Communications biology, 4(1), 185.

1071 Wang, S., Zhang, J., Jiao, W., Li, J. I., Xun, X., Sun, Y., ... & Bao, Z. (2017). Scallop genome

1072  provides insights into evolution of bilaterian karyotype and development. Nature ecology &

1073  evolution, 1(5), 0120.

1074 Wang, Q., Cao, T., & Wang, C. (2023). Genome-wide identification and expression analysis of

1075  Dmrt genes in bivalves. BMC genomics, 24(1), 457.

1076 Wang, J., & Nie, H. (2024). Genome-wide identification and expression analysis of Sox gene

1077  family in the Manila clam (Ruditapes philippinarum). Comparative Biochemistry and

1078  Physiology Part D: Genomics and Proteomics, 50, 101244.

1079 Wei, M., Ge, H., Shao, C., Yan, X., Nie, H., Duan, H., ... & Dong, Z. (2020). Chromosome-level

1080  clam genome helps elucidate the molecular basis of adaptation to a buried lifestyle. IScience,

1081  23(6).

1082 Wexler, J. R., Plachetzki, D. C., & Kopp, A. (2014). Pan-metazoan phylogeny of the DMRT gene

1083  family: a framework for functional studies. Development Genes and Evolution, 224, 175-181.

1084 Whitfield, L. S., Lovell-Badge, R., & Goodfellow, P. N. (1993). Rapid sequence evolution of the

1085  mammalian sex-determining gene SRY. Nature, 364(6439), 713-715.

1086 Wilkins, A. S. (1995). Moving up the hierarchy: a hypothesis on the evolution of a genetic sex

1087  determination pathway. Bioessays, 17(1), 71-77.

1088 Willson, J., Roddur, M. S., Liu, B., Zaharias, P., & Warnow, T. (2022). DISCO: species tree

1089  inference using multicopy gene family tree decomposition. Systematic Biology, 71(3), 610-

1090  629.

Wu, S., Zhang, Y., Li, Y., Wei, H., Guo, Z., Wang, S., ... & Bao, Z. (2020). Identification and

expression profiles of Fox transcription factors in the Yesso scallop (Patinopecten

yessoensis). Gene, 733, 144387.

Xiong, Y. E., Brandley, M. C., Xu, S., Zhou, K., & Yang, G. (2009). Seven new dolphin

mitochondrial genomes and a time-calibrated phylogeny of whales. BMC Evolutionary

Biology, 9, 1-13.

Xu, R., Martelossi, J., Smits, M., Iannello, M., Peruzza, L., Babbucci, M., ... & Ghiselli, F. (2022).

Multi-tissue RNA-Seq analysis and long-read-based genome assembly reveal complex sex-

specific gene regulation and molecular evolution in the Manila clam. Genome Biology and

Evolution, 14(12), evac171.

Yang, M., Xu, F., Liu, J., Que, H., Li, L., & Zhang, G. (2014). Phylogeny of forkhead genes in

three spiralians and their expression in Pacific oyster Crassostrea gigas. Chinese journal of

oceanology and limnology, 32(6), 1207-1223.

Yang, J. L., Feng, D. D., Liu, J., Xu, J. K., Chen, K., Li, Y. F., ... & Lu, Y. (2021). Chromosome-

level genome assembly of the hard-shelled mussel Mytilus coruscus, a widely distributed

species from the temperate areas of East Asia. Gigascience, 10(4), giab024.

Yu, J. K., Mazet, F., Chen, Y. T., Huang, S. W., Jung, K. C., & Shimeld, S. M. (2008). The Fox

genes of Branchiostoma floridae. Development genes and evolution, 218, 629-638.

Yu, G., Smith, D. K., Zhu, H., Guan, Y., & Lam, T. T. Y. (2017). ggtree: an R package for

visualization and annotation of phylogenetic trees with their covariates and other associated

data. Methods in Ecology and Evolution, 8(1), 28-36.

Yu, J., Zhang, L., Li, Y., Li, R., Zhang, M., Li, W., ... & Bao, Z. (2017). Genome-wide

identification and expression profiling of the SOX gene family in a bivalve mollusc

Patinopecten yessoensis. Gene, 627, 530-537.

Yue, C., Li, Q., & Yu, H. (2021). Variance in expression and localization of sex-related genes

CgDsx, CgBHMG1 and CgFoxl2 during diploid and triploid Pacific oyster Crassostrea gigas

gonad differentiation. Gene, 790, 145692.

Zeng, Y., Zheng, H., He, C., Zhang, C., Zhang, H., & Zheng, H. (2024). Genome-wide

identification and expression analysis of Dmrt gene family and their role in gonad

development of Pacific oyster (Crassostrea gigas). Comparative Biochemistry and Physiology

Part B: Biochemistry and Molecular Biology, 269, 110904.

Zhang, N., Xu, F., & Guo, X. (2014). Genomic analysis of the Pacific oyster (*Crassostrea gigas*)

reveals possible conservation of vertebrate sex determination in a mollusc. G3: Genes,

Genomes, Genetics, 4(11), 2207-2217.

Zhou, Y., Shearwin-Whyatt, L., Li, J., Song, Z., Hayakawa, T., Stevens, D., ... & Zhang, G.

(2021). Platypus and echidna genomes reveal mammalian biology and evolution. Nature,

592(7856), 756-762.

1128   DATA ACCESSIBILITY AND BENEFIT-SHARING

Data accessibility statement

The data underlying this article and the codes used to run the analyses are available on GitHub

(https://github.com/filonico/bivalvia_SRGs).

Benefit-sharing statement

Benefits from this research accrue from the sharing of our data, codes, and results on public

databases as described above.

1135   AUTHOR CONTIRBUTIONS

1136   FN, AL, FG, and LM conceived the study. FN performed research and analysed data. AL, FG and

1137   SN provided computational resources. FN wrote the paper. All authors edited and approved the

1138   manuscript.

1139

TABLES

Tab. 1. Top enriched GO terms for highly divergent genes of bivalves, mammals, and *Drosophila***.** The extended version of the table, which includes also the expected number of annotated genes per GO term and all the other enriched GO terms, can be accessed in **Supp. Tab. S8**.

Dataset	GO.ID	Term	Annotated genes	Significant genes	Corrected p-value
Bivalvia	GO:0060255	regulation of macromolecule metabolic process	737	59	0.04525
	GO:0080090	regulation of primary metabolic process	673	53	0.01818
	GO:0019219	regulation of nucleobase-containing compound metabolic process	541	41	0.02388
	GO:0006351	DNA-templated transcription	571	39	0.03767
	GO:0032774	RNA biosynthetic process	579	39	0.04490
	GO:0051252	regulation of RNA metabolic process	517	37	0.02719
	GO:0006355	regulation of DNA-templated transcription	490	35	0.03751
	GO:2001141	regulation of RNA biosynthetic process	491	35	0.03844
	GO:0006950	response to stress	370	33	0.01949
	GO:0032502	developmental process	261	27	0.04445
	GO:0006468	protein phosphorylation	345	23	0.02483
	GO:0031325	positive regulation of cellular metabolic process	125	17	0.00801
	GO:0010604	positive regulation of macromolecule metabolic process	151	17	0.04047
	GO:0051172	negative regulation of nitrogen compound metabolic process	117	16	0.00814
	GO:0051173	positive regulation of nitrogen compound metabolic process	137	15	0.02454
	GO:0006310	DNA recombination	66	14	0.00087
	GO:0048513	animal organ development	83	12	0.04088
	GO:0010629	negative regulation of gene expression	78	11	0.00048
	GO:0023051	regulation of signaling	133	11	0.02872
	GO:0045934	negative regulation of nucleobase-containing compound metabolic process	64	11	0.03637
	GO:0009605	response to external stimulus	90	11	0.04544
	GO:0044419	biological process involved in interspecies interaction between organisms	63	11	0.04761
Mammalia	GO:0006955	immune response	1297	145	0.00061
	GO:0098542	defense response to other organism	853	112	0.02066
	GO:0045087	innate immune response	647	82	8.5e-10
	GO:0001817	regulation of cytokine production	630	51	0.04660
	GO:0042742	defense response to bacterium	233	45	1.7e-07
	GO:0006954	inflammatory response	642	45	0.01735
	GO:0019221	cytokine-mediated signaling pathway	382	44	3.9e-07
	GO:0002250	adaptive immune response	342	44	1.3e-05
	GO:0001819	positive regulation of cytokine production	402	41	0.02723
	GO:0002697	regulation of immune effector process	308	37	0.04426
	GO:0042110	T cell activation	432	35	0.02564
	GO:0051607	defense response to virus	257	34	1.9e-07
	GO:0048232	male gamete generation	491	32	0.02255

	GO:0007283	spermatogenesis	478	31	0.02801
	GO:0070661	leukocyte proliferation	273	29	0.01285
	GO:0002449	lymphocyte mediated immunity	221	29	0.04833
	GO:0070663	regulation of leukocyte proliferation	212	25	0.01870
	GO:0050727	regulation of inflammatory response	300	24	0.00235
	GO:0031349	positive regulation of defense response	240	24	0.01239
	GO:0002768	immune response-regulating cell surface receptor signaling pathway	177	22	0.00336
	GO:0050829	defense response to Gram-negative bacterium	66	17	1.7e-10
	GO:0071222	cellular response to lipopolysaccharide	164	17	0.00012
	GO:0010466	negative regulation of peptidase activity	163	16	0.00036
	GO:0002429	immune response-activating cell surface receptor signaling pathway	164	16	0.00243
	GO:1903555	regulation of tumor necrosis factor superfamily cytokine production	137	16	0.01244
	GO:0071706	tumor necrosis factor superfamily cytokine production	137	16	0.01244
	GO:0070665	positive regulation of leukocyte proliferation	132	16	0.02765
	GO:0045089	positive regulation of innate immune response	113	16	0.03224
	GO:0071356	cellular response to tumor necrosis factor	175	15	0.00219
	GO:0002695	negative regulation of leukocyte activation	148	15	0.01151
	GO:0002456	T cell mediated immunity	82	15	0.01605
	GO:0002705	positive regulation of leukocyte mediated immunity	113	15	0.01837
	GO:0032680	regulation of tumor necrosis factor production	133	15	0.03262
	GO:0032640	tumor necrosis factor production	133	15	0.03262
	GO:0050866	negative regulation of cell activation	165	15	0.04048
	GO:0000819	sister chromatid segregation	140	11	0.02927
	GO:0070192	chromosome organization involved in meiotic cell cycle	54	9	0.00849
	GO:0007131	reciprocal meiotic recombination	37	7	0.00066
	GO:0007143	female meiotic nuclear division	54	6	0.02270
	GO:0035967	cellular response to topologically incorrect protein	44	5	0.03334
	GO:0035966	response to topologically incorrect protein	47	5	0.04266
Drosophila	GO:0007141	male meiosis I	13	4	0.00150
	GO:0140543	positive regulation of piRNA transcription	3	3	6.9e-05
	GO:0010526	retrotransposon silencing	8	3	0.00331
	GO:0007130	synaptonemal complex assembly	10	3	0.00666
	GO:0030719	P granule organization	11	3	0.00888
	GO:0071218	cellular response to misfolded protein	12	3	0.01149
	GO:0051788	response to misfolded protein	12	3	0.01149
	GO:0007135	meiosis II	15	3	0.02169
	GO:0034508	centromere complex assembly	19	3	0.04094

1145

1146 **FIGURES**

1147

Fig. 1. Workflow for the bivalve dataset. Starting from a set of both genomes and

transcriptomes covering a great portion of bivalve taxonomic diversity, we first characterized the

entire complement of Dmrt, Sox, and Fox genes (upper row). We used sequence annotation and

phylogenetic tools to obtain reliable sequences and filter out any putative mis-assembled or mis-

annotated sequence. Afterwards, we built a reduced set of transcriptomes and genomes (the

reduced bivalve dataset, where we minimized the redundancy of congeneric species) from which

to draw the molecular evolution patterns of orthologous genes (bottom row). After having

obtained gene single-copy orthologous groups, we calculated the amino acid distances within each

orthogroup and then we built the distribution of median values. The same pipeline was also

utilized for the mammal and the fruit fly datasets, with just two minor differences: the starting

dataset was composed of only genomes, and the reduction step (R) was not necessary.

1159

1160 **Fig. 2. Dmrt, Sox, and Fox gene (DSFG) complement in bivalves and their outgroups.** The

1161 presence of genes in various species are indicated by filled circles. Numbers inside each circle

1162 specify genes with 2 or more copies. The shaded area highlights non-bivalve species, belonging

1163 either to other molluscs or to the references. The phylogenetic tree of analyzed species, as inferred

1164 from literature, is shown on the left, while major taxonomic groups are reported on the right.

Species represented by transcriptomic data are marked with an asterisk ('*'), and species not present in the reduced bivalve dataset are marked with two asterisks ('**'; see main text and **Fig. 1**); note that the two categories do nor overlap. DSFG trees are shown on the bottom (full trees can be found in **Supp. Fig. S1–S3**). Full species names, along with all assembly and taxonomic information, can be found in **Supp. Tab. S1**. Ad.: Adapedonta; Ar.: Arcida; Ca.: Cardiida; Ce.: Cephalopoda; L.: Lucinida; My.: Myida; Ref.: reference genes; S.: Sphaeriida.

Fig. 3. Distribution of amino acid sequence divergence (AASD) of single-copy orthogroups in

bivalves (A), including Dmrt, Sox, and Fox genes (DSFGs; B), and their correlations with

tip-to-tip distances (C), alignment lengths (D), and number of species (E). The distribution of

AASD has been computed on the median values of pairwise distances of over 11k SCOs from the

reduced bivalve dataset (see main text and **Fig. 1**). Genes have been divided according to their

median AASD value into three different groups, which are indicated by different colours and

increasing numbers (Groups 1, 2, and 3). Genes from Group 1 and Group 2 are collectively named

'highly divergent genes'. Circle heights of DSFGs show the median value of their AASD, while

the size indicates the number of represented species. DSFG trees are shown on the bottom (full

trees can be found in **Supp. Fig. S1–S3**). Darker points in C-E indicate Dmrt, Sox, and Fox gene

SCOs. The correlation between the amino acid distance and the tip-to-tip distance has been

computed on 200 randomly selected orthogroups. **Insets:** scheme of the sex-determination

working model in *Crassostrea gigas* as hypothesised by **Zhang et al. (2014)**, showing the main

genes involved. Green arrows indicate transcription activations, red arrows indicate transcription

suppressions.

1188

Fig. 4. Distribution of amino acid divergence (AASD) of single-copy orthogroups in Mammalia (A) and *Drosophila* (C), including Dmrt, Sox, and Fox genes (DSFGs; B-D). The distributions of AASD in mammals and fruit flies have been computed on the median values of pairwise distances of over 16k and 12k SCOs, respectively. Genes have been divided according to their median AASD value into three different groups, which are indicated by different colors and

increasing numbers (Groups 1, 2, and 3). Genes from Group 1 and Group 2 are collectively named

'highly divergent genes'. Circle heights of DSFGs show the median value of their AASD, while

the size indicates the number of represented species. DSFG trees are shown on the bottom (full

trees can be found in **Supp. Fig. S5–S7** for mammals and in **Supp. Fig. S8–S10** for fruit flies). In

Drosophila, *Sxl* and *tra*, both involved in the SD pathway (inset) do not belong to the group of

highly divergent genes (mean amino acid divergence of 0.0923793827106834 and

0.978190195613595, respectively; that is, within the boundaries of Group 3). **Insets:** scheme of

the sex-determination molecular pathways in *Mus musculus* and in *Drosophila melanogaster*,

showing the main genes involved (adapted from **Beukeboom & Perrim, 2014**). Green arrows

indicate transcription activations, red arrows indicate transcription suppressions. X: sex

chromosomes; A: autosomal chromosomes; $DSX^{M/F}$: *DSX* splicing variants present in males or

females, respectively.

SUPPLEMENTARY TABLES

Supp. Tab. S1. Genomic and transcriptomic data of bivalves and other molluscs. For each species, the relative ID, taxonomic information, BUSCO statistics, NCBI accession number, and source publication are reported. Biv: Bivalvia; Ca: Caenogastropoda; Cep: Cephalopoda; Co: Coleoidea; Gas: Gastropoda; Gen: Genome; He: Heterobranchia; Im: Imparidentia; Ne: Neomphaliones; Pa: Palaeoheterodonta; Pt: Pteriomorpha; Tra: Transcriptome; Ve: Vetigastropoda.

Supp. Tab. S2. Dmrt, Sox, and Fox gene (DSFG) family and domain identifiers (IDs) in PANTHER and CDD, respectively. After having retrieved putative DSFGs based on HMM profiles, IDs have been used to retain only reliable hits.

Supp. Tab. S3. List of DSFGs from reference species used to assess the identity of DSFGs in molluscs. NCBI accession numbers are reported in parenthesis. Each row represents an orthology group.

Supp. Tab. S4. Genomic data of mammals used to retrieve DSFGs and compute amino acid sequence divergence (AASD) of single copy orthogroups (SCOs). For each species, the relative ID, taxonomic information, BUSCO statistics, NCBI accession number, and source publication are reported.

Supp. Tab. S5. Genomic data of *Drosophila* used to retrieve DSFGs and compute AASD of SCOs. For each species, the relative ID, taxonomic information, BUSCO statistics, NCBI accession number, and source publication are reported.

1231 **Supp. Tab. S6. Complete set of DSFGs in bivalves.** For each gene, the species ID (Sp. ID) as in

1232 **Supp. Tab. S1**, the accession number (Gene ID), the Possvm-based annotation, and the CDD

1233 domains (including their Pssm-ID) are indicated.

1234

1235 **Supp. Tab. S7. Proportions of missing data in bivalve DSFGs.**

1236

1237 **Supp. Tab. S8. All the enriched GO terms for Group 1 and Group 2 genes of bivalves,**

1238 **mammals, and *Drosophila*.**

1239

1240 **Supp. Tab. S9. Complete set of DSFGs in mammals.** For each gene, the species ID (Sp. ID) as

1241 in **Supp. Tab. S4**, the accession number (Gene ID), and the Possvm-based annotation are

1242 indicated.

1243

1244 **Supp. Tab. S10. Complete set of DSFGs in *Drosophila*.** For each gene, the species ID (Sp. ID)

1245 as in **Supp. Tab. S5**, the accession number (Gene ID), and the Possvm-based annotation are

1246 indicated.

1247

1248 # SUPPLEMENTARY FIGURES

1249 **Supp. Fig. S1. ML phylogenetic tree of the Dmrt gene family in molluscs, including the**

1250 **Possvm orthology inference.** Reference genes from *H. sapiens*, *C. elegans*, and *D. melanogaster*

1251 are marked with an asterisk at the beginning of the tip. Species ID can be found in **Supp. Tab. S1**.

1252 Bootstrap values are shown for each node.

1253

1254 **Supp. Fig. S2. ML phylogenetic tree of the Sox gene family in molluscs, including the Possvm**

1255 **orthology inference.** Reference genes from *H. sapiens*, *C. elegans*, and *D. melanogaster* are

1256   marked with an asterisk at the beginning of the tip. Species ID can be found in **Supp. Tab. S1**.

1257   Bootstrap values are shown for each node.

1258

1259   **Supp. Fig. S3. ML phylogenetic tree of the Fox gene family in molluscs, including the**

1260   **Possvm orthology inference.** Reference genes from *H. sapiens*, *C. elegans*, and *D. melanogaster*

1261   are marked with an asterisk at the beginning of the tip. Species ID can be found in **Supp. Tab. S1**.

1262   Bootstrap values are shown for each node.

1263

Supp. Fig. S4. Dmrt, Sox, and Fox gene (DSFG) complement in Mammalia and *Drosophila*

spp. Presence/absence of genes in various species are indicated by filled circles. Numbers inside

each circle specify genes with 2 or more copies. The shaded area highlights outgroup species,

Gallus gallus (Aves) for mammals and *Anopheles gambiae* (Culicidae) for fruit flies. The

phylogenetic tree of analyzed species, as inferred from literature, is shown on the left, while major

taxonomic groups are reported on the right. All species are represented by genomic data. DSFG

trees are shown on the bottom (full trees can be found in **Supp. Fig. S5–S7**). Full species names

for both mammals and fruit flies, along with all assembly and taxonomic information, can be

found in **Supp. Tab. S4** and **Supp. Tab. S5**, respectively. A.: Aves; Chirop.: Chiroptera; L.:

Lagomorpha; M.: Monotremata; Me.: Metatheria; P.: Pholidota; Pe.: Perissodactyla; Prim.:

Primates; Roden.: Rodentia; X.: Xenarthra; C.: Culicidae.

1275

1276   **Supp. Fig. S5. ML phylogenetic tree of the Dmrt gene family in mammals, including the**

1277   **Possvm orthology inference.** Reference genes from *H. sapiens*, *Mus musculus*, *Elephas maximus*

1278   *indicus*, and *Ornithorhynchus anatinus* are marked with an asterisk at the beginning of the tip.

1279   Species ID can be found in **Supp. Tab. S4**. Bootstrap values are shown for each node.

1280

1281 **Supp. Fig. S6. ML phylogenetic tree of the Sox gene family in mammals, including the**

1282 **Possvm orthology inference.** Reference genes from *H. sapiens*, *Mus musculus*, *Elephas maximus*

1283 *indicus*, and *Ornithorhynchus anatinus* are marked with an asterisk at the beginning of the tip.

1284 Species ID can be found in **Supp. Tab. S4**. Bootstrap values are shown for each node.

1285

1286 **Supp. Fig. S7. ML phylogenetic tree of the Fox gene family in mammals, including the**

1287 **Possvm orthology inference.** Reference genes from *H. sapiens*, *Mus musculus*, *Elephas maximus*

1288 *indicus*, and *Ornithorhynchus anatinus* are marked with an asterisk at the beginning of the tip.

1289 Species ID can be found in **Supp. Tab. S4**. Bootstrap values are shown for each node.

1290

1291 **Supp. Fig. S8. ML phylogenetic tree of the Dmrt gene family in fruit flies, including the**

1292 **Possvm orthology inference.** Reference genes from *Drosophila melanogaster*, *Drosophila hydei*,

1293 *Drosophila pseudoobscura*, and *Drosophila suzukii* are marked with an asterisk at the beginning

1294 of the tip. Species ID can be found in **Supp. Tab. S5**. Bootstrap values are shown for each node.

1295

1296 **Supp. Fig. S9. ML phylogenetic tree of the Sox gene family in fruit flies, including the**

1297 **Possvm orthology inference.** Reference genes from *Drosophila melanogaster*, *Drosophila hydei*,

1298 *Drosophila pseudoobscura*, and *Drosophila suzukii* are marked with an asterisk at the beginning

1299 of the tip. Species ID can be found in **Supp. Tab. S5**. Bootstrap values are shown for each node.

1300

1301 **Supp. Fig. S10. ML phylogenetic tree of the Fox gene family in fruit flies, including the**

1302 **Possvm orthology inference.** Reference genes from *Drosophila melanogaster*, *Drosophila hydei*,

1303 *Drosophila pseudoobscura*, and *Drosophila suzukii* are marked with an asterisk at the beginning

1304 of the tip. Species ID can be found in **Supp. Tab. S5**. Bootstrap values are shown for each node.

1305

1306   **Supp. Fig. S11. ML phylogenetic tree of the Dmrt gene family in mollusc species.** Species ID

1307   can be found in **Supp. Tab. S1**. The tree has been midpoint rooted. Bootstrap values are shown

1308   for each node.

1309

1310   **Supp. Fig. S12. ML phylogenetic tree of *Sox-B1* and *Sox-B2* genes in mollusc and reference**

1311   **species**. Reference genes from *H. sapiens*, *C. elegans*, and *D. melanogaster* are marked with an

1312   asterisk at the beginning of the tip names. Species ID can be found in **Supp. Tab. S1**. Bootstrap

1313   values are shown for each node.

1314

1315   **Supp. Fig. S13. ML phylogenetic tree of *Fox-J2*, *Fox-M*, *Fox-O*, and *Fox-P* genes in mollusc**

1316   **and reference species.** Reference genes from *H. sapiens*, *C. elegans*, and *D. melanogaster* are

1317   marked with an asterisk at the beginning of the tip names. Species ID can be found in **Supp. Tab.**

1318   **S1**. Bootstrap values are shown for each node.

1319

1320   **Supp. Fig. S14. ML phylogenetic tree of the Fox gene family in bivalves and the sea urchin**

1321   ***Strongylocentrotus purpuratus* (Spur).** Reference genes from *S. purpuratus* are marked with an

1322   asterisk at the beginning of the tip names. Species ID can be found in **Supp. Tab. S1**. *S.*

1323   *purpuratus* genes are those given by **Tu et al., 2006**. Bootstrap values are shown for each node.

1324

Supp. Fig. S15. Distribution of amino acid sequence divergence (AASD) of single-copy

orthogroups in *Crassostrea gigas*, *Crassostrea angulata*, *Crassostrea ariakensis*, and

***Crassostrea virginica* (A), including DSFG (B)**. The distribution of AASD in *Crassostrea* has

been computed on the median values of pairwise distances of over 14 k single-copy orthogroups

(SCOs). Circle heights of DSFGs show the median value of their AASD. *Dmrt-1L* genes are

indicated as 'dmrt_disco_tree3'.