Exploring the evolution of sex determination related genes through a cross-species genomics analysis

Filippo Nicolini[1,2,3]*, Sergey Nuzhdin[4], Fabrizio Ghiselli[1], Andrea Luchetti[1], Liliana Milani[1]

[1]*Department of Biological, Geological and Environmental Sciences, University of Bologna, Bologna (BO), Italy.*

[2]*Fano Marine Center, Fano (PU), Italy.*

[3]*Department of Genetics and Genome Biology, University of Leicester, Leicester, UK.*

[4]*Department of Molecular and Computational Biology, University of Southern California, Los Angeles, CA, USA.*

*Author for correspondence: fn76@leicester.ac.uk

Keywords: bivalves; comparative genomics; Dmrt genes; *Drosophila*; Fox genes; mammals; molecular evolution; molluscs; Sox genes.

Abstract

The molecular basis of sex determination (SD), while being extensively studied in model organisms, remains poorly understood in many animal groups. Bivalves, a diverse class of molluscs with a variety of reproductive modes, represent an ideal yet challenging clade for investigating SD and the evolution of sexual systems. However, the absence of a comprehensive framework has limited progress in this field, particularly regarding the study of sex-determination related genes (SRGs). In this study, we performed a genome-wide sequence evolutionary analysis of the Dmrt, Sox, and Fox gene families in more than 40 bivalve species. For the first time, we provide an extensive and phylogenetic-aware dataset of these SRGs, and we find support to the hypothesis that *Dmrt-1L* and *Sox-H* may act as primary sex-determining genes, by showing their high levels of sequence diversity

within the bivalve genomic context. To validate our findings, we studied the same gene families in

two well-characterized systems, mammals and *Drosophila*. In the former, we found that the male

sex-determining gene *Sry* exhibits a pattern of amino acid sequence diversity similar to that of *Dmrt-*

1L and *Sox-H* in bivalves, consistent with its role of master SD regulator. In contrast, no such pattern

was observed among genes of the fruit fly SD cascade, which is controlled by a chromosomic

mechanism. Overall, our findings highlight similarities in the sequence evolution of some mammal

and bivalve SRGs, possibly driven by a comparable architecture of SD cascades. This work

underscores once again the importance of employing a comparative approach when investigating

understudied and non-model systems.

INTRODUCTION

In sexually reproducing organisms, the mechanism of sex determination (SD), i.e., the process by which the male or female identity of an organism (or gonadic tissue) is established, is highly diverse, ranging from strictly genetic systems to environmentally-dependent processes (**Haag & Doty, 2005**; **Uller & Helanterä, 2011**; **Bachtrog et al., 2014**; **Beukeboom & Perrin, 2014**). Characterising the molecular basis of SD is crucial for understanding not only the reproductive biology but also the evolutionary pressures shaping these systems (**Wilkins et al., 1995**; **Ellegren & Parsch, 2007**; **Grath & Parsch, 2016**; **Nicolini et al., 2023a**), as sex-determination related genes (SRGs; including primary sex-determining genes [SDGs]) are those responsible for the phenotypic differences between males and females, thanks to their sex-biased expression and interactions (**Ellegren & Parsch, 2007**; **Beukeboom & Perrin, 2014**; **Grath & Parsch, 2016**). One key aspect of SRGs is that they often exhibit accelerated rates of sequence evolution, due to their involvement in sex-related traits and reproduction. This represents the effect of sexual selection and/or adaptation, which act on sex-biased genes and produce highly divergent proteins at the interspecific level (**Civetta & Singh, 1998**; **Ellegren & Parsch, 2007**; **Meisel, 2011**; **Grath & Parsch, 2016**). Rapid sequence evolution is known for *Sex-determining Region Y* (*Sry*) of therians (**Pamilo & O'Neill, 1997**; **Mawaribuchi et al. 2012**), *Doublesex and mab-3 related gene W* (*Dm-W*) of the African clawed frog *Xenopus laevis*, and *Doublesex and mab-3 related gene Y* (*Dmy*) of the medaka fish *Oryzias latipes* (**Mawaribuchi et al. 2012**), all of which are master SDGs, that is, genes whose expression is primarily responsible for the establishment of the sex of the organism. Evolution under episodic diversifying selection has been detected also in *Drosophila* for genes involved in the SD cascade (e.g., *Sex-lethal* [*Sxl*], *transformer* [*tra*], and *doublesex* [*dsx*]), in correspondence with its establishment in the genus common ancestor (**Mullon et al., 2012**; **Baral et al., 2019**); though, rapid sequence evolution does not seem to concern extant amino acid sequences (**Haerty et al., 2007**; **Baral et al., 2019**), as they are globally evolving under purifying selection, especially in their

catalytic domain (**Mullon et al., 2012**; **Baral et al., 2019**). However, concerning the *dsx* genes,

higher rates of nucleotide and amino acid sequence evolution can be observed for male-specific

regions, if compared to female-specific regions and dimerization domains (**Baral et al., 2019**).

While SD has been extensively studied in model organisms, like mammals, insects, and

roundworms, comparatively little is known about the molecular mechanisms in non-model

organisms. A remarkable example is represented by bivalve molluscs, which exhibit a wide variety

of reproductive modes and sexual systems (**Breton et al., 2018**). Notwithstanding the considerable

importance in the human socio-economic landscape (reviewed in **Haszprunar & Wanninger,**

2012; **Gomes-dos-Santos et al., 2020**), the study of SD mechanisms in bivalves has been hampered

by the striking divergence among species (**Li et al., 2022**), and thus largely overlooked and limited

to few case studies (**Breton et al., 2018**; **Nicolini et al., 2023a**). So far, no master SDG has been

unambiguously identified, and the only working hypothesis on the functioning of the SD gene

regulatory network is available for the Pacific oyster *Magallana* (formerly *Crassostrea*) *gigas*

(**Zhang et al., 2014**). Nonetheless, the field still lacks both a robust functional investigation and an

evolutionary framework in which to place the current knowledge (**Nicolini et al., 2023a**). As a

matter of fact, major efforts have been dedicated to identify sex-biased genes through differential

gene expression (DGE) analyses (e.g., **Ghiselli et al., 2012**; **Milani et al., 2013**;

Teaniniuraitemoana et al., 2014; **Zhang et al., 2014**; **Capt et al., 2018**; **Afonso et al., 2019**), but

very few have leveraged cutting-edge techniques to investigate their actual role in SD and/or gonad

differentiation and development (e.g., **Liang et al., 2019**; **Sun et al., 2022**).

Components of the *doublesex* and *mab-3* related transcription factor (Dmrt), *Sry*-related HMG-box

(Sox), and forkhead box (Fox) gene families (henceforth referred to as DSFG) are notoriously

known as key actors in several developmental processes across Metazoa (**Benayoun et al., 2011**;

Matson & Zarkower, 2012; **Sarkar & Hochedlinger, 2013**; **Mawaribuchi et al., 2019**), including

SD in certain clades: the aforementioned *Dm-W*, *Dmy*, and *dsx* all belong to the Dmrt gene family,

while *Sry* belongs to the Sox gene family; *Fox-L2*, which takes part in most of the vertebrate SD

processes as a downstream effector of the female pathway, belongs to the Fox gene families.

Members of the DSFGs have been identified as putative SRGs also in bivalves, thanks to both DGE

analyses and *in-situ* hybridization (ISH; e.g., **Naimi et al., 2009**; **Li et al., 2018**; **Liang et al., 2019**;

Yue et al., 2021), suggesting that their role in morphological and sexual development is maintained

also in the clade. However, the clear role of DSFGs has yet to be elucidated, probably as a

consequence to the lack of (i) a systematic classification of the gene families and (ii) a

comprehensive understanding of their evolutionary history.

To overcome such limitations, this study aims to perform a thorough investigation of the DSFG

families in bivalves, with the attempt to provide a high-quality resource to be used as a reference

for future studies. Through the analysis of more than 40 annotated bivalve genomes and

transcriptomes, we aim (i) to describe the complete set and evolutionary history of DSFGs in

bivalves by means of phylogenetic inferences, manual curation, and orthology prediction; (ii) to

identify DSFGs potentially involved in bivalve SD by investigating their sequence evolution in a

genome-wide context. Our hypothesis is that, if any of the DSFGs is directly involved in SD (i.e.,

is a SDG), then we should expect it to show a higher rate of sequence evolution, as already found

in previous studies (**Pamilo & O'Neill, 1997**; **Mawaribuchi et al. 2012**) and discussed earlier; this

characteristic, in turn, would be reflected in a high diversity of the extant amino acid sequences

across the bivalve clade. To assess the robustness and reliability of our approach, we additionally

applied our pipeline to two non-bivalve datasets, composed of mammal and *Drosophila* species,

respectively (hereon referred to as the 'mammal dataset' and the 'fruit fly dataset'). By choosing

two clades for which SD is well characterised, we wanted to compare our results with those obtained

on taxa for which a deeper and more detailed knowledge is available. In particular, mammals and

Drosophila provide two different frameworks to study the patterns of molecular evolution in SDGs:

the former is a system where SD is completely genetic (i.e., the development into a male or into a

female is triggered by the up- or downregulation of *Sry* in undifferentiated gonads, respectively),

and for which a high rate of sequence evolution is already known for the male SDG *Sry*; the latter

is instead a system where SD is chromosomic, thus lacks a master SDG (the sexual fate of the

individual is determined by the ratio between autosomes and X chromosomes), and no high rate of

sequence evolution has been detected in extant amino acid sequences of SRGs (which are globally

evolving under purifying selection). In this sense, the mammal and the fruit fly datasets represent

opposing control datasets to be compared to bivalves, as it is expected that a higher rate of sequence

evolution concerns only master SDGs (like *Sry* in therians; i.e., the top regulatory part of the SD

cascade), but not the downstream genes (i.e., the bottom effectors). Therefore, we test our pipeline

on mammals and fruit flies with the following expectations: (i) in the mammalian dataset, *Sry* should

be detected as fast-evolving (**Pamilo & O'Neill, 1997**; **Mawaribuchi et al. 2012**); while (ii) in the

fruit fly dataset, no gene among those working within the sex-determining cascade should result as

evolving at a higher pace (**Haerty et al., 2007**; **Mullon et al., 2012**; **Baral et al., 2019**). In this way,

we are able to assess the reliability of results in bivalves, by comparing them to a null hypothesis.

This work offers novel insights into the evolutionary dynamics of SRGs and contributes a valuable

genomic resource for understanding SD in bivalves, one of the most ecologically and economically

important groups of marine organisms. In particular, here we provide the first extensive

phylogenetic-based classification of DSFGs in bivalves, covering many species from the major

bivalve orders, along with a comprehensive investigation of their sequence evolution.

RESULTS

Assembly of the bivalve, mammal, and fruit fly datasets

The bivalve dataset (hereafter referred to as the 'comprehensive bivalve dataset'; **Fig. 1**) on which

we retrieved and analysed DSFGs consists of 29 bivalve genomes, 14 bivalve transcriptomes, and

7 outgroup genomes (5 gastropods and 2 *Octopus* spp.; **Supp. Tab. S1**). Annotated genome

assemblies of bivalves were obtained from various publicly available resources, while reference

genome assemblies for gastropods and cephalopods were downloaded from NCBI (**Supp. Tab. S1**).

Transcriptomes of several additional bivalve species were also included to provide an extensive

identification of SRGs in underrepresented bivalve orders (mainly belonging to the Heterodonta

clade). BUSCO statistics for complete single-copy genes spanned from the 64.9% in *Modiolus*

modiolus to the 99.4% of *Perna viridis*, with a median value of 94.7%. We were able to get at least

one representative species for 11 different bivalve orders, covering a good proportion of the

phylogenetic diversity of the clades Pteriomorphia, Palaeoheterodonta, and Imparidentia, and thus

building the most extensive genomic and transcriptomic dataset for bivalve comparative analyses

so far (**Supp. Tab. S1**). Unfortunately, no genomes or transcriptomes for Protobranchia,

Archiheterodonta, and Anomalodesmata were available at the time of the project, thus we were not

able to include any of those clades in our analysis.

The mammal dataset consists of 32 species and 1 outgroup (*Gallus gallus*, Aves; **Supp. Tab. S2**),

and covers 12 major orders, while the fruit fly dataset consists of 17 species and 1 outgroup

(*Anopheles gambiae*, Culicidae; **Supp. Tab. S3**), and covers 2 *Drosophila* subgenera (i.e.,

Drosophila and *Sophophora*). BUSCO statistics for complete single-copy genes were generally

higher than those of bivalves, with a median of 98.3% for mammals and of 99.8% for fruit flies

(**Supp. Tab. S2–S3**).

Dmrt, Sox, and Fox complements in bivalves

Members of DSFG families were retrieved from the comprehensive bivalve dataset through the

HMM identification of their signature catalytic domains (the *dsx* and *mab-3* [DM] for Dmrt genes;

the high mobility group (HMG) box for Sox genes; and the forkhead domain for Fox genes).

Obtained hits were then annotated with the PANTHER sequence scoring and the Conserved Domain

Dataset (CDD), followed by a phylogeny-based orthology prediction including DSFGs from *Homo*

sapiens, *Drosophila melanogaster*, and *Caenorhabditis elegans* (hereafter referred to as 'reference

species'). Our pipeline managed to successfully identify and annotate DSFGs in bivalves, as proved

by the same analysis in mammals and fruit flies (see the paragraph **Dmrt, Sox, and Fox genes, and**

amino acid divergence in the testing datasets).

We retrieved four main orthology groups of Dmrt genes in bivalves (**Fig. 2**; **Supp. Fig. S1**; **Supp.**

Tab. S4), three corresponding to the groups present in the Bilateria common ancestor (*Dmrt-2*,

Dmrt-3, and *Dmrt-4/5*; **Mawaribuchi et al., 2019**), and one additional group with no clear ortholog

among reference genes, and thus putatively specific to molluscs (named *Dmrt-1L*, as per **Li et al.,**

2018b; **Evensen et al., 2022**). The majority of identified Dmrt genes are present in single-copy in

each species, but *Dmrt-4/5*s show a group-specific expansion in Palaeoheterodonta and

Heterodonta, while *Dmrt-1L* is completely absent from Heterodonta. The degree of missing data for

Dmrt genes in bivalves is ~35%, with *Dmrt-2* having the highest (~56%) and *Dmrt-4/5* the lowest

(~7%; **Supp. Tab. S5**). The ubiquitin-binding CUE-like DMA domain has been annotated in most

of the *Dmrt-3* and *Dmrt-4/5* genes, while an additional DM domain has been annotated in *Dmrt-1L*

genes in Mytilida and the gastropod *Pomacea canaliculata* (**Supp. Tab. S6**). Additionally, we

retrieved six main orthology groups of Sox genes, none of which is restricted to molluscs or bivalves

(**Fig. 2**; **Supp. Fig. S2**; **Supp. Tab. S4**). Five Sox groups (*Sox-B1/2*, *Sox-C*, *Sox-D*, *Sox-E*, and *Sox-*

F) are those traditionally considered to be present in the Bilateria common ancestor (**Phochanukul**

& Russell, 2010), while one has been identified outside mammals only recently (*Sox-H*, or *Sox-30*;

Han et al., 2010). *Sox-B2* and *Sox-B1* have been grouped in the same clade, as in our phylogenetic

reconstruction *Sox-B2* is paraphyletic with respect to *Sox-B1* (**Supp. Fig. S2**), despite being

traditionally recognised as a separate paralogy group in humans, fruit flies, and roundworms. The

degree of missing data for Sox genes in bivalves is ~8%, with *Sox-H* having the highest (~21%) and

Sox-B1/2 and *Sox-C* both having no missing genes (**Supp. Tab. S5**). The Sox N-terminal signature

domain was annotated for *Sox-E* genes (**Supp. Tab. S4**). Concerning Fox genes, we retrieved 27

main orthology groups (**Fig. 2**; **Supp. Fig. S3**; **Supp. Tab. S4**), two of which are specific to molluscs

(*Fox-OG13/NA*, *Fox-OG16/NA*). Additionally, other potential mollusc-specific Fox groups have

been identified, but these have been excluded from the final orthology analysis as they are present

in less than half of bivalve species (see **Materials and Methods**; **Supp. Tab. S4**). The two major

Fox gene subgroups, Group I (monophyletic, specific to Metazoa; includes *Fox-A*, *Fox-B*, *Fox-C*,

Fox-D, *Fox-E*, *Fox-F*, *Fox-G*, *Fox-H*, *Fox-L1*, *Fox-L2*, *Fox-Q2*) and Group II (paraphyletic, specific

to Opisthokonta; includes *Fox-O*, *Fox-P*, *Fox-J2*, *Fox-J1*, *Fox-K*, *Fox-N2/3*, *Fox-N1/4*; **Larroux et**

al., 2008), have been recovered, including the four Fox genes that were present in the Bilateria

common ancestor (*Fox-C*, *Fox-F*, *Fox-L1*, and *Fox-Q1*; **Shimeld et al., 2010**). Two putative

lineage-specific expansions have been recovered for *Fox-OG28/NA*, one regarding *Mytilus* spp. and

one regarding the two Myida species (**Fig. 2**; **Supp. Fig. S3**). The degree of missing data for Fox

genes in bivalves is ~22%, with *Fox-H* having the highest (~42%) and *Fox-J1* having no missing

genes (**Supp. Tab. S5**). The forkhead-associated (FHA) domain was annotated for *Fox-K* genes, the

Fox-P coiled-coil signature domain was annotated for *Fox-P* genes, while both the forkhead N- and

C-terminal signature domains were annotated for *Fox-A* genes (**Supp. Tab. S4**).

Amino acid sequence divergence of Dmrt, Sox, and Fox genes in bivalves

To test whether DSFGs potentially involved in bivalve SD have a high rate of sequence evolution

within a genome-wide context, we used the amino acid sequence divergence (AASD) on single-

copy orthogroups (SCOs) as a metrics of sequence diversity. This metric is pretty fast and

straightforward to obtain and allow to save computational time, as it only requires the amino acid

alignment and the corresponding best-fit substitution mode, while preserving enough evolutionary

information (see below). To further save computational time and also prevent over-represented

bivalve clades (such as Ostreida and Mytilia) to bias the analysis, SCOs were obtained after a

reduced dataset comprised of 34 bivalve species (hereafter referred to as the 'reduced bivalve

dataset'; **Fig. 1**; **Supp. Tab. S1**) which included, for each bivalve genus, only the best genomes and

transcriptomes in terms of either BUSCO scores or assembly statistics (**Supp. Tab. S1**). *Archivesica*

marissinica and *Saccostrea glomerata* were removed, as their annotated coding sequences contain

many stop codons, which prevent accurate amino acid guided alignments. We produced amino acid

alignments of ~11k SCOs (32 of which belongs to the DSFG families), calculated the pairwise

AASD within each of them, and then built the distribution of the median values (**Fig. 1**). Afterwards,

genes were categorised into three groups: Group 1, consisting of genes from the 1% upper quantile

of the distribution; Group 2, consisting of genes between the 1% and 5% upper quantiles; and Group

3, consisting of all the remaining genes. Group 1 and Group 2 genes will be referred to as 'highly

divergent genes'. In particular, 112 genes were assigned to Group 1 (1% upper quantile), 447 to

Group 2 (5% upper quantile), and 10,603 to Group 3. Most of the DSFGs (29/32) fell in Group 3

(**Fig. 3**), which means they have a median AASD comparable to the vast majority of other genes in

bivalves (median level of the genomes). Just *Dmrt-1L*, *Sox-H*, and *Sox-F* showed higher

divergences, and have been accordingly placed in Group 2. Overall, pairwise AASD proved to be a

good approximation of the tip-to-tip distances calculated on the corresponding ML phylogenetic

trees ($R = 0.84$, $p < 2.2E{-}16$, considering 200 randomly-selected trees; **Fig. 3C**), and it also shows

no influence from the alignment length ($R = 0.11$) or the number of represented species ($R = -0.23$;

Fig. 3D–E). Genes from Group 1 and Group 2 are strongly involved in cellular regulatory processes

(such as those related to the metabolism of nucleic acids, proteins, and other macromolecules), but

also in development and response to external stimuli, as shown by the GO-term enrichment analysis

(**Tab. 1**; **Supp. Tab. S6**).

Dmrt, Sox, and Fox genes, and amino acid sequence divergence in the test datasets

To validate our approach for the study of bivalve SRG molecular evolution, we ran the same analysis

as before on two additional datasets, consisting of reference genomes of mammals and *Drosophila*

species (**Supp. Tab. S2–S3**, respectively), whose sex-determining mechanisms are well studied and

characterised. As a matter of fact, despite it is well known that SDGs tend to evolve faster than

genes not involved in SD, the hypothesis has never been tested extensively across the entire

phylogenetic diversity of a group: molecular evolution of SDGs and SRGs has mainly been tested

on single species or inside the boundaries of taxonomic genera (**Stothard et al., 2003; Haerty et**

al., 2007; Mank et al., 2007; Mullon et al., 2012; Papa et al., 2017; Ghiselli e al., 2018).

Overall, the DSFG datasets retrieved in mammals and fruit flies are far more complete than those

in bivalves, and most of the already-recognised orthology groups have been identified. In mammals,

we retrieved 7 Dmrt orthology groups with ~3.1% of missing data, 20 Sox orthology groups with

~8.1% of missing data, and 42 Fox orthology groups with ~4.6% of missing data (**Supp. Fig. S4A,**

S5–S7; **Supp. Tab. S7**). Of these, just *Sox-5* was not included in the subsequent AASD analysis, as

it did not meet the 50%-species occupancy threshold. OrthoFinder analysed ~650M genes, and the

number of SCOs used in the AASD analysis (thus resulting from the DISCO-based orthogroup

decomposition pipeline) is >16k (**Fig. 4A**). From the distribution of median AASD, 163 genes were

assigned to Group 1, 649 to Group 2, and 15,355 to Group 3. Most of the DSFGs (66/68) fell in

Group 3 (**Fig. 4B**), while *Sry* and *Fox-D4* showed higher divergences, and have been accordingly

placed in Group 1 and 2, respectively. Genes from Group 1 and Group 2 show a strong enrichment

in immune-related functions (such as innate and adaptive immune response, defence response to

bacteria and viruses, lymphocyte metabolism, etc.), but also in reproductive processes (such as

spermatogenesis; **Tab. 1**; **Supp. Tab. S6**). Concerning *Drosophila*, we retrieved 4 Dmrt orthology

groups with ~1.7% of missing data, 7 *Sox* orthology groups with ~3.9% of missing data, and 17 Fox

genes with ~8.3% of missing data (**Supp. Fig. S4B, S9–S10**; **Supp. Tab. S8**). OrthoFinder analysed

~240M, and the distribution of median AASD was built after >12k SCOs (**Fig. 4C**). 126 genes were

assigned to Group 1, 501 to Group 2, and 11,880 to Group 3. All the DSFGs have been used in the

AASD analysis, but none of them have been placed in Group 1 or 2, that is, all the DSFGs in

Drosophila have an AASD comparable to the median level of the genome (**Fig. 4D**). Genes of

Group 1 and Group 2 show a GO-term enrichment in meiotic processes, such as

chromosome/chromatid organisation, and retrotransposon silencing (**Tab. 1**; **Supp. Tab. S6**).

DISCUSSION

A new manually-curated and phylogenetic-based reference dataset of Dmrt, Sox, and Fox genes in bivalves

The annotation and characterisation process of a gene family may harbour many overlooked challenges in a certain clade of organisms (**Vizueta et al., 2020**). For example, the presence of highly conserved catalytic domains may hamper the correct identification of the components of a gene family because of insufficient phylogenetic signal, as it is the case for Hox and ParaHox genes and their homeobox motif (**Baldwin-Brown et al., 2018**; **Nicolini et al., 2023b**). Conversely, the components of dynamic gene families characterised by abrupt and sequential duplication events may be difficult to sort into separate groups. As a matter of fact, varying levels of sequence heterogeneity and gene copy numbers makes the inference of orthologous groups hard, as for certain clans of the P450 family (**Dermauw et al., 2020**). Regardless of the causes, having a solid and wide phylogenetic context, in which to study gene duplications/losses and orthology relationships, is crucial to overcome these difficulties. In the same way, manual curation and visual inspection of multiple sequence alignments, phylogenetic trees, and gene structures (in terms of domain composition, start and stop codons, and other feature representations) is helpful, despite being time-demanding and possibly less reproducible. In this study, we characterised the full complement of DSFGs in the vast class of bivalves, by leveraging sequence domain annotation, phylogenetics, and manual curation of the dataset. Our aim was to obtain the most reliable gene complements as possible, combined with a vast taxonomic dataset, a solid phylogenetic inference, an openly available dataset of gene sequences, and a reproducible pipeline for the annotation of gene identity. By doing so, we want to provide a reliable resource for future studies of DSFGs, either focused on bivalves or on Metazoa in general.

Concerning the Dmrt gene family, we identified orthologs of the vertebrate *Dmrt-2*, *Dmrt-3*, and *Dmrt-4/5* (*A1/A2;* **Fig. 2**; **Supp. Fig. S1**; **Supp. Tab. S4**), which are also expected to have been

present in the Bilateria common ancestor (**Mawaribuchi et al., 2019**). **Wang et al. (2023)** found that *Dmrt-4/5* is duplicated in *Mercenaria mercenaria* and *Cyclina sinensis* (Venerida), and in *Dreissena polymorpha* (Myida), and we confirm this result by tracing back the duplication event to the split between Palaeoheterdonta (here represented by Unionida) and Heterodonta (here represented by Venerida, Myida, Sphaeriida, Adapedonta, Cardiida, and Lucinida; **Fig. 2**). Furthermore, we confirm *Dmrt-1L* to be present in many bivalve species (mainly belonging to the Ostreida, Pectinida, Mytilida, and Unionida orders; **Fig. 2**), as well as in gastropods and *Octopus*. Though, our phylogenetic analysis did not retrieve any unambiguous orthology relationship among *Dmrt-1L* and either vertebrate *Dmrt-1* or *Drosophila dsx* genes, as instead it was proposed in previous works (**Li et al., 2018b**; **Evensen et al., 2022**). As a matter of fact, the amino acid sequence of the *Dmrt-1L* DM domain does not recall that of any other Dmrt gene. Furthermore, it must be considered that various phylogenetic analyses have recovered both *Dmrt-1* and *dsx* genes to be restricted to vertebrates and arthropods, respectively (**Wexler et al., 2014**; **Mawaribuchi et al., 2019**; **Panara et al., 2019**), that is, they do not have any direct ortholog outside their relative clades. Thus, if *Dmrt-1L*, *dsx*, and *Dmrt-1* were true orthologs, their origin would need to be placed at least in the Bilateria common ancestor, which seems however to be not the case. All considered, we thus confirm that *Dmrt-1L* is not orthologous to *Dmrt-1* and *dsx* and is rather a mollusc-specific gene (**Evensen et al., 2022**). The monophyly of the *Dmrt-1L* group is not supported by the phylogenetic tree inferred with Dmrt genes from molluscs and the reference species (**Supp. Fig. S1**); though, it is recovered when analysing just genes from mollusc species (**Supp Fig. S11**). To this regard, we speculate that in our analysis, the difficulty in obtaining the monophyly of *Dmrt-1L* genes may have arisen primarily because of the many *C. elegans*-restricted genes (**Supp. Tab. S10**), which are placed among bivalve genes (**Supp. Fig. S1**), but also because of the high AASD of *Dmrt-1L* genes (see **High amino acid sequence divergence identifies putative sex-determining genes**), which hampers a straight-forward phylogenetic reconstruction. Furthermore, our broad-context analysis allowed us to identify some cases of incorrect gene identification in bivalves, which have arisen

because of erroneous or ambiguous annotations in previous works, as a result of limited datasets or analyses. For example, (i) the scallop-specific cluster of Dmrt genes retrieved by **Wang et al. (2023)** rather belongs to the *Dmrt-1L* group, and (ii) the classification of Dmrt genes in *Crassostrea* species provided by **Zeng et al. (2024)** needs to be revised following as follows: *Dmrt-1* genes are *Dmrt-4/5*; *Dmrt-2* genes are *Dmrt-3*; *Dmrt-3* genes are *Dmrt-1L*; hence, *Crassostrea* species do not have *Dmrt-2* genes.

For what concerns the Sox gene family, bivalves (or molluscs) do not show any major clade-restricted gene, as only the five Bilateria-specific Sox groups (*Sox-B1/2*, *Sox-C*, *Sox-D*, *Sox-E*, and *Sox-F*) and *Sox-H* have been identified (**Fig. 2**; **Supp. Fig. S2**; **Supp. Tab. S4**), in accordance with previous findings (**Yu et al., 2017b**; **Evensen et al., 2022**; **Wang & Nie , 2024**). *Sox-B1/2* is clearly made up of two subgroups (i.e., *Sox-B1* and *Sox-B2*), as expected, but their respective identity could not be unambiguously established, as *Sox-B1/2* genes of reference species do not form separate clusters (**Supp. Fig. S2**), even when inferring the phylogenetic tree only of components of the *Sox-B1/2* group from molluscs and reference species (**Supp. Fig. S12**).

Compared to Dmrt and Sox genes, the Fox gene family appears as the most dynamic in terms of gene presence/absence, as already shown by other works (**Wu et al., 2020**; **Schomburg et al., 2022**; **Seudre et al., 2022**). Our phylogenetic analysis successfully recovered Group I and Group II of Fox genes (**Larroux et al., 2008**), which include the four Fox genes that were present in the Bilateria common ancestor (*Fox-C*, *Fox-F*, *Fox-L1*, and *Fox-Q1*; **Fig. 2**; **Supp. Fig. S3**; **Supp. Tab. S4**; **Shimeld et al., 2010**). To our knowledge, this is the first broad-taxonomic identification and classification of Fox genes in bivalves, as up to now they have been systematically characterised only in *C. gigas* (**Yang et al., 2014**), *Patinopecten yessoensis* (**Wu et al., 2020**), and *Ruditapes philippinarum* (**Liu et al., 2024**). Firstly, our analysis confirms the absence in molluscs of *Fox-I*, *Fox-Q1*, *Fox-R*, *Fox-S* (**Supp. Fig. S3**), which are in fact thought to have emerged with the diversification of deuterostomes or vertebrates (**Yang et al., 2014**; **Wu et al., 2020**; **Schomburg et al., 2022**; **Seudre et al., 2022**). Furthermore, we have found many Fox groups that appeared as

mollusc-specific and/or still-unnamed at a first analysis. However, a more in-depth investigation revealed a different scenario. *Fox-OG2/NA* appears close to the human *Fox-M* gene in the phylogenetic tree, but they do not form a monophyletic group (**Supp. Fig. S3**). However, by comparing *Fox-OG2/NA* sequences and phylogenetic tree with those analysed by **Yang et al. (2014)**, **Wu et al., (2020)**, **Schomburg et al. (2022)**, **Seudre et al. (2022)**, it appears clear that this group of Fox genes is indeed *Fox-M*. However, our analysis has failed to retrieve a monophyletic relationship among bivalve and human *Fox-M* genes, even when inferring a tree with just *Fox-J2*, *Fox-M*, *Fox-O*, and *Fox-P* complements (**Supp. Fig. S13**), which belong to the same Fox group. Regarding the *Fox-OG39/NA* group, it does not have any homolog in reference species (**Supp. Fig. S3**) but is found to belong to the *Fox-AB* group by sequence comparison with previous works (**Yang et al., 2014**; **Wu et al., 2020**; **Seudre et al., 2022**). *Fox-AB* was formerly described only in the sea urchin *Strongylocentrotus purpuratus* and the lancelet *Branchiostoma floridae* (**Tu et al., 2006**; **Yu et al., 2008**), but was later identified also in several Spiralia lineages, including molluscs (e.g., **Yang et al., 2014**; **Wu et al., 2020**; **Seudre et al., 2022**). A similar situation concerns *Fox-OG15/NA* and *Fox-OG28/NA*, which again could not be named based on orthology relationships with the reference species genes (**Supp. Fig. S3**), but actually represent two lineage-specific expansions of the *Fox-Q2* group (named *Fox-Q2b* and *Fox-Q2c*), as already appointed in previous studies (**Yang et al., 2014**; **Wu et al., 2020**). This observation fits within the wider context of the *Fox-Q2* group expansion in Bilateria and, particularly, in Spiralia, that led to remarkable differences in their gene copy numbers across various clades (**Seudre et al., 2022**). Two additional Fox genes have been previously identified in bivalves and were named *Fox-Y* and *Fox-Z* (**Yang et al., 2014**; **Wu et al., 2020**). In our analysis, these Fox genes were identified as *Fox-OG13/NA* and *Fox-OG16/NA*, after sequence comparison with Fox genes from *C. gigas* and *P. yessoensis*. On one hand, *Fox-Y* was firstly identified in *S. purpuratus* (**Tu et al., 2006**) and only recently in a few bivalve species (**Yang et al., 2014**; **Wu et al., 2020**). However, when analysing bivalve and *S. purpuratus* Fox genes, we failed in retrieving such a clear orthology relationship, as *S. purpuratus Fox-Y* does not fall within

the phylogenetic range of bivalve *Fox-OG13/NA*, which contains the supposed *Fox-Y* orthologs

(**Supp. Fig. S14**). Also, the forkhead domains of *Fox-OG13/NA* genes were annotated as 'forkhead

domain P' (**Supp. Tab. S4**). On the other hand, *Fox-Z* was firstly identified in bivalves and in

several other protostomes, thanks to a phylogenetic work including the brachiopod *Lingula unguis*,

the annelid *Capitella teleta*, the scorpion *Centruroides sculpturatus*, and the centipede *Strigamia*

maritima (**Wu et al., 2020**). However, later works have not recovered this Fox gene, even when

analysing annelids (**Seudre et al., 2022**) and panarthropods (**Schomburg et al., 2022**) in a more

focused effort. In this case, the forkhead domains were annotated as either a generic 'forkhead

domain' or a 'forkhead domain Q2' (**Supp. Tab. S4**). All considered, we argue that bivalves possess

two additional Fox groups (here *Fox-OG13/NA* and *Fox-OG16/NA*; **Fig. 2**; **Supp. Fig. S3**; **Supp.**

Tab. S4) which are shared with other mollusc species, as revealed also by other authors. However,

given the discordant results of the phylogenetic hypothesis and domain annotation, we think that a

more thorough investigation on their orthology relationships with Fox genes from other Metazoa is

needed, and thus we chose to not employ their former names *Fox-Y* and *Fox-Z*.

Besides the DSFG groups discussed so far, it must be also considered that many orphan genes have

been identified (**Supp. Fig. S1–S3**; **Supp. Tab. S4**). For example, **Wu et al. (2020)** identified a

duplication event of *Fox-H* genes in *C. gigas*, which has been recovered also in our analysis for the

entire Ostreida clade (*Fox-OG36/NA*; **Supp. Fig. S3**). Similarly, a gene orthology group putatively

specific to Pteriomorphia has been identified among Sox genes (*Sox-OG1/NA*). Of course, these

genes deserve as much attention as their widely distributed paralogs, as they may constitute true

group-specific expansions and may play fundamental roles in some biological processes. However,

they have not been discussed here or included in **Fig. 2** for clarity purposes, but their sequences are

available in on GitHub (see the **Data Availability Statement**).

Overall, our analysis clearly shows the importance of adopting a wide-angle approach when

characterising the members of a gene family, especially for large ones such as the Fox genes

(**Schomburg et al., 2022**). As a matter of fact, the presence of duplication events and orphan genes

needs to be addressed with a broad taxonomic dataset, in order to account for possible mis-

annotations, gene phylogenetic misplacements, and sequence heterogeneity. Additionally, many

reference species need to be included for the gene identification process, to consider distantly related

genes and obtain a solid annotation. Our gene annotation pipeline also resulted to be very solid,

even with non-model organisms and sub-optimal genomic and transcriptomic resources as those of

bivalves. As a matter of fact, by running the same pipeline on two additional datasets composed of

mammal and fruit fly genomes, we were able to obtain high-quality orthology groups in accordance

with previous knowledge on the clades (**Supp. Fig. S4-S10**; **Supp. Tab. S7–S8**), with little or no

need of manual curation. Furthermore, the present work represents the first broad analysis of DSFGs

in both mammals and fruit flies, as so far attention has been mainly dedicated to single well-studied

organisms or little clades (e.g., **Jackson et al., 2010**).

High amino acid sequence divergence identifies putative sex-determining genes

Sex-biased genes tend to evolve more rapidly than unbiased genes at the level of their protein

sequences. Accelerated rates have been observed in both male-biased genes (reviewed in **Parsch**

and Ellegren 2013; **Grath and Parsch 2016**) and female-biased genes (e.g., **Papa et al. 2017**;

Ghiselli et al. 2018), but also in SRGs and SDGs (**O'Neil & Belote, 1992**; **Whitfield et al., 1993**;

de Bono & Hodgkin, 1996). For example, it has been shown that *Dm-W*, *Dmy*, and *Sry* (which are

SDGs in the African clawed frog *X. laevis*, in the medaka fish *O. latipes*, and in eutherians,

respectively) all have higher substitution rates than their paralogues (*Dmrt-1* for *Dm-W* and *Dmy*,

Sox-3 for *Sry*), particularly when considering their DNA-binding domains (**Mawaribuchi et al.**

2012). Similarly, both a burst of positive selection and a relaxation of purifying selection has been

detected in *Drosophila Sxl* in correspondence with its recruitment at the top of the sex-determining

cascade. The same signs of relaxed purifying selection have been found in the downstream targets

of *Sxl*, that is, *transformer* (*tra*) and *dsx*, despite no evidence of positive selection has been detected

(**Mullon et al., 2012**).

Considering these shared features of SRGs and SDGs, we decided to look for signs of accelerated

sequence evolution in DSFGs of bivalves to evaluate if any of them could be *a priori* associated

with SD by employing the tools of molecular evolution. However, we analysed patterns of sequence

evolution not only among putative SRGs and their close paralogs, but also considering the genomic

context in which these genes evolve. Our aim was to check whether SRGs show higher rates of

sequence evolution when compared to other genes not involved in SD and not belonging to the same

gene family. To do so, we obtained the AASD median values of more than 11k SCOs from bivalve

genomes (**Fig. 3A**), in order to build a statistical distribution to be used as a reference: if SRGs/SDGs

(in this case, DSFGs) truly evolve faster than other genes, we may expect them to fall within the 5%

(or even 1%) upper quantile of the distribution (**Fig. 3B**), i.e., within highly divergent genes (Group

1 and Group 2 genes of the distribution). We chose to use the AASD as a metric of sequence

evolution (instead of the tip-to-tip distances of phylogenetic trees, which account for more

comprehensive evolutionary models) to save computational time. As a matter of fact, the AASD

median values proved to be a good approximation of the tip-to-tip median distances in 200 randomly

selected genes (**Fig. 3C**; R = 0.84, p < 2.2E−6).

Among DSFGs, three fell within the 5% upper quantile, namely *Dmrt-1L*, *Sox-H*, and *Sox-F*.

Interestingly, on the basis of DGE analyses, *Dmrt-1L* and *Sox-H* have been already proposed to be

involved in the male SD pathway of *C. gigas* (inset in **Fig. 3B**; **Zhang et al., 2014**), while *Sox-H* to

be also a candidate gene involved in the development of male germ cells in the Manila clam

Ruditapes philippinarum (**Ghiselli et al., 2012**). Specifically, *Sox-H* would play a major role in *C.*

gigas SD, by interacting with *Dmrt-1L* and determining the onset of the male phenotype

development; at the same time, both *Sox-H* and *Dmrt-1L* would inhibit *Fox-L2*, which instead is

necessary to start the female phenotype development. *Dmrt-1L* and *Sox-H* have been appointed

several other times to be involved in male-gonad development and differentiation, through DGE

(e.g., **Teaniniuraitemoana et al., 2014**; **Capt et al, 2018**; **Afonso et al., 2019**), ISH (e.g., **Naimi**

et al., 2009; **Li et al., 2018**; **Liang et al., 2019**; **Yue et al., 2021**), and RNA interference (**Liang et**

al., 2019; **Sun et al., 2022**). Therefore, the high AASD of *Dmrt-1L* and *Sox-H* is coherent with

previous works, strengthening their role as putative SRGs.

The relationship between high gene AASD and the involvement in SD is particularly strengthened

when looking at the patterns of AASD in the test datasets, which corroborates the solidity of our

analysis: (i) in the mammal dataset—which represents a strictly genetic SD system with a master

and rapidly-evolving SDG—one of the genes from the 5% upper quantile of the distribution is *Sry*

(**Fig. 4A–B**), the male sex-determining gene in eutherians (inset in **Fig. 4B**); (ii) in the fruit fly

dataset—which represents a chromosomic SD system and with no differences in the rates of

sequence evolution among SRGs—none of the DSFG exhibit significantly higher AASD (**Fig. 4C–**

D), including the downstream effector *dsx* (inset in **Fig. 4D**). Also, *Sxl* and *tra*, both involved in the

SD pathway of *Drosophila* (inset in **Fig. 4D**) do not belong to the group of highly divergent genes,

as they have a mean amino acid divergence of about 0.09 and 0.9, respectively (Group 3; **Fig. 4D**).

Therefore, it can be argued that both *Dmrt-1L* and *Sox-H* may not only be SRGs, but may participate

in bivalve SD as primary SDGs, which is reflected in their high AASD, as it is observed for *Sry* in

mammals. As a matter of fact, if they were involved in SD just as intermediate actors of the

signalling cascade, we should have not observed a high AASD, as *Drosophila Sxl*, *tra*, and *dsx* seem

to suggest. Overall, these patterns of molecular evolution concerning SRGs and SDGs are also

supported by the way SD regulatory networks evolve. It has been proposed that the sex-determining

cascades tend to arise and be established with a bottom-up mechanism (**Wilkins et al., 1995**;

Mullon et al., 2012; **Beukeboom & Perrim, 2014**; **Capel, 2017**). This means that the regulatory

relationships among genes at the bottom of the cascade are settled prior to the regulatory

relationships among genes at the top and, consequently, upstream regulators are progressively

recruited to fine-tune diverse SD signals. These evolutionary patterns eventually produce gene

regulatory networks in which the divergence of the upstream triggers is higher than that of

downstream effectors, in terms of both identity and sequence composition (**Beukeboom & Perrim,**

2014). This mechanism has been proposed for *Drosophila* species (**Mullon et al., 2012**), *C. elegans*

(**Stothard & Pilgrim, 2003**), and vertebrates, despite in the latter case it has been questioned several

times (reviewed in **Capel, 2017**).

Two main objections can be moved against our approach: (1) the distribution of AASD is not

appropriate for this kind of inference, as it does not represent the true gene evolutionary (or

substitution) rates (which instead are those usually utilized when dealing with SRGs and SDGs); (2)

the three datasets are not comparable one to each other, as they take into consideration very different

animal groups, with different taxonomic rankings and different divergence times (thus, the patterns

of AASD may be the products of other confounding factors not directly related to SD). Concerning

the first objection, we are aware that the AASD does not represent the evolutionary rate itself, but

rather its product. However, the two features are tightly linked, as on the long term highly divergent

proteins tend to be produced by genes with high evolutionary (or substitution) rates (**Echave et al.,**

2016). By performing a GO term enrichment, it emerged that highly divergent genes of the mammal

dataset are mainly involved in the immune response and male spermatogenesis (**Tab. 1**; **Supp. Tab.**

S10), which are two processes notoriously connected with rapid sequence evolution (i.e., higher

evolutionary rates; **Swanson & Vacquier, 2002**; **Murat et al., 2023**; **Vinkler et al., 2023**).

Similarly, highly divergent genes from the fruit fly dataset show an enrichment for GO terms

associated with meiotic-related functions (such as the formation of the synaptonemal complex by

the products of *c(2)M*, *c(3)G*, *corona*, and *corolla* genes; **Tab. 1**; **Supp. Tab. S10**), which again are

known to evolve rapidly (**Hemmer et al., 2016**). In other words, the test datasets—which include

well-studied and characterised model systems—allow us to directly link the high AASD (as

computed in this work) with high rates of sequence evolution (as found in previous works). This

consideration can thus be extended also to the bivalve dataset: highly divergent genes in terms of

AASD, which include some DSFGs and show an enrichment for GO-terms associated to

macromolecule metabolism and morphological development (**Tab. 1**; **Supp. Tab. S10**), are also

genes with accelerated substitution rates (**Ghiselli et al., 2018**; **Iannello et al., 2023**).

Concerning the second objection, we chose two test datasets with different characteristics as we

wanted to check the extent of our hypothesis, that is, molecular evolution can be used to look for

putative primary SDGs in taxonomic-wide analyses. The difference in divergence times and

taxonomy ranks for bivalves and therians (Late Cambrian [about 498 million years ago, Mya; **Song**

et al., 2023] and Early Mesozoic [166–123 Mya; **Álvarez-Carretero et al., 2022**], respectively) do

not seem to influence the sequence diversity of SRGs, as both *Dmrt-1L*/*Sox-H* for bivalves and *Sry*

for mammals exhibit high AASD with respect to their own distributions, regardless of their age.

Dmrt-1L and *Sox-H* (which are mollusc- and Bilateria-specific, respectively) are undoubtedly older

than *Sry,* which emerged in the Theria common ancestor (**Foster et al., 1992**), but each one of them

can be considered a highly divergent gene in bivalves and mammals, respectively (i.e., genes that

are included in the 5% upper quantile of bivalve and mammal AASD distributions). Conversely, the

difference in divergence times and taxonomic ranks for *Drosophila* (Paleocene/Eocene boundary

[about 56 Mya; **Russo et al., 2013**]) may seem to be influencing the results for the dataset, resulting

in a false negative. In other words, it can be argued that: (i) the genes included in the SD cascade of

Drosophila (such as *Sxl, tra*, and *dsx*; inset in **Fig. 4D**) may have a high AASD which, however,

has not been detected by our methodological approach (for example, this may be traced back to the

young diversification age of *Drosophila* species if compared to bivalves); (ii) the species included

in the analysis are congeneric, thus the sequence differentiation of SRGs may exist not at the amino

acid level but at the nucleotide one. To better disentangle this issue and further discuss the fruit fly

dataset, we repeated the analysis of the AASD only on species of the *Crassostrea* genus (*C. gigas*,

Crassostrea angulata, *Crassostrea ariakensis*, and *Crassostrea virginica*), which are much younger

(Middle Cretaceous [less than 100 Mya; **Qi et al., 2023**]), thus comparable to *Drosophila*. Results

showed that, even when analysing a smaller bivalve dataset, encompassing only 4 species of recent

origin, the high AASD of *Dmrt-1L* persists, that is, *Dmrt-1L* is still grouped together with highly

divergent genes (**Supp. Fig. S15**). The same has not been recovered for *Sox-H*, which fell in genes

from Group 3 (the group corresponding to the 95% interval of the AASD distribution; see **Materials**

and Methods) but still have the second highest AASD median value among DSFGs (**Supp. Fig.**

S15).

Of course, we should not expect that highly divergent genes are only those involved in SD but may

also participate in other processes (as discussed earlier and shown by GO term enrichments; **Tab.**

1; **Supp. Tab. S10**). Besides the genes of interest for SD (*Dmrt-1L*/*Sox-H* for bivalves, and *Sry* for

mammals), other components of the DSFG families have been retrieved with a high AASD, despite

they have never been linked directly to SD so far: *Sox-F* in bivalves (**Fig. 3B**) and *Fox-D4* in

mammals (**Fig. 4B**). This implies that our approach cannot be used to unambiguously identify SDGs

alone. Instead, the analysis is meant to be used to detect highly divergent genes and, subsequently,

by comparison with literature and a more thorough and focused functional investigation, detect

putative SDGs among them. In this view, the mammal dataset exemplifies the importance of putting

the results of our pipeline (as those of any other comparative genomics analysis) into the correct

evolutionary and genomic context: among DSFGs of mammals, two genes exhibit high AASD, one

of which is directly related to SD (*Sry*), while the other has a function connected with neural

development (*Fox-D4*; **Klein et al., 2013**). Thus, the high AASD may arise either because of the

involvement in the upper SD pathway or because of other life-history traits connected with the gene,

respectively. Regarding bivalves, *Dmrt-1L* and *Sox-H* show a sharp connection with SD as a putative

primary SDG, either when considering their molecular evolutionary features or when looking at

their gene expression and possible function in gonad development (**Naimi et al., 2009;**

Teaniniuraitemoana et al., 2014; Zhang et al., 2014; Capt et al, 2018; Li et al., 2018; Afonso et

al., 2019; Liang et al., 2019; Yue et al., 2021). It is difficult to further speculate on the actual

involvement in SD of *Dmrt-1L* and *Sox-H* without any additional information on their biology, also

considering that genes evolve not only at the level of their amino acid sequences, but also at the

levels of exon-intron structures, indels, and regulatory relationships (among others). Nonetheless,

molecular evolution proves to be a valuable tool to investigate genes putatively involved in SD, and

to identify major targets onto which dedicate future research effort.

CONCLUSIONS

Genes functioning in reproductive processes, and particularly SDGs, are often among the most variable in animal genomes, in terms of both sequence composition and regulatory interactions (**Swanson et al., 2001**; **Bachtrong et al., 2014**). Such high evolutionary rates may be traced back both to adaptive evolution (either as natural or sexual selection) or to non-adaptive processes (**Vicoso & Charlesworth, 2006**; **Meisel & Connallon, 2013**; **Parsch & Ellegren, 2013**; **Grath & Parsch, 2016**) and often results in striking differences in reproductive and sexual systems even among closely-related species. In the present work we took advantage of this characteristic to identify SDGs among the DSFG families in bivalves. By comprehensively analysing the phylogenetic history and AASD in a broad taxonomic dataset, we appointed *Dmrt-1L* and *Sox-H* as putative SDGs, thus confirming results in previous works that found them to be transcribed in a male-biased manner and/or strongly involved in male-gonad formation (**Naimi et al., 2009; Teaniniuraitemoana et al., 2014; Zhang et al., 2014; Capt et al, 2018; Li et al., 2018; Afonso et al., 2019; Liang et al., 2019; Yue et al., 2021**). Future studies would now need to further investigate their evolutionary history. For example, considering that SRGs tend to accumulate in the genomic neighbourhood where primary SDGs are located (**Capel, 2017**), analysing the genomic location of DSFGs in bivalve genomes may provide enlightening results. Similarly, revealing the genetic interactions of *Dmrt-1L* and *Sox-H*, through functional and genome editing assays, would undoubtedly benefit our understanding of their role in the sexual processes of bivalves.

MATERIALS AND METHODS

Dataset of bivalve annotated genomes and transcriptomes

Annotated genome assemblies of bivalves were obtained from various publicly available resources, while reference genome assemblies for gastropods and cephalopods were downloaded from NCBI (**Supp. Tab. S1**). Isoforms from genome annotations were removed from genome annotations using

a perl script from the AGAT toolkit (v0.8.0; **Dainat, unpublished**). The nucleotide coding sequence

fasta file for *Sinonovacula constricta* (Adapedonta), was not available for download. To avoid

excluding the species from our analyses, the file was generated in-house by mapping the annotated

protein sequences on the reference genome using miniprot (v0.13-0; **Li, 2023**). Then, the

corresponding nucleotide sequences were extracted using AGAT on the resulting gff annotation file.

Assembled and annotated transcriptomes were obtained from **Piccinini et al., 2021, Iannello et al.,**

2023 and **Iannello et al., in preparation**. The resulting set of annotated genomes and

transcriptomes was checked for completeness using BUSCO with the Metazoa reference dataset

(v5.2.2; **Manni et al., 2021**).

Identification and classification of Dmrt, Sox and Fox genes in bivalves

Members of DSFG families were retrieved from the comprehensive bivalve dataset with hmmsearch

from the HMMER package (v3.3.2; http://hmmer.org/). The signature catalytic domains of each

family were used as queries. Specifically, hidden Markov model (HMM) profiles were built after

the Pfam databases for the DM domain (PF00751), the HMG box (PF00505) and the forkhead

domain (PF00250) to retrieve members of the DSFG families, respectively. The E-value for both

the per-target and the per-domain inclusion threshold was set to 10^{-5}. Obtained hits were then

annotated using (i) the PANTHER HMM standalone sequence scoring against the PANTHER

library v18.0 and (ii) RPS-BLAST (v2.5.0+) against CDD (pre-compiled version, downloaded from

https://ftp.ncbi.nih.gov/pub/mmdb/cdd/little_endian/ on 09/11/23). In both cases, hits with an E-

value of 10^{-5} were retained. Genes which were correctly annotated by both systems (on the basis of

the PANTHER gene family and CDD domain identifiers; **Supp. Tab. S9**) were kept for subsequent

analyses. DSFGs from *Homo sapiens*, *Drosophila melanogaster*, and *Caenorhabditis elegans*

(**Supp. Tab. S10**; the reference species) were retrieved from NCBI and were used as reference genes

for annotation (see below). Classification and nomenclature of each family was retrieved from:

Mawaribuchi et al., 2019 for Dmrt genes; **Phochanukul & Russell, 2010** and **Sarkar &**

Hochedlinger, 2013 for Sox genes; **Mazet et al., 2003** for Fox genes. The alignments of mollusc

and reference DSFGs were guided by the aforementioned Pfam HMM profiles and performed with

Clustal Omega (v1.2.3; **Sievers et al., 2011**), then trimmed with trimAl (v1.4.rev15; **Capella-**

Gutiérrez et al., 2009) with a gap threshold of 40%. Resulting alignments were manually inspected

to remove sequences with incomplete catalytic domains, then aligned and trimmed again as before.

Phylogenetic trees were inferred using IQ-TREE (v2.1.4-beta COVID-edition; **Minh et al., 2020**)

with automatic model selection (**Kalyaanamoorthy et al., 2017**), 1,000 bootstrap replicates and 5

independent runs. The phylogenetic tree of Dmrt genes was midpoint rooted, as no clear homology

relationship has been found with other gene families or zinc-finger proteins so far (**Wexler et al.,**

2014). Phylogenetic trees of Sox and Fox gene families were rooted using two fungi MATA-1

sequences (XP_62685912.1, CCD57795.1) and two Amoebozoa forkhead-like domains

(XP_004368148.1, XP_004333268.1), respectively (**Heenan et al., 2016**; **Nakagawa et al., 2016**).

The rooting was performed with Gotree (v0.4.5; Lemoine & Gascuel, 2021). To identify and

annotate bivalve homology groups within each gene family, we utilized a species overlap algorithm

followed by an MCL clustering weighted by node supports as implemented in Possvm (v1.2; **Grau-**

Bové & Sebé-Pedrós, 2021). DSFGs from the reference species were used as reference annotation.

To better establish the orthology relationships among ambiguous groups of Dmrt and Fox genes,

we ran a series of other phylogenetic reconstructions (see **Discussion**), by using the same pipeline

as before. In the case of *Fox-Y* genes, we also used Fox gene sequences from the sea urchin

Strongylocentrotus purpuratus, as given by **Tu et al. (2006)**. All the phylogenetic trees were plotted

using the R package 'ggtree' (**Yu et al., 2017a**).

Sequence diversity of bivalve single-copy orthogroups

For the analysis of AASD in bivalves, we assembled a second dataset (hereafter referred to as the

'reduced bivalve dataset'; **Fig. 1**) composed of 34 genomes and transcriptomes, to save

computational time and prevent over-represented taxa to bias the analysis. For each bivalve genus,

only the best genome and/or transcriptome in term of BUSCO and assembly statistics was kept

(**Supp. Tab. S1**). Genes were clustered in orthologous groups using OrthoFinder (v2.5.5; **Emms &**

Kelly, 2019) with DIAMOND ultra-sensitive and default parameters. The number of retrieved

SCOs was 5, which is drastically low but can be explained considering the mixed nature of the

dataset, that is, it includes both genomes and transcriptomes with highly different BUSCO scores

(**Supp. Tab. S1**). OrthoFinder orthogroups were therefore splitted into SCOs using DISCO (v1.3.1;

Willson et al., 2022), and orthogroups with at least 17 species (50% of the species included in the

bivalve reduced dataset) were retained. Amino acid and nucleotide sequences of SCOs were then

aligned using Clustal Omega as implemented in TranslatorX (v1.1; **Abascal et al., 2010**), and jointly

trimmed using trimAl with a gap threshold of 40% and the removal of spurious sequences (-

resoerlap 50 -seqoverlap 50). Eventually, orthogroups containing (i) internal stop codons, (ii) with

less than 17 species left (50% of the species included in the bivalve reduced dataset), or (iii)

containing DSFGs were removed from downstream analyses. The best amino acid substitution

model was inferred for each trimmed alignment using ModelFinder as implemented in IQTREE2

(model search was restricted to matrices accepted by the 'phangorn' R library, i.e., Blosum62,

cpREV, Dayhoff, DCMut, FLU, HIVb, HIVw, JTT, JTTDCMut, LG, mtART, mtMAM, mtREV,

mtZOA, rtREV, VT, WAG) and the corresponding pairwise amino acid distances were computed

with the function 'dist.ml' from the 'phangorn' R package (**Schliep 2011**). We decided to use the

pairwise amino acid distance instead of the tip-to-tip phylogenetic distance (which accounts for a

more comprehensive evolutionary signal) to save computational time. However, to check whether

the two metrics were comparable to each other, we randomly selected 200 decomposed orthogroups

(including orthogroups from the DSFGs) and computed the maximum likelihood (ML) trees using

IQTREE2, with ModelSelection restricted as before. Then, the tip-to-tip pairwise distances were

obtained with the R package 'adephylo' (**Jombart & Dray, 2010**). The same pipeline to obtain

SCOs and pairwise amino acid distances was also utilized on the DSFG families.

Mammals and *Drosophila* spp. as test datasets

For both mammals and fruit flies, annotated genomes were downloaded from NCBI using the command-line tool 'datasets', then processed using the same pipeline and scripts as before (**Fig. 1**).

GO term enrichment

After having obtained the distributions of AASD in the three datasets (Bivalvia, Mammalia, and *Drosophila*) and having sorted SCOs genes up into 3 groups (Group 1, Group 2, and Group 3), we performed a gene ontology (GO) enrichment analysis of genes from Group 1 and genes from Group 1 + Group 2. To do so, we firstly selected one gene per SCO, giving priority to few chosen species: (i) for bivalves, we selected genes from *Pecten maximus*, or alternatively from *Crassostrea gigas*, *Hyriopsis bialata*, *Tridacna squamosa*, and *Solen grandis*; (ii) for mammals, we selected genes from *H. sapiens*, or alternatively from *Bubalus bubalis*, *Panthera tigris*, *Camelus dromedarius*, and *Monodelphis domestica*; (iii) for fruit flies, we selected genes from *D. melanogaster*, or alternatively from *Drosophila hydei*, *Drosophila pseudoobscura*, and *Drosophila suzukii*. By doing so, we ensured that each SCO was represented by one gene. Afterwards, we annotated the obtained datasets with the corresponding GO terms using the OMA browser (accessed 18/09/2024; **Altenhoff et al., 2024**). The GO-term enrichment of Group 1 genes and Group 1 + Group 2 genes was performed with the R package 'topGO' using the Fisher's exact test (**Alexa et al., 2024**).

LIST OF ABBREVIATIONS

AASD: amino acid sequence divergence

DGE: differential gene expression

DM [domain]: *dsx* and *mab-3* [domain]

Dmrt: *doublesex* and *mab-3* related transcription factor

Dmrt-1L: *doublesex and mab-3 related transcription factor 1-like*

Dm-W: *Doublesex and mab-3 related gene W*

Dmy: *Doublesex and mab-3 related gene Y*

DSFGs: Dmrt, Sox, and Fox genes.

dsx: *doublesex*

Fox: forkhead box

ISH: *in-situ* hybridization

Mya: million years ago

SCO: single copy orthogroup

SD: sex determination

SDG: sex-determining gene

Sox: *Sry*-related HMG-box

SRG: sex-determination related gene

Sry: *Sex-determining region Y*

DATA AVAILABILITY STATEMENT

681   The data underlying this article and the codes used to run the analyses are available on GitHub

682   (https://github.com/filonico/bivalvia_SRGs).

ACKOWLEDGEMENTS

We would like to thank all the people who helped improving this work, by having promoted open discussions and/or exchanged ideas. But above all, we would like to thank Giobbe Forni, Mariangela Iannello and Giovanni Piccinini for always having offered their support whenever it was needed. This work was supported by the Canziani bequest funded to F.G. and A.L., by the European Union - NextGenerationEU under the National Recovery and Resilience Plan (PNRR) – Mission 4 Education and research – Component 2 From research to business - Investment 1.1, Notice Prin 2022 - DD N. 104 del 2/2/2022, entitled "Unraveling Sex Determination in Bivalves: An Integrative Single-Cell Analysis during the Early Embryo Development of *Mytilus galloprovincialis*", proposal code 2022CYM559 - CUP J53D23006200006 to L.M., and by the Saltonstall-Kennedy Program by NOAA Fisheries (SK-NOAA) funded to S.N.

REFERENCES

Abascal, F., Zardoya, R., & Telford, M. J. (2010). TranslatorX: multiple alignment of nucleotide sequences guided by amino acid translations. Nucleic acids research, 38(suppl_2), W7-W13.

Afonso, L. F., Americo, J. A., Soares-Souza, G. B., Torres, A. L. Q., Wajsenzon, I. J. R., & Rebelo, M. D. F. (2019). Gonad transcriptome of golden mussel Limnoperna fortunei reveals potential sex differentiation genes. biorxiv, 818757.

Albertin, C. B., Simakov, O., Mitros, T., Wang, Z. Y., Pungor, J. R., Edsinger-Gonzales, E., ... & Rokhsar, D. S. (2015). The octopus genome and the evolution of cephalopod neural and morphological novelties. Nature, 524(7564), 220-224.

Alexa, A., Rahnenfuhrer, J. (2024). topGO: Enrichment Analysis for Gene Ontology. R package version 2.56.0.

Allen, S. L., Delaney, E. K., Kopp, A., & Chenoweth, S. F. (2017). Single-molecule sequencing of the Drosophila serrata genome. G3: Genes, Genomes, Genetics, 7(3), 781-788.

Altenhoff, A. M., Warwick Vesztrocy, A., Bernard, C., Train, C. M., Nicheperovich, A., Prieto Baños, S., ... & Glover, N. M. (2024). OMA orthology in 2024: improved prokaryote coverage, ancestral and extant GO enrichment, a revamped synteny viewer and more in the OMA Ecosystem. Nucleic Acids Research, 52(D1), D513-D521.

Álvarez-Carretero, S., Tamuri, A. U., Battini, M., Nascimento, F. F., Carlisle, E., Asher, R. J., ... & Dos Reis, M. (2022). A species-level timeline of mammal evolution integrating phylogenomic data. Nature, 602(7896), 263-267.

Bachtrog, D., Mank, J. E., Peichel, C. L., Kirkpatrick, M., Otto, S. P., Ashman, T. L., ... & Tree of Sex Consortium. (2014). Sex determination: why so many ways of doing it?. PLoS biology, 12(7), e1001899.

Bai, C. M., Xin, L. S., Rosani, U., Wu, B., Wang, Q. C., Duan, X. K., ... & Wang, C. M. (2019). Chromosomal-level assembly of the blood clam, Scapharca (Anadara) broughtonii, using long sequence reads and Hi-C. GigaScience, 8(7), giz067.

Baldwin-Brown, J. G., Weeks, S. C., & Long, A. D. (2018). A new standard for crustacean genomes: the highly contiguous, annotated genome assembly of the clam shrimp Eulimnadia texana reveals HOX gene order and identifies the sex chromosome. Genome biology and evolution, 10(1), 143-156.

Baral, S., Arumugam, G., Deshmukh, R., & Kunte, K. (2019). Genetic architecture and sex-specific selection govern modular, male-biased evolution of doublesex. Science advances, 5(5), eaau3753.

Benayoun, B. A., Caburet, S., & Veitia, R. A. (2011). Forkhead transcription factors: key players in health and disease. Trends in Genetics, 27(6), 224-232.

Beukeboom, L.W., Perrin, N. (2014). The evolution of sex determination. Oxford: Oxford University Press.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics, 30(15), 2114-2120.

Bredemeyer, K. R., Hillier, L., Harris, A. J., Hughes, G. M., Foley, N. M., Lawless, C., ... & Murphy,

W. J. (2023). Single-haplotype comparative genomics provides insights into lineage-specific

structural variation during cat evolution. Nature genetics, 55(11), 1953-1963.

Breton, S., Capt, C., Guerra, D., & Stewart, D. (2018). Sex-determining mechanisms in bivalves.

Transitions between sexual systems: Understanding the mechanisms of, and pathways between,

dioecy, hermaphroditism and other sexual systems, 165-192.

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using

DIAMOND. Nature methods, 12(1), 59-60.

Capella-Gutiérrez, Salvador, José M. Silla-Martínez, and Toni Gabaldón. "trimAl: a tool for

automated alignment trimming in large-scale phylogenetic analyses." Bioinformatics 25.15

(2009): 1972-1973.

Capt, C., Renaut, S., Ghiselli, F., Milani, L., Johnson, N. A., Sietman, B. E., ... & Breton, S. (2018).

Deciphering the link between doubly uniparental inheritance of mtDNA and sex determination

in bivalves: clues from comparative transcriptomics. Genome biology and evolution, 10(2),

577-590.

Chakraborty, M., Chang, C. H., Khost, D. E., Vedanayagam, J., Adrion, J. R., Liao, Y., ... &

Emerson, J. J. (2021). Evolution of genome structure in the Drosophila simulans species

complex. Genome research, 31(3), 380-396.

Civetta, A., & Singh, R. S. (1998). Sex-related genes, directional sexual selection, and speciation.

Molecular biology and evolution, 15(7), 901-909.

Corrochano-Fraile, A., Davie, A., Carboni, S., & Bekaert, M. (2022). Evidence of multiple genome

duplication events in Mytilus evolution. BMC genomics, 23(1), 340.

Dainat J. AGAT: Another Gff Analysis Toolkit to handle annotations in any GTF/GFF format.

(Version v0.8.0). Zenodo.

de Bono, M., & Hodgkin, J. (1996). Evolution of sex determination in Caenorhabditis: unusually

high divergence of tra-1 and its functional consequences. Genetics, 144(2), 587-595.

Deng, T., Pang, C., Lu, X., Zhu, P., Duan, A., Tan, Z., ... & Liang, X. (2016). De novo transcriptome
assembly of the Chinese swamp buffalo by RNA sequencing and SSR marker discovery. PLoS
One, 11(1), e0147132.

Dermauw, W., Van Leeuwen, T., & Feyereisen, R. (2020). Diversity and evolution of the P450
family in arthropods. Insect biochemistry and molecular biology, 127, 103490.

Dong, C., Zhang, L., Xia, S., Sosa, D., Arsala, D., & Long, M. (2022). New gene evolution with
subcellular expression patterns detected in PacBio-sequenced genomes of Drosophila genus.
bioRxiv, 2022-11.

Echave, J., Spielman, S. J., & Wilke, C. O. (2016). Causes of evolutionary rate variation among
protein sites. Nature Reviews Genetics, 17(2), 109-121.

Elbers, J. P., Rogers, M. F., Perelman, P. L., Proskuryakova, A. A., Serdyukova, N. A., Johnson,
W. E., ... & Burger, P. A. (2019). Improving Illumina assemblies with Hi-C and long reads: An
example with the North African dromedary. Molecular Ecology Resources, 19(4), 1015-1026.

Ellegren, H., & Parsch, J. (2007). The evolution of sex-biased genes and sex-biased gene expression.
Nature Reviews Genetics, 8(9), 689-698.

Emms, D. M., & Kelly, S. (2019). OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome biology, 20, 1-14.

Evensen, K. G., Robinson, W. E., Krick, K., Murray, H. M., & Poynton, H. C. (2022). Comparative
phylotranscriptomics reveals putative sex differentiating genes across eight diverse bivalve
species. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 41,
100952.

Fan, H., Wu, Q., Wei, F., Yang, F., Ng, B. L., & Hu, Y. (2019). Chromosome-level genome
assembly for giant panda provides novel insights into Carnivora chromosome evolution.
Genome biology, 20, 1-12.

Farhat, S., Bonnivard, E., Pales Espinosa, E., Tanguy, A., Boutet, I., Guiglielmoni, N., ... & Allam,
B. (2022). Comparative analysis of the Mercenaria mercenaria genome provides insights into

the diversity of transposable elements and immune molecules in bivalve mollusks. BMC

genomics, 23(1), 192.

Foster, J. W., Brennan, F. E., Hampikian, G. K., Goodfellow, P. N., Sinclair, A. H., Lovell-Badge,

R., ... & Marshall Graves, J. A. (1992). Evolution of sex determination and the Y chromosome:

SRY-related sequences in marsupials. Nature, 359(6395), 531-533.

Foote, A. D., Liu, Y., Thomas, G. W., Vinař, T., Alföldi, J., Deng, J., ... & Gibbs, R. A. (2015).

Convergent evolution of the genomes of marine mammals. Nature genetics, 47(3), 272-275.

Fouret, J., Brunet, F. G., Binet, M., Aurine, N., Enchéry, F., Croze, S., ... & Legras-Lachuer, C.

(2020). Sequencing the genome of Indian flying fox, natural reservoir of nipah virus, using

hybrid assembly and conservative secondary scaffolding. Frontiers in Microbiology, 11, 1807.

Gerdol, M., Moreira, R., Cruz, F., Gómez-Garrido, J., Vlasova, A., Rosani, U., ... & Figueras, A.

(2020). Massive gene presence-absence variation shapes an open pan-genome in the

Mediterranean mussel. Genome biology, 21, 1-21.

Ghiselli, F., Milani, L., Chang, P. L., Hedgecock, D., Davis, J. P., Nuzhdin, S. V., & Passamonti,

M. (2012). De novo assembly of the Manila clam Ruditapes philippinarum transcriptome

provides new insights into expression bias, mitochondrial doubly uniparental inheritance and

sex determination. Molecular biology and evolution, 29(2), 771-786.

Ghiselli, F., Iannello, M., Puccio, G., Chang, P. L., Plazzi, F., Nuzhdin, S. V., & Passamonti, M.

(2018). Comparative transcriptomics in two bivalve species offers different perspectives on the

evolution of sex-biased genes. Genome biology and evolution, 10(6), 1389-1402.

Gomes-dos-Santos, A., Lopes-Lima, M., Castro, L. F. C., & Froufe, E. (2020). Molluscan genomics:

the road so far and the way forward. Hydrobiologia, 847(7), 1705-1726.

Gomes-dos-Santos, A., Lopes-Lima, M., Machado, A. M., Marcos Ramos, A., Usié, A., Bolotov, I.

N., ... & Froufe, E. (2021). The Crown Pearl: a draft genome assembly of the European

freshwater pearl mussel Margaritifera margaritifera (Linnaeus, 1758). DNA research, 28(2),

dsab002.

Gomes-dos-Santos, A., Lopes-Lima, M., Machado, M. A., Teixeira, A., C. Castro, L. F., & Froufe,

E. (2023). PacBio Hi-Fi genome assembly of the Iberian dolphin freshwater mussel Unio

delphinus Spengler, 1793. Scientific Data, 10(1), 340.

Gómez-Chiarri, M., Warren, W. C., Guo, X., & Proestou, D. (2015). Developing tools for the study

of molluscan immunity: the sequencing of the genome of the eastern oyster, Crassostrea

virginica. Fish & shellfish immunology, 46(1), 2-4.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., ... & Regev, A.

(2011). Trinity: reconstructing a full-length transcriptome without a genome from RNA-Seq

data. Nature biotechnology, 29(7), 644.

Grath, S., & Parsch, J. (2016). Sex-biased gene expression. Annual review of genetics, 50(1), 29-

44.

Grau-Bové, X., & Sebé-Pedrós, A. (2021). Orthology clusters from gene trees with Possvm.

Molecular Biology and Evolution, 38(11), 5204-5208.

Haag, E. S., & Doty, A. V. (2005). Sex determination across evolution: connecting the dots. PLoS

biology, 3(1), e21.

Haas, BJ. https://github.com/TransDecoder/TransDecoder.

Habtewold, T., Wagah, M., Tambwe, M. M., Moore, S., Windbichler, N., Christophides, G., ... &

Lawniczak, M. (2023). A chromosomal reference genome sequence for the malaria mosquito,

Anopheles gambiae, Giles, 1902, Ifakara strain. Wellcome Open Research, 8.

Han, F., Wang, Z., Wu, F., Liu, Z., Huang, B., & Wang, D. (2010). Characterization, phylogeny,

alternative splicing and expression of Sox30 gene. BMC Molecular Biology, 11, 1-11.

Hart, S. F., Yonemitsu, M. A., Giersch, R. M., Garrett, F. E., Beal, B. F., Arriagada, G., ... &

Metzger, M. J. (2023). Centuries of genome instability and evolution in soft-shell clam, Mya

arenaria, bivalve transmissible neoplasia. Nature Cancer, 4(11), 1561-1574.

Haszprunar, G., & Wanninger, A. (2012). Molluscs. Current Biology, 22(13), R510-R514.

Haerty, W., Jagadeeshan, S., Kulathinal, R. J., Wong, A., Ravi Ram, K., Sirot, L. K., ... & Singh, R.
S. (2007). Evolution in the fast lane: rapidly evolving sex-related genes in Drosophila. Genetics,
177(3), 1321-1335.

Heenan, P., Zondag, L., & Wilson, M. J. (2016). Evolution of the Sox gene family within the
chordate phylum. Gene, 575(2), 385-392.

Hemmer, L. W., & Blumenstiel, J. P. (2016). Holding it together: rapid evolution and positive
selection in the synaptonemal complex of Drosophila. BMC evolutionary biology, 16, 1-17.

Hoskins, R. A., Carlson, J. W., Wan, K. H., Park, S., Mendez, I., Galle, S. E., ... & Celniker, S. E.
(2015). The Release 6 reference sequence of the Drosophila melanogaster genome. Genome
research, 25(3), 445-458.

Iannello, M., Forni, G., Piccinini, G., Xu, R., Martelossi, J., Ghiselli, F., & Milani, L. (2023).
Signatures of extreme longevity: a perspective from bivalve molecular evolution. Genome
Biology and Evolution, 15(11), evad159.

Inoue, K., Yoshioka, Y., Tanaka, H., Kinjo, A., Sassa, M., Ueda, I., ... & Itoh, T. (2021). Genomics
and transcriptomics of the green mussel explain the durability of its byssus. Scientific Reports,
11(1), 5992.

Ip, J. C. H., Xu, T., Sun, J., Li, R., Chen, C., Lan, Y., ... & Qiu, J. W. (2021). Host–endosymbiont
genome integration in a deep-sea chemosymbiotic clam. Molecular Biology and Evolution,
38(2), 502-518.

Jackson, B. C., Carpenter, C., Nebert, D. W., & Vasiliou, V. (2010). Update of human and mouse
forkhead box (FOX) gene families. Human genomics, 4, 1-8.

Jombart, T., & Dray, S. (2010). Adephylo: exploratory analyses for the phylogenetic comparative
method. Bioinformatics, 26(15), 1-21.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K., Von Haeseler, A., & Jermiin, L. S. (2017).
ModelFinder: fast model selection for accurate phylogenetic estimates. Nature methods, 14(6),
587-589.

Kenny, N. J., McCarthy, S. A., Dudchenko, O., James, K., Betteridge, E., Corton, C., ... & Williams,

S. T. (2020). The gene-rich genome of the scallop Pecten maximus. Gigascience, 9(5), giaa037.

Kim, B. Y., Wang, J. R., Miller, D. E., Barmina, O., Delaney, E., Thompson, A., ... & Petrov, D. A.

(2021). Highly contiguous assemblies of 101 drosophilid genomes. Elife, 10, e66405.

Klein, S. L., Neilson, K. M., Orban, J., Yaklichkin, S., Hoffbauer, J., Mood, K., ... & Moody, S. A.

(2013). Conserved structural domains in FoxD4L1, a neural forkhead box transcription factor,

are required to repress or activate target genes. PloS one, 8(4), e61845.

Knudsen, B., Kohn, A. B., Nahir, B., McFadden, C. S., & Moroz, L. L. (2006). Complete DNA

sequence of the mitochondrial genome of the sea-slug, Aplysia californica: conservation of the

gene order in Euthyneura. Molecular Phylogenetics and Evolution, 38(2), 459-469.

Lan, Y., Sun, J., Chen, C., Sun, Y., Zhou, Y., Yang, Y., ... & Qian, P. Y. (2021). Hologenome

analysis reveals dual symbiosis in the deep-sea hydrothermal vent snail Gigantopelta aegis.

Nature communications, 12(1), 1165.

Larroux, C., Luke, G. N., Koopman, P., Rokhsar, D. S., Shimeld, S. M., & Degnan, B. M. (2008).

Genesis and expansion of metazoan transcription factor gene classes. Molecular biology and

evolution, 25(5), 980-996.

Lemoine, F., & Gascuel, O. (2021). Gotree/Goalign: toolkit and Go API to facilitate the

development of phylogenetic workflows. NAR Genomics and Bioinformatics, 3(3), lqab075.

Li, C., Liu, X., Liu, B., Ma, B., Liu, F., Liu, G., ... & Wang, C. (2018). Draft genome of the Peruvian

scallop Argopecten purpuratus. GigaScience, 7(4), giy031.

Li, R., Zhang, L., Li, W., Zhang, Y., Li, Y., Zhang, M., ... & Bao, Z. (2018). FOXL2 and DMRT1L

are yin and yang genes for determining timing of sex differentiation in the bivalve mollusk

Patinopecten yessoensis. Frontiers in Physiology, 9, 1166.

Li, F., Bian, L., Ge, J., Han, F., Liu, Z., Li, X., ... & Chen, S. (2020). Chromosome-level genome

assembly of the East Asian common octopus (Octopus sinensis) using PacBio sequencing and

Hi-C technology. Molecular ecology resources, 20(6), 1572-1582.

Li, A., Dai, H., Guo, X., Zhang, Z., Zhang, K., Wang, C., ... & Zhang, G. (2021). Genome of the

estuarine oyster provides insights into climate impact and adaptive plasticity. Communications

Biology, 4(1), 1287.

Li, X. Y., Mei, J., Ge, C. T., Liu, X. L., & Gui, J. F. (2022). Sex determination mechanisms and sex

control approaches in aquaculture animals. Science China Life Sciences, 65(6), 1091-1122.

Li, H. (2023). Protein-to-genome alignment with miniprot. Bioinformatics, 39(1), btad014.

Liang, S., Liu, D., Li, X., Wei, M., Yu, X., Li, Q., ... & Qin, Z. (2019). SOX2 participates in

spermatogenesis of Zhikong scallop Chlamys farreri. Scientific Reports, 9(1), 76.

Liao, Y., Zhang, X., Chakraborty, M., & Emerson, J. J. (2021). Topologically associating domains

and their role in the evolution of genome structure and function in Drosophila. Genome

research, 31(3), 397-410.

Liu, X. L., Zhang, Z. F., Shao, M. Y., Liu, J. G., & Muhammad, F. (2012). Sexually dimorphic

expression of foxl2 during gametogenesis in scallop Chlamys farreri, conserved with

vertebrates. Development Genes and Evolution, 222, 279-286.

Liu, C., Zhang, Y., Ren, Y., Wang, H., Li, S., Jiang, F., ... & Fan, W. (2018). The genome of the

golden apple snail Pomacea canaliculata provides insight into stress tolerance and invasive

adaptation. Gigascience, 7(9), giy101.

Liu, X., Li, C., Chen, M., Liu, B., Yan, X., Ning, J., ... & Wang, C. (2020). Draft genomes of two

Atlantic bay scallop subspecies Argopecten irradians irradians and A. i. concentricus. Scientific

Data, 7(1), 99.

Liu, T., Zhang, Y., Nie, H., Sun, J., & Yan, X. (2024). Characterization and expression patterns of

the Fox gene family under heat and cold stress in Manila clam Ruditapes philippinarum based

on genome-wide identification. Comparative Biochemistry and Physiology Part D: Genomics

and Proteomics, 101313.

Mahajan, S., Wei, K. H. C., Nalley, M. J., Gibilisco, L., & Bachtrog, D. (2018). De novo assembly
of a young Drosophila Y chromosome using single-molecule sequencing and chromatin
conformation capture. PLoS biology, 16(7), e2006348.

Mai, D., Nalley, M. J., & Bachtrog, D. (2020). Patterns of genomic differentiation in the Drosophila
nasuta species complex. Molecular biology and evolution, 37(1), 208-220.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO update:
novel and streamlined workflows along with broader and deeper phylogenetic coverage for
scoring of eukaryotic, prokaryotic, and viral genomes. Molecular biology and evolution,
38(10), 4647-4654.

Matson, C. K., & Zarkower, D. (2012). Sex and the singular DM domain: insights into sexual
regulation, evolution and plasticity. Nature Reviews Genetics, 13(3), 163-174.

Mawaribuchi, S., Yoshimoto, S., Ohashi, S., Takamatsu, N., & Ito, M. (2012). Molecular evolution
of vertebrate sex-determining genes. Chromosome Research, 20, 139-151.

Mawaribuchi, S., Ito, Y., & Ito, M. (2019). Independent evolution for sex determination and
differentiation in the DMRT family in animals. Biology Open, 8(8), bio041962.

Mazet, F., Yu, J. K., Liberles, D. A., Holland, L. Z., & Shimeld, S. M. (2003). Phylogenetic
relationships of the Fox (Forkhead) gene family in the Bilateria. Gene, 316, 79-89.

McCartney, M. A., Auch, B., Kono, T., Mallez, S., Zhang, Y., Obille, A., ... & Gohl, D. M. (2022).
The genome of the zebra mussel, Dreissena polymorpha: a resource for comparative genomics,
invasion genetics, and biocontrol. G3, 12(2), jkab423.

Mead, D., Fingland, K., Cripps, R., Miguez, R. P., Smith, M., Corton, C., ... & Blaxter, M. (2020).
The genome sequence of the eastern grey squirrel, Sciurus carolinensis Gmelin, 1788.
Wellcome Open Research, 5.

Meisel, R. P. (2011). Towards a more nuanced understanding of the relationship between sex-biased
gene expression and rates of protein-coding sequence evolution. Molecular Biology and
Evolution, 28(6), 1893-1900.

Meisel, R. P., & Connallon, T. (2013). The faster-X effect: integrating theory and data. Trends in

genetics, 29(9), 537-544.

Milani, L., Ghiselli, F., Nuzhdin, S. V., & Passamonti, M. (2013). Nuclear genes with sex bias in

Ruditapes philippinarum (Bivalvia, Veneridae): mitochondrial inheritance and sex

determination in DUI species. Journal of Experimental Zoology Part B: Molecular and

Developmental Evolution, 320(7), 442-454.

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., Von Haeseler, A.,

& Lanfear, R. (2020). IQ-TREE 2: new models and efficient methods for phylogenetic

inference in the genomic era. Molecular biology and evolution, 37(5), 1530-1534.

Moreno, J. A., Dudchenko, O., Feigin, C. Y., Mereby, S. A., Chen, Z., Ramos, R., ... & Mallarino,

R. (2024). Emx2 underlies the development and evolution of marsupial gliding membranes.

Nature, 1-9.

Mullon, C., Pomiankowski, A., & Reuter, M. (2012). Molecular evolution of Drosophila Sex-lethal

and related sex determining genes. BMC evolutionary biology, 12, 1-11.

Murat, F., Mbengue, N., Winge, S. B., Trefzer, T., Leushkin, E., Sepp, M., ... & Kaessmann, H.

(2023). The molecular evolution of spermatogenesis across mammals. Nature, 613(7943), 308-

316.

Murata, Y., Nikaido, M., Sasaki, T., Cao, Y., Fukumoto, Y., Hasegawa, M., & Okada, N. (2003).

Afrotherian phylogeny as inferred from complete mitochondrial genomes. Molecular

phylogenetics and evolution, 28(2), 253-260.

Naimi, A., Martinez, A. S., Specq, M. L., Diss, B., Mathieu, M., & Sourdaine, P. (2009). Molecular

cloning and gene expression of *Cg-Foxl2* during the development and the adult gametogenetic

cycle in the oyster *Crassostrea gigas*. Comparative Biochemistry and Physiology Part B:

Biochemistry and Molecular Biology, 154(1), 134-142.

Nakagawa, S., Gisselbrecht, S. S., Rogers, J. M., Hartl, D. L., & Bulyk, M. L. (2013). DNA-binding specificity changes in the evolution of forkhead transcription factors. Proceedings of the National Academy of Sciences, 110(30), 12349-12354.

Nicolini, F., Ghiselli, F., Luchetti, A., & Milani, L. (2023). Bivalves as emerging model systems to study the mechanisms and evolution of sex determination: a genomic point of view. Genome Biology and Evolution, 15(10), evad181.

Nicolini, F., Martelossi, J., Forni, G., Savojardo, C., Mantovani, B., & Luchetti, A. (2023). Comparative genomics of Hox and ParaHox genes among major lineages of Branchiopoda with emphasis on tadpole shrimps. Frontiers in Ecology and Evolution, 11, 1046960.

O'Neil, M. T., & Belote, J. M. (1992). Interspecific comparison of the transformer gene of Drosophila reveals an unusually high degree of evolutionary divergence. Genetics, 131(1), 113-128.

Orkin, J. D., Montague, M. J., Tejada-Martinez, D., De Manuel, M., Del Campo, J., Cheves Hernandez, S., ... & Melin, A. D. (2021). The genomics of ecological flexibility, large brains, and long lives in capuchin monkeys revealed with fecalFACS. Proceedings of the National Academy of Sciences, 118(7), e2010632118.

Paggeot, L. X., DeBiasse, M. B., Escalona, M., Fairbairn, C., Marimuthu, M. P., Nguyen, O., ... & Dawson, M. N. (2022). Reference genome for the California ribbed mussel, Mytilus californianus, an ecosystem engineer. Journal of Heredity, 113(6), 681-688.

Pamilo, P., & O'Neill, R. J. (1997). Evolution of the Sry genes. Molecular biology and evolution, 14(1), 49-55.

Panara, V., Budd, G. E., & Janssen, R. (2019). Phylogenetic analysis and embryonic expression of panarthropod Dmrt genes. Frontiers in zoology, 16, 1-18.

Papa, F., Windbichler, N., Waterhouse, R. M., Cagnetti, A., D'Amato, R., Persampieri, T., ... & Papathanos, P. A. (2017). Rapid evolution of female-biased genes among four species of Anopheles malaria mosquitoes. Genome research, 27(9), 1536-1548.

Paris, M., Boyer, R., Jaenichen, R., Wolf, J., Karageorgi, M., Green, J., ... & Prud'homme, B.

(2020). Near-chromosome level genome assembly of the fruit pest Drosophila suzukii using

long-read sequencing. Scientific reports, 10(1), 11227.

Parsch, J., & Ellegren, H. (2013). The evolutionary causes and consequences of sex-biased gene

expression. Nature Reviews Genetics, 14(2), 83-87.

Penaloza, C., Gutierrez, A. P., Eöry, L., Wang, S., Guo, X., Archibald, A. L., ... & Houston, R. D.

(2021). A chromosome-level genome assembly for the Pacific oyster Crassostrea gigas.

GigaScience, 10(3), giab020.

Phochanukul, N., & Russell, S. (2010). No backbone but lots of Sox: Invertebrate Sox genes. The

international journal of biochemistry & cell biology, 42(3), 453-464.

Piccinini, G., Iannello, M., Puccio, G., Plazzi, F., Havird, J. C., & Ghiselli, F. (2021). Mitonuclear

coevolution, but not nuclear compensation, drives evolution of OXPHOS complexes in

bivalves. Molecular Biology and Evolution, 38(6), 2597-2614.

Powell, D., Subramanian, S., Suwansa-Ard, S., Zhao, M., O'Connor, W., Raftos, D., & Elizur, A.

(2018). The genome of the oyster Saccostrea offers insight into the environmental resilience of

bivalves. DNA Research, 25(6), 655-665.

Qi, H., Cong, R., Wang, Y., Li, L., & Zhang, G. (2023). Construction and analysis of the

chromosome-level haplotype-resolved genomes of two Crassostrea oyster congeners:

Crassostrea angulata and Crassostrea gigas. GigaScience, 12, giad077.

Ran, Z., Li, Z., Yan, X., Liao, K., Kong, F., Zhang, L., ... & Xu, J. (2019). Chromosome-level

genome assembly of the razor clam Sinonovacula constricta (Lamarck, 1818). Molecular

ecology resources, 19(6), 1647-1658.

Ranz, J. M., Go, A. C., González, P. M., Clifton, B. D., Gomes, S., Jaberyzadeh, A., ... & Civetta,

A. (2023). Gene expression differentiation in the reproductive tissues of Drosophila willistoni

subspecies and their hybrids. Molecular Ecology, 32(13), 3605-3623.

Renschler, G., Richard, G., Valsecchi, C. I. K., Toscano, S., Arrigoni, L., Ramírez, F., & Akhtar, A.

(2019). Hi-C guided assemblies reveal conserved regulatory topologies on X and autosomes

despite extensive genome shuffling. Genes & development, 33(21-22), 1591-1612.

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., ... & Jarvis, E. D. (2021).

Towards complete and error-free genome assemblies of all vertebrate species. Nature,

592(7856), 737-746.

Rogers, R. L., Grizzard, S. L., Titus-McQuillan, J. E., Bockrath, K., Patel, S., Wares, J. P., ... &

Moore, C. C. (2021). Gene family amplification facilitates adaptation in freshwater unionid

bivalve Megalonaias nervosa. Molecular Ecology, 30(5), 1155-1173.

Russo, C. A., Mello, B., Frazão, A., & Voloch, C. M. (2013). Phylogenetic analysis and a time tree

for a large drosophilid data set (Diptera: Drosophilidae). Zoological Journal of the Linnean

Society, 169(4), 765-775.

Sanchez-Flores, A., Peñaloza, F., Carpinteyro-Ponce, J., Nazario-Yepiz, N., Abreu-Goodger, C.,

Machado, C. A., & Markow, T. A. (2016). Genome evolution in three species of cactophilic

Drosophila. G3: Genes, Genomes, Genetics, 6(10), 3097-3105.

Sarkar, A., & Hochedlinger, K. (2013). The sox family of transcription factors: versatile regulators

of stem and progenitor cell fate. Cell stem cell, 12(1), 15-30.

Schliep, K. P. (2011). phangorn: phylogenetic analysis in R. Bioinformatics, 27(4), 592-593.

Schomburg, C., Janssen, R., & Prpic, N. M. (2022). Phylogenetic analysis of forkhead transcription

factors in the Panarthropoda. Development genes and evolution, 232(1), 39-48.

Seudre, O., Martín-Zamora, F. M., Rapisarda, V., Luqman, I., Carrillo-Baltodano, A. M., & Martín-

Durán, J. M. (2022). The Fox gene repertoire in the annelid Owenia fusiformis reveals multiple

expansions of the foxQ2 class in Spiralia. Genome Biology and Evolution, 14(10), evac139.

Shimeld, S. M., Boyle, M. J., Brunet, T., Luke, G. N., & Seaver, E. C. (2010). Clustered Fox genes

in lophotrochozoans and the evolution of the bilaterian Fox gene cluster. Developmental

biology, 340(2), 234-248.

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., ... & Higgins, D. G. (2011).

Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal

Omega. Molecular systems biology, 7(1), 539.

Smith, C. H. (2021). A high-quality reference genome for a parasitic bivalve with doubly uniparental

inheritance (Bivalvia: Unionida). Genome Biology and Evolution, 13(3), evab029.

Song, H., Wang, Y., Shao, H., Li, Z., Hu, P., Yap-Chiongco, M. K., ... & Kocot, K. M. (2023).

Scaphopoda is the sister taxon to Bivalvia: Evidence of ancient incomplete lineage sorting.

Proceedings of the National Academy of Sciences, 120(40), e2302361120.

Stammnitz, M. R., Gori, K., Kwon, Y. M., Harry, E., Martin, F. J., Billis, K., ... & Murchison, E. P.

(2023). The evolution of two transmissible cancers in Tasmanian devils. Science, 380(6642),

283-293.

Stothard, P., & Pilgrim, D. (2003). Sex-determination gene and pathway evolution in nematodes.

Bioessays, 25(3), 221-231.

Sun, J., Zhang, Y. U., Xu, T., Zhang, Y., Mu, H., Zhang, Y., ... & Qian, P. Y. (2017). Adaptation to

deep-sea chemosynthetic environments as revealed by mussel genomes. Nature ecology &

evolution, 1(5), 0121.

Sun, D., Yu, H., & Li, Q. (2022). Examination of the roles of Foxl2 and Dmrt1 in sex differentiation

and gonadal development of oysters by using RNA interference. Aquaculture, 548, 737732.

Swanson, W. J., & Vacquier, V. D. (2002). The rapid evolution of reproductive proteins. Nature

reviews genetics, 3(2), 137-144.

Teaniniuraitemoana, V., Huvet, A., Levy, P., Klopp, C., Lhuillier, E., Gaertner-Mazouni, N., ... &

Le Moullac, G. (2014). Gonad transcriptome analysis of pearl oyster Pinctada margaritifera:

identification of potential sex differentiation and sex determining genes. BMC genomics, 15,

1-20.

Teng, W., Fu, H., Li, Z., Zhang, Q., Xu, C., Yu, H., ... & Li, Q. (2023). Parallel evolution in
Crassostrea oysters along the latitudinal gradient is associated with variation in multiple genes
involved in adipogenesis. Molecular Ecology, 32(19), 5276-5287.

Tu, Q., Brown, C. T., Davidson, E. H., & Oliveri, P. (2006). Sea urchin Forkhead gene family:
phylogeny and embryonic expression. Developmental biology, 300(1), 49-62.

Tvedte, E. S., Gasser, M., Sparklin, B. C., Michalski, J., Hjelmen, C. E., Johnston, J. S., ... &
Dunning Hotopp, J. C. (2021). Comparison of long-read sequencing technologies in
interrogating bacteria and fly genomes. G3, 11(6), jkab083.

Uller, T., & Helanterä, H. (2011). From the origin of sex-determining factors to the evolution of
sex-determining systems. The Quarterly review of biology, 86(3), 163-180.

Vicoso, B., & Charlesworth, B. (2006). Evolution on the X chromosome: unusual patterns and
processes. Nature Reviews Genetics, 7(8), 645-653.

Vilstrup, J. T., Seguin-Orlando, A., Stiller, M., Ginolhac, A., Raghavan, M., Nielsen, S. C., ... &
Orlando, L. (2013). Mitochondrial phylogenomics of modern and ancient equids. PloS one,
8(2), e55950.

Vinkler, M., Fiddaman, S. R., Těšický, M., O'Connor, E. A., Savage, A. E., Lenz, T. L., ... &
Westerdahl, H. (2023). Understanding the evolution of immune genes in jawed vertebrates.
Journal of evolutionary biology, 36(6), 847-873.

Vizueta, J., Sánchez-Gracia, A., & Rozas, J. (2020). BITACORA: A comprehensive tool for the
identification and annotation of gene families in genome assemblies.

Wang, G., Dong, S., Guo, P., Cui, X., Duan, S., & Li, J. (2020). Identification of Foxl2 in freshwater
mussel Hyriopsis cumingii and its involvement in sex differentiation. Gene, 754, 144853.

Wang, C., Wallerman, O., Arendt, M. L., Sundström, E., Karlsson, Å., Nordin, J., ... & Lindblad-
Toh, K. (2021). A novel canine reference genome resolves genomic architecture and uncovers
transcript complexity. Communications biology, 4(1), 185.

Wang, S., Zhang, J., Jiao, W., Li, J. I., Xun, X., Sun, Y., ... & Bao, Z. (2017). Scallop genome provides insights into evolution of bilaterian karyotype and development. Nature ecology & evolution, 1(5), 0120.

Wang, Q., Cao, T., & Wang, C. (2023). Genome-wide identification and expression analysis of Dmrt genes in bivalves. BMC genomics, 24(1), 457.

Wang, J., & Nie, H. (2024). Genome-wide identification and expression analysis of Sox gene family in the Manila clam (Ruditapes philippinarum). Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 50, 101244.

Wei, M., Ge, H., Shao, C., Yan, X., Nie, H., Duan, H., ... & Dong, Z. (2020). Chromosome-level clam genome helps elucidate the molecular basis of adaptation to a buried lifestyle. IScience, 23(6).

Wexler, J. R., Plachetzki, D. C., & Kopp, A. (2014). Pan-metazoan phylogeny of the DMRT gene family: a framework for functional studies. Development Genes and Evolution, 224, 175-181.

Whitfield, L. S., Lovell-Badge, R., & Goodfellow, P. N. (1993). Rapid sequence evolution of the mammalian sex-determining gene SRY. Nature, 364(6439), 713-715.

Wilkins, A. S. (1995). Moving up the hierarchy: a hypothesis on the evolution of a genetic sex determination pathway. Bioessays, 17(1), 71-77.

Willson, J., Roddur, M. S., Liu, B., Zaharias, P., & Warnow, T. (2022). DISCO: species tree inference using multicopy gene family tree decomposition. Systematic Biology, 71(3), 610-629.

Wu, S., Zhang, Y., Li, Y., Wei, H., Guo, Z., Wang, S., ... & Bao, Z. (2020). Identification and expression profiles of Fox transcription factors in the Yesso scallop (Patinopecten yessoensis). Gene, 733, 144387.

Xiong, Y. E., Brandley, M. C., Xu, S., Zhou, K., & Yang, G. (2009). Seven new dolphin mitochondrial genomes and a time-calibrated phylogeny of whales. BMC Evolutionary Biology, 9, 1-13.

Xu, R., Martelossi, J., Smits, M., Iannello, M., Peruzza, L., Babbucci, M., ... & Ghiselli, F. (2022).

Multi-tissue RNA-Seq analysis and long-read-based genome assembly reveal complex sex-

specific gene regulation and molecular evolution in the Manila clam. Genome Biology and

Evolution, 14(12), evac171.

Yang, M., Xu, F., Liu, J., Que, H., Li, L., & Zhang, G. (2014). Phylogeny of forkhead genes in three

spiralians and their expression in Pacific oyster Crassostrea gigas. Chinese journal of

oceanology and limnology, 32(6), 1207-1223.

Yang, J. L., Feng, D. D., Liu, J., Xu, J. K., Chen, K., Li, Y. F., ... & Lu, Y. (2021). Chromosome-

level genome assembly of the hard-shelled mussel Mytilus coruscus, a widely distributed

species from the temperate areas of East Asia. Gigascience, 10(4), giab024.

Yu, J. K., Mazet, F., Chen, Y. T., Huang, S. W., Jung, K. C., & Shimeld, S. M. (2008). The Fox

genes of Branchiostoma floridae. Development genes and evolution, 218, 629-638.

Yu, G., Smith, D. K., Zhu, H., Guan, Y., & Lam, T. T. Y. (2017). ggtree: an R package for

visualization and annotation of phylogenetic trees with their covariates and other associated

data. Methods in Ecology and Evolution, 8(1), 28-36.

Yu, J., Zhang, L., Li, Y., Li, R., Zhang, M., Li, W., ... & Bao, Z. (2017). Genome-wide identification

and expression profiling of the SOX gene family in a bivalve mollusc Patinopecten yessoensis.

Gene, 627, 530-537.

Yue, C., Li, Q., & Yu, H. (2021). Variance in expression and localization of sex-related genes

CgDsx, CgBHMG1 and CgFoxl2 during diploid and triploid Pacific oyster Crassostrea gigas

gonad differentiation. Gene, 790, 145692.

Zeng, Y., Zheng, H., He, C., Zhang, C., Zhang, H., & Zheng, H. (2024). Genome-wide identification

and expression analysis of Dmrt gene family and their role in gonad development of Pacific

oyster (Crassostrea gigas). Comparative Biochemistry and Physiology Part B: Biochemistry

and Molecular Biology, 269, 110904.

Zhang, N., Xu, F., & Guo, X. (2014). Genomic analysis of the Pacific oyster (*Crassostrea gigas*)

reveals possible conservation of vertebrate sex determination in a mollusc. G3: Genes,

Genomes, Genetics, 4(11), 2207-2217.

Zhou, Y., Shearwin-Whyatt, L., Li, J., Song, Z., Hayakawa, T., Stevens, D., ... & Zhang, G. (2021).

Platypus and echidna genomes reveal mammalian biology and evolution. Nature, 592(7856),

756-762.

FIGURES

1146

Fig. 1. Workflow for the bivalve dataset. Starting from a set of both genomes and transcriptomes covering a great portion of bivalve taxonomic diversity, we first characterized the entire complement of Dmrt, Sox, and Fox genes (upper row). We used sequence annotation and phylogenetic tools to obtain reliable sequences and filter out any putative mis-assembled or mis-annotated sequence. Afterwards, we built a reduced set of transcriptomes and genomes (the reduced bivalve dataset, where we minimized the redundancy of congeneric species) from which to draw the molecular evolution patterns of orthologous genes (bottom row). After having obtained gene single-copy orthologous groups, we calculated the amino acid distances within each orthogroup and then we built the distribution of median values. The same pipeline was also utilized for the mammal and the fruit fly datasets, with just two minor differences: the starting dataset was composed of only genomes, and the reduction step (R) was not necessary.

1158

Fig. 2. Dmrt, Sox, and Fox gene (DSFG) complement in bivalves and their outgroups. The presence of genes in various species are indicated by filled circles. Numbers inside each circle specify genes with 2 or more copies. The shaded area highlights non-bivalve species, belonging either to other molluscs or to the references. The phylogenetic tree of analyzed species, as inferred from literature, is shown on the left, while major taxonomic groups are reported on the right. Species

represented by transcriptomic data are marked with an asterisk ('*'), and species not present in the

reduced bivalve dataset are marked with two asterisks ('**'; see main text and **Fig. 1**); note that the

two categories do nor overlap. DSFG trees are shown on the bottom (full trees can be found in **Supp.**

Fig. S1–S3). Full species names, along with all assembly and taxonomic information, can be found

in **Supp. Tab. S1**. Ad.: Adapedonta; Ar.: Arcida; Ca.: Cardiida; Ce.: Cephalopoda; L.: Lucinida;

My.: Myida; Ref.: reference genes; S.: Sphaeriida.

Fig. 3. Distribution of amino acid sequence divergence (AASD) of single-copy orthogroups in

bivalves (A), including Dmrt, Sox, and Fox genes (DSFGs; B), and their correlations with tip-

to-tip distances (C), alignment lengths (D), and number of species (E). The distribution of AASD

has been computed on the median values of pairwise distances of over 11k SCOs from the reduced

bivalve dataset (see main text and **Fig. 1**). Genes have been divided according to their median AASD

value into three different groups, which are indicated by different colours and increasing numbers

(Groups 1, 2, and 3). Genes from Group 1 and Group 2 are collectively named 'highly divergent

genes'. Circle heights of DSFGs show the median value of their AASD, while the size indicates the

number of represented species. DSFG trees are shown on the bottom (full trees can be found in

Supp. Fig. S1–S3). Darker points in C-E indicate Dmrt, Sox, and Fox gene SCOs. The correlation

between the amino acid distance and the tip-to-tip distance has been computed on 200 randomly

selected orthogroups. **Insets:** scheme of the sex-determination working model in *Crassostrea gigas*

as hypothesised by **Zhang et al. (2014)**, showing the main genes involved. Green arrows indicate

transcription activations, red arrows indicate transcription suppressions.

Fig. 4. Distribution of amino acid divergence (AASD) of single-copy orthogroups in Mammalia (A) and *Drosophila* (C), including Dmrt, Sox, and Fox genes (DSFGs; B-D). The distributions of AASD in mammals and fruit flies have been computed on the median values of pairwise distances of over 16k and 12k SCOs, respectively. Genes have been divided according to their median AASD value into three different groups, which are indicated by different colors and increasing numbers

(Groups 1, 2, and 3). Genes from Group 1 and Group 2 are collectively named 'highly divergent

genes'. Circle heights of DSFGs show the median value of their AASD, while the size indicates the

number of represented species. DSFG trees are shown on the bottom (full trees can be found in

Supp. Fig. S5–S7 for mammals and in **Supp. Fig. S8–S10** for fruit flies). In *Drosophila*, *Sxl* and

tra, both involved in the SD pathway (inset) do not belong to the group of highly divergent genes

(mean amino acid divergence of 0.0923793827106834 and 0.978190195613595, respectively; that

is, within the boundaries of Group 3). **Insets:** scheme of the sex-determination molecular pathways

in *Mus musculus* and in *Drosophila melanogaster*, showing the main genes involved (adapted from

Beukeboom & Perrim, 2014). Green arrows indicate transcription activations, red arrows indicate

transcription suppressions. X: sex chromosomes; A: autosomal chromosomes; $DSX^{M/F}$: *DSX*

splicing variants present in males or females, respectively.

TABLES

Tab. 1. Top enriched GO terms for highly divergent genes of bivalves, mammals, and *Drosophila*. The extended version of the table, which includes also the expected number of annotated genes per GO term and all the other enriched GO terms, can be accessed in **Supp. Tab. S6**.

Dataset	GO.ID	Term	Annotated genes	Significant genes	Corrected p-value
Bivalvia	GO:0060255	regulation of macromolecule metabolic process	737	59	0.04525
	GO:0080090	regulation of primary metabolic process	673	53	0.01818
	GO:0019219	regulation of nucleobase-containing compound metabolic process	541	41	0.02388
	GO:0006351	DNA-templated transcription	571	39	0.03767
	GO:0032774	RNA biosynthetic process	579	39	0.04490
	GO:0051252	regulation of RNA metabolic process	517	37	0.02719
	GO:0006355	regulation of DNA-templated transcription	490	35	0.03751
	GO:2001141	regulation of RNA biosynthetic process	491	35	0.03844
	GO:0006950	response to stress	370	33	0.01949
	GO:0032502	developmental process	261	27	0.04445
	GO:0006468	protein phosphorylation	345	23	0.02483
	GO:0031325	positive regulation of cellular metabolic process	125	17	0.00801
	GO:0010604	positive regulation of macromolecule metabolic process	151	17	0.04047
	GO:0051172	negative regulation of nitrogen compound metabolic process	117	16	0.00814
	GO:0051173	positive regulation of nitrogen compound metabolic process	137	15	0.02454
	GO:0006310	DNA recombination	66	14	0.00087
	GO:0048513	animal organ development	83	12	0.04088
	GO:0010629	negative regulation of gene expression	78	11	0.00048
	GO:0023051	regulation of signaling	133	11	0.02872
	GO:0045934	negative regulation of nucleobase-containing compound metabolic process	64	11	0.03637
	GO:0009605	response to external stimulus	90	11	0.04544
	GO:0044419	biological process involved in interspecies interaction between organisms	63	11	0.04761
Mammalia	GO:0006955	immune response	1297	145	0.00061
	GO:0098542	defense response to other organism	853	112	0.02066
	GO:0045087	innate immune response	647	82	8.5e-10
	GO:0001817	regulation of cytokine production	630	51	0.04660
	GO:0042742	defense response to bacterium	233	45	1.7e-07
	GO:0006954	inflammatory response	642	45	0.01735
	GO:0019221	cytokine-mediated signaling pathway	382	44	3.9e-07
	GO:0002250	adaptive immune response	342	44	1.3e-05
	GO:0001819	positive regulation of cytokine production	402	41	0.02723
	GO:0002697	regulation of immune effector process	308	37	0.04426
	GO:0042110	T cell activation	432	35	0.02564
	GO:0051607	defense response to virus	257	34	1.9e-07
	GO:0048232	male gamete generation	491	32	0.02255

	GO:0007283	spermatogenesis	478	31	0.02801
	GO:0070661	leukocyte proliferation	273	29	0.01285
	GO:0002449	lymphocyte mediated immunity	221	29	0.04833
	GO:0070663	regulation of leukocyte proliferation	212	25	0.01870
	GO:0050727	regulation of inflammatory response	300	24	0.00235
	GO:0031349	positive regulation of defense response	240	24	0.01239
	GO:0002768	immune response-regulating cell surface receptor signaling pathway	177	22	0.00336
	GO:0050829	defense response to Gram-negative bacterium	66	17	1.7e-10
	GO:0071222	cellular response to lipopolysaccharide	164	17	0.00012
	GO:0010466	negative regulation of peptidase activity	163	16	0.00036
	GO:0002429	immune response-activating cell surface receptor signaling pathway	164	16	0.00243
	GO:1903555	regulation of tumor necrosis factor superfamily cytokine production	137	16	0.01244
	GO:0071706	tumor necrosis factor superfamily cytokine production	137	16	0.01244
	GO:0070665	positive regulation of leukocyte proliferation	132	16	0.02765
	GO:0045089	positive regulation of innate immune response	113	16	0.03224
	GO:0071356	cellular response to tumor necrosis factor	175	15	0.00219
	GO:0002695	negative regulation of leukocyte activation	148	15	0.01151
	GO:0002456	T cell mediated immunity	82	15	0.01605
	GO:0002705	positive regulation of leukocyte mediated immunity	113	15	0.01837
	GO:0032680	regulation of tumor necrosis factor production	133	15	0.03262
	GO:0032640	tumor necrosis factor production	133	15	0.03262
	GO:0050866	negative regulation of cell activation	165	15	0.04048
	GO:0000819	sister chromatid segregation	140	11	0.02927
	GO:0070192	chromosome organization involved in meiotic cell cycle	54	9	0.00849
	GO:0007131	reciprocal meiotic recombination	37	7	0.00066
	GO:0007143	female meiotic nuclear division	54	6	0.02270
	GO:0035967	cellular response to topologically incorrect protein	44	5	0.03334
	GO:0035966	response to topologically incorrect protein	47	5	0.04266
Drosophila	GO:0007141	male meiosis I	13	4	0.00150
	GO:0140543	positive regulation of piRNA transcription	3	3	6.9e-05
	GO:0010526	retrotransposon silencing	8	3	0.00331
	GO:0007130	synaptonemal complex assembly	10	3	0.00666
	GO:0030719	P granule organization	11	3	0.00888
	GO:0071218	cellular response to misfolded protein	12	3	0.01149
	GO:0051788	response to misfolded protein	12	3	0.01149
	GO:0007135	meiosis II	15	3	0.02169
	GO:0034508	centromere complex assembly	19	3	0.04094

1208

SUPPLEMENTARY FIGURES

Supp. Fig. S1. ML phylogenetic tree of the Dmrt gene family in molluscs, including the Possvm orthology inference. Reference genes from *H. sapiens*, *C. elegans*, and *D. melanogaster* are marked with an asterisk at the beginning of the tip. Species ID can be found in **Supp. Tab. S1**. Bootstrap values are shown for each node.

Supp. Fig. S2. ML phylogenetic tree of the Sox gene family in molluscs, including the Possvm orthology inference. Reference genes from *H. sapiens*, *C. elegans*, and *D. melanogaster* are marked with an asterisk at the beginning of the tip. Species ID can be found in **Supp. Tab. S1**. Bootstrap values are shown for each node.

Supp. Fig. S3. ML phylogenetic tree of the Fox gene family in molluscs, including the Possvm orthology inference. Reference genes from *H. sapiens*, *C. elegans*, and *D. melanogaster* are marked with an asterisk at the beginning of the tip. Species ID can be found in **Supp. Tab. S1**. Bootstrap values are shown for each node.

Supp. Fig. S4. Dmrt, Sox, and Fox gene (DSFG) complement in Mammalia and *Drosophila* spp. Presence/absence of genes in various species are indicated by filled circles. Numbers inside each circle specify genes with 2 or more copies. The shaded area highlights outgroup species, *Gallus gallus* (Aves) for mammals and *Anopheles gambiae* (Culicidae) for fruit flies. The phylogenetic tree of analyzed species, as inferred from literature, is shown on the left, while major taxonomic groups are reported on the right. All species are represented by genomic data. DSFG trees are shown on the bottom (full trees can be found in **Supp. Fig. S5–S7**). Full species names for both mammals and fruit flies, along with all assembly and taxonomic information, can be found in **Supp. Tab. S4** and **Supp. Tab. S5**, respectively. A.: Aves; Chirop.: Chiroptera; L.: Lagomorpha; M.: Monotremata;

1234   Me.: Metatheria; P.: Pholidota; Pe.: Perissodactyla; Prim.: Primates; Roden.: Rodentia; X.:

1235   Xenarthra; C.: Culicidae.

1236

1237   **Supp. Fig. S5. ML phylogenetic tree of the Dmrt gene family in mammals, including the**

1238   **Possvm orthology inference.** Reference genes from *H. sapiens*, *Mus musculus*, *Elephas maximus*

1239   *indicus*, and *Ornithorhynchus anatinus* are marked with an asterisk at the beginning of the tip.

1240   Species ID can be found in **Supp. Tab. S4**. Bootstrap values are shown for each node.

1241

1242   **Supp. Fig. S6. ML phylogenetic tree of the Sox gene family in mammals, including the Possvm**

1243   **orthology inference.** Reference genes from *H. sapiens*, *Mus musculus*, *Elephas maximus indicus*,

1244   and *Ornithorhynchus anatinus* are marked with an asterisk at the beginning of the tip. Species ID

1245   can be found in **Supp. Tab. S4**. Bootstrap values are shown for each node.

1246

1247   **Supp. Fig. S7. ML phylogenetic tree of the Fox gene family in mammals, including the Possvm**

1248   **orthology inference.** Reference genes from *H. sapiens*, *Mus musculus*, *Elephas maximus indicus*,

1249   and *Ornithorhynchus anatinus* are marked with an asterisk at the beginning of the tip. Species ID

1250   can be found in **Supp. Tab. S4**. Bootstrap values are shown for each node.

1251

1252   **Supp. Fig. S8. ML phylogenetic tree of the Dmrt gene family in fruit flies, including the Possvm**

1253   **orthology inference.** Reference genes from *Drosophila melanogaster*, *Drosophila hydei*,

1254   *Drosophila pseudoobscura*, and *Drosophila suzukii* are marked with an asterisk at the beginning of

1255   the tip. Species ID can be found in **Supp. Tab. S5**. Bootstrap values are shown for each node.

1256

1257   **Supp. Fig. S9. ML phylogenetic tree of the Sox gene family in fruit flies, including the Possvm**

1258   **orthology inference.** Reference genes from *Drosophila melanogaster*, *Drosophila hydei*,

1259 *Drosophila pseudoobscura*, and *Drosophila suzukii* are marked with an asterisk at the beginning of

1260 the tip. Species ID can be found in **Supp. Tab. S5**. Bootstrap values are shown for each node.

1261

1262 **Supp. Fig. S10. ML phylogenetic tree of the Fox gene family in fruit flies, including the Possvm**

1263 **orthology inference.** Reference genes from *Drosophila melanogaster*, *Drosophila hydei*,

1264 *Drosophila pseudoobscura*, and *Drosophila suzukii* are marked with an asterisk at the beginning of

1265 the tip. Species ID can be found in **Supp. Tab. S5**. Bootstrap values are shown for each node.

1266

SUPPLEMENTARY TABLES

Supp. Tab. S1. Genomic and transcriptomic data of bivalves and other molluscs. For each

species, the relative ID, taxonomic information, BUSCO statistics, NCBI accession number, and

source publication are reported. Biv: Bivalvia; Ca: Caenogastropoda; Cep: Cephalopoda; Co:

Coleoidea; Gas: Gastropoda; Gen: Genome; He: Heterobranchia; Im: Imparidentia; Ne:

Neomphaliones; Pa: Palaeoheterodonta; Pt: Pteriomorpha; Tra: Transcriptome; Ve: Vetigastropoda.

1273

1274 **Supp. Tab. S2. Genomic data of mammals used to retrieve DSFGs and compute amino acid**

1275 **sequence divergence (AASD) of single copy orthogroups (SCOs).** For each species, the relative

1276 ID, taxonomic information, BUSCO statistics, NCBI accession number, and source publication are

1277 reported.

1278

1279 **Supp. Tab. S3. Genomic data of Drosophila used to retrieve DSFGs and compute AASD of**

1280 **SCOs.** For each species, the relative ID, taxonomic information, BUSCO statistics, NCBI accession

1281 number, and source publication are reported.

1282

1283 **Supp. Tab. S4. Complete set of DSFGs in bivalves.** For each gene, the species ID (Sp. ID) as in

1284 **Supp. Tab. S1**, the accession number (Gene ID), the Possvm-based annotation, and the CDD

1285 domains (including their Pssm-ID) are indicated.

1286

1287 **Supp. Tab. S5. Proportions of missing data in bivalve DSFGs.**

1288

1289 **Supp. Tab. S6. All the enriched GO terms for Group 1 and Group 2 genes of bivalves,**

1290 **mammals, and *Drosophila*.**

1291

1292 **Supp. Tab. S7. Complete set of DSFGs in mammals.** For each gene, the species ID (Sp. ID) as in

1293 **Supp. Tab. S4**, the accession number (Gene ID), and the Possvm-based annotation are indicated.

1294

1295 **Supp. Tab. S8. Complete set of DSFGs in *Drosophila*.** For each gene, the species ID (Sp. ID) as

1296 in **Supp. Tab. S5**, the accession number (Gene ID), and the Possvm-based annotation are indicated.

1297

1298 **Supp. Tab. S9. Dmrt, Sox, and Fox gene (DSFG) family and domain identifiers (IDs) in**

1299 **PANTHER and CDD, respectively.** After having retrieved putative DSFGs based on HMM

1300 profiles, IDs have been used to retain only reliable hits.

1301

1302 **Supp. Tab. S10. List of DSFGs from reference species used to assess the identity of DSFGs in**

1303 **molluscs.** NCBI accession numbers are reported in parenthesis. Each row represents an orthology

1304 group.

59