Corticospinal Excitability in Response to Mediolateral Gait Instability

AUTHORS: Raven O. Huiberts[1], Sjoerd M. Bruijn[1] & Jennifer L. Davies[2]

AFFILIATIONS: *[1] Department of Human Movement Sciences, Faculty of Behavioural and Movement Sciences, Vrije Universiteit Amsterdam and Amsterdam Movement Sciences, Amsterdam, The Netherlands*

[2] School of Healthcare Sciences, Cardiff University, United Kingdom

Correspondence: Jennifer Davies, School of Healthcare Sciences, Ty Dewi Sant, Cardiff University Heath Park Campus, Cardiff, CF14 4XN; DaviesJ@cardiff.ac.uk; 02920 688581

Word count abstract: 278

Word count manuscript: 4391 (excluding Summary, figure legends, acknowledgements, disclosures and references)

Figures and tables: 10 figures

SUMMARY

Unpredictable gait disturbances, particularly in the mediolateral direction, pose a significant challenge to stability and are a common contributor to falls. Although the corticospinal tract is critical for gait and postural control, its response to such instabilities remains unclear. To investigate if corticospinal excitability increases during unpredictable lateral destabilisations, single-pulse transcranial magnetic stimulations were delivered over the primary motor cortex of 15 healthy individuals during steady-state and randomly laterally destabilised treadmill gait. Full-body kinematics were recorded using an optoelectronic motion capture system. Stimulations where the coil was >5 mm from the target location on the head were excluded. Corticospinal excitability was quantified for four upper- and three lower-leg muscles as the amplitude of the motor evoked potential (MEP) and compared between steady-state and destabilised gait. Lateral destabilisation resulted in a wider step width and shorter stride duration with increased variability and greater dynamic instability. Foot placement control during laterally destabilised gait was increased at midstance and there was also an increased average foot placement error in this condition. No differences in corticospinal excitability were observed in the lower-leg muscles. All upper-leg muscles demonstrated greater absolute MEPs in destabilised relative to steady-state gait. After normalising to muscle activity these periods became less pronounced, however, increases in corticospinal excitability during destabilised gait were observed in all but the gastrocnemius muscles. These finding suggest a heightened readiness of the corticospinal tract projecting to upper-leg muscles during destabilised gait, which could reflect general stabilising strategies such as decreasing stride time and increasing step width. Increases in the relative contribution of the corticospinal tract (i.e. normalised MEPs) were primarily noted during periods of low muscle activity, possibly hinting towards greater motor planning.

46 **KEY WORDS:**

47 *Mediolateral gait stability, Motor cortex, Corticospinal excitability, Transcranial magnetic*

48 *stimulation, Human gait*

INTRODUCTION

Human gait is a complex task that requires coordinated activity of around 200 muscles.

Despite the complexity of human gait, it can be seemingly effortless. Historically, it was

assumed that this was achieved primarily through spinal control systems, a notion based on

observations that quadrupedal mammals are capable of basic locomotion rhythmicity even in

the absence of supraspinal processing [1–6]. While there are shared control mechanisms

between human and quadrupedal gait [7,8], cortical lesions in humans can drastically

debilitate gait [9,10] indicating an important role for supraspinal control. This likely reflects

the greater stability demands of bipedal gait [11]. Understanding these supraspinal processes

is essential, particularly for advancing fall prevention strategies and neurorehabilitation.

Transcranial magnetic stimulation (TMS) can be used to probe supraspinal processes. TMS of

the motor cortex depolarises pyramidal tract cells and, with sufficient excitation, this results in

the generation of action potentials that descend along the corticospinal tract to depolarise

alpha motor neurones in the spinal cord [12–14]. The subsequent firing of motor units can be

measured using surface electromyography (EMG) as a motor evoked potential (MEP). The

size of the MEP reflects the excitability of the corticospinal tract [15,16].

Using TMS, it has been demonstrated that corticospinal excitability changes throughout the

gait cycle, generally aligning with phasic muscle activity [17–19]. However, these excitability

patterns do not always mirror ongoing muscle activity. Schubert and colleagues [20] observed

increased excitability prior to phasic muscle activation and suggested that this reflects

increased supraspinal input as an anticipatory adjustment aimed to preserve gait stability.

Gait stability is the ability to counteract the forces that induce moments to accelerate our

centre of mass (CoM) away from the base of support, such as gravity and internal and external

perturbations, by generating appropriate muscles forces, in order to remain upright [21–23].

Gait stability requires active control in the mediolateral direction [24,25], which is achieved

through different strategies encompassing many muscles. Among these, the gluteus medius is

often depicted as a primary contributor due to its important role in foot placement control

[26–30]. Mediolateral gait stability control deteriorates with ageing [31,32] and is a strong

predictor of fall risk [33,34]. Improving the understanding of how mediolateral gait stability is

achieved will allow the development of strategies to prevent and reverse this deterioration,

thereby reducing risk of falls.

During standing balance, a positive correlation between mediolateral sway velocity and the

excitability of the corticospinal tract projecting to tibialis anterior has been reported [35].

However, less is known about how excitability of the corticospinal tract projecting to leg

muscles is modulated during mediolaterally unstable gait, particularly when faced with

unpredictable perturbations, such as missteps or lateral pushes in daily life. Since falls often

result from such unpredictable disturbances, we test if corticospinal excitability increases in

response to unpredictable mediolateral gait instabilities.

We hypothesise that corticospinal excitability will increase in accordance with mediolateral

stability demands. Increases in corticospinal excitability will be observed throughout the gait

cycle, reflecting heightened readiness to respond to unpredictable perturbations.

METHODS

Participants

Twenty-three participants (seven male, 16 females aged 18-40 years) provided written

informed consent to participate. All participants reported that they were healthy adults without

any contraindications to TMS or other risk factors that could impact the safety or reliability of

this study (https://doi.org/10.17605/OSF.IO/7AJUZ and [36]). Participants were informed that

they could withdraw from the experiment at any time. All procedures were conducted in

accordance with the Cardiff University School of Healthcare Sciences Research Ethics

Committee. There were no adverse events. Six participants were excluded because no MEP

could be observed within a comfortable stimulation intensity. One participant was excluded

because they had to leave early. After processing, another participant was excluded because

the TMS coil moved too much during the walking trials, resulting in less than 100 useful

stimulations per condition. Data from the remaining 15 participants are reported here.

Procedures

The participants performed both steady-state and laterally destabilised treadmill gait within a

Gait Realtime Analysis Interactive Lab (GRAIL) system (Motek Medical BV, Houten,

Netherlands). Destabilisation was induced by continuous pseudorandom mediolateral

oscillations of the treadmill with a peak-to-peak amplitude of 100 mm. The velocity of the

treadmill was set at a constant speed of 4.0 km/h. Each condition (steady-state and laterally

destabilised) consisted of a 200-s familiarisation period followed by a 20-min trial. The two

conditions were performed in a single session with a break in between. The order of the

conditions was randomised.

Motion capture

During the familiarisation and the trials, full-body kinematics were captured at 100 Hz using

Vicon Nexus software (version 2.12, Oxford, UK). Spherical retroreflective markers (12-mm

diameter) were placed in accordance with the full-body Plug-in Gait model (Vicon 2.12,

Oxford, UK) with some modifications: the markers on the side/back of the head were placed

on the side/back of the helmet (see below), and additional markers were placed on the nasal

bridge and the left and right tragus.

For real-time detection of heel strike, the HBM 2.0 model (Motek Medical BV,

https://knowledge.motekmedical.com/wp-content/uploads/2019/07/HBM2-Reference-Manual-

Full-Body.pdf) was used. Kinematic data were streamed into D-flow software (version 3.36,

Motek Medical BV) and processed in real time.

EMG

125 Before the trials, surface electrodes (Trigno Avanti Sensor, Delsys, Boston, MA, USA) were

126 placed on the following muscles on the right leg according to the SENIAM guidelines

127 (SENIAM): adductor magnus, biceps femoris, gluteus medius, rectus femoris, tibialis anterior,

128 gastrocnemius medialis, gastrocnemius lateralis. EMG data was collected at a sampling

129 frequency of 2000 Hz in Vicon Nexus software and was synchronised with kinematic data.

TMS

TMS (posterior-anterior current) was administered using a custom-made bent batwing coil

(Magstim, Whitland, Wales). The coil was positioned on the scalp over the left motor cortex

and position was adjusted until MEPs could be observed in the tibialis anterior of the right

leg. The position of the coil on the head of the participant was fixed using of a custom-

designed helmet (Figure 1). The weight of the coil and helmet was supported by two spring-

coil balancers that were attached the ceiling over the centre of the treadmill. The tension in the

spring-coil balancers was adjusted for each participant so that the coil rested on the scalp with

enough weight to ensure stability of coil position throughout the gait cycle (i.e., to prevent the

coil 'bouncing' with vertical head movement) but enough weight removed so that this would

not become uncomfortable over the duration of the experiment.

[insert figure 1]

For each participant, motor threshold was determined as the minimal stimulation intensity

required to evoke an MEP in tibialis anterior while the participant was standing quietly. This

was done according to the following procedure: First, the intensity of the stimulator was set to

35% of the maximal stimulation output, which (for most individuals) is a subthreshold

intensity within the limits of tolerability. Thereafter, the intensity was progressively increased

in steps of 5% until a discernible MEP could be observed in at least 5 out of 10 stimulations,

or the participant reported that they found the stimulation intolerable. If the latter occurred,

the session was terminated (n = 6/23 participants). If the former occurred, the intensity was

decreased in steps of 2% until the simulation did not evoke an MEP in 5 out of 10

stimulations. Lastly, the intensity was increased again in steps of 1% until the lowest intensity

that consistently evoked a MEP was found.

During the experimental trials, the TMS intensity was set at 110% of motor threshold.

Stimulations were given with random delays after heel strike with a varying number of strides

between the stimulations (ranging from 3 to 5 strides). This resulted in approximately 200

stimulations over the 20-minute trial, evenly distributed across the stride cycle. The

stimulation instant was recorded from the output of Dflow via a Phidget Interface Kit 8/8/8

(Phidgets Inc. Calgary, AB, Canada) with a Trigno Analog Input Adaptor (Delsys) that was

sampled at 2000 Hz in Vicon Nexus software (version 2.12) synchronised with EMG data.

Relative coil displacement

Prior to the experiment, a static measurement was conducted using an additional marker

placed at a central point within the helmet. During the experimental trials, the virtual marker's

position was calculated using both the fixed helmet markers and the head markers. To

determine the displacement of the helmet relative to the head, the deviation of the virtual

marker calculated from the head markers was compared to its intended position, as defined by

the rigid markers on the helmet.

Data processing

Gait events for offline analysis were calculated with a custom code where heel strike was

identified as the moment the heel marker reach its lowest point and toe-off was identified as

the moment the heel marker reached its highest upwards acceleration

(https://github.com/SjoerdBruijn/GaitEvents).

Gait pattern

The gait pattern in each condition was described using step width (average and variance),

stride duration (average and variance), dynamic stability and quality of foot placement control

in the last 150 strides of the 200-s familiarisation period.

Step width was defined as the mediolateral distance between the heels at heel strike. Stride

duration was defined as the time interval between two consecutive right heel strikes.

Dynamic stability was calculated with the local divergence exponent (λ_s), a measure that

quantifies the resilience to infinitesimal perturbations that occur during gait [37]. It is

computed by determining the average exponential rate of divergence of neighbouring

trajectories in state space [38]. Higher values indicate greater local instability. For these

calculations, Rosenstein's algorithm was used [39]. The state space was created from time-

normalised mediolateral velocity time series of the averaged thorax markers (C7 and left and

right acromion) and five time-delayed copies. The λ_s was estimated as the slope of the mean

divergence curve, with a normalised stride time from 0 to 0.5 stride [40,41].

The quality of foot placement control was calculated using the model of Wang and Srinivasan

[28] (utilising codes from [42]).This model assumes a linear relation between the deviations

in CoM state and subsequent foot placement (Equation below). Within our analysis the pelvis

(PEL) was used instead of the CoM.

$$Foot\ Placement = \alpha(i) + \beta(i)_{position} * PEL(i)_{position} + \beta(i)_{velocity} * PEL(i)_{velocity} + \epsilon(i)$$

• $\beta_{position} = position\ regression\ coefficient$

• $\beta_{velocity} = velocity\ regression\ coefficient$

• $\epsilon = residual\ variance$ (i.e. foot placement error)

The foot placement was calculated in the single-leg stance phase, defined as the time between

contralateral toe off and heel strike. To account for displacement, such as wandering

backward, forward, left, or right, the coordinate system was taken from the previous stance

foot. Additionally, within the laterally oscillating treadmill condition, the $PEL_{velocity}$ was

corrected by adding the treadmill's lateral velocity to the derivative of the $PEL_{position}$. The

degree of foot placement control was quantified by the relative explained variance (R^2). To

attain normality in the data fisher transformation was performed on the R^2 values. Lastly, as

the R^2 indicates the percentage of the foot placement variance explained by the variance in

PEL states, we also calculated the average foot placement error as a measure of foot

placement precision. This was calculated as the standard deviation of ϵ at the instant of foot

placement.

Ongoing EMG and MEPs

EMG signals were bandpass filtered between 20 Hz and 500 Hz using a second-order bi-

directional Butterworth filter. Ongoing EMG was calculated by rectifying and then averaging

the filtered signal from 20 ms prior to stimulation to 10 ms after stimulation. The stimulation

artifact was excluded by removing the data point at the time of stimulation (Figure 2).

Absolute MEP amplitudes were calculated as the peak-to-peak amplitude of the filtered signal

within the time window of 10 to 60 ms post-stimulation. Normalised MEPs were determined

by dividing the absolute MEP by the ongoing EMG. MEPs were excluded if the relative

helmet-head position at the time of stimulation was >5 mm along the vertical axis or >10 mm

in the horizontal plane (combined displacement) from the original relative position. For each

stimulation, the corresponding percentage of the gait cycle was determined based on the

percentage of the total time between consecutive right heel strikes.

[insert figure 2]

For the ongoing EMG, non-normalised MEPs and normalised MEPs, all data points were

weight based smoothed utilising SMART [43] with adjustments. First, all data points were

linked to their corresponding gait cycle percentage (Figure 3A). Thereafter, the estimated

time-series was reconstructed for each participant by convolving the kernel with the

outcomes. This was then divided by the sum of the kernel density (weights) to maintain the

proper scaling (Figure 3B & C). Lastly, normalised weights were calculated and averaged

across participants for each time point in the gait cycle. This method ensures that participants

with higher sample density at specific percentages of the gait cycle have a greater impact on

the smoothed outcome at those corresponding time points.

[insert figure 3]

To accommodate the cyclical nature of the data, the SMART code was modified by replacing

the standard Gaussian distribution, which assumes linear data, with the Von Mises distribution

(Figure 4). To prevent bias by outliers, a smoothing kernel containing sufficient data points

was required. For this at least six data points had to be contained within 2 standard deviations

(σ) of each weight distribution (Figure 3B and 4). As the data displayed an average density of

$1.99 \pm 0.58/1\%$ (steady-state) and $2.02 \pm 0.55/1\%$ (laterally destabilised) for each participant,

a σ of 2% was chosen. Consequently, the Von Mises distribution with a kappa (κ) of 60,

which contained the same amount of datapoints as a normal distribution with a σ of 2%

(Figure 4A), was used.

[insert figure 4]

Statistics

Statistics were performed in Python (version 3.12.0). All kinematic outcomes, apart from the

relative explained variance of the foot placement model, were compared between steady-state

and laterally destabilised gait using paired t-statistics.

The relative explained variance (R^2) of the foot placement model over a step, ongoing EMG, non-normalised MEPs and normalised MEPs were compared between steady-state and laterally destabilised gait using cluster based-permutation testing according to the following procedure: First, t-tests were applied to each time point. However, since a difference at one time point is not independent of its neighbouring time points, condition differences were tested over clusters of consecutive time points, all of which exhibited statistically significant difference. This was done by summing the T values of a cluster and comparing this to summed T values of the permuted data's cluster (i.e. shuffling the data in a random order followed by smoothing it). Clusters were considered significant if their summed T value was greater than the 95th percentile of the 1000 permutations. This procedure was performed according to the code of Leeuwen and colleagues [43]. For all muscle activity and corticospinal excitability measures, this method was adjusted to account for the cyclic nature of the data, by wrapping the clusters around, connecting the clusters at the start and end of the gait cycle (https://doi.org/10.17605/OSF.IO/7AJUZ). For all statistical tests a significance threshold α of 0.05 was used.

RESULTS:

Of the 16 participants who completed the full experiment, one was excluded due to excessive relative movement between their head and the coil (i.e. fewer than 100 remaining stimulations per condition). Data are presented from the remaining 15 participants (four males and 11 females; age 18–40 years). Seven participants performed the laterally destabilised condition first and seven performed the steady-state condition first. Two participants received TMS stimulation at 100% of motor threshold instead of the targeted 110% due to reporting discomfort at the higher intensity. However, MEPs were still observed in these participants and their results are included.

Gait stability and quality of foot placement control

The lateral destabilisation evoked a significant increase in step width, step width variability

and stride duration variability and a significant decrease in stride duration and the local

dynamic stability, demonstrated by increased λ_s (Figure 5).

[insert figure 5]

The relative explained variance (Fisher-transformed R) of the foot placement model was

greater in the laterally destabilised condition than in the steady-state condition at midstance.

However, this difference was not observed in the rest of the stepcycle. Additionally, there was

a significantly greater average foot placement error in the laterally destabilised condition

(Figure 6).

[insert figure 6]

Ongoing EMG and MEPs

In total, more than 3100 stimulations (average 212 steady-state and 215 destabilised

stimulations per participant) per condition were within the relative displacement threshold and

included in analysis. This was 97.76% of the total stimulations. Steady-state and laterally

destabilised gait were compared using a temporal resolution of 0.5% and a κ of 60. Our

normalised weight distribution indicate standard deviations of sample spread of $7.7*10^{-4}$ and

$4.9*10^{-4}$ for the steady-state and the laterally destabilised condition, respectively (Figure 7).

[insert figure 7]

Ongoing EMG

Compared to normal walking, destabilised walking increased muscle activity during some

phase of the gait cycle in all muscles except the gastrocnemius medialis (Figure 8).

[insert figure 8]

Absolute MEPs

No differences in absolute MEP amplitude were detected in the lower-leg muscles (gastrocnemius medialis, gastrocnemius lateralis, and tibialis anterior). However, all upper-leg muscles demonstrated greater absolute MEPs during laterally destabilised gait than during steady-state walking (Figure 9). In the rectus femoris, MEP amplitude was larger over almost the entire gait cycle. A similar consistent increase in absolute MEPs was observed in the biceps femoris, with no observed differences during the terminal swing and a short period during the stance phase. The adductor magnus exhibited increased absolute MEPs from terminal swing into the early stance phase. In the gluteus medius, three periods of significantly increased absolute MEPs were observed: during early stance, from late stance into early swing, and during the late swing phase.

[insert figure 9]

Normalised MEPs

After normalising the absolute MEPs to the ongoing EMG, a short period of significantly increased MEPs during destabilised as compared to steady-state walking became apparent in the tibialis anterior, and increases remained significant in all upper-leg muscles (Figure 10).

In tibialis anterior, normalised MEP was greater during laterally destabilised gait than during steady-state walking during the transition from stance to the double support phase. In the rectus femoris, increases were noted during the stance and mid-swing phases. The biceps femoris demonstrated increased normalised MEPs during the stance phase and from double support into the mid-swing phase. The adductor magnus muscle exhibited two periods of significantly increased normalised MEPs: both during mid stance. Lastly, one period of increased normalised MEP was observed in the gluteus medius during the second double support phase.

[insert figure 10]

DISCUSSION:

We investigated whether unpredictable mediolateral gait instability was associated with

greater excitability of the corticospinal tract to multiple leg muscles. For this purpose, we

compared MEPs elicited during steady-state treadmill gait to those elicited during laterally

destabilised treadmill gait. Our perturbation increased gait instability and caused participants

to adopt a more cautious walking strategy. This was accompanied by increased activity of

proximal and distal leg muscles and increased excitability of the corticospinal tract,

particularly to proximal leg muscles.

Gait stability and quality of foot placement control

Similar to other studies [44,45], we observed a larger step width, step width variability, stride

duration variability and shorter stride duration during laterally destabilised gait than during

steady-state gait. Furthermore, we observed increased λ_s in the laterally destabilised condition,

indicating poorer local dynamic stability, i.e. more susceptibility to small perturbations [46].

Increased variability in stride duration and step width is often associated with increased risk

of falls [47]; however, it can also be attributed to a purposeful adjustment response to

overcome the stability demands [11,44]. One of the main mechanisms behind this adjustment

is foot placement control. While we did observe a tightening of this foot placement control

during midstance, this did not result in a tightening of the actual foot placement (i.e. reduction

in foot placement error at the end of the step cycle). The quality of foot placement control can

increase due to a greater variance in the CoM_{states}, which most likely will happen due to the

treadmill's oscillations. Thus, we also estimated the foot placement control error [48], and

found that this increased during lateral destabilised gait (Figure 5), indicating a decreased

precision in foot placement. This likely results from the constant nature of the perturbations,

as foot placement strategies require time to be executed [27,49,50]. Altogether participants

widened their steps and increased stride frequency to maintain a stable gait in the presence of

perturbations [51]. Increases in step width and reductions in step length are general strategies

characterising "cautious walking" in unpredictable situations [51–53].

It is important to consider that the gait parameters measured during familiarisation might

deviate from those during the actual trial due to the introduction of TMS. TMS itself already

works as a central perturbation, leading to anticipation and compensatory adjustments for the

evoked movements [54]. While both conditions should be equally affected by the additional

stimulation, the gait patterns during the familiarisation might not perfectly align those during

the trials.

Muscle activity and absolute MEPs

Muscle activity and absolute MEP amplitudes followed a similar phasic modulation over the

gait cycle, which is in line with previous studies [17–19]. The cautious walking pattern

employed in laterally destabilised gait was associated with greater activity of all muscles

except gastrocnemius medialis. In the upper-leg muscles, this was mirrored by greater

absolute MEPs. However, in lower-leg muscles there was no change in the amplitude of

absolute MEPs. This was unexpected, but may be due to the method of destabilisation

employed.

During steady-state gait, mediolateral stability is the dynamic equilibrium between the

moments created by gravity and the forces generated by the muscles [55,56]. Primarily, the

hip abductors, plantar flexors and to a lesser extent the dorsiflexors produce forces counteract

the destabilisation due to gravity [22,55,57]. However, we suggest that in our laterally

destabilised gait condition there was little demand on lower-leg muscles to compensate for the

perturbations. Although lower-leg muscles play an important role in steady-state gait stability

[58] and ankle moments have the capacity to correct for preceding foot placement errors [59],

the unpredictability of constant and random surface shifts during gait may have necessitated a

greater reliance on the upper-leg muscles. This is reflected by the rectus femoris and

bicepsfemoris demonstrating increases in MEP amplitude over a large portion of the gait

cycle, which likely served as a proactive control strategy [15], aligning with the 'cautious

walking' strategy. For the gluteus medius and adductor magnus the responses seem more

specific, with the increases aligning the muscles contributions to mediolateral acceleration

[55]. Therefore, we suggest that the method of destabilisation limited the applicability of

lower-leg muscles strategies, forcing participants to rely on upper-leg muscles to maintain gait

stability.

Normalised MEPs

The clear link between ongoing (pre-stimulus) EMG and absolute MEP amplitude is

consistent with the established linear dependency between ongoing voluntary muscle activity

and MEP amplitude [60]. However, this relation can become uncoupled if the corticospinal

tract gets more (or less) access to the spinal motor neurone pool at a given level of muscle

activity. Schubert and colleagues [20] reported early increases in tibialis anterior and

gastrocnemius medialis MEPs prior to changes in phasic muscle activity during gait, which

they interpreted as a preparatory mechanism to preserve gait stability. Although we observed

some increases in normalised MEPs prior to the muscle activity (Figure 10: tibialis anterior

from 50 to 60% , biceps femoris 70 to 85% and gluteus medius 95 to 5%), which were greater

in laterally destabilised gait for the tibialis anterior and biceps femoris, this effect was not

consistently observed across all muscles, nor was it amplified in the muscles known to be

most involved in stabilising gait [11], such as the gluteus medius. Therefore, we suspect that

early (normalised) MEP facilitation does not universally respond to (external) preparatory

stability demands. Further research is required to understand why early facilitations occur.

Normalised MEPs were greater during destabilised gait than during steady-state gait in all

upper-leg muscles and the TA. This indicates that the changes in excitability of the

corticospinal tract were not solely driven by changes in muscle activity. Highlighting that

excitability changes serve additional roles besides modulating the ongoing muscle activity

[35].

No changes in normalised MEPs were observed across conditions for the gastrocnemii, this

again likely reflects the limited use of ankle strategies to counteract our method of

destabilisation [61,62]. The facilitation in normalised MEPs in the TA during the second

double support phase may represent a preparatory effect to ensure toe clearance [63].The

gluteus medius plays a predominant role in mediolateral gait stability yet there was only a

short period around the second double support phase where normalised MEPs in this muscle

were increased in the destabilised condition [11]. Although this is a transitional phase that

does demand stabilisation [64], it does not reflect the phase with the greatest muscle potential

to control this destabilisation, which is mostly seen as swing leg's midstance or the stance

leg's late stance phase [11,50]. Similarly, the timepoints at which normalised MEPs were

increased in the destabilised condition in the other muscles, although potentially functional,

had no clear connection with specific stabilisation strategies [55,58]. This raises the question

whether these changes are linked to stability demands, phasic postural adjustments or other

factors.

We should take into account that our results could be affected by the inherent variability in

both ongoing EMG and MEP signals. MEPs are subject to rapid, spontaneous fluctuations in

corticospinal excitability, especially during periods of low excitability, which complicates

their interpretation [65,66]. Additionally, during periods of low muscle activity, EMG signals

become more susceptible to noise, further complicating the interpretation of normalised MEPs

[67]. Therefore, caution is advised in interpreting normalised MEPs, especially during periods

of low excitability or muscle activity. Interestingly, regardless of the higher variability across

stimuli within a condition during these phases of low muscle activity, most of the differences

between conditions that we found were within periods of low excitability and muscle activity.

This potentially indicates that the observed differences in normalised MEPs might be

reflective of motor planning, typically occurring during pre-active states [68]. Such planning-

related increases in MEP amplitude align with the idea that preparatory excitability reflects

task complexity, even without concurrent increases in EMG activity [69].

Conclusions

In conclusion, we investigated the effects of unpredictable lateral destabilisation on the

excitability of the corticospinal tract to multiple leg muscles. We found that the imposed

perturbations increased instability and led participants to adopt a plethora of "cautious

walking" strategies. This was associated with periods of increased absolute corticospinal

excitability in all upper-leg muscles along with sustained increases in normalised MEPs,

particularly at times of low muscle activity. These findings suggest that constant postural

destabilisations heighten the responsiveness of the corticospinal tract to proximal leg muscles,

suggesting that the corticospinal tract is involved in the adaptations to these conditions.

Acknowledgments

This research was funded by a Royal Society International Exchange Award to JD and SB and

a Cardiff University Taith Student Mobility Award to RH.

Supplementary material

Opensource material of the raw data/codes and the questionnaire can be found at:

https://doi.org/10.17605/OSF.IO/7AJUZ.

REFERENCES

1. Kiehn O. LOCOMOTOR CIRCUITS IN THE MAMMALIAN SPINAL CORD. Annu Rev
Neurosci. 2006;29:279–306.

2. Fedirchuk B, Nielsen J, Petersen N, Hultborn H. Pharmacologically evoked fictive
motor patterns in the acutely spinalized marmoset monkey (Callithrix jacchus). Exp
Brain Res. 1998;122:351–61.

3. LIDDELL EGT, PHILLIPS CG. PYRAMIDAL SECTION IN THE CAT. Brain. 1944;67:1–9.

4. Shik ML, Orlovsky GN. Neurophysiology of locomotor automatism. Physiol Rev.
1976;56:465–501.

5. Duysens J, Van de Crommert HWAA. Neural control of locomotion; Part 1: The central
pattern generator from cats to humans. Gait Posture. 1998;7:131–41.

6. Muir GD, Whishaw IQ. Complete locomotor recovery following corticospinal tract
lesions: measurement of ground reaction forces during overground locomotion in rats.
Behavioural Brain Research. 1999;103:45–53.

7. Dietz V. Human neuronal control of automatic functional movements: interaction
between central programs and afferent input. Physiol Rev. 1992;72:33–69.

8. Pribe C, Grossberg S, Cohen MA. Neural control of interlimb oscillations. Biol Cybern.
1997;77:141–52.

9. KNUTSSON E, RICHARDS C. DIFFERENT TYPES OF DISTURBED MOTOR CONTROL IN
GAIT OF HEMIPARETIC PATIENTS. Brain. 1979;102:405–30.

10. Barthélemy D, Grey MJ, Nielsen JB, Bouyer L. Involvement of the corticospinal tract in
the control of human gait. 2011. p. 181–97.

11. Bruijn SM, van Dieën JH. Control of human gait stability through foot placement. J R
Soc Interface. 2018;15:20170816.

12. Siebner HR, Funke K, Aberra AS, Antal A, Bestmann S, Chen R, et al. Transcranial
magnetic stimulation of the brain: What is stimulated? – A consensus and critical
position paper. Clinical Neurophysiology. 2022;140:59–97.

13. Di Lazzaro V, Rothwell JC. Corticospinal activity evoked and modulated by non-
invasive stimulation of the intact human motor cortex. J Physiol. 2014;592:4115–28.

14. Groppa S, Schlaak BH, Münchau A, Werner-Petroll N, Dünnweber J, Bäumer T, et al.
The human dorsal premotor cortex facilitates the excitability of ipsilateral primary motor
cortex via a short latency cortico-cortical route. Hum Brain Mapp. 2012;33:419–30.

15. Bestmann S, Krakauer JW. The uses and interpretations of the motor-evoked
potential for understanding behaviour. Exp Brain Res. 2015;233:679–89.

16. Hupfeld KE, Swanson CW, Fling BW, Seidler RD. TMS-induced silent periods: A
review of methods and call for consistency. J Neurosci Methods. 2020;346:108950.

17. Petersen TH, Willerslev-Olsen M, Conway BA, Nielsen JB. The motor cortex drives the
muscles during walking in human subjects. J Physiol. 2012;590:2443–52.

18. Barthelemy D, Nielsen JB. Corticospinal contribution to arm muscle activity during
human walking. J Physiol. 2010;588:967–79.

19. Capaday C, Lavoie BA, Barbeau H, Schneider C, Bonnard M. Studies on the
Corticospinal Control of Human Walking. I. Responses to Focal Transcranial Magnetic
Stimulation of the Motor Cortex. J Neurophysiol. 1999;81:129–39.

20. Schubert M, Curt A, Jensen L, Dietz V. Corticospinal input in human gait: modulation
of magnetically evoked motor responses. Exp Brain Res. 1997;115:234–46.

21. Roelker SA, Kautz SA, Neptune RR. Muscle contributions to mediolateral and
anteroposterior foot placement during walking. J Biomech. 2019;95:109310.

22. Neptune RR, McGowan CP. Muscle contributions to frontal plane angular
momentum during walking. J Biomech. 2016;49:2975–81.

23. Hof AL. The equations of motion for a standing human reveal three mechanisms for
balance. J Biomech. 2007;40:451–7.

24. Bauby CE, Kuo AD. Active control of lateral balance in human walking. J Biomech.
2000;33:1433–40.

25. O'Connor SM, Kuo AD. Direction-Dependent Control of Balance During Walking and
Standing. J Neurophysiol. 2009;102:1411–9.

26. Rankin BL, Buffo SK, Dean JC. A neuromechanical strategy for mediolateral foot
placement in walking humans. J Neurophysiol. 2014;112:374–83.

27. Hof AL, Duysens J. Responses of human hip abductor muscles to lateral balance
perturbations during walking. Exp Brain Res. 2013;230:301–10.

28. Wang Y, Srinivasan M. Stepping in the direction of the fall: the next foot placement
can be predicted from current upper body state in steady-state walking. Biol Lett.
2014;10:20140405.

29. van Leeuwen AM, van Dieën JH, Daffertshofer A, Bruijn SM. Active foot placement
control ensures stable gait: Effect of constraints on foot placement and ankle moments.
PLoS One. 2020;15:e0242215.

30. Hurt CP, Rosenblatt N, Crenshaw JR, Grabiner MD. Variation in trunk kinematics
influences variation in step width during treadmill walking by older and younger adults.
Gait Posture. 2010;31:461–4.

31. Schrager MA, Kelly VE, Price R, Ferrucci L, Shumway-Cook A. The effects of age on
medio-lateral stability during normal and narrow base walking. Gait Posture.
2008;28:466–71.

32. Osoba MY, Rao AK, Agrawal SK, Lalwani AK. Balance and gait in the elderly: A
contemporary review. Laryngoscope Investig Otolaryngol. 2019;4:143–53.

33. Maki BE. Gait Changes in Older Adults: Predictors of Falls or Indicators of Fear? J Am
Geriatr Soc. 1997;45:313–20.

34. Brauer SG, Burns YR, Galley P. A Prospective Study of Laboratory and Clinical
Measures of Postural Stability to Predict Community-Dwelling Fallers. J Gerontol A Biol
Sci Med Sci. 2000;55:M469–76.

35. Nandi T, Fisher BE, Hortobágyi T, Salem GJ. Increasing mediolateral standing sway is
associated with increasing corticospinal excitability, and decreasing M1 inhibition and
facilitation. Gait Posture. 2018;60:135–40.

36. Rossi S, Hallett M, Rossini PM, Pascual-Leone A. Screening questionnaire before
TMS: An update. Clinical Neurophysiology. 2011;122:1686.

37. Dingwell JB, Cusumano JP. Nonlinear time series analysis of normal and pathological
human walking. Chaos: An Interdisciplinary Journal of Nonlinear Science. 2000;10:848–
63.

38. Dingwell JB, Marin LC. Kinematic variability and local dynamic stability of upper body
motions when walking at different speeds. J Biomech. 2006;39:444–52.

39. Rosenstein MT, Collins JJ, De Luca CJ. A practical method for calculating largest
Lyapunov exponents from small data sets. Physica D. 1993;65:117–34.

40. Stenum J, Bruijn SM, Jensen BR. The effect of walking speed on local dynamic
stability is sensitive to calculation methods. J Biomech. 2014;47:3776–9.

41. Bruijn SM. Local Dynamic Stability. Zenodo. 2017;

42. Afschrift M, Bruijn SM, Dieën JH van. ASSESSMENT OF STABILIZING FEEDBACK
CONTROL OF WALKING, A TUTORIAL. bioRxiv [Internet]. 2023;2023.09.28.559516.
Available from: http://biorxiv.org/content/early/2023/09/29/2023.09.28.559516.abstract

43. van Leeuwen J, Smeets JBJ, Belopolsky A V. Forget binning and get SMART: Getting
more out of the time-course of response data. Atten Percept Psychophys.
2019;81:2956–67.

44. McAndrew PM, Dingwell JB, Wilken JM. Walking variability during continuous
pseudo-random oscillations of the support surface and visual field. J Biomech.
2010;43:1470–5.

45. Onushko T, Boerger T, Van Dehy J, Schmit BD. Dynamic stability and stepping
strategies of young healthy adults walking on an oscillating treadmill. PLoS One.
2019;14:e0212207.

46. Bruijn SM, Bregman DJJ, Meijer OG, Beek PJ, van Dieën JH. Maximum Lyapunov
exponents as predictors of global gait stability: A modelling approach. Med Eng Phys.
2012;34:428–36.

47. Toebes MJP, Hoozemans MJM, Furrer R, Dekker J, van Dieën JH. Local dynamic
stability and variability of gait are associated with fall history in elderly subjects. Gait
Posture. 2012;36:527–31.

48. Mahaki M, van Leeuwen AM, Bruijn SM, van der Velde N, van Dieën JH. Mediolateral
foot placement control can be trained: Older adults learn to walk more stable, when
ankle moments are constrained. PLoS One. 2023;18:e0292449.

49. Vlutters M, Van Asseldonk EHF, van der Kooij H. Foot Placement Modulation
Diminishes for Perturbations Near Foot Contact. Front Bioeng Biotechnol. 2018;6.

50. Afschrift M, Pitto L, Aerts W, van Deursen R, Jonkers I, De Groote F. Modulation of
gluteus medius activity reflects the potential of the muscle to meet the mechanical
demands during perturbed walking. Sci Rep. 2018;8:11675.

51. Perry JA, Srinivasan M. Walking with wider steps changes foot placement control,
increases kinematic variability and does not improve linear stability. R Soc Open Sci.
2017;4:160627.

52. Wu M, Matsubara JH, Gordon KE. General and Specific Strategies Used to Facilitate
Locomotor Maneuvers. PLoS One. 2015;10:e0132707.

53. Hak L, Houdijk H, Steenbrink F, Mert A, van der Wurff P, Beek PJ, et al. Stepping
strategies for regulating gait adaptability and stability. J Biomech. 2013;46:905–11.

54. Camus M, Pailhous J, Bonnard M. On-line flexibility of the cognitive tuning of
corticospinal excitability: A TMS study in human gait. Brain Res. 2006;1076:144–9.

55. Pandy MG, Lin Y-C, Kim HJ. Muscle coordination of mediolateral balance in normal
walking. J Biomech. 2010;43:2055–64.

56. MacKinnon CD, Winter DA. Control of whole body balance in the frontal plane during
human walking. J Biomech. 1993;26:633–44.

57. Neptune RR, McGowan CP. Muscle contributions to whole-body sagittal plane
angular momentum during walking. J Biomech. 2011;44:6–12.

58. Anderson FC, Pandy MG. Individual muscle contributions to support in normal
walking. Gait Posture. 2003;17:159–69.

59. van Leeuwen AM, van Dieën JH, Daffertshofer A, Bruijn SM. Ankle muscles drive
mediolateral center of pressure control to ensure stable steady state gait. Sci Rep.
2021;11:21481.

60. Darling WG, Wolf SL, Butler AJ. Variability of motor potentials evoked by transcranial
magnetic stimulation depends on muscle activation. Exp Brain Res. 2006;174:376–85.

61. Hof AL, Duysens J. Responses of human ankle muscles to mediolateral balance
perturbations during walking. Hum Mov Sci. 2018;57:69–82.

62. Koch M, Eckardt N, Zech A, Hamacher D. Compensation of stochastic time-
continuous perturbations during walking in healthy young adults: An analysis of the
structure of gait variability. Gait Posture. 2020;80:253–9.

63. Perera CK, Gopalai AA, Ahmad SA, Gouwanda D. Muscles Affecting Minimum Toe
Clearance. Front Public Health. 2021;9.

64. Ciunelis K, Borkowski R, Błażkiewicz M. The Impact of Induced Acceleration
Perturbations in Selected Phases of the Gait Cycle on Kinematic and Kinetic Parameters.
Applied Sciences. 2024;14:4849.

65. Kiers L, Cros D, Chiappa KH, Fang J. Variability of motor potentials evoked by
transcranial magnetic stimulation. Electroencephalography and Clinical
Neurophysiology/Evoked Potentials Section. 1993;89:415–23.

66. Cuypers K, Thijs H, Meesen RLJ. Optimization of the Transcranial Magnetic
Stimulation Protocol by Defining a Reliable Estimate for Corticospinal Excitability. PLoS
One. 2014;9:e86380.

67. De Luca CJ. The Use of Surface Electromyography in Biomechanics. J Appl Biomech.
1997;13:135–63.

68. Wong AL, Haith AM, Krakauer JW. Motor Planning. The Neuroscientist. 2015;21:385–
98.

69. Kennefick M, Burma JS, van Donkelaar P, McNeil CJ. Corticospinal excitability is
enhanced while preparing for complex movements. Exp Brain Res. 2019;237:829–37.

Figures and legends

Figure 1

Figure 1. *The TMS-helmet with a built-in custom made bent batwing coil.*

614   Figure 2

Figure 2. *Calculation of the ongoing EMG and absolute MEPs. The blue dotted lines indicate the windows over which the data was rectified and averaged to calculate the ongoing EMG. The red dotted lines indicate the window over which the peak-to-peak value was calculated indicating the absolute MEP*

627   Figure 3

Figure 3. *Illustration of the process of weighted smoothing for data from one condition of one participant. A) Stimulations and how the number of stimulation was linked to their gait cycle percentage. B) Display of how for each time sample the weight was distributed. Repeating this over the new temporal axis set at a chosen temporal resolution results in a general weight distribution based on sample density. C) The smoothed results from convolving the weight distribution with the stimulation amplitude.*

639   Figure 4

Figure 4. *Smoothing differences between the Von Mises and Gaussian distributions. **A)** Differences at timepoints 0 and 50.0% for the probability density function (PDF) of both the Von Mises and Gaussian distribution. **B)** Effect of sample density on temporal weights and how this differs between linear and circular distributions. The sample density is displayed in black bars with the amount of samples per 0.5%. The amount if equal to the y-axis times 1000   **C)** Effect of the difference in weight distribution on the smoothed data outcome.*

647   Figure 5

Figure 5. The effect of lateral destabilisation on step width, step width variance, stride duration, stride duration variance, and the local divergence exponent (λs). The dots represent the outcomes of individual participants, connected between conditions by black dotted lines to show the changes for each participant.

656   Figure 6

Figure 6. A) fisher transformed explained variance of the foot placement as a function of step cycle. A step is defined from toe-off to subsequent heel strike. The coloured shaded area indicates the standard deviation of the corresponding condition. The black lines and grey shaded area indicate significant difference B) The average foot placement error (indicating how precise the actual foot placement is relative to the predicted foot placement) was increased during destabilised walking.

668   Figure 7

Figure 7. The amount samples at each temporal point (i.e. the amount of samples per 0.5%) summed over all participants. Purple is seen due to the overlapping bars of both conditions. The dotted lines indicate the normalised weight at each timepoint. This was calculated based on the sample density, a perfect homogeneous data spread would result in a flat line at 0.0050.

682   Figure 8

Figure 8. *Ongoing EMG's (uV) as function of the gait cycle for all muscles. 0%*
corresponds to right heel strike. The coloured shaded area's are the standard deviations of
the conditions. The black lines and with the grey shade around it indicate significant
difference. M. gastrocnemius medialis (MG), m. gastrocnemius lateralis (LG), m. rectus
femoris (RF), m. biceps femoris (BF), m. adductor magnus (AM)

687   Figure 9

Figure 9. *Absolute MEPs(uV) as function the gait cycle for all muscles. 0% corresponds to right heel strike. The coloured shaded area's are the standard deviations of the conditions. The black lines and with the grey shade around it indicate significant difference. M. gastrocnemius medialis (MG), m. gastrocnemius lateralis (LG), m. rectus femoris (RF), m. biceps femoris (BF), m. adductor magnus (AM)*

693 Figure 10

Figure 10. *Normalised MEPs are displayed over the gait cycle for all muscles. The plot starts at 0% of the gait cycle which corresponds to the right heel strike. The coloured shaded area's are the standard deviations of the conditions. The black lines and with the grey shade around it indicate significant difference. M. gastrocnemius medialis (MG), m. gastrocnemius lateralis (LG), m. rectus femoris (RF), m. biceps femoris (BF), m. adductor magnus (AM)*