ECD, a novel androgen receptor target promotes prostate cancer tumorigenesis by regulating glycolysis

Mohsin Raza[1#], Asher Rajkumar Rajan[1#], Benjamin B. Kennedy[1], Timothy E. Reznicek[1], Farshid Oruji[1], Sameer Mirza[1$], M Jordan Rowley[1], Glen Kristiansen[5], Kaustubh Datta[2@], Bhopal C. Mohapatra[1*], Hamid Band[1, 3, 4*], and Vimla Band[1, 4*]

Departments of [1]Genetics, Cell Biology and Anatomy, [2]Department of Biochemistry and Molecular Biology, [3]Eppley Institute for Research in Cancer, and [4]Fred & Pamela Buffett Cancer Center, University of Nebraska Medical Center, 985805 Nebraska Medical Center, Omaha, NE and [5]Institute for Pathology, University Hospital Carl-Gustav-Carus, University of Technology, Dresden, Fetscherstrasse 74, D-01307 Dresden, Germany.

#Both authors contributed equally to this manuscript.

***Corresponding authors**: Vimla Band, PhD, Department of Genetics Cell Biology & Anatomy, University of Nebraska Medical Center, 985805 Nebraska Medical Center, Omaha, NE, 68198, USA. vband@unmc.edu; Phone: 402-559-8565. Hamid Band, MD, PhD, Eppley Institute for Research in Cancer, and Allied Disease, 985950 Nebraska Medical Center, Omaha, NE 68198-5950, USA; Email: hband@unmc.edu; Phone: 402-559-8572. Bhopal C. Mohapatra, Department of Genetics, Cell Biology, and Anatomy, University of Nebraska Medical Center 985805 Nebraska Medical Center, Omaha, NE 68198. Phone: 402-559-8543; E-mail: bmohapat@unmc.edu.

Current Addresses: [$]Department of Chemistry, College of Science, UAEU University, Al Ain, Abu Dhabi, U. A. E. [@] Department of Biochemistry and Molecular Biology, Massy Cancer Center, Virginia Commonwealth University, Richmond, VA, United States.

Running Title: ECD, an AR target gene, regulates glycolysis to promote PC oncogenesis.

Keywords: Androgens, androgen receptor, ecdysoneless (ECD), glycolysis, prostate cancer, oncogenesis, enzalutamide

Abstract:

Androgen receptor (AR)-mediated signaling is essential for PC tumorigenesis. In TCGA database we observed a positive correlation between ECD and AR expression. Consistently, Dihydrotestosterone (DHT) treatment of PC cell lines increased ECD mRNA and protein levels, and AR knockdown (KD) reduced ECD expression. Bioinformatic analysis predicted three consensus androgen response elements in the ECD promoter, and DHT treatment increased AR occupancy at the ECD promoter, and enhanced ECD promoter activity. Enzalutamide treatment decreased ECD levels, and ECD knockout (KO) in PC cells reduced oncogenic traits, suggesting a functional role of ECD to maintain PC oncogenesis. ECD mRNA and protein are overexpressed in PC patient tissues, and its overexpression predicts shorter survival. Overexpression of ECD in PC cell lines enhanced the oncogenic traits *in vitro* and developed faster and larger highly proliferative xenograft tumors. RNA-seq analysis of mouse tumors revealed increase mRNA levels of several glycolytic genes. ECD associates with mRNA of several key glycolytic genes and is required for their stability, consistent with our recent demonstration of ECD as an RNA binding protein. Higher glucose uptake and glycolysis was seen upon ECD OE in PC cells. Together, we demonstrate role of a novel AR target gene ECD in PC tumorigenesis.

Introduction:

Prostate cancer (PC) is the second leading cause of cancer related adult males deaths [1]. Androgen receptor (AR) mediated signaling activated through binding of androgens is essential for PC tumorigenesis [2]. Initial androgen deprivation therapy is invariably followed by development of castration resistant PC (CRPC) [3]. AR inhibitors have proven effective but resistance to these inhibitors invariably develops within 2-3 years, even though most such patients retain a functional AR that sustains tumor growth and spread [4, 5]. Thus, understanding of novel molecular pathways that sustain the AR-driven PC tumorigenesis could open new approaches to therapy of AR-dependent PC.

Tumors, including PC, exhibit metabolic reprogramming to cope with increased macromolecular synthetic, redox and energetic demands for increased oncogenic traits [6]. A hallmark of cancer cells is the Warburg effect, which involves a switch to aerobic glycolysis instead of oxidative phosphorylation [7]. Increase in the rate of glycolytic flux has been reported during the late stage of high-grade primary PC, as well as CRPC, and PC with neuroendocrine differentiation [8, 9]. In addition, PC has been shown to maintain a unique androgen regulated metabolic profile with high rates of lipogenesis as well as oxidative phosphorylation [10]. While the role of glycolysis is well accepted across cancers, mechanisms that promote glycolysis in AR-dependent PC remain an area of active investigation. A unique property of prostate epithelial cells is that glycolytic metabolism sustains physiological citrate secretion and in early stages of PC, citrate is needed for oxidative phosphorylation and lipogenesis for tumor progression, however, in advanced and incurable CRPC, a metabolic shift towards choline, amino acid, and glycolytic metabolism takes over tumor progression [11]. Thus, inhibiting glycolysis in advanced PC provides a unique opportunity. Here, we identify ECD as a novel link between AR-dependent PC

tumorigenesis and modulation of glycolysis through a novel RNA binding role of ECD to stabilize

mRNAs of key glycolytic genes.

ECD is the highly conserved mammalian orthologue of *Drosophila* ecdysoneless (ECD)

gene [12]. *Ecd* in essential for mouse embryonic development and deletion of *Ecd* in *Ecd^flox/flox*

mouse embryonic fibroblasts (MEFs) led to G1 cell cycle arrest [13]. ECD plays a role in

mitigating endoplasmic reticulum stress through upregulation of GRP78 [14]. We and others have

shown that ECD interacts with several components of the mRNA spliceosome machinery, and

participates in pre-mRNA processing in both drosophila and mammalian cells [15, 16] as well as

in mRNA export in mammalian cells [17]. Our [18, 19] and others' studies have established that

ECD mRNA and protein are overexpressed in various solid cancers, and that ECD OE plays a

critical role in promoting tumorigenesis [20, 21]. Importantly, transgenic ECD OE targeted to

mouse mammary epithelium led to mammary epithelial hyperplasia and tumorigenesis [22].

However, role of ECD in PC tumorigenesis have remained unexplored until this study.

Here, we establish that ECD is a novel AR target gene, and its' expression correlates with

AR expression. ECD expression is regulated by AR through androgen-dependent AR recruitment

to the ECD promoter and androgen treatment leads to increased ECD promoter activity.

Enzalutamide treatment decreases ECD levels, and ECD KO decreases oncogenic traits,

suggesting a key role of ECD in PC oncogenesis. ECD mRNA and protein are overexpressed in

PC patient tissues, with ECD OE predicting short survival in PC patients. ECD OE enhanced *in*

vitro oncogenic traits as well as *in vivo* tumorigenicity. Using RNA-seq analysis of xenograft

tumors, RNA immunoprecipitation analyses, and functional assays we demonstrate that ECD binds

to and stabilizes the mRNAs of key glycolytic genes to promote glycolysis. Thus, our studies identify

a novel AR target gene ECD and its function in promoting PC oncogenesis through regulation of

glycolysis.

Results:

ECD expression in prostate cancer cells is regulated by androgen receptor.

All PC are initially androgen-driven [23] and most remain dependent on androgen receptor

signaling even after castration-resistant [24], Given the connection of AR in PC, we analyzed

TCGA-prostate adenocarcinoma dataset and observed a significant positive correlation between

AR and ECD mRNA expression **(Fig. 1A)**. Consistently, DHT dose-response **(Fig. 1B-E)**, as well

as time-course analyses **(Fig. 1F-I)** in two different cell lines LNCaP and C4-2B showed an

increase in ECD protein and mRNA levels occurred rapidly, plateaued at later time points and

decreased towards the end of the time course (more pronounced in C4-2B) **(Fig. 1C & 1G)**.

Further, the basal as well as DHT-induced increase in ECD protein levels were abrogated by

siRNA KD of AR, which was confirmed by anti-AR immunoblotting LNCaP **(Fig. 1J)** and C4-2B

(Fig. 1K), with or without the addition of exogenous DHT. These results show that ECD

expression in PC cell models is positively regulated by androgens and dependent on the AR,

suggesting the likelihood that ECD is transcriptionally regulated by AR.

AR binds to the ECD promoter to regulate ECD expression, and Enzalutamide treatment

decreases ECD expression.

Both androgens activated and constitutively active ARs function by binding to gene

promotors through defined androgen response elements (AREs) to regulate transcription of

downstream target genes [25]. Given that DHT induction, combined with AR dependence of *ECD*

mRNA expression, we examined the ECD promoter region for potential AR binding sites, using

three independent prediction tools, Pscan [26], EPD motif search tool [27] and de novo motif discovery algorithm Motif Based Sequence analysis tool [28]. These analyses revealed three potential androgen response elements (AREs) in the ECD promoter: nucleotides -698 to - 682 (site 1); -657 to -641 (site 2); and -640 to -624 (site 3) **(Fig. 2A & B)**. Indeed, chromatin immunoprecipitation (ChIP) assays demonstrated specific occupancy of AR at the ECD promoter (-712 to -565, encompassing all three sites) in both LNCaP and C4-2B PC cell lines compared to IgG control, with the occupancy significantly increased upon DHT treatment **(Fig. 2C & D)**. Next, we cloned a ~1,500 bp region encompassing the ECD promotor region in a reporter construct upstream of secreted Gaussia Luciferase. PC cells transduced with the ECD promotor Luciferase construct were left untreated or treated with DHT for 3 h, and secreted Gaussia Luciferase and alkaline phosphatase (SEAP; internal control) activities were measured in the media. Notably, compared to basal reporter activity (Ctrl; first bar), DHT treated cells expressing the Full length ECD promoter showed about 2.5-fold increase in luciferase activity **(Fig. 2E & F, lanes 2)**. Importantly, constructs with deletion of AREs either alone or in combination showed progressive diminution of luciferase activity while the construct with all three AREs deleted showed activity close to that of the control **(Fig. 2E & F lanes 3 to 9).** Collectively, our results identify ECD is an AR-regulated gene, with three functional ARE elements in its promoter region that mediate androgen-induced AR-dependent ECD expression.

Based on a published report GSE 48403 [29] showed ECD mRNA expression was decreased post-androgen-deprivation therapy as compared to pre-treatment **(Fig. 2G).** Thus, we treated LNCaP and C4-2B cell lines with different concentrations of Enzalutamide in the presence of DHT. Notably, Enzalutamide treatment dose-dependently decreased ECD protein levels in both cell lines **(Fig. 2H and 2I)**. Expected decrease in AR and PSA, an AR target gene was observed

(Fig. 2H and 2I). Enzalutamide induced reduction in ECD is at the transcriptional levels **(Fig. 2J**

and 2K). Together, these results conclusively demonstrate ECD is a novel AR target gene.

ECD expression is required to sustain oncogenic traits of prostate cancer cells.

Next, we engineered the androgen-dependent LNCaP and androgen-independent C4-2B

cell lines for doxycycline (Dox)-inducible CRISPR knockout with two distinct guide RNAs

(gRNA1 and gRNA2) against ECD. Dox-inducible depletion of ECD in gRNA1/2 expressing cell

lines compared to the non-targeted control (Cas9-NTC) was observed **(Fig. 3A).** Analysis of

colony forming capacity as a measure of cell growth demonstrated that ECD depletion led a

significant decrease in PC cell proliferation compared to NTC **(Fig. 3B & C)**. Further, ECD

depletion in comparison with NTC exhibited a significant decrease in oncogenic traits, such as

tumorsphere forming ability **(Fig. 3D & E),** a measurement of the stem cell ability of tumor cell

lines [30], anchorage independent growth using the soft agar colony forming assay (**Fig. 3F & G**),

three-dimensional growth in Matrigel **(Fig. 3H & I),** cell migration **(Fig. 3J & K),** as well as tumor

cell invasion **(Fig. 3L & M).** Taken together, these results establish a critical role of ECD

expression in sustaining the oncogenic traits of PC cells.

ECD protein is overexpressed in prostate cancer tumor tissues and its overexpression

predicts shorter patient survival.

To assess the expression levels of ECD in primary PC tumor samples, we carried out IHC

analysis of a PC tissue microarray (TMAs) comprising of 522 patient samples with available

clinical, pathological and survival data using a previously validated anti-ECD antibody [13]. PC

samples showed a range of ECD IHC staining. Based on the intensity of staining, the patients were

grouped into ECD-low (staining intensity of 0 or 1) and ECD-high (staining intensity of 2 or 3)

groups; representative examples of low IHC staining of hyperplastic tissue, and PC tumor samples vs.

high ECD IHC staining are presented in (**Fig. 4A & B**). Overall, ECD protein was overexpressed

in tumor samples of ~40% of PC patients **(Fig. 4C)**. Kaplan Meier analysis showed that ECD OE

correlated with shorter overall survival (p=0.024) **(Fig. 4C)**. Multivariate survival analysis using

Cox's regression model independently confirmed the ECD OE as a poor prognostic marker, with

a significantly elevated relative risk and confidence interval, independent of Gleason grade and

positive surgical resection margins **(Table S1).** In addition, analysis of two publicly available

mRNA expression datasets (GSE32269 & GSE35988) revealed *ECD* mRNA overexpression in

PC tissues as compared to normal prostate **(Fig. S1A & B).** Kaplan-Meier analysis of The Cancer

Genome Atlas (TCGA) dataset of 484 mixed prostate adenocarcinoma samples (2022-v32) using

the R2 Genomics platform showed that high ECD mRNA was associated with shorter patient

survival **(Fig. 4D)**. Collectively, our results show that ECD mRNA and protein levels are increased

in a significant subset of PC tumor samples, and that ECD OE correlates with negative prognostic

features and shorter survival of patients, suggesting that ECD OE may promote PC tumorigenesis.

ECD overexpression in PC cells promotes the *in vitro* oncogenic traits and *in vivo*

tumorigenesis.

Next, we established stable ECD-OE PC cell lines LNCaP (hormone sensitive) and its

derivatives C4-2B (hormone insensitive). Overexpression of ECD is observed in both LNCaP **(Fig.**

5A) and C4-2B **(Fig. S2A).** While there were no significant differences in proliferation of vector

vs ECD-OE PC cell lines when measured in two-dimensional culture on plastic either in the

absence or presence of DHT **(Fig. S2B & C),** a significant increase in anchorage independent

growth in soft agar **(Fig. 5B & C & Fig. S2D & E)** as well as in three-dimensional Matrigel

growth **(Fig. 5D & E & Fig. S2F & G)** was seen in ECD-overexpressing vs. vector control

LNCaP and C4-2B cells. Notably, ECD-OE LNCaP cells remained hormone sensitive like control

cells **(Fig. S2H).**

To assess the impact of ECD OE on *in vivo* oncogenesis of PC cells, we introduced a

mCherry-luciferase reporter into vector control and ECD-overexpressing LNCaP cells, and

subcutaneously injected these in the dorsal flank region of 8-weeks old male athymic nude mice.

Tumor growth was monitored twice a week using calipers and weekly using bioluminescence

imaging. Compared to the time course of tumor growth in mice implanted with control cells, those

implanted with ECD– overexpressing cells developed tumors that appeared earlier and grew to

be significantly larger (**Fig. 5F-J, Table S2).** IHC staining of tumor sections confirmed the

overexpression of ECD in ECD-OE tumors **(Fig. 5K)**. H&E-stained tumor sections showed that

ECD-OE tumors had more necrosis **(Fig. 5K)**, a characteristic of aggressive tumors [31]. Finally,

ECD-OE tumors showed increased Ki-67 staining **(Fig. 5K)**, a marker of tumor cell proliferation

[32]. Together, the combination of *in vitro* and *in vivo* analyses demonstrates that ECD OE

enhances the tumorigenic behavior of PC cells.

ECD overexpression in prostate cancer cells upregulates the glycolysis pathway.

To explore how ECD OE promotes the tumorigenic behavior of PC cells, we conducted

RNA-seq analysis of xenograft tumors formed by ECD-overexpressing vs. Vector-expressing

LNCaP cells followed by bioinformatics analyses. Principal component analysis (PCA)

demonstrated the distinct clusters of the four each of the two types of tumors **(Fig. 6A**). Based on

a log fold change >1 and false discovery rate (FDR) of <0.05, we found 520 differentially

expressed genes (DEGs) in ECD-overexpressing compared to vector control tumors. Gene set

enrichment analysis (GSEA) using the hallmark gene sets showed a significant upregulation of

hypoxia, glycolysis, angiogenesis, myogenesis, apical junctions, TNFα signaling via NFkB and

mTORC1 signaling in ECD-OE tumors compared to vector control tumors (**Fig. 6B & C**). These

signaling pathways are well-known to be upregulated in aggressive prostate cancers [33-35].

Kaplan-Meier plot for survival curve of prostate cancer patients shows short survival in patients with tumors

expressing high *ECD* mRNA (n=283) in comparison to low *ECD* mRNA (n=201). P-value is indicated in

the KM plot. Among the genes upregulated in ECD-OE LNCaP tumors were key glycolytic

pathway genes (*ALDOA, LDHA, PKM, PGK1, HK1, HK2, ENO1*), Epithelial to Mesenchymal

Transition (EMT) related genes (*VIM, SNAI1, TWIST2, TGFb1, TGFb2, HMGA2),* invasion and

migration associated genes (*MMP2, MMP13, MMP14, MMP15* and *MMP24*), and genes

associated with Notch signaling (NOTCH1, NOTCH2, NOTCH3 and NOTCH4), FGFR signaling

(SPRY2, FGFR1) [34] and a previously identified aggressive prostate cancer signature gene set

(*CCND2, RNF19A, DPP4* and *PRKDC*) [35] (**Fig. 6D**). Notably, ECD-OE tumors exhibited

increased expression of neuro-endocrine prostate cancer (NEPC) related genes (*SYP, EZH2,*

ENO2, NRP2, NCAM1 and SOX2), a features of aggressive metastatic castration resistant PC

(mCRPC) [36] (**Fig. 6D**). qRT-PCR analysis of mRNA isolated from xenograft tumors and *in vitro*

cultured LNCaP cell lines validated the significant upregulation of glycolytic genes *HK2, LDHA*

and PKM2 (**Fig. 6E-G**) and NEPC related *SYP, NCAM1* and *ENO2* (**Fig. S3A & B**) in ECD-OE

group compared to vector control. Together, these results suggested the upregulation of glycolysis

as a potential mechanism of ECD-promoted oncogenesis, consistent with the previously identified

role of ECD in promoting glycolysis related gene expressions in mammary tumors of *ECD*Tg mice

[22].

185   Given others and our recent findings that ECD regulates various aspects of mRNA

biogenesis [37] and our recent findings that ECD is a RNA binding protein [38], we examined if

ECD directly binds to and regulates glycolytic genes. LDHA and PKM2 were among the ECD

interacting mRNAs identified through our recent analyses using enhanced crosslinking and

immunoprecipitation (eCLIP) assay [38]. Using an RNA immunoprecipitation (RIP) assay, we

found that *LDHA*, *PKM2* and *HK2* mRNAs were indeed associated with ECD in both LNCaP (**Fig.**

7A) and C42-B (**Fig. 7B**) cell lines. Next, we analyzed the stability of these mRNAs in control vs.

ECD-overexpressing cells by determining mRNA half-lives following treatment with

actinomycin-D. These analyses revealed significantly longer (1.5 to 2-fold) half-lives of *PKM2*,

HK2 and *LDHA* in ECD-overexpressing vs. control LNCaP (**Fig. 7C-E**) and C4-2B (**Fig. 7F-H**)

PC cells, determining that ECD binds to and stabilizes the mRNAs of key glycolytic genes.

To functionally link the upregulation of glycolytic gene mRNA levels upon ECD OE to

glycolysis, a Warburg effect [39], we measured glucose uptake and glycolytic rates. Indeed, a

significant increase in glucose uptake in ECD-OE LNCaP (**Fig. 8A**) and C4-2B (**Fig. 8B**) cell lines

as compared to vector expressing cells. To directly assess the impact on glycolysis, we conducted

Seahorse XFe glycolytic rate assays to determine the proton efflux rate (PER) from glycolysis

(GlycoPER) and extracellular acidification rate (ECAR) as readouts of glycolysis. Notably, the

PER (**Fig. 8C & D**) and ECAR (**Fig. 8E & F**), measured in the presence of rotenone (Rot) and

antimycin A (AA) to inhibit mitochondrial respiration, were increased in ECD-OE cells compared

to vector cells. Additionally, ECD-OE cells showed increased levels of basal glycolysis (**Fig. 8G**

& H) as well as of glycolytic reserve/compensatory glycolysis (**Fig. 8I & J**) after adding 2-Deoxy-

D-glucose (2-DG) compared to their vector controls. Further, cellular oxygen consumption rate

(OCR) was increased in ECD-OE PC cells relative to their controls (**Fig. S4A & B**). These results

demonstrate that ECD OE in PC cells increases the expression of key glycolytic genes through

binding to and stabilization of their mRNAs to promote the Warburg effect.

Discussion:

While androgen signaling through AR is universally accepted as an early driver of prostate

tumorigenesis with well-established targeting agents, androgen and AR directed therapies

eventually fail in most patients [40]. Metabolic reprogramming represents a key molecular

determinant of the aggressive behavior of PC and therapeutic resistance, with enhanced glucose

utilization through aerobic glycolysis (Warburg effect) a major factor [39]. Molecular pathways

that link AR signaling and glycolytic metabolism in PC are of interest as these are potential

biomarkers, as well as for therapeutic targeting in conjunction with AR directed therapies. Here,

we identify the Ecdysoneless (ECD) protein as a novel link between AR signaling and promotion

of glycolysis and demonstrate that ECD, which is overexpressed in >40% of PC patient tumors

and predicts shorter survival, is an AR-responsive target gene whose increased expression

promotes glycolysis through ECD binding to glycolytic gene mRNAs to enhance their stability.

Our studies elucidate a novel mechanism to promote metabolic reprogramming and tumorigenesis

in PC and raise the possibility of glycolysis pathway co-targeting with AR in ECD-OE PC.

AR-mediated signaling is a critical element of the early progression of all PCs and remains

an oncogenic driver in most late-stage PCs [48]. Our comprehensive analyses establish that ECD

OE is directly regulated by AR signaling. We found a strong positive correlation between AR and

ECD mRNA expression levels across PCs (TCGA-PRAD dataset) **(Fig. 1A)** and cell-based

analyses of PC cell lines showed a dose- and time-dependent upregulation of ECD mRNA and

protein levels by exogeneous androgen treatment of PC cells **(Fig. 1B-I)** with siRNA KD

demonstrating a requirement of AR for basal as well as androgen-stimulated ECD expression **Fig.**

1J & K),. Direct evidence for ECD as a novel androgen-responsive AR target gene in PC was

provided by ChIP **(Fig. 2A-D)** that established the potential AREs identified by in-silico analysis

of the ECD promoter to be authentic AR binding sites and mutational analyses of the three AREs

using a luciferase reporter assay **(Fig. 2E & F)** demonstrated their combined requirement for

androgen-induced ECD expression. Consistently, treatment of PC cell lines with anti-androgen

Enzalutamide decreased ECD levels **(Fig. 2H-K)**. Further the role of ECD in PC was established

by a genetic loss as well as gain of function approaches in PC cell line models. Inducible ECD KO

led to a marked reduction in key in vitro oncogenic traits **(Fig. 3A-M)**.

While ECD plays a physiological role in cell cycle progression and ER stress mitigation,

and its overexpression correlates with poor prognosis and shorter survival in breast, pancreatic

and gastric cancers [20, 21], any role of ECD in PC has remained unexplored.   Our IHC analyses

of a large cohort of PC patient tumors using a well-authenticated anti-ECD monoclonal antibody

[41], we found ECD OE in nearly 43% of PC cases and such overexpression specified shorter

patient survival (**Fig**. **4A-C**). Analysis of publicly available TCGA PC adenocarcinoma dataset

demonstrating shorter survival of PC patients with higher ECD mRNA expression **(Fig. 4D)** lend

further support to our IHC results. Notably, high ECD OE was associated with key negative

prognosticators of higher Gleason score and positive resection margins **(Table 1)**. Thus, ECD OE

emerges as a novel biomarker of poor prognosis and shorter patient survival in PC.

These results support a novel mechanism whereby AR signaling induces the upregulation

of ECD expression in PC as one pro-oncogenic pathway. It is noteworthy that while we and others

have reported the upregulation of ECD expression in other cancers, mechanisms responsible for

such overexpression remain unknown. Given the androgen/AR mediated upregulation of ECD

expression in PC identified here, it will be of great interest to assess the potential regulation of

ECD OE across cancers through other nuclear hormone receptors, especially in hormonally

regulated malignancies. Notably, the AR requirement for ECD OE was observed in an androgen-

dependent as well as an androgen-independent PC cell line model, suggesting that AR signaling

rather than exogenous androgen requirement is a key determinant of ECD OE in PC. As significant

subsets of CRPCs are AR-independent [49], future studies across AR-dependent and AR-

independent PCs should help determine if ECD is primarily a feature of AR signaling driven PCs

or if ECD is also a player in AR-independent PCs, potentially downstream of other transcriptional

regulators.

263  Besides emerging as a correlative biomarker of aggressive PC, our studies clearly

demonstrate that ECD OE is biologically important in promoting the tumorigenic behavior of PC

cells (**Fig. 5A-E & Fig. S2).** Notably, both vector and ECD-OE PC cells remain hormone

responsive for their proliferation (**Fig. S3**). Importantly, ECD OE imparted a more aggressive *in*

vivo tumorigenic phenotype with earlier growth of xenografted tumors, significantly larger tumors

(**Fig 5F-J**), and higher levels of Ki-67 staining and necrotic lesions indicative of hyper-

proliferative and rapidly growing aggressive tumors (**Fig. 5K**), consistent with known association

of hyperproliferation and necrosis with aggressive prostate tumors in patients [42]. (**Fig. 5L**). Thus,

like the role of ECD OE in other cancers such as breast cancer [22], its overexpression appears to

propel PC towards increased aggressiveness. We note that while our studies in established PC cell

lines clearly reveal the impact of ECD OE to promote the tumorigenic phenotype, our studies do

not address any potential role of ECD OE in early malignant transformation driven by androgen-

AR axis. In our studies using immortalized mammary epithelial cell models, ECD OE alone did

not impart a tumorigenic phenotype [43]. However, since ECD plus mutant RAS co-

overexpression in immortal mammary epithelial cells promoted full tumorigenesis [43], and

transgenic overexpression of ECD in the mammary epithelium led to hyperplasia and late-onset

tumor induction [22], incorporation of ECD transgene or tissue-specific ECD KO genetic models

of prostate tumorigenesis should help assess a potential early pro-tumorigenic role of ECD in PC.

Our mechanistic studies provide a plausible explanation for the pro-oncogenic role of ECD

OE in PC. Our RNA-seq analyses identified a potential role of ECD OE in upregulating the mRNA

levels of key glycolytic genes, *HK2, LDHA* and *PKM2* (**Fig. 6**). Consistent with our recently

established biochemical function of ECD as an RNA-binding protein[38], these mRNAs are

physically associated with ECD **(Fig. 7A & B)** and ECD OE enhances their stability **(Fig. 7C-H).**

A significant increase in glucose uptake (**Fig. 8A & B**) and glycolysis measured with Seahorse

glycolytic assays directly established that the increased expression of glycolytic gene mRNAs

upon ECD OE translates into a biological impact to upregulate glycolysis **(Fig. 8C-J)**. Thus, we

posit that increased expression of ECD in PC, initiated by androgen-AR signaling, serves as one

mechanism to promote a glycolytic switch to enhance the aggressiveness of PC. This mechanism

is consistent with our analyses of ECD transgene driven mammary tumors which also exhibited

an upregulation of glycolysis enzyme mRNAs and increased glucose uptake [22]. A linkage of

ECD OE to increased glycolysis as a mechanism to enhance PC tumor aggressiveness is also

consistent with prior studies that have demonstrated a key role of aerobic glycolysis in promoting

cell survival [44], metastasis [45] and drug resistance in PC [46]. While our studies suggest the

ECD OE as a potential positive regulator of glycolysis in >40% of prostate tumors, clearly other

mechanisms are likely to play a role either concurrently with ECD OE or in patients without ECD

OE. Notably, AR signaling is known to control glycolysis directly in PC, [47] however other

mechanisms like aberrant FGFR tyrosine kinase signaling [48] and activation of Wnt/β-catenin

signaling [49] have also been linked to enhanced glycolysis in PC. Future studies will be needed

to assess how ECD OE may synergize with other mechanisms that promote glycolysis.

In conclusion our work reveals ECD as a novel androgen-regulated AR target gene and

demonstrate that ECD OE, a feature of >40% PC patients, promotes PC tumor aggressiveness

through upregulation of glycolysis. Our findings open the possibility of exploring glycolysis-

targeted therapeutic approaches for PC patients with high ECD expression. These results support

oncogenic role of ECD in both hormone dependent and hormone independent AR-expressing

prostate cancer.

Materials and Methods:

Antibodies and reagents

311   Antibodies: anti-AR (Invitrogen cat # MA5-13426), β-actin (Sigma cat # A5441), Ki67

312   (Abcam cat # ab16667), anti-ECD antibody generated by us has been described previously [13].

313   5α-Dihydrotestosterone solution (Millipore Sigma cat # D-073), Chemiluminescence signals were

314   detected by (ECL Plus; Amersham-Pharmacia cat # NEL-105001EA) and Matrigel (Coring Cat #

315   356230), and Isoflurane (cat # 502017) from MWI Animal Health.

Cell lines and medium.

PC cell lines LNCaP and C4-2B were obtained from the American Type Culture Collection.

Cell lines were maintained in complete RPMI medium [50]. For androgen deprivation

experiments, cells were cultured in phenol red-free complete RPMI medium supplemented with

10% charcoal stripped FBS (cat # S11650 Atlanta Biologicals). Lenti-X 293T cell line used for

lentiviral packaging was from Takara Bio (cat # 632180) and cultured in complete DMEM medium

[50]. Cells were cultured in a humidified atmosphere with 5% CO_2 at 37°C. Cell lines were

regularly evaluated for mycoplasma and continuously cultured for no more than 3 months.

Chromatin Immunoprecipitation assay.

Cells grown in charcoal stripped medium and were treated with DHT for 3 h, washed with

PBS and treated with 1% formaldehyde for 10 min to crosslink protein-DNA complexes. Cells

were rinsed and processed for ChIP with control IgG (Abcam, cat # ab172730) or anti-AR antibody

(Abcam, cat # ab185913). The following PCR Primers were used to determine the enrichment:

ECD AR Binding site Forward primer -TTAATTCTGGGATCTTTATCTGTAG; ECD AR

Binding site Reverse primer- GGCTCCAGGAAAGTGCTCACTTGAC. Real- time PCR

quantification was performed in triplicates using SYBR Green PCR master mix (Applied

Biosystems cat # 4309155). AR Binding percentage was calculated post normalization against

IgG.

Luciferase-promoter reporter assays.

The promoter sequences of GAPDH (Genome=hg38;chr12+: 6533244-6534713;

TSS=6534517; Upstream=1273, Downstream=196; Length=1470), and wildtype (WT) ECD

(Genome=hg38;chr10-:73169381-73167884; TSS=73168095; Upstream=1286,

Downstream=211; Length=1498) or its predicted AR binding site deletion mutants (Details in

Table S3) were synthesized and cloned upstream of the Gaussia luciferase in the lentiviral pEZX-

LvPG04 vector (sequence-confirmed constructs were custom-made by Genecopoeia; cat#

HPRM35342-LvPG04-WT, GAPDH-LvPGO4). The ECD promoter mutants and the sequences

deleted from TSS are listed in **Table S3**. The reporter constructs were introduced into LNCaP and

C4-2B cell lines using lentiviral supernatants. For promotor luciferase experiments, transduced

cells were cultured in charcoal stripped serum containing medium for 5 days and then treated with

100 nM DHT for 3 h. Post treatment, supernatant was collected and analyzed using the Secrete-

Pair Dual Luminescence Assay Kit (cat # LF031) to measure luciferase activity following the

manufacturer's protocol. The fold induction was calculated after normalizing the values against

secreted alkaline phosphatase luminescence, used as an internal control.

RNA isolation and quantitative real-time PCR.

Total RNA isolation and qRT-PCR analysis was conducted in indicated PC cell lines as

described previously [17]. The qRT-PCR primers are listed in the **Table S4**.

Generation of CRISPR-Cas9 mediated inducible ECD Knockout (KO) cell lines.

To generate inducible ECD KO cell lines, Lentiviral supernatants were generated as

mentioned above for Edit-R inducible Lentiviral hEF1a-Blast-Cas9 plasmid (cat # CAS11229) and

Edit-R lentiviral plasmids carrying guide RNAs against non-targeting control (NTC) (cat #

GSG11811), or human ECD gRNA1 (cat # GSGH11838-246505144, Seq-

TCCAAAGTCTCAACCCACAA) or gRNA2 (cat # GSGH11838-246505139, Seq-

CTTGGGTATACTTACCCTAT). All plasmids were procured from horizon discovery. Cells

were transduced first for cas-9 expression, then selected in medium containing 5 µg/ml blasticidin

for 5 days. Inducible Cas-9 expressing cells were then transduced with either NTC or gRNA1/2

lentiviral supernatants and selected in 5 µg/ml blasticidin + 1 µg/ml puromycin for additional 5

days.

Transwell migration and invasion assays:

370    Cell lines were subjected to serum deprivation (cultured in complete RPMI medium with

371  0.5 % FBS) for 24 h, trypsinized and seeded in transwell migration (Corning BioCoat 8.0 μm cat

372  # 354578) or invasion (BioCoat Matrigel 8.0 μm cat # 354480) chambers at 100,000 cells per well

373  in 0.5% FBS containing medium. After three hours of incubation at 37°C, complete RPMI medium

374  containing 10% FBS was added as a chemoattractant in the bottom chambers. After 24 h of

375  incubation, the cells on the top of filters were scrapped off and the migrated cells on the bottom of

376  filters were fixed in ice-cold methanol, stained with crystal violet, and rinsed in PBS. Cells in six

377  random fields per well were imaged at 100x magnification and quantified using the ImageJ

378  software. Experiments was repeated three times independently with triplicates each time.

380  **Anchorage-independent growth assays.**

381    20,000 vector-control or ECD-OE cells in complete RPMI medium containing 0.3%

382  agarose were seeded on top of a 0.6% bottom layer of agarose in 6-well plates, as described [43].

383  Cultures were fed every 2 days. Twenty-one days after cell seeding, the colonies were stained with

384  0.05% crystal violet in 25% methanol. Colonies were imaged under 10x magnification with phase

385  contrast microscope and counted using the image J software. Experiments were performed in

386  triplicates, and each experiment was repeated three times.

388  **Matrigel organoid cultures.**

389    2,000 vector-control or ECD-OE cells resuspended in the medium containing 4%

390  Matrigel were seeded per well on top of a base layer of 100% Matrigel in eight-well Nunc

391  chamber slides. The cells were allowed to form organoid colonies for 6 days and then imaged.

392  The ImageJ software was used for quantification with a size cut-off of 70 μm in diameter. All

393   experiments were performed in triplicate wells and repeated three times.

Prostate cancer patient tissue microarrays and immunohistochemical analysis.

The prostate cancer tissue specimens were obtained with informed consent from patients

who underwent radical prostatectomy at the Department of Urology, Institute for Pathology,

University Hospital "Carl-Gustav-Carus", University of Technology, Dresden between 1999 and

2005. Patient ages ranged between 43 and 74 years (median 62 years). Preoperative PSA levels

ranged from 0.8 to 39 ng/ml (median 7.2). Clinical follow-up data was assessed annually.

The median follow-up time of all cases was 17 months (range 1–68 months). The distribution of

Gleason scores (GS) in the cohort was as follows: GS 2–6: 217 (35.3%) GS 7: 291 (47.4%), GS

8–10: 106 (17.3%). Tissue microarrays (TMAs) composed of 556 cases of paired

normal/hyperplastic prostate and prostate cancer (one core per case) were analyzed for ECD

expression using an anti-ECD monoclonal antibody [13]. The intensity of staining was expressed

as a semiquantitative scoring system: negative = 0, low = 1, moderate = 2 and High =3.

Stratification into ECD-High vs ECD-Low expression status reflects cases with staining intensity

of 0 or 1 vs. 2 or 3, respectively. Kaplan-Meier survival analysis compared 224 ECD-High (2-3)

vs. 298 ECD-Low (< or = 1) cases. All slides were evaluated by a pathologist. Multivariate

survival analysis was calculated using Cox's regression model (**Fig. 4 and Table S1)**. The Charite

approved the use of PC tissue samples by the University Ethics Committee under the protocol

'Retrospective analysis of tissue samples by immunohistochemistry and molecular biological

techniques' (EA1/06/2004) on 20 September 2004.

ECD overexpression in prostate cancer cell lines.

The ECD (GeneID: 11319) CDS in the mammalian expression vector pLenti-C-Myc-

DDK-P2A-Puro and empty Vector control plasmids were obtained from (Origene Biotech cat #

PS100092). Lentiviral supernatants were generated by transient co-transfection of Lenti-X 293T

cells with ECD expression vector and packaging plasmids psPAX2 (Addgene, cat # 12260) and

pMD2.G (Addgene cat # 12259) using X-treme GENE HP DNA transfection reagent (Roche cat

06366236001). Collected lentiviral supernatant was added to PC cells in the presence of

polybrene (8 µg/ml, Sigma # H9268). Post-infection, transduced cells were selected in 1 µg/ml

puromycin for 5 days, and ECD levels assessed by Western blotting using an anti-ECD monoclonal

antibody [13]. For siRNA KD, cells were transfected with 40 nM of control or gene-specific

siRNAs (AR siRNA, Santa Cruz, cat # SC29204) using the DharmaFECT 1 transfection reagent (cat

T-2001-03, Dharmacon, Pittsburgh, PA).

Cell proliferation assay.

2,000 cells were plated per well in 96 well plates, with six replicates for each condition.

Medium was changed on alternate days. Cell proliferation was measured at the indicated times

using the Cell Titer- Glo® luminescent cell viability assay (Promega, cat # G7570) following the

manufacturer's protocol.

Xenograft studies and IVIS imaging.

All animal experiments were pre-approved by the UNMC Institutional Animal Care and

Use Committee (IACUC Protocol # 22-059-10-FC). To assess the effect of ECD OE on

tumorigenesis, $3x10^6$ vector control or ECD-OE LNCaP cells engineered with lentiviral

mCherry-luciferase in 0.1 ml 50% Matrigel (BD Biosciences) in cold PBS were injected

subcutaneously in 8-week-old athymic nude male mice (n=14) on both sides of dorsal flank. Tumor

growth was monitored twice a week using calipers and once a week by IVIS bioluminescence

imaging for up to 37 days. Tumor volume was calculated as length x width x depth/2 [22]. Mice

were euthanized when mice reached pre-determined humane endpoints or reached 2 cm^3 in

volume, as per the institutional IACUC guidelines. For IVIS imaging, mice received 200 μl D-

luciferin (15 mg/ml in PBS; Millipore Sigma #L9504) intraperitoneally 15 min before isoflurane

anesthesia and were placed dorsoventrally in the IVIS™ Imaging System (IVIS 2000). Images

were acquired using the IVIS Spectrum CT and analyzed using the Living Image 4.4 software

(PerkinElmer). The fold change in total Photon flux (p/s) over time was used to compare tumor

growth between the vector and ECD-OE cells groups.

Immunohistochemistry of mice tumors.

The mouse tumors harvested at necropsy were fixed in 10% neutral buffered formalin (cat

316- 155; PROTOCOL), paraffin-embedded and 5 mm sections deparaffinized in xylene and

rehydrated in descending alcohols followed by IHC staining of indicated antibodies as previously

described [22].

RNA sequencing, bioinformatics, and statistical analysis.

Trizol-isolated total RNA from the core piece of the isolated tumors was analyzed on a

Bioanalyzer for quality and purity. RNA-seq libraries were prepared and subjected to 100 bp

paired-end sequencing on an Illumina NovaSeq 6000 system (UNMC Genomics Core). Analysis

of RNA-seq results was done using the nf-core RNA-seq pipeline version 3.9 [51], accessed via

Nextflow version 22.10.3 [52] and executed with Singularity [53] to ensure reproducibility and

portability. Paired-end sequencing reads were aligned to the human reference genome (GRCh38)

using the Spliced Transcripts Alignment to a Reference (STAR) software version 2.7.11a [54],

incorporating Ensemble gene annotations for accurate mapping. Quantification of transcript

abundance was performed using StringTie2 version 1.3.6 [55], with transcripts per million (TPM)

normalization to facilitate comparison across samples. Differential expression analysis between

vector and ECD-OE xenograft tumors was conducted using DESeq2 [56]. Differentially expressed

genes were identified using criteria of an adjusted p-value < 0.05 and a minimum twofold change

in expression. For functional interpretation of the differential expression results, we utilized the

Gene Set Enrichment Analysis (GSEA) software from the Broad Institute [57]. Enrichment plots

were generated to visualize the distribution of gene sets and to identify pathways significantly

associated with the observed transcriptional changes.

RNA Immunoprecipitation (RIP) assay.

RIP assays were performed using the EZ-Magna RIP Kit (Millipore, cat #17-701). Briefly,

cells were harvested and lysed using RIP lysis buffer with 1 freeze-thaw cycle. Cellular debris was

removed by centrifuging the cell lysates at 13K for 15 min. 1 mg aliquots of clarified lysate (BCA

method) were incubated overnight at 4°C with protein A Dynabeads bound to 2.5 µg of polyclonal

rabbit anti-ECD (Proteintech, cat # 10192-1-AP) or control IgG (Abcam cat #

ab172730) antibodies. Post incubation, samples were subjected to five washes in wash buffer. The

bound RNAs were retrieved and subjected to qRT-PCR analysis using gene specific primers. 10%

of lysates served as input controls.

Analysis of mRNA stability.

cell lines were seeded in 6-well plates (200,000 cells per well). 24 h after plating, 5 µg/ml

actinomycin D (catalog # A1410; Sigma, St. Louis, MO) was added and total RNA isolated at the

indicated time points using the TRIzol method. cDNA was synthesized using iScript™ gDNA

Clear cDNA Synthesis Kit (Bio-Rad), as above, followed by qRT-PCR with the indicated gene

specific primers.

Glucose Uptake-Glo assay:

Cells were seeded at 25,000 cells per well in 96-well plates. 24 h later, the cells were

washed twice with PBS and cultured in glucose-free DMEM media (Thermo Scientific cat #

11966025) for 1 h (glucose starvation). Glucose uptake assay was then performed using the

Promega Glucose Uptake-Glo (J1341) assay kit according to the manufacturer's protocol. Parallel

CellTiter-Glo assays were used for normalization of cell numbers.

Seahorse XF Glycolytic Rate Assay.

The Oxygen Consumption Rate (OCR), Extracellular Acidification Rate (ECAR) and

Proton Efflux rate (PER) were measured using the Seahorse XFe Glycolytic rate assay Kit (Agilent

Technologies; cat #103344-100) according to the manufacturer's protocol. Briefly, 3.5×10^4 cells

were seeded per well in 96-well Seahorse XFe cell culture plates and incubated overnight. Cells

were washed and incubated with XF assay medium supplemented with 1 mM pyruvate (Sigma;

cat# S8636), 2 mM L-glutamine (Invitrogen; cat#25030081) and 10 mM glucose (Gibco;

cat#A24940-01) for 1 h at 37°C in a CO_2-free incubator. The OCR, ECAR and PER estimations

were performed as per the manufacturer's instructions. Basal OCR and ECAR was assessed in

response to rotenone/antimycin A (0.5 µM) and 2-deoxy-D-glucose (2-DG; 50 mM). ATP rate was

estimated in response to oligomycin (1.5 μM) and rotenone/antimycin A (0.5 μM). Basal

glycolysis was calculated from non-glycolytic acidification (background) ECAR/PER. Glycolytic

reserve (Compensatory) was calculated as the difference between glycolytic capacity and basal

glycolysis.

Statistical analysis.

Statistical analyses were done using the GraphPad Prism 11 software. Significance

between two groups was determined by the two-tailed Student's t test. Comparisons among three

or more groups were done using either one-way ANOVA or two-way ANOVA with post hoc

multiple comparisons test. A p-value of ≤ 0.05 was considered statistically significant. Survival

curves were analyzed using the Kaplan-Meier analysis with the log rank test. The results are

expressed as the mean $\pm$ SEM from three independent experiments, done in triplicates. The type

of statistical test used in individual experiments is indictated in figure legends.

Acknowledgements: This research was funded by Pilot grants from the Fred & Pamela Buffett

Cancer Center (HB & VB); Department of Defense grants W81XWH-17-1-0616 and W81XWH-

20-1-0058 to HB and W81XWH-20-1-0546, HT94252410337; and NIH grants R21CA241055

and R03CA253193 to VB; the NIH MIRA award R35 GM147467 (to M.J.R.); the Raphael Bonita

Memorial Fund, and support to UNMC core facilities from the NCI Cancer Center Support Grant

(P30CA036727) awarded to Fred & Pamela Buffett Cancer Center and from the Nebraska

Research Initiative.

Author Contributions:

Conceptualization and design; M.R., A.R.R., B.B.K., T.E.R., F.O., S.M., M.J.R., G.K., K.D.,

B.C.M., H.B., and V.B. Manuscript writing; M.R., A.R.R., B.C.M., T.E.R., K.D., M.J.R., H.B.,

and V.B. Data analysis and interpretation; M.R., A.R.R., B.B.K., T.E.R., F.O., S.M., M.J.R., G.K.,

K.D., B.C.M., H.B., and V.B. Manuscript editing; M.J.R., B.C.M., H.B., and V.B.

Bioinformatics analysis; M.R., T.E.R., M.J.R. All authors have read and approved the final

version of the manuscript and agree with the order of presentation of the authors.

Competing Interests:

Dr. H. Band and Dr. V. Band received funding from Nimbus Therapeutics for an unrelated

project. The remaining authors do not have any competing interest to declare.

Data Availability Statement:

The RNA-seq files generated in this study are available from the NCBI Gene Expression Omnibus

(GEO) under the accession number: GSE287898.

References:

543 1   Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. *CA Cancer J Clin* 2023; 73:
544   17-48.
545
546 2   Huggins C, Hodges CV. Studies on prostatic cancer. I. The effect of castration, of estrogen and
547   androgen injection on serum phosphatases in metastatic carcinoma of the prostate. *CA Cancer J*
548   *Clin* 1972; 22: 232-240.
549
550 3   Rathkopf D, Scher HI. Androgen receptor antagonists in castration-resistant prostate cancer.
551   *Cancer J* 2013; 19: 43-49.
552
553 4   Watson PA, Arora VK, Sawyers CL. Emerging mechanisms of resistance to androgen receptor
554   inhibitors in prostate cancer. *Nat Rev Cancer* 2015; 15: 701-711.
555
556 5   Lokeshwar SD, Klaassen Z, Saad F. Treatment and trials in non-metastatic castration-resistant
557   prostate cancer. *Nat Rev Urol* 2021; 18: 433-442.
558
559 6   Ahmad F, Cherukuri MK, Choyke PL. Metabolic reprogramming in prostate cancer. *Br J Cancer*
560   2021; 125: 1185-1196.

562 7   Lasorsa F, di Meo NA, Rutigliano M, Ferro M, Terracciano D, Tataru OS *et al*. Emerging
563   Hallmarks of Metabolic Reprogramming in Prostate Cancer. *Int J Mol Sci* 2023; 24.
564

565 8   de Wet L, Williams A, Gillard M, Kregel S, Lamperis S, Gutgesell LC *et al*. SOX2 mediates
566   metabolic reprogramming of prostate cancer cells. *Oncogene* 2022; 41: 1190-1202.
567

568 9   Mishra R, Haldar S, Placencio V, Madhav A, Rohena-Rivera K, Agarwal P *et al*. Stromal
569   epigenetic alterations drive metabolic and neuroendocrine prostate cancer reprogramming. *J Clin*
570   *Invest* 2018; 128: 4472-4484.
571

572 10   Flavin R, Zadra G, Loda M. Metabolic alterations and targeted therapies in prostate cancer. *J*
573   *Pathol* 2011; 223: 283-294.
574

575 11   Beier AK, Puhr M, Stope MB, Thomas C, Erb HHH. Metabolic changes during prostate cancer
576   development and progression. *J Cancer Res Clin Oncol* 2023; 149: 2259-2270.
577

578 12   Gaziova I, Bonnette PC, Henrich VC, Jindra M. Cell-autonomous roles of the ecdysoneless gene
579   in Drosophila development and oogenesis. *Development* 2004; 131: 2715-2725.
580

581 13   Kim JH, Gurumurthy CB, Naramura M, Zhang Y, Dudley AT, Doglio L *et al*. Role of
582   mammalian Ecdysoneless in cell cycle regulation. *J Biol Chem* 2009; 284: 26402-26410.
583

584 14   Olou AA, Sarkar A, Bele A, Gurumurthy CB, Mir RA, Ammons SA *et al*. Mammalian ECD
585   Protein Is a Novel Negative Regulator of the PERK Arm of the Unfolded Protein Response. *Mol*
586   *Cell Biol* 2017; 37.
587

588 15   Claudius AK, Romani P, Lamkemeyer T, Jindra M, Uhlirova M. Unexpected role of the steroid-
589   deficiency protein ecdysoneless in pre-mRNA splicing. *PLoS Genet* 2014; 10: e1004287.
590

591 16   Malinova A, Cvackova Z, Mateju D, Horejsi Z, Abeza C, Vandermoere F *et al*. Assembly of the
592   U5 snRNP component PRPF8 is controlled by the HSP90/R2TP chaperones. *J Cell Biol* 2017;
593   216: 1579-1596.
594

595 17   Saleem I, Mirza S, Sarkar A, Raza M, Mohapatra B, Mushtaq I *et al*. The Mammalian
596   Ecdysoneless Protein Interacts with RNA Helicase DDX39A To Regulate Nuclear mRNA
597   Export. *Mol Cell Biol* 2021; 41: e0010321.
598

599 18   Mirza S, Kalluchi A, Raza M, Saleem I, Mohapatra B, Pal D *et al*. Ecdysoneless Protein
600   Regulates Viral and Cellular mRNA Splicing to Promote Cervical Oncogenesis. *Mol Cancer Res*
601   2022; 20: 305-318.
602

603 19   Dey P, Rachagani S, Chakraborty S, Singh PK, Zhao X, Gurumurthy CB *et al*. Overexpression of
604   ecdysoneless in pancreatic cancer and its role in oncogenesis by regulating glycolysis. *Clin*
605   *Cancer Res* 2012; 18: 6188-6198.
606

607 20   Xu SH, Huang JZ, Xu ML, Yu G, Yin XF, Chen D *et al*. ACK1 promotes gastric cancer
608   epithelial-mesenchymal transition and metastasis through AKT-POU2F1-ECD signalling. *J*
609   *Pathol* 2015; 236: 175-185.
610

611 21   Xu SH, Huang JZ, Chen M, Zeng M, Zou FY, Chen D *et al*. Amplification of ACK1 promotes

612   gastric tumorigenesis via ECD-dependent p53 ubiquitination degradation. *Oncotarget* 2017; 8:
613   12705-12716.
614

615   22   Mohapatra BC, Mirza S, Bele A, Gurumurthy CB, Raza M, Saleem I *et al*. Ecdysoneless
616   Overexpression Drives Mammary Tumorigenesis through Upregulation of C-MYC and Glucose
617   Metabolism. *Mol Cancer Res* 2022; 20: 1391-1404.
618

619   23   Isaacs JT. Hormonally responsive versus unresponsive progression of prostatic cancer to
620   antiandrogen therapy as studied with the Dunning R-3327-AT and -G rat adenocarcinomas.
621   *Cancer Res* 1982; 42: 5010-5014.
622

623   24   Coutinho I, Day TK, Tilley WD, Selth LA. Androgen receptor signaling in castration-resistant
624   prostate cancer: a lesson in persistence. *Endocr Relat Cancer* 2016; 23: T179-T197.
625

626   25   Wang Q, Li W, Liu XS, Carroll JS, Janne OA, Keeton EK *et al*. A hierarchical network of
627   transcription factors governs androgen receptor-dependent prostate cancer growth. *Mol Cell*
628   2007; 27: 380-392.
629

630   26   Zambelli F, Pesole G, Pavesi G. Pscan: finding over-represented transcription factor binding site
631   motifs in sequences from co-regulated or co-expressed genes. *Nucleic Acids Res* 2009; 37: W247-
632   252.
633

634   27   Dreos R, Ambrosini G, Groux R, Cavin Perier R, Bucher P. The eukaryotic promoter database in
635   its 30th year: focus on non-vertebrate organisms. *Nucleic Acids Res* 2017; 45: D51-D55.
636

637   28   Bailey TL, Johnson J, Grant CE, Noble WS. The MEME Suite. *Nucleic Acids Res* 2015; 43:
638   W39-49.
639

640   29   Rajan P, Sudbery IM, Villasevil ME, Mui E, Fleming J, Davis M *et al*. Next-generation
641   sequencing of advanced prostate cancer treated with androgen-deprivation therapy. *Eur Urol*
642   2014; 66: 32-39.
643

644   30   Lee CH, Yu CC, Wang BY, Chang WW. Tumorsphere as an effective in vitro platform for
645   screening anti-cancer stem cell drugs. *Oncotarget* 2016; 7: 1215-1226.
646

647   31   Bredholt G, Mannelqvist M, Stefansson IM, Birkeland E, Bo TH, Oyan AM *et al*. Tumor necrosis
648   is an important hallmark of aggressive endometrial cancer and associates with hypoxia,
649   angiogenesis and inflammation responses. *Oncotarget* 2015; 6: 39676-39691.
650

651   32   Gerdes J, Li L, Schlueter C, Duchrow M, Wohlenberg C, Gerlach C *et al*. Immunobiochemical
652   and molecular biologic characterization of the cell proliferation-associated nuclear antigen that is
653   defined by monoclonal antibody Ki-67. *Am J Pathol* 1991; 138: 867-873.
654

655   33   Tabrizian N, Nouruzi S, Cui CJ, Kobelev M, Namekawa T, Lodhia I *et al*. ASCL1 is activated
656   downstream of the ROR2/CREB signaling pathway to support lineage plasticity in prostate
657   cancer. *Cell Rep* 2023; 42: 112937.
658

659   34   Smith BA, Sokolov A, Uzunangelov V, Baertsch R, Newton Y, Graim K *et al*. A basal stem cell
660   signature identifies aggressive prostate cancer phenotypes. *Proc Natl Acad Sci U S A* 2015; 112:
661   E6544-6552.
662

663 35 Agell L, Hernandez S, Nonell L, Lorenzo M, Puigdecanet E, de Muga S *et al*. A 12-gene
664   expression signature is associated with aggressive histological in prostate cancer: SEC14L1 and
665   TCEB1 genes are potential markers of progression. *Am J Pathol* 2012; 181: 1585-1594.
666
667 36 Islam R, Mishra J, Polavaram NS, Bhattacharya S, Hong Z, Bodas S *et al*. Neuropilin-2 axis in
668   regulating secretory phenotype of neuroendocrine-like prostate cancer cells and its implication in
669   therapy resistance. *Cell Rep* 2022; 40: 111097.
670
671 37 Erkelenz S, Stankovic D, Mundorf J, Bresser T, Claudius AK, Boehm V *et al*. Ecd promotes U5
672   snRNP maturation and Prp8 stability. *Nucleic Acids Res* 2021; 49: 1688-1707.
673
674 38 Raza M, Rajan AR, Kalluchi A, Saleem I, Kennedy BB, Bhakat KK *et al*. ECD functions as a
675   novel RNA-binding protein to regulate mRNA splicing. *bioRxiv* 2025: 2025.2001.2024.634785.
676
677 39 Warburg O, Wind F, Negelein E. The Metabolism of Tumors in the Body. *J Gen Physiol* 1927; 8:
678   519-530.
679
680 40 Chen Y, Zhou Q, Hankey W, Fang X, Yuan F. Second generation androgen receptor antagonists
681   and challenges in prostate cancer treatment. *Cell Death Dis* 2022; 13: 632.
682
683 41 Zhao X, Mirza S, Alshareeda A, Zhang Y, Gurumurthy CB, Bele A *et al*. Overexpression of a
684   novel cell cycle regulator ecdysoneless in breast cancer: a marker of poor prognosis in
685   HER2/neu-overexpressing breast cancer patients. *Breast Cancer Res Treat* 2012; 134: 171-180.
686
687 42 Acosta AM, Al Rasheed MRH, Rauscher GH, Vormittag E, Mon KS, Sharif A *et al*. Tumor
688   necrosis in radical prostatectomies with high-grade prostate cancer is associated with multiple
689   poor prognostic features and a high prevalence of residual disease. *Hum Pathol* 2018; 75: 1-9.
690
691 43 Bele A, Mirza S, Zhang Y, Ahmad Mir R, Lin S, Kim JH *et al*. The cell cycle regulator
692   ecdysoneless cooperates with H-Ras to promote oncogenic transformation of human mammary
693   epithelial cells. *Cell Cycle* 2015; 14: 990-1000.
694
695 44 Ma S, Chen Y, Quan P, Zhang J, Han S, Wang G *et al*. NPAS2 promotes aerobic glycolysis and
696   tumor growth in prostate cancer through HIF-1A signaling. *BMC Cancer* 2023; 23: 280.
697
698 45 Zhou Y, Zeng F, Richards GO, Wang N. ENO2, a Glycolytic Enzyme, Contributes to Prostate
699   Cancer Metastasis: A Systematic Review of Literature. *Cancers (Basel)* 2024; 16.
700
701 46 Cui Y, Nadiminty N, Liu C, Lou W, Schwartz CT, Gao AC. Upregulation of glucose metabolism
702   by NF-kappaB2/p52 mediates enzalutamide resistance in castration-resistant prostate cancer cells.
703   *Endocr Relat Cancer* 2014; 21: 435-442.
704
705 47 Gillis JL, Hinneh JA, Ryan NK, Irani S, Moldovan M, Quek LE *et al*. A feedback loop between
706   the androgen receptor and 6-phosphogluoconate dehydrogenase (6PGD) drives prostate cancer
707   growth. *Elife* 2021; 10.
708
709 48 Liu J, Chen G, Liu Z, Liu S, Cai Z, You P *et al*. Aberrant FGFR Tyrosine Kinase Signaling
710   Enhances the Warburg Effect by Reprogramming LDH Isoform Expression and Activity in
711   Prostate Cancer. *Cancer Res* 2018; 78: 4459-4470.
712
49 Yang B, Chen W, Tao T, Zhang J, Kong D, Hao J *et al*. UBE2N promotes cell viability and

glycolysis by promoting Axin1 ubiquitination in prostate cancer cells. *Biol Direct* 2024; 19: 35.

50   Chakraborty S, Bhat AM, Mushtaq I, Luan H, Kalluchi A, Mirza S *et al*. EHD1-dependent traffic
of IGF-1 receptor to the cell surface is essential for Ewing sarcoma tumorigenesis and metastasis.
Commun Biol 2023; 6: 758.

51   Ewels PA, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A *et al*. The nf-core framework for
community-curated bioinformatics pipelines. *Nat Biotechnol* 2020; 38: 276-278.

52   Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C. Nextflow enables
reproducible computational workflows. *Nat Biotechnol* 2017; 35: 316-319.

53   Kurtzer GM, Sochat V, Bauer MW. Singularity: Scientific containers for mobility of compute.
PLoS One 2017; 12: e0177459.

54   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S *et al*. STAR: ultrafast universal
RNA-seq aligner. *Bioinformatics* 2013; 29: 15-21.

55   Kovaka S, Zimin AV, Pertea GM, Razaghi R, Salzberg SL, Pertea M. Transcriptome assembly
from long-read RNA-seq alignments with StringTie2. *Genome Biol* 2019; 20: 278.

56   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq
data with DESeq2. *Genome Biol* 2014; 15: 550.

57   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA *et al*. Gene set
enrichment analysis: a knowledge-based approach for interpreting genome-wide expression
profiles. *Proc Natl Acad Sci U S A* 2005; 102: 15545-15550.

Fig. 1

Fig. 1. Correlation of ECD and AR expression; AR regulates ECD expression. (A) Spearman's pairwise correlation plot illustrates a positive correction of *AR* and *ECD* mRNA in TCGA Prostate Adenocarcinoma Firehose dataset of cBioPortal online platform. Number of patients (N), correlation coefficient (r) and p-value are indicated on the plot. Pairwise correlation analysis was performed to calculate statistical significance. **(B & C)** Indicated PC cell lines were cultured in steroid free conditions for 72 hours. Cells were then treated with indicated concentrations of DHT for 12 hours and immunoblotted with anti-ECD or anti-AR antibodies. β- actin was used as a loading control. **(D & E)** qRT-PCR was performed in the same samples; RNA was isolated by standard TRIzol phenol chloroform method. 18s rRNA was used for normalization. **(F & G)** Similar protocols as mentioned above was used to perform a time response with DHT. Western blotting **(F & G)** and qRT-PCR **(H & I)** were performed on cell lysates, as above. **(J & K)** Indicated PC cell lines transfected with control siRNA or AR siRNA and cultured for 72 hours. Cells were then treated with indicated concentrations of DHT for 12 hours and then lysates were western blotted with anti-ECD or anti-AR antibodies. β- actin was used as a loading control in all blots. Numbers below the blot show the quantification of band intensities after normalizing with their respective loading control, β- actin in comparison with control samples using ImageJ software. Bar graphs quantitation is from three independent experiments in triplicates. Student t- test was used to calculate statistical significance *** $p < 0.001$ ** $p < 0.01$, * $p < 0.05$, [ns] $p > 0.05$.

Fig. 2.

Fig. 2. Androgen treatment increased AR occupancy on ECD promoter and Enzalutamide treatment decreases ECD expression. (A & B) AR binding site; Top three predicted AR binding sites on the ECD promotor. **(C & D)** Chromatin immunoprecipitation (ChIP) analysis. Indicated cells were cultured in steroid-free conditions for 3 days, followed by dihydrotestosterone (DHT) treatment for 3 h. Cells were cross-linked with formaldehyde and chromatin was immunoprecipitated with antibody against AR. IgG, used as a non-binding Control. Immunoprecipitated DNA was quantified with qPCR for specific binding sites at ECD promoter using primers encompassing all three sites -712 to -565 from transcription start site. Student t- test was performed to calculate statistical significance *** $p < 0.001$. **(E & F)** DHT increases ECD promoter activity. ECD promoter spanning approx. 1500 bp was cloned upstream of secreted Gaussia Luciferase, Secreted Alkaline phosphatase (SEAP) was used as an internal control. Cells expressing WT promoter or various AR binding sites deleted mutants were treated with 100 nM DHT for 3 h. Supernatant was collected, and Luminescence was measured using Secrete-Pair Dual Luminescence Assay Kit. Data represent three independent experiments, each done in triplicates. *Student t- test* was performed to calculate statistical significance *** $p < 0.001$, ** $p < 0.01$. **(G)** Box plot depicting ECD mRNA expression in prostate tumors pre- and post-androgen deprivation therapy 7 samples in each case (GSE48403). A Wilcoxon matched-pairs signed-rank test was used to compare expression in the groups. **(H & I**) Indicated PC cell lines were cultured in steroid-free medium for 72 hours, cells were then treated with 10 nM DHT, followed by treatment with indicated concentration of MDV3100 for another 48 hours. Lysates were then western blotted with indicated antibodies. β-actin was used as a loading control. **(J & K)** qRT-PCR was performed in the one set of samples treated as above. RNA was isolated by standard TRIzol phenol chloroform method. 18s rRNA was used for normalization. Bar graphs represent fold change in AR and ECD

mRNA in LNCaP **(J)** and C4-2B cells **(K)** respect to vehicle treatment. Data represent three independent experiments, each done in triplicates. Student t- test was performed to calculate statistical significance *** $p < 0.001$, ** $p < 0.01$.

Fig. 3

Fig. 3. Depletion of ECD induces proliferation block and diminishes oncogenic traits in prostate cancer cells. LNCaP and C4-2B cells expressing doxycycline (Dox) regulated Cas9 and guide RNAs targeting either ECD (gRNA1 & 2) or non-targeting control (NTC) were cultured in the presence of Dox (1 µg/ml) for 96 hours to deplete ECD. **(A)** Western blot analysis. Numbers below the blot show the quantification of band intensities after normalizing with their respective loading control, β- actin in comparison with NTC using ImageJ software. Representative images and histograms depicting 2D (two dimensional) colony **(B & C),** tumorsphere **(D & E)**, soft agar colony **(F & G),** Matrigel organoid formation **(H & I),** migration **(J & K)** and invasion **(L & M),** of control and ECD depleted PC cells lines. The representative images of tumorspheres, soft agar colonies and Matrigel organoids are 200x magnification, scale bar, 250 µm. Migration and invasion images are 40x magnification, scale bar, 1000 µm. The Bar graph data presented as mean values ± SE of three experiments done in triplicates from 40x magnification images. Student t-test was used to calculate statistical significance *** $p < 0.001$, ** $p < 0.01$.

Fig. 4

(A)

Hyperplastic prostate tissue

(B)

Prostate cancer

Low ECD expression

High ECD expression

(C)

ECD-Low (n=298)

ECD-High (n=224)

Log Rank= 5.09; p=0.024

Cumulative Survival

Time after surgery(months)

(D)

ECD-Low (n=201)

ECD-High (n=283)

raw p 0.018
bonf p 1.000

Overall survival probability

Follow up in months

Fig. 4. Higher expression of ECD protein and mRNA in prostate cancer tissues correlate with short patient survival. (A & B) Representative IHC images of ECD expression in hyperplastic prostate tissue **(A),** and low-ECD (staining intensity 1) and high-ECD expression (staining intensity 3) in prostate cancer tissues **(B)** at magnification 200x in (**A & B**). **(C)** Kaplan-Meier cumulative survival curve after surgery with patient tumors expressing low (staining intensities 0 or 1) or high ECD protein levels (staining intensities 2 or 3). P-values were calculated with a log-rank test. **(D)** Mixed prostate adenocarcinoma (2022-v32) cohort (n=484) of The Cancer Genome Atlas (TCGA) dataset was used to stratify *ECD* mRNA expression as high (> 31.49 transcripts per million (TPM) and low (< 31.49 TPM). Analysis was performed using R2 Genomics platform. Kaplan-Meier plot for survival curve of prostate cancer patients shows short survival in patients with tumors expressing high *ECD* mRNA (n=283) in comparison to low *ECD* mRNA (n=201). P-value is indicated in the KM plot.

Fig. 5.

Fig. 5. ECD overexpression in LNCaP cells exhibit enhanced *in vitro* oncogenic traits and *in vivo* tumorigenesis. (A) Western blot depicts ECD overexpression; densitometry in respect to vector cells after normalizing with β-actin, shown below. **(B)** Representative images in 100x magnification (scale bar, 400 μm) show soft agar colony formation abilities of cells. **(C)** Bars depict the quantitation of soft agar colonies per 40x magnification **(D)** Representative images in 100x magnification (scale bar, 400 μm) depict 3D colony formation abilities in Matrigel. **(E)** Bar graphs show the quantitation of 3D colonies per 40x magnification field. Data represent three independent experiments, each done in triplicates. ***p < 0.001 Student t-test was used to calculate statistical significance in **(C)** and **(E)**. **(F)** mCherry-luciferase expressing LNCaP-vector and ECD-OE cells were injected subcutaneously in nude mice (14/group) in both sides. Primary tumor growth was monitored by bioluminescence imaging at the indicated time on day 7 and day 37. **(G)** Bioluminescence signals of tumors over time are shown as change in photon flux over the time. Data are mean ± SEM, n=14 (two sided tumors); Unpaired t-test; ***p<0.001. **(H)** Tumor growth is followed over days with vernier caliper and fold change in tumor volume is plotted. Data are mean ± SEM, n=14; Two-way ANOVA (mixed model); ***p<0.001. **(I)** Ten representative harvested tumors are displayed in each group. **(J)** Scattered plot represents the weights of harvested tumors. Data are mean ± SEM, n=14; unpaired t-test; **p<0.01. **(K)** H&E and IHC analysis with ECD and Ki-67. Representative images from three independent tumors (T1, T2 and T3) from both groups, in 400x magnification (scale bar, 200 μm).

Fig. 6

(A) PCA Plot

(B) ECD Vs. Vector Xenografts

HALLMARK MTORC1 SIGNALING
HALLMARK INTERFERON GAMMA RESPONSE
HALLMARK APICAL JUNCTION
HALLMARK TNFA SIGNALING VIA NFKB
HALLMARK INTERFERON_ALPHA_RESPONSE
HALLMARK ALLOGRAFT_REJECTION
HALLMARK ANGIOGENESIS
HALLMARK GLYCOLYSIS
HALLMARK MYOGENESIS
HALLMARK HYPOXIA

(C)

Enrichment plot: HALLMARK_GLYCOLYSIS

NES: 1.52
NOM p-value: 0.0018
FDR q-value: 0.06

(D)

(E) Mouse Xenograft

(F) LNCaP

(G) C4-2B

Fig. 6. ECD regulates mRNA expression of glycolytic and neuro endocrine signature genes.
(A), Principal component analysis (PCA) analysis of RNA-seq data from LNCaP-vector expressing and LNCaP-ECD-OE xenograft tumors shows distinct clustering of datasets. **(B)** Bar plot displays enriched hallmark GSEA (Gene set enrichment analysis) gene sets. X-axis represents normalized enrichment scores (NES) of the signaling pathways with significant nominal p-values (NOM p-val) and <25% false discover rate q-values (FDR q-val). Red arrow indicates glycolysis (followed further in this study). **(C)** Enrichment plot of glycolysis in ECD-OE xenograft tumors compared to vector expressing xenograft tumors. NES, NOM p-val and FDR q-val. are indicated on the plot. **(D)** Heatmap of key glycolytic genes and indicated signature genes of PC oncogenic pathways is presented. Yellow represents upregulated and blue as downregulated genes. Major glycolytic genes are marked as red arrows. **(E-G)** qRT-PCR confirms the overexpression of *ECD* in xenograft tumors and prostate cancer cell lines. 18s rRNA was used for normalization. qRT-PCR analyses of indicated glycolytic genes. mRNA quantitation data represents mean ± SEM with two-tailed unpaired *t* test. *n* = 3; **, *P* < 0.01; ***, *P* < 0.001.

Fig. 7.

Fig.7. ECD associates with mRNAs of glycolytic enzymes and regulates their stability. (A & B). Indicated PC cells were subjected to RNA immunoprecipitation using IgG or anti-ECD antibodies and processed as per instructions in Magna RIP™ RNA RIP kit. Co-precipitated RNAs were purified and the indicated target RNAs were detected by qRT-PCR using gene specific primers. Graph display mean fold enrichment in each respective gene, as compared to IgG control. **(C-H)** mRNA stabilities of various glycolytic enzymes after actinomycin D (5 µg/mL) treatment for indicated time points in vector or ECD expressing PC cells. Trend lines were created by measuring relative abundance of each transcript with respect to the 0-minute time point for vector and ECD-OE LNCaP and C4-2B cells and are expressed as percentage of mRNA remaining. Transcript abundance was measured by qRT-PCR using 18S rRNA as a housekeeping gene. Half-life (t1/2) for indicated mRNAs are displayed for vector and ECD-OE cells. Graph displays mean fold enrichment in each respective gene, as compared to IgG control. Quantitation data represents mean ± SEM with two-tailed unpaired *t* test. $n = 3$; ***, $P < 0.001$.

Fig. 8.

Fig. 8. ECD overexpression increases glucose uptake and glycolytic rate. Glucose uptake in LNCaP **(A)** and C4-2B **(B)** PC cell lines. The values were normalized with respective to cell counts and depicted as compared with vector control. Quantification of results from three biological experiments, each with six technical replicates is shown as a bar graph. Data represents as mean ± SEM and two-tailed unpaired Student's t-test (*p<0.05, **p<0.01, ***p<0.001). **(C-J)** Sea horse glycolytic rate was assessed in vector and ECD-OE LNCaP **(C)** and C4-2B **(E)** cells. Cells were seeded in 96-well plates and exposed to Rot/AA (rotenone and antimycin cocktail) and 2-DG to measure proton eflux rate **(PER, C & D)** and extracellular acidification rate **(ECAR, E & F)**. One representative PER and ECAR plots from three independent experiments are displayed (**C-F**). Basal glycolysis **(G & H)** and compensatory glycolysis **(I & J)** rates are presented as bar graphs, glycoPER in pmol/min calculated from subtracting ECAR (extra cellular acidification rate) from PER (proton efflux rates) (**G-J**). Quantitation data represents mean ± SEM with two-tailed unpaired *t* test. $n = 3$; **, $P < 0.01$; ***, $P < 0.001$.

Table S1

Multivariate Cox's regression analysis of prostate cancer patient tumors
RR: Relative risk; CI: Confidence interval.

Parameter	RR	95% CI	P-value
ECD expression (Low. vs. High)	1.689	1.085-2.629	0.02
Gleason grade (2-6 vs.7-10)	2.415	1.247-4.675	0.009
Resection Margins (Neg. vs. Pos.)	1.666	1.038-2.675	0.035

Table S2

Incidence of tumor formation in vector or ECD-overexpressing LNCaP xenograft model.

No. of Days	LNCaP Control group	ECD OE LNCaP Group
16	0 (0/26)/ 0%	6 (6/28)/ 21.4%
22	8 (8/26)/ 30.7%	20 (20/28)/ 71.4%
27	9 (9/26)/ 34.6%	23 (23/28)/ 82.1%
35	13 (13/26)/ 50%	26 (26/28)/ 92.8%

Table S3

List of ECD promoter mutants and the corresponding sequences deleted from TSS.

ECD Promotor Mutants	Sequence deleted from TSS
CS-HPRM35342-LvPG04-01- site-1Δ	698 bp – 682 bp
CS-HPRM35342-LvPG04-02-site-2Δ	657 bp – 641bp
CS-HPRM35342-LvPG04-03-site-3Δ	640 bp – 624 bp
CS- HPRM35342-LvPG04-04-site-1,2Δ	698 bp – 682 bp, 657 bp – 641bp
CS-HPRM35342-LvPG04-05-site-1,3Δ	698 bp – 682 bp, 640 bp – 624 bp
CS-HPRM35342- LvPG04-06 - site-2,3Δ	657 bp – 641bp, 640 bp – 624 bp
CS-HPRM35342-LvPG04-07 - site-1,2,3Δ	698 bp – 682 bp, 657 bp – 641bp, 640 bp – 624bp

Table S4

List of real time qRT-PCR primers of human genes.

Primer Name	Forward Sequence	Reverse Sequence
LDHA	TTGACCTACGTGGCTTGGAAG	GGTAACGGAATCGGGCTGAAT
HKII	GAGCCACCACTCACCCTACT	ACCCAAAGCACACACCGGAAGTT
PKM2	CCACTTGCAATTATTTGAGGAA	GTGAGCAGACCTGCCAGACT
ECD	ACTTTGAAACACACGAACCTGGCG	TGATGCAGGTGTGTGCTAGTTCCT
18S	GCTTAATTTGACTCAACACGGGA	AGCTATCAATCTGTCAATCCTGTC
AR	TTGCAAGAGAGCTGCATCAGT	ACTGTGTGTGGAAATAGATGGGC

Fig. S1

Fig. S1. PC tissue specimens show higher expression of *ECD* mRNA. Box plot showing mRNA expression of *ECD* from **(A)** 4 normal and 22 tumor samples from cohort 1 (GSE32269) and **(B)** 28 normal and 59 tumor samples from another independent cohort (GSE35988). The mentioned two publicly available datasets were analyzed for *ECD* mRNA expression. A *paired t* test was used to compare expression in normal versus cancer samples.

Fig. S2

Fig. S2. ECD-overexpressing PC cells exhibit enhanced anchorage independence and Matrigel three-dimensional (3D) growth but show no alteration in 2-D proliferation rate or estrogen dependence. (**A**) Western blot shows overexpression of ECD in C4-2B cell line. Numbers below the blot shows the quantification of band intensities after normalizing with their respective loading control, β- actin in comparison with vector cells using ImageJ software. (**B & C**) Cell proliferation curves showing proliferation rate of LNCaP (**B**) and C4-2B cells (**C**) overexpressing ECD. Assays were performed using Cell-Titer-Glo**.** Luminescence values were plotted against number of days and normalized to day 1. Quantitation data represents mean ± SEM with two-way ANOVA test. $n = 3$; ns, $P > 0.05$. (**D**) Representative images in 100x magnification (scale bar, 400 µm) depict soft agar colony formation abilities upon overexpression of ECD in C4-2B cells. Images were taken after fixing and staining the colonies with 0.05% crystal violet. (**E**) Bars depict the quantitation of soft agar colonies per 40x magnification field in vector and ECD overexpressing C4-2B cells. (**F**) Representative images in 100x magnification (scale bar, 400 µm) depict 3D colony formation abilities of C4-2B in Matrigel upon overexpression of ECD. (**G**) Bar graphs show the quantitation of 3D colonies per 40x magnification field in vector and ECD overexpressing C4-2B cells. Data represent three independent experiments, each done in triplicates. ***p < 0.001 Student t-test was used to calculate statistical significance. (**H**) LNCaP cells expressing vector or ECD were cultured in charcoal-stripped serum containing phenol red free medium for 72 hours. Cells were then treated with indicated concentrations of DHT. Growth Assays were performed using Cell-Titer-Glo**.** Luminescence values were plotted for both Day 1 and Day 5. Data represent three independent experiments, each done in triplicates. [ns]p > 0.05, Two-way ANOVA (mixed model) was used to calculate statistical significance.

Fig. S3

Fig. S3. Transcriptomic analysis of ECD overexpressing prostate xenograft tumors and LNCaP cell line confirm upregulation of Neuroendocrine PC markers. (A & B) qRT-PCR analysis of indicated neuroendocrine (NE) signature genes of LNCaP-vector expressing and LNCaP ECD-OE xenograft tumors **(A)** and vector expressing or ECD-OE LNCaP cells **(B).** 18s rRNA was used for normalization. mRNA quantitation data represents mean ± SEM with two-tailed unpaired *t* test. $n = 3$; ns, $P > 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Fig. S4

Fig. S4. Seahorse Glycolytic Rate assay in prostate cancer cell lines overexpressing ECD. (A & B) Oxygen consumption rates (OCR) at various time points followed by injections of Rot/AA (0.5 μM), and 2-DG (50 mM) in vector or ECD-OE cells.