Pectobacterium versatile* β-lactamase, a common good of the soft rot *Pectobacteriaceae

species complex

Camille Lorang [1#], Pierre-Yves Canto[1#], Erwan Gueguen[2], Jacques Pédron[1] and Marie-Anne Barny[1*]

[1]Sorbonne Université, INRAE, CNRS, Institut d'Ecologie et des Sciences de l'Environnement de Paris, F-75252

Paris, France.

[2] Université de Lyon, CNRS, UCBL, INSA, UMR5240 Microbiologie, Adaptation et Pathogénie, Villeurbanne,

France.

[*] Corresponding author: marie-anne.barny@sorbonne-universite.fr

[#] equal contributions

Running title: β-lactamase role in natural environment

Statements relating to our ethics and integrity policies:

The data shown in this paper are available within the article and supplementary materials. The funding

of the ANR, ANR-19-CE35-0016-03, is acknowledged. The funders had no role in study design, data

collection and interpretation, or the decision to submit the work for publication. The authors declare

that there are no conflicts of interest. All applicable local, national and international regulations and

conventions, as well as normal scientific ethical practices, were followed in the preparation of this

work. This manuscript has not been previously or simultaneously published or submitted elsewhere

and was critically reviewed and approved by all co-authors before submission. The CRediT of all

authors is provided.

Key words: β-lactamase, Pectobacteriaceae, ecology, species complex

Abstract

Little is known about the role of antibiotics and associated resistance in microbial ecosystems in the absence of clinical antibiotic pressure. The Soft Rot Pectobacteriaceae (SRP) species complex, which comprises 37 bacterial species that are collectively responsible for the severe rotting of many crops, is an interesting model to analyse the role of β-lactam and β-lactamases in natural ecosystems. In particular, within this complex, most *Pectobacterium versatile* strains harbour a β-lactamase called Bla$_{PEC-1}$. The aim of our work was to analyse the role of Bla$_{PEC-1}$ during infection. To this end, two *bla*$_{PEC-1}$-deleted strains were constructed and compared with their wild-type counterparts *in vitro* and in potato tuber infections. *In vitro*, the Bla$_{PEC-1}$ β-lactamase enables *P. versatile* to resist ampicillin or carbapenem produced by *Pectobacterium brasiliense*. In mono-infections on potato tubers, *bla*$_{PEC-1}$-deleted strains were unaffected in terms of virulence, fitness or association with commensal bacteria. In mixed infections, the Bla$_{PEC-1}$ β-lactamase proved necessary for the coexistence of *P. versatile* with the carbapenem-producing strain, and also for the protection of carbapenem-sensitive strains both *in vitro* and *in planta*. Interestingly, *in planta* protection was observed even if the *bla*$_{PEC-1}$ gene was repressed and bacteria expressing Bla$_{PEC-1}$ were in the minority within the symptom. These results indicate that Bla$_{PEC-1}$ exerts a true β-lactamase function during the infection process and acts as a public good of the SRP species complex. Finally, our results highlight the important role of β-lactamase in maintaining of strains diversity in natural ecosystem.

Introduction

Antibiotics are probably one of the most successful forms of chemotherapy in the history of medicine. The β-lactam antibiotics, which disrupt bacterial cell wall formation in both Gram-negative and Gram-positive bacteria, are currently the most widely used class of antibacterial agents in the treatment of infectious diseases (Bush and Bradford, 2016). All β-lactam antibiotics share the β-lactam ring as the essential core structure and are classified into four classes: penicillins, cephalosporins, carbapenems and monobactams (Kim *et al.*, 2023). The massive use of β-lactams in human and animal health is leading to the constant emergence of bacterial resistance. In particular, the regular emergence of new β-lactamases that cleave and inactivate different β-lactam rings is a cause for concern (Tooke *et al.*, 2019). To date, more than 2,700 unique β-lactamases have been described, they are classified into four molecular classes (classes A to D) based on their primary structure, which are further subdivided into functional groups based on their substrate preferences (Bush, 2018). For example, carbapenems are not hydrolysed by most β-lactamases, with the exception of class B and class D2df β-lactamases. It is recognized that β-lactamases are ancient enzymes that existed without the pressure of therapeutic antibiotics, and that they have emerged in clinically pathogenic species following transfer from environmental sources. The rapid spread of transmissible β-lactamases carried on plasmids or transposons that hydrolyze recently approved cephalosporins, monobactams and carbapenems is important in Gram-negative clinical pathogens (Cantón *et al.*, 2008). However, the wealth of data on clinical antibiotics, resistance mechanisms and spread of resistance through mobile genetic elements contrasts with the lack of knowledge about the evolutionary and ecological processes that take place in microbial ecosystems in the absence of antibiotic pressure. In natural ecosystems, antibiotics are not present at therapeutic dose and some authors have proposed that antibiotics act as regulatory or signaling molecules rather than weapons in this context (Romero *et al.*, 2011). In this scenario, β-lactamases in nature could disrupt such regulatory role or signaling. This scenario is supported by *in vitro* analysis of antibiotic role at sub-inhibitory concentration but is scarcely supported by experimental data *in natura*. To better understand the role and function of β-lactam and β-lactamases in natural ecosystems, it is important to develop experimental setups to study their role in the absence of clinical antibiotics selection pressure.

Soft rot *Pectobacteriaceae* (SRP) are interesting model to analyze the role of β-lactam and β-lactamases in natural ecosystems. SRP are plant pathogens belonging to the genera *Pectobacterium* and *Dickeya* within the order *Enterobacterale*. SRP disrupt the plant cell wall and induce rotting symptoms on infected host plants (Hugouvieux☐Cotte☐Pattat *et al.*, 2014; Charkowski, 2018). SRP can infect important crops such as potatoes, causing a disease called blackleg in potato fields and a severe rotting of potato tubers called soft rot, which can lead to severe post-harvest losses. SRP act as a species complex and several species are often associated within symptoms (Degefu, 2021; de Werra *et al.*, 2021; Ge *et al.*, 2021; Motyka-Pomagruk *et al.*, 2021; Barny *et al.*, 2024). Recently, one

species, *Pectobacterium versatile*, was described to harbor a β-lactamase, Bla_{PEC-1}, that is the closet relative of Bla_{TEM-1}, a class A2b β-lactamase plasmid-borne and widespread in clinical enterobacteria (Bush, 2018; Royer *et al.*, 2022). The oldest identified *P. versatile* strain carrying the Bla_{PEC-1} β-lactamase was isolated from *Solanum tuberosum* in Canada in 1918, before the discovery of the first β-lactam antibiotic in the early 20th century ruling out the possibility that the acquisition of this gene within the chromosome is linked to the clinical use of β-lactam antibiotic (Royer *et al.*, 2022). Phenotypic characterization *in vitro* revealed a low level of *P. versatile* Bla_{PEC-1} β-lactamase expression and it remains unclear whether Bla_{PEC-1} acts as a true β-lactamase during the infection process or whether it is merely a signaling molecules as previously proposed (Romero *et al.*, 2011; Royer *et al.*, 2022). Interestingly, within plant symptoms, *P. versatile* is often associated with other SRP species such as *Pectobacterium carotovorum* or *Pectobacterium brasiliense* and some strains of these latter species produce the structurally simplest member of the carbapenem class of β-lactam antibiotics, the (5R)-carbapen-2-em-3-carboxylic acid (Parker *et al.*, 1982; McGowan *et al.*, 1997, 2005). *In vitro*, this carbapenem is toxic to most *Pectobacterium* strains (Shyntum *et al.*, 2019). However, carbapenem production is repressed in anaerobic conditions *in vitro* (McGowan *et al.*, 2005; Shyntum *et al.*, 2019). Furthermore, this carbapenem is highly unstable (Parker *et al.*, 1982). Both the repression observed under anaerobic conditions and the instability of the carbapenem ring raise the question of carbapenem functional importance during potato infection.

The aim of the present work was to analyze the role of the *P. versatile* β-lactamase Bla_{PEC-1} in the context of plant infection when confronted with or not with *P. brasiliense* producing carbapenem. To this end, we constructed bla_{PEC-1}-deleted strains in two different *P. versatile* strain backgrounds and compared them with their wild-type counterparts *in vitro* and *in planta,* with mono infection or mixed infections with different *P. brasiliense* strains.

Experimental Procedures

Bacterial strains, plasmids and growth conditions

The bacterial strains, plasmids and oligonucleotides used in this study are described in Tables S1 and S2 respectively. Briefly, the following strains have been used in the study: *P. brasiliense* type strain CFBP6617 (here after 6617; other names 1692, BAA-417, BPBB-212, Ecbr-212), *P. brasiliense* CFBP5381 (here after 5381), *P. versatile* type strain CFBP6051 (here after 6051; other names ICMP9168, NCPPB3387, de Boer 21) and its Δbla_{PEC-1} mutant derivative CL1, *P. versatile* A73-S18-O15 (here after A73) and its Δbla_{PEC-1} mutant derivative CL2, *E. coli* DH5-a-l-pir, *E. coli* MFD-pir. Strains were grown routinely at 30°C in LB (10 g/L tryptone, 5 g/L yeast extract, NaCl 5g/L), in TSB (1,7 g/L casein peptone, 0.3 g/L soya peptone, 0.3 g/L NaCl, 0.25 g/L dipotassium hydrogen phosphate, 0.25 g/L glucose) or in CVP (1.02 g/L $CaCl_2$, 5 g tri-sodium citrate, 2.0 g/L NH_4NO_3, 4 g/L

agar, 2.8 ml NaOH 5M, 18 g/L pectine Dipecta-AG366-Agdia-biofords,-USA) prepared as described

by Ben Moussa *et al.*, 2023 for single layer CVP. When required, antibiotics were added at the

following concentrations: ampicillin (Amp), 200 μg/L; Chloramphenicol (Cm), 50 μg/L.;

Diaminopimelic acid (DAP) (57 μg/mL) was added for the growth of the *E. coli* MFD-pir strain.

Media were solidified with 15 g/L agar.

Construction of Δbla$_{pec-1}$ mutants

To construct in-frame deletion mutants of the *bla*$_{pec-1}$ gene, counter-selection method was used

(Edwards *et al.*, 1998). The cloning vector pRE112, that is an R6K-based suicide plasmid carrying the

*sac*B gene and the *cat* gene (CmR) was used to clone into *Sac*I/*Kpn*I digested pRE112 two PCR

fragments corresponding to the upstream and downstream 0.5-kbp DNA of the *bla*$_{pec-1}$ gene using the

T5 exonuclease-dependent assembly method (Xia *et al.*, 2019).

Chemical ultracompetent DH5α λpir cells were prepared with the Mix & Go! *E. coli* Transformation

Kit using standard procedures (Zymo Research). Transformants were selected onto LB plate

supplemented with Cm. Colonies with the correct plasmid were selected by colony PCR with oligo

pairs L762/L763 and DreamTaq DNA polymerase (Thermofisher, Waltham, MA, USA). Plasmids

were extracted with the NucleoSpin Plasmid kit (Macherey-Nagel), checked by restriction digestion

(NEB) and sequenced (Eurofins). Then, plasmids were transferred into competent *E. coli* strain

MFDpir (Ferrières *et al.*, 2010) prepared as described above for DH5α λpir. *E. coli* MFDpir produces

the RP4 conjugation machinery, which allows the transfer of the suicide plasmid into *Pectobacterium*

strains by conjugation. For conjugation, a few colonies of *P. versatile* A73 or *P. versatile* 6051 and

were mixed in the same proportion with MFDpir colonies carrying the plasmid of interest in 500 µl

LB and centrifuged for 2 min at 6000 rpm. The pellet was resuspended in 90 µl LB with 5 µl DAP at

57 mg/mL, and deposited onto a LB agar plate incubated at 28°C. After 24 h, the bacteria were

resuspended in 1 ml LB, diluted in 10-fold series from 10-1 to 10-7 and spread onto LB agar

supplemented with Cm at 4µg/l to select the first event of recombination. Transconjugants re-isolated

on this medium were then spread onto LB agar supplemented with 5% sucrose and incubated at 19°C

for 2-3 days to allow the second event of recombination. Sucrose-resistant colonies were then streak

on LB-Cm plate to check plasmid loss and to LB-Amp plate to check for *bla*$_{pec-1}$ gene loss. Total DNA

of sucrose resistant, Cm and Amp sensitive colonies was used to amplified the mutagenized locus with

the with the Prime start max DNA polymerase (Takara, kusatsu, Japan) and primers bla$_{pec-1}$-sacI-up-

fwd / bla$_{pec}$-1-KpnI-down-rev and checked by sequencing (Eurofins). The resulting Δbla$_{pec-1}$

mutants constructed in strain *P. versatile* CFPB6051 and *P. versatile* A73 were named CL1 and

CL2 respectively.

In vitro growth inhibition assay

Pectobacterium strains were grown with shaking for overnight at 28°C in TSB medium. The next day,

the optical density (OD_{600nm}) of the culture was adjusted to 0.15. The temperature of the molten LB

agar was lowered to around 40°C (just before the agar resolidified) and 25 mL of the LB agar were

mixed with 1ml of the bacteria to be tested in lawn culture. A spot of 5 μl of strain 6617 was then

added to the inoculated plates which were incubated at 30°C for 24–48 h before visualization of the

inhibition zone.

Potato tuber inoculations

Bacteria were plated on LB plates and incubated overnight at 28°C. A single colony was then used to

inoculate 2 ml of LB medium, which was incubated overnight at 28°C with agitation at 120 rpm. This

liquid culture (100 μl) was used to inoculate a 10 % TSB agar plate, which was incubated overnight at

28°C. The grown bacterial layer was then suspended in 50 mM phosphate buffer (pH 7) and adjusted

to OD_{600nm}=1. For co-inoculations, the strains involved were then mixed in equal volumes.

For each strain or strain mixture, 10 potato tubers were inoculated and each tuber was punctured with

pipette tips containing 10 μl of the inoculum. The tip was left in the tuber. As a negative control, a

pipette tip containing 10 µl of sterile phosphate buffer (50 mM) was inserted into a tuber. The

inoculated tubers were placed on paper towels in plastic boxes, 50 mL of distilled water was poured

into the bottom of the boxes and the boxes were carefully sealed to maintain relative humidity above

90%. After 5 days of incubation at 26°C, the tubers were cut and the entire macerating symptom was

collected and weighed. To count the bacterial load in the symptom, part of the symptom (50 to 300

mg) was weighed and suspended in 1 ml of sterile phosphate buffer (50 mM) and diluted in 10-fold

series and 100 µl of dilution 10^{-5} to 10^{-7} were spread on TSB plates and incubated 24h at 28°C. To

analyzed the fitness of the Δbla_{pec-1} mutant relative to the wild type strain, 100 individual colonies (per

analyzed potato tuber) were streak on CVP, TSB-Amp and TSB agar plates. Colonies that formed pits

on CVP, had the visual characteristics of *Pectobacterium* strains and grew on TSB-Amp agar plates

were defined as *P. versatile* wild-type strains. Conversely, colonies that formed pits on CVP, had the

visual characteristics of *Pectobacterium* strains and were unable to grow on TSB-Amp agar plates

were defined as *P. versatile* Δbla_{pec-1} mutant derivatives.

To recover the bacteria from rotten symptoms for DNA or RNA extraction, the rest of the symptom

was suspended with 7 ml of 50 mM phosphate buffer and stirred for 4 hours at ambient temperature.

The tubes were then left without shaking for 10 minutes to allow the excess starch to settle and 2 ml at

the surface of the suspension was recovered and centrifuged for 10 minutes at 10,000 g. The formed

pellet was kept at -80°C until DNA or RNA extraction.

RNA extraction and RT-qPCR analysis

To investigate *carA* and bla_{pec-1} expression, RNA was isolated using PureLink® RNA Mini Kit from

Thermo Scientific™, and 400 ng of RNA from potatoes symptoms or LB media were used to generate

cDNA using the Thermo-scientific maxima First Strand cDNA Synthesis Kit for RT-qPCR with
dsDNase. RT-qPCR experiments were performed with SsoAdvanced™ Universal SYBR® Green
Supermix from BIO-RAD™. The housekeeping gene *gapA* was used to normalize the expression data
for each gene of interest. The comparative quantitation method (ΔΔCt) was used to contrast the
different treatments (**Livak and Schmittgen, 2001**). Ct values quantify the number of PCR cycles
necessary to amplify a template to a chosen threshold concentration, ΔCt values quantify the
difference in Ct values between a target gene and a the *gapA* gene for a given sample, and ΔΔCt
values are used for the comparison between potato tuber symptoms and LB medium. Relative fold
changes were calculated using $2^{-\Delta\Delta Ct}$.

DNA extraction, DNA amplification and sequencing
DNA extraction was performed on 10 ml of the time zero inoculum, on 2 ml of 5 potato macerated
tubers symptoms for each inoculated strain mixture. The bacteria were pelleted by centrifugation and
DNA extraction was performed with the genomic DNA purification kit (Lucigen).
Amplification and sequencing were performed at MR DNA (www.mrdnalab.com, Shallowater, TX,
USA). Either the *gap*A 376 partial gene sequence was amplified using PCR primers gapAF376
(GCCCGTCTCACAAAGA) and gapAR (TCRTACCARGAAACCAGTT) or the 16SrRNA gene V3-
V4 region was amplified using PCR primers 341F (CCTACGGGNGGCWGCAG) and 805R
(GGACTACHVGGGTWTCTAAT) with barcodes on the forward primer. PCR using the HotStarTaq
Plus Master Mix Kit (Qiagen, USA) was performed under the following conditions: 94°C for 3 min,
followed by 28 cycles of 94°C for 30 s, 57°C for 40 s (for *gapA* amplicon) or 53°C for 40s (for 16S
amplicon) and 72°C for 1 min, after which a final elongation step at 72°C for 5 min was performed.
Amplified PCR products were checked in 2 % agarose gel to determine the success of amplification
and the relative intensity of bands. Multiple samples were pooled together in equal proportions based
on their molecular weight and DNA concentrations. Pooled samples were purified using calibrated
Ampure XP beads and used to prepare Illumina DNA library. Sequencing was performed on a MiSeq
sequencer following the manufacturer's guidelines. Sequence data were processed using MR DNA
analysis pipeline (MR DNA, Shallowater, TX, USA). In summary, sequences were joined, depleted of
barcodes and sequences <150bp or with ambiguous base calls were removed. Reads were filtered
based on Q score and expected error probability and any read with a number of expected errors greater
than 1.0 were discarded.

Illumina sequencing analysis
After quality trimming, a total of 1,296,980 and 1,238,179 reads were obtained for 30 *gap*A and 20
16S analyzed potato tubers (50 samples *in toto*) assays respectively, with an average of 43,232 ± 8,490
and 61,908 ± 18049 reads per samples respectively.

In order to quantify the distribution of the *Pectobacterium* strains at time zero within the inoculum and

5 days after potato inoculation, SRP distribution analysis was performed at the strain level as

previously described (Barny *et al.*, 2024). Briefly, following *gap*A amplification, reads were aligned

to the *gap*A sequences of the 6 strains analyzed (*P. brasiliense* 6617, *P. brasiliense* 5381, *P. versatile*

A73 wild-type or Δbla_{PEC-1} and *P. versatile* 6051 wild-type or Δbla_{PEC-1}), using the nucleotide-

nucleotide Blastn tool (version 2.15.0+, e-value threshold 10^{-5}). Only reads with 100% full-length

identity were used (616,137 reads, 47.5% of the total reads).

The clustering, alignment and phylogenetic analysis of 16S gene fragments was performed using the

QIIME2 (version 2023.5) (Bolyen *et al.*, 2019) pipeline v.1.39.5, dedicated to the study of microbial

communities. Demultiplexing of 16S rDNA gene sequences and quality control using DADA2

(Callahan *et al.*, 2016). SILVA (version 138) (Bokulich *et al.*, 2018) was used to perform taxonomic

classification of each ASV. The feature table of ASVs was rarified to a sampling depth of 32,191

sequences per sample prior to downstream analyses.

Results

***In vitro*, the β-lactamase Bla$_{PEC-1}$ allows *P. versatile* to resist to ampicillin or to the carbapenem**

produced by *P. brasiliense*

To analyze the role of Bla$_{PEC-1}$, we constructed mutants deleted of the bla_{PEC-1} gene in two strains

background of *P. versatile*, namely the type strain 6051 isolated from *Solanum tuberosum* and the

strain A73 isolated from water. As expected, the two wild-type strains were resistant to ampicillin and

the two Δbla_{PEC-1} mutants became sensitive to this antibiotic, confirming that the ß-lactamase Bla$_{PEC-1}$

confers resistance to the ß-lactam ring of the penicillin class as previously observed by Royer *et al*.

(2022) (Figure 1A). In contrast, the *P. brasiliense* 6617, which produces the carbapenem (5R)-

carbapen-2-em-3-carboxylic acid, was sensitive to ampicillin. This sensitivity indicates that the *carF*

and *carH* genes required for resistance to the carbapenem produced are inefficient against ampicillin

(McGowan *et al.*, 1997). We also tested another *P. brasiliense* strain, 5381, which was sensitive to

both ampicillin and to the carbapenem produced by *P. brasiliense* 6617. When the *P. versatile* strains

A73 or 6051 were confronted in competition assays with the carbapenem-producing strain *P.*

brasiliense 6617, the two wild-type strains were resistant, while the two Δbla_{PEC-1} mutants were

sensitive, indicating that the β-lactamase Bla$_{PEC-1}$ confers resistance to the simple carbapenem

produced by *P. brasiliense* 6617 in addition to the penicillin class of ß-lactamase ring (Figure 1B).

Figure 1: Spectrum of activity of Bla$_{PEC-1}$ β-lactamase *in vitro*
A: Growth of the strains depicted in the left panel on LB and LB ampicillin (LB + Amp) media. *P. versatile* (Pv) strains: 6051 and A73; *P. brasiliense* (Pb) strains: 6617 and 5381. B: Growth inhibition assays between the carbapenem-producing strain Pb 6617 spotted and the lawn of strain Pb 6617, Pb 5381, Pb A73, Pv 6051, Pv A73 Δ*bla*$_{PEC-1}$ mutant, Pv 6051 Δ*bla*$_{PEC-1}$ mutant.

In the absence of carbapenem, maceration capacity and fitness of *P. versatile* Δ*bla*$_{PEC-1}$ mutants in potato tubers are not altered compared to WT strains

We then investigated whether deletion of the *bla*$_{PEC-1}$ gene altered the maceration capacity of the *P. versatile* strains in the absence of carbapenem. No difference in maceration capacity was observed between each wild-type strain and its Δ*bla*$_{PEC-1}$ mutant derivative (Figure 2A). Strain A73 appeared to be slightly more aggressive than strain 6051, but this was only statistically significant when comparing the two wild-type strains. We then analyzed the fitness of each Δ*bla*$_{PEC-1}$ mutant when co-inoculated with its wild-type counterpart (Figure 2B). The different susceptibilities to ampicillin of *P. versatile* wild type strains and their Δ*bla*$_{PEC-1}$ mutant derivatives (see Figure 1A) were used to quantify the respective presence of each strain in potato tubers at the end of the co-infection experiment (Figure 2B). Within mix-inoculation, each wild-type strain and its Δ*bla*$_{PEC-1}$ mutant derivative were present in the same proportion at the end of the experiment indicating that the β-lactamase Bla$_{PEC-1}$ neither affects nor enhances the fitness of both strains in potato tubers.

Figure 2: Maceration capacity and fitness of *P. versatile* Δ*bla*_{PEC-1} mutants

A: Weight of macerate symptom following inoculations of single species or mix infections. The species or mix that were inoculated on 10 potato tubers are indicated below each boxplot. B: The different susceptibilities of *P. versatile* 6051/A73 wild-type strains and 6051/A73 Δ*bla*_{PEC-1} mutant to ampicillin was used to assess the fate of each strain within mixed infections 5 days post infection (dpi). The boxplots represent the bacterial abundance of each strain at 5 dpi. The result of the statistical analysis is indicated by the letters (a, b,) at the top of the boxplots. The bacterial mixtures sharing the same letter(s) are not statistically different from each other (P>0.05 Kruskal Wallis followed by Dunn test with the Bonferroni correction).

Coexistence of *P. versatile* wild-type strains with the carbapenem producing strain *P. brasiliense* 6617 in potato tuber

The resistance of the β-lactamase-producing wild-type strain A 73 and 6051 of *P. versatile in vitro* to the carbapenem produced by *P. brasiliense* 6617 prompted us to confront the strains *in planta*. To do this, we set up potato tuber infections with either single infection of each strain or co-infection between *P. brasiliense* and each of the wild-type *P. versatile* strains (10 tubers per condition). There was considerable variation in maceration capacity (Figure 3A). Statistical analysis showed that *P. brasiliense* strain 6617 was more aggressive than *P. versatile* 6051. Strain A73 showed an intermediate level of aggressiveness that was not statistically different from either 6051 or 6617. Co-inoculation of *P. brasiliense* 6617 with any *P. versatile* resulted in symptoms similar to single inoculation with 6617 or A73 suggesting that the mixed inoculation does not affect disease progression. The different susceptibilities of *P. brasiliense* 6617 and *P. versatile* 6051/A73 strains to ampicillin (see Figure 1A) were used to quantify the respective presence of each strain in potato tubers at the end of the co-infection experiment. After co-inoculation, *P. brasiliense* 6617 dominated the population at 5 dpi, consistent with its high level of virulence. Variability between inoculated tuber

was important, however, both *P. versatile* strains were detected at 5 dpi in 8 out of 10 and 5 out of 10

inoculated potato tubers for A73 and 6051 respectively. Overall, the mean recovery of *P. versatile*

A73 was 14,5% indicating its ability to co-exist with *P. brasiliense* 6617. In contrast, *P. versatile* 6051

was strongly outcompeted by strain *P. brasiliense* 6617 with a mean recovery of 1,2% (Figure 3B)

Figure 3: *In planta* competition between *P. versatile* strain expressing Bla$_{PEC-1}$ and the *P. brasiliense* strain
expressing carbapenem.
A: Weight of macerate symptom following inoculations of single species or mix infections. The species or mix
that were inoculated on 10 potato tubers are indicated below each boxplot. B: The different susceptibilities of *P.*
brasiliense 6617 and *P. versatile* 6051/A73 wild-type strains to ampicillin was used to assess the fate of each
strain within mixed infections 5 days post infection. The boxplots represent the bacterial abundance of each
strain (% cfu 5 dpi). The result of the statistical analysis is indicated by the letters (a, b) at the top of the
boxplots. The boxplots sharing the same letter(s) are not statistically different from each other (P>0.05 Kruskal
Wallis followed by Dunn test with the Bonferroni correction).

No association was found between commensal abundance in potato soft rot symptoms and the

presence of the *P. versatile* bla$_{PEC-1}$ gene.

During infection of potato tubers, the important release of nutrients due to degradation of plant cell

walls may allow the development of associated commensals (Kõiv *et al.*, 2015). We tested whether the

β-lactamase Bla$_{PEC-1}$ affects commensals invasion. Using 16S metabarcoding, we compared the

abundance and diversity of commensals within symptoms at the end of the experiment between 20

tubers infected with either the wild-type strains or the Δ*bla*$_{PEC-1}$ mutants (5 tubers per condition). The

presence of commensals was often limited and was less than 1 % in 16 out of 20 tubers analyzed but a

slightly higher proportion of associated commensals (from 2.1% to 14%) was observed in the 4

remaining potato tubers (Table 1). NMDS analysis indicated that commensal abundance was not

statistically different between Δbla_{PEC-1} mutants and their wild-type counterparts (Fig S1).

Strain inoculated	potato	% *Pectobacterium*	% others
A73 WT	1	99.6	0.4
	2	99.6	0.4
	3	99.1	0.9
	4	99.6	0.4
	5	99.7	0.3
A73 Δbla_{PEC-1}	1	95.3	4.7
	2	99.4	0.6
	3	99.2	0.8
	4	97.8	2.2
	5	96.7	3.3
6051 WT	1	99.7	0.3
	2	99.7	0.3
	3	99.6	0.4
	4	99.7	0.3
	5	86.1	13.9
6051 Δbla_{PEC-1}	1	99.7	0.3
	2	99.6	0.4
	3	99.4	0.6
	4	99.7	0.3
	5	99.6	0.4

Table 1: Percentage of *Pectobacterium* and commensal at 5 dpi. Analysis was performed for each individual potato
inoculated 5dpi (five replicates).

We then analyzed the nature of the commensals recovered within the symptoms. The most common

commensals to the 20 tubers analyzed were unclassified *Enterobacteriaceae* and bacteria belonging to

the genus *Anaerocolumna* (Figure 4). Other commensals varied stochastically between potato tubers

(Figure 4). In conclusion, the presence of commensals within symptoms was variable between tubers

and did not correlate with the presence of the bla_{PEC-1} gene.

Figure 4: Commensal associated bacteria
Type of associated commensals in each analyzed potato tuber 5 days post infection following 16S barcode analysis. The inoculated strain and the potato number is indicated below each bar. The taxonomic assignment has been set at the genus level (g) or higher when this was not possible (f: family, o: order).

Expression of carbapenem and β-lactamase encoding genes within potato tuber

To compared the level of expression of the genes involved in the synthesis of carbapenem or Bla_{PEC-1} β-lactamase during infection of potato tubers and LB synthetic media, we set up a mono-infection experiment with either the *P. brasiliense* strain 6617 or the *P. versatile* strains 6051 and A73. The expression levels of the *bla*$_{PEC-1}$ and *carA* genes involved in the production of Bla_{PEC-1} β-lactamase and carbapenem respectively, were quantified by RT-PCR (McGowan *et al.*, 1997; Royer *et al.*, 2022). Large variations were observed between potato tubers. However, the *carA* gene of *P. brasiliense* 6617 was consistently strongly repressed within potato tubers as compared with LB media (Figure 5). Similarly, the *bla*$_{PEC-1}$ gene was also repressed within potato tubers as compared with LB media but to a lesser extent than the one observed with the *carA* gene.

Figure 5: Relative expression of carbapenem and β-lactamase encoding genes within potato tuber

$2^{-\Delta\Delta Ct}$ values were used to express the fold change in *car*A (Pb 6617 strain) and *bla*$_{PEC-1}$ (Pv A73 and 6051 strains) gene expression within potato tuber symptoms relative to expression in LB medium. A relative fold change of less than 1 indicates repression *in planta* compared to LB conditions *in vitro*. Expression data have been normalized using the housekeeping gene *gap*A as a reference gene.

***P. versatile bla*$_{PEC-1}$ gene is required during co-infection with carbapenem-producing *P. brasiliense* to protects another *P. brasiliense* strain sensitive to carbapenem.**

During infection, mixtures of SRP spp. can account for up to 20% of the observed symptoms (Degefu, 2021; de Werra *et al.*, 2021; Ge *et al.*, 2021; Motyka-Pomagruk *et al.*, 2021). We therefore wondered whether the *P. versatile* β-lactamase Bla$_{PEC-1}$ could protect other *Pectobacterium* strains from the carbapenem produced by the *P. brasiliense* strain 6617 (Shyntum *et al.*, 2019). *In vitro*, the carbapenem-sensitive strain *P. brasiliense* 5381 was exposed to the carbapenem-producing strain 6617 in the vicinity of either the *P. versatile* β-lactamase-producing strain 6051 or its Δ*bla*$_{PEC-1}$ mutants derivative. The presence of the *P. versatile* strain 6051, but not of the 6051 Δ*bla*$_{PEC-1}$ mutant, protected the sensitive strain 381 from the toxic effect of *P. brasiliense* strain 6617 (Figure 6). This indicates that, *in vitro*, the β-lactamase Bla$_{PEC-1}$ secreted by the WT strain of *P. versatile* is able to degrade the carbapenem produced and secreted by *P. brasiliense* 6617 efficiently enough to protect the sensitive strain 5381 when this latter strain is close enough to the β-lactamase PEC-1 produced by *P. versatile*.

Figure 6: *In vitro*, the β-lactamase Bla$_{PEC-1}$ confers protection against carbapenem
The *P. brasiliense* 5381 strain grown in the lawn is sensitive to the carbapenem produced by the *P. brasiliense* 6617 strain spotted in the middle of the Petri dish. The wild-type *P. versatile* strain 6051 spotted on the left, protects the sensitive strain only when the spot is in close proximity (compare upper and lower spots). None of the spots of the 6051 Δbla$_{PEC-1}$ mutant of *P. versatile*, on the right, confers protection.

We then ask whether the protection conferred by *P. versatile* β-lactamase producing strains is also observed *in planta*. Potato tuber co-infection experiments were set up with 1) the *P. brasiliense* 6617 strain producing carbapenem, 2) the *P. brasiliense* 5381 strain sensitive to carbapenem and 3) either the wild-type *P. versatile* strains A73 or 6051 or their Δ*bla*$_{PEC-1}$ mutant derivatives. As control, we also tested whether the carbapenem-sensitive strain 5381 is maintained when co-inoculated alone with the carbapenem producing strain 6617. The proportion of each strain at the beginning and end of the experiment was assessed by Illumina sequencing of the *gap*A discriminating gene marker (Barny *et al.*, 2024). When the carbapenem-sensitive 5381 strain was co-inoculated alone with the carbapenem-producing strain 6617, it was outcompeted and the only strain recovered at the end of the experiment was the carbapenem-producing strain 6617 (Figure 7A). As well, in the mixture involving 6617 + 5381 and either the A73 Δ*bla*$_{PEC-1}$ mutant or the 6051 Δ*bla*$_{PEC-1}$ mutant, only the carbapenem-producing 6617 was recovered at the end of the experiment (Figure 7 B and C). In contrast, in the presence of the wild type *P. versatile* strains A73 or 6051, both the *P. versatile* strains and the sensitive strain 5381 were detect at various level the end of the experiment in 9 out of the 10 analyzed potato tubers (Figure 7 B and C). This indicated that the β-lactamase Bla$_{PEC-1}$ protect both the *P. versatile* strain itself and the carbapenem sensitive strain 5381 from the toxic effect of the carbapenem produced by 6617 as observed *in vitro*. The protection conferred by strain 6051 was less than that

conferred by strain A73, consistent with its weaker ability to compete with strain 6617. However, no

correlation was observed between the proportion of *P. versatile* wild-type strains and the proportion of

the carbapenem-sensitive strain 5381 recovered within the symptoms at 5dpi (Figure 7 B and C). This

lack of correlation is likely explained by spatial effects within soft rot symptoms, as *in vitro* rescue of

the carbapenem-sensitive strain 5381 strongly depends on spatial effects as observed *in vitro* (Figure

6).

Figure 7: Fate of SRP consortium in mixed inoculation on potato tubers

The consortium inoculated (T0) is compared with the mix observed 5 days post inoculation in 5 independent

potato tubers. The strains involved in each consortium are followed by analysis of the *gap*A barcode (Barny *et*

al., 2024). Carbapenem producing strain are depicted in blue, carpapenem sensitive strain are depicted in red and

β-lactamase producing strains are depicted in yellow. A: mix between the carbapenem producing *P. brasiliense*
6617 strain and the carbapenem sensitive *P. brasiliense* 5381 strain. B: mix between the carbapenem-producing
P. brasiliense 6617 strain, the carbapenem-sensitive *P. brasiliense* 5381 strain and either the *P. versatile* A73
WT strain expressing the β-lactamase Bla_{PEC-1} or its Δbla_{PEC-1} mutant derivative. C: mix between the
carbapenem-producing *P. brasiliense* 6617 strain, the carbapenem-sensitive *P. brasiliense* 5381 strain and either
the *P. versatile* 6051 WT strain expressing the β-lactamase Bla_{PEC-1} or its Δbla_{PEC-1} mutant derivative.
448

Discussion

The β-lactamase Bla_{PEC-1} is the closest relative of Bla_{TEM-1}, a class A2b β-lactamase that is plasmid-
borne and widespread in clinical enterobacterales. Bla_{PEC-1}, like Bla_{TEM-1} cleave β-lactam rings of the
penicillin family but is inefficient on clinically used carbapenem such as meropenem or imipenem
(Royer *et al.*, 2022). Here we have shown, using *in vitro* competition assays, that the β-lactamase
Bla_{PEC-1} is nevertheless efficient in inactivating the simplest of the carbapenem molecule, produced by
P. brasiliense. This suggests that the inability of Bla_{PEC-1} to inactivate imipenem and meropenem is
probably due to the steric hindrance provided by the C-6 side chain of these clinically useful
carbapenems. It remains to be determined whether the closest Bla_{PEC-1} β-lactamase, Bla_{TEM-1} that is
clinically widespread around the world, is also able to cleave this simplest carbapenem molecule.
We then analyzed the role of the *P. versatile* β-lactamase Bla_{PEC-1} in the context of plant infection in
the absence of antibiotic pressure. *In planta*, during mono-infection experiments, we could not detect a
fitness cost associated with the presence of Bla_{PEC-1} as the wild-type and the bla_{PEC-1} -deleted strains
multiplied similarly within symptoms**.** The fitness cost of a resistance determinant has been shown to
partly depend on its genetic localization, such as a chromosomal mutation or a plasmid acquisition
(Vanacker *et al.*, 2023). The absence of fitness cost of associated with the bla_{PEC-1} gene could probably
be partly due to the chromosomal location of this gene within the *P. versatile* genomes (Royer *et al.*,
2022). It has also been shown that the fitness cost of antibiotic resistance genes is lower for β-
lactamases compared to mutations in core genes which have a more pleiotropic effect that could be
deleterious in the absence of antibiotic pressure (Vanacker *et al.*, 2023). The absence of detectable
fitness costs for chromosomally encoded β-lactamases in the absence of antibiotic pressure probably
explains the wide distribution of these enzymes within natural communities. For example, in remote,
undisturbed Alaskan soils with no antibiotic selection pressure, 5% of bacterial genomes carry diverse
β-lactamase genes that are readily expressed in *Escherichia coli* (Allen *et al.*, 2009). Similarly,
permafrost bacteria represent a vast reservoir of antibiotic resistance genes, among which β-lactamase
genes have been observed at high frequencies (Rigou *et al.*, 2022).
Whether β-lactamases play a true role in antibiotic resistance in natural environments has been
debated due to their low concentration (Davies *et al.*, 2006; Aminov, 2009). *In vitro* phenotypic
characterization of *P. versatile* revealed that Bla_{PEC-1} expression levels were low compared to that of
Bla_{TEM-1} (Royer *et al.*, 2022), suggesting that the role of Bla_{PEC-1} during infection could be merely a

signaling role as previously proposed (Romero *et al.*, 2011). Quantification of bla_{PEC-1} expression

within the potato tuber showed that bla_{PEC-1} is repressed compared to its LB medium expression level.

At the same time, the expression level of the *carA* gene, which is involved in the biosynthesis of the

carbapenem by strain 6617, is also strongly repressed within the potato tuber. This is consistent with

the dramatic repression of carbapenem expression observed *in vitro* in anaerobic medium (McGowan

et al., 2005; Shyntum *et al.*, 2019). Consistent with anaerobiosis, we found that commensals of the

genus *Anaerocolumna*, a strictly anaerobic genus (Ueki et al., 2016), are systematically found

associated with rotten symptoms in potato tubers. Surprisingly, despite the strong repression of *car*A

in potato tubers, we found, that a carbapenem-sensitive strain was strongly outcompeted and could not

be recovered when mixed inoculated with the carbapenem-producing strain. This indicates that

carbapenem production is still efficient during infection, even though its expression is severely

repressed compared to LB medium. The effectiveness of carbapenem in the potato tuber was also

unexpected, as the carbapenem produced is also a highly unstable molecule. (Parker et al., 1982). This

showed that lower expression level than the one observed *in vitro* and instability are not sufficient to

rule out functionality and highlights the need for careful analysis to understand the effect of antibiotic

production in natural environments.

When a *P. versatile* β-lactamase producing strain was added to the consortium, this third strain was

able to rescue the carbapenem sensitive strain from the deleterious effect of the carbapenem producing

strain. Such protection was space dependent and was observed *in vitro* only when the 3 strains were in

close proximity. *In planta*, the protection was effective even when the *P. versatile* strain was a

minority within the symptoms and the protection level, exemplified by the maintenance of the

carbapenem sensitive strain, did not correlate with the proportion of each strain (protective, sensitive

and antibiotic producing) within the symptoms. This likely highlights the formation of specific micro

niches within the symptoms. This is in agreement with the fact that stochasticity in the initial phase of

colony formation can be a crucial factor that determines the ensuing population dynamic (Von Bronk

et al., 2017).

Within the SRP species complex, *P. versatile* occupies a special place as it is the most abundant SRP

species, both on crops and in the environment, highlighting its ecological success (Portier *et al.*, 2020;

Ben Moussa *et al.*, 2022). Among SRP species, *P. versatile* is also the only species that harbors the β-

lactamase Bla_{PEC-1} at high frequency (Royer *et al.*, 2022), suggesting that Bla_{PEC-1} may be involved in

its ecological success. However, *P. versatile* is not considered to be a major pathogen of the SRP

species complex, but rather a companion species during epidemics (van der Wolf *et al.*, 2021). The

work performed here indicates that Bla_{PEC-1} exerts a true β-lactamase function that is efficient at

microscale during the infection process and highlights the role of *P. versatile* as a companion species

that maintains strain diversity within the SRP species complex. The maintenance of such diversity is

crucial to allow the regular emergence of new epidemic clones, such as those regularly observed on

the potato host (Jonkheer *et al.*, 2021). Beyond the SRP species complex, the high occurrence of

various β-lactamases in natural pristine environments has recently been described (Lima-Bittencourt *et*
al., 2007; Pedroso *et al.*, 2017; Fonseca *et al.*, 2018; Rigou *et al.*, 2022). The work performed here
unravels the role of β-lactamase in maintaining the diversity of β-lactam susceptible strains in the
environment.

Acknowledgments

521 Authors acknowledge Helena Dutey for her help in performing some plant experiments. The authors
522 also acknowledge the funding from the ANR project ANR-19-CE35-0016-03. The funders had no role
523 in study design, data collection and interpretation, or the decision to submit the work for publication.

References

Allen, H.K., Moe, L.A., Rodbumrer, J., Gaarder, A., and Handelsman, J. (2009) Functional metagenomics
reveals diverse β-lactamases in a remote Alaskan soil. *ISME J* **3**: 243–251.

Aminov, R.I. (2009) The role of antibiotics and antibiotic resistance in nature. *Environmental Microbiology* **11**:
2970–2988.

Barny, M.-A., Thieffry, S., Gomes de Faria, C., Thebault, E., and Pédron, J. (2024) Bacterial pathogens dynamic
during multi-species infections. *Peer Community Journal* **4**: e49.

Ben Moussa, H., Bertrand, C., Rochelle-Newall, E., Fiorini, S., Pédron, J., and Barny, M.-A. (2022) The
diversity of soft rot *Pectobacteriaceae* along the Durance river stream in the south-east of France revealed by
multiple seasonal surveys. *Phytopathology*® PHYTO-12-21-0515-R.

Ben Moussa, H., Pédron, J., Hugouvieux□Cotte□Pattat, N., and Barny, M. (2023) Two species with a peculiar
evolution within the genus *Pectobacterium* suggest adaptation to a new environmental niche. *Environmental*
Microbiology 1462-2920.16479.

Bokulich, N.A., Kaehler, B.D., Rideout, J.R., Dillon, M., Bolyen, E., Knight, R., et al. (2018) Optimizing
taxonomic classification of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin.
Microbiome **6**: 90.

Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., et al. (2019)
Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol*
37: 852–857.

Bush, K. (2018) Past and present perspectives on β-Lactamases. *Antimicrob Agents Chemother* **62**: e01076-18.

Bush, K. and Bradford, P.A. (2016) β-Lactams and β-Lactamase inhibitors: An overview. *Cold Spring Harb*
Perspect Med **6**: a025247.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., and Holmes, S.P. (2016) DADA2:
High-resolution sample inference from Illumina amplicon data. *Nat Methods* **13**: 581–583.

Cantón, R., Novais, A., Valverde, A., Machado, E., Peixe, L., Baquero, F., and Coque, T.M. (2008) Prevalence
and spread of extended-spectrum β-lactamase-producing *Enterobacteriaceae* in Europe. *Clinical*
Microbiology and Infection **14**: 144–153.

Charkowski, A.O. (2018) The changing face of bacterial soft-rot diseases. *Annu Rev Phytopathol* **56**: 269–288.

Davies, J., Spiegelman, G.B., and Yim, G. (2006) The world of subinhibitory antibiotic concentrations. *Current*
Opinion in Microbiology **9**: 445–453.

Degefu, Y. (2021) Co-occurrence of latent *Dickeya* and *Pectobacterium* species in potato seed tuber samples
from northern Finland. *AFSci* **30**:.

Edwards, R.A., Keller, L.H., and Schifferli, D.M. (1998) Improved allelic exchange vectors and their use to
analyze 987P fimbria gene expression. *Gene* **207**: 149–157.

Ferrières, L., Hémery, G., Nham, T., Guérout, A.-M., Mazel, D., Beloin, C., and Ghigo, J.-M. (2010) Silent
mischief: bacteriophage Mu Insertions contaminate products of *Escherichia coli* random mutagenesis
performed using suicidal transposon delivery plasmids mobilized by broad-host-range RP4 conjugative
machinery. *J Bacteriol* **192**: 6418–6427.

Fonseca, E.L., Andrade, B.G.N., and Vicente, A.C.P. (2018) The resistome of low-impacted marine
environments is composed by distant metallo-β-Lactamases homologs. *Front Microbiol* **9**: 677.

Ge, T., Ekbataniamiri, F., Johnson, S.B., Larkin, R.P., and Hao, J. (2021) Interaction between *Dickeya*
dianthicola and *Pectobacterium parmentieri* in potato infection under field conditions. *Microorganisms* **9**:
316.

Hugouvieux□Cotte□Pattat, N., Condemine, G., and Shevchik, V.E. (2014) Bacterial pectate lyases, structural
and functional diversity. *Environmental Microbiology Reports* **6**: 427–440.

Jonkheer, E.M., Brankovics, B., Houwers, I.M., van der Wolf, J.M., Bonants, P.J.M., Vreeburg, R.A.M., et al.
(2021) The *Pectobacterium* pangenome, with a focus on *Pectobacterium brasiliense*, shows a robust core and
extensive exchange of genes from a shared gene pool. *BMC Genomics* **22**: 265.

Kim, Dogyeoung, Kim, S., Kwon, Y., Kim, Y., Park, H., Kwak, K., et al. (2023) Structural Insights for β-
Lactam Antibiotics. *Biomol Ther (Seoul)* **31**: 141–147.

Kõiv, V., Roosaare, M., Vedler, E., Ann Kivistik, P., Toppi, K., Schryer, D.W., et al. (2015) Microbial
population dynamics in response to *Pectobacterium atrosepticum* infection in potato tubers. *Sci Rep* **5**:
11606.

Lima-Bittencourt, C.I., Cursino, L., Gonçalves-Dornelas, H., Pontes, D.S., Nardi, R.M.D., Callisto, M., et al.
(2007) Multiple antimicrobial resistance in *Enterobacteriaceae* isolates from pristine freshwater. *Genet Mol*
Res **6**: 510–521.

Livak, K.J. and Schmittgen, T.D. (2001) Analysis of relative gene expression data using real-time quantitative
PCR and the 2−ΔΔCT Method. *Methods* **25**: 402–408.

McGowan, S.J., Barnard, A.M.L., Bosgelmez, G., Sebaihia, M., Simpson, N.J.L., Thomson, N.R., et al. (2005)
Carbapenem antibiotic biosynthesis in *Erwinia carotovora* is regulated by physiological and genetic factors
modulating the quorum sensing□dependent control pathway. *Molecular Microbiology* **55**: 526–545.

McGowan, S.J., Sebaihia, M., O'Leary, S., Hardie, K.R., Williams, P., Stewart, G.S.A.B., et al. (1997) Analysis
of the carbapenem gene cluster of *Erwinia carotovora*□: definition of the antibiotic biosynthetic genes and
evidence for a novel β□lactam resistance mechanism. *Molecular Microbiology* **26**: 545–556.

Motyka-Pomagruk, A., Zoledowska, S., Sledz, W., and Lojkowska, E. (2021) The occurrence of bacteria from
different species of Pectobacteriaceae on seed potato plantations in Poland. *Eur J Plant Pathol* **159**: 309–325.

Parker, W.L., Rathnum, M.L., Wells, J.J.S., Trejo, W.H., Principe, P.A., and Sykes, R.B. (1982) SQ 27, 860, A
simple carbapenem produced by species of *Serratia* and *Erwinia*. *J Antibiot* **35**: 653–660.

Pedroso, M.M., Selleck, C., Enculescu, C., Harmer, J.R., Mitić, N., Craig, W.R., et al. (2017) Characterization of
a highly efficient antibiotic-degrading metallo-β-lactamase obtained from an uncultured member of a
permafrost community. *Metallomics* **9**: 1157–1168.

Portier, P., Pédron, J., Taghouti, G., Dutrieux, C., and Barny, M.-A. (2020) Updated taxonomy of
Pectobacterium genus in the CIRM-CFBP bacterial collection: when newly described species reveal "Old"
endemic population. *Microorganisms* **8**: 1441.

Rigou, S., Christo-Foroux, E., Santini, S., Goncharov, A., Strauss, J., Grosse, G., et al. (2022) Metagenomic
survey of the microbiome of ancient Siberian permafrost and modern Kamchatkan cryosols. *microLife* **3**:
uqac003.

Romero, D., Traxler, M.F., López, D., and Kolter, R. (2011) Antibiotics as signal molecules. *Chem Rev* **111**:
5492–5505.

Royer, G., Dixit, Z., Pédron, J., Pierrat, G., Demontant, V., Berçot, B., et al. (2022) Genetic and phenotypic
study of the *Pectobacterium versatile* beta-lactamase, the enzyme most similar to the plasmid-encoded TEM-
1. *Appl Environ Microbiol* **88**: e00220-22.

Shyntum, D.Y., Nkomo, N.P., Shingange, N.L., Gricia, A.R., Bellieny-Rabelo, D., and Moleleki, L.N. (2019)
The impact of type VI secretion system, bacteriocins and antibiotics on bacterial competition of
Pectobacterium carotovorum subsp. *brasiliense* and the regulation of carbapenem biosynthesis by iron and
the ferric-uptake regulator. *Front Microbiol* **10**:.

Tooke, C.L., Hinchliffe, P., Bragginton, E.C., Colenso, C.K., Hirvonen, V.H.A., Takebayashi, Y., and Spencer,
J. (2019) β-Lactamases and β-Lactamase inhibitors in the 21st Century. *Journal of Molecular Biology* **431**:
3472–3500.

Vanacker, M., Lenuzza, N., and Rasigade, J.-P. (2023) The fitness cost of horizontally transferred and
mutational antimicrobial resistance in *Escherichia coli*. *Front Microbiol* **14**: 1186920.

Von Bronk, B., Schaffer, S.A., Götz, A., and Opitz, M. (2017) Effects of stochasticity and division of labor in
toxin production on two-strain bacterial competition in *Escherichia coli*. *PLoS Biol* **15**: e2001457.

de Werra, P., Debonneville, C., Kellenberger, I., and Dupuis, B. (2021) Pathogenicity and relative abundance of
Dickeya and *Pectobacterium* species in Switzerland: An Epidemiological Dichotomy. *Microorganisms* **9**:
2270.

van der Wolf, J.M., Acuña, I., De Boer, S.H., Brurberg, M.B., Cahill, G., Charkowski, A.O., et al. (2021)
Diseases caused by *Pectobacterium* and *Dickeya* species around the world. In *Plant Diseases Caused by*
Dickeya and Pectobacterium Species. Van Gijsegem, F., van der Wolf, J.M., and Toth, I.K. (eds). Cham:
Springer International Publishing, pp. 215–261.

Xia, Y., Li, K., Li, J., Wang, T., Gu, L., and Xun, L. (2019) T5 exonuclease-dependent assembly offers a low-
cost method for efficient cloning and site-directed mutagenesis. *Nucleic Acids Research* **47**: e15.

Data Accessibility

629 16S and *gap*A amplicon sequences are available at Zenodo (https://doi.org/10.5281/zenodo.14832342)

Author contributions

632 M.A.B. and J.P. designed the research. E.G, M.A.B, and J.P. supervised research. C.L and P.Y.C

633 performed the experiments. M.A.B and. J.P. analyzed the data. J.P. and P.Y. created the figures.

634 M.A.B. wrote the first draft and all authors reviewed and edited the manuscript. M.A.B. was in charge

635 of the funding acquisition.

Supplementary Figure and tables

Figure S1: Principal components analysis of commensals.

Principal component analysis was performed on the abundance matrix of commensal bacterial genera, 5

replicates per strain. Only 5 samples out of 20 (in blue) stand out, without forming a coherent group. For these

samples, 2 variables (in green) are explanatory (genera *Anaerocolumna* and *Enterobacteriaceae*).

Table S1: Bacterial strains and plasmids

Bacterial strains and plasmids	Description	Source and reference
Strains		
Escherichia coli K12		
DH5α	*supE44 lacU169 (Φ80lacZΔ M15) hsdR17 (rK mK) recA1 endA1 gyrA96 thi-1 relA1*	Laboratory collection
DH5α λpir	λpir phage lysogen of DH5α	E. Gueguen
MFD*pir*	*RP4-2-Tc::(ΔMu1::aac(3)IV-·ΔaphA-·Δnic35-ΔMu2::zeo) ΔdapA::erm-pir) ΔrecA*	E. Gueguen [1]
Pectobacterium versatile		
CFBP6051	Type strain, isolated from *solanum tuberosum* in Netherland, Amp^R	CIRM-CFBP [2]
CL1	CFBP6051Δ*bla*$_{pec-1}$, Amp*S*	This study

A73	Isolated from river water, Amp^R	Laboratory collection [3]
CL2	A73 Δ*bla*pec-1, Amp*S*	This study
Pectobacterium brasiliense		
CFBP6617	Type strain, isolated from *solanum tuberosum* in Brazil, Amp^S	CIRM-CFBP [4]
CFBP5381	Isolated from *solanum tuberosum* in Algeria, Amp$^{\underline{S}}$	CIRM-CFBP [4]
Plasmids		
pRE112	Suicide vector for allelic exchange, Cm^R, *sacB*, *oriT* RP4, *ori*R6K	E. Gueguen [5]
pCL1	pRE112- Δ*bla*pec-1 (ΔCL1_sequences à déposer), Cm^R	This study
pCL2	pRE112- Δ*bla*pec-1 (ΔCL2_ séquences à déposer), Cm^R	This study

1.   Ferrières L, Hémery G, Nham T, Guérout A-M, Mazel D, Beloin C, et al. Silent Mischief: Bacteriophage Mu
Insertions Contaminate Products of *Escherichia coli* Random Mutagenesis Performed Using Suicidal Transposon
Delivery Plasmids Mobilized by Broad-Host-Range RP4 Conjugative Machinery. J Bacteriol. 2010;192: 6418–6427.
doi:10.1128/JB.00621-10
2.   Portier P, Pédron J, Taghouti G, Fischer-Le Saux M, Caullireau E, Bertrand C, et al. Elevation of
Pectobacterium carotovorum subsp. *odoriferum* to species level as *Pectobacterium odoriferum* sp. nov., proposal
of *Pectobacterium brasiliense* sp. nov. and *Pectobacterium actinidiae* sp. nov., emended description of
Pectobacterium carotovorum and description of *Pectobacterium versatile* sp. nov., isolated from streams and
symptoms on diverse plants. Int J Syst Evol Microbiol. 2019;69: 3207–3216. doi:10.1099/ijsem.0.003611
3.   Ben Moussa H, Bertrand C, Rochelle-Newall E, Fiorini S, Pédron J, Barny M-A. The diversity of soft rot
Pectobacteriaceae along the Durance River stream in the south-east of France revealed by multiple seasonal
surveys. Phytopathology. 2022;. doi:10.1094/PHYTO-12-21-0515-R
4.   Liu Y, Filiatrault MJ. Complete Genome Sequence of the Necrotrophic Plant-Pathogenic Bacterium
Pectobacterium brasiliense 1692. Baltrus DA, editor. Microbiol Resour Announc. 2020;9: e00037-20.
doi:10.1128/MRA.00037-20
5.   Edwards RA, Keller LH, Schifferli DM. Improved allelic exchange vectors and their use to analyze 987P
fimbria gene expression. Gene. 1998;207: 149–157. doi:10.1016/S0378-1119(97)00619-7

Table S2: Oligonucleotides used in this study

Oligonucleotide	Sequence (5'-3')	Use
L762	gttattggtgcccttaaacg	Primers to check the correct cloning into pRE112. Amplification of the cloned region send to sanger sequencing (eurofin)
L763	gcatccaacgccattcatgg	
gapAF376	GCCCGTCTCACAAAGA	Amplification of *gapA* 376bp partial gene sequence for Illumina sequencing
gapAR	TCRTACCARGAAACCAGTT	
341F	CCTACGGGNGGCWGCAG	Amplification of the V3-V4 region of the 16SrRNA gene for iIlumina sequencing
805R	GGACTACHVGGGTWTCTAAT	
blapec-1**-sacI-up-fwd**	AACTGCATGaattcccgggagagctcttcg	Amplification of the upstream 0.5-kb DNA fragments of *P. versatile A73 and CFBP6051 for* cloning into pRE112
blapec-1**-up-rev**	gTTACCAGTGCTTGAAATGTTGCATattaa	
blapec-1**-down-fwd**	ATGCAACATTTCAAGCACTGGTAAcaatca	Amplification of the downstream 0.5-kb DNA fragments of *P. versatile A73 and CFBP6051* and cloning into pRE112
blapec-1**-KpnI-down-rev**	gatcccaagcttcttctagaggtaccagat	
***bla*PEC-1_fwd**	AAA GAT ACT TAC TGT CGC C	RT-qPCR - *bla*PEC-1 gene

*bla*_{PEC-1}**rev**	TAT CCG CAA TAA ACC AGC C	
carA-**fwd**	AGT GTC AGA TGG AAA GAG GA	RT-qPCR-*carA* gene
carA-**rev**	GAC ATG AAC CGG CGA GAA	
gapA-**fwd**	GTA TCG GCC GTA TTG TTT T	RT-qPCR-*gapA* gene
gapA-**rev**	TTC GTT CCA CTT CAG GTT	