Behavioural evidence of a humidistat: a temperature-compensating mechanism of

hydroregulation

Danilo Giacometti[1,*] and Glenn J. Tattersall[1]

[1] Department of Biological Sciences, Brock University, St. Catharines, ON L2S 3A1, Canada

[*] d.giacometti96@gmail.com

Abstract

The ability to control hydration state essential for terrestrial species, especially amphibians,

which are highly susceptible to dehydration. Here, we examined how temperature (17°C vs.

22°C) influenced behavioral hydroregulation in spotted salamanders (*Ambystoma maculatum*)

using a laboratory humidity gradient. Salamanders defended a constant vapour pressure deficit

(VPD) between temperatures by targeting higher RH at 22°C than at 17°C, possibly to

compensate for increased evaporative demand at warmer temperatures. Individuals selecting

higher VPDs experienced greater evaporative water loss (EWL), with larger salamanders losing

more water than smaller ones after accounting for temperature. Together, these results highlight

a trade-off among body size, humidity preference, and desiccation tolerance. Salamanders also

rehydrated faster at 22°C than 17°C, highlighting temperature-dependent water uptake rates. Our

finding that salamanders regulated a constant driving force of evaporation between temperatures

suggests the ability to detect rates of EWL. Local cooling of the skin arising from evaporation is

a plausible mechanism: if moist-skinned ectotherms allow for a specific, fixed decline in skin

relative to core temperature, they can effectively detect the drive for evaporation. Ultimately, our

study underscores the complexity of amphibian hydroregulation and emphasises the role of

behaviour in maintaining hydration state.

Keywords: amphibian, behaviour, ecophysiology, hydroregulation, temperature, water balance

Introduction

The transition from aquatic to terrestrial environments was one of the most significant milestones in vertebrate evolution (Laurin, 2010). On land, water availability fluctuates both spatially and temporally, meaning that organisms face constant challenges related to the maintenance of water balance (Takei, 2015). Therefore, the ability to control hydration state is essential for terrestrial species. Effective hydroregulation may be achieved through a combination of physiological and behavioural strategies. Physiological processes include the enhancement of skin resistance to water loss (Tracy et al., 2008), the presence of waterproofing lipids in the integument (Lillywhite, 2006), or the modulation of kidney and bladder function to increase water retention (Schmidt-Nielsen and Schmidt-Nielsen, 1952; Shoemaker and Bickler, 1979). Hydroregulatory behaviours may comprise the adoption of water-conserving postures (Pough et al., 1983), aggregation (Broly et al., 2014), or burrowing (Bulova, 2002). Often, however, behavioural hydroregulation is oversimplified to the selection of humid microhabitats (but see Dezetter et al., 2023). As a result, we still have limited information about how animals perceive and process changes in habitat humidity, and how behaviour can be used to mitigate potential disruptions to the maintenance of water balance (Hillyard and Willumsen, 2011; Hillyard et al., 1998).

Among vertebrate ectotherms, amphibians in particular face elevated hydroregulatory challenges. Truly, amphibians have a relatively thin skin that is supported by blood vessels in the epidermis, which together facilitate cutaneous gas exchange (Tattersall, 2007). However, the ability to exchange gases through the skin also makes amphibians highly susceptible to evaporative water loss (EWL), especially at high temperatures (Lertzman-Lepofsky et al., 2020). The extent to which temperature impacts EWL depends on the vapour pressure deficit (VPD), which measures the drying power of air, and is determined by the difference between the

saturated water vapour pressure (WVP_{sat}) and the actual water vapour pressure (WVP_{act})

(Anderson, 1936). Since warm temperatures typically produce high VPD, amphibians facing

these conditions should suffer high evaporative demand and pay high hydroregulatory costs

(Riddell et al., 2019; Riddell et al., 2024). Because of this, many have argued that thermal

constraints over hydroregulation explained the recurrent pattern of low body temperatures (T_bs)

in amphibians, on the basis that sustaining low T_bs would minimise EWL (Brattstrom, 1979).

This view has been recently expanded into a framework that posits that amphibians constantly

balance thermoregulation and hydroregulation to sustain performance (Greenberg and Palen,

2021; Titon and Gomes, 2017). However, while this framework has both thermosensation and

hydrosensation components, our current knowledge of the former outweighs the latter (Ortega et

al., 2023; Tracy et al., 2008), particularly in terms of the behaviours that contribute to the

maintenance of hydration state.

One possible explanation for this knowledge gap is the difficulty in disentangling

temperature and humidity effects, since most laboratory studies test amphibians in combined

humidity and thermal gradients (hereafter "hydrothermal gradients") (Bundy and Tracy, 1977;

Galindo et al., 2018). While one may argue that the use of hydrothermal gradients provides a

realistic scenario, any response measured in such an apparatus is by definition confounded by

temperature and humidity (Mitchell and Bergmann, 2016). Because one cannot properly gauge

whether the observed behaviours are innately hydroregulatory, the extensive use of hydrothermal

gradients resulted in the simplistic yet prevalent view that, for amphibians, "wetter is better" (see

Brattstrom, 1979). However, if one is interested in assessing humidity preference and the

behaviours that underpin hydroregulation, one should test amphibians in a temperature-

controlled humidity gradient (Heatwole, 1962; Shelford, 1913). For example, Spotila (Spotila,

1972) demonstrated interspecific differences in humidity preference (RH selection range: 60.4–

90% at 15°C) across 12 closely-related species of salamanders. Humidity preference was

associated with microhabitat use, suggesting that salamanders were able to detect differences in

air moisture content and respond in a manner that maximised hydration state (Spotila, 1972).

Despite their potential, temperature-controlled humidity gradients have seldom been used in

amphibian research since the 1970s (but see Chapman and Bidwell, 2023). As such, our

understanding of behavioural hydroregulation remains limited.

In this study, we used a laboratory humidity gradient to assess how temperature (17°C vs.

22°C) affected behavioural hydroregulation in the spotted salamander (*Ambystoma maculatum*).

We focussed on two main behavioural metrics: VPD selection and RH selection. By measuring

VPD selection, one is able to obtain behavioural information based on a temperature-

independent humidity gradient and gauge behavioural hydroregulation *sensu stricto* (Dezetter et

al., 2023; Wolcott and Wolcott, 2001). In contrast, RH selection provides information on how

changes in temperature impact the behaviours that contribute to the maintenance of hydration

state (Galindo et al., 2018; Spotila, 1972). Based on this, we predicted that *A. maculatum* would

alter RH selection to maintain a constant VPD between temperatures (Spotila, 1972). To further

understand how behaviour contributes to the maintenance of hydration state, we tested for a

functional trade-off among EWL, VPD selection, activity level, and temperature. We predicted

that highly active salamanders that also selected high VPD would be the ones with the highest

EWL; these effects should be greater at 22°C than at 17°C (Heatwole and Newby, 1972; Spight,

1968). Finally, we also tested whether rehydration rates (ReR) increased as a function of EWL

and test temperature (Claussen, 1969; Spight, 1967). With this integrative approach, we aim to

provide a broader understanding of the role of behaviour to the maintenance of hydration state in

amphibians.

Material and methods

Salamander collection and husbandry

In May 2022, we collected 44 post-breeding, adult *A. maculatum* (22 females + 22 males) with a

drift fence installed around Bat Lake, Algonquin Provincial Park, ON, Canada (45.5857°N,

78.5185°W). We recorded the sex of each salamander based on cloacal morphology (Petranka,

1998) and identified each individual based on their unique dorsal spot pattern. To transfer the

salamanders to Brock University, we allocated them into plastic containers with ventilated fitted

lids (34 cm x 19.6 cm x 12 cm). These containers had *Sphagnum* moss, pine needles, and water,

and were placed in a transport box at ~ 4°C to avoid overheating and dehydration during

transportation.

In the lab, we housed salamanders in pairs within ventilated tanks containing coconut

husk fibre, *Sphagnum* moss, PVC pipe refuges, and a water dish. These tanks were kept in a

facility with controlled temperature, RH, and photoperiod. We adjusted temperature and

photoperiod seasonally to simulate native habitat conditions while keeping RH constantly at 70%

(Giacometti and Tattersall, 2024). We fed salamanders twice a week with mealworms,

nightcrawlers, and isopods dusted in multivitamin and calcium powder. We weighed

salamanders weekly to the nearest 0.01 g using an analytical scale (Mettler Toledo, model

PB602-S) to monitor changes in body mass (M_b) as a proxy of well-being; all animals

maintained M_b throughout their period in the lab. The current study was conducted between late

June and August 2024 with salamanders that had been acclimatised to summer conditions (14°C

and a 14:12 light:dark cycle) for at least four weeks.

Laboratory humidity gradient

To assess behavioural hydroregulation, we used an annular humidity gradient (total length = 120

cm, outer radius = 45 cm, inner radius = 30 cm) built by Brock University Technical Services

(Figure S1). The thermally conductive floor of the gradient had copper pipes connected to a

water bath (Haake™, model DC10) kept at either 17°C or 22°C. These temperatures represent,

respectively, the median and maximum upper selected temperatures (T_{sel}) by *A. maculatum* in the

summer (Giacometti and Tattersall, 2024). The gradient chamber was divided into an inner and

outer lane by a 10.5 cm tall partition attached to the gradient lid, allowing the simultaneous study

of two individuals per experiment (visually isolated). To create a humidity gradient, we delivered

air of different moisture content to compartments (dry, intermediate, and humid) in the gradient

chamber. We created these compartments by attaching sponges (4 cm x 3cm x 9 cm) to the

gradient wall every 30 cm. To ensure the sponges remained in place and allowed for salamander

navigation during the experiments, we sewed the hook side of Velcro tape onto the sponges and

used aquarium safe silicon sealant (Adhesive Guru, product AG310) to attach the loop side of

Velcro tape to the gradient wall. The gradient lid had two venting holes every 90° relative to the

horizontal axis, such that the venting holes were positioned above the centre of the dry

compartment at 0°, the intermediate compartments at 90° and 270°, and the humid compartment

at 180° (Figure S1).

To generate dry air at 0°, we used tubing to direct compressed air through a column of

desiccant and then into the gradient at 1900 mL/min (split to direct evenly to both lanes). We

generated humid air using a water vapour bubbling system connected to the venting holes at

180°. This system consisted of an 1800 mL flask filled with dechlorinated water and kept 3°C

warmer than the prevailing experimental temperature. We aerated the water in the flask with an

aquarium air stone (15 cm diameter) attached to tubing that received compressed air at 2100

mL/min. The pressure created inside the flask directed water vapour from the flask to the venting

holes in the gradient lid (split evenly to both lanes). To generate intermediate moisture levels at

90° and 270°, we used a manifold to create a subsample line that combined air from the dry and

humid lines and directed mixed air into the gradient chamber at 1000 mL/min. Since dry air is

heavier than humid air (Haynes, 2016), we drew air at 250 mL/min from the dry line, and 750

mL/min from the humid line. We verified flow rates with flow meters (Aalborg Instruments &

Controls Inc., model PMR1-010972) at the beginning and end of each experiment to ensure

proper functioning of the humidity gradient.

With this setup, we created a stable, near-linear humidity gradient at 17°C ($R^2 = 0.85$)

and 22°C ($R^2 = 0.84$) (Figure S2). We determined this with hygro-thermometers (Inkbird, model

IBS-TH1; accuracy ± 3%) placed every 22.5° and set to record temperature and RH every 60 s

for 24 h in the absence of any animals in the gradient. Since RH is a temperature-dependent

measurement and therefore not readily comparable between 17°C and 22°C, we used VPD as a

measure of gradient humidity between temperatures (Table S1). Following Haynes (Haynes,

2016), where T is temperature (°C), we calculated:

$$WVP_{sat} = 6.11 \times 10^{\left(\frac{7.5 \times T}{237.3 + T}\right)} \qquad (1)$$

$$159 \qquad WVP_{act} = WVP_{sat} \times \left(\frac{RH}{100}\right) \qquad (2)$$

$$160 \qquad VPD = WVP_{sat} - WVP_{act} \qquad (3)$$

We conducted our experiments in total darkness inside a temperature-controlled room

(maintained at either 17 or 22°C). We placed the humidity gradient on top of 15 cm of insulation

foam board and padding so that inadvertent building vibrations would not disturb the

salamanders during the experiments. We positioned an infrared illuminator (wavelength = 850

nm; TVPSii, model TP-IRBP15) and a high-resolution infrared webcam (Agama, model V-

1325R) 180 cm above the centre of the gradient to record the salamanders via a time-lapse image

acquisition software (HandyAVI®) set to capture an image every 30 s.

Experimental design

We allowed the salamanders a total of 12 h inside the humidity gradient, from ~21:00 h to ~9:00

h. Each salamander was randomly tested twice (once under each experimental temperature), with

a minimum interval of 10 days between experiments. We considered the initial 3 h as the

habituation period and the subsequent 9 h as the experimental period. All individuals were fasted

for at least a week prior to experimentation to ensure a post-absorptive state (Secor and Boehm,

2006). We always allowed a minimum interval of 10 h between experiments after disinfecting

the humidity gradient with 70% ethanol at the end of an experiment.

We always handled the salamanders using nitrile gloves. Before introducing salamanders

into the gradient, we placed them individually into a container filled with 30 mL of dechlorinated

water for 15 min so that water could be absorbed through the skin and animals could be

introduced in a high hydration state. Then, we blotted the salamanders with paper towel, weighed

them, and placed them into the gradient. We always introduced the salamanders into the gradient

through the intermediate compartment, determining at random whether animals would be

initially facing the dry or humid end, and whether they would be tested in the inner or outer

gradient lanes. To minimise handling stress over hydroregulatory behaviours, we did not handle

or manipulate the salamanders during the experiments. After finishing an experiment, we

removed the salamanders from the gradient, blotted them with paper towel, and weighed them.

We considered the difference in M_b before and after an experiment divided by the length of our

experiment as a proxy for evaporative water loss (EWL; mL/h) (Ortega-Chinchilla et al., 2023).

We determined ReR (mL/h) by placing the salamanders individually in a container with 30 mL

of dechlorinated water for 30 min. Then, we removed the salamanders from the containers,

blotted them with paper towel, and reweighed them to determine ReR based on M_b gained per

half hour (Sherman and Stadlen, 1986). After this, we placed the salamanders back into their

housing tanks.

Data processing

We obtained a total of 1440 images for each individual over the course of each 12 h long

experiment. We imported image sequences into Fiji (Schindelin et al., 2012), and for each

experiment we recorded the identity (ID) and sex of the individual, as well as the test

temperature (17°C or 22°C). We used the manual tracking plug-in in Fiji to track the position of

the mid-body of the salamander within the gradient for each image in the sequence. As an output,

we obtained a set of Cartesian (*x*, *y*) coordinates that had *x*, *y* = 0,0 as the top left of the image.

We then used calculations derived from Giacometti et al. (Giacometti et al., 2021) to convert

salamander body position into selected gradient humidity, which we determined by calculating

median selected VPD and median selected RH.

To quantify activity in the gradient, we measured the distance moved (d_t) by each

individual every 30 s based on the distance defined by the arc of a circle:

$$d_t = \frac{0.5 \times (r_t + r_{t-1}) \times \pi \times 2 \times |\theta_t - \theta_{t-1}|}{360} \qquad (4)$$

where r is the mean radius, and θ are polar coordinates between adjacent time points t and t-1 for

all n time points. We then obtained total distance moved (D_t) by taking the sum of d_t:

$$D_t = \sum_1^{n-1} d_t \qquad (5)$$

where n is the total number of images ($n = 1440$).

Finally, we assessed the temperature dependency of ReR, using the Van't Hoff equation

to calculate the temperature coefficient (Q_{10}) for ReR between 17°C (T1) and 22°C (T2):

$$Q_{10} = (ReR_2/ReR_1)^{\frac{10}{T_2 - T_1}} \qquad (6)$$

Data analyses

We performed all analyses using R (version 4.4.1) in RStudio (version 2024-06-14) (R Core

Team, 2024) considering a significance level of 0.05. We built linear mixed-effects models using

the "lmer" function from the lme4 package (Bates et al., 2015), and we included ID as a random

term in all models to account for repeated measures. We tested our hypotheses using data from

the experimental period. We fit two individual models to test how humidity selection differed

between temperatures. To this end, we considered either median VPD or RH as the response

variables, experimental temperature as the predictor, and log-transformed M_b as a covariate. To

assess how EWL differed between temperatures, we fit a model that had log-transformed EWL

as the response variable, experimental temperature, median VPD, and D_t as the predictor

variables, and log-transformed M_b as a covariate. We also evaluated how ReR differed between

temperatures with a model that had log-transformed ReR as the response variable, experimental

temperature and log-transformed EWL as the predictors, and log-transformed M_b as a covariate.

For each model, we assessed residual autocorrelation with the "checkresiduals" function from

the forecast package (Hyndman et al., 2020), and the "acf" and "qqnorm" functions from the

stats package (R Core Team, 2024). We evaluated model fit with the "check_model" function

from the performance package (Lüdecke et al., 2021), and computed marginal fixed effects

through the "ggeffect" function from the ggeffects package (Lüdecke, 2018). We created figures

using the ggplot2 (Wickham, 2016), Thermimage (Tattersall, 2021), and ggpubr (Kassambara

and Kassambara, 2020) packages.

Results

During the experimental period, salamanders defended a constant VPD regardless of test

temperature (Table 1) (Figure 1a). This effect was mediated by M_b, with smaller individuals

selecting lower VPD than larger individuals. Importantly, salamanders avoided high VPD at both

test temperatures (Figure S3). Salamanders selected different RH between temperatures (Table 1)

(Figure 1b), with smaller individuals selecting higher RH than larger individuals. Regardless of

temperature and activity levels, salamanders with high EWL were those that selected higher

VPD and were larger (Table S2) (Figure 2). Test temperature was the only factor impacting ReR

($Q_{10}= 2.54 \pm 2.26$), with salamanders rehydrating faster at 22°C than at 17°C (Table S3) (Figure

3). Salamanders moved similar total distances within the gradient regardless of temperature

(Table 2).

Discussion

We used an integrative approach to evaluate the effect of temperature on behavioural

hydroregulation in *A*. *maculatum*. Our findings suggest that salamanders behaviourally defended

a constant VPD by altering RH selection between temperatures. By targeting higher RH at 22°C

than at 17°C, salamanders potentially compensate for increased evaporative drive at warmer

temperatures. We also found evidence for a trade-off among EWL, body size, and humidity

selection. Specifically, larger individuals that selected high VPDs experienced greater EWL than

smaller individuals that selected low VPDs. Temperature was the only factor explaining ReR,

with salamanders rehydrating faster at 22°C than at 17°C. Together, our findings underscore the

crucial role of behavioural strategies in the maintenance of hydration state in amphibians.

We found that salamanders defended a constant selected VPD between temperatures.

Since VPD represents the driving force for EWL (Riddell et al., 2024), our data suggest that *A*.

maculatum can detect their rate of water loss and alter their behaviour patterns to control

hydration state. To maintain homeostasis of VPD, salamanders selected higher RH at 22°C than

at 17°C. The maintenance of a constant VPD across temperatures illustrates an underappreciated

facet of amphibian behaviour: the ability to sense, process, and respond to moisture content in

the air in a physiologically relevant manner (Shelford, 1913), and thereby maintain an effectively

constant vapour pressure gradient and EWL. Under the "wetter is better" paradigm, amphibians

are depicted as poor thermoregulators that sustain low T_bs and always select the wettest

microhabitats available to minimise dehydration (reviewed in Brattstrom, 1979). Our data

challenge this view by demonstrating that *A. maculatum* not only showed evidence of humidity

preference, but also employed behavioural adjustments to mitigate increased evaporative demand

at warmer temperatures.

Our humidity preference values for *A. maculatum* were similar to those reported in the

congener *A. opacum* (VPD = 0.22 kPa, or RH = 74% at 13°C) (Marangio and Anderson, 1977).

Humidity preference in *Ambystoma* species, however, appears to be lower than those reported for

other amphibians (Bellis, 1962; Galindo et al., 2018; Shoemaker et al., 1992). The fossorial habit

of most *Ambystoma* species coupled with their relatively large body size could explain this

pattern. Truly, fossorial amphibians typically bear behavioural and morphophysiological

adaptations that allow for an increased independence from standing water in the environment

compared to non-fossorial species (Shoemaker et al., 1992). For example, *Ceratophrys*

stolzmanni uses soil humidity as a cue to adjust burrowing depth and maintain water balance

(Székely et al., 2018), and *A. maculatum* can tolerate up to 41% of M_b loss through evaporation

before losing function (Hall, 1922). Furthermore, our analyses showed that larger salamanders

selected overall higher VPD than smaller ones at both test temperatures, suggesting that

individuals with greater water content may be able to withstand greater VPDs (MacCracken and

Stebbings, 2012). Small-bodied salamanders often coil to decrease the surface area available for

EWL (e.g., Heatwole, 1960); however, we never observed *A. maculatum* assuming water-

conserving postures despite the relatively wide range of body size in our study animals (M_b

range = 4.66–25.80 g). Besides possessing a heightened desiccation tolerance (Giacometti and

Tattersall, 2025) and being able to maintain hydration state behaviourally, *A. maculatum* may

also benefit from the presence of cutaneous grooves in the venter and flanks which aid in

transferring water to different parts of the body by capillarity (Lopez and Brodie, 1977; Toledo

and Jared, 1993).

Our study also highlights how body size may influence behavioural hydroregulation in *A.*

maculatum. Larger salamanders selected higher VPD and exhibited greater EWL than smaller

salamanders, indicating a size-dependent trade-off between evaporation and humidity selection.

Contrary to our prediction, however, the relationship between EWL and VPD was not mediated

by temperature nor activity level. The allometric scaling between body size and EWL suggests

that, relative to their size, smaller individuals should evaporate faster than larger individuals

because small individuals have a high surface area-to-volume ratio (Jørgensen, 1997). Therefore,

smaller salamanders are predicted to experience greater hydric costs than larger salamanders, and

behavioural adjustments should be crucial to mitigate EWL (Riddell et al., 2024). Our results

supported this notion, as evidenced by the fact that smaller *A. maculatum* prioritised the selection

of low VPDs that culminated in overall low EWL. By contrast, larger *A. maculatum* selected

relatively high VPDs that resulted in increased EWL. In nature, size-specific habitat use is an

important behavioural strategy to minimise dehydration in aboveground-dwelling salamanders

(Heatwole, 1962; Spotila, 1972). However, the extent to which size-dependent habitat selection

contributes to hydroregulation in fossorial amphibians still warrants further research.

Consistent with previous work (Claussen, 1969; Spight, 1967), we found that *A.*

maculatum rehydrated faster at 22°C than at 17°C. This thermal dependency of rehydration

highlights the dual role of temperature as both a driver of EWL (Giacometti and Tattersall, 2025)

and a modulator of ReR. Notably, however, we did not find a relationship between EWL and

ReR in the current study. A possible reason for this lies in the fact that we allowed *A. maculatum*

to hydroregulate at will prior to ReR assessments. Evidence from *Rhinella marina* indicates that

toads may gain water at rates as fast as 30 mL/h when dehydrated to 30% of their initial M_b

(Kosmala et al., 2020). However, water uptake is minimal when *R. marina* have access to water

(Juarez et al., 2024). Our results back up this pattern, as rehydration in *A. maculatum* mostly

consisted of salamanders regaining their initial M_b. Indeed, *A. maculatum* lost a similar

proportion of M_b between temperatures (6.69 ± 3.57% at 17°C and 6.01 ± 3.81% at 22°C), and

thus our reported values might not represent their maximum ReRs. Despite the integumentary

and vascular adaptations that favour water retention in salamanders, ReRs in this group are still

lower than those of anurans (Marangio and Anderson, 1977; Spight, 1967; Toledo and Jared,

1993). Nonetheless, our calculated Q_{10} value for ReR suggests that at least some degree of

temperature-dependent physiological regulation of water uptake exists in *A. maculatum*. The

mechanisms behind such regulation are still unclear, but may involve hormonal control of skin

permeability to enhance water uptake (Alvarado and Johnson, 1965), along with an increase in

blood circulation at warmer temperatures (Jørgensen, 1997).

Conclusions

The ability to maintain hydration state under varying thermal conditions is crucial to amphibians

(Ortega-Chinchilla et al., 2023). By integrating behavioural and physiological perspectives, our

study demonstrates the critical role of temperature-compensating behaviours to the maintenance

of hydration state in *A. maculatum*. Our finding that salamanders maintained a constant VPD

between temperatures challenges the traditional use of RH measurements to assess

hydroregulation and habitat suitability in amphibians (Galindo et al., 2018). Thus, to properly

understand amphibian hydroregulation, future studies should incorporate information on how

behaviour is used to manage water loss, rather than rely on simple physical measurements.

Although our study provides insight into the behaviours involved in hydroregulation, we still

have limited information about hydrosensation compared to thermal sensation. In this context,

clarifying how amphibians use water-sensing units in the skin to detect water content in the air is

invaluable (Spotila, 1972). Likewise, the observation that salamanders regulate a constant

driving force of evaporation under changing thermal regimes suggests a mechanism for detecting

rates of EWL. Local cooling of the skin arising from evaporation (Andrade, 2016; Sinsch, 1984)

is a plausible mechanism; if moist-skinned ectotherms allow for a specific, fixed decline in skin

relative to core temperature, they can effectively detect the drive for evaporative water loss.

Ultimately, our study underscores how behavioural adjustments are central to creating conditions

conducive to the maintenance of a homeostatic hydration state in amphibians.

Acknowledgements: We thank Ontario Parks, the Ministry of Northern Development, Mines,

Natural Resources, and Forestry, the Algonquin Wildlife Research Station, Kevin Kemmish,

Njal Rollinson, and Patrick Moldowan for facilitating access to Bat Lake and our study animals.

We are grateful to Shawn Bukovac, Kristin Bray, Sarah Kehoe, Tom Eles, and Natasha Hearn

for assistance with animal care. We thank Stephen Renda, Mitch Sillaste and Art Reimer from

Brock University's Machine Shop for building the apparatus used in our study.

Funding sources: Funding for this study was provided by a Doris White Memorial Graduate

Scholarship and a Gervan Fearon Graduate Scholarship to Danilo Giacometti and a Natural

Sciences and Engineering Research Council of Canada Discovery Grant to Glenn Tattersall

(RGPIN-2020-05089).

Author contributions: Danilo Giacometti and Glenn Tattersall conceived the ideas and

designed methodology; Danilo Giacometti collected the data; Danilo Giacometti analysed the

data with input from Glenn Tattersall; Danilo Giacometti led the writing of the manuscript. Both

authors contributed critically to the drafts and gave final approval for publication.

Conflict of interest: None declared.

Data availability statement: Our data and code can be accessed from the Brock University

Dataverse: https://doi.org/10.5683/SP3/IKHBTP.

References

Alvarado, R. H. and Johnson, S. R. (1965). The effects of arginine vasotocin and oxytocin on sodium and water balance in *Ambystoma*. *Comparative Biochemistry and Physiology* **16**, 531–546.

Anderson, D. B. (1936). Relative humidity or vapor pressure deficit. *Ecology* **17**, 277–282.

Andrade, D. V. (2016). Temperature effects on the metabolism of amphibians and reptiles. *Amphibian and Reptile Adaptations to the Environment; Interplay between Physiology and Behavior; de Andrade, DV, Bevier, CR, de Carvalho, JE, Eds* 129–254.

Bates, D., Mächler, M., Bolker, B. and Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* **67**, 1–48.

Bellis, E. D. (1962). The influence of humidity on wood frog activity. *American Midland Naturalist* 139–148.

Brattstrom, B. H. (1979). Amphibian temperature regulation studies in the field and laboratory. *American Zoologist* **19**, 345–356.

Broly, P., Devigne, L., Deneubourg, J. and Devigne, C. (2014). Effects of group size on
aggregation against desiccation in woodlice (Isopoda: Oniscidea). *Physiological*
Entomology **39**, 165–171.

Bulova, S. J. (2002). How temperature, humidity, and burrow selection affect evaporative water
loss in desert tortoises. *Journal of Thermal Biology* **27**, 175–189.

Bundy, D. and Tracy, C. R. (1977). Behavioral response of American toads (*Bufo americanus*)
to stressful thermal and hydric environments. *Herpetologica* **5**, 455–458.

Chapman, T. L. and Bidwell, J. R. (2023). Behavioral preference for microclimate conditions
across elevation in Plethodon montanus. *Journal of Experimental Zoology Part A:*
Ecological and Integrative Physiology **339**, 967–977.

Claussen, D. L. (1969). Studies on water loss and rehydration in anurans. *Physiological Zoology*
42, 1–14.

Dezetter, M., Le Galliard, J.-F. and Lourdais, O. (2023). Behavioural hydroregulation
protects against acute effects of drought in a dry-skinned ectotherm. *Oecologia* **201**, 355–
367.

Galindo, C. A., Cruz, E. X. and Bernal, M. H. (2018). Evaluation of the combined temperature
and relative humidity preferences of the Colombian terrestrial salamander *Bolitoglossa*
ramosi (Amphibia: Plethodontidae). *Canadian Journal of Zoology* **96**, 1230–1235.

Giacometti, D. and Tattersall, G. J. (2024). Seasonal variation of behavioural thermoregulation
in a fossorial salamander (*Ambystoma maculatum*). *Royal Society Open Science* **11**,
240537.

Giacometti, D. and Tattersall, G. J. (2025). Seasonal plasticity in the thermal sensitivity of
metabolism but not water loss in a fossorial ectotherm. *ResearchSquare*.

Giacometti, D., Yagi, K. T., Abney, C. R., Jung, M. P. and Tattersall, G. J. (2021). Staying
warm is not always the norm: behavioural differences in thermoregulation of two snake
species. *Canadian Journal of Zoology* **99**, 974–983.

Greenberg, D. A. and Palen, W. J. (2021). Hydrothermal physiology and climate vulnerability
in amphibians. *Proceedings of the Royal Society B.* **288**, 20202273.

Hall, F. G. (1922). The vital limit of exsiccation of certain animals. *The Biological Bulletin* **42**,
31–51.

Haynes, W. M. ed. (2016). *CRC Handbook of Chemistry and Physics*. 97th ed. Boca Raton:
CRC Press.

Heatwole, H. (1960). Burrowing ability and behavioral responses to desiccation of the
salamander, Plethodon cinereus. *Ecology* **41**, 661–668.

Heatwole, H. (1962). Environmental factors influencing local distribution and activity of the
salamander, *Plethodon cinereus*. *Ecology* **43**, 460–472.

Heatwole, H. and Newby, R. C. (1972). Interaction of internal rhythm and loss of body water in
influencing activity levels of amphibians. *Herpetologica* **28**, 156–162.

Hillyard, S. D. and Willumsen, N. J. (2011). Chemosensory function of amphibian skin:
integrating epithelial transport, capillary blood flow and behaviour. *Acta Physiologica*
202, 533–548.

Hillyard, S. D., Hoff, K. von S. and Propper, C. (1998). The water absorption response: a
behavioral assay for physiological processes in terrestrial amphibians. *Physiological*
Zoology **71**, 127–138.

Hyndman, R. J., Athanasopoulos, G., Bergmeir, C., Caceres, G., Chhay, L., O'Hara-Wild,
M., Petropoulos, F., Razbash, S. and Wang, E. (2020). forecast: Forecasting functions
for time series and linear models. R package version 8.23.0.

Jørgensen, C. B. (1997). 200 years of amphibian water economy: from Robert Townson to the
present. *Biological Reviews* **72**, 153–237.

Juarez, B. H., Quintanilla-Salinas, I., Lacey, M. P. and O'Connell, L. A. (2024). Water
availability and temperature as modifiers of evaporative water loss in tropical frogs.
Integrative and Comparative Biology **64**, 354–365.

Kassambara, A. and Kassambara, M. A. (2020). Package 'ggpubr'. *R package version 0.6.0.*

Kosmala, G. K., Brown, G. P., Shine, R. and Christian, K. (2020). Skin resistance to water
gain and loss has changed in cane toads (*Rhinella marina*) during their Australian
invasion. *Ecology and Evolution* **10**, 13071–13079.

Laurin, M. (2010). *How vertebrates left the water*. University of California Press.

Lertzman-Lepofsky, G. F., Kissel, A. M., Sinervo, B. and Palen, W. J. (2020). Water loss and
temperature interact to compound amphibian vulnerability to climate change. *Global*
Change Biology **26**, 4868–4879.

Lillywhite, H. B. (2006). Water relations of tetrapod integument. *Journal of Experimental*
Biology **209**, 202–226.

Lopez, C. H. and Brodie, E. D. (1977). The function of costal grooves in salamanders
(Amphibia, Urodela). *Journal of Herpetology* **11**, 372–374.

Lüdecke, D. (2018). ggeffects: Tidy Data Frames of Marginal Effects from Regression Models.
Journal of Open Source Software **3**, 772.

Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P. and Makowski, D. (2021).
performance: An R package for assessment, comparison and testing of statistical models.
Journal of Open Source Software **6**,.

MacCracken, J. G. and Stebbings, J. L. (2012). Test of a body condition index with
amphibians. *Journal of Herpetology* **46**, 346–350.

Marangio, M. S. and Anderson, J. D. (1977). Soil moisture preference and water relations of
the marbled salamander, *Ambystoma opacum* (Amphibia, Urodela, Ambystomatidae).
Journal of Herpetology **11**, 169–176.

Mitchell, A. and Bergmann, P. J. (2016). Thermal and moisture habitat preferences do not
maximize jumping performance in frogs. *Functional Ecology* **30**, 733–742.

Ortega, Z., Ganci, C. C. and Rivas, M. L. (2023). Thermoregulation and hydric balance in
amphibians. *Evolutionary Ecology of Amphibians* 103–119.

Ortega-Chinchilla, J. E., Cabanzo-Olarte, L. C., Vaz, R., Lamadrid-Feris, F., Bevier, C. R.
and Navas, C. A. (2023). Behavioral models of hydrothermal regulation in anurans: A
field study in the Atlantic Forest, Brazil. *Biotropica* **55**, 329–338.

Petranka, J. W. (1998). *Salamanders of the United States and Canada*. Smithsonian Institution
Press.

Pough, F. H., Taigen, T. L., Stewart, M. M. and Brussard, P. F. (1983). Behavioral
modification of evaporative water loss by a puerto rican frog. *Ecology* **64**, 244–252.

R Core Team (2024). R: A language and environment for statistical computing. *R Foundation*
for Statistical Computing. Vienna, Austria.

Riddell, E. A., Roback, E. Y., Wells, C. E., Zamudio, K. R. and Sears, M. W. (2019).
Thermal cues drive plasticity of desiccation resistance in montane salamanders with
implications for climate change. *Nature Communications* **10**, 4091.

Riddell, E. A., Burger, I. J., Muñoz, M. M., Weaver, S. J. and Womack, M. C. (2024).
Amphibians exhibit extremely high hydric costs of respiration. *Integrative and*
Comparative Biology **64**, 366–376.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. *Nature Methods* **9**, 676–682.

Schmidt-Nielsen, K. and Schmidt-Nielsen, B. (1952). Water metabolism of desert mammals.
Physiological Reviews **32**, 135–166.

Secor, S. M. and Boehm, M. (2006). Specific dynamic action of Ambystomatid salamanders
and the effects of meal size, meal type, and body temperature. *Physiological and*
Biochemical Zoology **79**, 720–735.

Shelford, V. E. (1913). The reactions of certain animals to gradients of evaporating power of air:
A study in experimental ecology. *The Biological Bulletin* **25**, 79–120.

Sherman, E. and Stadlen, S. G. (1986). The effect of dehydration on rehydration and metabolic
rate in a lunged and a lungless salamander. *Comparative Biochemistry and Physiology*
Part A: Physiology **85**, 483–487.

Shoemaker, V. H. and Bickler, P. E. (1979). Kidney and bladder function in a uricotelic
treefrog (*Phyllomedusa sauvagei*). *Journal of comparative physiology* **133**, 211–218.

Shoemaker, V. H., Hillman, S. S., Hillyard, S. D., Jackson, D. C., Mcclanahan, L. L.,
Withers, P. and Wygoda, M. (1992). Exchange of water, ions and respiratory gases in
terrestrial amphibians. In *Environmental Physiology of the Amphibians*, pp. 125–150. The
University of Chicago Press.

Sinsch, U. (1984). Thermal influences on the habitat preference and the diurnal activity in three
European Rana species. *Oecologia* **64**, 125–131.

Spight, T. M. (1967). The water economy of salamanders: water uptake after dehydration.
Comparative Biochemistry and Physiology **20**, 767–771.

Spight, T. M. (1968). The water economy of salamanders: evaporative water loss. *Physiological*
Zoology **41**, 195–203.

Spotila, J. R. (1972). Role of temperature and water in the ecology of lungless salamanders.
Ecological Monographs **42**, 95–125.

Székely, D., Cogălniceanu, D., Székely, P. and Denoël, M. (2018). Dryness affects burrowing
depth in a semi-fossorial amphibian. *Journal of Arid Environments* **155**, 79–81.

Takei, Y. (2015). From aquatic to terrestrial life: evolution of the mechanisms for water
acquisition. *Zoological science* **32**, 1–7.

Tattersall, G. (2007). Skin Breathing in Amphibians. In *Endothelial Biomedicine*, pp. 85–91.

Tattersall, G. J. (2021). Thermimage: Thermal image analysis.

Titon, B. and Gomes, F. R. (2017). Associations of water balance and thermal sensitivity of
toads with macroclimatic characteristics of geographical distribution. *Comparative*
Biochemistry and Physiology Part A: Molecular & Integrative Physiology **208**, 54–60.

Toledo, R. C. and Jared, C. (1993). Cutaneous adaptations to water balance in amphibians.
Comparative Biochemistry and Physiology Part A: Physiology **105**, 593–608.

Tracy, C. R., Christian, K. A., Betts, G. and Tracy, C. R. (2008). Body temperature and
resistance to evaporative water loss in tropical Australian frogs. *Comparative*
Biochemistry and Physiology Part A: Molecular & Integrative Physiology **150**, 102–108.

Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York.

Wolcott, T. G. and Wolcott, D. L. (2001). Role of behavior in meeting osmotic challenges.
American Zoologist **41**, 795–806.

Tables

Table 1. Behavioural, body size, and physiological parameters of *Ambystoma maculatum* tested

in a humidity gradient kept at either 17°C or 22°C. Values are shown as mean ± SD.

	17°C (N=44)	22°C (N=44)
Selected VPD (kPa)		
Mean ± SD	0.56 ± 0.16	0.57 ± 0.22
(min, max)	(0.27, 0.95)	(0.12, 1.06)
Selected RH (%)		
Mean ± SD	71.20 ± 8.31	78.48 ± 8.34
(min, max)	(51.01, 85.88)	(60.09, 95.30)
M_b (g)		
Mean ± SD	12.00 ± 3.91	12.36 ± 4.62
(min, max)	(5.02, 23.70)	(4.66, 25.80)
EWL (mL/h)		
Mean ± SD	0.07 ± 0.05	0.06 ± 0.06
(min, max)	(0.01, 0.19)	(0.01, 0.30)
ReR (mL/h)		
Mean ± SD	0.18 ± 0.13	0.31 ± 0.25
(min, max)	(0.02, 0.56)	(0.02, 1.14)
Total distance moved (m)		
Mean ± SD	40.00 ± 20.80	40.31 ± 26.94
(min, max)	(0.11, 90.62)	(0.34, 105.01)

524   VPD = vapour pressure deficit, RH = relative humidity, M_b = body mass, EWL = rates of

525   evaporative water loss, ReR = rehydration rates.

Table 2. Parameter estimates (β), 95% confidence intervals (95% CI), and *p*-values for the models testing how humidity selection differed between temperatures in *Ambystoma maculatum* (N = 44). We considered either vapour pressure deficit (VPD) or relative humidity (RH) as response variables, temperature and log-transformed body mass ($\log M_b$) as predictors, and ID as a random term. Significant parameters are shown in bold.

	VPD ~ temperature + $\log M_b$ + (1|ID)		
Predictors	*Estimates*	*95% CI*	*p*
Intercept	0.40	0.26 – 0.53	**<0.001**
temperature (22ºC)	0.01	-0.06 – 0.07	0.883
$\log M_b$	0.01	0.00 – 0.02	**0.011**
Random Effects			
σ^2	0.03		
$\tau_{00\ ID}$	0.01		
ICC	0.22		
Marginal R^2 / Conditional R^2	0.10 / 0.30		
	RH ~ temperature + $\log M_b$ + (1|ID)		
Predictors	*Estimates*	*95% CI*	*p*
Intercept	87.31	73.65 – 100.98	**<0.001**
temperature (22ºC)	7.42	4.45 – 10.39	**<0.001**
$\log M_b$	-15.22	-27.96 – -2.49	**0.020**
Random Effects			
σ^2	48.92		
$\tau_{00\ ID}$	16.14		
ICC	0.25		
Marginal R^2 / Conditional R^2	0.22 / 0.42		

σ^2 = residual variance; $\tau_{00\ ID}$ = individual variance; ICC = intraclass correlation coefficient.

Figures

Figure 1. a. Salamanders defended a constant vapour pressure deficit (VPD) through

behavioural selection of different relative humidity (RH) levels (**b**) between 17°C and 22°C. In

both panels, large dots show mean marginal effects and bars indicate the corresponding 95%

confidence intervals. Small dots show the mean predicted value for each individual salamander

at a given temperature (17°C = blue, 22°C = red).

Figure 2. Relationship between log-transformed rates of evaporative water loss (logEWL) and

log-transformed body mass (logM$_b$) in *Ambystoma maculatum* tested in a humidity gradient at

either 17°C or 22°C. Larger individuals had higher EWL, and individuals that selected high

VPDs were also the ones with higher EWL at both temperatures. In both panels, each dot

represents an individual salamander. The solid lines and shaded areas indicate the predicted

relationship between logEWL and logM$_b$, and the 95% confidence interval, respectively. Vapour

pressure deficit (VPD) is colour-coded, with colder colours indicating wetter conditions and

warmer colours indicating dryer conditions.

550   **Figure 3.** Difference in log-transformed rehydration rates (logReR) between temperatures.

551   Salamanders rehydrated faster at 22°C than at 17°C. Large dots show mean marginal effects and

552   bars show the corresponding 95% confidence intervals. Small dots indicate the mean predicted

553   value for each individual salamander between temperatures (17°C in blue and 22°C in red).