Endogenous RALF peptide function is required for powdery mildew host colonization

Henriette Leicher[1], Sebastian Schade[1,#], Jan W. Huebbers[2], Kristina S. Munzert-Eberlein[3], Genc Haljiti[4],

David Biermann[5], Athanasios Makris[6], Xiaoxuan Zhu[7], Yashank Chauhan[1], Christina Ludwig[4], Marion

C. Müller[1], Toshinori Kinoshita[8,9], Timo Engelsdorf[3], Julien Gronnier[5], Martina K. Ried-Lasi[6], Aurélien

Boisson-Dernier[7], Ralph Hückelhoven[1] and Martin Stegmann[1,#]

[1]Chair of Phytopathology, TUM School of Life Sciences, Technical University of Munich (TUM), Freising,
Germany

[2]Unit of Plant Molecular Cell Biology, Institute for Biology I, RWTH Aachen University, Aachen, Germany

[3]Molecular Plant Physiology, Department of Biology, Philipps-Universität Marburg, Marburg, Germany

[4]Bavarian Center for Biomolecular Mass Spectrometry (BayBioMS), TUM School of Life Science, Technical
University of Munich (TUM), Freising, Germany

[5]Plant Cell Biology, TUM School of Life Sciences, Technical University of Munich (TUM), Freising, Germany

[6]Leibniz-Institute of Plant Biochemistry, Halle, Germany

[7]Université Côte d'Azur, INRAE, CNRS, Institut Sophia Agrobiotech, Sophia Antipolis, France

[8]Graduate School of Science, Nagoya University, Chikusa, Nagoya, Japan

[9]Institute of Transformative Bio-Molecules (WPI-ITbM), Nagoya University, Chikusa, Nagoya, Japan

[#]Institute of Botany, Molecular Botany, Ulm University, Ulm, Germany

*Corresponding author: Email: martin.stegmann@tum.de
20

Abstract

The receptor kinase FERONIA (FER) is a susceptibility factor for biotrophic powdery mildew fungal

pathogens in *Arabidopsis thaliana*, but the underlying molecular mechanisms remain largely unknown.

FER is required for the perception of endogenous RAPID ALKALINIZATION FACTOR (RALF) peptides

to control various aspects of plant growth, development and immunity. RALFs are perceived by FER-

LORELEI-LIKE GPI-ANCHORED PROTEIN (LLG) heterocomplexes to induce cellular responses and

bind to LEUCINE-RICH REPEAT EXTENSIN (LRX) proteins as structural components of the cell wall.

Combining genetics, cell biology and biochemistry, we found that FER`s endogenous RALF ligands are

necessary for full colonization success of the powdery mildew species *Erysiphe cruciferarum*. We reveal

that LLGs and LRXs are also powdery mildew susceptibility factors. We show that cell wall remodeling

and apoplastic pH homeostasis, hallmark features of RALF function, support powdery mildew

reproductive success. Moreover, we provide data that RALF-dependent powdery mildew pathogenesis

is partially independent of FER. Powdery mildew fungi likely do not produce RALF peptide mimics,

suggesting their reliance on endogenous RALFs for successful host colonization. We propose that

powdery mildew fungi require RALF-mediated modulation of apoplastic pH and pectin re-modelling for

successful host colonization, highlighting a new susceptibility mechanism by obligate biotrophic fungi.

Introduction

Powdery mildew is one of the most widespread fungal plant diseases affecting several economically relevant crop species (Glawe, 2008), with infection being characterized by continuous interaction between the biotrophic fungus and its host plant. Susceptibility genes (S-genes) can affect fungal infection success and are utilized by pathogens for their own benefit. In consequence, mutating these genes usually confers pathogen and genotype-independent and hence durable disease resistance (van Schie & Takken, 2014; Engelhardt *et al.*, 2018) . The first well-characterized powdery mildew susceptibility factor was the barley MILDEW LOCUS O (MLO) that confers barley susceptibility towards adapted *Blumeria hordei*. Mutants in orthologous *MLO* genes are powdery mildew resistant in phylogenetically diverse plant species, including *Arabidopsis thaliana* (hereafter Arabidopsis) and tomato (Kusch & Panstruga, 2017).

Next to susceptibility factors, the outcome of plant-pathogen interactions is shaped by their immune system. Pattern-triggered immunity (PTI) is the first line of induced resistance and activated by the perception of microbe-associated molecular patterns (MAMPs) by pattern recognition receptors (PRRs), which are often plasma membrane localized receptor-like kinases (RLKs). To subvert PTI, adapted pathogens evolved effectors that are mainly perceived by intracellular nucleotide-binding leucine-rich repeat receptors (NB-LRRs) activating pathogen race-specific effector triggered immunity (Ngou *et al.*, 2022).

In addition to their function as PRRs, RLKs are important sensors for endogenous signals regulating growth, abiotic stress and/or modulating immunity (Rzemieniewski & Stegmann, 2022; Zecua-Ramirez *et al.*, 2026). Adapted pathogens can hijack RLK pathways to support accommodation (Hok *et al.*, 2011; Hok *et al.*, 2014; Ried *et al.*, 2019). The conserved RLK FERONIA (FER) and related CATHARANTUS ROSEUS RECEPTOR-LIKE KINASE 1-LIKEs (CrRLK1Ls) perceive endogenous RAPID ALKALINIZATION FACTOR (RALF) peptides to regulate multiple plant physiological processes (Ge *et al.*, 2017; Gonneau *et al.*, 2018; Zhu *et al.*, 2021; Zhong *et al.*, 2022; Lan *et al.*, 2023). FER recognizes RALFs in a heteromeric complex with LORELEI-LIKE GPI ANCHORED PROTEIN 1 (LLG1) (Li *et al.*, 2015; Xiao *et al.*, 2019) and controls PTI as a RALF-regulated scaffold for PRR complexes and their nanoscale organization (Stegmann *et al.*, 2017; Gronnier *et al.*, 2022). In addition to FER-dependent signaling, recent reports show that RALFs have a structural role in organizing the patterning of demethylesterified pectins in the cell wall (Moussu *et al.*, 2023; Schoenaers *et al.*, 2024). Here, RALFs bind to cell wall bound LEUCINE-RICH REPEAT EXTENSIN (LRX) proteins but the biochemical interplay between LRX and FER-LLG complexes remains little understood (Mecchia *et al.*, 2017; Zhao *et al.*, 2018; Dünser *et al.*, 2019; Moussu *et al.*, 2020; Gronnier *et al.*, 2022; Schade *et al.*, 2025). Adapted pathogens can hijack RALF signaling for their own benefit. *Fusarium oxysporum* and root knot nematodes produce RALF mimics to enhance infection success on tomato and Arabidopsis (Masachis *et al.*, 2016; Thynne *et al.*, 2017; Zhang *et al.*, 2020). Furthermore, the Ler-0 *fer-1* mutant is more resistant to the powdery mildew species *Golovinomyces orontii*, suggesting that FER acts as an *S*-factor (Kessler *et al.*, 2010). Yet, how FER confers powdery mildew susceptibility remains largely unknown.

Here, we investigated the molecular mechanism of FER-mediated powdery mildew susceptibility, using the Arabidopsis-adapted powdery mildew fungus *Erysiphe cruciferarum* (*Ecr)*. We show that LLG1 and

LRX1-LRX5 are required for successful powdery mildew infection. By loss of function studies, we identify
RALF peptides as important susceptibility factors for powdery mildew. CRISPR Cas9-generated
ralf22/23/33 (CRISPR *ralf3x*) and *ralf1/18/22/23/31/33/34 (*CRISPR *ralf7x*) mutants show increasing
resistance to *Ecr* infection depending on the number of eliminated RALFs. Our bioinformatic analysis
indicates that powdery mildew fungi do not encode functional RALF peptide mimics, supporting that they
require host-endogenous RALFs for their own benefit (Masachis *et al.*, 2016; Thynne *et al.*, 2017). We
suggest that the enhanced resistance of *fer* and CRISPR *ralf7x* is not associated with defects in MLO-
mediated penetration, or deregulation of defense-related phytohormone signaling. Instead, powdery
mildew might require FER-RALF-mediated apoplastic pH regulation and cell wall remodeling for
successful colonization. Surprisingly, our data indicates that RALF-mediated powdery mildew
susceptibility is partially independent on FER and exerts its effect in a combination of signaling and
structural functions. We propose that powdery mildew fungi require RALFs to provide a favorable growth
environment, highlighting a new mechanism of host subversion by adapted plant pathogens.

Results

FER, LLGs and LRX are powdery mildew susceptibility factors

Ler-0 *fer-1* is more resistant to *Golovinomyces orontii* (Kessler *et al.*, 2010). We confirmed this
phenotype in Arabidopsis Col-0 using the *fer-4* mutant allele (Duan *et al.*, 2010) and *Ecr* (Micali *et al.*,
2008). The *fer-4* mutant showed slightly weaker visible symptoms (Fig. 1A), reduced conidiophore
production and less relative fungal DNA content compared to Col-0 5 dpi (Fig. 1B, C, Fig. S1A).
Expression of pFER::FER-GFP complemented the *fer-4* resistance phenotype (Fig. 1C).

Next, we investigated whether FER-related RALF receptor complex components are involved in *Ecr*
susceptibility. The *llg1-2* mutant showed reduced conidiophore production (Fig. 1D). A similar effect was
observed in the *llg1-3* allele, in which LLG1 is impaired in RALF binding but mutants do not show growth
phenotypes (Fig. 1D, Fig. S11) (Shen *et al.*, 2017; Xiao *et al.*, 2019). The resistance phenotype of the
llg1 mutants, however, was not clear in qPCR quantification (Fig. S1B), indicating potential redundancy
of LLG1 with related members of the LRE/LLG gene family (Xiao *et al.*, 2019; Noble *et al.*, 2022).
Surprisingly, *llg1-2* and *llg1-3* were previously described as more susceptible to *Golonivomyces*
cichoracearum (Shen *et al.*, 2017), raising the question of mildew-species-dependent differences in *llg1*
pathophenotypes, unlike *fer* (Fig. 1C, D) (Kessler *et al.*, 2010). RALFs also bind to cell wall localized
LRX proteins (Moussu *et al.*, 2020). A *lrx3 lrx4 lrx5* (*lrx3/4/5*) triple mutant phenocopies *fer* in immune
signaling, salt stress responses and plant morphology (Zhao *et al.*, 2018; Dünser *et al.*, 2019; Moussu
et al., 2020; Gronnier *et al.*, 2022). This *lrx3/4/5* mutant displayed reduced conidiophore production upon
Ecr infection (Fig. 1E). Yet, relative fungal DNA content was unaltered (Fig. S1C). Further mutating
LRX1 and *LRX2* in *lrx3/4/5* (*lrx1/2/3/4/5*, hereafter *lrx5x*) (Herger *et al.*, 2020) promoted the resistance
phenotype (Fig. 1E, Fig. S1C). Overexpression of LRR4-citrine (LRR4$^{\Delta E}$/LRX4 lacking the extensin
domain) has a dominant negative effect on plant growth, while PTI responses are not affected (Dünser
et al., 2019; Gronnier *et al.*, 2022). The p35S::LRR4-citrine lines phenocopied *lrx5x* and *fer-4* in showing
reduced *Ecr* conidiophore production (Fig. 1F).

Powdery mildew requires endogenous RALF peptides for successful colonization

We hypothesized that RALF signaling through FER-LLG1 and/or their binding to LRXs is required for *Ecr* colonization. Many plant pathogens secrete RALF peptide mimics to promote virulence (Masachis *et al.*, 2016; Thynne *et al.*, 2017; Zhang *et al.*, 2020). We searched available genomes of powdery mildew species for the presence of potential RALFs. However, consistent with previous reports (Thynne *et al.*, 2017) we could not detect evidence for RALFs among different *Erysiphales* species (Table S1, method section for details on our bioinformatic search pipeline). Thus, we hypothesized that endogenous RALFs are involved in the *Ecr*-Arabidopsis interaction. Arabidopsis encodes for 37 RALFs (Abarca *et al.*, 2021). We anticipated genetic redundancy and generated CRISPR mutants in which we targeted *RALF23*, a described FER-LLG1 ligand (Stegmann *et al.*, 2017; Xiao *et al.*, 2019; Tang, J *et al.*, 2022) and seven additional RALF-peptides with predicted leaf expression (*RALF1*, *RALF14*, *RALF18*, *RALF22*, *RALF23*, *RALF31*, *RALF33*, *RALF34*) (Fig. S2A). We could not obtain an octuple mutant but generated two higher order mutants, *ralf22/23/33* (CRISPR *ralf3x*) and *ralf1/18/22/23/31/33/34* (CRISPR *ralf7x*) (Fig. 2A, Fig. S2B, C). CRISPR *ralf3x* displayed a mild growth phenotype with slightly shortened petioles, while CRISPR *ralf7x* was strongly impaired in shoot growth (Fig. 2B, Fig. S2D, S11). Compared to *fer-4*, however, CRISPR *ralf7x* showed less pronounced shoot growth defects with a trend towards longer petioles (Fig. S2D) and an overall more wildtype-like morphology (Fig. 2B, Fig. S11). *Ecr* infection was reduced on both CRISPR *ralf3x* and CRISPR *ralf7x* (Fig. 2B-E). Notably, CRISPR *ralf7x* displayed slightly elevated resistance compared to *fer-4* (Fig. 2D). Expression of *p35S::RALF23* complemented CRISPR *ralf7x* growth and *Ecr* infection phenotypes (Fig. 2F, Fig. S2E, F, S11). Additionally, the previously reported *ralf1/22/23/33* quadruple mutant (Lan *et al.*, 2023) was more resistant to *Ecr* with slightly less condiophore production compared to *fer-4* (Fig. 2A, G). We analyzed expression of the seven *RALFs* mutated in CRISPR *ralf7x* in Col-0 upon *Ecr* infection by qPCR. *RALF18* and *RALF23* were significantly upregulated at 4 and 5 dpi (Fig. S3). Together with our genetic data, this underlines the role of endogenous RALF peptides in *Ecr* host colonization.

FER and RALF mutants show post-penetration powdery mildew resistance

We next sought out to unravel cellular mechanisms underlying RALF-mediated *Ecr* susceptibility. RALF-induced calcium influx in synergid cells and pollen tubes is facilitated by FER-LRE-mediated activation of the NTA/MLO7 calcium channel (Gao *et al.*, 2022). The MLO family encodes important powdery mildew susceptibility factors across families (Büschges *et al.*, 1997; van Schie & Takken, 2014; Acevedo-Garcia *et al.*, 2017; Nekrasov *et al.*, 2017) . In Arabidopsis, mutation of three MLOs is required to confer full penetration resistance (Consonni *et al.*, 2006).

MLO-YFP in barley and MLO2-GFP in Arabidopsis focally accumulate at powdery mildew penetration sites (Bhat *et al.*, 2005; Huebbers *et al.*, 2024). We confirmed MLO2 localization to the fungal penetration site with p35S::MLO2-mCherry lines (Fig. S4A, left panel). We tested whether FER-GFP accumulates at *Ecr* penetration sites using *fer-4 pFER::FER-GFP* (Duan *et al.*, 2010). Although FER-GFP signal outlined the plasma membrane, specific penetration site recruitment was not detected (Fig. S4A, right panel). The *fer-4* mutant showed strongly reduced production of conidiophores on infected tissue (Fig. 1C), but initial hyphal growth was largely unaffected (Fig. S4B). Also, epidermal cell penetration was

unaffected in *fer-4* and CRISPR *ralf7x*, unlike *mlo2/6/12* (Fig. S4C). Callose deposition was not
enhanced at the penetration site on *fer-4* or CRISPR *ralf7x* compared to Col-0 (Fig. S4 D, E). This
suggests that mutating RALF-related components primarily affects *Ecr* reproductive success and not
fungal entry or epiphytic growth. We also tested whether *Ecr* spores derived from colonies grown on *fer-*
4, CRISPR *ralf7x* and *llg1-2* mutants show compromised infection potential upon re-inoculation of a Col-
0 host. However, neither of the spores displayed altered pathogenicity compared to Col-0-derived
spores (Fig. S4F). This indicates that, although colonies grown on the mutant plants have a reduced
number of spores, their quality is not impaired.

The *mlo2/6/12* mutant displays aberrant callose deposition in trichomes with delocalized or absent
accumulation (Huebbers *et al.*, 2024). Likewise, *fer-4* mutants are impaired in trichome formation (Duan
et al., 2010). Consistently, we observed less trichomes on *fer-4* but not on CRISPR *ralf7x* (Fig. S4G).
Unlike *mlo2/6/12*, neither CRISPR *ralf7x* nor *fer-4* showed mis-regulated trichome callose deposition
(Fig. S4H) (Huebbers *et al.*, 2024). A fraction of trichomes on the CRISPR *ralf7x* mutant were enlarged
in the base region which was not observed on Col-0 and *fer-4* (Fig. S4H). Further, trichomes on *fer-4*
and CRISPR *ralf7x* had fewer and more branches, respectively, compared to Col-0 (Fig. S4I).
Collectively, this data suggests that FER-RALF-dependent *Ecr* susceptibility and trichome development
is most likely not directly linked to MLO function.

The powdery mildew resistance phenotype of FER pathway mutants is not associated with de-
regulated defense phytohormone signaling

Salicylic acid (SA) is an important defense hormone promoting resistance to biotrophic and
hemibiotrophic pathogens (Peng *et al.*, 2021). Ler-0 *fer-1* seedlings did not show elevated basal
PATHOGENESIS-RELATED 1 (*PR1*) expression, an SA marker gene (Kessler *et al.*, 2010). Yet, the
weak *fer-5* allele in Col-0 showed mildly constitutively elevated SA levels in seedlings (Duan *et al.*, 2010;
Engelsdorf *et al.*, 2018), prompting us to test whether *fer-4*`s *Ecr* resistance phenotype may be
associated with elevated SA levels. We quantified free and glycosylated SA by high performance liquid
chromatography (HPLC) upon *Ecr* infection. *Ecr* infection led to mildly enhanced SA levels (Fig. 3A).
However, neither *fer-4* nor CRISPR *ralf7x* showed higher SA levels before or after infection (Fig. 3A).
We also measured the major catabolic SA products 2,3-DHBA and 2,5-DHBA to investigate the flux
through the SA pathway (Zhang *et al.*, 2013; Zhang *et al.*, 2017) as well as camalexin levels associated
with anti-fungal resistance (Liu *et al.*, 2016; Mucha *et al.*, 2019). Neither of those compounds were
enhanced before or after infection in *fer-4* and CRISPR *ralf7x* (Fig. 3B, C, Fig. S5A). *PR1* expression
was also not elevated in any of the mutants in control conditions. *Ecr* infection resulted in enhanced
PR1 expression. The *lrx5x* mutant showed enhanced *Ecr*-induced *PR1* expression, which, however,
was not observed in the other mutants (Fig. 3D). When infected with the biotrophic oomycete pathogen
Hpa, fer-4 and CRISPR *ralf7x* were more susceptible than Col-0 or *fer*-4/pFER::FER-GFP (Fig. S5B).
This further supports the hypothesis that reduced *Ecr* sporulation observed on the tested mutants is not
caused by enhanced SA signaling.

Jasmonic acid (JA) is usually implicated in defense against insect herbivory and necrotrophic fungi
(Bürger & Chory, 2019), but JA treatment can also inhibit colonization by the powdery mildew fungus

Erysiphe cichoracearum (Zimmerli *et al.*, 2004). FER inhibits JA signaling by destabilizing the
transcription factor MYC2 (Guo *et al.*, 2018). We generated a CRISPR-Cas9 *myc2* mutant in *fer-4* that
abolished *MYC2* expression (Fig. S5C, D). Similar to previous reports, *fer-4* CRISPR *myc2* partially
restored the *fer-4* growth phenotype (Fig. S5E) (Guo *et al.*, 2018) . In *fer-4*, the CRISPR *myc2* mutation
showed a clear trend towards complementation of deregulated JA-responsive gene expression (Guo *et*
al., 2018), a phenotype that was also observed in CRISPR *ralf7x* and *lrx5x* (Fig. 3E, F, Fig. S5D).
Importantly, mutating *MYC2* did not restore *Ecr* susceptibility in *fer-4* (Fig. 3G, Fig. S5F). This suggests
that the resistance phenotype of *fer-4* is independent of MYC2-mediated JA signaling.

Apoplastic pH modulations affect powdery mildew infection success

To determine potential protein interaction partners of FER, we performed co-immunoprecipitation-mass
spectrometry (CoIP-MS) in mature leaves of *fer-4* pFER::FER-GFP lines. We identified 68 potential
interaction partners of FER-GFP with the majority being RLKs or transport-related proteins (Fig. S6A,
B). We identified four CrRLK1L members (HERK1, MDS1/LET2, THE1 and LET1). In response to *Ecr*
infection, we could only detect six differential FER-GFP interactors, which may be related to the early
experimental time point chosen (1 dpi) (Fig. S7A, B).

Additional interactors include the plasma membrane AUTOINHIBITED H+-ATPase 1 (AHA1) and AHA3
(Fig. S6A, B). Compared to the Lti6b-GFP control, MS intensity of AHA1 was 2,7-fold enriched (Fig. 4A),
confirming previously published AHA1-FER interaction in roots (Haruta *et al.*, 2014; Du *et al.*, 2016). We
also found AHA3 to be significantly enriched with peptides only detected in FER-GFP samples (Fig. 4A,
Fig. S6A, B). Perception of RALF by FER induces phosphorylation of an inhibitory site on AHA1,
resulting in apoplastic alkalinization (Pearce *et al.*, 2010; Haruta *et al.*, 2014). Since some fungal
pathogens modify the apoplastic pH for successful infection (Kesten *et al.*, 2019), we hypothesized that
FER-RALF-induced pH modulations may affect powdery mildew infection success. An *aha1* single
mutant did not display *Ecr* colonization differences (Weis *et al.*, 2013). While this might be explained by
genetic redundancy, *aha1 aha2* double mutants are seedling lethal, making genetic analysis difficult
(Haruta *et al.*, 2010).

Phosphorylation of the C-terminal penultimate threonine residue in AHA1/AHA2 (AHA1^{T947}/AHA2^{T948})
activates AHA proton pump activity (Falhof *et al.*, 2016). We tested whether *Ecr* infection affects
AHA1^{T947}/AHA2^{T948} phosphorylation by using a phosphosite-specific antibody. However, pT947/pT948
phosphorylation was not altered upon *Ecr* infection (Fig. S8A, B). RALF perception by FER promotes
phosphorylation of serine 899 (pS899) to inhibit AHA1/AHA2 proton pump activity (Haruta *et al.*, 2014;
Liu *et al.*, 2025; Sun *et al.*, 2025). Importantly, pS899 does not correlate with reduced pT947/pT948 and
RALF23/RALF33 treatment does not affect pT947/pT948 (Haruta *et al.*, 2014; Liu *et al.*, 2025).
Therefore, the results do not exclude a FER-dependent regulation of AHA activity correlated with altered
Ecr infection success. To investigate a potential role of AHA1 in *Ecr* susceptibility, we tested the *ost2-*
2D mutant producing a constitutive active AHA1 (Merlot *et al.*, 2007). The *ost2-2D* mutant displayed
enhanced *Ecr* resistance (Fig. 4B). Interestingly, the related Ler *ost2-1D* mutant displays reduced PTI
responses, similar to *fer*, thus uncoupling PTI efficiency from *Ecr* infection success (Stegmann *et al.*,
2017; Zhai *et al.*, 2025). The fungal toxin fusicoccin induces constitutive AHA activation and promotes

powdery mildew resistance in barley (Baunsgaard *et al.*, 1998; Zhou *et al.*, 2000). Indeed, fusicoccin

treatment could also induce *Ecr* resistance and promote the *fer-4* resistance phenotype (Fig. 4C, D).

We hypothesized that alkaline pH infiltration may enhance susceptibility and/or break the *fer-4*

resistance phenotype, whereas acidic buffers may promote resistance. We infiltrated MES buffer at pH

5.5 and pH 6.5 into the leaf apoplast prior to infection. Infiltration of a 15 mM $MgCl_2$ solution (pH 7)

served as an osmolarity control. Surprisingly, both infiltration of the more acidic and more alkaline buffer

reduced conidiophore production in comparison to the controls, both in Col-0 and *fer*-4 (Fig. 4E, F). We

next tested a more alkaline buffer and infiltrated pH 7.5 buffered HEPES into the apoplast before *Ecr*

infection. In line with pH 5.5 and pH 7.5 buffered MES, pH 7.5 buffered HEPES induced strong *Ecr*

resistance in both Col-0 and *fer-4* (Fig. S8C). When analyzing individual fungal colonies upon infection

on Col-0 MES pH 5.5 pre-buffered leaves we observed a morphological similarity to fungal colonies on

fer-4 (Fig. 4F). This observation suggests the importance of apoplastic pH modulation for successful

conidiophore production of *Ecr* at later time points of infection, which may be regulated through FER-

RALF-dependent signaling.

Among other physiological outputs, apoplastic pH is involved in growth regulation by controlling PECTIN

METHYL ESTERASE (PME) activity (Hocq *et al.*, 2016; Sénéchal *et al.*, 2017; Hocq *et al.*, 2023). PMEs

de-esterify cell wall pectins, which increases the negative charge of homogalacturonans resulting in

enhanced Ca^{2+} binding and pectin cross-linking for cell wall stiffening (Willats *et al.*, 2001). Importantly,

demethylated pectin interacts with RALFs and LRX-RALF complexes to structurally organize the cell

wall of root hairs and pollen tubes (Moussu *et al.*, 2023; Schoenaers *et al.*, 2024). Recently, *fer-4*

mutants were shown to have constitutively increased PME activity in seedlings, which we confirmed in

leaf tissue (Fig. 4G) (Biermann *et al.*, 2025). Similarly, PME activity was enhanced in *lrx5x* and CRISPR

ralf7x. *Ecr* infection, however, had no effect on PME activity in the harvested leaf samples (Fig. 4G). In

accordance with previous observations in the hypocotyl (Biermann *et al.*, 2025), leaf staining with the

fluorescently labelled chitosan oligosaccharide probe (COS-488), which binds to de-esterified pectin

oligomers (Mravec *et al.,* 2017*)*, displayed higher signals in *fer-4* compared to Col-0 (Fig. 4H, I).

Unfortunately, we could not analyze changes in the pectin methylesterification-status around *Ecr*

infection sites, because of strong COS-488 staining of fungal structures (Fig. S8D). To test whether

PME activity affects *Ecr* infection, we infiltrated leaves with epigallocatechin gallate (EGCG) that inhibits

PMEs (Lewis *et al.*, 2008). EGCG pre-treatment promoted *Ecr* resistance on Col-0 but had no effect in

fer-4 (Fig. 4J, K). To complement the inhibitor treatment, we tested the effect of PME-INHIBITOR 3

(PMEI3) overexpression on *Ecr* colonization. Overexpression of *PMEI3* inhibits RALF1 signaling in roots

(Rößling *et al.*, 2024). *PMEI3* overexpression resulted in reduced fungal colonization (Fig. 4L). We also

tested *pme3* and *pme17* mutants because of their role in *B. cinerea* infection and FER-dependent

epidermal cell lobing (Raiola *et al.*, 2011; Del Corpo *et al.*, 2020; Tang, W *et al.*, 2022). Knockout of

single *PMEs,* however, had no effect on conidiophore production (Fig. S8E). Taken together, full PME

activity, which affects FER-dependent RALF signaling, is required for powdery mildew susceptibility.

RALF-mediated powdery mildew susceptibility is partially independent of FER and FER-signaling

The reduced *Ecr* infection success upon pH buffering prompted us to resolve potential infection-induced apoplastic pH changes with cellular resolution with the ratio-metric pH sensor pUBQ10::SYP122-pHusion (Kesten *et al.*, 2019). We detected apoplastic pH increases close to *Ecr* penetration sites at 2 dpi. Surprisingly, infection-related pH increases were also observed in pUBQ10::SYP122-pHusion CRISPR *fer* (Fig. 5A, B, Fig. S9A).

To assess whether FER recruitment to RALF-LLG1 complexes is required for *Ecr* susceptibility, we infected *fer-4* lines expressing structure-guided FER mutants disrupting the LLG binding interface (Xiao *et al.*, 2019). Both FER^{N303Y} and FER^{G257A} restored powdery mildew susceptibility, indicating that RALF-induced recruitment of FER to LLG1 is not essential for *Ecr* infection (Fig. 5C). We thus asked the question whether the susceptibility function of RALFs is linked to cell wall structural functions. We introduced mutations into RALF23, either disrupting the predicted LRX-binding interface (RALF23$^{Y113A/Y114A}$), analogous to RALF4$^{Y83A/Y84A}$ (Moussu *et al.*, 2020) or the YISY motif required for LLG-FER-dependent signaling (RALF23$^{Y93A/Y96A}$) (Xiao *et al.*, 2019). We expressed both variants under control of the 35S promoter in CRISPR *ralf7x*. We tested two and three lines, respectively, with CRISPR *ralf7x* p35S::RALF23$^{Y93A/Y96A}$ displaying similar expression levels as the WT complementation lines. CRISPR *ralf7x* p35S::RALF23$^{Y113A/Y114A}$ lines showed weaker expression compared to the WT construct (Fig. S2F, G). RALF23$^{Y113A/Y114A}$ could not complement the CRISPR *ralf7x* growth and enhanced *Ecr* resistance phenotype (Fig. 5D, Fig. S11). RALF23$^{Y93A/Y96A}$ largely complemented the growth phenotype (Fig. S11) but only one out of three tested lines fully restored *Ecr* susceptibility (Fig. 5D). We also generated pRALF23 (pR23)::mCherry-RALF23 lines, with the mCherry tag located between the signal peptide and RALF23, carrying the same mutations. All RALF variants marked the outline of epidermal cells (Fig. S10, panel 1). Upon plasmolysis, a strong protoplastic signal for all mCherry-RALF23 variants was detected (Fig. S10, panel 2). However, mCherry-RALF23^{WT} and mCherry-RALF23$^{Y93A/Y96A}$ displayed strong remaining signal at the cell wall, while mCherry-RALF23$^{Y113A/Y114A}$ was fully depleted (Fig. S10, panel 1-4). This suggests loss of cell wall binding in RALF23$^{Y113A/Y114A}$ (Fig. S10). RALF23 cell wall localization is consistent with previous reports on RALF22 and RALF4 (Moussu *et al.*, 2023; Schoenaers *et al.*, 2024). RALF22 secretion is impaired in *lrx1/2* (Schoenaers *et al.,* 2024). By contrast, RALF23 secretion is neither affected by disrupting LRX nor LLG1 binding motives. RALF23 and RALF23$^{Y93A/Y96A}$, but not RALF23$^{Y113A/Y114A}$, could rescue enhanced PME activity of CRISPR *ralf7x* (Fig. 5 E), suggesting that RALF23 interaction with LRXs is essential to inhibit PME function. Together, this data implicates that both interaction of RALF23 with LLG and LRX complexes is required to support full *Ecr* colonization with the latter playing a predominant role for RALF-regulated PME activity and powdery mildew susceptibility.

We next tested whether application of synthetic RALF23 peptide can affect powdery mildew infection success. Surprisingly, infiltration of 1 µM RALF23 into single leaves 24 h prior to *Ecr* infection had a mild resistance-inducing effect in Col-0 (Fig. 5F), while the infiltration of lower concentrations (10 nM and 100 nM) had no significant effect (Fig. S9B). Remarkably, RALF23 infiltration increased conidiophore production in the *fer-4* mutant background, suggesting that RALF23 exerts FER-independent effects

(Fig. 5F). Interestingly, infiltration with RALF23 promoted alkalinization near the infection site in pUBQ10::SYP122-pHusion, but not in CRISPR *fer* (Fig. 5G). Surprisingly, water infiltration (control) increased *Ecr*-induced alkalinization in CRISPR *fer.* This raises the question whether FER is additionally involved in buffering *Ecr*-induced pH changes upon apoplast infiltration. Also, RALF23 infiltration promoted GFP fluorescence in pUBQ::SYP122-pHusion in a FER-dependent manner at early time points post treatment. Buffering abolished the recorded pH shifts (Fig. S9C), suggesting that it effectively compromises RALF23 pH changes. We continued testing the effect of RALF23-GFP overexpression on *Ecr* infection outcome. Similar to exogenous RALF23 treatment (Fig. 5F), RALF23-GFP overexpression displayed a trend towards enhanced *Ecr* resistance in Col-0 (Fig. 5H). In *fer-2,* RALF23-GFP overexpression promoted susceptibility (Fig. 5H) and displayed a clear trend to reduce the mutant`s enhanced PME activity (Fig. S9D). This observation is in line with previous findings that RALF23 treatment significantly reduces PME activity in *fer-4*, but not in Col-0 (Biermann *et al.*, 2025). *RALF23-GFP* showed overexpression in both Col-0 and *fer-2* (Fig. S9D).

Collectively, our data suggests that RALF-mediated powdery mildew susceptibility is at least partially FER-independent. Moreover, it indicates that likely appropriate spatio-temporal availability of RALFs affect *Ecr* reproductive success in a combination of partially FER-dependent effects on alkalinization and FER-independent responses, potentially related to RALF-regulated PME activity.

Discussion

This work provides first molecular insights into the mechanism of FER-mediated powdery mildew susceptibility. We revealed that FER`s RALF peptide ligands are critical for the completion of *Ecr*`s asexual life cycle upon host colonization.

RALFs are perceived by LLG-FER/CrRLK1L complexes to induce downstream signaling and bind to cell wall localized LRX proteins and demethylated pectin for cell wall organization. Our data suggests that powdery mildew sporulation requires functional apoplastic pH homeostasis and modifications of the cell wall pectin methyl-esterification status, both processes in which FER-RALF and/or LRX-RALF modules fulfil central regulatory functions. Earlier work revealed that barley powdery mildew infection is accompanied by pH shifts and the induction of apoplastic alkalinization at early time points post inoculation (2h) (Felle *et al.*, 2004). Importantly though, mildew-induced pH shifts were dynamic, with fluctuations between alkaline and more acidic conditions during the recorded time period (50 h) (Felle *et al.*, 2004). In light of our observations, disrupting apoplastic pH homeostasis by buffering, RALF23 peptide treatments or RALF23 overexpression, may circumvent dynamic endogenous regulatory circuits and inhibit *Ecr* host colonization. A challenge for the future will be to reveal spatio-temporal pH fluctuations and RALF secretion during powdery mildew infection.

In addition, the pectin methyl-esterification status contributes to powdery mildew pathogenesis. Both EGCG treatment and PMEI3 overexpression promoted *Ecr* resistance. Notably, RALF-signaling outputs and the methyl-esterification status of the cell wall are tightly interconnected. PMEI3 is strongly pH dependent with the highest activity at acidic conditions (Xu *et al.*, 2022) RALF-induced apoplastic alkalinization may inhibit PMEI3 which increases PME activity for cell wall remodeling. RALF22 and RALF4 bind to LRXs to affect pectin compaction in the cell wall (Moussu *et al.*, 2023; Schoenaers *et al.*,

2024). In return, PME activity modulates RALF1/RALF23 signaling and the formation of RALF1-pectin
molecular condensates required for FER-LLG1-dependent responses (Liu *et al.*, 2024; Rößling *et al.*,
2024; Biermann *et al.*, 2025). Consistently, EGCG treatment enhanced powdery mildew resistance in
Col-0 but had no additive effect in *fer-4*. Considering that *fer-4* is strongly compromised in RALF
perception, our data indicates that PME inhibition through EGCG treatment affects the same FER-
dependent pathway. We propose that EGCG treatment and PMEI3 overexpression block sustained *Ecr*
growth by reducing the availability of demethylated pectin for RALF signaling through FER. Fusicoccin
and buffer treatment could also induce resistance in *fer-4*, suggesting additional effects that do not solely
rely on pH-dependent PME regulation. Our work revealed that *fer*, CRISPR *ralf7x* and *lrx5x* display
elevated steady-state PME activity, in accordance with previous findings (Biermann *et al.*, 2025).
Moreover, PME activity is primarily regulated by RALF binding to LRXs. This raises the question whether
disruption of the FER-RALF-LRX pathway activates feedback mechanisms to promote PME activity in
an effort to re-establish RALF cell wall binding and signaling. Interestingly, RALF23 treatment and
overexpression in *fer* can reduce PME activity (Biermann *et al.*, 2025) and enhance powdery mildew
susceptibility. This indicates FER independent effects of RALF23 on cell wall organization and powdery
mildew infection. FER is dispensable for RALF22-induced root hair tip alkalinization, calcium influx and
cell wall modification (Schoenaers *et al.*, 2024) and RALF23 affects cell wall pectin status in a partially
FER-independent and LRX-dependent manner (Biermann *et al.*, 2025). Our SYP122-pHusion results
suggest that *Ecr* induces apoplastic pH shifts independent of FER, raising the question of receptor
redundancy or RALF-independent pH modulations, possibly through PTI (Zhai *et al.*, 2025; Wang *et al.*,
2026) or, alternatively, actively triggered by the fungus. Interestingly though, RALF23-induced
enhancement of *Ecr*-triggered apoplastic pH modulation is FER dependent. This is in line with our
findings that RALF23 interaction with FER-LLG1 supports full colonization. However, RALF23 binding
to LRXs is important for cell wall binding and essential for *Ecr* infection success. This suggests a
predominant role for RALF23 structural function and/or effects on cell wall chemistry to support *Ecr*
colonization. Yet, the interplay between RALF structural and signaling function remains largely unknown
(Schade *et al.*, 2025) . RALFs bind to LRX with high affinity at acidic pH, while LLGs require alkaline
conditions for optimal RALF recognition (Moussu *et al.*, 2020). It is possible that dynamic apoplastic pH
fluctuations induced by *Ecr* infection releases RALFs from LRXs to trigger CrRLK1L-dependent
signaling that further affects its reproductive success. Our data also raises the question whether
additional FER-related receptors are required for RALF perception to establish full *Ecr* susceptibility.
FER can interact with related RALF-binding CrRLK1Ls, including HERK1 and ANJ, to form higher order
complexes (Galindo-Trigo *et al.*, 2020; Zhong *et al.*, 2022; Lan *et al.*, 2023). This is supported by our
CoIP-MS data according to which FER interacted with several members of the CrRLK1L family, namely
HERK1, MDS1, THE1 and LET1.

The fungal haustorium is surrounded by the extrahaustorial matrix (EHMx) and the extrahaustorial
membrane (EHM), a plant derived membrane that is continuous with the plasma membrane but has
distinctive properties (Hückelhoven & Panstruga, 2011; Polonio *et al.*, 2021). At the site of invasion the
EHMx is sealed with densely packed cell wall depositions forming the haustorial neck, rendering the
EHMx a separate apoplastic compartment (Micali *et al.*, 2011). Nutrients taken up from the EHMx should
derive from the invaded plant cell and require prior export. Plant cell sugar uptake is facilitated by

SUGAR TRANSPORT PROTEINs (STP), which are associated with plant microbe interactions
(Bezrutczyk *et al.*, 2018). In Arabidopsis, STP8-GFP, upon overexpression, localizes to the EHM and
increases susceptibility to powdery mildew (Liu *et al.*, 2021) and transport-compromised STP13 alleles
are associated with resistance to powdery mildew in wheat and barley (Milne *et al.*, 2019). STPs are H^+
symporters, and STP8 has a pH optimum at 5.5 for cellular import (Liu *et al.*, 2021). It is possible that
Ecr infection induces RALF-CrRLK1L-dependent changes to the apoplastic pH that reprograms
transporter topology to favor haustorial sugar uptake over plant cell import. In accordance, *fer* mutants
show altered sucrose uptake and starch accumulation (Yang *et al.*, 2015; Yeats *et al.*, 2016). Supporting
our hypothesis, we identified several transporter proteins as FER interactors in our CoIP-MS experiment,
including STP1, involved in resistance against bacterial infection (Yamada *et al.*, 2016). We did not
identify many differential FER-GFP interactors upon *Ecr* infection, likely due to the early experimental
time point chosen (1 dpi).

We propose a model in which RALF-mediated apoplastic pH-modulation and pectin de-
methylesterification are a prerequisite for sustained powdery mildew host colonization (Figure 6). In
accordance, endogenous pectin modification affects powdery mildew infection outcome. Mutants in the
pectate lyase *POWDERY MILDEW RESISTANT 6 (PMR6)* and the pectin acetyltransferase *PMR5* are
more resistant to powdery mildew without altered fungal penetration or de-regulated SA/JA signaling
(Vogel *et al.*, 2002; Vogel *et al.*, 2004). Our data shows that RALF pathway mutants display a similar
phenotype. Moreover, the *fer-4* mutant has reduced cell wall cellulose content (Yeats *et al.*, 2016) and
dysfunctional apoplastic pH modulation which could interfere with fungal nutrient uptake. This supports
a previously coined hypothesis of altered fungal nutrition as a consequence of cell wall defects (Vogel
et al., 2002; Vogel *et al.*, 2004).

Other plant pathogens produce RALFs to establish host colonization, including *F. oxysporum* and root
knot nematodes (Masachis *et al.*, 2016; Zhang *et al.*, 2020). Interestingly, these RALF peptide mimics
lack conserved sequence elements required for LRX binding, but contain a modified YISY motif required
for LLG-CrRLK1L-dependent signaling (Pearce *et al.*, 2010). This suggests that fungal RALFs support
virulence mainly through CrRLK1L-induced signaling and less through cell wall binding. None of the
analyzed powdery mildew genomes contained RALF peptide mimics, suggesting that *Ecr* likely relies
on the combination of endogenous RALF signaling and cell wall remodeling to support sustained host
interaction. Our work thus highlights a new mode of RALF function, combining signaling and structural
roles for the control of plant microbe interactions.

Materials and Methods

Powdery mildew propagation

Erysiphe cruciferarum was propagated weekly by transferring spores to 4 – 5-week-old uninfected Col-
0 plants. Spores were used for further inoculation 3 – 4 weeks after propagation. Propagation and
experiments were performed in an environmentally controlled growth cabinet (22 °C, 70 % relative
humidity, 12 h photoperiod).

Molecular cloning

To generate p35S::RALF23 and p35::MLO2-mCherry lines gene sequences were cloned from gDNA using primers listed in Table S3 and assembled into a GoldenGate-modified pCB302 binary vector for plant expression. The p35S::RALF23$^{Y93A/Y96A}$ and p35S::RALF23$^{Y113A/Y114A}$ constructs were obtained by site-directed mutagenesis using primers listed in Table S3 and likewise assembled into a GoldenGate-modified pCB302 binary vector. For pR23::mCherry-RALF23 constructs, pR23 was cloned using primers listed in Table S3. Gene fragments for RALF23^{WT}, RALF23$^{Y93A/Y96A}$ and RALF23$^{Y113A/Y11A}$ with an N-terminal mCherry-tag located between the signal peptide and the mature RALF23 were obtained by gene synthesis (Twist Bioscience, USA). pR23 and the synthesized gene fragments were assembled into a GoldenGate-modified pCB302 binary vector for plant expression. To generate CRISPR-Cas9 mutants the software tool chopchop (https://chopchop.cbu.uib.no/) was used to design target sites (1 to 2 per gene of interest). All target sequences are listed in Table S4. Individual guide RNAs with gene specific target sites were obtained by gene synthesis (Twist Bioscience, USA). Higher order gRNA stacks were cloned into pICSL4723OD with FastRed-pRPS5::Cas9 (Castel *et al.*, 2019). All generated plant expression constructs were transformed into *Agrobacterium tumefaciens* strain GV3101 before floral dip transformation.

Plant material and growth conditions

Plant lines used in this study are listed in Table S2. All plants were grown with two plants per pot in environmentally controlled conditions (21 °C, 55 – 65% relative humidity, 8 h photoperiod). 5 – 6-week-old plants were used for *Ecr* infection.

Conidiophore counting and fungal penetration assay

Conidiophore production was analyzed 5 days after *Ecr* infection. Leaves were destained using EtOH:Acetic Acid (6:1) and subsequently stained using ink (ink: 25 % acetic acid, 9:1). Fungal structures were visualized using a Zeiss AXIO imager Z1.m microscope. Fungal penetration success was similarly analyzed and quantified at 2 dpi. A penetration attempt was scored as successful when the spore formed a penetration tube and first hyphal structures.

Quantification of fungal DNA by qPCR

For quantification of fungal DNA, two infected plants per genotype were harvested at 5 dpi before DNA extraction with standard protocols. RT-qPCR experiments were performed using Maxima SYBR green mix (Thermo Fisher Scientific, USA) on an AriaMx Real-Time PCR system (Agilent Technologies, USA). The amount of fungal DNA was calculated relative to plant DNA using primers for the Arabidopsis small RubisCO subunit (AtRbcS) and *Ecr* tubulin (Table S3).

Reinoculation experiments

5 weeks old Col-0, *fer-4*, *llg1-2* and CRISPR *ralf7x* plants were highly infected with powdery mildew. 11 days post infection spores from these plants were used to infect 5 weeks old Col-0.

Hyaloperonospora arabidopsidis (*Hpa*) infection and quantification

For *Hpa* infection assays, plants were vernalized for 2 – 3 days in the dark at 4 °C and grown in environmentally controlled conditions (22 °C, 60% relative humidity, 10 h photoperiod). 14-day-old seedlings were spray-inoculated with a spore suspension of *Hpa* isolate Noco2 (5 × 104 spores/mL) (Asai *et al.*, 2015). After inoculation, plants were kept in trays covered with sealed lids to maintain high humidity (18 °C, near-saturation humidity). At 7 days post inoculation, infected aerial parts were harvested in water and vortexed to release spores. Spores were counted using a hemocytometer and normalized to leaf fresh weight.

Infiltration experiments

Individual leaves were infiltrated with synthetic RALF23 (Sequence: ATRRYISYGALRRNTIPCSRRGASYYNCRRGAQANPYSRGCSAITRCRRS) 24 h prior to infection. Fusicoccin (10 µM), EGCG (50 µM) and MES buffers were infiltrated 2 h before *Ecr* infection.

Gene expression analysis

For RNA extraction, TRIzol reagent (Roche, Switzerland) and purification with Direct-zol™ RNA Miniprep Plus kit (Zymo Research, Germany) with on column DNAse I digestion was used. RNA was reverse transcribed with random hexamer primer and Revert Aid reverse transcriptase (Thermo Fisher Scientific, USA). RT-qPCR experiments were performed using Maxima SYBR green mix (Thermo Fisher Scientific, USA) on an AriaMx Real-Time PCR system (Agilent Technologies, USA). Gene expression levels were normalized to *UBIQUITIN5* (*UBQ5*). All primers are listed in Table S3.

PME activity assay

PME activity assays were performed as previously described (Bethke *et al.*, 2014). In brief, plant material of untreated and infected plants was harvested 5 dpi and ground in liquid nitrogen. Plant powder was incubated with 500 µL protein extraction buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 10% glycerol, 5 mM dithiothreitol, 1% protease inhibitor cocktail, 2 mM Na2MoO4, 2.5 mM NaF, 1.5 mM activated Na3VO4, 1mM phenylmethanesulfonyl fluoride, 0.5% IGEPAL) for 30 min at 8°C before centrifugation at 16 000 g for 30 min. The supernatant was transferred into a fresh 1,5 mL tube and protein concentration was determined by Bradford assay. The protein concentration was adjusted to 3 µg/µL. To assess PME activity, the 15 µL liquid was loaded into holes in an agarose gel plate containing esterified pectin (1.2 % (w/v) agarose, 0.1 % (w/v) pectin from apple, 12 mM citric acid, 50 mM Na_2HPO_4 pH 7) before incubation at 37°C for 16 h. Afterwards, plates were briefly washed with water and stained with 0.05 % ruthenium red for 30 min. Residual dye was washed off with water and images of the plates were captured by scanning. Darker stained areas correlating with PME activity were quantified using Fiji ImageJ software (Schindelin *et al.*, 2012).

AHA-phosphorylation analysis

Infected plant material was ground in liquid nitrogen. Protein extraction buffer (3 % [w/v] SDS, 30 mM TRIS-HCl pH 8.0, 10 mM EDTA. 10 mM NaF, 30 % [w/v] Sucrose, 0.012% [w/v] Coomassie Brilliant Blue, 15 % [v/v] 2-Mercaptoethanol) and extracted proteins were immediately used for SDS-

506 PAGE/western blot. Proteins were detected using AHA/pThr947 antibodies (Hayashi *et al.*, 2010). Band

507 intensities were quantified using Fiji ImageJ software.

Quantification of salicylic acid, dihydroxybenzoic acids and camalexin

SA, DHBAs and camalexin were extracted as described (Nawrath & Métraux, 1999) with minor

modifications. Ground fine leaf powder was spiked with 250 ng *ortho*-anisic acid (oANI) as internal

standard and sequentially extracted with 70% and 90% methanol for 1 hour at 65°C. Combined

supernatants were concentrated, macromolecules precipitated with 5 % trichloroacetic acid and

supernatants partitioned two times against cyclohexane / ethyl acetate (1:1) to obtain free SA, DHBAs

and camalexin. The aqueous phase was acidified with 8 M HCl (1:1) and incubated for 1 hour at 80°C.

Hydrolyzed extracts were spiked with oANI and partitioned as above to obtain glycosylated SA and

DHBAs. Organic phases of free and glycosylated fractions were concentrated and dissolved in 90% 25

mM KH_2PO_4, pH 2.6 / 10% acetonitrile (ACN). HPLC analysis was performed on an Agilent 1260 Infinity

II system with fluorescence detection. SA, DHBAs, camalexin and oANI were separated on an Agilent

InfinityLab Poroshell 120 SB-C18 column (3 x 100 mm, 2.7 µm) at 40°C, using a solvent gradient of (A)

25mM KH_2PO_4, pH 2.6 and (B) $ACN:H_2O$ (99:1) at 1 mL/min flow rate (all steps in v/v): 0 to 1 min: 10%

B; 1 to 6 min: 10 – 25 %B; 6 to 8.5 min: 25 – 80 %B; 8.5 to 10.5 min: 80% B; 10.5 to 11 min: 80 – 10%

B; 11 to 14 min 10% B. SA (Ex 305 nm, Em 407 nm), DHBAs (Ex 320 nm, Em 450 nm), camalexin (Ex

318 nm, Em 370 nm) and oANI (Ex 305 nm, Em 365 nm) were calibrated against a standard dilution

series of all metabolites of interest in the range of 1 to 100 ng.

Histochemical staining of callose deposition in rosette leaves and trichomes

Three rosette leaves per 35-d-old Arabidopsis plant were collected and destained in 70% ethanol (v/v).

Afterwards, leaves were incubated over-night in aniline blue solution (0.01% Aniline Blue (m/v) in

150 mM K_2HPO_4 buffer) and mounted on a microscopy slide. Micrographs were captured by a BZ-9000

microscope (Keyence, Osaka, Japan) using UV illumination. Images were optimized by Adobe

Photoshop 2024 in an identical manner for all micrographs. Trichome and trichome branch number were

scored by counting.

Callose quantification at the fungal penetration site

Leaves were harvested and destained in EtOH:Acetic Acid (6:1). Next, leaves were incubated for 10

min in 0.067 mM K_2HPO_4. Subsequently, leaves were transferred into staining solution with 0.05%

methyl blue in 0.067 M K_2HPO_4 and incubated overnight. Fungal structures were stained before analysis

with 0.05 % propidium iodide (PI). Methyl blues fluorescence was detected between 480 - 530 nm with

a laser at 458 nm for excitation. PI fluorescence was excited with a 510 nm laser and detected at 600 –

620 nm. Z-Projections were created and mean grey values were measured using Fiji (Schindelin *et al.*,

2012).

COS-488 synthesis, imaging and quantification

COS-488 was synthesized as described before (Mravec *et al.*, 2017). Chitosan oligosaccharides

(Carbosynth OC09272) were dissolved in 100 mM $NaCH_3CO_2$ (pH 4.9) to 1 mg/mL. AlexFluor 488

hydroxylamine was prepared in DMSO (10 mg/mL).16 µL were added to 500 µL COS solution and

incubated at 37°C in dark for two days under gentle shaking. Staining of leaf epidermis cells was
performed by harvesting leaf disks from mature Arabidopsis plants. COS-488 was infiltrated at 1:1000
dilution in ½ MS + 1% Sucrose pH 5.8 under mild vacuum. Afterwards, leaf disks were incubated in
COS-488 solution for 2 h and washed with ½ MS + 1 % Sucrose pH 5.8 three times under mild vacuum.
Samples were mounted and z-stacks acquired using a Leica SP8 confocal microscope equipped with
HC PL APO 40x water objective, argon laser and HyD detectors. COS-488 was excited at 488 nm.
Emission was detected between 490-560 nm. Z-Projections were created and mean grey values were
measured using Fiji (Schindelin *et al.*, 2012).

Confocal microscopy

Confocal laser-scanning microscopy was performed using a Leica TCS SP5 (Leica, Germany)
microscope (with Leica Application Suite X 3.7.4.23463). All images were taken with a 20 x water
immersion objective. To visualize GFP fluorescence an argon laser with HyD detector was used
(emission at 488 nm, detection 500-550 nm). mCherry fluorophore was excited with a DPSS laser
(emission 561 nm, detection 600-640 nm). and a detection window of 600 - 640 nm. Changes in the
apoplastic pH around fungal infection sites were analyzed using pUBQ10::SYP122-pHusion expressing
plants. GFP and RFP were excited with a 488 nm and a 561 nm laser, respectively. Signals were
detected at 500 – 545 nm (GFP) and 650 – 695 nm (RFP). Z-stacks with identical settings were taken
at the penetration site and in unpenetrated areas. The GFP/RFP ratios were determined using Fiji Image
J.

CoIP-MS

Sample preparation and mass spectrometric data acquisition for all CoIP samples was carried out as
previously described (Scheinost *et al.*, 2026) . Protein identification and quantification were performed
with MaxQuant software, using parameters previously described (Scheinost *et al.*, 2026) and the
following exceptions: Proteins were identified by searching MS2 spectra against the TAIR protein
database for *Arabidopsis thaliana* (downloaded March 2022, 48231 protein entries). Phosphorylation of
serine/threonine/tyrosine, oxidation of methionine and acetylation at the protein N-terminus were
specified as variable modifications. The "match-between-run" functionality was enabled (matching time
window 0.7 min, alignment time window 20 min). For protein quantification the MaxQuant output table
proteinGroups.txt and Label-Free Quantification (LFQ) (Cox et al., 2014) intensities were used. Only
proteins detected in at least two biological replicates in at least one experimental condition were further
processed. Missing values were imputed by a protein-specific constant value, which was defined as the
lowest detected protein-specific LFQ-value over all samples divided by two, using the data visualization
platform omicsViewer (DOI: 10.1101/2022.03.10.483845). Additionally, a maximal imputed LFQ value
was defined as 15% quantile of the protein distribution from the dataset. The volcano plots (Figure S6A,
S7A) were generated using RStudio. Interactions were marked as significant if they passed the threshold
of FDR < 0.05 and fold change > 2. The FER-GFP infected with *Ecr* and untreated experiments were
performed in biological triplicates, while all other experiments were performed in biological quadruplets.

Identification of putative powdery mildew RALF-likes

To identify putative RALF-like peptides in the predicted proteins and genome assemblies of selected phytopathogenic fungi, we used the hmmsearch program from the hmmer suite v3.3.2 (www.hmmer.org) with default parameters. The hidden markov model profile (.hmm) for the RALF motif (PF05498.15) was obtained from the PFAM website (https://ftp.ebi.ac.uk/pub/databases/Pfam/current_release/Pfam-A.hmm.gz), access data 19.11.2023). RALF HMM profile was used to search predicted protein sequences from the selected fungal species (see Supplementary Table S1) using the hmmsearch function. Hits with E-values below $1e^{-05}$ were retained. To perform the hmmsearch analysis on the genome assemblies, we used a custom Python script to split the contigs into 300 bp fragments with a sliding window of 50bp. All six open reading frames of these split sequences were translated and stop codons were removed to generate a multi-FASTA file suitable for be processed with hmmsearch. Search hits with E-values < 1e-05 were retained. Accession number of genome assemblies used in this analysis are reported in Supplementary Table S1.

Quantification and statistical analysis

GraphPad Prism (Version 8.0.1) was used to perform statistical analysis. Detailed descriptions of the sample size, p-values and statistical methods are indicated in the respective figure legends.

Acknowledgements

We thank Stefanie Ranf for providing GoldenGate vectors for molecular cloning and Mark Youles (TSL Norwich) and Laurence Tomlinson (TSL Norwich) for providing the pICSL4723OD vector for CRISPR-Cas9 cloning. We thank Hannah König for supporting the collection of micrographs to assess callose depositions in rosette leaves and trichomes. The authors thank Verena Breitner and Franziska Hackbarth for excellent technical assistance and maintenance of mass spectrometers at the BayBioMS. Confocal imaging was performed at the Center for Advanced Light Microscopy (CALM) of the TUM School of Life Sciences. This work was funded by the Deutsche Forschungsgemeinschaft (DFG) grants STE2448/4-1 (HL and MS), TRR356 TP B09 (HL, SS and MS), TRR356 TP B01 (JG), TRR356 TP B08 (MKRL and AM), SFB1101 TP A08 (JG), individual research grant 451218338 (MKRL), EN 1071/3-1 (TE) and PA861/20-1 (JH). The Q Exactive HFX mass spectrometer was funded in part by the DFG grant INST 95/1435-1 FUGG (CL). The work was further supported by the French government through the France 2030 investment plan managed by the National Research Agency (ANR), as part of the Initiative of Excellence of Université Côte d'Azur under reference number ANR-15-IDEX-01 (ABD), the Technical University of Munich (MS, HL, DB, JG, SS, MM, RH) and Ulm University (MS).

Author contributions

Conceptualization: MS; Investigation: HL, SS, JH, KM, GH, DB, AM, XZ, YC, MM, ABD, MS; Funding acquisition: MS, RH, ABD, MKRL, JG, TE, TK, MM, CL; Project administration: MS; Supervision: MS, HL, RH, ABD, MKRL, JG, TE, CL; Writing – original draft: HL, MS; Writing – review & editing: all authors.

Data availability

The mass spectrometric raw files as well as the MaxQuant output files have been deposited to the

ProteomeXchange Consortium via the PRIDE partner repository and can be accessed using the

identifier PXD059737 (https://proteomecentral.proteomexchange.org/cgi/GetDataset?ID=PXD059737).

Reviewer account username: reviewer_pxd059737@ebi.ac.uk, password: lumM0fo5d8wq.

Competing interests

The authors declare no competing interests.

References

Abarca A, Franck CM, Zipfel C. 2021. Family-wide evaluation of RAPID ALKALINIZATION
FACTOR peptides. *Plant Physiology* **187**: 996-1010.
Acevedo-Garcia J, Spencer D, Thieron H, Reinstädler A, Hammond-Kosack K, Phillips
AL, Panstruga R. 2017. mlo-based powdery mildew resistance in hexaploid bread
wheat generated by a non-transgenic TILLING approach. *Plant Biotechnology Journal*
15: 367-378.
Asai S, Shirasu K, Jones JDG. 2015. Hyaloperonospora arabidopsidis (Downy Mildew)
Infection Assay in Arabidopsis. *Bio-protocol* **5**: e1627.
Baunsgaard L, Fuglsang AT, Jahn T, Korthout HAAJ, De Boer AH, Palmgren MG. 1998.
The 14–3-3 proteins associate with the plant plasma membrane H1-ATPase to
generate a fusicoccin binding complex and a fusicoccin responsive system. *The Plant*
Journal **13**: 661-671.
Bethke G, Grundman RE, Sreekanta S, Truman W, Katagiri F, Glazebrook J. 2014.
Arabidopsis PECTIN METHYLESTERASEs contribute to immunity against
Pseudomonas syringae. *Plant Physiol* **164**: 1093-1107.
Bezrutczyk M, Yang J, Eom J-S, Prior M, Sosso D, Hartwig T, Szurek B, Oliva R, Vera-
Cruz C, White FF, et al. 2018. Sugar flux and signaling in plant–microbe interactions.
The Plant Journal **93**: 675-685.
Bhat RA, Miklis M, Schmelzer E, Schulze-Lefert P, Panstruga R. 2005. Recruitment and
interaction dynamics of plant penetration resistance components in a plasma
membrane microdomain. *Proceedings of the National Academy of Sciences* **102**: 3135-
3140.
Biermann D, von Arx M, Munzert-Eberlein KS, Xhelilaj K, Séré D, Stegmann M, Vert G,
Wolf S, Engelsdorf T, Zipfel C, et al. 2025. A RALF-brassinosteroid signaling circuit
regulates Arabidopsis hypocotyl cell shape. *Current Biology* **35**: 5002-5017.e5005.
Bürger M, Chory J. 2019. Stressed Out About Hormones: How Plants Orchestrate Immunity.
Cell Host & Microbe **26**: 163-172.
Büschges R, Hollricher K, Panstruga R, Simons G, Wolter M, Frijters A, van Daelen R,
van der Lee T, Diergaarde P, Groenendijk J, et al. 1997. The Barley *Mlo* Gene: A
Novel Control Element of Plant Pathogen Resistance. *Cell* **88**: 695-705.
Castel B, Tomlinson L, Locci F, Yang Y, Jones JDG. 2019. Optimization of T-DNA
architecture for Cas9-mediated mutagenesis in Arabidopsis. *PLOS ONE* **14**: e0204778.
Chakravorty D, Yu Y, Assmann SM. 2018. A kinase-dead version of FERONIA receptor-like
kinase has dose-dependent impacts on rosette morphology and RALF1-mediated
stomatal movements. *FEBS Letters* **592**: 3429-3437.
Consonni C, Humphry ME, Hartmann HA, Livaja M, Durner J, Westphal L, Vogel J, Lipka
V, Kemmerling B, Schulze-Lefert P, et al. 2006. Conserved requirement for a plant
host cell protein in powdery mildew pathogenesis. *Nat Genet* **38**: 716-720.
Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. 2014. Accurate Proteome-wide
Label-free Quantification by Delayed Normalization and Maximal Peptide Ratio
Extraction, Termed MaxLFQ*. *Molecular & Cellular Proteomics* **13**: 2513-2526.
Delulio GA, Guo L, Zhang Y, Goldberg JM, Kistler HC, Ma L-J. 2018. Kinome Expansion in
the Fusarium oxysporum Species Complex Driven by Accessory Chromosomes.
mSphere **3**: 10.1128/msphere.00231-00218.
Del Corpo D, Fullone MR, Miele R, Lafond M, Pontiggia D, Grisel S, Kieffer-Jaquinod S,
Giardina T, Bellincampi D, Lionetti V. 2020. AtPME17 is a functional Arabidopsis
thaliana pectin methylesterase regulated by its PRO region that triggers PME activity
in the resistance to Botrytis cinerea. *Molecular Plant Pathology* **21**: 1620-1633.
Deslauriers SD, Larsen PB. 2010. FERONIA Is a Key Modulator of Brassinosteroid and
Ethylene Responsiveness in Arabidopsis Hypocotyls. *Molecular Plant* **3**: 626-640.
Dobón A, Canet JV, García-Andrade J, Angulo C, Neumetzler L, Persson S, Vera P. 2015.
Novel Disease Susceptibility Factors for Fungal Necrotrophic Pathogens in
Arabidopsis. *PLOS Pathogens* **11**: e1004800.

Draeger C, Ndinyanka Fabrice T, Gineau E, Mouille G, Kuhn BM, Moller I, Abdou MT, Frey B, Pauly M, Bacic A, et al. 2015. Arabidopsis leucine-rich repeat extensin (LRX) proteins modify cell wall composition and influence plant growth. *BMC Plant Biol* **15**: 155.

Du C, Li X, Chen J, Chen W, Li B, Li C, Wang L, Li J, Zhao X, Lin J, et al. 2016. Receptor kinase complex transmits RALF peptide signal to inhibit root growth in Arabidopsis. *Proceedings of the National Academy of Sciences* **113**: E8326-E8334.

Duan Q, Kita D, Li C, Cheung AY, Wu H-M. 2010. FERONIA receptor-like kinase regulates RHO GTPase signaling of root hair development. *Proceedings of the National Academy of Sciences* **107**: 17821-17826.

Dünser K, Gupta S, Herger A, Feraru MI, Ringli C, Kleine-Vehn J. 2019. Extracellular matrix sensing by FERONIA and Leucine Rich Repeat Extensins controls vacuolar expansion during cellular elongation in Arabidopsis thaliana. *The EMBO Journal* **38**: e100353.

Engelhardt S, Stam R, Hückelhoven R. 2018. Good Riddance? Breaking Disease Susceptibility in the Era of New Breeding Technologies. *Agronomy* **8**: 114.

Engelsdorf T, Gigli-Bisceglia N, Veerabagu M, McKenna JF, Vaahtera L, Augstein F, Van der Does D, Zipfel C, Hamann T. 2018. The plant cell wall integrity maintenance and immune signaling systems cooperate to control stress responses in Arabidopsis thaliana. *Science Signaling* **11**: eaao3070.

Fabrice TN, Vogler H, Draeger C, Munglani G, Gupta S, Herger AG, Knox P, Grossniklaus U, Ringli C. 2018. LRX Proteins Play a Crucial Role in Pollen Grain and Pollen Tube Cell Wall Development. *Plant Physiol* **176**: 1981-1992.

Falhof J, Pedersen Jesper T, Fuglsang Anja T, Palmgren M. 2016. Plasma Membrane H+-ATPase Regulation in the Center of Plant Physiology. *Molecular Plant* **9**: 323-337.

Felle HH, Herrmann A, Hanstein S, Hückelhoven R, Kogel K-H. 2004. Apoplastic pH Signaling in Barley Leaves Attacked by the Powdery Mildew Fungus Blumeria graminis f. sp. hordei. *Molecular Plant-Microbe Interactions* **17**: 118-123.

Galindo-Trigo S, Blanco-Touriñán N, DeFalco TA, Wells ES, Gray JE, Zipfel C, Smith LM. 2020. *Cr*RLK1L receptor-like kinases HERK1 and ANJEA are female determinants of pollen tube reception. *EMBO reports* **21**: e48466.

Gao Q, Wang C, Xi Y, Shao Q, Li L, Luan S. 2022. A receptor–channel trio conducts Ca2+ signalling for pollen tube reception. *Nature* **607**: 534-539.

Ge Z, Bergonci T, Zhao Y, Zou Y, Du S, Liu M-C, Luo X, Ruan H, García-Valencia LE, Zhong S, et al. 2017. Arabidopsis pollen tube integrity and sperm release are regulated by RALF-mediated signaling. *Science* **358**: 1596-1600.

Glawe DA. 2008. The Powdery Mildews: A Review of the World's Most Familiar (Yet Poorly Known) Plant Pathogens. *Annual Review of Phytopathology* **46**: 27-51.

Gonneau M, Desprez T, Martin M, Doblas VG, Bacete L, Miart F, Sormani R, Hématy K, Renou J, Landrein B, et al. 2018. Receptor Kinase THESEUS1 Is a Rapid Alkalinization Factor 34 Receptor in Arabidopsis. *Current Biology* **28**: 2452-2458.e2454.

Gronnier J, Franck CM, Stegmann M, DeFalco TA, Abarca A, von Arx M, Dünser K, Lin W, Yang Z, Kleine-Vehn J, et al. 2022. Regulation of immune receptor kinase plasma membrane nanoscale organization by a plant peptide hormone and its receptors. *Elife* **11**: e74162.

Guénin S, Mareck A, Rayon C, Lamour R, Assoumou Ndong Y, Domon J-M, Sénéchal F, Fournet F, Jamet E, Canut H, et al. 2011. Identification of pectin methylesterase 3 as a basic pectin methylesterase isoform involved in adventitious rooting in Arabidopsis thaliana. *New Phytologist* **192**: 114-126.

Guo H, Nolan TM, Song G, Liu S, Xie Z, Chen J, Schnable PS, Walley JW, Yin Y. 2018. FERONIA Receptor Kinase Contributes to Plant Immunity by Suppressing Jasmonic Acid Signaling in Arabidopsis thaliana. *Curr Biol* **28**: 3316-3324.e3316.

Haruta M, Burch HL, Nelson RB, Barrett-Wilt G, Kline KG, Mohsin SB, Young JC, Otegui MS, Sussman MR. 2010. Molecular Characterization of Mutant Arabidopsis Plants with Reduced Plasma Membrane Proton Pump Activity *Journal of Biological Chemistry* **285**: 17918-17929.

Haruta M, Sabat G, Stecker K, Minkoff BB, Sussman MR. 2014. A peptide hormone and its
receptor protein kinase regulate plant cell expansion. *Science* **343**: 408-411.
Hayashi Y, Nakamura S, Takemiya A, Takahashi Y, Shimazaki K-i, Kinoshita T. 2010.
Biochemical Characterization of In Vitro Phosphorylation and Dephosphorylation of the
Plasma Membrane H+-ATPase. *Plant and Cell Physiology* **51**: 1186-1196.
Herger A, Gupta S, Kadler G, Franck CM, Boisson-Dernier A, Ringli C. 2020. Overlapping
functions and protein-protein interactions of LRR-extensins in Arabidopsis. *PLOS*
Genetics **16**: e1008847.
Hocq L, Habrylo O, Sénéchal F, Voxeur A, Pau-Roblot C, Safran J, Fournet F, Bassard
S, Battu V, Demailly H, et al. 2023. Mutation of AtPME2, a pH-Dependent Pectin
Methylesterase, Affects Cell Wall Structure and Hypocotyl Elongation. *Plant and Cell*
Physiology **65**: 301-318.
Hocq L, Sénéchal F, Lefebvre V, Lehner A, Domon J-M, Mollet J-C, Dehors J, Pageau K,
Marcelo P, Guérineau F, et al. 2016. Combined Experimental and Computational
Approaches Reveal Distinct pH Dependence of Pectin Methylesterase Inhibitors. *Plant*
Physiology **173**: 1075-1093.
Hok S, Allasia V, Andrio E, Naessens E, Ribes E, Panabières F, Attard A, Ris N, Clément
M, Barlet X, et al. 2014. The Receptor Kinase IMPAIRED OOMYCETE
SUSCEPTIBILITY1 Attenuates Abscisic Acid Responses in Arabidopsis   *Plant*
Physiology **166**: 1506-1518.
Hok S, Danchin EGJ, Allasia V, Panabières F, Attard A, Keller H. 2011. An Arabidopsis
(malectin-like) leucine-rich repeat receptor-like kinase contributes to downy mildew
disease. *Plant, Cell & Environment* **34**: 1944-1957.
Hückelhoven R, Panstruga R. 2011. Cell biology of the plant–powdery mildew interaction.
Current Opinion in Plant Biology **14**: 738-746.
Huebbers JW, Caldarescu GA, Kubátová Z, Sabol P, Levecque SCJ, Kuhn H, Kulich I,
Reinstädler A, Büttgen K, Manga-Robles A, et al. 2024. Interplay of EXO70 and
MLO proteins modulates trichome cell wall composition and susceptibility to powdery
mildew. *The Plant Cell* **36**: 1007-1035.
Kadota Y, Sklenar J, Derbyshire P, Stransfeld L, Asai S, Ntoukakis V, Jones Jonathan D,
Shirasu K, Menke F, Jones A, et al. 2014. Direct Regulation of the NADPH Oxidase
RBOHD by the PRR-Associated Kinase BIK1 during Plant Immunity. *Molecular Cell* **54**:
43-55.
Kessler SA, Shimosato-Asano H, Keinath NF, Wuest SE, Ingram G, Panstruga R,
Grossniklaus U. 2010. Conserved Molecular Components for Pollen Tube Reception
and Fungal Invasion. *Science* **330**: 968-971.
Kesten C, Gámez-Arjona FM, Menna A, Scholl S, Dora S, Huerta AI, Huang HY, Tintor N,
Kinoshita T, Rep M, et al. 2019. Pathogen induced pH changes regulate the growth
defense balance in plants. *The EMBO Journal* **38**: e101822.
Kusch S, Panstruga R. 2017. mlo-Based Resistance: An Apparently Universal "Weapon" to
Defeat Powdery Mildew Disease. *Molecular Plant-Microbe Interactions®* **30**: 179-189.
Lan Z, Song Z, Wang Z, Li L, Liu Y, Zhi S, Wang R, Wang J, Li Q, Bleckmann A, et al.
2023. Antagonistic RALF peptides control an intergeneric hybridization barrier on
Brassicaceae stigmas. *Cell* **186**: 4773-4787.e4712.
Lewis KC, Selzer T, Shahar C, Udi Y, Tworowski D, Sagi I. 2008. Inhibition of pectin methyl
esterase activity by green tea catechins. *Phytochemistry* **69**: 2586-2592.
Li C, Yeh F-L, Cheung AY, Duan Q, Kita D, Liu M-C, Maman J, Luu EJ, Wu BW, Gates L,
et al. 2015. Glycosylphosphatidylinositol-anchored proteins as chaperones and co-
receptors for FERONIA receptor kinase signaling in Arabidopsis. *Elife* **4**: e06587.
Liu J, Liu M, Tan L, Huai B, Ma X, Pan Q, Zheng P, Wen Y, Zhang Q, Zhao Q, et al. 2021.
AtSTP8, an endoplasmic reticulum-localised monosaccharide transporter from
Arabidopsis, is recruited to the extrahaustorial membrane during powdery mildew
infection. *New Phytologist* **230**: 2404-2419.
Liu M-CJ, Yeh F-LJ, Yvon R, Simpson K, Jordan S, Chambers J, Wu H-M, Cheung AY.
2024. Extracellular pectin-RALF phase separation mediates FERONIA global signaling
function. *Cell* **187**: 312-330.e322.

Liu S, Bartnikas LM, Volko SM, Ausubel FM, Tang D. 2016. Mutation of the Glucosinolate
Biosynthesis Enzyme Cytochrome P450 83A1 Monooxygenase Increases Camalexin
Accumulation and Powdery Mildew Resistance. *Frontiers in Plant Science* **7**.
Liu Z, Chu X, Ren W, Cheng L, Liu C, Wang C, Gao S, Dai S, Li C. 2025. PCP-B peptides
and CrRLK1L receptor kinases control pollination via pH gating of aquaporins in
Arabidopsis. *Developmental Cell* **60**: 1336-1347.e1335.
**Manning VA, Pandelova I, Dhillon B, Wilhelm LJ, Goodwin SB, Berlin AM, Figueroa M,
Freitag M, Hane JK, Henrissat B, et al. 2013.** Comparative Genomics of a Plant-
Pathogenic Fungus, Pyrenophora tritici-repentis, Reveals Transduplication and the
Impact of Repeat Elements on Pathogenicity and Population Divergence. *G3
Genes|Genomes|Genetics* **3**: 41-63.
**Masachis S, Segorbe D, Turrà D, Leon-Ruiz M, Fürst U, El Ghalid M, Leonard G, López-
Berges MS, Richards TA, Felix G, et al. 2016.** A fungal pathogen secretes plant
alkalinizing peptides to increase infection. *Nature Microbiology* **1**: 16043.
**Mecchia MA, Santos-Fernandez G, Duss NN, Somoza SC, Boisson-Dernier A,
Gagliardini V, Martínez-Bernardini A, Fabrice TN, Ringli C, Muschietti JP, et al.
2017.** RALF4/19 peptides interact with LRX proteins to control pollen tube growth in
Arabidopsis. *Science* **358**: 1600-1603.
**Merlot S, Leonhardt N, Fenzi F, Valon C, Costa M, Piette L, Vavasseur A, Genty B, Boivin
K, Müller A, et al. 2007.** Constitutive activation of a plasma membrane H+ATPase
prevents abscisic acid-mediated stomatal closure. *The EMBO Journal* **26**: 3216-3226.
Micali C, Göllner K, Humphry M, Consonni C, Panstruga R. 2008. The Powdery Mildew
Disease of Arabidopsis: A Paradigm for the Interaction between Plants and Biotrophic
Fungi. *Arabidopsis Book* **6**: e0115.
Micali CO, Neumann U, Grunewald D, Panstruga R, O'Connell R. 2011. Biogenesis of a
specialized plant–fungal interface during host cell internalization of Golovinomyces
orontii haustoria. *Cellular Microbiology* **13**: 210-226.
**Milne RJ, Dibley KE, Schnippenkoetter W, Mascher M, Lui ACW, Wang L, Lo C, Ashton
AR, Ryan PR, Lagudah ES. 2019.** The Wheat Lr67 Gene from the Sugar Transport
Protein 13 Family Confers Multipathogen Resistance in Barley. *Plant Physiol* **179**:
1285-1297.
**Moussu S, Broyart C, Santos-Fernandez G, Augustin S, Wehrle S, Grossniklaus U,
Santiago J. 2020.** Structural basis for recognition of RALF peptides by LRX proteins
during pollen tube growth. *Proceedings of the National Academy of Sciences* **117**:
7494-7503.
**Moussu S, Lee HK, Haas KT, Broyart C, Rathgeb U, De Bellis D, Levasseur T,
Schoenaers S, Fernandez GS, Grossniklaus U, et al. 2023.** Plant cell wall patterning
and expansion mediated by protein-peptide-polysaccharide interaction. *Science* **382**:
719-725.
**Mravec J, Kračun SK, Rydahl MG, Westereng B, Pontiggia D, De Lorenzo G, Domozych
DS, Willats WGT. 2017.** An oligogalacturonide-derived molecular probe demonstrates
the dynamics of calcium-mediated pectin complexation in cell walls of tip-growing
structures. *Plant J* **91**: 534-546.
**Mucha S, Heinzlmeir S, Kriechbaumer V, Strickland B, Kirchhelle C, Choudhary M,
Kowalski N, Eichmann R, Hückelhoven R, Grill E, et al. 2019.** The Formation of a
Camalexin Biosynthetic Metabolon. *The Plant Cell* **31**: 2697-2710.
**Müller MC, Praz CR, Sotiropoulos AG, Menardo F, Kunz L, Schudel S, Oberhänsli S,
Poretti M, Wehrli A, Bourras S, et al. 2019.** A chromosome-scale genome assembly
reveals a highly dynamic effector repertoire of wheat powdery mildew. *New Phytologist*
221: 2176-2189.
Nawrath C, Métraux J-P. 1999. Salicylic Acid Induction–Deficient Mutants of Arabidopsis
Express PR-2 and PR-5 and Accumulate High Levels of Camalexin after Pathogen
Inoculation. *The Plant Cell* **11**: 1393-1404.
Nekrasov V, Wang C, Win J, Lanz C, Weigel D, Kamoun S. 2017. Rapid generation of a
transgene-free powdery mildew resistant tomato by genome deletion. *Scientific
Reports* **7**: 482.

Ngou BPM, Jones JDG, Ding P. 2022. Plant immune networks. *Trends in Plant Science* **27**:
255-273.
Noble JA, Bielski NV, Liu M-CJ, DeFalco TA, Stegmann M, Nelson ADL, McNamara K,
Sullivan B, Dinh KK, Khuu N, et al. 2022. Evolutionary analysis of the LORELEI gene
family in plants reveals regulatory subfunctionalization. *Plant Physiology* **190**: 2539-
2556.
Pearce G, Yamaguchi Y, Munske G, Ryan CA. 2010. Structure–activity studies of RALF,
Rapid Alkalinization Factor, reveal an essential – YISY – motif. *Peptides* **31**: 1973-
1977.
Peaucelle A, Louvet R, Johansen JN, Höfte H, Laufs P, Pelloux J, Mouille G. 2008.
Arabidopsis Phyllotaxis Is Controlled by the Methyl-Esterification Status of Cell-Wall
Pectins. *Current Biology* **18**: 1943-1948.
Peng Y, Yang J, Li X, Zhang Y. 2021. Salicylic Acid: Biosynthesis and Signaling. *Annual*
Review of Plant Biology **72**: 761-791.
Polonio Á, Pérez-García A, Martínez-Cruz J, Fernández-Ortuño D, de Vicente A 2021. The
Haustorium of Phytopathogenic Fungi: A Short Overview of a Specialized Cell of
Obligate Biotrophic Plant Parasites. In: Cánovas FM, Lüttge U, Risueño M-C, Pretzsch
H eds. *Progress in Botany Vol. 82*. Cham: Springer International Publishing, 337-355.
Raiola A, Lionetti V, Elmaghraby I, Immerzeel P, Mellerowicz EJ, Salvi G, Cervone F,
Bellincampi D. 2011. Pectin Methylesterase Is Induced in Arabidopsis upon Infection
and Is Necessary for a Successful Colonization by Necrotrophic Pathogens. *Molecular*
Plant-Microbe Interactions® **24**: 432-440.
Ried MK, Banhara A, Hwu F-Y, Binder A, Gust AA, Höfle C, Hückelhoven R, Nürnberger
T, Parniske M. 2019. A set of Arabidopsis genes involved in the accommodation of the
downy mildew pathogen Hyaloperonospora arabidopsidis. *PLOS Pathogens* **15**:
e1007747.
Rößling A-K, Dünser K, Liu C, Lauw S, Rodriguez-Franco M, Kalmbach L, Barbez E,
Kleine-Vehn J. 2024. Pectin methylesterase activity is required for RALF1 peptide
signalling output. *Elife* **13**: RP96943.
Rzemieniewski J, Stegmann M. 2022. Regulation of pattern-triggered immunity and growth
by phytocytokines. *Current Opinion in Plant Biology* **68**: 102230.
Schade S, Leicher H, von Arx M, Monte I, Gronnier J, Stegmann M. 2025. The interplay of
RALF structural and signaling functions in plant-microbe interactions. *PLOS Pathogens*
21: e1013588.
Scheinost L, Ludwig C, Höfflin N, Taskin AA, Marada A, Vögtle F-N, Meisinger C, Köhn
M. 2026. Synthetic trap-peptides identify a TOM complex phosphatase – PP2A
dephosphorylates Tom6. *The FEBS Journal* **293**: 271-294.
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, et al. 2012. Fiji: an open-source platform for
biological-image analysis. *Nature Methods* **9**: 676-682.
Schmid M, Davison TS, Henz SR, Pape UJ, Demar M, Vingron M, Schölkopf B, Weigel D,
Lohmann JU. 2005. A gene expression map of Arabidopsis thaliana development. *Nat*
Genet **37**: 501-506.
Schoenaers S, Lee HK, Gonneau M, Faucher E, Levasseur T, Akary E, Claeijs N, Moussu
S, Broyart C, Balcerowicz D, et al. 2024. Rapid alkalinization factor 22 has a structural
and signalling role in root hair cell wall assembly. *Nature Plants* **10**: 494-511.
Sénéchal F, Habrylo O, Hocq L, Domon J-M, Marcelo P, Lefebvre V, Pelloux J,
Mercadante D. 2017. Structural and dynamical characterization of the pH-dependence
of the pectin methylesterase-pectin methylesterase inhibitor complex. *Journal of*
Biological Chemistry **292**: 21538-21547.
Shen Q, Bourdais G, Pan H, Robatzek S, Tang D. 2017. *Arabidopsis*
glycosylphosphatidylinositol-anchored protein LLG1 associates with and modulates
FLS2 to regulate innate immunity. *Proceedings of the National Academy of Sciences*
114: 5749-5754.

Stegmann M, Monaghan J, Smakowska-Luzan E, Rovenich H, Lehner A, Holton N,
Belkhadir Y, Zipfel C. 2017. The receptor kinase FER is a RALF-regulated scaffold
controlling plant immune signaling. *Science* **355**: 287-289.
Sun M, Lu B, Yang Y, Fan J, Ren W, Chu X, Gao Y, Wu J, Wang J, Ke H, et al. 2025.
Unveiling FERONIA receptor kinase–mediated cellular mechanisms with a small-
molecule inhibitor. *Proceedings of the National Academy of Sciences* **122**:
e2515322122.
Tang J, Wu D, Li X, Wang L, Xu L, Zhang Y, Xu F, Liu H, Xie Q, Dai S, et al. 2022. Plant
immunity suppression via PHR1-RALF-FERONIA shapes the root microbiome to
alleviate phosphate starvation. *The EMBO Journal* **41**: e109102.
Tang W, Lin W, Zhou X, Guo J, Dang X, Li B, Lin D, Yang Z. 2022. Mechano-transduction
via the pectin-FERONIA complex activates ROP6 GTPase signaling in
Arabidopsis pavement cell morphogenesis. *Current Biology* **32**: 508-
517.e503.
Thynne E, Saur IML, Simbaqueba J, Ogilvie HA, Gonzalez-Cendales Y, Mead O, Taranto
A, Catanzariti A-M, McDonald MC, Schwessinger B, et al. 2017. Fungal
phytopathogens encode functional homologues of plant rapid alkalinization factor
(RALF) peptides. *Molecular Plant Pathology* **18**: 811-824.
van Schie CCN, Takken FLW. 2014. Susceptibility Genes 101: How to Be a Good Host.
Annual Review of Phytopathology **52**: 551-581.
Vogel JP, Raab TK, Schiff C, Somerville SC. 2002. PMR6, a Pectate Lyase–Like Gene
Required for Powdery Mildew Susceptibility in Arabidopsis. *The Plant Cell* **14**: 2095-
2106.
Vogel JP, Raab TK, Somerville CR, Somerville SC. 2004. Mutations in PMR5 result in
powdery mildew resistance and altered cell wall composition. *The Plant Journal* **40**:
968-978.
Wang H, Li X, Zhai K, Rhodes J, Sang T, Zhao J, Gao Y, Ma S, Song B, Pan Q, et al. 2026.
A regulatory network promotes apoplastic alkalinization to prime plant immunity in
tissues distal to site of infection. *Cell*.
Weis C, Pfeilmeier S, Glawischnig E, Isono E, Pachl F, Hahne H, Kuster B, Eichmann R,
Hückelhoven R. 2013. Co-immunoprecipitation-based identification of putative BAX
INHIBITOR-1-interacting proteins involved in cell death regulation and plant–powdery
mildew interactions. *Molecular Plant Pathology* **14**: 791-802.
Willats WGT, Orfila C, Limberg G, Buchholt HC, van Alebeek G-JWM, Voragen AG,
Marcus SE, Christensen TMIE, Mikkelsen JD, Murray BS, et al. 2001. Modulation
of the Degree and Pattern of Methyl-esterification of Pectic Homogalacturonan in Plant
Cell Walls: IMPLICATIONS FOR PECTIN METHYL ESTERASE ACTION, MATRIX
PROPERTIES, AND CELL ADHESION *Journal of Biological Chemistry* **276**: 19404-
19413.
Wu Y, Ma X, Pan Z, Kale SD, Song Y, King H, Zhang Q, Presley C, Deng X, Wei C-I, et al.
2018. Comparative genome analyses reveal sequence features reflecting distinct
modes of host-adaptation between dicot and monocot powdery mildew. *BMC*
Genomics **19**: 705.
Xiao Y, Stegmann M, Han Z, DeFalco TA, Parys K, Xu L, Belkhadir Y, Zipfel C, Chai J.
2019. Mechanisms of RALF peptide perception by a heterotypic receptor complex.
Nature **572**: 270-274.
Xu F, Gonneau M, Faucher E, Habrylo O, Lefebvre V, Domon J-M, Martin M, Sénéchal F,
Peaucelle A, Pelloux J, et al. 2022. Biochemical characterization of Pectin
Methylesterase Inhibitor 3 from Arabidopsis thaliana. *The Cell Surface* **8**: 100080.
Xu X, Chen X. 2023. A complete genome sequence of Podosphaera xanthii race 2 F, the
causal agent of powdery mildew isolated from cucumber in China. *Pathogens* **12**: 561.
Yamada K, Saijo Y, Nakagami H, Takano Y. 2016. Regulation of sugar transporter activity
for antibacterial defense in Arabidopsis. *Science* **354**: 1427-1430.
Yang T, Wang L, Li C, Liu Y, Zhu S, Qi Y, Liu X, Lin Q, Luan S, Yu F. 2015. Receptor protein
kinase FERONIA controls leaf starch accumulation by interacting with glyceraldehyde-

3-phosphate dehydrogenase. *Biochemical and Biophysical Research Communications*
465: 77-82.

Yeats TH, Sorek H, Wemmer DE, Somerville CR. 2016. Cellulose Deficiency Is Enhanced
on Hyper Accumulation of Sucrose by a H+-Coupled Sucrose Symporter *Plant*
Physiology **171**: 110-124.

Zaccaron AZ, Neill T, Corcoran J, Mahaffee WF, Stergiopoulos I. 2023. A chromosome-
scale genome assembly of the grape powdery mildew pathogen *Erysiphe necator*
reveals its genomic architecture and previously unknown features of its biology. *mBio*
14: e00645-00623.

Zecua-Ramirez P, Jha V, Stegmann M. 2026. Peptide signaling at the intersection of growth,
nutrient sensing, and stress responses. *The Plant Journal* **125**: e70733.

Zhai K, Derbyshire P, Zhang S, Choi S, Wang L, Song B, Kinoshita T, Zhou J-M, Menke
FLH, Bender KW, et al. 2025. A Phosphorelay Circuit Drives Extracellular
Alkalinization in Plant Receptor Kinase Signaling. *bioRxiv*: 2025.2008.2016.670655.

Zhang K, Halitschke R, Yin C, Liu C-J, Gan S-S. 2013. Salicylic acid 3-hydroxylase regulates
Arabidopsis leaf longevity by mediating salicylic acid catabolism. *Proceedings of the*
National Academy of Sciences **110**: 14807-14812.

Zhang X, Peng H, Zhu S, Xing J, Li X, Zhu Z, Zheng J, Wang L, Wang B, Chen J, et al.
2020. Nematode-Encoded RALF Peptide Mimics Facilitate Parasitism of Plants
through the FERONIA Receptor Kinase. *Molecular Plant* **13**: 1434-1454.

Zhang Y, Zhao L, Zhao J, Li Y, Wang J, Guo R, Gan S, Liu C-J, Zhang K. 2017. S5H/DMR6
Encodes a Salicylic Acid 5-Hydroxylase That Fine-Tunes Salicylic Acid
Homeostasis *Plant Physiology* **175**: 1082-1093.

Zhao C, Zayed O, Yu Z, Jiang W, Zhu P, Hsu C-C, Zhang L, Tao WA, Lozano-Durán R,
Zhu J-K. 2018. Leucine-rich repeat extensin proteins regulate plant salt tolerance in
Arabidopsis. *Proceedings of the National Academy of Sciences* **115**: 13123-13128.

Zhong S, Li L, Wang Z, Ge Z, Li Q, Bleckmann A, Wang J, Song Z, Shi Y, Liu T, et al.
2022. RALF peptide signaling controls the polytubey block in Arabidopsis. *Science* **375**:
290-296.

Zhou F, Andersen CH, Burhenne K, Fischer PH, Collinge DB, Thordal-Christensen H.
2000. Proton extrusion is an essential signalling component in the HR of epidermal
single cells in the barley-powdery mildew interaction. *Plant J* **23**: 245-254.

Zhu S, Fu Q, Xu F, Zheng H, Yu F. 2021. New paradigms in cell adaptation: decades of
discoveries on the CrRLK1L receptor kinase signalling network. *New Phytologist* **232**:
1168-1183.

Zimmerli L, Stein M, Lipka V, Schulze-Lefert P, Somerville S. 2004. Host and non-host
pathogens elicit different jasmonate/ethylene responses in Arabidopsis. *The Plant*
Journal **40**: 633-646.

Figures

Figure 1: FER and additional RALF binding proteins are powdery mildew susceptibility factors.
A) Images of Col-0 and *fer-4* plants infected with *Ecr* (14 dpi). B) Ink stained *Ecr* colonies (5 dpi) grown on Col-0 and *fer-4* plants. Black arrows indicate conidiophores. Scale bar represents 50 μm. C) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=223-392 pooled from three independent experiments (Dunn's multiple comparisons test, a-b, b-c p<0.0001; a-c p=0.0106). D) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=345-665 pooled from five independent experiments (Dunn's multiple comparisons test, a-b, a-c p<0.0001; b-c p=0.0235). E) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=285-453 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p=0.0150; a-c p<0.0001; b-c p=0.0408). F) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes (LRR4$^{\Delta E}$: LRX4 lacking the extensin domain). Mean ± SD, n=113-159 pooled from two independent experiments (Dunn's multiple comparisons test, a-b p<0.0001). All experiments were done at least three times with similar results, except F), which was repeated once with identical results.

1017

Figure 2: Specific RALFs are powdery mildew susceptibility factors. A) Table of RALF peptides mutated in CRISPR *ralf3x*, CRISPR *ralf7x* and *ralf1/22/23/33.* B) Images of Col-0*, fer-4*, CRISPR *ralf3x* and CRISPR *ralf7x* plants infected with *Ecr* (14 dpi). C) Ink stained *Ecr* colonies (5 dpi) grown on Col-0, *fer-4*, CRISPR *ralf3x* and CRISPR *ralf7x* plants. Black arrows indicate conidiophores. Scale bar represents 50 µm. D) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=98-456 pooled from three independent experiments (Dunn's multiple comparisons test, a-b, a-c p<0.0001, b-c p<0.0002. Comparison between *fer-4* and CRISPR *ralf7x*: Mann Whitney test, ** p=0.007). E) Amount of fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection of the indicated genotypes, n=4, data points with different symbols indicate independent biological replicates. F) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=101-286 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p<0.0001). G) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=226-382 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p<0.0001. Comparison between *fer-4* and *ralf1/22/23/33*: Mann Whitney test, ** p=0.0132). All experiments were performed at least three times with similar results.

1034

1035

1036

Figure 3: FER-RALF mediated powdery mildew susceptibility is independent of JA and SA signaling. A) Quantification of free (blue) and glycosylated (green) SA in untreated and *Ecr* infected (5 dpi) leaves of the indicated genotypes. Mean ± SD, n=3 pooled from three independent experiments (Tukey's multiple comparisons test, free and glycosylated SA was analyzed individually, free: a-bc p=0.0423; a-c p=0.0006; ab-c p≤0.0013, glycosylated: A-C p≤0.0029; A-BC p≤0.0383). B) Quantification of free (blue) and glycosylated (green) 2,5-DHBA in untreated and *Ecr* infected (5 dpi) leaves of the indicated genotypes. Mean ± SD, n=3 pooled from three independent experiments (Tukey's multiple comparisons test, free and glycosylated 2,5-DHBA was analyzed individually, free: not significant, glycosylated: A-C p=0.0153; A-BC p=0.0342; AB-C p=0.0345). C) Quantification of free (blue) Camalexin in untreated and *Ecr* infected (5 dpi) leaves of the indicated genotypes. Mean ± SD, n=3 pooled from three independent experiments (Tukey's multiple comparisons test, a-bc p=0.0486; ab-c p≤0.0319; a-c p=0.0137. D) RT-qPCR of *PR1* in untreated (grey) adult leaves and upon infection with *Ecr* (5 dpi, green). Housekeeping gene *UBQ5*. Mean ± SD, n=3, data points with different symbols indicate independent biological replicates (Tukey's multiple comparisons test, a-d p<0.0001; a-cb p≤0.0464; a-cd p≤0.0137; ab-cd p≤0.0161). E) RT-qPCR of *JAZ10* in untreated (grey) adult leaves and upon infection with *Ecr* (5 dpi, green). Housekeeping gene *UBQ5*. Mean ± SD, n=3, data points with different symbols indicate independent biological replicates (Tukey's multiple comparisons test, not significant) F) RT-qPCR of *VSP2* in untreated (grey) adult leaves and upon infection with *Ecr* (5 dpi, green). Housekeeping gene *UBQ5*. Mean ± SD, n=3, data points with different symbols indicate independent biological replicates (Tukey's multiple comparisons test, not significant). G) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=152-209 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p<0.0001). All experiments were performed three times with similar results.

1060

Figure 4: Apoplastic pH modulation and PME activity determine powdery mildew infection success. A) MS Intensities measured after Co-immunoprecipitation with untreated pFER::FER-GFP and p35S::Lti6b-GFP plants using GFP-TRAP beads. n=3 for pFER::FER-GFP, n=4 for p35S::Lti6b-GFP, data points indicate independent biological replicates. FDR values indicate statistical differences between pFER::FER-GFP and p35S::Lti6b-GFP. No AHA3 derived peptides were detected in p35S::Lti6b-GFP samples. B) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=49-91 pooled from two independent experiments (Mann Whitney test, a-b p<0.0001). C) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of leaves pretreated with H_2O (blue) or Fusicoccin (green). Mean ± SD, n=232-316 pooled from seven independent experiments (Mann-Whitney test, genotypes were analyzed separately, Col-0: a-b p=0.0058, *fer-4*: A-B p=0.002). D) Amount of fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection and pretreatment with H_2O (blue) or Fusicoccin (green). n=3, data points with different symbols indicate independent biological replicates. All data is normalized to Col-0 (H_2O). E) Conidiophores per field of view (5 dpi) after pretreatment with H_2O (blue), $MgCl_2$ 15 mM pH 7 (grey), MES buffer 10 mM pH 5.5 (green) and MES buffer 10 mM pH 6.5 (red) before *Ecr* infection. Mean ± SD, n=126-195 pooled from three independent experiments (Dunn's multiple comparison, a-bc, a-dc, a-d, a-e, ab-d, ab-e, bc-3 p<0.0001, ab-cd p≤0.0086, bc-d p=0.0049, d-e p≤0.034). F) Ink stained *Ecr* colonies (5 dpi) grown on Col-0 and *fer-4* plants after pretreatment with 15 mM $MgCl_2$, 10 mM MES buffer (pH 5.5) and 10 mM MES buffer (pH 6.5). Black arrows indicate conidiophores. Scale bar represents 50 µm. G) PME activity as ruthenium red stained halo size in pixel in untreated leaves (grey) and upon *Ecr* infection (5 dpi, green) of the indicated genotypes. Mean ± SD, n=3, data points with different symbols indicate independent biological replicates (Tukey's multiple comparisons test, a-c p≤0.0292, a-bc p=0.0484, ab-c p≤0.0312). H) Representative confocal images of epidermal cells of Col-0 and *fer-4* leaves stained with COS-488 labeling de-esterified pectin. Scale bar represents 50 µm. I) Quantification of COS-488 fluorescence intensity in Col-0 and *fer-4* leaves. Mean ± SD, n = 100–110 pooled from two independent experiments (Unpaired t-test, a-b p<0.0001). J) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of leaves pretreated with H_2O (blue) or EGCG (red). Mean ± SD, n=235-376 pooled from seven independent experiments (Mann-Whitney test, genotypes were analyzed separately, Col-0: a-b p<0.0001, *fer-4*: ns). K) Amount of fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection and pretreatment with H_2O (blue) or EGCG (red). n=3, data points with different symbols indicate independent biological replicates. All data is normalized to Col-0 (H_2O). L) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of

1094   the indicated genotypes. Mean ± SD, n=182-309 pooled from four independent experiments (Dunn's
1095   multiple comparisons test, a-b, b-c, a-c p<0.0001). All experiments were performed at least three times
1096   with similar results.

1097

Figure 5: RALF-mediated powdery mildew susceptibility is partially independent of FER and FER
signaling. A) Confocal images of pUBQ10::SYP122-pHusion lines in the Col-0 and CRISPR *fer*
background (2 dpi). Dotted lines indicate the location of the fungal spore. Scale bar represents 50 µm.
B) Quantification of the EGFP/mRFP1 ratio near the penetration site (green) and in uninfected areas
(grey). Mean ± SD, n=24 pooled from three independent experiments (Mann Whitney test, a-b
p<0.0001). C) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the
indicated genotypes. Mean ± SD, n=131-210 pooled from four independent experiments (Dunn's
multiple comparisons test, a-b p<0.0001). D) Conidiophores per field of view (5 dpi) of fungal colonies
grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=168-326 pooled from six
independent experiments (Dunn's multiple comparisons test, a-bc p=0.0469, a-cd p<0.0001, a-d
p<0.0001, ab-cc p≤0.0004, bc-dc p≤0.0019). E) PME activity normalized to Col-0. Mean ± SD, n=4, data
points with different symbols indicate independent biological replicates (Tukey's multiple comparisons
test, a-e p≤0.0003, a-d p≤0.0034, a-cd p=0.0058, ab-e≤0.0012, ab-d≤0.0138, ab-cd≤0.0227, a-
cbde≤0.0326). F) Conidiophores per field of view (5 dpi) after pretreatment with H₂O (blue) or 1µM
RALF23 before *Ecr* infection. Mean ± SD, n=485-970 pooled from 11 independent experiments (Mann
Whitney test, statistical analysis was performed on each genotype individually, Col-0: a-b p=0.0008, *fer-*
4: A-B p=0.0064). G) Quantification of the EGFP/mRFP1 ratio near the penetration site (green) and in
uninfected areas (grey). 1 µM synthetic RALF peptide or H₂O were infiltrated 2 h before infection. Mean
± SD, n=18 pooled from three independent experiments (Dunn's multiple comparisons test a-b
p≤0.0412, Mann Whitney test, **p=0.0064, ns=not significant). H) Conidiophores per field of view (5 dpi)
of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=95-137 pooled
from three independent experiments (Dunn's multiple comparisons test, a-b p=0.0076, a-c p<0.0001, b-
c p=0.0017). All experiments were performed at least three times with similar results.

1121

1122

Figure 6: Model of RALF-FER-dependent effects on powdery mildew sporulation. RALF peptides bind to negatively charged pectins and LRXs in the cell wall, as well as the FER-LLG1 complex at the plasma membrane. RALF binding to FER-LLG1 results in inhibition of AHA1/2 activity and apoplastic pH changes. The apoplastic pH modulates PMEI/PME activity and with this likely the methyl-esterification status of cell wall pectins. Both PME activity (1) and apoplastic pH modulation (2) affect powdery mildew sporulation. RALF-peptides appear to affect sporulation in a FER-independent (3) and FER-dependent (4) manner, potentially through regulating cell wall structure and the apoplastic pH. CW: cell wall; PM: Plasma membrane, EHM: Extrahaustorial membrane. Created with BioRender.com.

1131

1132

1133

1134 **Figure S1: Quantification of fungal growth on mutants of RALF binding proteins.** A-C) Amount of
1135 fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection of the indicated genotypes. Mean ± SD,
1136 data points indicate independent biological replicates. A) n=14, B) n=7–9, C) n=3. C) Data points with
1137 different symbols indicate independent biological replicates. All experiments were done at least three
1138 times with similar results.

1139

Figure S2: Characterization of CRISPR *ralf* mutants. A) Expression levels of the indicated *RALF*
genes in the vegetative rosette. Data was obtained using ePlant browser (https://bar.utoronto.ca/eplant/)
and is based on (Schmid *et al.*, 2005). B) Characterization of CRISPR *ralf3x*. Schematic diagram of
RALF22, RALF23 and *RALF33* gene structure and the CRISPR-Cas9-mediated mutation pattern
detected by DNA sequencing. The mature RALF peptide is indicated in blue. C) Characterization of
CRISPR *ralf7x*. Schematic diagram of *RALF1, RALF18, RALF22, RALF23*, *RALF31, RALF33* and
RALF34 gene structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA
sequencing. The mature RALF peptide is indicated in blue. D) Quantification of the petiole length of 4-
week-old Col-0, *fer-4*, CRISPR *ralf3x* and CRISPR *ralf7x* plants. Each data point represents the average
length of petioles on a fully grown rosette. Mean ± SD, n=7 pooled from two independent experiments
(Tukey's multiple comparisons test, a-b p<0.0001). E) Amount of fungal DNA normalized to plant DNA
(5 dpi) upon *Ecr* infection of the indicated genotypes. Data points with different symbols indicate
independent biological replicates n=3-5, F-G) RT-qPCR of *RALF23* in adult leaves of the indicated plant
lines. Housekeeping gene *UBQ5,* n=2. Data points with different symbols indicate independent
biological replicates. D-E) were performed at least two times with identical results. F–G) were performed
twice with identical results.

1156

1157

Figure S3: Regulation of RALF peptides during powdery mildew infection. RT-qPCR of the indicated *RALF* genes in Col-0 (blue) and *fer-4* (red) at 0 dpi (circle), 1 dpi (square), 4 dpi (triangle), 5 dpi (diamond). Housekeeping gene *UBQ5*. Mean ± SD, n=4-5, data points indicate independent biological replicates. (Dunn's multiple comparison test against 0 dpi within the respective genotype, p-values < 0.05 are indicated in the graph, all other comparisons were not significant.

1163

1164

Figure S4: FER- and RALF-mediated powdery mildew susceptibility is most likely not directly linked to MLO. A) Confocal microscopy images of fungal spores penetrating p35S::MLO2-mCherry and pFER::FER-GFP lines (2 dpi). Dotted lines indicate fungal spores, white arrow marks the penetration site. Scale bar represents 50 µm. B) Quantification of hyphal branching (2 dpi). Mean ± SD, n=77–98 pooled from two independent experiments (Mann Whitney test, ns: not significant). C) Penetration efficiency of *Ecr* on the indicated genotypes (calculated as the percentage of successful penetrations from all counted interactions, 2 dpi). Mean ± SD, n=2–4. D) Confocal microscopy images of fungal spores at 1 dpi. Callose depositions were stained using methyl blue (turquoise). Fungal structures were stained with propidium iodide (red). Scale bar represents 50 µm. E) Quantification of methyl blue fluorescence around the fungal penetration site, indicating the amount of callose deposition. Mean ± SD, n=43-45 pooled from three independent experiments (Dunn's multiple comparisons test). F) Conidiophores per field of view (5 dpi) of fungal colonies grown on Col-0 plants. Spores from the indicated genotypes were used for infection. Mean ± SD, n=218-336 pooled from three independent experiments (Dunn's multiple comparisons test). G) Quantification of trichome numbers per leaf area on the indicated genotypes. Mean ± SD, n=27 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p<0.0001). H) Microscopic images of aniline blue-stained trichomes of Col-0, *fer-4* and CRISPR *ralf7x*. Scale bar in the upper row represents 50 µm, scale bar in the lower panel represents 500 µm. I) Quantification of trichome branching on the indicated genotypes. Mean ± SD, n=27 pooled from three independent experiments (Dunn's multiple comparisons test, statistical analysis was performed on each category individually, 2 branches: a-b p=0.0002; 3 branches: a-b p=0.0259; a-c, b-c p<0.0001; 4 branches: a-b p<0.0001). All experiments were performed three times with similar results, except B), which was performed twice with identical results.

1187

1188

Figure S5: Characterization of *fer-4* CRISPR *myc2*. A) Quantification of glycosylated (green) 2,3-DHBA in untreated and *Ecr* infected (5 dpi) leaves of the indicated genotypes. Mean ± SD, n=3 pooled from three independent experiments (Tukey's multiple comparisons test, glycosylated: a-b p≤0.0324). B) Fungal spores/mg fresh weight (FW) of *Hyaloperonospora arabidopsidis* (*Hpa*) grown on the indicted genotypes. Mean ± SD, n=19-23 pooled from three independent experiments (Dunn's multiple comparison test, a-b p≤0.0129). All experiments were repeated at least three times with similar results. C) Characterization of *fer-4* CRISPR *myc2*. Schematic diagram of *MYC2* gene structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. D) RT-qPCR of *MYC2* in untreated (grey) adult leaves and upon infection with *Ecr* (5 dpi, green). Housekeeping gene *UBQ5*. Mean ± SD, n=3, data points with different symbols indicate independent biological replicates (Tukey's multiple comparisons test, a-b p≤0.0358). E) Images of 4-week-old Col-0, *fer-4* and *fer-4* CRISPR *myc2* plants. F) Amount of fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection of the indicated genotypes, n=3. Data points with different symbols indicate independent biological replicates. A, B, D and F were performed three times with similar results.

1203

1204

Figure S6: FER interactors detected by CoIP-MS. A) Volcano plot of detected FER-GFP interactors. Significantly enriched proteins (FDR=0.05 and fold change (FER-GFP/Lti6b-GFP) >2) are marked with colored circles according to the legend. Non-significant proteins are colored grey. B) List of significant FER-GFP interactors compared to Lti6b-GFP control. Data was obtained from biological triplicates for FER-GFP and biological quadruplets for Lti6b-GFP.

1210

A

B

Identifier	Molecular function	FDR	log(10)FC
AT1G32900.1 |GBSS1	Glycosyltransferase	0,0062	-1,223
AT2G36830.1 |TIP1;1	Water channel	0,0007	-0,459
AT3G59770.2 |SAC9	Phosphatase activity	0,0114	0,425
AT4G02030.1 |UNH	Protein binding	0,0062	0,668
AT5G35620.3 |EIF4E2	Translation initiation factor	0,0150	-0,354
AT3G55620.1 |eIF6A	Translation initiation factor	0,0142	0,330

1211

1212 **Figure S7: Powdery mildew dependent FER interactors detected by CoIP-MS.** A) Volcano plot of
1213 detected FER-GFP interactors after *Ecr* infection (1 dpi). Significantly enriched proteins (FDR=0.05 and
1214 fold change (*Ecr* infection (1dpi)/Untreated) >2) are marked in black. Non-significant proteins are colored
1215 grey. B) List of significant FER-GFP interactors at 1 dpi with *Ecr*, compared to the untreated FER-GFP
1216 control. Data was obtained from biological triplicates for each treatment.

1217

1218

Figure S8: Role of apoplastic pH and PMEs for *Ecr* infection success. A) Quantification of the band
intensities detected in western blot probed with α-AHA and α-Thr^{947P} antibodies. Band intensities were
measured using Image J software. Data points indicate individual experiments. Data points from the
same experiment are marked with the same color. B) AHA phosphorylation upon infection with *Ecr* (3
dpi). Western blots were probed with α-AHA and α-Thr^{947P} antibodies. CBB = Coomassie brilliant blue.
C) Conidiophores per field of view (5 dpi) after pretreatment with H_2O (blue), $MgCl_2$ 10 mM pH 7 (grey),
HEPES buffer 10 mM pH 7.5 (red) before *Ecr* infection. Mean ± SD, n=53-153 pooled from three
independent experiments (Dunn's multiple comparison, a-b, a-dc, a-bc, a-cd, a-d, b-d p<0.0001, b-cd
p=0.0245). D) Confocal images of fungal structures stained with COS-488. Scale bar represents 50 μm
E) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated
genotypes. Mean ± SD, n=161-338 pooled from three independent experiments (Dunn's multiple
comparisons test, a-b p<0.0001).   All experiments were repeated at least three times with similar results.

1231

1232

Figure S9: Analysis of FER-independent RALF23 functions during *Ecr* infection. A)
Characterization of pUBQ10::SYP122-pHusion CRISPR *fer-4*. Schematic diagram of *FERONIA* gene
structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. B)
Conidiophores per field of view (5 dpi) after pretreatment with H_2O (blue), 100 nM RALF23 (turquoise),
or 10 nM RALF23 (green). Mean ± SD, n=210-259 pooled from four independent experiments (Dunn's
multiple comparisons test, genotypes were analyzed separately. C) Quantification of the EGFP/mRFP1
3 h post infiltration of 1 µM RALF23 in H_2O or MES pH 5.5. Mean ± SD, n=21-24 pooled from three
independent experiments (Dunn's multiple comparisons test a-b≤0.009, Mann Whitney test, **p=0.0017,
ns=not significant). D) PME activity normalized to Col-0. Mean ± SD, n=4-5, data points with different
symbols indicate independent biological replicates (Tukey's multiple comparisons test, not significant).
(E) RT-qPCR of *RALF23* in untreated adult leaves of the indicated plant lines. Housekeeping gene
UBQ5. n=3, data points with different symbols indicate independent biological replicates. B-E) All
experiments were performed at least three times with similar results.

1246

1247

Figure S10: Cell wall localization of mCherry-RALF23 peptide variants. A) Representative z-stack images of epidermal cell expressing different variants of pR23::mCherry-RALF23. Panel 1: mCherry-RALF23 localization in untreated epidermal cells, panel 2-4: Epidermal cells after mannitol treatment (0.5 M), panel 3 and 4 show the image section marked in panel 2. Scale bar represents 50 µm. cw: Cell wall, pm: Plasma membrane.

1253

1254 **Figure S11: Images of plant lines used in this study.** Representative images of 5-week-old plants
1255 used in this study.

1256 **Tables**

1257 **Table S1: Identification of RALF-like peptides from phytopathogenic fungi species reported to**
1258 **be able to infect Arabidopsis.**

Species	Significant hits in predicted proteins (E-values < 10e^{-05})	Significant hits in translated genome assembly (E-values < 10e^{-05})	GenBank assembly accession number	Reference
Fusarium oxysporum NRRL 32931	1	2	GCA_000271745.2	(DeIulio *et al.*, 2018)
Pyrenophora tritici-repentis Pt-1C-BFP	1	1	GCA_000149985.1	(Manning *et al.*, 2013)
Golovinomyces cichoracearum[a] UMSG1	0	0	GCA_003611235.1	(Wu *et al.*, 2018)
Erysiphe neolycopersicum[a] UMSG2	0	0	GCA_003610855	(Wu *et al.*, 2018)
Erysiphe necator EnFRAME01	0	0	GCA_024703715.1	(Zaccaron *et al.*, 2023)
Blumeria graminis f.sp. *tritici* CHE_96224	0	0	GCA_900519115.1	(Müller *et al.*, 2019)
Podosphaera xanthii YZU573	0	0	GCA_028751805.1	(Xu & Chen, 2023)

1259

1260

Table S2: Plant lines used in this study.

Plant line	Genetic Background	Reference
fer-4	Col-0	(Duan *et al.*, 2010)
fer-2	Col-0	(Deslauriers & Larsen, 2010)
pFER::FER-GFP	*fer-4*	(Duan *et al.*, 2010)
llg1-2	Col-0	(Li *et al.*, 2015)
llg1-3	Col-0	(Shen *et al.*, 2017)
lrx1/2/3	Col-0	(Draeger *et al.*, 2015)
lrx1/2/3/4/5	Col-0	(Herger *et al.*, 2020)
p35S::LRR4-citrine	Col-0	(Fabrice *et al.*, 2018)
ralf1/22/23/33	Col-0	(Lan *et al.*, 2023)
CRISPR *ralf3x*	Col-0	Generated in this study
CRISPR *ralf7x*	Col-0	Generated in this study
p35S::RALF23	CRISPR *ralf7x*	Generated in this study
p35S::RALF23$^{Y93A/Y96A}$	CRISPR *ralf7x*	Generated in this study
p35S::RALF23$^{Y113A/Y114A}$	CRISPR *ralf7x*	Generated in this study
pR23::mCherry-RALF23	CRISPR *ralf7x*	Generated in this study
pR23::mCherry-RALF23$^{Y93A/Y96A}$	CRISPR *ralf7x*	Generated in this study
pR23::mCherry-RALF23$^{Y113A/Y114A}$	CRISPR *ralf7x*	Generated in this study
mlo2/6/12	Col-0	(Consonni *et al.*, 2006)
CRISPR *myc2*	*fer-4*	Generated in this study
PMEI3-OE	Col-0	(Peaucelle *et al.*, 2008)
ost2-2D	Col-0	(Merlot *et al.*, 2007)
pme3	Col-0	(Guénin *et al.*, 2011)
pme17	Col-0	(Del Corpo *et al.*, 2020)
pUBQ10::SYP122-pHusion	Col-0	(Kesten *et al.*, 2019)
pUBQ10::SYP122-pHusion/CRISPR *fer*	pUBQ10::SYP122-pHusion	Generated in this study
pFER:FER-GFP	*fer-4*	(Chakravorty *et al.*, 2018)
pFER:FER^{K565R}-GFP	*fer-4*	(Chakravorty *et al.*, 2018)
p35S::RALF23-GFP	Col-0	(Dobón *et al.*, 2015)
fer-2 x p35S::RALF23-GFP	*fer-2*	(Stegmann *et al.*, 2017)
pFER::FER^{WT}-GFP	*fer-4*	(Xiao *et al.*, 2019)
pFER::FER^{N303Y}-GFP	*fer-4*	(Xiao *et al.*, 2019)
pFER::FER^{G257A}-GFP	*fer-4*	(Xiao *et al.*, 2019)
p35S::Lti6b-GFP	Col-0	(Kadota *et al.*, 2014)

1264 **Table S3: Primers used in this study.**

Primers used for cloning		
Name	**Sequence (5' - 3')**	**Purpose**
RALF23_Esp3I_F	ttCGTCTCaaatgagaggactctccagaaac	p35S::RALF23
RALF23_Esp3I_R	ttCGTCTCaactatgagcgccggcagcgag	
$R23^{Y113AY114A}$_F	ttCGTCTCctgccgccaattgtcgacgtg	p35S::RALF23$^{Y113A/Y1114A}$
$R23^{Y113AY114A}$_R	ttCGTCTCcggcagatgcgccgcgacgtg	
$R23^{Y93AY96A}$_F	ttCGTCTCatcagcgctggtgcgctgaggag	p35S::RALF23$^{Y93A/Y96A}$
$R23^{Y93AY96A}$_R	ttCGTCTCgctgatgtacctcctcgtagc	
MLO2_Esp3I_F	ttCGTCTCaaatggcagatcaagtaaaag	p35S::MLO2-mCherry
MLO2_Blunt_R	tttcttaaaagaaaaatctcttaa (5'Phosphorylated)	
pR23_BpiI_F	ttGAAGACaacgagtggagaacctacgatttc	pR23::mCherryRALF23
pR23_BpiI_R	ttGAAGACaacattttcttctgtacactgtag	
Primers used for RT-qPCR		
Name	**Sequence (5' - 3')**	**Purpose**
Ath_Rubisco_F	TGATGGACGGTACTGGACAA	Quantification of plant DNA
Ath_Rubsico_R	GAAGCTTGGTGGCTTGTAGG	
Ecr_Tubulin_F	TGACAGCCCGGAATGAGT	Quantification of fungal DNA
Ecr_Tubulin_R	TTGTCTTCGTTTCCCAGGTC	
RALF23_qPCR_F	TTCATAACTGGTCCGTCGCC	RALF23
RALF23_qPCR_R	AGCTAATATGCGCCGGTTGA	
RALF1_qPCR_F	CATCTCTTCACCTCCGGTCC	RALF1
RALF1_qPCR_R	CTATTGAACCGTGGCAGCCT	
RALF18_qPCR_F	GTCGGATCTCGTCCGGTAAA	RALF18
RALF18_qPCR_R	CGGTTTACCATCTGGTTTGGC	
RALF22_qPCR_F	GCGGAAGAAGAAGAGATGGAG	RALF22
RALF22_qPCR_R	TTGTAGTACGACGCACCAC	
RALF31_qPCR_F	ACGGAGAGATCGACGCAATG	RALF31
RALF31_qPCR_R	CCATCAGTACTCTCCGGCTG	
RALF33_qPCR_F	CTCACCGTCCACTTCTTATTC	RALF33
RALF33_qPCR_R	CGTTGTTGCTAAAATACGCC	
RALF34_qPCR_F	CTCCTCCTCATTCTCTCTCTTC	RALF34
RALF34_qPCR_R	TCGTCTTCCTCCGTAACTTC	
UBQ5_qPCR_F	ACTCCTTCCTCAAACGCTGA	UBQ5
UBQ5_qPCR_R	CCAAGCCGAAGAAGATCAAG	
MYC2_qPCR_F	GTGCGGGATTAGCTGGTAAA	MYC2
MYC2_qPCR_R	ATGCATCCCAAACACTCCTC	
PR1_qPCR_F	CGGAGCTACGCAGAACAACT	PR1
PR1_qPCR_R	CAGACAAGTCACCGCTACCC	
JAZ10_qPCR_F	GAGAAGCGCAAGGAGAGATTAG	JAZ10
JAZ10_qPCR_R	CTTAGTAGGTAACGTAATCTCC	
PDF1.2_qPCR_F	CTG CTT TCG ACG CAC CGG CA	PDF1.2
PDF1.2_qPCR_R	TGA TCC ATG TTT GGC TCC TTC	
Primers used for genotyping		
Name	**Sequence (5' - 3')**	**Purpose**
GG pUC18 Seq-F	GTATCACGAGGCCCTTTCGT	Inserts in GoldenGate pUC18
GG pUC18 Seq-R	TAATGAATCGGCCAACGC	
pGGPlant-XL-AB-insert_Fwd	AAACGACAATCTGAGCTCCACC	Insert in GoldenGate pGGPlantXL-AB
pGGPlantXL-AB-insert_Rev	CGGTGTCATCTATGTTACTAGAT	
RALF1_F	CAAAATTAAAAAATCGAGTCCAC	CRISPR RALF1
RALF1_R	GCTTGATGGATCAAAGAAAAC	
RALF14_F	GCTCTTAATCTTCGCCG	CRISPR RALF14
RALF14_R	GTAAACCGGTAACAATGTG	
RALF18_F	CATCACAAACCATTTCAACG	CRISPR RALF18
RALF18_R	CACACCAACGAAAGGAAG	
RALF22_F	GTTTCCAAGAGAGAAAGAGAG	CRISPR RALF22
RALF22_R	GAAAATCGAGAAGAGAGACAG	
RALF23_F	CTTCATAGTTCGTGCACAG	CRISPR RALF23
RALF23_R	CTAATAATCCGCAATGAACTTAG	
RALF31_F	CTTCTTCTGCTTAGAGACAG	CRISPR RALF31
RALF31_R	CAGCAATGCAAAAGCTTAATC	
RALF33_F	CCTCAGAGAAAGAAGAAGAAAG	CRISPR RALF33

RALF33_R	GCAATGCAAAGTATAAACATAG	
RALF34_F	CCTCTTTACATGTGACTTCC	CRISPR RALF34
RALF34_R	CACGGTTTACAAACAAACATC	
MYC2_F	GGAACACACGAAGGTTGGAC	CRISPR MYC2
MYC2_R	CTGCTACTTCTTTCACGACGG	
FER_F	CGTAACCAAACGAACATG	CRISPR FER
FER_R	GGTAAGTGAAACCAGAGTC	

1265

1266

1267 **Table S4: CRISPR target sites.**

Target site	Sequence	Purpose
RALF1.1	ACACTCTGCTATTGAACCG	Generation of CRISPR *ralf* mutants
RALF1.2	AATTGTCAGAACGGAGCTC	
RALF14.1	GATGCATTTTCCACTCAAC	
RALF18.1	GTGGTTGCAGTGCCGCTAC	
RALF22.1	GTTCGATTCAGACATAAGC	
RALF22.2	CGTACTACAACTGCCAGCG	
RALF23.1	CCTCGTAGCTAATATGCGC	
RALF23.2	CAGCCACGAGAGTAAGGAT	
RALF31.1	TGGCGCAGAAACGGTACAT	
RALF31.2	TTTGTGTCTCTAGCGCATC	
RALF33.1	CGATCGAGTCGAAATGCAA	
RALF33.2	TAGCGCTATTACTCGTTGC	
RALF34.1	CTCTCACTCTTCTACCCGA	
RALF34.2	CCGTTCATCCGTATAGCCG	
MYC2.1	CCGCCGGAGTAGAAAACGG	Generation of *fer*-4 CRISPR *myc2*
FER.1	GACGATTAAGTATCCTACG	Generation of pUBQ10::SYP122-apopHusion CRISPR *fer*-4
FER.2	TGATCGATGAAGATCACAG	

1268