Endogenous RALF peptide function is required for powdery mildew host colonization

Henriette Leicher[1], Sebastian Schade[1,#], Jan W. Huebbers[2], Kristina S. Munzert-Eberlein[3], Genc Haljiti[4],

Christina Ludwig[4], Marion Müller[1], Toshinori Kinoshita[5,6], Timo Engelsdorf[3], Ralph Hückelhoven[1] and

Martin Stegmann[1,#]

[1]Chair of Phytopathology, TUM School of Life Sciences, Technical University of Munich, Freising, Germany
[2]Unit of Plant Molecular Cell Biology, Institute for Biology I, RWTH Aachen University, Aachen, Germany
[3]Molecular Plant Physiology, Department of Biology, Philipps-Universität Marburg, Marburg, Germany
[4]Bavarian Center for Biomolecular Mass Spectrometry (BayBioMS), TUM School of Life Science, Technical
University of Munich (TUM), Freising, Germany
[5]Graduate School of Science, Nagoya University, Chikusa, Nagoya, Japan
[6]Institute of Transformative Bio-Molecules (WPI-ITbM), Nagoya University, Chikusa, Nagoya, Japan
[#]present address: Institute of Botany, Molecular Botany, Ulm University, Germany
13
14 *Corresponding author: Email: martin.stegmann@tum.de
15

Abstract

The receptor kinase FERONIA (FER) is a susceptibility factor for biotrophic powdery mildew fungal
pathogens in *Arabidopsis thaliana*, but the underlying molecular mechanisms remain largely unknown.
FER is required for the perception of endogenous RAPID ALKALINIZATION FACTOR (RALF) peptides
to control various aspects of plant growth, development and immunity. RALFs are perceived by FER-
LORELEI-LIKE GPI-ANCHORED PROTEIN (LLG) heterocomplexes to induce cellular responses and
bind to LEUCINE-RICH REPEAT EXTENSIN (LRX) proteins as structural components of the cell wall.
We show that FER`s RALF ligands are necessary for full success of powdery mildew colonization. Our
data reveals that LLGs and LRXs are also powdery mildew susceptibility factors, and that the
susceptibility function of RALFs is best explained by a combination of peptide signalling and cell wall-
related effects. We further reveal that cell wall and apoplastic pH homeostasis, hallmark features of
RALF function, support powdery mildew reproductive success. The function of host RALF peptides is
underlined by the fact that, in contrast to previously described pathogens that hijack FER for
colonization, powdery mildew fungi do not produce functional RALF mimics. Surprisingly, RALF-
dependent powdery mildew pathogenesis is partially independent of FER. The resistance phenotype of
ralf and *fer* mutants is not associated with de-regulated salicylic acid and jasmonic acid signalling,
important defense-related phytohormones. Collectively, powdery mildew fungi require RALF signalling
for apoplastic pH regulation and cell wall structural organization for successful colonization, highlighting
a novel mechanism for host susceptibility by obligate biotrophic fungi.

Key words: RALF peptides, powdery mildew, FER, peptide signalling, cell wall, pH homeostasis

Introduction

Plants have evolved a multi-layered immune system to defend against potential pathogens. Pattern-triggered immunity (PTI) is activated through the perception of conserved microbe-derived molecular signatures (microbe-associated molecular patterns, MAMPs) or host derived damage-associated molecular patterns (DAMPs) by plasma membrane localized pattern recognition receptors (PRRs), which are either receptor-like kinases (RLKs) or receptor-like proteins (RLPs). To subvert PTI, host adapted pathogens evolved effectors that again can be perceived by specific resistance (R) proteins (encoded by R-genes) including intracellular localized nucleotide-binding leucine-rich repeat receptors (NB-LRRs). Effector recognition activates a strong second layer of immunity culminating in the hypersensitive response (HR) involving cell death to inhibit proliferation of biotrophic pathogens (Ngou et al., 2022) .

Other genetic factors affecting infection success are susceptibility genes (S-genes). These encode plant factors with various endogenous functions that can be utilized by pathogens for their own benefit. In consequence, mutating S-genes usually confers pathogen genotype-independent and hence durable disease resistance (Engelhardt et al., 2018; van Schie and Takken, 2014).

With over 400 different species that infect nearly 10 000 possible host plants, powdery mildew is one of the most widespread fungal plant diseases, also affecting a variety of economically relevant crop species (Glawe, 2008). During the powdery mildew infection cycle, a continuous interaction occurs between the obligate biotrophic fungus and its host plant, which determines the infection success at multiple levels. The first well-characterized powdery mildew susceptibility factor is the barley MILDEW LOCUS O (MLO) protein, that confers barley susceptibility towards adapted *Blumeria hordei*. The *mlo* mutants show race-unspecific resistance against powdery mildew even after decades of field application. Mutants in *mlo* are powdery mildew resistant in phylogenetically diverse plant species, including *Arabidopsis* and tomato, suggesting that *S*-genes are functionally conserved (Kusch and Panstruga, 2017).

In addition to their function as PRRs, RLKs are important sensors for endogenous signals regulating growth and/or modulating immunity (Rzemieniewski and Stegmann, 2022). Adapted pathogens can hijack these RLK pathways to facilitate successful host accommodation. In *Arabidopsis*, IMPAIRED OOMYCETE SUSCEPTIBILITY 1 (IOS1) and SYMRK-HOMOLOGOUES RKs (ShRKs) are susceptibility factors for *Hyaloperonospora arabidopsidis* (*Hpa*) infection but they show mild or no susceptibility function in interaction with the powdery mildew fungus *Erysiphe cruciferarum* (Hok et al., 2014; Hok et al., 2011; Ried et al., 2019). Silencing of specific powdery mildew infection-regulated RLKs enhances penetration resistance to *Blumeria hordei* (*Bh*) in single cell transformation experiments (Douchkov et al., 2014; Schnepf et al., 2018).

The RLK FERONIA (FER) is an important regulator of diverse physiological responses. FER and related CATHARANTUS ROSEUS RECEPTOR-LIKE KINASE 1-LIKEs (CrRLK1Ls) perceive endogenous RAPID ALKALINIZATION FACTOR (RALF) peptides to regulate multiple plant processes, including pollen hydration, pollen tube growth, polytubey block, intergeneric hybridization and root hair growth (Ge et al., 2017; Gonneau et al., 2018; Lan et al., 2023; Zhong et al., 2022; Zhu et al., 2021). FER contributes to PTI by acting as a RALF-regulated scaffold protein for PRR complexes (Stegmann et al., 2017). FER perceives RALF peptides in a heteromeric complex with the GPI anchored protein LORELEI-LIKE GPI

ANCHORED PROTEIN 1 (LLG1) (Li et al., 2015; Xiao et al., 2019). In controlling antibacterial
resistance, FER´s role is associated with a function in jasmonic acid (JA) signalling. Here, FER
phosphorylates and destabilize the JA downstream transcription factor MYELOCYTOMATOSIS 2
(MYC2) to enhance resistance to *Pseudomonas syringae* pv. *tomato* (*Pto*) (Guo et al., 2018). RALF
peptides also bind to LRX proteins but the biochemical interplay between LRX and FER-LLG complexes
remains little understood (Dünser et al., 2019; Gronnier et al., 2022; Mecchia et al., 2017; Moussu et al.,
2020; Zhao et al., 2018). Recent reports show that RALF peptides (via binding to LRXs) have a structural
role in organizing the patterning of demethylesterified pectins in the cell wall. This suggests that pectin
may connect cell wall structural LRX-RALF complexes with plasma membrane-localized CrRLK1Ls that
in turn monitor cell wall integrity (Moussu et al., 2023; Schoenaers et al., 2024). The central role of FER-
LLG/LRX-RALF modules for plant physiology is highlighted by the fact that adapted pathogens can
hijack RALF signalling for their own benefit. *Fusarium oxysporum* and root knot nematodes produce a
RALF mimic to enhance infection success on tomato and *Arabidopsis* (Masachis et al., 2016; Thynne
et al., 2017; Zhang et al., 2020). This function is likely related to RALF-mediated immune suppression
and physiological reprogramming. Furthermore, the *fer-1* mutant in the Ler-0 *Arabidopsis* ecotype is less
susceptible to the powdery mildew disease caused by *Erysiphe orontii*, suggesting that FER acts as an
S-factor for powdery mildew (Kessler et al., 2010). Yet, how FER confers susceptibility remains little
understood.

Here, we sought out to unravel the molecular mechanism of FER-mediated powdery mildew
susceptibility. We confirmed FER`s function in powdery mildew colonization in the Col-0 *fer-4* mutant
using the adapted Arabidopsis-infecting powdery mildew species *Erysiphe cruciferarum*. We show that
other RALF binding and receptor complex-related proteins, including LLGs and LRX, are required for
successful *Ecr* colonization. By loss of function studies, we unravel RALF peptides as important
susceptibility factors for powdery mildew. CRISPR Cas9-generated *ralf22/23/33* (CRISPR *ralf3x*) and
*ralf1/18/22/23/31/33/34 (*CRISPR *ralf7x*) show increasing resistance to powdery mildew infection
depending on the number of eliminated RALFs. Our bioinformatic analysis suggests that powdery
mildew fungi do not encode functional RALF peptide mimics, supporting that mildew requires host-
endogenous RALF signalling for its own benefit (Masachis et al., 2016; Thynne et al., 2017). We suggest
that the enhanced resistance of *fer* and CRISPR *ralf7x* is not associated with defects in MLO-mediated
fungal penetration or deregulation of defense-related phytohormone signalling. Instead, powdery mildew
might require FER-RALF-mediated apoplastic pH regulation and cell wall remodelling for successful
colonization. Surprisingly, our data indicates that RALF-mediated powdery mildew susceptibility is only
partially dependent on FER and exerts its effect in a combination of signalling and structural functions.
We propose that powdery mildew fungi require RALFs to provide a favourable growth environment,
highlighting a novel mechanism of host subversion by adapted plant pathogens.

Results

FER, LLGs and LRX are powdery mildew susceptibility factors

The Ler-0 *fer-1* mutant was demonstrated to be more resistant to infection with the powdery mildew
species *Golovinomyces orontii* (Kessler et al., 2010). We confirmed this phenotype in the Arabidopsis

Col-0 ecotype using the well described *fer-4* mutant allele (Duan et al., 2010) and the powdery mildew

species *Erysiphe cruciferarum* (*Ecr*), which is also able to infect Arabidopsis (Micali et al., 2008).

Macroscopically the *fer-4* mutant showed slightly reduced visible disease symptoms compared to Col-0

(Fig. 1A). We quantified the resistance by counting conidiophores (the asexual reproductive structures

of this fungus) of individual colonies 5 days post *Ecr* inoculation. The *fer-4* mutant showed strongly

reduced conidiophore production compared to the WT control (Fig. 1B, C). Further quantification by

qPCR of relative fungal DNA, normalized to plant DNA, demonstrated reduced fungal growth on *fer-4*

(Fig. S1A). Expression of FER-GFP under the control of its native promoter complemented the

resistance phenotype of *fer-4* (Fig. 1C). This data indicates that *fer* loss of function mutants, across

ecotypes and fungal species, are more resistant to powdery mildew infection.

Next, we investigated whether FER-related RALF receptor complex components are involved in

powdery mildew susceptibility. We tested a loss of function mutant in the RALF receptor LLG1 (*llg1-2*)

which showed reduced conidiophore production (Fig. 1D). A similar effect was observed in the *llg1-3*

allele, in which LLG1 is impaired in RALF binding but mutants do not show *fer*-like growth phenotypes

(Fig. 1D, Fig. S8) (Shen et al., 2017). The resistance phenotype of the *llg1* mutants, however, was not

clear in qPCR quantification (Fig. S1B). Since qPCR gives an average infection level over the whole

plant, this method is more prone to uncontrollable differences in infection densities and is less sensitive

to mild phenotypes, unlike conidiophore counting, which evaluates the reproductive success of

individual colonies. Surprisingly, *llg1-2* and *llg1-3* were both previously described to be more susceptible

to the powdery mildew species *Golonivomyces cichoracearum* (Shen et al., 2017), raising the question

of mildew-species-dependent differences in *llg1* pathophenotypes, unlike *fer* (Fig. 1) (Kessler et al.,

2010) and potential redundancy of LLG1 with related members of the LRE/LLG gene family (Noble et

al., 2022; Xiao et al., 2019). Both *LLG2* and *LLG3* show very weak predicted expression in leaves (Xiao

et al., 2019), which does not exclude a role in leaf-related immune responses, as previously described

for the CrRLK1L ANXUR1 (Mang et al., 2017). In addition to the plasma membrane localized FER-LLG1

receptor complex, RALF peptides bind to cell wall localized LRX proteins with high affinity (Moussu et

al., 2020). A *lrx3 lrx4 lrx5* (*lrx3/4/5*) triple mutant phenocopies *fer* in immune signalling, salt stress

responses and overall plant morphology (Dünser et al., 2019; Gronnier et al., 2022; Moussu et al., 2020;

Zhao et al., 2018). This *lrx3/4/5* mutant displayed reduced conidiophore production upon *Ecr* infection

(Fig. 1E). Yet, the mutant did not show reduced relative fungal DNA content (Fig. S1C). Further mutating

LRX1 and *LRX2* in *lrx3/4/5* (*lrx1/2/3/4/5*, hereafter referred to as *lrx5x* mutant) (Herger et al., 2020)

promoted the enhanced resistance phenotype (Fig. 1E, Fig. S1C). Altered plant morphology and PTI

responses of *lrx3/4/5* are indistinguishable from those in *fer-4* (Gronnier et al., 2022; Zhao et al., 2018),

suggesting that LRXs display an increased level of redundancy in the context of powdery mildew

susceptibility control. Overexpression of LRR4-citrine (LRX4 lacking the extensin domain) has a

dominant negative effect on plant growth, while PTI responses are not affected (Dünser et al., 2019;

Gronnier et al., 2022). The p35S::LRR4-citrine lines phenocopied *lrx5x* mutants and *fer-4* in showing

enhanced powdery mildew resistance (Fig. 1F).

Powdery mildew requires endogenous RALF peptides for successful colonization

Given the involvement of FER-LLG1 and LRXs in the establishment of powdery mildew pathogenesis
on Col-0, we hypothesized that RALF peptide signalling through FER-LLG1 and/or their binding to LRXs
is required for powdery mildew colonization. Many plant pathogens, including fungi, secrete functional
RALF peptide mimics to suppress immunity or reprogram host physiology (Masachis et al., 2016;
Thynne et al., 2017; Zhang et al., 2020). We searched available genomes of powdery mildew species
for the presence of potential RALF peptide mimics. However, consistent with previous reports (Thynne
et al., 2017) we could not detect evidence for RALF peptide mimics among different species of the order
Erysiphales (Table S1, method section for details on our bioinformatic search pipeline). Based on this,
we hypothesized that endogenous RALFs are involved in the *Ecr*-Arabidopsis interaction. To test this
hypothesis, we generated CRISPR mutants of leaf-expressed RALF peptides. The Arabidopsis genome
encodes for at least 37 RALF members (Abarca et al., 2021). We anticipated genetic redundancy and
generated CRISPR mutants in which we targeted *RALF23*, a well-described ligand of FER-LLG1
(Stegmann et al., 2017; Tang et al., 2022; Xiao et al., 2019) and seven additional RALF-peptides with
structural similarity to RALF23 and predicted leaf expression (*RALF1*, *RALF14*, *RALF18*, *RALF22*,
RALF33, *RALF23*, *RALF31*, *RALF34*) (Fig. S2A). We could not obtain an octuple mutant, raising the
question whether loss of multiple RALFs may cause severe developmental defects and/or lethality.
Nevertheless, two higher order mutants, a knockout of *ralf22/23/33* (hereafter CRISPR *ralf3x*) and a
ralf1/18/22/23/31/33/34 mutant (hereafter CRISPR *ralf7x*) (Fig. 2A, Fig. S2B, C), were generated. While
the CRISPR *ralf3x* mutant only displayed a mild growth phenotype with slightly shortened petioles, the
CRISPR *ralf7x* mutant was strongly impaired in shoot growth (Fig. 2B, Fig. S2D, S8). Compared to *fer-
4*, however, CRISPR *ralf7x* showed less pronounced shoot growth defects with a trend towards longer
petioles (Fig. S2D) and an overall more wildtype-like morphology (Fig. 2B, Fig. S8). Interestingly, the
powdery mildew infection success was significantly reduced on CRISPR *ralf3x* and CRISPR *ralf7x*,
demonstrated by both a reduced number of conidiophores and decreased relative fungal DNA contents
(Fig. 2B-E). Importantly, CRISPR *ralf7x* displayed elevated resistance compared to *fer-4* (Fig. 2D, E).
Expression of p35S::RALF23 complemented the CRISPR *ralf7x* growth and powdery mildew infection
phenotype (Fig. 2F, Fig. S2E, S8). We also observed a strong powdery mildew resistance phenotype
with the previously reported *ralf1/22/23/31* quadruple mutant (Fig. 2G) (Lan et al., 2023). Collectively,
this data show that multiple FER-LLG1 and LRX-binding RALFs genetically contribute to powdery
mildew pathogenesis.

FER and RALF mutants show post-penetration powdery mildew resistance

We next sought out to unravel the cellular mechanism underlying RALF-mediated powdery mildew
susceptibility. Recent evidence showed that RALF-induced calcium influx in synergid cells and pollen
tubes is facilitated by FER-LRE-mediated activation of the NTA/MLO7 calcium channel (Gao et al.,
2022). The MLO family encodes for important powdery mildew susceptibility factors across different
families (Acevedo-Garcia et al., 2017; Büschges et al., 1997; Engelhardt et al., 2018; Li et al., 2022b;
Nekrasov et al., 2017; van Schie and Takken, 2014). In Arabidopsis, mutation of three partially
redundant MLOs is required to confer full penetration resistance (Consonni et al., 2006). Genetically,
MLO2 is the main susceptibility factor and active as a calcium channel in COS7 cell culture (Consonni

et al., 2006; Gao et al., 2022). This raises the possibility that RALF perception by FER-LLG complexes
activates MLO channels to facilitate mildew colonization.

MLO-YFP in barley and MLO2-GFP in Arabidopsis focally accumulate at powdery mildew penetration
sites (Bhat et al., 2005; Huebbers et al., 2024). MLO2 localization to the fungal penetration site was
confirmed with p35S::MLO2-mCherry lines (Fig. 3A, left panel). Further, we tested whether FER-GFP
accumulates at *Ecr* penetration sites using *fer-4 pFER::FER-GFP* (Duan et al., 2010). The FER-GFP
signal outlined the plasma membrane, but specific recruitment to the site of fungal attack, haustorial
neck and/or extrahaustorial membrane could not be detected (Fig. 3A, right panel). The *fer-4* mutant
shows strongly reduced production of conidiophores on infected tissue (Fig. 1C), but initial hyphal growth
is largely unaffected (Fig. 3B). This prompted us to assess whether *fer-4* or CRISPR *ralf7x* mutants
show any defect in epidermal cell penetration. Host cell entry by *Ecr* was successful in ~80% of
penetration attempts on Col-0. Surprisingly, we did not observe any difference in epidermal cell entry in
fer-4 and CRISPR *ralf7x*. As previously reported (Consonni et al., 2006), the *mlo2/6/12* mutant showed
complete penetration resistance (Fig. 3C). This suggests that mutating RALF-related components
primarily affects reproductive success and not epiphytic growth, which may explain the overall milder
phenotypes we observed in qPCR-based quantification of *Ecr* infection success.

While callose deposition is mostly confined on Col-0 trichomes, the *mlo2/6/12* mutant displays aberrant
callose deposition in trichomes with delocalized or absent accumulation (Huebbers et al., 2024).
Likewise, *fer-4* mutants are impaired in trichome formation (Duan et al., 2010) which prompted us to test
whether *fer-4*, CRISPR *ralf7x* and *mlo2/6/12* show phenotypic similarities. We observed less trichomes
on *fer-4*, while the number of trichomes on CRISPR *ralf7x* was not significantly reduced (Fig. 3D).
Further, trichomes on *fer-4* tended to have fewer branches than trichomes on Col-0 while trichomes on
CRISPR *ralf7x* had more branches (Fig. 3E). Moreover, a fraction of trichomes on the CRISPR *ralf7x*
mutant were enlarged in the base region (≈ 22 %), which was not observed on Col-0 and *fer*-4 (Fig. 3F).
Unlike *mlo2/6/12*, neither CRISPR *ralf7x* nor *fer*-4 showed mis-regulated callose deposition in trichomes
(Fig. 3F)(Huebbers et al., 2024). Collectively, this data suggests that FER-RALF-dependent powdery
mildew susceptibility and trichome development is not directly linked to MLO function. Our data suggests
that mutation in FER-RALF primarily affects post-penetration fungal colonization.

The powdery mildew resistance phenotype of *fer-4* is not associated with de-regulated salicylic
acid and jasmonic acid signalling

Salicylic acid (SA) is an important defence hormone promoting resistance to biotrophic and
hemibiotrophic pathogens (Peng et al., 2021). For example, reduced powdery mildew colonization in
POWDERY MILDEW RESISTANCE 4 (*PMR4*) mutants is dependent on enhanced SA signalling and
expression of the SA hydrolysing enzyme NahG enhances powdery mildew colonization (Nishimura et
al., 2003). Ler-0 *fer-1* seedlings did not show elevated *PATHOGENESIS-RELATED 1* (*PR1*) expression,
a marker gene of the SA pathway (Kessler et al., 2010). Yet, the weak *fer-5* allele in Col-0 (Duan et al.,
2010) showed mildly elevated SA levels in seedlings (Engelsdorf et al., 2018), prompting us to test
whether the mildew resistance phenotype in *fer-4* may be associated with elevated SA levels. We
quantified free and glycosylated SA by high pressure liquid chromatography (HPLC) upon *Ecr* infection.

As expected *Ecr* infection led to mildly enhanced levels of SA (Fig. 4A). However, neither *fer-4* nor

CRISPR *ralf7x* showed higher SA levels in the untreated state or after infection (Fig. 4A). We also

measured the major catabolic SA products 2,3-DHBA and 2,5-DHBA to investigate the flux through the

SA pathway (Zhang et al., 2013; Zhang et al., 2017) as well as camalexin levels, an indole alkaloid

associated with anti-fungal resistance (Liu et al., 2016; Mucha et al., 2019). Neither of those compounds

were significantly enhanced before or after infection in *fer-4* and CRISPR *ralf7x* compared to the Col-0

control (Fig. 4B, C, Fig. S3A). In accordance with the measured SA levels, *PR1* expression was not

elevated in any of the tested mutants in control conditions. *Ecr* infection resulted in enhanced *PR1*

expression levels which was observed in all tested genotypes (Fig. 4D). This suggests that the *fer-4* and

CRISPR *ralf7x* resistance phenotype is likely independent of altered SA signalling.

Jasmonic acid (JA), in concert with ethylene, is usually implicated in defence against insect herbivory

and necrotrophic fungi (Bürger and Chory, 2019), but JA treatment can also inhibit colonization by the

powdery mildew fungus *Erysiphe cichoracearum* in Arabidopsis (Zimmerli et al., 2004). Importantly, FER

inhibits JA signalling by phosphorylating and destabilizing the central JA-associated transcription factor

MYC2 (Guo et al., 2018). To test whether FER`s susceptibility role is linked to MYC2-mediated JA

regulation, we generated a *fer-4 myc2* mutant via CRISPR-Cas9 in the *fer-4* background that fully

diminished MYC2 expression (Fig. S3B, C). Similar to previous reports, mutating *myc2* in *fer-4* partially

restored the *fer-4* growth phenotype (Fig. S3D) (Guo et al., 2018). Furthermore, the *fer-4* CRISPR *myc2*

mutant partially complemented the deregulation of JA-responsive genes in *fer-4*, a phenotype that was

also observed in CRISPR *ralf7x* and *lrx5x* (Fig. 4E, F, Fig. S3C). Importantly though, mutating *MYC2*

did not restore powdery mildew susceptibility in *fer-4* (Fig. 4G, Fig. S3E). This suggests that the

resistance phenotype of *fer-4* is independent of MYC2-mediated JA signalling.

Apoplastic pH modulations affect powdery mildew infection success

RALF peptides were first described as potent inducers of apoplastic alkalinisation (Pearce et al., 2010).

Similarly, powdery mildew infection of susceptible barley results in transient pH spiking with initial peaks

towards the alkaline (Felle et al., 2004). Infiltration of acidic buffers into barley leaves before inoculation

with *Blumeria hordei* increases the incidence of papilla formation, indicative of enhanced resistance

(Wiese et al., 2004). Since other fungal pathogens also modify the apoplastic pH environment for

successful infection (Kesten et al., 2019), we hypothesized that FER-RALF-induced pH modulations

affect powdery mildew infection success.

Perception of RALF by FER induces the phosphorylation of an inhibitory site on the major Arabidopsis

plasma membrane H^+ ATPase AHA1 (Haruta et al., 2014), resulting in its inhibition and apoplastic

alkalinisation. AHA1 is the strongest expressed AHA in leaf tissue (Haruta et al., 2010), but the FER-

dependent phosphorylation site is conserved in AHA1 and AHA2, the two major isoforms in Arabidopsis.

An *aha1* single mutant did not display colonization difference upon *Ecr* infection (Weis et al., 2013).

While this might be explained by genetic redundancy, *aha1 aha2* double mutants are seedling lethal,

making genetic analysis difficult (Haruta et al., 2010).

To determine potential protein interaction partners of FER, we performed a co-immunoprecipiation

experiment coupled with a mass spectrometric readout (CoIP-MS). For that we genetically tagged FER

with GFP and used a fusion of Lti6b-GFP as negative control, representing a membrane-anchored
protein. In this experiment, we could identify 68 potential protein interaction partners of FER, including
AHA1 and AHA3 (Fig. S4A, B). Compared to the Lti6b-GFP control sample, the mass spectrometric
intensity of AHA1 was 2,7-fold enriched (Fig. 5A), confirming previously published data on AHA1-FER
interaction (Du et al., 2016; Haruta et al., 2014). We also found AHA3 to be significantly enriched, since
AHA3 derived peptides were only detected in FER-GFP but not in Lti6b-GFP samples. AHA2 was found
to be 2,3 -fold enriched (Fig. 5A), but below the significance threshold (FDR<0.05).

The majority of the detected protein interaction partners of FER were either RLKs or proteins involved
in transmembrane transport (Fig. S4A, B). In particular, we identified four CrRLK1L members (HERK1,
MDS1/LET2, THE1 and LET1). We also detected peptides corresponding to the GPI-anchored protein
SHAVEN 3 (SHV3) and the closely related SHAVEN3-LIKE1 *(SVL1)*. Double mutants of *SHV3* and
SVL1 (*shv3 svl1*) phenocopy *fer* for increased apoplastic pH, reduced hypocotyl elongation dependent
on external sucrose supply, increased cellular sucrose uptake and sucrose-dependent alterations in cell
wall contents (Yeats et al., 2016). In response to *Ecr* infection, we could only detect six differential FER-
GFP interactors, which may be related to the early experimental time point chosen (1 dpi).

Phosphorylation of the C-terminal penultimate threonine residue in AHA1/AHA2 (AHA1^{T947}/AHA2^{T948})
activates AHA proton pump activity (Falhof et al., 2016). For example, auxin promotes
AHA1^{T947}/AHA2^{T948} phosphorylation in a TRANSMEMBRANE KINASE 1-dependent manner (Lin et al.,
2021). We tested whether *Ecr* infection affects AHA1^{T947}/AHA2^{T948} by making use of a phosphosite-
specific antibody. However, pT947/pT948 phosphorylation was not altered upon *Ecr* infection, neither
in Col-0 nor *in fer-4* (Fig. S6A, B). RALF perception by FER, including RALF1 and RALF23, promotes
phosphorylation of serine 899 (pS899) to inhibit AHA1/AHA2 proton pump activity (Haruta et al., 2014;
Liu et al., 2025). Importantly, pS899 does not correlate with reduced pT947/pT948 and a recent report
showed that RALF23/RALF33 treatment does not induce pT947/pT948 (Haruta et al., 2014; Liu et al.,
2025). Therefore, the results do not exclude a FER-dependent regulation of AHA activity and/or proton
flux across the plasma membrane correlated with altered *Ecr* infection success. Therefore, we decided
to test the *ost2-2D* mutant, which produces a constitutive active AHA1 (Merlot et al., 2007), for altered
powdery mildew susceptibility. The *ost2-2D* mutant displayed enhanced *Ecr* resistance (Fig. 5B). The
fungal toxin fusicoccin stabilizes 14-3-3 protein interaction with phosphorylation-activated AHA1/AHA2
to induce constitutive proton extrusion and can support powdery mildew resistance of barley
(Baunsgaard et al., 1998; Zhou et al., 2000). Treatment of plants with fusicoccin before *Ecr* inoculation
induced resistance in a partially FER-dependent manner (Fig. 5C, D). Next, we tested whether buffering
the apoplastic pH can create an environment that is favourable for the fungus. We hypothesized that
alkaline pH infiltration may promote susceptibility and/or break the *fer-4* resistance phenotype, whereas
acidic buffers may promote resistance. We infiltrated MES buffer at pH 5,8 (resembling the apoplastic
pH) and pH 7,5 into the leaf apoplast prior to infection. Infiltration of a 15 mM MgCl$_2$ solution served as
an osmolarity control to exclude effects of high salt concentrations. Surprisingly, both infiltration of the
acidic and the alkaline buffers reduced conidiophore production in comparison to the MgCl$_2$ control (Fig.
5E). This indicates that *Ecr* requires dynamic pH modulations rather than a fixed (e.g. alkaline) pH for
successful host colonization. When analysing individual fungal colonies upon infection on Col-0 pre-

buffered leaves we observed a morphological similarity between fungal colonies on *fer-4* with normal

hyphal growth but strongly reduced conidiophore production at 5 dpi (Fig. 5F). This observation suggests

the importance of apoplastic pH modulations for successful conidiophore production of *Ecr* at later time

points of the infection, which may be regulated through FER-RALF-dependent signalling.

Among other physiological outputs, apoplastic pH modulation is involved in plant growth regulation by

controlling the activity of PECTIN METHYL ESTERASES (PMEs) (Hocq et al., 2023; Hocq et al., 2016;

Sénéchal et al., 2017). PMEs de-esterify cell wall pectins, which increases the negative charge of

homogalacturonans resulting in enhanced Ca^{2+} binding and pectin cross-linking which stiffens the cell

wall (Willats et al., 2001). Importantly, demethylated pectin interacts with LRX-RALF complexes to

structurally organize the cell wall of root hairs and pollen tubes to support polar growth (Moussu et al.,

2023; Schoenaers et al., 2024). PMEs are also involved in biotic interactions. For example, *AtPME17*

expression is strongly up-regulated during infection with the necrotrophic fungus *Botrytis cinerea* and

pme17 mutants are more susceptible to infection (Del Corpo et al., 2020). Recently, *fer-4* mutants were

shown to have constitutively increased PME activity (Biermann et al., 2024). In accordance with LRX

and RALF-peptides being involved in the same physiological pathway as FER, PME activity was

similarly enhanced in *lrx5x* and CRISPR *ralf7x*. *Ecr* infection, however, had no effect on PME activity in

the harvested leaf samples at 5 dpi (Fig. 5G). To test whether PME-mediated pectin demethylation

affects powdery mildew infection, we infiltrated single leaves with epigallocatechin gallate (EGCG), a

compound from green tea (Camellia sinensis) that inhibits PMEs (Lewis et al., 2008). qPCR analysis

of the infiltrated leaves at 5 dpi revealed reduced fungal DNA content on Col-0 leaves infiltrated with

EGCG, indicating that PME activity is potentially required for successful powdery mildew colonization.

Interestingly, EGCG treatment had no additive effect on the resistance of *fer-4* mutants (Fig. 5H). To

complement the inhibitor treatment experiment, we tested the effect of PME-inhibitor (PMEI)

overexpression on *Ecr* colonization. PMEIs inhibit PMEs *in vivo* in a pH dependent manner to fine-tune

cell wall remodelling. Overexpression of *PMEI3* was recently shown do inhibit RALF1 signalling in roots

(Rößling et al., 2024). In line with the EGCG treatment, inhibition of PME activity by *PMEI3*

overexpression resulted in reduced fungal colonization (Fig. 5I). Knockout of single *PMEs,* however,

had not effect on conidiophore production, most likely due to high redundancy of PMEs in Arabidopsis

(Fig. 5J). Taken together, we could show that PME activity, which also affects FER-dependent RALF

signalling, is required for full powdery mildew susceptibility.

RALF-mediated powdery mildew susceptibility is partially independent of FER and FER-signalling

The reduced *Ecr* infection success upon pH buffering prompted us to resolve potential infection-induced

apoplastic pH changes with cellular resolution. For this we used the ratio-metric pH sensor

pUBQ10::SYP122-pHusion (Kesten et al., 2019). We detected an increase in the apoplastic pH close to

Ecr penetration sites at 2 dpi. Surprisingly, similar infection-related pH increase was observed in a

CRISPR-Cas9-generated *fer* mutant (CRISPR *fer*) in the pUBQ10::SYP122-pHusion line (Fig. 6A, B).

When mining the CoIP-MS data for changes in protein phosphorylation upon *Ecr* infection we found the

tendency to enhanced phosphorylation in two areas of the FER intracellular kinase domain, including

the area around amino acid position 511-518 as well as 861-874 (Fig. S7A). The phosphorylation sites
pS500 and pS881, on the other hand, were not regulated (Fig. S7A). The site pS874 has been previously
reported to be phosphorylated upon RALF1 treatment (Haruta et al., 2014). This provides indications
that FER undergoes *Ecr*-dependent phosphorylation, which prompted us to test whether FER kinase
activity is required for *Ecr* colonization. Experiments using a kinase dead FER variant (FER^{K565R})
complemented the powdery mildew resistance phenotype of *fer-4*, suggesting that FER kinase activity
is dispensable for powdery mildew susceptibility (Fig. 6C).

To assess whether FER recruitment to RALF-LLG1 complexes is required for *Ecr* susceptibility, we
infected *fer-4* lines expressing structure-guided FER mutants disrupting the LLG binding interface (Xiao
et al., 2019). Both FER^{N303Y} and FER^{G257A} fully restored powdery mildew susceptibility, indicating that
RALF-induced recruitment of FER to LLG1 is not essential for powdery mildew infection (Fig. 6D). We
thus asked the question whether the susceptibility function of RALFs is primarily attributed to structural
functions rather than FER-LLG-dependent signalling. Therefore, we introduced structure-guided
mutations into RALF23, either disrupting the predicted LRX-binding interface (RALF23$^{Y113A/Y114A}$),
analogous to RALF4$^{Y83A/Y84A}$ (Moussu et al., 2020) or the YISY motif required for LLG binding and FER-
dependent signalling (RALF23$^{Y93A/Y96A}$) (Xiao et al., 2019) and expressed both variants under control of
the 35S promoter in CRISPR *ralf7x*. We tested two lines each with CRISPR *ralf7x* p35S::RALF23$^{Y93A/Y96A}$
displaying similar expression levels as the WT complementation lines. CRISPR *ralf7x*
35S::RALF23$^{Y113A/Y114A}$ lines showed weaker expression compared to the WT construct (Fig. S2E, F).
The weaker expressing RALF23$^{Y113A/Y114A}$ lines could not complement the CRISPR *ralf7x* growth and
enhanced *Ecr* resistance phenotype (Fig. 6E, Fig. S8A). RALF23$^{Y93A/Y96A}$ largely complemented the
growth phenotype (Fig. S8A). However, RALF23$^{Y93A/Y96A}$ showed moderate but incomplete
complementation in powdery mildew infection (Fig. 6E). Together, this data suggests that both
interaction of RALF23 with LLG and LRX complexes is required to support *Ecr* colonization.

We next tested whether application of synthetic RALF23 peptide can affect powdery mildew infection
success in a FER-dependent manner. In plant-bacterial interaction, co-treatment with RALF23 reduced
PAMP-triggered resistance to *Pto* infection (Stegmann et al., 2017). Surprisingly, RALF23 infiltration into
single leaves 24 h prior to *Ecr* infection had a mild resistance-inducing effect in Col-0 (Fig. 6F). This
indicates that infiltration of high doses of a peptide at a fixed time point induces antagonistic responses,
in line with a recent report showing differential concentration-dependent effects of RALF23 on root
immunity (Chen et al., 2024). Remarkably, RALF23 infiltration increased conidiophore production in the
fer-4 mutant background, suggesting that RALF23 exerts FER-independent effects (Fig. 6F). Based on
this we continued testing the effect of RALF23-GFP overexpression on *Ecr* infection outcome,
comparing Col-0 with the *fer-2* mutant. RALF23-GFP overexpression suppresses PAMP-triggered
responses in a FER-dependent manner (Stegmann et al., 2017). Similar to exogenous RALF23 peptide
treatment (Fig. 6F), RALF23-GFP overexpression mildly promoted resistance to *Ecr*. Strikingly,
RALF23-GFP overexpression restored susceptibility in the *fer-2* mutant background (Fig. 6G). *RALF23-*
GFP showed similar transcript abundance in Col-0 and *fer-2* (Fig. S7B). Collectively, based on these
data we hypothesize that RALF-mediated powdery mildew susceptibility is at least partially FER-
independent.

Discussion

This work provides first molecular insights into the mechanism of FER-mediated powdery mildew susceptibility. We revealed that FER`s RALF peptide ligands are critical for the completion of *Ecr*`s asexual life cycle upon host colonisation.

RALFs are perceived by LLG-FER/CrRLK1L complexes to induce downstream signalling. The kinase domain of FER displayed a tendency towards increased phosphorylation of several residues upon *Ecr* infection, but a kinase dead FERK^{565R} fully complemented the *Ecr* colonization phenotype of *fer-4.* FER kinase activity is not required for all FER-related outputs. The *fer-4* defects in ovule fertilization and FLS2-BAK1 complex formation are fully complemented by FER kinase dead variants (Gronnier et al., 2022; Haruta et al., 2018). RALF1-induced calcium influx and root growth inhibition is dependent on FER kinase activity, but *fer-4* FERK^{565R} lines retain mild RALF1 sensitivity (Haruta et al., 2018). Moreover, FER^{K565R} confers sensitivity to RALF1-induced stomatal closure in a protein accumulation-dependent manner, suggesting that FER kinase activity is required for RALF signalling, but can be compensated by higher protein doses (Chakravorty et al., 2018). Disrupting FER`s LLG1 binding interface was also dispensable for *Ecr* colonization. Yet, RALF binding to LLG1 contributes to susceptibility, since *llg1-3* mutants with disrupted RALF binding were more resistant to *Ecr* colonization and, importantly, RALF23 mutants with compromised YISY motif required for LLG binding and CrRLK1L-dependent signalling (Xiao et al., 2019) failed to fully complement the *Ecr* resistance phenotype of CRISPR *ralf7x*. This suggests that RALF-mediated *Ecr* susceptibility is associated with RALF-induced signalling but not strictly dependent on FER and/or FER kinase activity. Consistently, the CRISPR *ralf7x* mutant was more resistant than *fer-4,* and *fer* loss of function mutants displayed increased *Ecr* susceptibility upon RALF23 treatment or RALF23-GFP overexpression, reinforcing at least partially FER-independent functions of RALF-peptides during *Ecr* colonization. FER is dispensable for RALF22-induced root hair tip alkalinisation, calcium influx and cell wall modification (Schoenaers et al., 2024) and RALF23 affects cell wall pectin status in a partially FER-independent manner (Biermann et al., 2024), further supporting FER-independent functions of RALF peptides in vegetative physiological outputs. Our data raises the question whether additional FER-related receptors are required to establish full susceptibility. We speculate that LLGs may recruit further FER-related CrRLK1L proteins that redundantly support host mildew colonization.

FER can interact with related RALF-binding CrRLK1Ls, including HERK1 and ANJ, to form higher order complexes. This was confirmed by our CoIP-MS data according to which FER interacted with several members of the CrRLK1L family (HERK1, MDS1, THE1 and LET1) in mature leaves. HERK1 associates with FER on papilla and synergid cells to control RALF-induced pollen tube penetration (Galindo-Trigo et al., 2020; Lan et al., 2023). MDS1 (also known as LET2) and LET1 have a role in SUMM2-mediated cell death regulation (Huang et al., 2020; Liu et al., 2020). A LET1/LET2 function in RALF perception remains unknown, but LET2 and LET1 form heteromers and associate with LLG1, suggesting participation in higher order CrRLK1L-RALF complexes (Huang et al., 2020). THE1 perceives RALF34 (mutated in CRISPR *ralf7x*) and RALF34's seedling growth inhibition and alkalinisation activity does not require FER (Abarca et al., 2021). Our CoIP-MS data revealed also other FER-associated RLKs, in line with the proposed role of FER as a RLK scaffold (Stegmann et al., 2017). Future work should analyse

combinatorial CrRLK1L mutants for increased *Ecr* resistance and subsequent biochemical analysis for
Ecr-dependent CrRLK1L association and phosphorylation.

RALF binding to LRXs shapes the organization of the cell wall in root hair cells and growing pollen tubes
(Moussu et al., 2023; Schoenaers et al., 2024). We identified LRX1-LRX5 as powdery mildew
susceptibility factors, suggesting that the structural role of LRX-RALF complexes contributes to fungal
host establishment. Consistently, a RALF23 LRX binding interface mutant failed to complement the
CRISPR *ralf7x Ecr* phenotype and overexpression of LRR4-citrine, which lacks the LRX4 extensin
domain, conferred enhanced powdery mildew resistance. LRR4-citrine imposes a dominant negative
phenotype in several LRX-related vegetative physiological outputs (Dünser et al., 2019; Zhao et al.,
2018), likely by sequestration of RALFs, outcompeting of endogenous LRXs and interference with their
cell wall organizing properties. However, LRX1/LRX2 are also required for RALF22 secretion
(Schoenaers et al., 2024). We hence cannot exclude that the *lrx5x Ecr* phenotype is explained by the
lack of signalling competent RALF accumulation in the apoplast. RALFs bind to LRX with high affinity at
acidic pH, while LLG2 requires alkaline conditions for optimal RALF recognition (Moussu et al., 2020),
indicating that pH-dependent coordinated binding and release of RALFs by LRXs could be involved in
CrRLK1L-dependent signalling. It is thus possible that dynamic apoplastic alkalinisation induced by *Ecr*
infection releases RALFs from LRXs to trigger CrRLK1L-dependent signalling events that further support
its reproductive success.

Our work also revealed first downstream mechanistic insights into RALF-FER/LLG1 and/or RALF-LRX-
dependent pathways required for successful *Ecr*-Arabidopsis interaction. The powdery mildew
resistance phenotype of *fer-4* was not attributed to de-regulated SA or JA signalling, suggesting a
different type of underlying mechanism. Our genetic and cell biological data indicate that FER and RALF-
mediated powdery mildew colonization is not directly linked to *MLO*-mediated penetration success. Our
data rather supports that FER-RALF is required for maintenance of compatibility and post-penetration
completion of *Ecr*`s asexual life cycle. The *mlo6* single mutants do not show defects in *Golovinomyces*
orontii epidermal cell penetration, but a trend towards reduced conidiophore production. The *mlo2* single
mutant still maintains ~50% of host cell entry, but abolished conidiophore production (Consonni et al.,
2006), raising the question whether MLO also contributes to long-term host association. Therefore, our
data does not exclude a role of RALF-induced and MLO-dependent signalling required for *Ecr*
sporulation. We also cannot fully rule out that FER and/or RALFs may act redundantly with other
CrRLK1Ls/RALFs to contribute to *Ecr* epidermal cell penetration.

Our data suggests that powdery mildew susceptibility requires functional apoplastic pH homeostasis
and modifications of the cell wall pectin methyl-esterification status, both processes in which FER-RALF
and/or LRX-RALF modules fulfil central regulatory functions. Constitutive AHA1 activity in *ost2-2D*
mutants or fusicoccin treatment promoted powdery mildew resistance. Importantly, fusicoccin treatment
was less effective in *fer-4*, suggesting that mutating FER and chemically activating AHA1 affects the
same pathway. It is also possible that fusicoccin treatment cannot further promote apoplastic
acidification in *fer-4*, which displays constitutively elevated proton extrusion. Earlier work revealed that
barley powdery mildew infection is accompanied by pH shifts and the induction of apoplastic
alkalinisation (Felle et al., 2004). Importantly though, mildew-induced pH shifts were dynamic, with

fluctuations between alkaline and more acidic conditions (Felle et al., 2004). This supports our notion
that disrupting apoplastic pH homeostasis by buffering inhibits *Ecr* host colonization.

Our data further reveals that the pectin methylation status determines powdery mildew pathogenesis.
Overexpression of PMEI3 and EGCG treatment, a well-established chemical PME inhibitor, promoted
Ecr resistance. PMEI3 inhibits PME3, which is involved in FER-mediated epidermal cell lobing (Lin et
al., 2022). We could not resolve *Ecr* phenotypes in *pme3* loss of function mutants, suggesting
redundancy with other PMEs in this 66 member multi-gene family in Arabidopsis (Harholt et al., 2010;
Pelloux et al., 2007). PMEI3 activity is strongly pH dependent with the highest activity at acidic conditions
(Xu et al., 2022). RALF-induced apoplastic alkalinisation may inhibit PMEI3, promoting PME activity for
cell wall remodelling. RALF22, and RALF4 bind to LRXs to regulate pectin compaction in the cell wall.
LRXs shape RALFs to expose a polycationic region for interaction with de-methylated and acidic pectin
(Moussu et al., 2023; Schoenaers et al., 2024). Moreover, PME activity is required for RALF1-induced
root growth inhibition independent of LRX1-LRX5 (Rößling et al., 2024) and RALF1 forms molecular
condensates with pectin required for downstream FER-LLG1-dependent responses (Liu et al., 2024).
Strikingly, a RALF1 mutant that is compromised in binding to FER-LLG1 and does not induce
alkalinisation (Liu et al., 2024; Pelloux et al., 2007; Stegmann et al., 2017) fails to form RALF-pectin
condensates (Liu et al., 2024), suggesting that RALF-pectin association and signalling outputs are
intimately interconnected. Our work revealed that *fer*, CRISPR *ralf7x* and *lrx5x* display elevated steady-
state PME activity, an output that was unaffected by *Ecr* infection and is in accordance with previous
findings (Biermann et al., 2024). This may be explained by the requirement of de-methylated pectin for
RALF signalling and that the absence of LRX-RALFs and CrRLK1L-LLG-RALFs might activate plant
feedback mechanisms to promote PME activity. Pectin de-methylation is an irreversible process. The
high steady-state PME activity in *fer-4* may thus explain the reduced effect of EGCG-promoted *Ecr*
resistance. The *fer-4* mutant may accumulate high levels of de-methylated pectin in the cell wall,
abrogating a detectable effect by apoplastic PME inhibition. FER is required for expression of the
polygalacturonase ADPG1 upon disruption of lignin biosynthesis (Liu et al., 2023), reinforcing the
function of FER-dependent cell wall remodelling during stress. We propose a mechanism in which pH-
dependent pectin de-methylesterification is a prerequisite for RALF-mediated sustained powdery mildew
host colonization. In accordance, endogenous pectin modification affects powdery mildew infection
outcome. Mutants in the pectate *lyase POWDERY MILDEW RESISTANT 6 (PMR6)* are more resistant
to *Golonivomyces cichoracearum* with normal host penetration, delayed fungal spread, strongly reduced
sporulation and normal SA/JA signalling (Vogel et al., 2002), resembling phenotypes of CRISPR *ralf7x*
and *fer-4*. PME-induced pectin demethylation is a prerequisite for pectate lyase-mediated degradation
(Wormit and Usadel, 2018). Yet, PME inhibition, which should reduce PMR6 activity, and PMEI over
accumulation, which should promote pectate lyase activity, limit fungal colonization, reinforcing the
notion that fine-tuned pectin remodelling, in concert with dynamic apoplastic pH modulations, underlies
powdery mildew reproductive success.

Ecr conidiation likely requires significant energy input and it remains unclear how disruption of the RALF
pathway and/or pH and cell wall dynamics affects this process. The fungal haustorium is surrounded by
the extrahaustorial matrix (EHMx) and the extrahaustorial membrane (EHM), a plant membrane that is

continuous with the plasma membrane but has distinctive properties (Hückelhoven and Panstruga,
2011; Polonio et al., 2021). At the site of invasion the extrahaustorial matrix is sealed with densely
packed cell wall depositions forming the haustorial neck region (Micali et al., 2011). This renders the
EHMx a separate apoplastic compartment that can have different conditions compared to the
neighbouring plant cell wall (Hückelhoven and Panstruga, 2011). Nutrient uptake from the EHMx is most
likely dependent on the compartments pH. From rust fungi it is known that ATPase activity is enhanced
in haustorium derived membrane vesicles compared to other fungal structures, indicating a role of fungal
ATPases in nutrient uptake via H^+ symporters (Voegele and Mendgen, 2011). The nutrients that are
found in the EHMx and taken up by the fungus should derive from the invaded plant cell. They must first
be exported from the host cell before they are available to the fungus. Nutrient export from the plant cell
involves SWEET proteins and the uptake is facilitated by sucrose transport proteins (STP), both of which
control apoplastic sugar levels and affect plant microbe interactions (Bezrutczyk et al., 2018). STPs
usually transport sugars from the apoplast into the plant cell and control available sugars for microbial
invaders (Geiger, 2020). In Arabidopsis, AtSTP8 localizes to the EHM and overexpression of AtSTP8
increases susceptibility to powdery mildew, indicating that STP8 is involved in nutrient supply (Liu et al.,
2021). *STP4* expression in leaves is enhanced after powdery mildew infection (Fotopoulos et al., 2003)
and STP13 is associated with resistance to both powdery mildew and rust fungi in wheat and barley
(Milne et al., 2019). Perception of flg22 by FLS2-BAK1 results in STP13 phosphorylation to enhance
cellular uptake capacity and limit apoplastic sugar availability for bacterial pathogens (Yamada et al.,
2016). STPs are H^+ symporters, and STP8 has a pH optimum at 5,5 for cellular import (Liu et al., 2021).
Disruption of pH dynamics through mutating RALF-CrRLK1L could impair nutrient supply by affecting
spatiotemporal export of nutrients to the EHMx and interfering with fungal uptake. In accordance, *fer*
mutants show enhanced sucrose uptake, starch accumulation and hypersensitivity to exogenous
sucrose treatment, a phenotype shared with *shv3 svl1* mutants (Yang et al., 2015; Yeats et al., 2016).
The *shv3 svl1* mutants also display reduced apoplastic pH (Yeats et al., 2016) and we identified SHV3
as an FER interactor. Since RALF1`s effects on pH-dependent root growth likely do not involve direct
FER-mediated AHA regulation, FER may also induce cellular H^+ import to counteract apoplastic
acidification via yet unknown transporters (Li et al., 2022a). It is possible that *Ecr* infection induces
RALF-CrRLK1L-dependent changes to the apoplastic pH that will reprogram transporter topology to
favour haustorial sugar uptake over plant cell import. Supporting our hypothesis, we identified several
proteins associated with transmembrane transporter activity as FER interactors in our Co-IP MS
experiment. This includes STP1, involved in resistance against bacterial infection, the ABC transporter
PEN3, associated with Arabidopsis host and non-host resistance against powdery mildew, and the
PLASMA MEMBRANE INTRINSIC 2;2 (PIP2;2) aquaporin, a recently discovered FER target (Collins et
al., 2003; Liu et al., 2025; Stein et al., 2006; Yamada et al., 2016). We propose that FER`s interaction
with membrane localized transport proteins and the pH-regulatory function of RALF-CrRLK1L modules
shape the transporter landscape at the host-*Ecr* interface to support colonization. Moreover, RALFs may
control the pectin methylesterification status of the cell wall surrounding powdery mildew penetration
sites. Methylated or demethylated pectins have not been detected at the haustorial neck region, which
does not exclude a role for fungal accommodation in the neighbouring cell wall (Micali et al., 2011).
Moreover, *fer* mutants show reduced cell wall cellulose content, further suggesting structural defects

that may impact *Ecr* host interaction. We speculate that this could control the ability of the fungus to
establish a defined pH environment for optimising transporter-mediated nutrient uptake. Future studies
should dissect the plant-*Ecr* interface in *FER/LRX-RALF*, pH and cell-wall related mutants with
ultrastructural analysis by electron microscopy.

Other plant pathogens produce RALFs to establish host colonization, including *Fusarium oxysporum*
and root knot nematodes (Masachis et al., 2016; Zhang et al., 2020). Interestingly, these RALF peptide
mimics lack conserved sequence elements required for LRX binding, but contain a modified YISY motif
required for LLG-CrRLK1L-dependent signalling (Pearce et al., 2010). This suggests that fungal RALFs
solely rely on CrRLK1L-induced signalling outputs and not on structurally re-shaping cell wall
architecture for their virulence function. None of the analyzed powdery mildew genomes contained RALF
peptide mimics, suggesting that *Ecr* likely relies on the combination of endogenous RALF signalling and
cell wall remodelling to support sustained host interaction. Our work thus highlights a novel mode of
RALF function, combining signalling and structural roles for the control of plant microbe interactions.

Materials and Methods

Powdery mildew propagation

Powdery mildew fungi of the species *Erysiphe cruciferarum* were propagated weekly by transferring
spores to 4 – 5 week old uninfected Col-0 plants. Powdery mildew spores were used for further
inoculation 3 – 4 weeks after propagation. Powdery mildew propagation and infection experiments were
performed in an environmentally controlled growth cabined (22 °C, 70 % relative humidity, 12 h
photoperiod).

Molecular cloning

To generate p35S::RALF23 and p35::MLO2-mCherry lines the respective gene sequences were cloned
from gDNA using the primers listed in Table S3 and assembled into a GoldenGate-modified pCB302
binary vector for plant expression. The structure guided RALF23 mutant lines p35S::RALF23$^{Y93A/Y96A}$
and p35S::RALF23$^{Y113A/Y114A}$ were obtained by site-directed mutagenesis using the respective primers
listed in Table S3 and likewise assembled into a GoldenGate-modified pCB302 binary vector for plant
expression. To generate CRISPR-Cas9 mutants the software tool chopchop
(https://chopchop.cbu.uib.no/) was used to design target sites in the respective genes (1 to 2 per gene
of interest). All target sequences used in this study are listed in Table S4. Individual guide RNAs with
gene specific target sites were obtained by gene synthesis (Twist Bioscience, USA). Using a
GoldenGate-adapted pUC18-based vector the guide RNA fragments were assembled. Higher order
gRNA stacks were cloned into pICSL4723OD with FastRed-pRPS5::Cas9 (Castel et al., 2019). For final
floral dip transformation of Arabidopsis all generated plant expression constructs were transformed into
Agrobacterium tumefaciens strain GV3101.

Plant material and growth conditions

Plant lines used in this study are listed in Table S2 with the according references.
Isolation of homozygous CRISPR mutants was done based on red fluorescence of transformed seeds.
In the T1 generation red fluorescent seeds were selected and grown on soil for genotyping with gene
specific primers. In the T2 generation mutants lacking the transgene were identified based on the loss
of fluorescence. Simliarly, homozygous p35S::MLO2-mCherry lines were selected based on red
fluorescence. Homozygous p35S::RALF23 lines were selected by transgene mediated BASTA
resistance. All plants were vernalized for 2 – 3 days in the dark at 4 °C and grown with two plants per
pot in environmentally controlled conditions (21 °C, 55 – 65% relative humidity, 8 h photoperiod). Five
to six week-old plants were used for infection.

Conidiophore counting and fungal penetration assay.

Conidiophore production was analysed 5 days after infection with *Ecr*. Four leaves per genotype were
harvested from 4 individual infected plants. The leaves were destained using EtOH:Acetic Acid (6:1)
and subsequently stained using ink (ink: 25 % acetic acid, 9:1). Ink-stained fungal structures were
visualized using the Zeiss AXIO imager Z1.m microscope with a 20x magnification for conidiophore
counting. Fungal penetration success was similarly analysed and quantified at 2 dpi. A penetration
attempt was scored as successful when the spore formed a penetration tube and first hyphal structures.
The host cell entry rate was determined by the number of successful penetrations per attempts (in %).

Quantification of fungal DNA by qPCR

For quantification of fungal DNA two infected plants per genotype were harvested at 5 dpi. Plant material
was ground in liquid nitrogen and equal amounts of plant powder was transferred into a 2 ml Eppendorf
tube. 500 µL extraction buffer (200 mM Tris-HCl pH 7,5, 250 mM NaCl, 25 mM EDTA, 0,5 % SDS) were
added per sample. After vortexing, 500 µL chloroform was added with subsequent mixing. After
centrifugation, the top layer was transferred into 500 µL isopropanol and mixed. The DNA pellet was
precipitated by centrifugation and washed with 70 % EtOH before resuspension in 50 µL water and
measurement of DNA concentration. For qPCR, the DNA concentration was adjusted to 20 ng/µL for all
samples. RT-qPCR experiments were performed using Maxima SYBR green mix (Thermo Fisher
Scientific, USA) on an AriaMx Real-Time PCR system (Agilent Technologies, USA). The amount of
fungal DNA was calculated relative to the amount of plant DNA using primers for the Arabidopsis small
RubisCO subunit (AtRbcS) and *Ecr* tubulin (Table S3). All samples were normalized to Col-0.

Infiltration experiments

622 For RALF peptide treatment individual leaves were infiltrated with 1 µM synthetic RALF23 peptide
623 (Sequence: ATRRYISYGALRRNTIPCSRRGASYYNCRRGAQANPYSRGCSAITRCRRS) 24 h prior to
624 infection. Fusicoccin (10 µM) and EGCG (50 µM) and MES buffers were infiltrated 2 h before *Ecr*
625 infection and subsequent microscopical analysis or qPCR-mediated assessment of infection success at
626 5 dpi.

Gene expression analysis

Expression of SA and JA responsive genes was analysed at 5 dpi upon *Ecr* infection. Four individual
plants were harvested per genotype and immediately frozen in liquid nitrogen. The plant material was

ground and equal amounts of plant powder were transferred into individual 2 ml Eppendorf tubes. For
RNA extraction, TRIzol reagent (Roche, Switzerland) and subsequent purification with Direct-zol™ RNA
Miniprep Plus kit (Zymo Research, Germany) with on column DNAse I digestion was used. 1.5 – 2 µg
of total RNA was reverse transcribed with random hexamer primer and Revert Aid reverse transcriptase
(Thermo Fisher Scientific, USA). RT-qPCR experiments were performed using Maxima SYBR green
mix (Thermo Fisher Scientific, USA) on an AriaMx Real-Time PCR system (Agilent Technologies, USA).
Expression levels of the tested genes were normalized to the house-keeping gene *UBIQUITIN 5*
(*UBQ5*). All primers used for gene expression analysis are listed in Table S3.

PME activity assay

PME activity assays were performed as previously described (Bethke et al., 2014). In brief, plant material
of untreated and infected plants was harvested 5 dpi and ground in liquid nitrogen. Equal amounts of
plant material were transferred per genotype and treatment into individual 2 ml tube. The plant powder
was incubated with 500 µL protein extraction buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 10%
glycerol, 5 mM dithiothreitol, 1% protease inhibitor cocktail, 2 mM Na2MoO4, 2.5 mM NaF, 1.5 mM
activated Na3VO4, 1mM phenylmethanesulfonyl fluoride, 0.5% IGEPAL) for 30 min at 8°C before
centrifugation at 16,000 g for 30 min. The supernatant was transferred into a fresh 1,5 ml tube and the
protein concentration was determined by Bradford assay. The protein concentration was adjusted to 3
µg/µL. To assess PME activity, the liquid was loaded into holes in an agarose gel plate containing
esterified pectin (1.2 % (w/v) agarose, 0.1 % (w/v) pectin from apple, 12 mM citric acid, 50 mM Na_2HPO_4
pH 7). Per hole 15 µL liquid were loaded and 4 technical replicates were performed per genotype and
treatment. The plates were incubated at 37°C for 16 h. After incubation, plates were briefly washed with
water and stained with 0.05 % ruthenium red for 30 min. Residual dye was washed off with water and
images of the plates were captured by scanning. Darker stained areas correlating with PME activity
were quantified using Fiji ImageJ software.

AHA-phosphorylation analysis

Infected plant material was harvested 3 dpi and ground in liquid nitrogen. 20 µL extraction buffer (3 %
[w/v] SDS, 30 mM TRIS-HCl pH 8.0, 10 mM EDTA. 10 mM NaF, 30 % [w/v] Sucrose, 0.012% [w/v]
Coomassie Brilliant Blue, 15 % [v/v] 2-Mercaptoethanol) per mg plant material was added and samples
were vortexed two times for 30 sec. Subsequently, samples were centrifuged at 12,000 x g for 1 min.
The supernatant was immediately used for SDS PAGE and western blot. Protein bands were detected
using polyclonal AHA and pThr947 antibodies (Hayashi et al., 2010). Band intensities were quantified
using Fiji ImageJ software.

Quantification of salicylic acid, dihydroxybenzoic acids and camalexin

For the quantification of salicylic acid (SA), dihydroxybenzoic acids (DHBAs) and camalexin 40-60 mg
of ground fine leaf powder was extracted as described by (Nawrath and Métraux, 1999) with minor
modifications. First, 600 µl 70% methanol containing 250 ng of *ortho*-anisic acid (oANI) as internal
standard were added and samples were incubated for 1 hour at 65°C. The samples were centrifuged
for 1 min at 14,000 x g and the supernatant was collected in a fresh reaction tube. The powder was re-

extracted with 600 µl of 90% methanol for 1 hour at 65°C, centrifuged as before and the supernatants

were combined. The solvent was evaporated using a rotational vacuum concentrator for 2.5 hours at

35°C and 10 mbar. After a precipitation step with 500 µl of 5 % trichloroacetic acid and centrifugation

for 10 min at 3,500 x g, the supernatant was transferred in a fresh reaction tube and partitioned against

600 µl cyclohexane / ethyl acetate (1:1). The samples were shaken, centrifuged for 10 min at 3,500 x g

and the organic (upper) phase was transferred in a fresh reaction tube. The partitioning step was

repeated and the organic phases combined to obtain free SA, DHBAs and camalexin. For the extraction

of the glycosylated SA and DHBAs, the aqueous phase was acidified by adding 8 M hydrochloric acid

(1:1) and incubated for 1 hour at 80°C. Again, 250 ng of oANI were added and the hydrolyzed extracts

partitioned two times against 600 µl cyclohexane / ethyl acetate (1:1) as described above to obtain

glycosylated SA and DHBAs. All organic phases (free and glycosylated fractions) were evaporated using

a rotational vacuum concentrator to near dryness to prevent sublimation of SA. The reduced extracts

were dissolved in 400 µl of 90% 25 mM potassium dihydrogen phosphate buffer (KH_2PO_4), pH 2.6, and

10% acetonitrile (ACN). HPLC analysis was performed on an Agilent Technologies 1260 Infinity II

system equipped with an G7121B fluorescence detector (Agilent Technologies, USA). SA, DHBAs,

camalexin and oANI were separated on a SecurityGuard Ultra guard column (3 mm; Phenomenex, USA)

followed by an InfinityLab Poroshell 120 SB-C18 analytical column (3 x 100 mm, 2.7 µm; Agilent

Technologies, USA) at 40°C. The separation was performed with a solvent gradient of eluent (A) 25mM

KH_2PO_4, pH 2.6 and eluent (B) ACN:H_2O (99:1) at 1 ml/min flow rate (all steps in v/v): 0 to 1 min: 10%

B; 1 to 6 min: 10 – 25 %B; 6 to 8.5 min: 25 – 80 %B; 8.5 to 10.5 min: 80% B; 10.5 to 11 min: 80 – 10%

B; 11 to 14 min 10% B. The fluorescence detector was used to detect oANI (excitation 305 nm, emission

365 nm), SA (excitation 305 nm, emission 407 nm), DHBAs (excitation 320 nm, emission 450 nm) and

camalexin (excitation 318 nm, emission 370 nm). For quantification, standard dilution series of oANI,

SA, 2,3-DHBA, 2,5-DHBA and camalexin were measured in the range of 1 to 100 ng.

Histochemical staining of callose deposition in rosette leaves and trichomes

Three rosette leaves per 35-d-old Arabidopsis plant were collected in 2-mL reaction tubes and destained

twice for 24 h each in 70% ethanol (v/v). Afterwards, rosette leaves were incubated over-night in aniline

blue solution (0.01% Aniline Blue (m/v) in 150 mM dipotassium hydrogen phosphate buffer) and

mounted on a microscopy slide. Micrographs were captured by a BZ-9000 transmitted light and

fluorescence microscope (Keyence, Osaka, Japan) using UV illumination. Images were optimized by

Adobe Photoshop 2024 in an identical manner for all micrographs. Trichome and trichome branch

number were scored by counting.

Confocal microscopy

Confocal laser-scanning microscopy was performed using a Leica TCS SP5 (Leica, Germany)

microscope (with Leica Application Suite X 3.7.4.23463). All images were taken with a 20 x water

immersion objective. To analyse the localization of FER-GFP and MLO2-mCherry, adult plants were

infected with *Ecr* and imaged at 2 dpi. To visualize GFP fluorescence an argon laser with an emitting

wavelength of 488 nm was used and the GFP signal was detected from 500 nm to 550 nm with a HyD

detector. The mCherry fluorophore was excited with a DPSS 561 laser emitting at 561 nm and a

detection window of 600 - 640 nm. Changes in the apoplastic pH around fungal infection sites were
analysed using plant lines expressing the pUBQ10::SYP122-pHusion. Four to five week-old plants were
infected with *Ecr* and confocal imaging was performed at 2 dpi. The GFP signal was detected with an
emitting wavelength of 488 nm and a detection window of 500 – 545 nm. The RFP signal was excited
with 561 nm and detected between 650 nm and 695 nm. To compare the fluorescence signal close to
the penetration site with signal in unpenetrated areas, zoomed in Z-stacks with identical settings were
taken at both locations. The fluorescence intensities and GFP/RFP ratios were determined using Fiji
Image J.

Co-IP MS/MS

Five week-old FER-GFP and Lti6b-GFP plants were infected with powdery mildew and harvested,
together with untreated samples at 24 h post infection. Per genotype and treatment, the rosettes of 12
plants were harvested and ground in liquid nitrogen. All plant material was used for further protein
extraction. Proteins were extracted with 1 ml extraction buffer (50 mM TRIS-HCl pH 7,5, 100 mM NaCl,
10 % glycerol, 2mM EDTA, 2mM DTT, 1 % protease inhibitor cocktail, 1 % PMSF, 1% IGEPAL) per 1
ml plant material. For immunoprecipitation, GFP-Trap agarose beads (ChromoTek) were used and
incubated with the crude extract for 3 h at 4 °C. Beads were washed 3 times with wash buffer (50 mM
Tris-HCl pH 7.5, 50 mM NaCl, 1 mM phenylmethanesulfonyl fluoride, 0.1 % IGEPAL). Subsequently,
washed beads were incubated in 50 µL of Laemmli sample buffer for 10 min at 95 °C. Next, in-gel trypsin
digestion was performed according to standard procedures (Shevchenko et al., 2006). Briefly, samples
were run on a Nu-PAGE™ 4%–12% Bis-Tris protein gel (ThermoFisher Scientific) for about 1 cm.
Subsequently, the still not size-separated single protein band per sample was cut out, reduced (50mM
dithiothreitol), alkylated (55mm chloroacetamide) and digested overnight with trypsin (Trypsin Gold,
mass spectrometry grade, Promega). The peptides obtained were dried to completeness and
resuspended in 24 µL of 2% acetonitrile, 0.1% formic acid in HPLC grade water. Finally, 5 µL of sample
were injected per mass spectrometric (MS) measurement.

LC-MS/MS data acquisition was carried out on a Dionex Ultimate 3000 RSLCnano system coupled to a
Q-Exactive HF-X mass spectrometer (ThermoFisher Scientific, Bremen). Peptides were delivered to a
trap column (ReproSil-pur C18-AQ, 5 µm, Dr. Maisch, 20 mm × 75 µm, self-packed) at a flow rate of 5
µL/min in 0.1% formic acid in HPLC grade water. After 10 minutes of loading, peptides were transferred
to an analytical column (ReproSil Gold C18-AQ, 3 µm, Dr. Maisch, 450 mm × 75 µm, self-packed) and
separated using a 50 min gradient from 4% to 32% of solvent B (0.1% FA, 5% DMSO in acetonitrile) in
solvent A (0.1% FA, 5% DMSO in HPLC grade water) at 300 nL/min flow rate. The Q-Exactive HF-X
mass spectrometer was operated in data dependent acquisition (DDA) and positive ionization mode.
MS1 spectra (360–1300 m/z) were recorded at a resolution of 60k using an automatic gain control (AGC)
target value of 3 e^6 and maximum injection time (maxIT) of 45 msec. Up to 18 peptide precursors were
selected for fragmentation. Only precursors with charge state 2 to 6 were selected and dynamic
exclusion of 30 sec was enabled. Peptide fragmentation was performed using higher energy collision
induced dissociation (HCD) and a normalized collision energy (NCE) of 26%. The precursor isolation
window width was set to 1.3 m/z. MS2 Resolution was 15.000 with an automatic gain control (AGC)
target value of 1e^5 and maximum injection time (maxIT) of 25 msec.

Protein and phosphopeptide identification and quantification were performed using the software
MaxQuant (version 1.6.3.4) (Tyanova et al., 2016) with its built-in search engine Andromeda. MS2
spectra were searched against the TAIR *Arabidopsis thaliana* protein database (downloaded March
2022, 48231 protein entries) supplemented with the GFP protein sequence as well as with common
contaminants (built-in option in MaxQuant). Trypsin/P was specified as proteolytic enzyme.
Carbamidomethylated cysteine was set as fixed modification. Phosphorylation of
serine/threonine/tyrosine, oxidation of methionine and acetylation at the protein N-terminus were
specified as variable modifications. Results were adjusted to a 1% false discovery rate on peptide
spectrum match (PSM) level and protein level employing a target-decoy approach using reversed
protein sequences. The minimal peptide length was defined as 7 amino acids and the "match-between-
run" functionality was enabled (matching time window 0.7 min, alignment time window 20 min). For
protein quantification the MaxQuant output table proteinGroups.txt and Label-Free Quantification (LFQ)
(Cox et al., 2014) intensities were used. Only proteins detected in at least two biological replicates in at
least one experimental condition were further processed. Missing values were imputed by a protein-
specific constant value, which was defined as the lowest detected protein-specific LFQ-value over all
samples divided by two, using the data visualisation platform omicsViewer (DOI:
10.1101/2022.03.10.483845). Additionally, a maximal imputed LFQ value was defined as 15% quantile
of the protein distribution from the dataset. The volcano plots (Figure S4A, S5A) were generated using
RStudio. Interactions were marked as significant if they passed the threshold of FDR < 0.05 and fold
change > 2. For phosphopeptide detection the MaxQuant output table Phospho(STY)Sites.txt was used.
Only phosphosites detected on FERONIA in at least two biological replicates in at least one experimental
condition were considered. The FER-GFP infected with powdery mildew and untreated experiments
were performed in biological triplicates, while all other experiments were performed in biological
quadruplets.

Petiole length measurements

Petioles were measured from rosettes of 4 week-old plants. Individual plants were cut off from the root
and leaves were detached close to the base. Individual leaves were imaged, and petiole length was
quantified using Fiji ImageJ software.

Identification of putative powdery mildew RALF-likes

To identify putative RALF-like peptides in the predicted proteins and genome assemblies of selected
phytopathogenic fungi, we used the hmmsearch program from the hmmer suite v3.3.2 (www.hmmer.org)
with default parameters. The hidden markov model profile (.hmm) for the RALF motif (PF05498.15) was
obtained from the PFAM website (https://ftp.ebi.ac.uk/pub/databases/Pfam/current_release/Pfam-
A.hmm.gz), access data 19.11.2023). The RALF HMM profile was used to search predicted protein
sequences from the selected fungal species (see Supplementary Table X) using the hmmsearch
function. Hits with E-values below $1e^{-05}$ were retained. To perform the hmmsearch analysis on the
genome assemblies, we used a custom Python script to split the contigs into 300 bp fragments with a
sliding window of 50bp. All six open reading frames of these split sequences were translated, and stop
codons were removed to generate a multi-FASTA file suitable for be processed with hmmsearch. Search

hits with E-values < 1e-05 were retained. Accession number of genome assemblies used in this analysis
are reported in Supplementary Table S1.

Quantification and statistical analysis

789 GraphPad Prism (Version 8.0.1) was used to perform statistical analysis. Detailed descriptions of the
790 sample size, p-values and statistical methods are indicated in the respective figure legends.

Funding

792 This work was funded by the following agencies: Deutsche Forschungsgemeinschaft (DFG) grants
793 STE2448/4-1 (HL and MS), TRR356 TP B09 (HL, SS and MS), EN 1071/3-1 (TE) and the Technical
794 University of Munich (MS, HL, SS, MM, RH).

Author contributions

796 Conceptualization: MS; Investigation: HL, MS, SS, JH, KM, MM, TL, GH; Funding acquisition: MS, RH,
797 JH, TE, TK; Project administration: MS; Supervision: MS, HL, TE, JH, RH; Writing – original draft: HL,
798 MS; Writing – review & editing: SS, MM, KM, TK, TE, JH, RH.

Acknowledgements

We thank Stefanie Ranf for providing GoldenGate vectors for molecular cloning and Mark Youles (TSL
Norwich) and Laurence Tomlinson (TSL Norwich) for providing the pICSL4723OD vector for CRISPR-
Cas9 cloning. We thank Hannah König for supporting the collection of micrographs to assess callose
depositions in rosette leaves and trichomes. The authors thank Verena Breitner and Franziska
Hackbarth for excellent technical assistance and maintenance of mass spectrometers at the BayBioMS.
The Q Exactive HFX mass spectrometer was funded in part by the German Research Foundation (INST
95/1435-1 FUGG).

Declaration of interests

808 The authors declare no competing interests.

Data availability

810 The mass spectrometric raw files as well as the MaxQuant output files have been deposited to the
811 ProteomeXchange Consortium via the PRIDE partner repository and can be accessed using the
812 identifier PXD059737 (https://proteomecentral.proteomexchange.org/cgi/GetDataset?ID=PXD059737).
813 Reviewer account username: reviewer_pxd059737@ebi.ac.uk, password: lumM0fo5d8wq.

References

Abarca, A., Franck, C.M., and Zipfel, C. (2021). Family-wide evaluation of RAPID ALKALINIZATION FACTOR peptides. Plant Physiology **187**:996-1010. doi:10.1093/plphys/kiab308.

Acevedo-Garcia, J., Spencer, D., Thieron, H., Reinstädler, A., Hammond-Kosack, K., Phillips, A.L., and Panstruga, R. (2017). mlo-based powdery mildew resistance in hexaploid bread wheat generated by a non-transgenic TILLING approach. Plant Biotechnology Journal **15**:367-378. doi:10.1111/pbi.12631.

Baunsgaard, L., Fuglsang, A.T., Jahn, T., Korthout, H.A.A.J., De Boer, A.H., and Palmgren, M.G. (1998). The 14–3-3 proteins associate with the plant plasma membrane H1-ATPase to generate a fusicoccin binding complex and a fusicoccin responsive system. The Plant Journal **13**:661-671. doi:10.1046/j.1365-313X.1998.00083.x.

Bethke, G., Grundman, R.E., Sreekanta, S., Truman, W., Katagiri, F., and Glazebrook, J. (2014). Arabidopsis PECTIN METHYLESTERASEs contribute to immunity against Pseudomonas syringae. Plant Physiol **164**:1093-1107. doi:10.1104/pp.113.227637.

Bezrutczyk, M., Yang, J., Eom, J.-S., Prior, M., Sosso, D., Hartwig, T., Szurek, B., Oliva, R., Vera-Cruz, C., White, F.F., et al. (2018). Sugar flux and signaling in plant–microbe interactions. The Plant Journal **93**:675-685. doi:10.1111/tpj.13775.

Bhat, R.A., Miklis, M., Schmelzer, E., Schulze-Lefert, P., and Panstruga, R. (2005). Recruitment and interaction dynamics of plant penetration resistance components in a plasma membrane microdomain. Proceedings of the National Academy of Sciences **102**:3135-3140. doi:10.1073/pnas.0500012102.

Biermann, D., von Arx, M., Xhelilaj, K., Séré, D., Stegmann, M., Wolf, S., Zipfel, C., and Gronnier, J. (2024). A RALF-Brassinosteroid morpho-signaling circuit regulates *Arabidopsis* hypocotyl cell shape. bioRxiv:2024.2003.2028.587199. doi:10.1101/2024.03.28.587199.

Bürger, M., and Chory, J. (2019). Stressed Out About Hormones: How Plants Orchestrate Immunity. Cell Host & Microbe **26**:163-172. doi:10.1016/j.chom.2019.07.006.

Büschges, R., Hollricher, K., Panstruga, R., Simons, G., Wolter, M., Frijters, A., van Daelen, R., van der Lee, T., Diergaarde, P., Groenendijk, J., et al. (1997). The Barley *Mlo* Gene: A Novel Control Element of Plant Pathogen Resistance. Cell **88**:695-705. doi:10.1016/S0092-8674(00)81912-1.

Castel, B., Tomlinson, L., Locci, F., Yang, Y., and Jones, J.D.G. (2019). Optimization of T-DNA architecture for Cas9-mediated mutagenesis in Arabidopsis. PLOS ONE **14**:e0204778. doi:10.1371/journal.pone.0204778.

Chakravorty, D., Yu, Y., and Assmann, S.M. (2018). A kinase-dead version of FERONIA receptor-like kinase has dose-dependent impacts on rosette morphology and RALF1-mediated stomatal movements. FEBS Letters **592**:3429-3437. doi:10.1002/1873-3468.13249.

Chen, J., Xu, F., Qiang, X., Liu, H., Wang, L., Jiang, L., Li, C., Wang, B., Luan, S., Wu, D., et al. (2024). Regulated cleavage and translocation of FERONIA control immunity in Arabidopsis roots. Nature Plants **10**:1761-1774. doi:10.1038/s41477-024-01823-8.

Collins, N.C., Thordal-Christensen, H., Lipka, V., Bau, S., Kombrink, E., Qiu, J.-L., Hückelhoven, R., Stein, M., Freialdenhoven, A., Somerville, S.C., et al. (2003). SNARE-protein-mediated disease resistance at the plant cell wall. Nature **425**:973-977. doi:10.1038/nature02076.

Consonni, C., Humphry, M.E., Hartmann, H.A., Livaja, M., Durner, J., Westphal, L., Vogel, J., Lipka, V., Kemmerling, B., Schulze-Lefert, P., et al. (2006). Conserved requirement for a plant host cell protein in powdery mildew pathogenesis. Nat Genet **38**:716-720. doi:10.1038/ng1806.

Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M. (2014). Accurate Proteome-wide Label-free Quantification by Delayed Normalization and Maximal Peptide Ratio Extraction, Termed MaxLFQ*. Molecular & Cellular Proteomics **13**:2513-2526. doi:10.1074/mcp.M113.031591.

Delulio, G.A., Guo, L., Zhang, Y., Goldberg, J.M., Kistler, H.C., and Ma, L.-J. (2018). Kinome Expansion in the Fusarium oxysporum Species Complex Driven by Accessory Chromosomes. mSphere **3**:10.1128/msphere.00231-00218. doi:10.1128/msphere.00231-18.

Del Corpo, D., Fullone, M.R., Miele, R., Lafond, M., Pontiggia, D., Grisel, S., Kieffer-Jaquinod, S., Giardina, T., Bellincampi, D., and Lionetti, V. (2020). AtPME17 is a functional Arabidopsis thaliana pectin methylesterase regulated by its PRO region that triggers PME activity in the resistance to Botrytis cinerea. Molecular Plant Pathology **21**:1620-1633. doi:10.1111/mpp.13002.

Deslauriers, S.D., and Larsen, P.B. (2010). FERONIA Is a Key Modulator of Brassinosteroid and Ethylene Responsiveness in Arabidopsis Hypocotyls. Molecular Plant **3**:626-640. doi:10.1093/mp/ssq015.

Dobón, A., Canet, J.V., García-Andrade, J., Angulo, C., Neumetzler, L., Persson, S., and Vera, P. (2015). Novel Disease Susceptibility Factors for Fungal Necrotrophic Pathogens in Arabidopsis. PLOS Pathogens **11**:e1004800. doi:10.1371/journal.ppat.1004800.

Douchkov, D., Lück, S., Johrde, A., Nowara, D., Himmelbach, A., Rajaraman, J., Stein, N., Sharma, R., Kilian, B., and Schweizer, P. (2014). Discovery of genes affecting resistance of barley to adapted and non-adapted powdery mildew fungi. Genome Biology **15**:518. doi:10.1186/s13059-014-0518-8.

Draeger, C., Ndinyanka Fabrice, T., Gineau, E., Mouille, G., Kuhn, B.M., Moller, I., Abdou, M.T., Frey, B., Pauly, M., Bacic, A., et al. (2015). Arabidopsis leucine-rich repeat extensin (LRX) proteins modify cell wall composition and influence plant growth. BMC Plant Biol **15**:155. doi:10.1186/s12870-015-0548-8.

Du, C., Li, X., Chen, J., Chen, W., Li, B., Li, C., Wang, L., Li, J., Zhao, X., Lin, J., et al. (2016). Receptor kinase complex transmits RALF peptide signal to inhibit root growth in *Arabidopsis*. Proceedings of the National Academy of Sciences **113**:E8326-E8334. doi:10.1073/pnas.1609626113.

Duan, Q., Kita, D., Li, C., Cheung, A.Y., and Wu, H.-M. (2010). FERONIA receptor-like kinase regulates RHO GTPase signaling of root hair development. Proceedings of the National Academy of Sciences **107**:17821-17826. doi:10.1073/pnas.1005366107.

Dünser, K., Gupta, S., Herger, A., Feraru, M.I., Ringli, C., and Kleine-Vehn, J. (2019). Extracellular matrix sensing by FERONIA and Leucine Rich Repeat Extensins controls vacuolar expansion during cellular elongation in Arabidopsis thaliana. The EMBO Journal **38**:e100353. doi:10.15252/embj.2018100353.

Engelhardt, S., Stam, R., and Hückelhoven, R. (2018). Good Riddance? Breaking Disease Susceptibility in the Era of New Breeding Technologies. Agronomy **8**:114. doi:10.3390/agronomy8070114

Engelsdorf, T., Gigli-Bisceglia, N., Veerabagu, M., McKenna, J.F., Vaahtera, L., Augstein, F., Van der Does, D., Zipfel, C., and Hamann, T. (2018). The plant cell wall integrity maintenance and immune signaling systems cooperate to control stress responses in Arabidopsis thaliana. Science Signaling **11**:eaao3070. doi:10.1126/scisignal.aao3070.

Fabrice, T.N., Vogler, H., Draeger, C., Munglani, G., Gupta, S., Herger, A.G., Knox, P., Grossniklaus, U., and Ringli, C. (2018). LRX Proteins Play a Crucial Role in Pollen Grain and Pollen Tube Cell Wall Development. Plant Physiol **176**:1981-1992. doi:10.1104/pp.17.01374.

Falhof, J., Pedersen, Jesper T., Fuglsang, Anja T., and Palmgren, M. (2016). Plasma Membrane H+-ATPase Regulation in the Center of Plant Physiology. Molecular Plant **9**:323-337. doi:10.1016/j.molp.2015.11.002.

Felle, H.H., Herrmann, A., Hanstein, S., Hückelhoven, R., and Kogel, K.-H. (2004). Apoplastic pH Signaling in Barley Leaves Attacked by the Powdery Mildew Fungus Blumeria graminis f. sp. hordei. Molecular Plant-Microbe Interactions® **17**:118-123. doi:10.1094/mpmi.2004.17.1.118.

Fotopoulos, V., Gilbert, M.J., Pittman, J.K., Marvier, A.C., Buchanan, A.J., Sauer, N., Hall, J.L., and Williams, L.E. (2003). The monosaccharide transporter gene, AtSTP4, and the cell-wall invertase, Atbetafruct1, are induced in Arabidopsis during infection with the fungal biotroph Erysiphe cichoracearum. Plant Physiol **132**:821-829. doi:10.1104/pp.103.021428.

Galindo-Trigo, S., Blanco-Touriñán, N., DeFalco, T.A., Wells, E.S., Gray, J.E., Zipfel, C., and
Smith, L.M. (2020). *Cr*RLK1L receptor-like kinases HERK1 and ANJEA are female determinants of
pollen tube reception. EMBO reports **21**:e48466. doi:10.15252/embr.201948466.
Gao, Q., Wang, C., Xi, Y., Shao, Q., Li, L., and Luan, S. (2022). A receptor–channel trio conducts
Ca2+ signalling for pollen tube reception. Nature **607**:534-539. doi:10.1038/s41586-022-04923-
7.
Ge, Z., Bergonci, T., Zhao, Y., Zou, Y., Du, S., Liu, M.-C., Luo, X., Ruan, H., García-Valencia,
L.E., Zhong, S., et al. (2017). Arabidopsis pollen tube integrity and sperm release are regulated
by RALF-mediated signaling. Science **358**:1596-1600. doi:10.1126/science.aao3642.
Geiger, D. (2020). Plant glucose transporter structure and function. Pflügers Archiv - European
Journal of Physiology **472**:1111-1128. doi:10.1007/s00424-020-02449-3.
Glawe, D.A. (2008). The Powdery Mildews: A Review of the World's Most Familiar (Yet Poorly
Known) Plant Pathogens. Annual Review of Phytopathology **46**:27-51.
doi:10.1146/annurev.phyto.46.081407.104740.
Gonneau, M., Desprez, T., Martin, M., Doblas, V.G., Bacete, L., Miart, F., Sormani, R., Hématy,
K., Renou, J., Landrein, B., et al. (2018). Receptor Kinase THESEUS1 Is a Rapid Alkalinization
Factor 34 Receptor in Arabidopsis. Current Biology **28**:2452-2458.e2454.
doi:10.1016/j.cub.2018.05.075.
Gronnier, J., Franck, C.M., Stegmann, M., DeFalco, T.A., Abarca, A., von Arx, M., Dünser, K.,
Lin, W., Yang, Z., Kleine-Vehn, J., et al. (2022). Regulation of immune receptor kinase plasma
membrane nanoscale organization by a plant peptide hormone and its receptors. eLife
11:e74162. doi:10.7554/eLife.74162.
Guénin, S., Mareck, A., Rayon, C., Lamour, R., Assoumou Ndong, Y., Domon, J.-M., Sénéchal,
F., Fournet, F., Jamet, E., Canut, H., et al. (2011). Identification of pectin methylesterase 3 as a
basic pectin methylesterase isoform involved in adventitious rooting in Arabidopsis thaliana. New
Phytologist **192**:114-126. doi:10.1111/j.1469-8137.2011.03797.x.
Guo, H., Nolan, T.M., Song, G., Liu, S., Xie, Z., Chen, J., Schnable, P.S., Walley, J.W., and Yin,
Y. (2018). FERONIA Receptor Kinase Contributes to Plant Immunity by Suppressing Jasmonic
Acid Signaling in *Arabidopsis thaliana*. Current Biology **28**:3316-3324.e3316.
doi:10.1016/j.cub.2018.07.078.
Harholt, J., Suttangkakul, A., and Vibe Scheller, H. (2010). Biosynthesis of Pectin. Plant
Physiology **153**:384-395. doi:10.1104/pp.110.156588.
Haruta, M., Sabat, G., Stecker, K., Minkoff, B.B., and Sussman, M.R. (2014). A Peptide
Hormone and Its Receptor Protein Kinase Regulate Plant Cell Expansion. Science **343**:408-411.
doi:10.1126/science.1244454.
Haruta, M., Gaddameedi, V., Burch, H., Fernandez, D., and Sussman, M.R. (2018).
Comparison of the effects of a kinase-dead mutation of FERONIA on ovule fertilization and root
growth of Arabidopsis. FEBS Letters **592**:2395-2402. doi:10.1002/1873-3468.13157.
Haruta, M., Burch, H.L., Nelson, R.B., Barrett-Wilt, G., Kline, K.G., Mohsin, S.B., Young, J.C.,
Otegui, M.S., and Sussman, M.R. (2010). Molecular Characterization of Mutant Arabidopsis
Plants with Reduced Plasma Membrane Proton Pump Activity Journal of Biological Chemistry
285:17918-17929. doi:10.1074/jbc.M110.101733.
Hayashi, Y., Nakamura, S., Takemiya, A., Takahashi, Y., Shimazaki, K.-i., and Kinoshita, T.
(2010). Biochemical Characterization of In Vitro Phosphorylation and Dephosphorylation of the
Plasma Membrane H+-ATPase. Plant and Cell Physiology **51**:1186-1196.
doi:10.1093/pcp/pcq078.
Herger, A., Gupta, S., Kadler, G., Franck, C.M., Boisson-Dernier, A., and Ringli, C. (2020).
Overlapping functions and protein-protein interactions of LRR-extensins in Arabidopsis. PLOS
Genetics **16**:e1008847. doi:10.1371/journal.pgen.1008847.
Hocq, L., Sénéchal, F., Lefebvre, V., Lehner, A., Domon, J.-M., Mollet, J.-C., Dehors, J.,
Pageau, K., Marcelo, P., Guérineau, F., et al. (2016). Combined Experimental and
Computational Approaches Reveal Distinct pH Dependence of Pectin Methylesterase Inhibitors.
Plant Physiology **173**:1075-1093. doi:10.1104/pp.16.01790.

Hocq, L., Habrylo, O., Sénéchal, F., Voxeur, A., Pau-Roblot, C., Safran, J., Fournet, F., Bassard, S., Battu, V., Demailly, H., et al. (2023). Mutation of AtPME2, a pH-Dependent Pectin Methylesterase, Affects Cell Wall Structure and Hypocotyl Elongation. Plant and Cell Physiology **65**:301-318. doi:10.1093/pcp/pcad154.

Hok, S., Danchin, E.G.J., Allasia, V., Panabières, F., Attard, A., and Keller, H. (2011). An Arabidopsis (malectin-like) leucine-rich repeat receptor-like kinase contributes to downy mildew disease. Plant, Cell & Environment **34**:1944-1957. doi:10.1111/j.1365-3040.2011.02390.x.

Hok, S., Allasia, V., Andrio, E., Naessens, E., Ribes, E., Panabières, F., Attard, A., Ris, N., Clément, M., Barlet, X., et al. (2014). The Receptor Kinase IMPAIRED OOMYCETE SUSCEPTIBILITY1 Attenuates Abscisic Acid Responses in Arabidopsis   Plant Physiology **166**:1506-1518. doi:10.1104/pp.114.248518.

Huang, Y., Yin, C., Liu, J., Feng, B., Ge, D., Kong, L., Ortiz-Morea, F.A., Richter, J., Hauser, M.-T., Wang, W.-M., et al. (2020). A trimeric CrRLK1L-LLG1 complex genetically modulates SUMM2-mediated autoimmunity. Nature Communications **11**:4859. doi:10.1038/s41467-020-18600-8.

Hückelhoven, R., and Panstruga, R. (2011). Cell biology of the plant–powdery mildew interaction. Current Opinion in Plant Biology **14**:738-746. doi:10.1016/j.pbi.2011.08.002.

Huebbers, J.W., Caldarescu, G.A., Kubátová, Z., Sabol, P., Levecque, S.C.J., Kuhn, H., Kulich, I., Reinstädler, A., Büttgen, K., Manga-Robles, A., et al. (2024). Interplay of EXO70 and MLO proteins modulates trichome cell wall composition and susceptibility to powdery mildew. The Plant Cell **36**:1007-1035. doi:10.1093/plcell/koad319.

Kadota, Y., Sklenar, J., Derbyshire, P., Stransfeld, L., Asai, S., Ntoukakis, V., Jones, Jonathan D., Shirasu, K., Menke, F., Jones, A., et al. (2014). Direct Regulation of the NADPH Oxidase RBOHD by the PRR-Associated Kinase BIK1 during Plant Immunity. Molecular Cell **54**:43-55. doi:10.1016/j.molcel.2014.02.021.

Kessler, S.A., Shimosato-Asano, H., Keinath, N.F., Wuest, S.E., Ingram, G., Panstruga, R., and Grossniklaus, U. (2010). Conserved Molecular Components for Pollen Tube Reception and Fungal Invasion. Science **330**:968-971. doi:10.1126/science.1195211.

Kesten, C., Gámez-Arjona, F.M., Menna, A., Scholl, S., Dora, S., Huerta, A.I., Huang, H.Y., Tintor, N., Kinoshita, T., Rep, M., et al. (2019). Pathogen induced pH changes regulate the growth defense balance in plants. The EMBO Journal **38**:e101822. doi:10.15252/embj.2019101822.

Kusch, S., and Panstruga, R. (2017). mlo-Based Resistance: An Apparently Universal "Weapon" to Defeat Powdery Mildew Disease. Molecular Plant-Microbe Interactions® **30**:179-189. doi:10.1094/mpmi-12-16-0255-cr.

Lan, Z., Song, Z., Wang, Z., Li, L., Liu, Y., Zhi, S., Wang, R., Wang, J., Li, Q., Bleckmann, A., et al. (2023). Antagonistic RALF peptides control an intergeneric hybridization barrier on Brassicaceae stigmas. Cell **186**:4773-4787.e4712. doi:10.1016/j.cell.2023.09.003.

Levesque-Tremblay, G., Pelloux, J., Braybrook, S.A., and Müller, K. (2015). Tuning of pectin methylesterification: consequences for cell wall biomechanics and development. Planta **242**:791-811. doi:10.1007/s00425-015-2358-5.

Lewis, K.C., Selzer, T., Shahar, C., Udi, Y., Tworowski, D., and Sagi, I. (2008). Inhibition of pectin methyl esterase activity by green tea catechins. Phytochemistry **69**:2586-2592. doi:10.1016/j.phytochem.2008.08.012.

Li, C., Yeh, F.-L., Cheung, A.Y., Duan, Q., Kita, D., Liu, M.-C., Maman, J., Luu, E.J., Wu, B.W., Gates, L., et al. (2015). Glycosylphosphatidylinositol-anchored proteins as chaperones and co-receptors for FERONIA receptor kinase signaling in Arabidopsis. eLife **4**:e06587. doi:10.7554/eLife.06587.

Li, L., Chen, H., Alotaibi, S.S., Pěnčík, A., Adamowski, M., Novák, O., and Friml, J. (2022a). RALF1 peptide triggers biphasic root growth inhibition upstream of auxin biosynthesis. Proceedings of the National Academy of Sciences **119**:e2121058119. doi:10.1073/pnas.2121058119.

Li, S., Lin, D., Zhang, Y., Deng, M., Chen, Y., Lv, B., Li, B., Lei, Y., Wang, Y., Zhao, L., et al. (2022b). Genome-edited powdery mildew resistance in wheat without growth penalties. Nature **602**:455-460. doi:10.1038/s41586-022-04395-9.

Lin, W., Tang, W., Pan, X., Huang, A., Gao, X., Anderson, C.T., and Yang, Z. (2022). *Arabidopsis* pavement cell morphogenesis requires FERONIA binding to pectin for activation of ROP GTPase signaling. Current Biology **32**:497-507.e494. doi:10.1016/j.cub.2021.11.030.

Lin, W., Zhou, X., Tang, W., Takahashi, K., Pan, X., Dai, J., Ren, H., Zhu, X., Pan, S., Zheng, H., et al. (2021). TMK-based cell-surface auxin signalling activates cell-wall acidification. Nature **599**:278-282. doi:10.1038/s41586-021-03976-4.

Liu, C., Yu, H., Voxeur, A., Rao, X., and Dixon, R.A. (2023). FERONIA and wall-associated kinases coordinate defense induced by lignin modification in plant cell walls. Science Advances **9**:eadf7714. doi:10.1126/sciadv.adf7714.

Liu, J., Liu, M., Tan, L., Huai, B., Ma, X., Pan, Q., Zheng, P., Wen, Y., Zhang, Q., Zhao, Q., et al. (2021). AtSTP8, an endoplasmic reticulum-localised monosaccharide transporter from Arabidopsis, is recruited to the extrahaustorial membrane during powdery mildew infection. New Phytologist **230**:2404-2419. doi:10.1111/nph.17347.

Liu, J., Huang, Y., Kong, L., Yu, X., Feng, B., Liu, D., Zhao, B., Mendes, G.C., Yuan, P., Ge, D., et al. (2020). The malectin-like receptor-like kinase LETUM1 modulates NLR protein SUMM2 activation via MEKK2 scaffolding. Nature Plants **6**:1106-1115. doi:10.1038/s41477-020-0748-6.

Liu, M.-C.J., Yeh, F.-L.J., Yvon, R., Simpson, K., Jordan, S., Chambers, J., Wu, H.-M., and Cheung, A.Y. (2024). Extracellular pectin-RALF phase separation mediates FERONIA global signaling function. Cell **187**:312-330.e322. doi:10.1016/j.cell.2023.11.038.

Liu, S., Bartnikas, L.M., Volko, S.M., Ausubel, F.M., and Tang, D. (2016). Mutation of the Glucosinolate Biosynthesis Enzyme Cytochrome P450 83A1 Monooxygenase Increases Camalexin Accumulation and Powdery Mildew Resistance. Frontiers in Plant Science **7**doi:10.3389/fpls.2016.00227.

Liu, Z., Chu, X., Ren, W., Cheng, L., Liu, C., Wang, C., Gao, S., Dai, S., and Li, C. (2025). PCP-B peptides and CrRLK1L receptor kinases control pollination via pH gating of aquaporins in Arabidopsis. Developmental Cell doi:10.1016/j.devcel.2024.12.026.

Mang, H., Feng, B., Hu, Z., Boisson-Dernier, A., Franck, C.M., Meng, X., Huang, Y., Zhou, J., Xu, G., Wang, T., et al. (2017). Differential Regulation of Two-Tiered Plant Immunity and Sexual Reproduction by ANXUR Receptor-Like Kinases. The Plant Cell **29**:3140-3156. doi:10.1105/tpc.17.00464.

Manning, V.A., Pandelova, I., Dhillon, B., Wilhelm, L.J., Goodwin, S.B., Berlin, A.M., Figueroa, M., Freitag, M., Hane, J.K., Henrissat, B., et al. (2013). Comparative Genomics of a Plant-Pathogenic Fungus, Pyrenophora tritici-repentis, Reveals Transduplication and the Impact of Repeat Elements on Pathogenicity and Population Divergence. G3 Genes|Genomes|Genetics **3**:41-63. doi:10.1534/g3.112.004044.

Masachis, S., Segorbe, D., Turrà, D., Leon-Ruiz, M., Fürst, U., El Ghalid, M., Leonard, G., López-Berges, M.S., Richards, T.A., Felix, G., et al. (2016). A fungal pathogen secretes plant alkalinizing peptides to increase infection. Nature Microbiology **1**:16043. doi:10.1038/nmicrobiol.2016.43.

Mecchia, M.A., Santos-Fernandez, G., Duss, N.N., Somoza, S.C., Boisson-Dernier, A., Gagliardini, V., Martínez-Bernardini, A., Fabrice, T.N., Ringli, C., Muschietti, J.P., et al. (2017). RALF4/19 peptides interact with LRX proteins to control pollen tube growth in Arabidopsis. Science **358**:1600-1603. doi:10.1126/science.aao5467.

Merlot, S., Leonhardt, N., Fenzi, F., Valon, C., Costa, M., Piette, L., Vavasseur, A., Genty, B., Boivin, K., Müller, A., et al. (2007). Constitutive activation of a plasma membrane H+ATPase prevents abscisic acid-mediated stomatal closure. The EMBO Journal **26**:3216-3226. doi:10.1038/sj.emboj.7601750.

Micali, C., Göllner, K., Humphry, M., Consonni, C., and Panstruga, R. (2008). The Powdery Mildew Disease of Arabidopsis: A Paradigm for the Interaction between Plants and Biotrophic Fungi. Arabidopsis Book **6**:e0115. doi:10.1199/tab.0115.

Micali, C.O., Neumann, U., Grunewald, D., Panstruga, R., and O'Connell, R. (2011). Biogenesis of a specialized plant–fungal interface during host cell internalization of Golovinomyces orontii haustoria. Cellular Microbiology **13**:210-226. doi:10.1111/j.1462-5822.2010.01530.x.

Milne, R.J., Dibley, K.E., Schnippenkoetter, W., Mascher, M., Lui, A.C.W., Wang, L., Lo, C., Ashton, A.R., Ryan, P.R., and Lagudah, E.S. (2019). The Wheat Lr67 Gene from the Sugar Transport Protein 13 Family Confers Multipathogen Resistance in Barley. Plant Physiol **179**:1285-1297. doi:10.1104/pp.18.00945.

Moussu, S., Broyart, C., Santos-Fernandez, G., Augustin, S., Wehrle, S., Grossniklaus, U., and Santiago, J. (2020). Structural basis for recognition of RALF peptides by LRX proteins during pollen tube growth. Proceedings of the National Academy of Sciences **117**:7494-7503. doi:10.1073/pnas.2000100117.

Moussu, S., Lee, H.K., Haas, K.T., Broyart, C., Rathgeb, U., De Bellis, D., Levasseur, T., Schoenaers, S., Fernandez, G.S., Grossniklaus, U., et al. (2023). Plant cell wall patterning and expansion mediated by protein-peptide-polysaccharide interaction. Science **382**:719-725. doi:10.1126/science.adi4720.

Mucha, S., Heinzlmeir, S., Kriechbaumer, V., Strickland, B., Kirchhelle, C., Choudhary, M., Kowalski, N., Eichmann, R., Hückelhoven, R., Grill, E., et al. (2019). The Formation of a Camalexin Biosynthetic Metabolon. The Plant Cell **31**:2697-2710. doi:10.1105/tpc.19.00403.

Müller, M.C., Praz, C.R., Sotiropoulos, A.G., Menardo, F., Kunz, L., Schudel, S., Oberhänsli, S., Poretti, M., Wehrli, A., Bourras, S., et al. (2019). A chromosome-scale genome assembly reveals a highly dynamic effector repertoire of wheat powdery mildew. New Phytologist **221**:2176-2189. doi:10.1111/nph.15529.

Nawrath, C., and Métraux, J.-P. (1999). Salicylic Acid Induction–Deficient Mutants of Arabidopsis Express PR-2 and PR-5 and Accumulate High Levels of Camalexin after Pathogen Inoculation. The Plant Cell **11**:1393-1404. doi:10.1105/tpc.11.8.1393.

Nekrasov, V., Wang, C., Win, J., Lanz, C., Weigel, D., and Kamoun, S. (2017). Rapid generation of a transgene-free powdery mildew resistant tomato by genome deletion. Scientific Reports **7**:482. doi:10.1038/s41598-017-00578-x.

Ngou, B.P.M., Jones, J.D.G., and Ding, P. (2022). Plant immune networks. Trends in Plant Science **27**:255-273. doi:10.1016/j.tplants.2021.08.012.

Nishimura, M.T., Stein, M., Hou, B.-H., Vogel, J.P., Edwards, H., and Somerville, S.C. (2003). Loss of a Callose Synthase Results in Salicylic Acid-Dependent Disease Resistance. Science **301**:969-972. doi:10.1126/science.1086716.

Noble, J.A., Bielski, N.V., Liu, M.-C.J., DeFalco, T.A., Stegmann, M., Nelson, A.D.L., McNamara, K., Sullivan, B., Dinh, K.K., Khuu, N., et al. (2022). Evolutionary analysis of the LORELEI gene family in plants reveals regulatory subfunctionalization. Plant Physiology **190**:2539-2556. doi:10.1093/plphys/kiac444.

Pearce, G., Yamaguchi, Y., Munske, G., and Ryan, C.A. (2010). Structure–activity studies of RALF, Rapid Alkalinization Factor, reveal an essential – YISY – motif. Peptides **31**:1973-1977. doi:10.1016/j.peptides.2010.08.012.

Peaucelle, A., Louvet, R., Johansen, J.N., Höfte, H., Laufs, P., Pelloux, J., and Mouille, G. (2008). Arabidopsis Phyllotaxis Is Controlled by the Methyl-Esterification Status of Cell-Wall Pectins. Current Biology **18**:1943-1948. doi:10.1016/j.cub.2008.10.065.

Pelloux, J., Rustérucci, C., and Mellerowicz, E.J. (2007). New insights into pectin methylesterase structure and function. Trends in Plant Science **12**:267-277. doi:10.1016/j.tplants.2007.04.001.

Peng, Y., Yang, J., Li, X., and Zhang, Y. (2021). Salicylic Acid: Biosynthesis and Signaling. Annual Review of Plant Biology **72**:761-791. doi:10.1146/annurev-arplant-081320-092855.

Polonio, Á., Pérez-García, A., Martínez-Cruz, J., Fernández-Ortuño, D., and de Vicente, A. (2021). The Haustorium of Phytopathogenic Fungi: A Short Overview of a Specialized Cell of Obligate Biotrophic Plant Parasites. In Progress in Botany Vol. 82, F.M. Cánovas and U. Lüttge and

M.-C. Risueño and H. Pretzsch, eds. (Springer International Publishing: Cham), pp. 337-355.
doi:10.1007/124_2020_45.

Ried, M.K., Banhara, A., Hwu, F.-Y., Binder, A., Gust, A.A., Höfle, C., Hückelhoven, R.,
Nürnberger, T., and Parniske, M. (2019). A set of Arabidopsis genes involved in the
accommodation of the downy mildew pathogen Hyaloperonospora arabidopsidis. PLOS
Pathogens **15**:e1007747. doi:10.1371/journal.ppat.1007747.

Rößling, A.-K., Dünser, K., Liu, C., Lauw, S., Rodriguez-Franco, M., Kalmbach, L., Barbez, E.,
and Kleine-Vehn, J. (2024). Pectin methylesterase activity is required for RALF1 peptide
signalling output. eLife **13**:RP96943. doi:10.7554/eLife.96943.

Rzemieniewski, J., and Stegmann, M. (2022). Regulation of pattern-triggered immunity and
growth by phytocytokines. Current Opinion in Plant Biology **68**:102230.
doi:10.1016/j.pbi.2022.102230.

Schmid, M., Davison, T.S., Henz, S.R., Pape, U.J., Demar, M., Vingron, M., Schölkopf, B.,
Weigel, D., and Lohmann, J.U. (2005). A gene expression map of Arabidopsis thaliana
development. Nat Genet **37**:501-506. doi:10.1038/ng1543.

Schnepf, V., Vlot, A.C., Kugler, K., and Hückelhoven, R. (2018). Barley susceptibility factor
RACB modulates transcript levels of signalling protein genes in compatible interaction with
Blumeria graminis f.sp. hordei. Molecular Plant Pathology **19**:393-404. doi:10.1111/mpp.12531.

Schoenaers, S., Lee, H.K., Gonneau, M., Faucher, E., Levasseur, T., Akary, E., Claeijs, N.,
Moussu, S., Broyart, C., Balcerowicz, D., et al. (2024). Rapid alkalinization factor 22 has a
structural and signalling role in root hair cell wall assembly. Nature Plants **10**:494-511.
doi:10.1038/s41477-024-01637-8.

Sénéchal, F., Habrylo, O., Hocq, L., Domon, J.-M., Marcelo, P., Lefebvre, V., Pelloux, J., and
Mercadante, D. (2017). Structural and dynamical characterization of the pH-dependence of the
pectin methylesterase-pectin methylesterase inhibitor complex. Journal of Biological Chemistry
292:21538-21547. doi:10.1074/jbc.RA117.000197.

Shen, Q., Bourdais, G., Pan, H., Robatzek, S., and Tang, D. (2017). *Arabidopsis*
glycosylphosphatidylinositol-anchored protein LLG1 associates with and modulates FLS2 to
regulate innate immunity. Proceedings of the National Academy of Sciences **114**:5749-5754.
doi:10.1073/pnas.1614468114.

Shevchenko, A., Tomas, H., Havli, J., Olsen, J.V., and Mann, M. (2006). In-gel digestion for mass
spectrometric characterization of proteins and proteomes. Nature Protocols **1**:2856-2860.
doi:10.1038/nprot.2006.468.

Stegmann, M., Monaghan, J., Smakowska-Luzan, E., Rovenich, H., Lehner, A., Holton, N.,
Belkhadir, Y., and Zipfel, C. (2017). The receptor kinase FER is a RALF-regulated scaffold
controlling plant immune signaling. Science **355**:287-289. doi:10.1126/science.aal2541.

Stein, M.n., Dittgen, J., Sánchez-Rodríguez, C., Hou, B.-H., Molina, A., Schulze-Lefert, P.,
Lipka, V., and Somerville, S. (2006). Arabidopsis PEN3/PDR8, an ATP Binding Cassette
Transporter, Contributes to Nonhost Resistance to Inappropriate Pathogens That Enter by Direct
Penetration. The Plant Cell **18**:731-746. doi:10.1105/tpc.105.038372.

Takahashi, K., Hayashi, K.-i., and Kinoshita, T. (2012). Auxin Activates the Plasma Membrane
H+-ATPase by Phosphorylation during Hypocotyl Elongation in Arabidopsis   Plant Physiology
159:632-641. doi:10.1104/pp.112.196428.

Tang, J., Wu, D., Li, X., Wang, L., Xu, L., Zhang, Y., Xu, F., Liu, H., Xie, Q., Dai, S., et al. (2022).
Plant immunity suppression via PHR1-RALF-FERONIA shapes the root microbiome to alleviate
phosphate starvation. The EMBO Journal **41**:e109102. doi:10.15252/embj.2021109102.

Thynne, E., Saur, I.M.L., Simbaqueba, J., Ogilvie, H.A., Gonzalez-Cendales, Y., Mead, O.,
Taranto, A., Catanzariti, A.-M., McDonald, M.C., Schwessinger, B., et al. (2017). Fungal
phytopathogens encode functional homologues of plant rapid alkalinization factor (RALF)
peptides. Molecular Plant Pathology **18**:811-824. doi:10.1111/mpp.12444.

Tyanova, S., Temu, T., and Cox, J. (2016). The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. Nature Protocols **11**:2301-2319.
doi:10.1038/nprot.2016.136.

van Schie, C.C.N., and Takken, F.L.W. (2014). Susceptibility Genes 101: How to Be a Good Host.
Annual Review of Phytopathology **52**:551-581. doi:10.1146/annurev-phyto-102313-045854.
Voegele, R.T., and Mendgen, K.W. (2011). Nutrient uptake in rust fungi: how sweet is parasitic
life? Euphytica **179**:41-55. doi:10.1007/s10681-011-0358-5.
Vogel, J.P., Raab, T.K., Schiff, C., and Somerville, S.C. (2002). PMR6, a Pectate Lyase–Like
Gene Required for Powdery Mildew Susceptibility in Arabidopsis. The Plant Cell **14**:2095-2106.
doi:10.1105/tpc.003509.
Weis, C., Pfeilmeier, S., Glawischnig, E., Isono, E., Pachl, F., Hahne, H., Kuster, B.,
Eichmann, R., and Hückelhoven, R. (2013). Co-immunoprecipitation-based identification of
putative BAX INHIBITOR-1-interacting proteins involved in cell death regulation and plant–
powdery mildew interactions. Molecular Plant Pathology **14**:791-802. doi:10.1111/mpp.12050.
Wiese, J., Kranz, T., and Schubert, S. (2004). Induction of Pathogen Resistance in Barley by
Abiotic Stress. Plant Biol (Stuttg) **6**:529-536. doi:10.1055/s-2004-821176.
Willats, W.G.T., Orfila, C., Limberg, G., Buchholt, H.C., van Alebeek, G.-J.W.M., Voragen,
A.G., Marcus, S.E., Christensen, T.M.I.E., Mikkelsen, J.D., Murray, B.S., et al. (2001).
Modulation of the Degree and Pattern of Methyl-esterification of Pectic Homogalacturonan in
Plant Cell Walls: IMPLICATIONS FOR PECTIN METHYL ESTERASE ACTION, MATRIX PROPERTIES,
AND CELL ADHESION Journal of Biological Chemistry **276**:19404-19413.
doi:10.1074/jbc.M011242200.
Wormit, A., and Usadel, B. (2018). The Multifaceted Role of Pectin Methylesterase Inhibitors
(PMEIs). International Journal of Molecular Sciences **19**:2878.
Wu, Y., Ma, X., Pan, Z., Kale, S.D., Song, Y., King, H., Zhang, Q., Presley, C., Deng, X., Wei, C.-
I., et al. (2018). Comparative genome analyses reveal sequence features reflecting distinct
modes of host-adaptation between dicot and monocot powdery mildew. BMC Genomics **19**:705.
doi:10.1186/s12864-018-5069-z.
Xiao, Y., Stegmann, M., Han, Z., DeFalco, T.A., Parys, K., Xu, L., Belkhadir, Y., Zipfel, C., and
Chai, J. (2019). Mechanisms of RALF peptide perception by a heterotypic receptor complex.
Nature **572**:270-274. doi:10.1038/s41586-019-1409-7.
Xu, F., Gonneau, M., Faucher, E., Habrylo, O., Lefebvre, V., Domon, J.-M., Martin, M.,
Sénéchal, F., Peaucelle, A., Pelloux, J., et al. (2022). Biochemical characterization of Pectin
Methylesterase Inhibitor 3 from Arabidopsis thaliana. The Cell Surface **8**:100080.
doi:10.1016/j.tcsw.2022.100080.
Xu, X., and Chen, X. (2023). A complete genome sequence of Podosphaera xanthii race 2 F, the
causal agent of powdery mildew isolated from cucumber in China. figshare.
Yamada, K., Saijo, Y., Nakagami, H., and Takano, Y. (2016). Regulation of sugar transporter
activity for antibacterial defense in Arabidopsis. Science **354**:1427-1430.
doi:10.1126/science.aah5692.
Yang, T., Wang, L., Li, C., Liu, Y., Zhu, S., Qi, Y., Liu, X., Lin, Q., Luan, S., and Yu, F. (2015).
Receptor protein kinase FERONIA controls leaf starch accumulation by interacting with
glyceraldehyde-3-phosphate dehydrogenase. Biochemical and Biophysical Research
Communications **465**:77-82. doi:10.1016/j.bbrc.2015.07.132.
Yeats, T.H., Sorek, H., Wemmer, D.E., and Somerville, C.R. (2016). Cellulose Deficiency Is
Enhanced on Hyper Accumulation of Sucrose by a H+-Coupled Sucrose Symporter Plant
Physiology **171**:110-124. doi:10.1104/pp.16.00302.
Zaccaron, A.Z., Neill, T., Corcoran, J., Mahaffee, W.F., and Stergiopoulos, I. (2023). A
chromosome-scale genome assembly of the grape powdery mildew pathogen *Erysiphe necator*
reveals its genomic architecture and previously unknown features of its biology. mBio **14**:e00645-
00623. doi:10.1128/mbio.00645-23.
Zhang, K., Halitschke, R., Yin, C., Liu, C.-J., and Gan, S.-S. (2013). Salicylic acid 3-hydroxylase
regulates *Arabidopsis* leaf longevity by mediating salicylic acid catabolism. Proceedings of the
National Academy of Sciences **110**:14807-14812. doi:10.1073/pnas.1302702110.

Zhang, X., Peng, H., Zhu, S., Xing, J., Li, X., Zhu, Z., Zheng, J., Wang, L., Wang, B., Chen, J., et
al. (2020). Nematode-Encoded RALF Peptide Mimics Facilitate Parasitism of Plants through the
FERONIA Receptor Kinase. Molecular Plant **13**:1434-1454. doi:10.1016/j.molp.2020.08.014.
Zhang, Y., Zhao, L., Zhao, J., Li, Y., Wang, J., Guo, R., Gan, S., Liu, C.-J., and Zhang, K. (2017).
S5H/DMR6 Encodes a Salicylic Acid 5-Hydroxylase That Fine-Tunes Salicylic Acid
Homeostasis Plant Physiology **175**:1082-1093. doi:10.1104/pp.17.00695.
Zhao, C., Zayed, O., Yu, Z., Jiang, W., Zhu, P., Hsu, C.-C., Zhang, L., Tao, W.A., Lozano-Durán,
R., and Zhu, J.-K. (2018). Leucine-rich repeat extensin proteins regulate plant salt tolerance in
Arabidopsis. Proceedings of the National Academy of Sciences **115**:13123-13128.
doi:10.1073/pnas.1816991115.
Zhong, S., Li, L., Wang, Z., Ge, Z., Li, Q., Bleckmann, A., Wang, J., Song, Z., Shi, Y., Liu, T., et
al. (2022). RALF peptide signaling controls the polytubey block in Arabidopsis. Science **375**:290-
296. doi:10.1126/science.abl4683.
Zhou, F., Andersen, C.H., Burhenne, K., Fischer, P.H., Collinge, D.B., and Thordal-
Christensen, H. (2000). Proton extrusion is an essential signalling component in the HR of
epidermal single cells in the barley-powdery mildew interaction. Plant J **23**:245-254.
doi:10.1046/j.1365-313x.2000.00777.x.
Zhu, S., Fu, Q., Xu, F., Zheng, H., and Yu, F. (2021). New paradigms in cell adaptation: decades
of discoveries on the CrRLK1L receptor kinase signalling network. New Phytologist **232**:1168-
1183. doi:10.1111/nph.17683.
Zimmerli, L., Stein, M., Lipka, V., Schulze-Lefert, P., and Somerville, S. (2004). Host and non-
host pathogens elicit different jasmonate/ethylene responses in Arabidopsis. The Plant Journal
40:633-646. doi:10.1111/j.1365-313X.2004.02236.x.

1255

1256

Figures

1258

Figure 1: FER and additional RALF binding proteins are *Ecr* susceptibility factors. A) Images of Col-0 and *fer-4* plants infected with *Ecr* (14 dpi). B) Ink stained *Ecr* colonies (5 dpi) grown on Col-0 and *fer-4* plants. Black arrows indicate conidiophores. Scale bar represents 50 µm. C) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=223-392 pooled from three independent experiments (Dunn's multiple comparisons test, a-b, b-c p<0.0001; a-c p=0.0106). D) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=345-665 pooled from five independent experiments (Dunn's multiple comparisons test, a-b, a-c p<0.0001; b-c p=0.0235). E) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=285-453 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p=0.0150; a-c p<0.0001; b-c p=0.0408). F) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=113-159 pooled from two independent experiments (Dunn's multiple comparisons test, a-b p<0.0001). All experiments were done at least three times with similar results, except (F), which was repeated once with identical results.

1273

Figure 2: Specific RALFs are powdery mildew susceptibility factors. A) Table of RALF peptides mutated in CRISPR *ralf3x*, CRISPR *ralf7x* and *ralf1/22/23/31* B) Images of Col-0, *fer-4*, CRISPR *ralf3x* and CRISPR *ralf7x* plants infected with *Ecr* (14 dpi). C) Ink stained *Ecr* colonies (5 dpi) grown on Col-0, *fer-4*, CRISPR *ralf3x* and CRISPR *ralf7x* plants. Black arrows indicate conidiophores. Scale bar represents 50 μm. D) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=98-456 pooled from three independent experiments (Dunn's multiple comparisons test, a-b, a-c p<0.0001, b-c p<0.0002. Comparison between *fer-4* and CRISPR *ralf7x*: Mann Whitney test, ** p=0.007). E) Amount of fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=4, data points indicate independent biological replicates. F) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=101-286 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p<0.0001). G) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=226-382 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p<0.0001. Comparison between *fer-4* and *ralf1/22/23/31*: Mann Whitney test, ** p=0.0132). All experiments were done at least three times with similar results.

1294

1295

**Figure 3: FER and RALF mediated powdery mildew susceptibility is most likely not directly linked
to MLO.** A) Confocal microscopy images of fungal spores penetrating p35S::MLO2-mCherry and
pFER::FER-GFP lines (2 dpi). Dotted lines indicate fungal spores, white arrow marks the penetration
site. Scale bar represents 50 μm. B) Quantification of hyphal branching (2 dpi). Mean ± SD, n=77–98
pooled from two independent experiments (Mann Whitney test, ns: not significant). C) Penetration
efficiency of *Ecr* on the indicated genotypes (calculated as the percentage of successful penetrations
from all counted interactions, 2 dpi). Mean ± SD, n=2–4. D) Quantification of trichome numbers per leaf
area on the indicated genotypes. Mean ± SD, n=27 pooled from three independent experiments (Dunn's
multiple comparisons test, a-b p<0.0001). E) Quantification of trichome branching on the indicated
genotypes. Mean ± SD, n=27 pooled from three independent experiments (Dunn's multiple comparisons
test, statistical analysis was performed on each category individually, 2 branches: a-b p=0.0002; 3
branches: a-b p=0.0259; a-c, b-c p<0.0001; 4 branches: a-b p<0.0001). F) Microscopic images of aniline
blue-stained trichomes of Col-0, *fer-4* and CRISPR *ralf7x*. Scale bar in the upper row represents 50 μm,
scale bar in the lower panel represents 500 μm. All experiments were done three times with similar
results, except (B), which was repeated once with identical results.

1311

1312

1313

Figure 4: FER-RALF mediated powdery mildew susceptibility is independent of JA and SA signaling. A) Quantification of free (blue) and glycosylated (green) SA in untreated and *Ecr* infected (5 dpi) leaves of the indicated genotypes. Mean ± SD, n=3 pooled from three independent experiments (Tukey's multiple comparisons test, free and glycosylated SA was analysed individually, free: a-bc p=0.0423; a-c p=0.0006; ab-c p≤0.0013, glycosylated: A-C p≤0.0029; A-BC p≤0.0383) B) Quantification of free (blue) and glycosylated (green) 2,5-DHBA in untreated and *Ecr* infected (5 dpi) leaves of the indicated genotypes. Mean ± SD, n=3 pooled from three independent experiments (Tukey's multiple comparisons test, all comparisons were not significant) C) Quantification of free (blue) Camalexin in untreated and *Ecr* infected (5 dpi) leaves of the indicated genotypes. Mean ± SD, n=3 pooled from three independent experiments (Tukey's multiple comparisons test, a-bc p=0.0486; ab-c p≤0.0319; a-c p=0.0137. D) RT-qPCR of *PR1* in untreated (grey) adult leaves and upon infection with *Ecr* (5 dpi, green). Housekeeping gene *UBQ5*. Mean ± SD, n=3, data points indicate independent biological replicates. E) RT-qPCR of *JAZ10* in untreated (grey) adult leaves and upon infection with *Ecr* (5 dpi, green). Housekeeping gene *UBQ5*. Mean ± SD, n=3, data points indicate independent biological replicates. F) RT-qPCR of *PDF1.2* in untreated (grey) adult leaves and upon infection with *Ecr* (5 dpi, green). Housekeeping gene *UBQ5*. Mean ± SD, n=3, data points indicate independent biological replicates. G) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=152-209 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p<0.0001). All experiments were done three times with similar results.

1333

Figure 5: Apoplastic pH modulation and PME activity determine powdery mildew infection success. A) MS Intensities measured after Co-immunoprecipitation with untreated pFER::FER-GFP and p35S::Lti6b-GFP plants using GFP-TRAP beads. n=3 for pFER::FER-GFP, n=4 for p35S::Lti6b-GFP, data points indicate independent biological replicates. FDR values indicate statistical differences between pFER::FER-GFP and p35S::Lti6b-GFP. No AHA3 derived peptides were detected in p35S::Lti6b-GFP samples. B) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=49-91 pooled from two independent experiments (Mann Whitney test, ****p<0.0001). C) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of leaves pretreated with H_2O (blue) or Fusicoccin (green). Mean ± SD, n=71-113 pooled from four independent experiments (Dunn's multiple comparisons test, a-b p≤0.0008). D) Amount of fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection and pretreatment with H_2O (blue) or Fusicoccin (green). Mean ± SD, n=3, data points indicate independent biological replicates. E) Conidiophores per field of view (5 dpi) after pretreatment with H_2O (blue), $MgCl_2$ 15 mM pH 5.8 (grey), MES buffer 10 mM pH 5.8 (green) and MES buffer 10 mM pH 7.5 (red) before *Ecr* infection. Mean ± SD, n=52-347 pooled from four independent experiments (Dunn's multiple comparisons test, a-b p<0,005, A-B p=0.0381, B-C p=0.0015, A-C p<0.0001). F) Ink stained *Ecr* colonies (5 dpi) grown on Col-0 and *fer-4* plants after pretreatment with 15 mM $MgCl_2$, 10 mM MES buffer (pH 5.8) and 10 mM MES buffer (pH 7.5). Black arrows indicate conidiophores. Scale bar represents 50 μm. F) PME activity in untreated leaves (grey) and upon *Ecr* infection (5 dpi, green) normalized to Col-0 PME activity in untreated leaves. Mean ± SD, n=3, data points indicate independent biological replicates. G) Amount of fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection and pretreatment with H_2O (blue) or EGCG (red). Mean ± SD, n=3, data points indicate independent biological replicates. H) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=182-309 pooled from four independent experiments (Dunn's multiple comparisons test, a-b, b-c, a-c p<0.0001). I) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=161-338 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p<0.0001). All experiments were done at least three times with similar results.

1362

Figure 6: RALF-mediated powdery mildew susceptibility is partially independent of FER and FER signaling. A) Confocal images of pUBQ10::SYP122-pHusion lines in the Col-0 and CRISPR *fer* background (2 dpi). Dotted lines indicate the location of the fungal spore. Scale bar represents 50 µm. B) Quantification of the EGFP/mRFP1 ratio near the penetration site (green) and in uninfected areas (grey). Mean ± SD, n=24 pooled from three independent experiments (Mann Whitney test, a-b p<0.0001). C) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=95-137 pooled from three independent experiments (Dunn's multiple comparisons test, a-b p=0.0277, a-c p<0.0001). D) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=131-210 pooled from four independent experiments (Dunn's multiple comparisons test, a-b p<0.0001). E) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=48-189 pooled from two independent experiments (Dunn's multiple comparisons test, a-bc p≤0.0031, ab-c ≤0,0049, a-c<0.0001). F) Conidiophores per field of view (5 dpi) after pretreatment with H₂O (blue) or 1µM RALF23 before *Ecr* infection. Mean ± SD, n=485-970 pooled from 11 independent experiments (Mann Whitney test, statistical analysis was performed on each genotype individually, Col-0: ***p=0.0008, *fer-4* ** p=0.0064). G) Conidiophores per field of view (5 dpi) of fungal colonies grown upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=95 - 137 pooled from 3 independent experiments (Dunn's multiple comparisons test, a-b p=0.0076, a-c p<0.0001, b-c p=0.0017). All experiments were done at least three times with similar results.

1382

1383   **Supplemental Figures**

1384

1385   **Figure S1: Quantification of fungal growth on mutants of RALF binding proteins.** A – D) Amount
1386   of fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection of the indicated genotypes. Mean ±
1387   SD, data points indicate independent biological replicates A) n=14, B) n=7–9, C) n=3, D) n =3–4. All
1388   experiments were done at least three times with similar results.

1389

1390

Figure S2: Characterization of CRISPR *ralf* mutants. A) Expression levels of the indicated *RALF* genes in the vegetative rosette. Data was obtained using ePlant browser (https://bar.utoronto.ca/eplant/) and is based on (Schmid et al., 2005). B) Characterization of CRISPR *ralf3x*. Schematic diagram of *RALF22, RALF23* and *RALF33* gene structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. The mature RALF peptide is indicated in blue. C) Characterization of CRISPR *ralf7x*. Schematic diagram of *RALF1, RALF18, RALF22, RALF31, RALF23*, *RALF33* and *RALF34* gene structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. The mature RALF peptide is indicated in blue. D) Quantification of the petiole length of 5 week-old Col-0, *fer-4*, CRISPR *ralf3x* and CRISPR *ralf7x* plants. Each data point represents the average length of petioles on a fully grown rosette. Mean ± SD, n=7 pooled from two independent experiments (Tukey's multiple comparisons test, a-b p<0.0001). E – F) RT-qPCR of *RALF23* in adult leaves of the indicated plant lines. Housekeeping gene *UBQ5*. Mean ± SD, n=2, data points indicate independent biological replicates. E) – F) were repeated once with identical results.

1404

1405

Figure S3: Characterization of *fer*-4 CRISPR *myc2*. A) Quantification of free (blue) and glycosylated (green) 2,3-DHBA in untreated and *Ecr* infected (5 dpi) leaves of the indicated genotypes. Mean ± SD, n=3 pooled from three independent experiments (Tukey's multiple comparisons test, glycosylated: a-b p≤0.0324). (B) Characterization of *fer*-4 CRISPR *myc2*. Schematic diagram of *MYC2* gene structure and the CRISPR-Cas9-mediated mutation pattern detected by DNA sequencing. C) RT-qPCR of *MYC2* in untreated (grey) adult leaves and upon infection with *Ecr* (5 dpi, green). Housekeeping gene *UBQ5*. Mean ± SD, n=3, data points indicate independent biological replicates. (D) Images of 4 week-old Col-0, *fer*-4 and *fer*-4 CRISPR *myc2* plants. (D) Amount of fungal DNA normalized to plant DNA (5 dpi) upon *Ecr* infection of the indicated genotypes. Mean ± SD, n=3, data points indicate independent biological replicates. A), C) and E) were done three times with similar results.

1416

1417

Figure S4: FER interactors detected by Co-IP MS/MS. (A) Vulcano plot of detected FER-GFP interactors. Significantly enriched proteins (FDR=0.05 and fold change (FER-GFP/Lti6b-GFP) >2) are marked with colored circles according to the legend. Non-significant proteins are colored grey. (B) List of significant FER-GFP interactors compared to Lti6b-GFP control. Data was obtained from biological triplicates for FER-GFP and biological quadruplets for Lti6b-GFP.

1423

B

Identifier	Molecular function	FDR	log10(fold change)
AT1G32900.1 |GBSS1	Glycosyltransferase	0,0062	-1,223
AT2G36830.1 |TIP1;1	Water channel	0,0007	-0,459
AT3G59770.2 |SAC9	Phosphatase activity	0,0114	0,425
AT4G02030.1 |UNH	Protein binding	0,0062	0,668
AT5G35620.3 |EIF4E2	Translation initiation factor	0,0150	-0,354
AT3G55620.1 |eIF6A	Translation initiation factor	0,0142	0,330

1424

1425 **Figure S5: Powdery mildew dependent FER interactors detected by Co-IP MS/MS.** (A) Vulcano plot
1426 of detected FER-GFP interactors after *Ecr* infection (1 dpi). Significantly enriched proteins (FDR=0.05
1427 and fold change (*Ecr* infection (1dpi)/Untreated) >2) are marked in black. Non-significant proteins are
1428 colored grey. (B) List of significant FER-GFP interactors at 1 dpi with *Ecr*, compared to the untreated
1429 FER-GFP control. Data was obtained from biological triplicates for each treatment.

1430

1431

Figure S6: AHA-Thr947 phosphorylation status during powdery mildew infection. (A)
Quantification of the band intensities detected in western blot probed with α-AHA and α-Thr^{947P}
antibodies. Band intensities were measured using Image J software. Data points indicate individual
experiments. Data points from the same experiment are marked with the same color. (B) AHA
phosphorylation upon infection with *Ecr* (3 dpi). Western blots were probed with α-AHA and α-Thr^{947P}
antibodies. CBB = Coomassie brilliant blue. The experiment was repeated at least three times with
similar results.

1439

1440

Figure S7: Phosphorylation changes on FER-GFP upon *Ecr* infection and characterization of
RALF23-GFP overexpression in Col-0 and *fer-2*. (A) MS Intensities of phosphorylated FER peptides
after GFP co-immunoprecipitation of FER-GFP (Untreated and 1 dpi). Mean ± SD, n=3, missing data
points occur from missed identification in the LC-MS/MS measurement. (B) RT-qPCR of *RALF23* in
untreated adult leaves of the indicated plant lines. Housekeeping gene *UBQ5*. n=3, data points indicate
independent biological replicates. A) The data was obtained from biological triplicates for both
treatments. B) The experiment was done three times with similar results.

1448

1449

1450 **Figure S8: Images of all plant lines used in this study.** (A) Representative images of 5 week-old

1451 plants used in this study.

1452   **Tables**

1453   **Table S1: Identification of RALF-like peptides from phytopathogenic fungi** [a]species reported to be
1454   able to infect *Arabidopsis thaliana*.

Species	Significant hits in predicted proteins (E-values < 10e[-05])	Significant hits in translated genome assembly (E-values < 10e[-05])	GenBank assembly accession number	Reference
Fusarium oxysporum NRRL 32931	1	2	GCA_000271745.2	(DeIulio et al., 2018)
Pyrenophora tritici-repentis Pt-1C-BFP	1	1	GCA_000149985.1	(Manning et al., 2013)
Golovinomyces cichoracearum[a] UMSG1	0	0	GCA_003611235.1	(Wu et al., 2018)
Erysiphe neolycopersicum[a] UMSG2	0	0	GCA_003610855	(Wu et al., 2018)
Erysiphe necator EnFRAME01	0	0	GCA_024703715.1	(Zaccaron et al., 2023)
Blumeria graminis f.sp. *tritici* CHE_96224	0	0	GCA_900519115.1	(Müller et al., 2019)
Podosphera xanthii YZU573	0	0	GCA_028751805.1	(Xu and Chen, 2023)

1455

1456

1457 **Table S2: Plant lines used in this study.**

Plant line	Genetic Backgorund	Reference
fer-4	Col-0	(Duan et al., 2010)
fer-2	Col-0	(Deslauriers and Larsen, 2010)
pFER::FER-GFP	*fer-4*	(Duan et al., 2010)
llg1-2	Col-0	(Li et al., 2015)
llg1-3	Col-0	(Shen et al., 2017)
lrx1/2/3	Col-0	(Draeger et al., 2015)
lrx1/2/3/4/5	Col-0	(Herger et al., 2020)
p35S::LRR4-citrine	Col-0	(Fabrice et al., 2018)
ralf1/22/23/31	Col-0	(Lan et al., 2023)
CRISPR *ralf3x*	Col-0	Generated in this study
CRISPR *ralf7x*	Col-0	Generated in this study
p35S::RALF23	CRISPR *ralf7x*	Generated in this study
p35S::RALF23$^{Y93A/Y96A}$	CRISPR *ralf7x*	Generated in this study
p35S::RALF23$^{Y113A/Y114A}$	CRISPR *ralf7x*	Generated in this study
mlo2/6/12	Col-0	(Consonni et al., 2006)
CRISPR *myc2*	*fer-4*	Generated in this study
PMEI3-OE	Col-0	(Peaucelle et al., 2008)
ost2-2D	Col-0	(Merlot et al., 2007)
pme3	Col-0	(Guénin et al., 2011)
pme17	Col-0	(Del Corpo et al., 2020)
pUBQ10::SYP122-pHusion	Col-0	(Kesten et al., 2019)
pUBQ10::SYP122-pHusion/CRISPR *fer*	pUBQ10::SYP122-pHusion	Generated in this study
pFER:FER-GFP	*fer-4*	(Chakravorty et al., 2018)
pFER:FER^{K565R}-GFP	*fer-4*	(Chakravorty et al., 2018)
p35S::RALF23-GFP	Col-0	(Dobón et al., 2015)
fer-2 x p35S::RALF23-GFP	*fer-2*	(Stegmann et al., 2017)
pFER:FER^{WT}-GFP	*fer-4*	(Xiao et al., 2019)
pFER:FER^{N303Y}-GFP	*fer-4*	(Xiao et al., 2019)
pFER:FER^{G257A}-GFP	*fer-4*	(Xiao et al., 2019)
p35S::Lti6b-GFP	Col-0	(Kadota et al., 2014)

1458

1459

1460   **Table S3: Primers used in this study**

Primers used for cloning		
Name	**Sequence (5' - 3')**	**Purpose**
RALF23_Esp3I_F	ttCGTCTCaaatgagaggactctccagaaac	p35S::RALF23
RALF23_Esp3I_R	ttCGTCTCaactatgagcgccggcagcgag	
R23^{Y113AY114A}_F	ttCGTCTCctgccgccaattgtcgacgtg	p35S::RALF23$^{Y113A/Y1114A}$
R23^{Y113AY114A}_R	ttCGTCTCcggcagatgcgccgcgacgtg	
R23^{Y93AY96A}_F	ttCGTCTCatcagcgctggtgcgctgaggag	p35S::RALF23$^{Y93A/Y96A}$
R23^{Y93AY96A}_R	ttCGTCTCgctgatgtacctcctcgtagc	
MLO2_Esp3I_F	ttCGTCTCaaatggcagatcaagtaaaag	p35S::MLO2-mCherry
MLO2_Blunt_R	tttcttaaaagaaaaatctcttaa (5'Phosphorylated)	
Primers used RT-qPCR		
Name	**Sequence (5' - 3')**	**Purpose**
Ath_Rubisco_F	TGATGGACGGTACTGGACAA	Quantification of plant DNA
Ath_Rubsico_R	GAAGCTTGGTGGCTTGTAGG	
Ecr_Tubulin_F	TGACAGCCCGGAATGAGT	Quantification of fungal DNA
Ecr_Tubulin_R	TTGTCTTCGTTTCCCAGGTC	
RALF23_qPCR_F	TTCATAACTGGTCCGTCGCC	RALF23
RALF23_qPCR_R	AGCTAATATGCGCCGGTTGA	
UBQ5_qPCR_F	ACTCCTTCCTCAAACGCTGA	UBQ5
UBQ5_qPCR_R	CCAAGCCGAAGAAGATCAAG	
MYC2_qPCR_F	GTGCGGGATTAGCTGGTAAA	MYC2
MYC2_qPCR_R	ATGCATCCCAAACACTCCTC	
PR1_qPCR_F	CGGAGCTACGCAGAACAACT	PR1
PR1_qPCR_R	CAGACAAGTCACCGCTACCC	
JAZ10_qPCR_F	GAGAAGCGCAAGGAGAGATTAG	JAZ10
JAZ10_qPCR_R	CTTAGTAGGTAACGTAATCTCC	
PDF1.2_qPCR_F	CTG CTT TCG ACG CAC CGG CA	PDF1.2
PDF1.2_qPCR_R	TGA TCC ATG TTT GGC TCC TTC	
Primers used for genotyping		
Name	**Sequence (5' - 3')**	**Purpose**
GG pUC18 Seq-F	GTATCACGAGGCCCTTTCGT	Inserts in GoldenGate pUC18
GG pUC18 Seq-R	TAATGAATCGGCCAACGC	
pGGPlant-XL-AB-insert_Fwd	AAACGACAATCTGAGCTCCACC	Insert in GoldenGate pGGPlantXL-AB
pGGPlantXL-AB-insert_Rev	CGGTGTCATCTATGTTACTAGAT	
RALF1_F	CAAAATTAAAAAATCGAGTCCAC	CRISPR RALF1
RALF1_R	GCTTGATGGATCAAAGAAAAC	
RALF14_F	GCTCTTAATCTTCGCCG	CRISPR RALF14
RALF14_R	GTAAACCGGTAACAATGTG	
RALF18_F	CATCACAAACCATTTCAACG	CRISPR RALF18
RALF18_R	CACACCAACGAAAGGAAG	
RALF22_F	GTTTCCAAGAGAGAAAGAGAG	CRISPR RALF22
RALF22_R	GAAAATCGAGAAGAGAGACAG	
RALF23_F	CTTCATAGTTCGTGCACAG	CRISPR RALF23
RALF23_R	CTAATAATCCGCAATGAACTTAG	
RALF31_F	CTTCTTCTGCTTAGAGACAG	CRISPR RALF31
RALF31_R	CAGCAATGCAAAAGCTTAATC	
RALF33_F	CCTCAGAGAAAGAAGAAGAAAG	CRISPR RALF33
RALF33_R	GCAATGCAAAGTATAAACATAG	
RALF34_F	CCTCTTTACATGTGACTTCC	CRISPR RALF34
RALF34_R	CACGGTTTACAAACAAACATC	
MYC2_F	GGAACACACGAAGGTTGGAC	CRISPR MYC2
MYC2_R	CTGCTACTTCTTTCACGACGG	
FER_F	CGTAACCAAACGAACATG	CRISPR FER
FER_R	GGTAAGTGAAACCAGAGTC	

1461

1462

1463 **Table S4: CRISPR target sites**

Target site	Sequence	Purpose
RALF1.1	ACACTCTGCTATTGAACCG	Generation of CRISPR *ralf* mutants
RALF1.2	AATTGTCAGAACGGAGCTC	
RALF14.1	GATGCATTTTCCACTCAAC	
RALF18.1	GTGGTTGCAGTGCCGCTAC	
RALF22.1	GTTCGATTCAGACATAAGC	
RALF22.2	CGTACTACAACTGCCAGCG	
RALF23.1	CCTCGTAGCTAATATGCGC	
RALF23.2	CAGCCACGAGAGTAAGGAT	
RALF31.1	TGGCGCAGAAACGGTACAT	
RALF31.2	TTTGTGTCTCTAGCGCATC	
RALF33.1	CGATCGAGTCGAAATGCAA	
RALF33.2	TAGCGCTATTACTCGTTGC	
RALF34.1	CTCTCACTCTTCTACCCGA	
RALF34.2	CCGTTCATCCGTATAGCCG	
MYC2.1	CCGCCGGAGTAGAAAACGG	Generation of *fer*-4 CRISPR *myc2*
FER.1	GACGATTAAGTATCCTACG	Generation of pUBQ10::SYP122-apopHusion CRISPR *fer*-4
FER.2	TGATCGATGAAGATCACAG	

1464