1 **Fermentative yeast diversity at the northern range limit of their oak tree hosts**
2

3 Javier Pinto[1+], Chloé Haberkorn[1+], Markus Franzén[2], Ayco J. M. Tack[3], Rike Stelkens[1*]
4
5 **Running title:** Fermentative yeast diversity in northern latitudes
6
[1]Department of Zoology, Stockholm University, Svante Arrheniusväg 18 B, 106 91 Stockholm,
Sweden;
[2]Center for Ecology and Evolution in Microbial Model Systems, Department of Biology and
Environmental Science, Linnaeus University, Kalmar, Sweden; markus.franzen@lnu.se
[3]Department of Ecology, Environment and Plant Sciences, Stockholm University, SE-106 91
Stockholm, Sweden; ayco.tack@su.se
13
14 [+] J.P. and C.H. contributed equally to this work; chloe.haberkorn@zoologi.su.se;
15 javier.pinto@zoologi.su.se
16
17 [*]Corresponding author: Rike Stelkens, Department of Zoology, Stockholm University, Sweden;
18 rike.stelkens@zoologi.su.se
19

20 **Data availability statement**
21 The data that support the findings of this study are openly availab tle in NCBI's Sequence Read
22 Archive (SRA) database under BioProject PRJNA1114957. All the scripts used for data processing,
23 analyses and figures are also openly available on GitHub: https://github.com/chaberko-lbbe/yeast-
24 oaktree.
25

Acknowledgements
We thank the oak bark collection team, including Genoveva Elisabeth Zimmermann, Jonas Lundqvist,
and Sofia Rouot for her help with processing samples in the lab. The work was funded by the
Swedish Research Council (grant 2022-03427) and the Knut and Alice Wallenberg Foundation
(2017.0163) to RS. The authors acknowledge support from the National Genomics Infrastructure in
Stockholm funded by the Science for Life Laboratory, and the SNIC/Uppsala Multidisciplinary Center
for Advanced Computational Science for assistance with sequencing and access to the UPPMAX
computational infrastructure. Computation and data handling were enabled by resources in project
NAISS 2023/22-116 provided by the National Academic Infrastructure for Supercomputing in Sweden
(NAISS) at UPPMAX, funded by the Swedish Research Council through grant agreement no. 2022-
06725.
37
38 **Funding**
39 This work was supported by research grants from the Swedish Research Council (2022-03427) and
40 the Knut and Alice Wallenberg Foundation (2017.0163) to RS.
41

42 **Conflict of interest statement**
43 The authors declare no conflict of interest.
44
Author contributions
Conceptualization: JP, CH, RS, and AT; Methodology, formal analysis, investigation, visualization, and
data curation: CH and JP; Writing (original draft): RS, CH and JP; Writing (review & editing): RS, CH,
JP, AT, and MF; Resources: RS and MF; Supervision, project administration, and funding acquisition:
RS

Abstract

Fermentative yeasts play important roles in both ecological and industrial processes but their

distribution and abundance in natural environments is not well understood. We investigated the

diversity of yeast species associated with oak trees (*Quercus spp.*) in the northern range limit of oak

in Sweden, and identified climatic and ecological conditions governing their distribution. We isolated

yeasts from oak bark collected from 28 forests and identified them to the species level using

metabarcoding. Most communities were dominated by species in the Saccharomycetaceae family,

especially by species of *Saccharomyces, Kluyveromyces,* and *Pichia*. Each of these genera showed a

distinct latitudinal and longitudinal distribution, and both temperature and precipitation metrics

predicted significant variation in their abundance. Consistent with this, laboratory assays revealed

significant effects of temperature on the growth of strains collected from different longitudes and

latitudes. We further found that older trees harbour more diverse and more balanced microbial

communities with more evenly distributed species abundances, and that communities across trees

were more similar when sharing a common dominant genus. This work provides a baseline for future

studies on the impact of climate change on the microbial biodiversity of temperate forests in

northern latitudes, and contributes to a growing collection of fermentative yeasts isolated from the

wild with potential for biotechnological applications.

Keywords: climate data, fermentative yeast, microbial biodiversity, metabarcoding, oak, distribution

patterns, northern range limit

Introduction

Fermentative yeasts are abundant in temperate forests where they play a crucial role in ecological processes (Mozzachiodi et al., 2022). They recycle organic matter, mediate nitrogen and carbon cycles, and serve as a food source for insects (Botha, 2011). They also impact microbial community composition, e.g. by inhibiting the growth of competing microorganisms by producing ethanol (Viljoen, 2006). Fermentative yeasts are also known for their high stress tolerance and their efficient use of monosaccharides and nitrogen (Treseder & Lennon, 2015; Romero-Olivares et al., 2021), making them well-suited for industrial applications, particularly for the production of alcoholic beverages and biofuels. Recent studies suggest that the metabolic diversity of natural yeasts, their variation in niche breadth and stress tolerance, is largely shaped by genetic factors (Opulente et al., 2024). A recent global distribution analysis showed that species richness is highest in mixed, montane forests in temperate climates, and is best predicted by microhabitat, vegetation type, and topography (David et al., 2024). Species ranges are strongly influenced by overlaps with other yeast species, indicating that niche partitioning plays an important role in their biogeography. The distribution of budding yeast in the genus *Saccharomyces* is known to be associated with environmental temperature (David et al., 2024; Mozzachiodi et al., 2022; Sweeney et al., 2004).

Oak trees (*Quercus spp.*) have been described as a frequent habitat for fermentative yeast, in different parts of the world (Mozzachiodi et al., 2022; Robinson et al., 2016; Sampaio & Gonçalves, 2008; Sniegowski et al., 2002). Oak trees, especially their bark and sap, provide natural sugars (Ferreira et al., 2018) and can accumulate decaying plant matter and moisture, which creates suitable growth environments for yeast. Once the polysaccharides in the bark are broken down into smaller sugars, e.g. by filamentous fungi (Battaglia et al., 2011; de Vries & Visser, 2001), they are available as a carbon source for fermentative yeasts. Forests in the north of Europe represent the northern range limit of oak, which is marked by long, cold winters and warm (but not hot) summers, a shorter growing season, and more coniferous vegetation. Historically, oak forests were widespread in southern Scandinavia. Over the past centuries, a combination of biogeographical shifts and human activities have significantly reduced the volume of oak forests (Löf et al., 2016). Tree health has also substantially deteriorated, especially in *Quercus robur*, which has suffered from crown defoliation and a general decline in recent decades (Drobyshev et al., 2007). Today, the remaining oak forests are concentrated along the coastlines in southern Sweden in the temperate and hemiboreal climate zones, delimited in the north by the subarctic climate of the boreal zone (Drobyshev et al., 2008). Although remaining oak forests in Sweden are likely important reservoirs of microbial diversity, the current lack of microbial data has precluded a systematic investigation of this ecosystem. Whether

fermentative yeasts are part of the oak microbiome also in the northernmost range of the *Quercus*
distribution, is so far unknown.

Our overarching aim was to investigate the diversity of fermentative yeasts found in the
northern range limit of oak, and identify climatic and environmental drivers of their distribution. For
this, we isolated yeasts from oak bark samples collected across southern Sweden, specifically from
the nemoral and hemiboreal climate zones (Jonsson et al., 2016). Our sampling area covered 583 km
from south to north, along latitudes between 55°52'33''N and 60°76'78''N (from southern Sweden
to as far north as the Swedish oak tree line allowed), and 379 km from west to east, along longitudes
between 12°48'38''E and 18°63'15''E (from the North sea to the Baltic sea). We used enrichment
protocols designed for fermentative yeast in the genera *Saccharomyces*, *Komagataella*, *Lachancea*,
and *Candida* (Cubillos et al., 2019; Sampaio & Gonçalves, 2008; Sniegowski et al., 2002; Villarreal et
al., 2022). For biodiversity and abundance assessment, we used metabarcode sequencing of the
internal transcribed spacer (ITS) region, allowing identification down to the species level (Alsammar
et al., 2019; Větrovský et al., 2019). Specifically, we asked 1) if microbial communities differ in their
composition and diversity, 2) if there are spatial patterns in microbial community structure along
latitudinal and longitudinal gradients, reflecting variation in temperature and precipitation, 3) if oak
tree hosts characteristics (e.g. tree age) predict yeast diversity, and 4) if the growth of the yeast
strains we isolated is affected by temperature using laboratory assays.
121

Materials and Methods

Yeast sampling from oak tree bark

Eight oak trees were sampled in each of 28 stands, at 23 different locations across the south of
Sweden, in both nemoral and hemiboreal vegetation zones (**Figure 1**). Trees included 67 *Q. robur*,
nine *Q. petraea*, and five oak trees (*Q. sp*) with unidentified species status. Sampling period was
during the mid and late summer, from July to September 2023. Approximately 15g of bark (ca. 4 × 1
× 0.5 cm), including both the outer and inner bark layers, was collected from each tree using an
ethanol-sterilized knife. Where single pieces of this size were not obtainable, multiple smaller bark
pieces were collected. Samples were collected using nitrile gloves and stored individually in sterile
plastic bags. All sampling tools were sterilized with 90% ethanol before and after sampling. Samples
were transported in a cooled container and stored refrigerated until processing.

Bark samples from each tree were grown in 25 mL of liquid enrichment medium (a modified
version of the medium described by Sampaio & Gonçalves, 2008), consisting of YNB (yeast nitrogen
base) supplemented with 1% (wt/vol) raffinose and 4% (vol/vol) ethanol. Instead of using 8%
ethanol, we reduced ethanol to 4% to increase our chances of uncovering more biodiversity,

including groups of yeast that are not able to tolerate high alcohol concentrations. We inoculated for

15 days in 50 ml Falcon tubes at room temperature. From each tube, 100 μL of liquid was then

sampled and diluted 4X in sterilised water. Fifty microliters of diluted sample were then plated onto

solid YMA media (1% glucose, 0.5% peptone, 0.3% yeast extract, 0.3% malt extract, 2% agar) with 4%

ethanol using glass beads, and grown for five days at 30°C. To confirm the presence of yeast, a PCR

of the ITS region was performed for three colonies per stand (**Figure S1**). Two to three individual

trees per stand with growth of putative yeasts were then randomly selected for DNA extraction

(total number of trees n = 81).

Figure 1. Schematic of the sampling, enrichment, and isolation of yeast from oak trees.

DNA Extraction

From each enrichment plate (one per tree) we randomly selected ~20 colonies, scraped the colonies

with a sterile pipette tip, and added them to 250 μL of 1X PBS. After vortexing, samples were spun

down at 3000 rpm for three minutes and the supernatant was discarded. Fifty microliters of

zymolyase solution (1 mg/mL zymolyase, 1 M sorbitol, pH 7) were added to the pellet and mixed by

pipetting. Cells were then incubated at 37°C for 2h before adding 1 mL of sterilised water. Cells were

then spun down at 13 000 rpm for 30 seconds to discard the supernatant and resuspended in 250 μL

PBS. After vortexing, 20 μL of proteinase K solution and 290 μL of AL buffer were added to each

sample, and mixed thoroughly by pipetting. Samples were then incubated at 60°C for 20 minutes

and cellular debris was removed by centrifugation at 3,000 rpm for 3 minutes. A KingFisher 96 deep-

well plate was prepared following the OMEGA Bio-Tek protocol to perform the DNA extraction. PCR1

was performed to enrich the samples for putatively present yeast DNA, by amplifying the ITS region

using primers for ITS4 and ITS5 (**Figure 2**, primer sequences in **Table S1**) (White et al., 1990;

Alsammar et al., 2019). Amplification was carried out using the Takara Taq polymerase kit. PCR1

amplicon concentrations were assessed using Qubit dsDNA HS Assay Kits. All concentrations were

normalised to 2ng/µL by diluting samples in MilliQ water, up to a volume of 50 µL per sample.

Sequencing

PCR1 amplicons with normalised concentrations were cleaned up using MagSI-DNA NGS PREP Plus

magnetic beads at the National Genomics Infrastructure (NGI, Solna, Sweden), followed by an

indexing PCR2 with Adaptamera indices and a second cleanup using magnetic beads. Quality control

was performed on the final library before sequencing on the NovaSeq 6000 platform. Raw reads

were deposited on NCBI's Sequence Read Archive (SRA) database under BioProject PRJNA1114957.

Figure 2. Schematic of ITS amplification using PCR1 with primer overhangs (**Table S1**), followed by indexing in

PCR2 for Illumina library preparation and sequencing.

Read Processing

Reads were processed to identify yeast species using the nf-core/ampliseq pipeline (Straub et al.,

2020) with default parameters and option 'illumina_pe_its'. Amplicon Sequence Variants (ASVs), i.e.

unique DNA sequences obtained from amplicon sequencing, were inferred and annotated with

DADA2 against the UNITE general FASTA release for Fungi v9.0 (Abarenkov et al., 2024). ASVs were

kept when their sequence similarity was above a common 97% confidence threshold (Kauserud,

2023). All the scripts used for data processing, analyses and figures are openly available on GitHub:

https://github.com/chaberko-lbbe/yeast-oaktree.

Environmental predictors

Longitudes and latitudes were extracted from GPS coordinates for each sampling location. Using

these coordinates, temperature and precipitation data were extracted from WorldClim version 2.1

(Fick & Hijmans, 2017) with QGIS software (Geographic Information System). Average monthly

values were computed, and used to extract annual mean, maxima, and minima for both

temperature and precipitation at each location. Because means followed the same trends as

minima, mean data were not used. Mean annual temperatures ranged between -8°C and 25°C, and

precipitation ranged between 13.3mm and 136.7mm, averaged across monthly mean minima and

maxima.

Individual host tree metrics

Tree height was measured using a Suunto PM-5/1520 clinometer. Diameter at 1.3 m breast height

(DBH) was assessed using Haglöf Mantax Blue Klave callipers. Bark depth was measured as the mean

depth of bark crevices in the four cardinal directions (North, South, East, and West) using a metal

ruler, following the methodology of Johansson et al. (2009). Tree age was determined by counting

the total number of annual growth rings in each core sample obtained at breast height, following

standard dendrochronological procedures and cross-validation (Holmes, 1983). This provided a

minimum age estimate, as core sampling at 1.3 m height does not account for the time required for

trees to reach breast height. Annual radial growth rates were quantified through tree-ring widths

(TRW) measurements using a digital LINTAB positioning table connected to an Olympus

stereomicroscope. These measurements were recorded to the nearest 0.01 mm using TSAPWin

Scientific software (Rinn, 2003). Each annual ring width represents the radial stem increment for

that particular year, providing a high-resolution time series of growth patterns. Growth rates were

calculated as the annual increment in ring width, with the mean annual growth rate determined by

averaging these measurements across the entire core sample. To account for age-related growth

trends and ensure data quality, all TRW series underwent standardization and cross-dating

procedures using COFECHA software (Johnson & Abrams, 2009).

Growth-at-temperature assays

To test whether temperature has an effect on the growth of the strains we isolated from the wild,

we assessed the genus affiliation of three colonies per site by Sanger sequencing. Since the strains

were collected blindly from the remaining colonies on the plates after amplicon sequencing, some

did not match the dominant species identified for each location by amplicon sequencing. Thus, for

the growth assays, we only retained strains identified by Sanger sequencing as *Saccharomyces*,

Kluyveromyces and *Pichia*, from a total of 18 locations. Sanger sequencing confirmed the presence of

Pichia in only one sample (Halmstad 1), which precluded us from testing for effects of latitude and

longitude on growth-at-temperature for this genus.

We measured the maximum biomass of each strain as a proxy for growth at three different

temperatures: 5°C (below 4°C growth is usually no longer observed in yeast; Salvadó Z. et al., 2011),

16°C, which is close to the mean temperature during Swedish summer; and 35°C, which is close to

the highest temperature ever recorded in Sweden (SMHI, 2023). Growth assays were performed on

a BioTek Epoch 2 microplate spectrophotometer (Agilent, USA) using maximum biomass, measured

as optical density (OD_{600nm}). We standardised all inoculates to an initial OD_{600nm} of 0.1 (approximately

10^6 cells) in 200 μL of YPD media (20g/L peptone, 10g/L yeast extract, 2% glucose). Before final

growth measurements were taken, each strain was grown for 24 hours (for measurements at 16°C

and 35°C) and for 120 hours (for measurements at 5°C) in 96-well plates. We used 10 technical OD

replicates per strain. Biomass was blank-corrected using negative controls (6 wells with media but no

yeast).

Statistical analysis

Abundance and taxonomy tables based on the ASV read data were used to compute alpha diversity

indices: ASV richness (the number of different ASVs), Shannon (accounting for both ASV richness and

their relative abundance; Shannon & Weaver, 1963), and Evenness indices (how evenly distributed

ASV abundance is among species within a community; Wilsey & Potvin, 2000) with the package

R/vegan v2.5-7 (Oksanen et al., 2020). A non-metric multidimensional scaling (NMDS) ordination was

computed based on ASV richness using the vegan function metaMDS (distance "bray", 2 dimensions,

try max 1000). To evaluate whether pairs of species co-occur more or less frequently than would be

expected by chance across a set of locations, R/cooccur v1.3 was used (Veech, 2013; Griffith et al.,

2016). To test if ASV abundance across the three dominant genera was predicted by geospatial and

environmental metrics, we ran linear models using the function lm in R/stats v4.2.3. Pairwise

comparisons of slopes between genera were made within each spatial gradient, using the function

emtrends (t-test) in the package R/emmeans v1.8.1 (Lenth, 2022). To assess differences in

community composition across tree species and insularity, a Bray-Curtis distance matrix was

computed based on genus-level abundance data using the vegdist function from the package

R/vegan (method "bray"). Permutational multivariate analysis of variance (PERMANOVA) was then

conducted with the adonis2 function (R/vegan), using either tree species or insularity as the

explanatory variable (999 permutations). Growth-at-temperature assay data were analysed by

running linear models against longitude and latitude for each genus at each temperature.

Results

A total of 149 ASVs were detected, of which 69 were attributed to the genus *Saccharomyces*. Up to

26 different ASVs were detected per tree with on average 166 620 ASV reads per tree. No ASVs were

detected in 3 out of 81 trees (one tree each in Blå Jungfrun, Björnstorp and Vårgårda-1).

1. Microbial communities across the sampling area differ in composition and diversity

First, we set out to test if microbial communities across the sampling area differ in their composition

and diversity. We identified a total of 14 genera of yeasts (**Figure 3A**) belonging to five families

across all samples (Debaryomycetaceae, Pichiaceae, Saccharomycetaceae, Saccharomycetales fam.

incertae sedis, and Saccharomycodaceae). Computing the sum of ASV reads detected across all

samples for each family revealed that species belonging to Saccharomycetaceae were the most

frequently detected (71.88%), followed by Pichiaceae (24.94%) (**Figure S2**). Among genera,

Saccharomyces was the most common (34.06%), followed closely by two other genera of the

Saccharomycetaceae family: *Kluyveromyces* (33.91%) and *Pichia* (22.39%) (**Figure 3A**). These three

genera represented up to 90.36% of the total ASV reads, while each of the eleven remaining genera

represented less than 3% of ASV reads.

Samples were plotted using non-metric multidimensional scaling (NMDS) and coloured

according to the most frequent genus (based on the number of ASV reads) within each sample

(**Figure 3B**). Three main clusters were revealed, which grouped together the samples that were

dominated by species of either *Saccharomyces*, *Kluyveromyces* or *Pichia* (with a larger spread in the

Pichia cluster). The proximity of points within each cluster suggests that microbial communities are

more similar when sharing a common dominant genus. While there was no significant difference

between the three clusters in species richness (median of 2.5 species), the number of yeast species

per tree was slightly higher when the dominant genus was *Pichia* (**Figure S3**).

Figure 3. Overview of the 14 yeast genera detected on oak trees in Sweden. **A)** Percentage of ASVs mapping to each genus across all samples. **B)** Projection using non-metric multidimensional scaling (NMDS), where each circle represents a single tree. Ordination is based on ASVs richness with colours indicating the dominant genus. Outlier samples LANY-1, TAN-1 and VASV-1 were removed for display purposes.

Up to 18 species were found across the 14 genera (**Table 1**). Within the genus *Saccharomyces*, only *S. paradoxus* was detected. In the genus *Pichia*, we identified *P. mandshurica*, *P. membranifaciens,* and *Pichia sp..* In the genus *Kluyveromyces*, we found *K. lactis* and *K. dobzhanskii*.

Table 1. Species detection across all genera. Occurrence represents the number of trees (out of 81) where species were detected with at least one ASV. Species names were assessed using Fungi v9.0, and current names were extracted from MycoBank (Robert et al., 2013).

Family	Genus	Species name detected	Current species name	Occurrence
Saccharomycetales fam. Incertae sedis	*Candida*	*Candida castellii*	*Oligophagozyma castellii* (2024)	1
Saccharomycetaceae	*Citeromyces*	*Citeromyces matritensis*	-	1
Debaryomycetaceae	*Debaryomyces*	*Debaryomyces delbrueckii*	*Torulaspora delbrueckii* (1961)	11
Saccharomycodaceae	*Hanseniaspora*	*Hanseniaspora osmophila*	-	3
Debaryomycetaceae	*Hyphopichia*	*Hyphopichia burtonii*	-	1
Saccharomycetaceae	*Issatchenkia*	*Issatchenkia orientalis*	*Pichia kudriavzevii* (1965)	7
Saccharomycetaceae	*Kazachstania*	*Kazachstania servazzii*	*Monosporozyma servazzii* (2024)	1
Saccharomycetaceae	*Kluyveromyces*	*Kluyveromyces dobzhanskii*	-	50
		Kluyveromyces lactis	-	6
Pichiaceae	*Kregervanrija*	*Kregervanrija fluxuum*	-	2
Saccharomycetaceae	*Lachancea*	*Lachancea kluyveri*	-	1

Saccharomycetaceae	Ogataea	Ogataea dorogensis	-	2
		Ogataea nonfermentans	-	1
Pichiaceae	Pichia	Pichia mandshurica	-	1
		Pichia membranifaciens	-	37
		Pichia sp		4
Saccharomycetaceae	Saccharomyces	Saccharomyces paradoxus	-	55
Saccharomycodaceae	Saccharomycodes	Saccharomycodes ludwigii	-	3

Co-occurrences of the 18 yeast species detected were visualised using a chord diagram
(**Figure 4**). The thickness of the connecting lines between species reflects the frequency with which
they co-occur across the sampled locations (between zero and 34 times; **Figure S4**). The species that
co-occurred most often were *Saccharomyces paradoxus*, *Kluyveromyces dobzhanskii* and *Pichia*
membranifaciens. However, when comparing the observed frequency of co-occurrence to what
would be expected under a random null model (binomial test), only a single significant positive co-
occurrence was detected (p-value 'greater than expected' < 0.05) between *Debaryomyces*
delbrueckii and *Saccharomyces paradoxus,* suggesting that these species are more likely to co-occur,
possibly sharing compatible ecological requirements or forming a symbiotic relationship. On the
other hand, *Pichia sp.* and *Saccharomyces paradoxus*, as well as *Debaryomyces delbrueckii* and
Kluyveromyces dobzhanskii, exhibited significant negative co-occurrence (p-value smaller than
expected' < 0.05) suggesting that these species are less likely to be found in the same environment
due distinct habitat preferences or niche competition.

Figure 4. Chord diagram depicting yeast species co-occurrence, based on shared presence across sampled trees. The connections (arcs) between species indicate how frequently they co-occur, with thicker arcs representing more frequent co-occurrences.

2. Spatial patterns: Yeast genus dominance is correlated to environmental gradients

Next, we tested for spatial patterns in the abundance of the three most dominant yeast genera along latitudinal and longitudinal gradients. We found that the abundance of ASVs significantly decreased with longitude (going from west to east) in the genus *Kluyveromyces* (linear model: R^2 = 0.19, p = 0.024), but not in *Saccharomyces* (R^2 = 0.03, p = 0.370) or *Pichia* (R^2 = 0.10, p = 0.138; **Figure 5B**). ASV abundance in *Pichia* significantly increased with latitude (going from south to north; linear model: R^2 = 0.40, p = 0.001), but not in *Saccharomyces* (R^2 = 0.18, p = 0.519) or *Kluyveromyces* (R^2 = 0.04, p = 0.309; **Figure 5C**).

We then used a linear model to test whether variation in ASV abundance in *Saccharomyces*, *Kluyveromyces*, and *Pichia*, i.e. 'genus', was explained by 'longitude', 'latitude' and their interactions (minus the interactions Long:Lat and Genus:Long:Lat), which was significant ($F_{8,67}$ = 2.99, p = 0.006). Of the individual coefficients, ASV abundance was significantly affected by longitude overall (t = -2.45, p = 0.017) with significant differences between *Pichia* and *Kluyveromyces* (t = -2.66, p = 0.010).

Interactions between *Pichia* vs. *Kluyveromyces* with latitude (t = 2.03, p = 0.047) and *Saccharomyces*
vs. *Kluyveromyces* with longitude were also found to be significant (t = 2.57, p = 0.013).
Overall, these results suggest that species of *Kluyveromyces* are significantly more abundant
in the south-west, that *Pichia* species are most common in the north-east, and that *Saccharomyces*
species were most common in the centre of Sweden (**Figure 5**). Despite the frequent co-occurrence
of *S. paradoxus*, *K. dobzhanskii*, and *P. membranifaciens* (**Figure 4**), this analysis highlights the
respective dominance of different genera over others along geospatial gradients.
333

Figure 5. A) Map of sampling locations with pie charts showing the proportion of ASVs found per yeast genus.
For display purposes, only genera with >2% of ASVs are shown in the legend. Dashed lines indicate the borders
of hemiboreal and nemoral zones, in which oak trees were sampled. The upper line follows the northern
latitudinal distribution limit of oak. **B)** Longitude, and **C)** Latitude correlated with the ASV abundance of the
three most common genera isolated across all locations (*Kluyveromyces*, *Saccharomyces*, and *Pichia*). R-
squared and p-values are indicated for each linear model per genus (p < 0.05 *, <0.01 **).

As latitude increases going from south to north in our sampling area, minimum temperature

decreases (r = -0.95, df = 71, p < 2.2e-16) and maximum precipitation increases (r = 0.64, df = 71, p =

9.05e-10; **Figure 6**). As longitude increases (going from west to east), maximum temperature (r =

0.86, df = 71, p < 2.2e-16) and minimum precipitation decrease (r = -0.73, df = 71, p = 1.74e-13). We

therefore tested whether these climate metrics (maximum/minimum temperature and

precipitation) also predicted variation in yeast diversity (including ASV richness, Shannon, Evenness,

and the raw number of yeast species) and visualised the strength and direction of these correlations

using a heatmap (**Figure 6**). We found that lower minimum precipitation negatively predicted ASV

richness, suggesting that diversity is larger at drier sampling sites (r = -0.24, df = 71, p = 0.04, **Figure**

6, panel A). Note here, that this pattern may be largely caused by the high ASV richness observed for

Saccharomyces (**Figure S5**).

Figure 6. Correlation patterns between environmental variables, tree metrics and yeast diversity indices.

Significance levels of Pearson correlations are indicated by asterisks (p < 0.05*, 0.01**, and 0.001***).

Analysis of the impact of temperature and precipitation on ASV abundance at the 'genus'

level showed that lower annual minimum temperatures predicted higher ASV abundance in *Pichia*

species (linear model: R^2 = 0.26, p = 0.012; **Figure 7B**), while lower maximum temperature predicted

an increase in *Kluyveromyces* ASV abundance nearly significantly (R^2 = 0.16, p = 0.050; **Figure 7A**).

Saccharomyces ASV abundance did not depend on either temperature metric. Higher minimum

precipitation predicted higher ASV abundance in *Kluyveromyces* (R^2 = 0.18, p = 0.037; **Figure 7D**),

while lower maximum precipitation predicted *Saccharomyces* ASV abundance (R^2 = 0.17, p = 0.046,

Figure 7C). Together, these patterns suggest that the colder temperatures and higher precipitation

levels in the north-eastern range limit of oak trees are well-suited for *Pichia* species, while the lower

maximum temperatures and higher precipitation levels on the Swedish west coast are better suited

for *Kluyveromyces* species. The absence of temperature effects on the distribution of *Saccharomyces*

suggests that *S. paradoxus* is more thermo-generalist than the species in the other two genera.

Figure 7. Annual **A)** maximum temperature, **B)** minimum temperature, **C)** maximum precipitation and **D)**

minimum precipitation correlated with the ASV abundance (total number of ASV reads) of the three most

common genera isolated across all locations (*Kluyveromyces, Saccharomyces*, and *Pichia*). R-squared and p-

values are indicated for each linear model per genus if significant (or approaching significance).

3. Ecological drivers of yeast diversity: Older trees have richer and more balanced microbial

communities

Third, we wanted to know if any ecological aspects of the oak tree hosts predicted yeast diversity.

We tested for correlations between yeast diversity measures (including ASV richness, Shannon,

Evenness, and the raw number of yeast species per bark sample) and several oak tree variables (tree

growth, age, height, diameter, bark depth, and tree species (either *Q. robur* or *Q. petraea*)). Tree

metrics are gathered in the bottom left corner of the heatmap (**Figure 6**). As expected, tree diameter

was positively correlated with bark depth (r = 0.81, df = 57, p = 6.98e-15) and tree age (r = 0.41, df =

56, p =0.001). A strong negative correlation was also found between tree age and tree growth (r = −

0.77, df = 56, p = 9.17e-12), which indicates that older trees grow more slowly.

Species diversity (Shannon index, r = 0.28, df = 52, p = 0.042) and species richness (number

of yeast species: r = 0.27, df = 52, p = 0.045) increased with tree age (**Figure 6**, panel B). We also

found that older, more slowly growing trees have a more balanced community composition, i.e.

species in the microbial communities on older trees have more evenly distributed ASV abundances

(correlation between tree growth and evenness: r = -0.32, df = 45, p = 0.03).

Although only 9 trees out of the 81 sampled were identified as *Q. petraea*, they show an

interesting difference in yeast diversity, although not significantly ($F_{1,63}$ = 1.56, p = 0.156), driven by

one particular tree on the island of Gotland. Of the three trees sampled on Gotland, two were *Q.*

robur and one was *Q. petraea*. The latter shows a very high abundance of *Kregervanrija fluxuum*,

which, to our best knowledge, is the only reported occurrence so far of this species in Sweden

(**Figure 8A**). The yeast diversity on islands off Sweden's east coast (Gotland, Gårdby, and Blå

Jungfrun) was significantly different from the diversity found on the mainland ($F_{1,76}$ = 2.27, p =

0.030). We found a strong enrichment on these islands for *Kregervanrija*, but also *Debaryomyces*

delbrueckii, which was found to be highly abundant on a single tree sampled on the island of Blå

Jungfrun (**Figure 8B**).

Figure 8. Relative abundance of yeast genera depending on **A)** host tree species and **B)** insularity. Significant differences in community composition depending on x-axis are indicated by asterisks (p < 0.05*).

4. The impact of temperature on the growth performance of wild yeast isolates

Finally, we tested whether the growth (maximum biomass) of the yeast strains we isolated across the sampling area was affected by temperature, using laboratory growth assays at three temperatures: 5°C, 16°C, and 35°C. We found that culturing temperature significantly affected the growth of *Kluyveromyces* collected from different longitudes and latitudes. *Kluyveromyces* strains from western Sweden grew better than strains from the east at 5°C (R^2 =0.132, p = 0.002; **Figure 9A**), while strains collected in southern Sweden grew better than strains from the north at 35°C (R^2 = 0.114, p = 0.005; **Figure 9B**). For *Saccharomyces,* southern strains grew better than strains from the north at 16°C (R^2 = 0.96, p = 0.002; **Figure 9B)** but no significant effects of latitude and longitude were found when growing *Saccharomyces* at extreme cold or hot temperatures.

Figure 9. Growth (maximum biomass) of yeast strains at 5°C, 16°C and 35°C. Strains were identified by Sanger sequencing to belong to *Kluyveromyces* and *Saccharomyces*, and plotted according to their sampling location's **A)** longitude and **B)** latitude. Each data point represents one of 10 technical replicate optical density measurements per strain and location. R-squared and p-values are indicated for each linear model per genus if significant (p < 0.01**).

Discussion

Temperate forests are a widespread global biome and a diversity hotspot for yeast (Sniegowski et al., 2002; Sampaio & Gonçalves, 2008; Robinson et al., 2016; Mozzachiodi et al., 2022; David et al., 2024). Fermentative yeasts play crucial roles in these ecosystems. They break down organic matter and contribute to nutrient recycling by converting sugars into ethanol, carbon dioxide, and other metabolites such as esters and glycerol (Dashko et al., 2014). Despite their importance for nature and society, our knowledge of the distribution and diversity of fermentative yeast across geographically and ecologically diverse, natural environments is still scarce. Previous work suggests that oak trees are a frequent habitat for yeast (Sniegowski et al., 2002; Sampaio & Gonçalves, 2008; Robinson et al., 2016; Mozzachiodi et al., 2022) but the diversity and abundance of yeast in the northern range limit of their oak hosts has not been investigated. The temperate forests of Scandinavia are marked by a climate with strong seasonal variation between long, cold winters and mild summers. So far, most work in Scandinavia has focused on fungal communities on spruce (*Picea*) (Müller & Hallaksela, 2000) and beech (*Fagus*) (Kubart et al., 2016; Asplund et al., 2019), but

no previous study has investigated whether oak trees in this region carry similar yeast communities

as trees further south in the *Quercus* range. Using metabarcoding, we describe the species diversity

and community composition of fermentative yeasts isolated from oak trees across a large area

(covering 221,000 km²) in the temperate climate zone of Sweden, and test whether climatic and

ecological variables predict their distribution.

The diversity and community composition of oak-associated yeasts in Sweden

We found three main clusters of oak-associated yeast communities across southern Sweden, each

dominated by species of either *Saccharomyces*, *Kluyveromyces* or *Pichia* (**Figure 3B**). Interestingly,

microbial communities across trees were more similar when sharing a common dominant genus.

Generally, the dominance of one genus did not exclude a species from another genus to co-occur in

the same location, i.e. there was no strong pattern of competitive exclusion. However, in some

locations, only a single genus was detected (small dots in **Figure 3B**).

One species pair co-occurred more frequently than expected by chance: *Debaryomyces*

delbrueckii and *Saccharomyces paradoxus* (**Figure S4**), which suggests they have compatible

ecological requirements or enter beneficial metabolic interactions. A study from temperate

woodlands in North America has found *S. paradoxus* to often co-occur with *S. cerevisiae* (Sweeney et

al., 2004). Although *S. cerevisiae* is a cosmopolitan species due to its long history of domestication

for alcoholic beverage production, we did not detect *S. cerevisiae* in Sweden. However, just as *S.*

cerevisiae, D. delbrueckii (also called *Torulaspora delbrueckii*) has been domesticated for winemaking

(Albertin et al., 2014) and has high tolerance to extremely low temperatures and freezing (Alves-

Araújo et al., 2004). Thus, *D. delbrueckii* may occupy a similar ecological niche in Sweden as *S.*

cerevisiae elsewhere in the world, but may have adaptive advantages at higher latitudes due to its

extreme cold tolerance. Other species pairs in our collection rarely or never overlapped in the same

location (e.g. *D. delbrueckii* with *K. dobzhanskii* or *S. paradoxus* with *Pichia sp.*), perhaps due to

competition or distinct habitat preferences. The positive co-occurence of *D. delbrueckii* with

S.paradoxus, and its negative co-occurence with *K. dobzhanskii,* reinforce our hypothesis that

Saccharomyces and *Kluyveromyces* have different niches in Sweden (**Figure 5**).

We also identified non-conventional yeast such as *Saccharomycodes*, used in the production

of alcohol-free beer (Montanari et al., 2009), *Kazachstania*, a yeast with potential use in

biorefineries (Balarezo-Cisneros et al., 2023), *Isaatchenkia*, a halophilic yeast with the capacity to

grow at low-pH conditions (Matsushika et al., 2016), and *Kregervanrija*, a yeast that weakly ferments

sugars (Kurtzman, 2011). Some of the species we detected are also found in human-associated

environments, e.g. during the spontaneous fermentation of grape juice. Non-*Saccharomyces* yeasts

such as *Hanseniospora*, *Candida*, *Pichia*, and *Kluyveromyces* are predominant during the initial phase

of alcoholic fermentation, before the conversion of sugars into ethanol by *Saccharomyces* (Jolly et

al., 2014). For instance, *P. membranifaciens* is commonly used in industrial applications because it

produces "killer toxins" that help control the growth of spoilage yeast and filamentous fungi (Santos

& Marquina, 2004).

Since the enrichment medium we used for strain isolation prior to metabarcoding favours

the growth of Saccharomycodaceae yeasts, and ITS sequencing protocols can add additional bias (De

Filippis et al., 2017), we cannot rule out the possibility that other yeast species are also present on

Swedish oak. Our collection of oak-associated fermentative yeast from southern Sweden, including

1150 cryo-preserved strains across all sampling locations, is available upon request for future

experimental work in ecology and evolution, or for potential uses in industry.

The abundance and distribution of oak-associated yeasts in Sweden

Our results reveal interesting patterns in the abundance and distribution of oak-associated yeast

across a longitudinal gradient in southern Sweden, and suggest that climate variation from west to

east and south to north, especially in temperature and precipitation (**Figure 7**), contribute to this.

Both coasts experience an oceanic climate, with more extreme seasonal temperature fluctuations in

the east than on the milder west coast (Köppen & Geiger, 1930). The west coast in particular is

characterised by a more humid climate. Going northwards, the climate gets colder and wetter, and

the vegetation changes from agricultural lands to deciduous forests, mixed forests, and even some

taiga. Our findings suggest genus-specific responses to these climatic gradients (**Figure 5B**).

Kluyveromyces showed a strong negative correlation with longitude with higher abundance in

western regions, while *Pichia* dominated in the east, and *Saccharomyces* was more common in

central areas. Compared to *Pichia, Kluyveromyces* may be less tolerant to the larger temperature

swings on the east coast. These differences in the spatial distribution of yeast genera across Sweden,

together with their overlap in some locations (**Figure 3B**), suggest that niche-defining ecological and

climate factors are more likely to shape microbial community composition than competitive

exclusion. In addition to climate, factors such as soil composition and competition from other

microbes, which we have not assessed here, also likely influence the observed distribution patterns

(Brockett et al., 2012; Kowallik et al., 2015; Mundra et al., 2021).

Effects of temperature on the growth of yeast isolates from the wild

Previous studies have shown that species of both *Kluyveromyces* and *Pichia* have a wide range of

critical thermal minima and maxima, from 5°C to 45°C (Dickson et al., 1979; Slininger et al., 1990;

Nambu-Nishida et al., 2017) and their frequent isolation from plants and insects in North America

and Russia (Sukhotina et al., 2006) suggests that they are resilient to a large range of temperatures.

Our assays revealed significant effects of temperature on the growth of fermentative yeast collected

from different longitudes and latitudes in Sweden, especially in the genus *Kluyveromyces* (**Figure 9**).

The better cold performance (at 5°C) of *Kluyveromyces* strains collected in the west compared to

strains from the east, suggests they are able to perform well in the relatively milder winter climate

on the west coast, while the better performance of southern vs. northern strains at hot conditions

(35°C) indicates that southern strains are well adapted to the warm summer temperatures typical

for this region, including occasional heat waves. This pattern is consistent with our finding that the

more moderate climate in the southwest predicts a significantly higher abundance of *Kluyveromyces*

(**Figure 5**). In agreement with this, a study in North America showed that *K. dobzhanskii* was most

frequently isolated at moderate temperatures (10 - 20°C) but less often at 30°C (Sylvester et al.,

2015).

In *Saccharomyces*, the absence of significant effects of longitude and latitude on growth at

extreme temperatures (**Figure 9**) is in line with the previously reported thermogeneralist

performance of *S. paradoxus* (Sniegowski et al., 2002; Robinson et al., 2016) - the most abundant

yeast species in our sampling area (**Table 1, Figure 3**). *S. paradoxus* is, after *S. cerevisiae,* the most

cosmopolitan species of *Saccharomyces* and is frequently isolated from trees, soil, and bark of

deciduous trees in the Northern Hemisphere (Charron et al., 2014). The ability of *Saccharomyces* to

grow both at warm and cold temperatures, for instance through the production and subsequent use

of glycerol as a cryo-preservative (Hohmann, 2002; Koh, 2013), may give them a selective advantage

over other fermentative yeasts in the generally cold climate of Sweden (Sweeney et al., 2004;

Salvadó Z. et al., 2011).

Conclusions

Our study provides insights into the diversity and distribution of fermentative yeasts in the northern

range limit of their oak tree hosts. To our knowledge, our data provide the first evidence that i) older

oak trees supported greater yeast species richness, and ii) that the relative strain frequencies in each

tree-specific microbial community are more balanced, the older the tree is. This suggests that tree

age plays an important role in establishing more stable microbial communities, allowing multiple

species to coexist in similar abundance. We found that communities were largely dominated by

species of the Saccharomycetaceae family, with notable longitudinal and latitudinal gradients in

their composition, which are associated with climate variables, especially temperature and

precipitation. Future research on the underlying genetic basis of thermal tolerance of these wild

yeast isolates may shed light on their vulnerability to future climate change. This is important as

changes in microbial community compositions can have cascading effects through the food web, and

potentially affect the diversity of entire ecosystems (Pörtner & Farrell, 2008) and their services

(Cavicchioli et al., 2019). Together, our findings contribute to a better understanding of the diversity

of natural yeast communities and the drivers of their distribution in the environment.

References

Abarenkov, K., Nilsson, R. H., Larsson, K.-H., Taylor, A. F. S., May, T. W., Frøslev, T. G., Pawlowska, J.,
Lindahl, B., Põldmaa, K., Truong, C., Vu, D., Hosoya, T., Niskanen, T., Piirmann, T., Ivanov, F.,
Zirk, A., Peterson, M., Cheeke, T. E., Ishigami, Y., … Kõljalg, U. (2024). The UNITE database for
molecular identification and taxonomic communication of fungi and other eukaryotes:
Sequences, taxa and classifications reconsidered. *Nucleic Acids Research*, *52*(D1), D791–
D797. https://doi.org/10.1093/nar/gkad1039

Albertin, W., Chasseriaud, L., Comte, G., Panfili, A., Delcamp, A., Salin, F., Marullo, P., & Bely, M.
(2014). Winemaking and Bioprocesses Strongly Shaped the Genetic Diversity of the
Ubiquitous Yeast Torulaspora delbrueckii. *PLOS ONE*, *9*(4), e94246.
https://doi.org/10.1371/journal.pone.0094246

Alsammar, H. F., Naseeb, S., Brancia, L. B., Gilman, R. T., Wang, P., & Delneri, D. (2019). Targeted
metagenomics approach to capture the biodiversity of Saccharomyces genus in wild
environments. *Environmental Microbiology Reports*, *11*(2), 206–214.
https://doi.org/10.1111/1758-2229.12724

Alves-Araújo, C., Almeida, M. J., Sousa, M. J., & Leão, C. (2004). Freeze tolerance of the yeast
Torulaspora delbrueckii: Cellular and biochemical basis. *FEMS Microbiology Letters*, *240*(1),
7–14. https://doi.org/10.1016/j.femsle.2004.09.008

Asplund, J., Kauserud, H., Ohlson, M., & Nybakken, L. (2019). Spruce and beech as local determinants
of forest fungal community structure in litter, humus and mineral soil. *FEMS Microbiology
Ecology*, *95*(2), fiy232. https://doi.org/10.1093/femsec/fiy232

Balarezo-Cisneros, L. N., Timouma, S., Hanak, A., Currin, A., Valle, F., & Delneri, D. (2023). High
quality de novo genome assembly of the non-conventional yeast Kazachstania bulderi
describes a potential low pH production host for biorefineries. *Communications Biology*,
6(1), 918. https://doi.org/10.1038/s42003-023-05285-0

Battaglia, E., Benoit, I., van den Brink, J., Wiebenga, A., Coutinho, P. M., Henrissat, B., & de Vries, R.
P. (2011). Carbohydrate-active enzymes from the zygomycete fungus Rhizopus oryzae: A
highly specialized approach to carbohydrate degradation depicted at genome level. *BMC
Genomics*, *12*, 1–12.

Botha, A. (2011). The importance and ecology of yeasts in soil. *Soil Biology and Biochemistry*, *43*(1),
1–8. https://doi.org/10.1016/j.soilbio.2010.10.001

Brockett, B. F., Prescott, C. E., & Grayston, S. J. (2012). Soil moisture is the major factor influencing
microbial community structure and enzyme activities across seven biogeoclimatic zones in
western Canada. *Soil Biology and Biochemistry*, *44*(1), 9–20.

Cavicchioli, R., Ripple, W. J., Timmis, K. N., Azam, F., Bakken, L. R., Baylis, M., Behrenfeld, M. J.,
Boetius, A., Boyd, P. W., & Classen, A. T. (2019). Scientists' warning to humanity:
Microorganisms and climate change. *Nature Reviews Microbiology*, *17*(9), 569–586.

Charron, G., Leducq, J.-B., Bertin, C., Dubé, A. K., & Landry, C. R. (2014). Exploring the northern limit
of the distribution of Saccharomyces cerevisiae and Saccharomyces paradoxus in North
America. *FEMS Yeast Research*, *14*(2), 281–288.

Cubillos, F. A., Gibson, B., Grijalva-Vallejos, N., Krogerus, K., & Nikulin, J. (2019). Bioprospecting for
brewers: Exploiting natural diversity for naturally diverse beers. *Yeast*, *36*(6), 383–398.

Dashko, S., Zhou, N., Compagno, C., & Piškur, J. (2014). Why, when, and how did yeast evolve
alcoholic fermentation? *FEMS Yeast Research*, *14*(6), 826–832.

David, K. T., Harrison, M.-C., Opulente, D. A., LaBella, A. L., Wolters, J. F., Zhou, X., Shen, X.-X.,
Groenewald, M., Pennell, M., & Hittinger, C. T. (2024). Saccharomycotina yeasts defy long-
standing macroecological patterns. *Proceedings of the National Academy of Sciences*,
121(10), e2316031121.

De Filippis, F., Laiola, M., Blaiotta, G., & Ercolini, D. (2017). Different amplicon targets for
sequencing-based studies of fungal diversity. *Applied and Environmental Microbiology*,
83(17), e00905-17.

de Vries, R. P., & Visser, J. (2001). Aspergillus enzymes involved in degradation of plant cell wall
polysaccharides. *Microbiology and Molecular Biology Reviews*, *65*(4), 497–522.

Dickson, R. C., Dickson, L. R., & Markin, J. S. (1979). Purification and properties of an inducible beta-
galactosidase isolated from the yeast Kluyveromyces lactis. *Journal of Bacteriology*, *137*(1),
51–61.

Drobyshev, I., Anderson, S., & Sonesson, K. (2007). Crown condition dynamics of oak in southern
Sweden 1988-1999. *Environmental Monitoring and Assessment*, *134*, 199–210.

Drobyshev, I., Niklasson, M., Eggertsson, O., Linderson, H., & Sonesson, K. (2008). Influence of annual
weather on growth of pedunculate oak in southern Sweden. *Annals of Forest Science*, *65*(5),
1.

Ferreira, J. P., Miranda, I., Sousa, V. B., & Pereira, H. (2018). Chemical composition of barks from
Quercus faginea trees and characterization of their lipophilic and polar extracts. *PLoS One*,
13(5), e0197135.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for
global land areas. *International Journal of Climatology*, *37*(12), 4302–4315.

Griffith, D. M., Veech, J. A., & Marsh, C. J. (2016). Cooccur: Probabilistic species co-occurrence
analysis in R. *Journal of Statistical Software*, *69*, 1–17.

Hohmann, S. (2002). Osmotic stress signaling and osmoadaptation in yeasts. *Microbiology and
Molecular Biology Reviews*, *66*(2), 300–372.

Holmes, R. L. (1983). *Computer-assisted quality control in tree-ring dating and measurement*.

Johansson, V., Bergman, K.-O., Lättman, H., & Milberg, P. (2009). *Tree and site quality preferences of*
six epiphytic lichens growing on oaks in southeastern Sweden. *46*(6), 496–506.

Johnson, S. E., & Abrams, M. D. (2009). Age class, longevity and growth rate relationships: Protracted
growth increases in old trees in the eastern United States. *Tree Physiology*, *29*(11), 1317–
1328.

Jolly, N. P., Varela, C., & Pretorius, I. S. (2014). Not your ordinary yeast: Non-Saccharomyces yeasts in
wine production uncovered. *FEMS Yeast Research*, *14*(2), 215–237.

Jonsson, B. G., Ekström, M., Esseen, P.-A., Grafström, A., Ståhl, G., & Westerlund, B. (2016). Dead
wood availability in managed Swedish forests–Policy outcomes and implications for
biodiversity. *Forest Ecology and Management*, *376*, 174–182.

Kauserud, H. (2023). ITS alchemy: On the use of ITS as a DNA marker in fungal ecology. *Fungal*
Ecology, *65*, 101274.

Koh, C. M. (2013). Storage of bacteria and yeast. In *Methods in enzymology* (Vol. 533, pp. 15–21).
Elsevier.

Köppen, W., & Geiger, R. (1930). *Handbuch der klimatologie* (Vol. 1). Gebrüder Borntraeger Berlin.

Kowallik, V., Miller, E., & Greig, D. (2015). The interaction of Saccharomyces paradoxus with its
natural competitors on oak bark. *Molecular Ecology*, *24*(7), 1596–1610.

Kubart, A., Vasaitis, R., Stenlid, J., & Dahlberg, A. (2016). Fungal communities in Norway spruce
stumps along a latitudinal gradient in Sweden. *Forest Ecology and Management*, *371*, 50–58.

Kurtzman, C. P. (2011). Kregervanrija Kurtzman (2006). In *The Yeasts* (pp. 497–501). Elsevier.

Lenth, R. V. (2022). *emmeans: Estimated Marginal Means, aka Least-Squares   Means*. R package
version 1.8.1-1. https://github.com/rvlenth/emmeans

Löf, M., Brunet, J., Filyushkina, A., Lindbladh, M., Skovsgaard, J. P., & Felton, A. (2016). Management
of oak forests: Striking a balance between timber production, biodiversity and cultural
services. *International Journal of Biodiversity Science, Ecosystem Services & Management*,
12(1–2), 59–73.

Matsushika, A., Negi, K., Suzuki, T., Goshima, T., & Hoshino, T. (2016). Identification and
characterization of a novel Issatchenkia orientalis GPI-anchored protein, IoGas1, required for
resistance to low pH and salt stress. *PloS One*, *11*(9), e0161888.

Montanari, L., Marconi, O., Mayer, H., & Fantozzi, P. (2009). Production of alcohol-free beer. In *Beer*
in health and disease prevention (pp. 61–75). Elsevier.

Mozzachiodi, S., Bai, F., Baldrian, P., Bell, G., Boundy-Mills, K., Buzzini, P., Čadež, N., Cubillos, F. A.,
Dashko, S., & Dimitrov, R. (2022). Yeasts from temperate forests. *Yeast*, *39*(1–2), 4–24.

Müller, M. M., & Hallaksela, A.-M. (2000). Fungal diversity in Norway spruce: A case study.
Mycological Research, *104*(9), 1139–1145.

Mundra, S., Kjønaas, O. J., Morgado, L. N., Krabberød, A. K., Ransedokken, Y., & Kauserud, H. (2021).
Soil depth matters: Shift in composition and inter-kingdom co-occurrence patterns of
microorganisms in forest soils. *FEMS Microbiology Ecology*, *97*(3), fiab022.

Nambu-Nishida, Y., Nishida, K., Hasunuma, T., & Kondo, A. (2017). Development of a comprehensive
set of tools for genome engineering in a cold-and thermo-tolerant Kluyveromyces marxianus
yeast strain. *Scientific Reports*, *7*(1), 8993.

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R., Simpson, G. L.,
Solymos, P., Stevens, M. H. H., & Wagner, H. (2020). vegan: Community ecology package. *R*
Package Version 2.5-7. https://vegandevs.github.io/vegan/

Opulente, D. A., LaBella, A. L., Harrison, M.-C., Wolters, J. F., Liu, C., Li, Y., Kominek, J., Steenwyk, J. L.,
Stoneman, H. R., & VanDenAvond, J. (2024). Genomic factors shape carbon and nitrogen
metabolic niche breadth across Saccharomycotina yeasts. *Science*, *384*(6694), eadj4503.

Pörtner, H. O., & Farrell, A. P. (2008). Physiology and climate change. *Science*, *322*(5902), 690–692.

Rinn, F. (2003). TSAP-Win. Time series analysis and presentation for dendrochronology and 409
related applications. *User Reference*.

Robinson, H. A., Pinharanda, A., & Bensasson, D. (2016). Summer temperature can predict the
distribution of wild yeast populations. *Ecology and Evolution*, *6*(4), 1236–1250.

Romero-Olivares, A., Morrison, E. W., Pringle, A., & Frey, S. D. (2021). Linking genes to traits in fungi.
Microbial Ecology, *82*(1), 145–155.

Salvadó Z., Arroyo-López F. N., Guillamón J. M., Salazar G., Querol A., & Barrio E. (2011).
Temperature Adaptation Markedly Determines Evolution within the Genus Saccharomyces.
Applied and Environmental Microbiology, *77*(7), 2292–2302.
https://doi.org/10.1128/AEM.01861-10

Sampaio, J. P., & Gonçalves, P. (2008). Natural Populations of Saccharomyces kudriavzevii in Portugal
Are Associated with Oak Bark and Are Sympatric with S. cerevisiae and S. paradoxus. *Applied*
and Environmental Microbiology, *74*(7), 2144–2152. https://doi.org/10.1128/AEM.02396-07

Santos, A., & Marquina, D. (2004). Killer toxin of Pichia membranifaciens and its possible use as a
biocontrol agent against grey mould disease of grapevine. *Microbiology*, *150*(8), 2527–2534.

Shannon, C. E., & Weaver, W. (1963). The Mathematical Theory of Communication. *Urbana, IL:*
University of Illinois Press.

Slininger, P., Bothast, R., Ladisch, M., & Okos, M. (1990). Optimum pH and temperature conditions
for xylose fermentation by Pichia stipitis. *Biotechnology and Bioengineering*, *35*(7), 727–731.

SMHI. (2023). *Swedish Meteorological and Hydrological Institute open data, Swedish temperature*
record. https://www.smhi.se/kunskapsbanken/meteorologi/svenska-
temperaturrekord/svenska-temperaturrekord-1.5792

Sniegowski, P. D., Dombrowski, P. G., & Fingerman, E. (2002). Saccharomyces cerevisiae and
Saccharomyces paradoxus coexist in a natural woodland site in North America and display
different levels of reproductive isolation from European conspecifics. *FEMS Yeast Research*,
1(4), 299–306.

Straub, D., Blackwell, N., Langarica-Fuentes, A., Peltzer, A., Nahnsen, S., & Kleindienst, S. (2020).
Interpretations of environmental microbial community studies are biased by the selected
16S rRNA (gene) amplicon sequencing pipeline. *Frontiers in Microbiology*, *11*, 550420.

Sukhotina, N., Naumova, E., & Naumov, G. (2006). *Molecular polymorphism of the yeast*
Kluyveromyces dobzhanskii: Geographic populations. 409(1).

Sweeney, J. Y., Kuehne, H. A., & Sniegowski, P. D. (2004). Sympatric natural Saccharomyces
cerevisiae and S. paradoxus populations have different thermal growth profiles. *FEMS Yeast*
Research, *4*(4–5), 521–525.

Sylvester, K., Wang, Q.-M., James, B., Mendez, R., Hulfachor, A. B., & Hittinger, C. T. (2015).
Temperature and host preferences drive the diversification of Saccharomyces and other
yeasts: A survey and the discovery of eight new yeast species. *FEMS Yeast Research*, *15*(3),
fov002.

Treseder, K. K., & Lennon, J. T. (2015). Fungal traits that drive ecosystem dynamics on land.
Microbiology and Molecular Biology Reviews, *79*(2), 243–262.

Veech, J. A. (2013). A probabilistic model for analysing species co-occurrence. *Global Ecology and*
Biogeography, *22*(2), 252–260.

Větrovský, T., Kohout, P., Kopecký, M., Machac, A., Man, M., Bahnmann, B. D., Brabcová, V., Choi, J.,
Meszárošová, L., & Human, Z. R. (2019). A meta-analysis of global fungal distribution reveals
climate-driven patterns. *Nature Communications*, *10*(1), 5142.

Viljoen, B. C. (2006). Yeast ecological interactions. Yeast'Yeast, Yeast'Bacteria, Yeast'Fungi
interactions and yeasts as biocontrol agents. In *Yeasts in food and beverages* (pp. 83–110).
Springer.

Villarreal, P., Quintrel, P. A., Olivares-Muñoz, S., Ruiz, J. J., Nespolo, R. F., & Cubillos, F. A. (2022).
Identification of new ethanol-tolerant yeast strains with fermentation potential from central
Patagonia. *Yeast*, *39*(1–2), 128–140.

White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). 38—AMPLIFICATION AND DIRECT SEQUENCING OF
FUNGAL RIBOSOMAL RNA GENES FOR PHYLOGENETICS. In M. A. Innis, D. H. Gelfand, J. J.
Sninsky, & T. J. White (Eds.), *PCR Protocols* (pp. 315–322). Academic Press.
https://doi.org/10.1016/B978-0-12-372180-8.50042-1

Wilsey, B. J., & Potvin, C. (2000). Biodiversity and ecosystem functioning: Importance of species
evenness in an old field. *Ecology*, *81*(4), 887–892.