1

# A privileged ER compartment for post-translational heteromeric assembly of an ion channel

Sudharsan Kannan[1], William Kasberg[2], Liliana R. Ernandez[1], Anjon Audhya[2], and Gail A. Robertson[1,*]

1   Department of Neuroscience, School of Medicine and Public Health, University of Wisconsin–Madison, Madison, WI 53705

2   Department of Biomolecular Chemistry, School of Medicine and Public Health, University of Wisconsin–Madison, Madison, WI 53706

**\*Corresponding author:** Gail A. Robertson

**Email:** garobert@wisc.edu

**Author contributions:** S.K. and G.A.R. designed research; S.K. and W.K. performed experiments; S.K. and L.R.E. analyzed data; A.A. contributed technology and analytic tools; and S.K. and G.A.R. wrote the paper.

**Competing Interest Statement:** The authors declare no competing interest.

**Classification**: Cell Biology

**Keywords**: arrhythmia, hERG, $I_{Kr}$, long QT syndrome, condensate, phase separation, protein trafficking, proteostasis, cotranslational assembly, endoplasmic reticulum, oligomerization

## Abstract

Mechanisms underlying heterotypic subunit assembly of ion channels and other oligomeric complexes are poorly understood. In the human heart, heteromeric assembly of two isoforms encoded by the *human ether-à-go-go related gene* (*hERG*) is essential for the normal function of cardiac $I_{Kr}$ in ventricular repolarization, with loss of hERG1b contributing to arrhythmias associated with long QT-syndrome (LQTS). While hERG1a homomers traffic efficiently to the plasma membrane, hERG1b homomers are retained in the endoplasmic reticulum (ER). When expressed together, the two subunits avidly associate during biogenesis. Seeking rules specifying heteromeric association, we characterized the fate of hERG1b proteins using confocal and superresolution imaging in fixed and live HeLa cells. We found hERG1b sequestered in punctate intracellular structures when expressed alone in HeLa cells. These puncta, which depend on the presence of an N-terminal "RXR" ER retention signal, represent a privileged ER sub-compartment distinct from that containing ER-retained, type 2 (hERG-based) LQTS mutant proteins, which were rapidly degraded by the proteasome. Introducing hERG1a to cells with preformed hERG1b puncta dissolved these puncta by rescuing extant hERG1b. Rescue occurred by association of fully translated hERG1b with 1a, a surprising finding given previous studies demonstrating cotranslational heteromeric association. We propose that sequestration limits potentially deleterious surface expression of hERG1b homomeric channels while preserving hERG1b for an alternative mode of heteromeric hERG1a/1b channel assembly post-translationally. These findings reveal a surprising versatility of biosynthetic pathways promoting heteromeric assembly.

**Significance Statement**

hERG potassium channels are essential for repolarizing the ventricular action potential.

Heteromeric hERG channels, which conduct cardiac $I_{Kr}$, are formed by association of

hERG1a and 1b subunits. However, mechanisms governing the assembly of hERG and

other heteromeric protein complexes are poorly understood. We identify a noncanonical

ER compartment that sequesters hERG1b, distinct from previously described ER quality

control pathways that degrade misfolded proteins, including LQTS-associated hERG

variants. Instead, this compartment preserves hERG1b and facilitates its post-

translational assembly with hERG1a, challenging the current view of simultaneous,

cotranslational heteromeric assembly. Our findings redefine ER retention as an active

regulatory step in protein complex formation that may broadly govern hetero-oligomeric

assembly in many systems and facilitate therapeutic approaches to arrhythmia and

other diseases.

**Main Text**

**Introduction**

The *human ether-à-go-go-related gene* (*hERG*, or *KCNH2*) encodes the voltage-gated

potassium channel Kv11.1, critical for repolarizing the ventricular action potential (1, 2).

Loss of hERG function due to unintended channel block or genetic mutations in *KCNH2*

can prolong the ventricular action potential, leading to drug-induced or type 2 Long-QT

syndrome (LQT2) and sudden cardiac death (1-3). In addition to cardiac dysfunction,

central nervous system diseases such as epilepsy and schizophrenia can also arise

from mutations and mis-splicing of hERG (4, 5). The majority of LQT2 mutants and the

schizophrenia variant of hERG are trafficking-deficient (6). Thus, understanding the

mechanisms that regulate the expression, assembly, and trafficking of hERG channels

is of broad biological and medical significance.

In human ventricular cardiomyocytes, alternate transcripts encode hERG1a and 1b

subunits (or ERG1a and 1b in other organisms), which assemble to form

heterotetrameric channels producing the repolarizing $I_{Kr}$ current (7-12). The subunits are

identical except in their N-termini, which in ERG1a harbors a Per-Arnt-Sim (PAS)

domain (13) that is missing in ERG1b (7). In the mature channel, the PAS domain

interacts with the C-terminus of the adjacent subunit (14), and perturbation of this

interaction alters channel function (15). Expressed independently in heterologous

systems, ERG1a subunits form homomeric channels that efficiently traffic to the plasma

membrane and produce potassium currents. In contrast, ERG1b homomers traffic

inefficiently, producing little or no current (7, 16). When expressed together, ERG1b

markedly alters the gating properties of ERG1a currents, providing strong evidence for

their coassembly (7). This association appears to occur cotranslationally, based on the

following evidence: 1) isolated hERG1a and 1b N-terminal fragments interact in vitro in

a dose-dependent manner (17); 2) coexpression of a hERG1a N-terminal fragment

disrupts channel oligomerization in cells and impairs core glycosylation, a process

known to occur cotranslationally (17, 18); and 3) immunoprecipitation from HEK293 cell

or cardiomyocyte lysates using a hERG1a-specific antibody copurifies nascent hERG1a

and 1b subunits along with their encoding mRNAs (19). Together these findings indicate

that hERG1a/1b subunit assembly begins cotranslationally, early in biogenesis.

Several lines of evidence underscore the critical role of the heteromeric composition of

cardiac $I_{Kr}$. In canine adult cardiomyocytes, specific antibody staining reveals an

approximately equal distribution of the two subunits, localized primarily to the Z line (9).

Similarly, in both canine and human myocardium, western blot analysis using a pan-

hERG antibody detects roughly equal amounts of the proteins, supporting the

conclusion that cardiac $I_{Kr}$ is predominantly formed by heteromeric hERG1a/1b

channels (9). The native current also displays properties more consistent with the

heteromer than the hERG1a homomer (20). The hERG1b subunit modulates hERG1a

properties, yielding larger currents by accelerating the time course of activation and

recovery from inactivation, thus enhancing repolarization (10). Computational studies

incorporating these differences in channel gating behavior into a cardiomyocyte model

predicted that loss of hERG1b would be pro-arrhythmic (10). This prediction was later

validated experimentally: shRNA depletion of hERG1b in cardiomyocytes or introduction

of the hERG1a-specific N-terminal PAS domain in trans, which effectively converts

native heteromers into hERG1a homomers, leads to prolonged action potential duration

(APD) and increased APD variability, two cellular markers of pro-arrhythmia (8).

Previous studies identified an N-terminal, arginine-based endoplasmic reticulum (ER)

retention motif (RXR) in hERG1b that is critical for its retention in the ER in the absence

of hERG1a (16). In heterologous expression systems, mutating the RXR motif

effectively rescues hERG1b from ER retention: hERG1b proteins are fully glycosylated

(mature) as assayed by western blot, and they produce robust currents, indicating

successful plasma membrane expression (16). Thus, RXR-mediated ER retention

ensures that hERG1b subunits predominantly reach the plasma membrane as

components of heteromers, while restricting their expression as homomers. However,

the detailed mechanisms of ER retention and rescue of 1b remain unclear.

In this study, we discovered that retained hERG1b proteins are stably sequestered

within a privileged ER compartment, which we have termed the "ER sequestration and

assembly" (ERSA) compartment. This compartment provides a protected environment,

shielding hERG1b from degradation while it awaits assembly with hERG1a, adding a

new level of complexity to the previously reported mechanism of cotranslational

assembly (17, 19). These findings reveal that biogenesis of heteromeric $I_{Kr}$ channels is

prioritized through two distinct mechanisms.

## Results

**hERG1b is sequestered in intracellular puncta**

hERG1a and 1b isoforms exhibit distinct cellular distributions when expressed

heterologously. Immunofluorescence confocal microscopy in HeLa cells shows that

hERG1a displays a widespread ER signal along with discernible plasma membrane

localization, indicative of mature, functional membrane expression (Fig. 1*A*) (21). In

contrast, hERG1b forms numerous large intracellular puncta with fluorescence signals

measuring 300 nm or greater, detected using automated image analysis (Fig. 1 *B* and

*C*; see Methods). Many fewer puncta are observed for hERG1a (Fig. 1 *A* and *C*). A

corresponding difference is observed when the two isoforms are expressed

exogenously at low levels in the more native-like milieu of live cardiomyocytes derived

from iPSC cells (Fig. 1 *D-F*).

Control experiments confirm that these puncta are not an artifact of the C-terminal tags

used for detection. Untagged hERG1b puncta expressed in HeLa cells showed the

same distribution when stained with pan-hERG antibody (Fig. S1*A*), and native puncta

were also observed in iPSC-CMs using a pan-hERG antibody (Fig. S1 *B* and *C*),

although the lack of availability of a hERG1b-specific antibody precluded isoform

assignment. Notably, a previous study from our lab demonstrated a punctate

localization pattern for hERG1b in antibody-stained canine ventricular myocytes,

whereas hERG1a exhibited a more diffuse distribution (9). These findings suggest that

the puncta serve as the intracellular substrate for the well-documented ER retention of

homomeric hERG1b subunits (10, 16, 17).

**hERG1b puncta are localized to ER**

To determine the subcellular localization of the puncta at superresolution, we imaged

hERG1b in iPSC-CMs using two-color stimulated emission depletion (STED)

microscopy (Fig. 2). ER labeling was achieved by fusing HaloTag to an ER signal

sequence and the KDEL ER retention motif (Halo-KDEL) (22), while hERG1b was

tagged with a C-terminal SNAP tag ($1b_{SNAP}$). mRNA was transcribed in vitro from the

Halo-KDEL and $1b_{SNAP}$ plasmid constructs and transfected into iPSC-CMs. After 24h,

fluorescent labeling of the ER with the JFX650 HaloTag ligand and $1b_{SNAP}$ with the

JFX554 SNAP-tag ligand revealed that more than 95% of the puncta were centered

within a 50 nm distance from the ER membrane, confirming their ER localization (Fig. 2

*A-E*).

hERG1b puncta showed minimal colocalization with other subcellular compartments,

including COPII (ER exit), COPI (Golgi-to-ER retrieval), lysosomes, or autophagosomes

(Fig. S2). Interestingly, the presence of few autophagosomes (Fig. S2*C*) indicates that

hERG1b expression did not trigger significant cellular stress under these conditions

(23). These findings establish that hERG1b puncta represent an ER-resident

substructure.

**hERG1b is sequestered in a privileged ER compartment**

A majority of LQT2-mutant hERG channels fail to mature, accumulating in the ER where

they are ubiquitinated and targeted for proteasomal degradation (6, 24-27). To

determine whether ER-sequestered hERG1b meets a similar fate, we compared its

retention to that of the trafficking-deficient LQT2 mutant hERG1a-Y611H using HA-

tagged constructs transfected into HeLa cells and immuno-stained (25). As expected,

hERG1a-Y611H exhibited intracellular signal with no apparent membrane localization,

consistent with its ER retention phenotype; however, it exhibited relatively few puncta

compared with hERG1b (Fig. 3 *A-C*).

To determine relative stabilities, we designed a pulse-chase paradigm to monitor

degradation of SNAP-tagged constructs labeled with fluorescent ligand (Fig. 3 *D-F*).

Cells were identified with HaloTag expressed independently from an internal ribosome

entry site (IRES) in the plasmid encoding each isoform (upper right insets, yellow). After

24 h transfection, we labeled extant SNAP-tagged proteins with fluorescent ligand

JFX650 (cyan) and then blocked detection of all subsequently synthesized protein using

a non-fluorescent ligand (PA-JF-549); thus, proteins labeled and imaged at t=0 could be

imaged again at 2-h intervals after labeling (Fig. 3 *D*). As previously demonstrated,

hERG1a fluorescence exhibited only a modest reduction over the 8-h period whereas

hERG1a-Y611H fluorescence showed a relatively rapid, single-exponential decay,

presumably via the action of the proteasome on ER-retained protein (25). The

corresponding LQT2 mutation in hERG1b showed a similar decay. In contrast, hERG1b

exhibited the greatest stability over time (Fig. 3 *E-F*; Fig. S3). These findings indicate

that wild-type hERG1b is harbored in the ER in a privileged compartment we have

termed "ER sequestration and assembly compartment," or ERSA. These experiments

show that ERSA is distinct from ER-associated compartments mediating proteolysis and

cell quality control in a disease setting. Notably, introducing Y271H mutation reduced

the number of hERG1b puncta and led to a significant reduction in protein levels in 8 h

(Fig 3. *E-G*), suggesting hERG1b ERSA precludes misfolded proteins.

**An ER retention motif promotes sequestration of hERG1b in ERSA**

Previous studies showed that hERG1b trafficking is promoted when an N-terminal ER

retention motif, $R^{15}XR$, is mutated. Enhanced trafficking is reflected in the maturation of

a lower, core-glycosylated band on a western blot to a higher, fully-glycosylated band,

and by the expression of membrane currents (16). We determined the effect of this

mutation, $N^{15}XN$, on the punctal content of cells, finding the number reduced by about

half (Fig. 4 *A*, *B*, and *D*). A quantitatively similar result was observed upon coexpression

of hERG1b with hERG1a, known to "rescue" hERG1b in maturation assays (16, 28).

Indeed, we found that coexpression with hERG1a yields fewer hERG1b puncta and the

signal was instead distributed throughout the cytoplasm and plasma membrane (Fig. 4

*C* and *D*). Thus, association with hERG1a dissolves the 1b puncta or prevents their

formation. Moreover, these findings establish the puncta as the physical substrate of

previously described hERG1b ER retention.

**Are hERG1b puncta condensates?**

We hypothesized that hERG1b puncta could represent either membranous vesicles or

condensates arising from multivalent interactions of the cytosolic channel domains (29).

To test whether hERG1b puncta are enclosed by membrane, we labeled cellular lipids

with Potomac Yellow dye (30, 31). Amid the diffuse staining of organelles, we observed

puncta that we assume represent membranous vesicles (Fig. 5 *A*, left panel). However,

we observed only limited colocalization with fewer than 5% of the Potomac Yellow spots

overlapping with hERG1b puncta using a 300 nm center-to-center distance threshold

(Fig. 5 *A* and *B*). We conclude that hERG1b is not bound in membranous vesicles.

Instead, hERG1b proteins concentrate in puncta that we operationally term

condensates.

If ERSA condensates arise from liquid-liquid phase separation, we would expect to

observe dynamic movement of molecules, which can be assessed using fluorescence

recovery after photobleaching (FRAP). With this technique, recovery of fluorescence

would reveal the diffusion of fluorescently tagged protein into the bleached area (32).

However, we observed no FRAP of puncta for at least 5 min after photobleaching (Fig. 5

*C* and *D*). In addition, the puncta were insensitive to treatment with 1,6-hexanediol, a

treatment often used to disrupt weak hydrophobic interactions promoting phase

separation (33) (Fig. S4). Thus, hERG1b condensates may have non-diffusive, gel-like

properties like ER-based TIS granule mesh-like condensates (34, 35), or arise as a

consequence of association within a lipid raft or via protein scaffolding (36, 37).

Given that the differences between hERG1a and 1b occur solely within their N-terminal

domains, we assayed the behavior of GFP-tagged hERG1b N-terminal fragments

(1bNT). We found fluorescent signals in scattered cytosolic puncta and densely

concentrated in the nucleus (Fig. 5*E*). The cytosolic puncta were dynamic, exhibiting

FRAP with slow recovery (Fig. 5 *E* and *F*). In silico analysis of hERG1b via Fuzdrop, a

proteome-wide ranking of proteins according to their propensity to form biomolecular

condensates (38), reveals likely droplet-forming regions in several positions of hERG1b

including the 1b-specific N-terminus (Fig. 5*G*). Thus, the hERG1b N-termini have

characteristics that facilitate their dynamic homotypic association. Like the full-length

protein, mutating $R^{15}XR$ to $N^{15}XN$ reduced the propensity of 1bNT to form puncta, as

quantified by counting the percentage of cells with puncta (Fig. 5 *H and I*), mirroring its

effect on full-length hERG1b and highlighting a key role for this motif in the condensate.

Whether the N-terminal propensity to phase separate plays a role in hERG1b

condensation in puncta will require further investigation.

**Sequestered hERG1b is rescued by hERG1a post-translationally**

In previous studies we showed that hERG1a and 1b assemble cotranslationally as the

subunits emerge from the ribosome (17, 19). Based on this model, we predicted that

hERG1a rescues hERG1b cotranslationally by associating with newly synthesized

subunits, while the sequestered hERG1b would be degraded via quality control

mechanisms. However, our current findings that hERG1b is stably retained in the ERSA

compartment prompted us to test whether pre-existing hERG1b proteins could be

rescued by hERG1a.

To accomplish this experiment, we created a stable cell line inducibly expressing

hERG1b$_{SNAP}$ and tracked its fate in live cells. After inducing hERG1b$_{SNAP}$ for 18 h, we

labeled the existing hERG1b proteins using a 30-min pulse of fluorescent JFX650

SNAP-tag ligand. We then added a nonfluorescent PA-JF-549 SNAP-tag ligand to the

media to block labeling of newly synthesized proteins (Fig. 6*A*). Cells were then

transfected with hERG1a-GFP. After 24 h, we observed few hERG1b puncta and clear

membrane localization of both hERG1a and 1b compared to mock-transfected controls

(Fig. 6 *B-D*), indicating that, unexpectedly, existing hERG1b subunits were rescued by

hERG1a. Thus, the ERSA both protects proteins from degradation and ensures they are

readily available for association with newly synthesized hERG1a.

**Discussion**

In this study, we demonstrate that hERG1b proteins, in the absence of hERG1a, are

selectively retained in a punctate ER sequestration and assembly compartment, or

*ERSA*. ERSA shields wild-type hERG1b proteins from proteasomal degradation, unlike

LQT2 mutant hERG1a and 1b proteins, which are rapidly degraded. Mutating an N-

terminal ER retention signal or co-expressing with hERG1a reduces hERG1b

occupancy within ERSA, indicating that this compartment serves as the physical

substrate for retained hERG1b. Notably, hERG1a rescues fully synthesized hERG1b,

highlighting a novel role for ERSA in facilitating hERG channel heteromerization beyond

previous models of coordinate cotranslation (17, 19). ER sequestration thus acts as a

protective mechanism, safeguarding hERG1b from proteasomal degradation and

preserving it in space and time until hERG1a becomes available for heteromeric

assembly.

Recent studies describe macromolecular protein assembly via three pathways: (1) "co-

cotranslational" or simultaneous assembly, where nascent proteins interact as they

emerge from the ribosome; (2) "co-post-translational" or sequential assembly, where a

fully formed subunit assembles with a different, nascent protein; and (3) "post-

translational" assembly where fully formed subunits assemble after translation is

complete (39, 40). Previous evidence supports the idea that hERG1a and 1b assemble

through simultaneous cotranslation: in vitro, their heterotypic N-terminal domains

associate in a dose-dependent manner; and in cells, disrupting N-terminal interactions

disrupts oligomerization and concomitantly diminishes core glycosylation, which serves

as a time stamp for early biogenesis (17, 18). Here, we present a new paradigm in

which fully synthesized hERG1b can also associate with nascent or newly synthesized

1a, as demonstrated for soluble proteins assembling to form macromolecular structures

in yeast (41, 42). It is unclear whether this association follows a co-post-translational

mechanism where 1a is translated near hERG1b, or a strictly post-translational

mechanism, in which the hERG1b-containing ERSA compartment is dispersed when in

contact with full-length hERG1a subunits.

Findings from this study raise the question of stoichiometry of hERG1b subunits within

the ERSA compartment. One possibility is that hERG1b subunits assemble into

homotetramers, which are escorted out of the ER by tandem interactions with hERG1a

homotetramers. This model is reminiscent of the dimeric trafficking of sodium channel

pairs (43). Alternatively, hERG1b may reside as homodimers that interact with newly

synthesized hERG1a homodimers. This model is inspired by the "dimerization of

dimers" mechanism previously established for homomeric assembly of Kv1.3 channels

(44). When both hERG1a and 1b mRNAs are locally abundant, the ERSA may play less

of a role as cotranslational heterodimers are formed, which then assemble into

heterotetramers. Future work determining stoichiometry and composition under different

conditions will be required to further illuminate the complexities of assembly.

Liquid-liquid phase separation (LLPS) serves to concentrate proteins and nucleic acids

into condensates that regulate many cellular functions and play a critical role in

physiology and disease (45, 46). Although most studies have focused predominantly on

condensates of cytosolic proteins, growing evidence highlights phase separation of

membrane proteins, including those involved in cell signaling (47-49). Our findings

suggest that condensates may aid in preserving the integrity of heteromeric hERG

channels, creating a protected environment that shields proteins from diffusible

components of the proteasomal machinery. This environment, within ERSA, may help

manage asymmetric hERG1a and 1b subunit production arising from uneven

transcription initiated at their respective promoters (7). By serving as a reservoir, the

condensate increases the local concentration of hERG1b, maximizing likelihood of

successful association of these rare proteins in the crowded cellular environment.

Dynamics of the 1bNT fragment within condensates suggests this domain is capable of

phase separation, but the full-length hERG1b ERSA lack liquid properties and may

instead represent gel-like assemblies or arise from scaffolding by unidentified proteins

that are involved in RXR-type ER retention. Thus, although hERG1bNT molecules can

phase separate, we do not know whether this property is sufficient to drive

condensation of the attached transmembrane domains in the ER. The tools available to

test this hypothesis are limited (33), requiring future developments to determine whether

phase separation drives the homotypic association of hERG1b membrane protein in the

cellular environment.

Different sequence motifs regulate ER localization through either retention or retrieval

mechanisms. Soluble proteins such as the ER chaperones protein disulfide isomerase

(PDI) and binding immunoglobulin protein (BiP) can exit the ER but are later retrieved

from the Golgi via COPI-coated vesicles. These vesicles contain KDEL receptors that

recognize a KDEL motif harbored within the soluble proteins' sequences (50, 51).

Membrane proteins often use dibasic motifs such as KKXX, KXK and RXR to regulate

ER localization. These motifs can mediate either retention within the ER, as observed

for NMDA receptor subunits (52), or retrieval from the Golgi via COPI-coated vesicles,

as reported for the GABA$_B$ receptor and TREK2 K+ channel (53, 54). Our findings show

that hERG1b puncta do not colocalize with COPI-positive compartments, suggesting

that hERG1b sequestration by the N-terminal RXR motif does not involve Golgi-ER

retrieval. We propose that the RXR helps establish the ERSA compartment to stably

retain fully synthesized hERG1b proteins in the ER, shielding them from proteasomal

degradation. Introducing an LQT2 mutation in hERG1b that misfolds the protein (25)

precludes its localization to ERSA (and exposes it to the proteasome), suggesting that

correct folding is a prerequisite for ERSA formation (Fig. 3 *E*). This novel mechanism

suggests an expanded role for the RXR motif, linking classical quality control functions

to biophysical compartmentalization to promote successful heteromeric assembly. It

remains to be determined whether RXR-mediated ER retention promotes ERSA

formation in other membrane proteins including the NMDA receptor and $K_{ATP}$ channel

subunits, where the RXR governs the ER exit of only fully realized oligomers (52, 55).

Indeed, studies of such complexes could help establish a role for the ERSA

compartment as fundamental to hetero-oligomeric assembly.

Why does hERG1b require an ER retention motif? We argue that hERG1b homomeric

channels trafficking to the membrane of native cardiomyocytes could deleteriously affect

cardiac $I_{Kr}$. Cardiac $I_{Kr}$ exerts its repolarizing effect late in the ventricular action potential

(56). This critical delay is due to slow activation combined with rapid inactivation,

suppressing currents during early depolarization. Later, as inactivation recovers, the

current rebounds and facilitates repolarization (1, 2, 57). Homomeric hERG1b channels,

however, activate rapidly and exhibit little inactivation, producing a classic delayed

rectifier-type current (7). This current would counteract the depolarized plateau potential

and shorten action potential duration (APD). Similar effects are observed with hERG

agonists or mutants such as inactivation-impaired S631A, both of which are associated

with short QT syndrome, in which the shortened APD allows reentrant arrhythmias in

both the ventricle and atria (58-60). Although no pathogenic mutations in the hERG1b

RXR signal have been reported to date, the essential role of hERG1 hetero-

oligomerization underscores the importance of underlying mechanisms in maintaining

cardiac rhythm.

## Methods

### Cell culture

HeLa cells (ATCC, CCL-2) were cultured in 35mm #1.5 glass bottom dishes (Cellvis,

D35-14-1.5-N) using DMEM culture media (ThermoFisher, 11995065) supplemented

with 10% FBS (ATCC, SCRR-30-2020). HeLa cells were transfected with 500

nanograms of hERG1a or 1b plasmid DNA per 35mm dish with 1 µl of TransIT-X2

Transfection reagent (Mirus Bio, MIR6004), and cells were imaged 48 h after

transfection. Human iCell iPSC-CMs (CDI-FujiFilm) were thawed, plated, and cultured

with CDI media according to the manufacturer's specifications. After 10 to 14 days in

culture, cells were transfected or fixed for immunostaining followed by imaging. For

STED microscopy, mRNAs expressing SNAP tagged hERG1b and HaloTag containing

KDEL motif in its C terminus were transfected to iPSC-CM using Lipofectamine

MessengerMAX (LMRNA001). 100 nM SNAP-tag ligand JFX650 and 100 nM HaloTag

ligand JFX549 were added to the culture media for 30 min, followed by three washes 5

min each using pre-warmed culture media to remove the free ligand. For membrane

labeling, 100 nM Potomac Yellow dye was added to the cell culture media for 15 min,

and the cells were imaged live.

**Plasmids and Generation of inducible stable cell line**

Overexpression plasmids were made from pcDNA3.1 as a backbone. Truncations and

insertion of coding sequences of hERG1a and 1b subunits and fluorescent proteins

were performed using In-Fusion® Snap Assembly Master Mix (Takara, 638948). To

generate a doxycycline-inducible cell line to express SNAP-tagged hERG 1b, we used a

transcription activator-like effector nuclease (TALEN)-mediated targeting system, which

incorporated the transgene at the AAVS1 locus (Qian et al., 2014). The puromycin-

resistant donor cassette was used to express SNAP-tagged hERG1b, followed by an

internal ribosome binding site and a gene encoding mApple. The cassette was

transfected into HeLa cells with the two AAVS1-site specific TALEN plasmids at a ratio

of 8:1:1, and cells were treated with puromycin (0.75 µg/ml) for approximately 3 weeks.

Individual clones were picked and tested for doxycycline-induced expression of SNAP-

tagged hERG1b and mApple. For overexpression studies, cells were treated with

doxycycline (50 ng/ml) for 24-48 h prior to analysis.

**In vitro transcription of mRNAs**

pCDNA3.1 plasmid containing SNAP-tagged hERG1b and Halo-KDEL with T7 promotor

were linearized using EcoRI restriction enzyme at the 3' end. Linearized plasmids were

transcribed using mMESAGE mMACHINE™ T7 Transcription Kit (Invitrogen, AM1344)

and poly(A) tailed using E. coli Poly(A) Polymerase (NEB, M0276S).

**Immunostaining**

Cells were cultured in 12 mm coverslips with #1.5 thickness (Electron Microscopy

Sciences 7229004). For labeling using HA (1:1000, Cell Signaling, 3724S), FLAG

(1:1000, Sigma, F1804-1MG), Pan-hERG (1:200, Enzo Life Sciences, ALX-251-049-

R100), and LC3A/B (1:400, Invitrogen, PA5-22731) antibodies, cells were fixed with 4%

paraformaldehyde (PFA) followed by three 5-min washes using PBS. Cells were

permeabilized using 0.5% Triton-X-100 followed by 30 min incubation in blocking

solution (0.05% Tween-20 and 5% normal goat serum) and overnight incubation at $4^0$C

with primary antibodies diluted in blocking solution, then incubation with secondary

antibodies for 1 h at room temperature. For labeling using Sec24a (1:300, Santa Cruz

Biotechnology, sc-169279), COPB (1:300, Santa Cruz Biotechnology sc-393615) and

LAMP1(1:100, Invitrogen, eBioH4A3) antibodies, cells were fixed using 100% ice-cold

methanol for 10 min at $-20^0$C, followed by three 5-min washes in PBS and blocking and

incubation with primary antibody overnight at $4^0$C, then secondary antibody for 1 h at

room temperature.

**Microscopy and image analysis**

Live and fixed confocal imaging was performed using a Nikon Ti2 spinning disk confocal

microscope with a 60x (1.4NA) and 100x (1.45NA) oil immersion objective and

Hamamatsu ORCA-Flash4.0 sCMOS camera. Nikon Elements software was used for

image acquisition.

Puncta quantification and colocalization analysis were performed using the Imaris

(Oxford instruments) spots model. Spot detection was initialized using an average

estimated diameter of 500 nm and modeled PSF-elongation along the z-axis with an

estimated diameter of 1.2 μm. Initial filtering was applied using the *quality* parameter

with an empirically determined threshold to maximize the inclusion of true-positive

puncta while excluding background noise. Additionally, spot regions were thresholded

based on *local contrast* with an empirically determined threshold to refine spot regions.

To ensure consistency and reproducibility, configuration files containing all the

thresholding parameters were generated and applied across images, enabling unbiased

segmentation. Further, the spots were filtered for those greater than 300 nm in diameter

in the XY plane. Puncta were detected throughout the field of view of the microscopy

images and assigned within individual cellular boundaries to quantify puncta on a per-

cell basis.

For 1bNT condensate experiments, the number of cells with condensates was

quantified manually in a blinded manner. All the quantifications were conducted on raw

images, and sample images provided in the figures were deconvoluted using Huygens

professional software (Scientific Volume Imaging), using the Classic Maximum

Likelihood Estimation (CMLE) algorithm with default settings. Deconvolution was

applied evenly to the images used for comparison.

Leica TCS SP8 STED system equipped with highly sensitive HyD detectors and an HC

Plan Apo CS2 100X, 1.4NA oil immersion objective lens was used to acquire STED

images. Random regions of interest (25 x 25 µm squares) were selected for image

acquisition in each cell. Acquisition parameters were controlled by the Leica Application

Suite platform (LASX) and images were deconvolved using Huygens CMLE algorithm.

Puncta with ER colocalization analysis was performed using the Imaris spots and

surface model. The number of puncta colocalized within the 50 nm, 100 nm, or 200 nm

threshold distance from the center of the puncta to the surface was quantified using

Imaris.

Fluorescence recovery after photobleaching (FRAP) studies were performed using a

Nikon Ti2 spinning disk confocal equipped with a $37^0$C incubator and 5% $CO_2$ supply.

100 µs stimulation with maximum power using a 488 nm stimulation laser was used to

photobleach the region of interest and images were acquired before and after

photobleaching in 2-sec intervals for 3-5 min. ImageJ was used to measure the intensity

of the photobleaching area and plotted against time.

**Pulse-chase analysis of hERG1 subunits**

In the pulse-chase analysis, we transfected plasmids containing either hERG1a-SNAP-

IRES-Halo, hERG1b-SNAP-IRES-Halo, hERG1a-Y611H-SNAP-IRES-Halo or hERG1b-

Y271H-SNAP-IRES-Halo. 24 h after transfection, the proteins were labeled with 100 nM

JFX650 SNAP-Tag ligand and 100 nM JF479 HaloTag ligand, followed by three 10-min

washes. Subsequently, 100 nM PA-JF549 SNAP-Tag ligand was added and left in the

media until the end of the experiment. The live cells were imaged immediately (0 h) and

at 2-h intervals up to 8 h after labeling using a Nikon Ti2 spinning disk confocal

microscope equipped with a $37^0$ C incubator and 5% $CO_2$ supply. The HaloTag

expression was used to identify transfected cells even after complete degradation of

hERG proteins. ImageJ was used to manually segment the cells, and the integrated

density values were exported per cell. Corrected total cell fluorescence (CTCF) was

calculated using the formula: CTCF = Integrated Density – (Area of selected cell X

Mean fluorescence of background). Mean CTCF values were normalized by setting the

baseline mean CTCF at 0 h to 100 for plotting one phase exponential decay.

**Statistical analysis**

Statistical tests were performed using GraphPad Prism software (Dotmatics) with

unpaired Student's t-tests for comparing two groups and one-way ANOVA tests for

comparing multiple groups. Statistical tests for each data set are listed in the figure

legends.

**Acknowledgments**

We thank members of the Robertson lab, especially Catherine Eichel, Lisandra Flores-

Aldama, and Taylor Voelker for critical discussion, Fang Liu for technical support, and

members of the Audhya lab for helpful advice and reagents. Zachary T. Campbell and

Cynthia M. Czajkowski for reviewing the manuscript and providing suggestions. We

thank Luke D. Lavis for providing Janelia Fluor dyes. The work was supported by NIH

grants R01-HL131403 and R01-NS081320 (G.A.R.), American Heart Association

Predoctoral Fellowship 24PRE1188958 (S.K.) and NIH grants R01 NS124165 and R35

GM134865 (A.A).

# References

1.  M. C. Sanguinetti, C. Jiang, M. E. Curran, M. T. Keating, A mechanistic link between an inherited and an acquired cardiac arrhythmia: HERG encodes the IKr potassium channel. *Cell* **81**, 299-307 (1995).
2.  M. C. Trudeau, J. W. Warmke, B. Ganetzky, G. A. Robertson, HERG, a human inward rectifier in the voltage-gated potassium channel family. *Science* **269**, 92-95 (1995).
3.  M. E. Curran *et al.*, A molecular basis for cardiac arrhythmia: HERG mutations cause long QT syndrome. *Cell* **80**, 795-803 (1995).
4.  R. D. Bagnall *et al.*, Exome-based analysis of cardiac arrhythmia, respiratory control, and epilepsy genes in sudden unexpected death in epilepsy. *Ann Neurol* **79**, 522-534 (2016).
5.  S. J. Huffaker *et al.*, A primate-specific, brain isoform of KCNH2 affects cortical physiology, cognition, neuronal repolarization and risk of schizophrenia. *Nature Medicine* **15**, 509-518 (2009).
6.  C. L. Anderson *et al.*, Large-scale mutational analysis of Kv11.1 reveals molecular insights into type 2 long QT syndrome. *Nat Commun* **5**, 5535 (2014).
7.  B. London *et al.*, Two isoforms of the mouse ether-a-go-go-related gene coassemble to form channels with properties similar to the rapidly activating component of the cardiac delayed rectifier K+ current. *Circ Res* **81**, 870-878 (1997).
8.  D. K. Jones *et al.*, hERG 1b is critical for human cardiac repolarization. *Proc Natl Acad Sci U S A* **111**, 18073-18077 (2014).
9.  E. M. C. Jones, E. C. Roti Roti, J. Wang, S. A. Delfosse, G. A. Robertson, Cardiac IKr Channels Minimally Comprise hERG 1a and 1b Subunits. *Journal of Biological Chemistry* **279**, 44690-44694 (2004).
10. H. Sale *et al.*, Physiological Properties of hERG 1a/1b Heteromeric Currents and a hERG 1b-Specific Mutation Associated With Long-QT Syndrome. *Circulation Research* **103**, e81-e95 (2008).
11. J. P. Lees-Miller *et al.*, Selective Knockout of Mouse ERG1 B Potassium Channel Eliminates $I_{Kr}$ in Adult Ventricular Myocytes and Elicits Episodes of Abrupt Sinus Bradycardia. *Molecular and Cellular Biology* **23**, 1856-1862 (2003).
12. J. P. Lees-Miller, C. Kondo, L. Wang, H. J. Duff, Electrophysiological characterization of an alternatively processed ERG K+ channel in mouse and human hearts. *Circ Res* **81**, 719-726 (1997).
13. J. H. Morais Cabral *et al.*, Crystal structure and functional analysis of the HERG potassium channel N terminus: a eukaryotic PAS domain. *Cell* **95**, 649-655 (1998).
14. W. Wang, R. MacKinnon, Cryo-EM Structure of the Open Human Ether-a-go-go-Related K(+) Channel hERG. *Cell* **169**, 422-430 e410 (2017).
15. C. A. Harley *et al.*, Conformation-sensitive antibody reveals an altered cytosolic PAS/CNBh assembly during hERG channel gating. *Proceedings of the National Academy of Sciences* **118**, e2108796118 (2021).

549    16.    P. Phartiyal, H. Sale, E. M. Jones, G. A. Robertson, Endoplasmic reticulum
550           retention and rescue by heteromeric assembly regulate human ERG 1a/1b
551           surface channel composition. *J Biol Chem* **283**, 3702-3707 (2008).
552    17.    P. Phartiyal, E. M. C. Jones, G. A. Robertson, Heteromeric Assembly of Human
553           Ether-à-go-go-related Gene (hERG) 1a/1b Channels Occurs Cotranslationally via
554           N-terminal Interactions. *Journal of Biological Chemistry* **282**, 9874-9882 (2007).
555    18.    G. Kreibich, B. L. Ulrich, D. D. Sabatini, Proteins of rough microsomal
556           membranes related to ribosome binding. I. Identification of ribophorins I and II,
557           membrane proteins characteristics of rough microsomes. *J Cell Biol* **77**, 464-487
558           (1978).
559    19.    F. Liu, D. K. Jones, W. J. de Lange, G. A. Robertson, Cotranslational association
560           of mRNA encoding subunits of heteromeric ion channels. *Proc Natl Acad Sci U S*
561           *A* **113**, 4859-4864 (2016).
562    20.    M. J. McPate *et al.*, hERG1a/1b heteromeric currents exhibit amplified
563           attenuation of inactivation in variant 1 short QT syndrome. *Biochem Biophys Res*
564           *Commun* **386**, 111-117 (2009).
565    21.    Q. Gong, C. L. Anderson, C. T. January, Z. Zhou, Pharmacological rescue of
566           trafficking defective HERG channels formed by coassembly of wild-type and long
567           QT mutant N470D subunits. *Am J Physiol Heart Circ Physiol* **287**, H652-658
568           (2004).
569    22.    R. M. Napier, L. C. Fowke, C. Hawes, M. Lewis, H. R. B. Pelham, Immunological
570           evidence that plants use both hdel and kdel for targeting proteins to the
571           endoplasmic reticulum. *Journal of Cell Science* **102**, 261-271 (1992).
572    23.    D. Gozuacik *et al.*, DAP-kinase is a mediator of endoplasmic reticulum stress-
573           induced caspase activation and autophagic cell death. *Cell Death &*
574           *Differentiation* **15**, 1875-1886 (2008).
575    24.    M. Furutani *et al.*, Novel mechanism associated with an inherited cardiac
576           arrhythmia: defective protein trafficking by the mutant HERG (G601S) potassium
577           channel. *Circulation* **99**, 2290-2294 (1999).
578    25.    Q. Gong, D. R. Keeney, M. Molinari, Z. Zhou, Degradation of trafficking-defective
579           long QT syndrome type II mutant channels by the ubiquitin-proteasome pathway.
580           *J Biol Chem* **280**, 19419-19425 (2005).
581    26.    B. Foo *et al.*, Mutation-specific peripheral and ER quality control of hERG
582           channel cell-surface expression. *Scientific Reports* **9** (2019).
583    27.    P. M. Apaja *et al.*, Ubiquitination-dependent quality control of hERG K$^+$ channel
584           with acquired and inherited conformational defect at the plasma membrane.
585           *Molecular Biology of the Cell* **24**, 3787-3804 (2013).
586    28.    E. B. Rios-Perez *et al.*, A stable cell line inducibly expressing hERG1a/1b
587           heteromeric channels. *J Pharmacol Toxicol Methods* **110**, 107081 (2021).
588    29.    Y. G. Zhao, H. Zhang, Phase Separation in Membrane Biology: The Interplay
589           between Membrane-Bound Organelles and Membraneless Condensates. *Dev*
590           *Cell* **55**, 30-44 (2020).
591    30.    W. R. Legant *et al.*, High-density three-dimensional localization microscopy
592           across large volumes. *Nature Methods* **13**, 359-365 (2016).
593    31.    J. B. Grimm, T. A. Brown, A. N. Tkachuk, L. D. Lavis, General Synthetic Method
594           for Si-Fluoresceins and Si-Rhodamines. *ACS Cent Sci* **3**, 975-985 (2017).

32.  N. O. Taylor, M. T. Wei, H. A. Stone, C. P. Brangwynne, Quantifying Dynamics in
Phase-Separated Condensates Using Fluorescence Recovery after
Photobleaching. *Biophys J* **117**, 1285-1300 (2019).
33.  S. Alberti, A. Gladfelter, T. Mittag, Considerations and Challenges in Studying
Liquid-Liquid Phase Separation and Biomolecular Condensates. *Cell* **176**, 419-
434 (2019).
34.  A. A. Hyman, C. A. Weber, F. Julicher, Liquid-liquid phase separation in biology.
*Annu Rev Cell Dev Biol* **30**, 39-58 (2014).
35.  W. Ma, C. Mayr, A Membraneless Organelle Associated with the Endoplasmic
Reticulum Enables 3'UTR-Mediated Protein-Protein Interactions. *Cell* **175**, 1492-
1506 e1419 (2018).
36.  E. Sezgin, I. Levental, S. Mayor, C. Eggeling, The mystery of membrane
organization: composition, regulation and roles of lipid rafts. *Nature Reviews*
*Molecular Cell Biology* **18**, 361-374 (2017).
37.  C. J. Dirusso, M. Dashtiahangar, T. D. Gilmore, Scaffold proteins as dynamic
integrators of biological processes. *Journal of Biological Chemistry* **298**, 102628
(2022).
38.  M. Hardenberg, A. Horvath, V. Ambrus, M. Fuxreiter, M. Vendruscolo,
Widespread occurrence of the droplet state of proteins in the human proteome.
*Proc Natl Acad Sci U S A* **117**, 33254-33262 (2020).
39.  M. Seidel *et al.*, Co-translational assembly orchestrates competing biogenesis
pathways. *Nature Communications* **13** (2022).
40.  F. Morales-Polanco, J. H. Lee, N. M. Barbosa, J. Frydman, Cotranslational
Mechanisms of Protein Biogenesis and Complex Assembly in Eukaryotes. *Annu*
*Rev Biomed Data Sci* **5**, 67-94 (2022).
41.  C. D. S. Duncan, J. Mata, Widespread Cotranslational Formation of Protein
Complexes. *PLoS Genetics* **7**, e1002398 (2011).
42.  C. D. Duncan, J. Mata, Cotranslational protein-RNA associations predict protein-
protein interactions. *BMC Genomics* **15**, 298 (2014).
43.  J. Clatot *et al.*, Voltage-gated sodium channels assemble and gate as dimers.
*Nature Communications* **8** (2017).
44.  L. Tu, C. Deutsch, Evidence for Dimerization of Dimers in K+ Channel Assembly.
*Biophysical Journal* **76**, 2004-2017 (1999).
45.  A. Patel *et al.*, A Liquid-to-Solid Phase Transition of the ALS Protein FUS
Accelerated by Disease Mutation. *Cell* **162**, 1066-1077 (2015).
46.  S. F. Banani, H. O. Lee, A. A. Hyman, M. K. Rosen, Biomolecular condensates:
organizers of cellular biochemistry. *Nature Reviews Molecular Cell Biology* **18**,
285-298 (2017).
47.  S. Banjade, M. K. Rosen, Phase transitions of multivalent proteins can promote
clustering of membrane receptors. *Elife* **3** (2014).
48.  X. Su *et al.*, Phase separation of signaling molecules promotes T cell receptor
signal transduction. *Science* **352**, 595-599 (2016).
49.  M. Zeng *et al.*, Phase Transition in Postsynaptic Densities Underlies Formation of
Synaptic Complexes and Synaptic Plasticity. *Cell* **166**, 1163-1175 e1112 (2016).
50.  S. Munro, H. R. Pelham, A C-terminal signal prevents secretion of luminal ER
proteins. *Cell* **48**, 899-907 (1987).

51.    L. Orci *et al.*, Bidirectional Transport by Distinct Populations of COPI-Coated
Vesicles. *Cell* **90**, 335-349 (1997).
52.    D. B. Scott, T. A. Blanpied, G. T. Swanson, C. Zhang, M. D. Ehlers, An NMDA
Receptor ER Retention Signal Regulated by Phosphorylation and Alternative
Splicing. *The Journal of Neuroscience* **21**, 3063-3072 (2001).
53.    S.-S. Kim, J. Park, E. Kim, E. M. Hwang, J.-Y. Park, β-COP Suppresses the
Surface Expression of the TREK2. *Cells* **12**, 1500 (2023).
54.    C. Brock, L. Boudier, D. Maurel, J. Blahos, J.-P. Pin, Assembly-dependent
Surface Targeting of the Heterodimeric GABA$_B$ Receptor Is Controlled by COPI
but Not 14-3-3. *Molecular Biology of the Cell* **16**, 5572-5578 (2005).
55.    N. Zerangue, B. Schwappach, Y. N. Jan, L. Y. Jan, A new ER trafficking signal
regulates the subunit stoichiometry of plasma membrane K(ATP) channels.
*Neuron* **22**, 537-548 (1999).
56.    M. C. Sanguinetti, N. K. Jurkiewicz, Two components of cardiac delayed rectifier
K+ current. Differential sensitivity to block by class III antiarrhythmic agents. *J*
*Gen Physiol* **96**, 195-215 (1990).
57.    T. Shibasaki, Conductance and kinetics of delayed rectifier potassium channels
in nodal cells of the rabbit heart. *J Physiol* **387**, 227-250 (1987).
58.    A. Zou, Q. P. Xu, M. C. Sanguinetti, A mutation in the pore region of HERG K[+]
channels expressed in *Xenopus* oocytes reduces rectification by shifting the
voltage dependence of inactivation. *The Journal of Physiology* **509**, 129-137
(1998).
59.    I. M. Herzberg, M. C. Trudeau, G. A. Robertson, Transfer of rapid inactivation and
sensitivity to the class III antiarrhythmic drug E-4031 from HERG to M-eag
channels. *The Journal of Physiology* **511**, 3-14 (1998).
60.    M. C. Sanguinetti, HERG1 channel agonists and cardiac arrhythmia. *Current*
*Opinion in Pharmacology* **15**, 22-27 (2014).
668

## Figure Legends

**Fig. 1.** hERG1b is sequestered in intracellular puncta. (*A*) Confocal images of fixed and immunostained HeLa cells transfected with *hERG1a* or (B) *hERG1b* plasmids; (*C*) quantification of the number of puncta per cell; data are mean ± s.d., n= 36 cells per condition, analyzed with Student's t-test (**** indicating p<0.0001); (*D*) confocal live cell images of iPSC-CM transfected with 1a-GFP or (E) 1b-GFP; (*F*) quantification of number of puncta per cell; data are mean ± s.d., n=18 cells for $1a_{GFP}$, 12 cells for $1b_{GFP}$, analyzed with Student's t-test (**** indicating p<0.0001); Scale bar: 10 μm in large image and 1 μm in the inset.

**Fig. 2.** hERG1b puncta are localized to ER. (*A-C*) STED microscopy of iPSC-CM transfected with mRNAs encoding $hERG1b_{SNAP}$ labeled with JFX554 (magenta) and ER marker Halo-KDEL labeled with JFX650 (cyan); (*D*) quantification of colocalization events thresholded with three different distances between the center of hERG1b puncta and the surface of the ER; data are mean ± s.d., n= 16 regions of interest (see Methods); (*E*) cartoon of ER localization of hERG1b puncta. The scale bar is 10 μm for *(A)* and 500 nm for (*B,C*).

**Fig. 3**. hERG1b is sequestered in a privileged compartment. (*A*) Immunostaining of HeLa cells transfected with $hERG1b_{HA}$ or (*B*) $hERG1a-Y611H_{HA}$; (*C*) quantification of the number of puncta per cell; data are mean ± s.d., n= 33 for hERG1b, n= 26 for 1a-Y611H, analyzed with a Student t-test (**** indicating p<0.0001) ; (*D*) experimental design for the pulse-chase experiment; (*E*) HeLa cells transfected with either $hERG1a_{SNAP}$, $1b_{SNAP}$, $1a-Y611H_{SNAP}$ or $1b-Y271H_{SNAP}$ imaged at various time points

provided in (*D*) and sample images provided for 0 hr and 8 hr after labeling; (*F*)

quantification of normalized corrected total fluorescence intensity (CTCF) plotted across

time; hERG1a-Y611H and hERG1b-Y611H data are fitted with single exponential decay;

hERG1a, hERG1b data are plotted with connected lines; data are mean ± SEM, n= 35-

45 cells per condition; data with individual data points are provided in Fig. S4; (*G*)

quantification of the number of puncta per cell; data are mean ± s.d., n=31 for hERG1b,

n=32 for 1a-Y611H, analyzed with a Student t-test (**** indicating $p < 0.0001$); inset on

the upper right of (*E*) shows the Halo-tag expression (yellow) used to identify the

transfected cells. Scale bar: 10 μm for large images (*A-E*) and 1 μm for insets (*A, B*).

**Fig. 4**. An ER retention motif promotes sequestration of hERG1b in ERSA. (*A*) confocal

images of HeLa cells expressing FLAG-tagged hERG1b alone, (*B*) hERG1b with N[15]XN

mutation, and (*C*) co-expression of WT hERG1a and hERG1b; (*D*) quantification of the

number of puncta per cell; data are mean ± s.d. (n= 30 cells per condition) analyzed

with a one-way ANOVA (**** indicating $p < 0.0001$); scale bar: 10 μm in large image and

1 μm in inset.

**Fig. 5.** Are hERG1b puncta are condensates? (*A*) Confocal images of live HeLa cells

expressing hERG1b (magenta) stained with Potomac-Yellow dye (cyan) labeling

membranes; scale bar: 10 μm in the large image and 2 μm in the insets (i, ii); (*B*)

quantification of percentage of number of hERG1b puncta colocalizing within 250 nm

with Potomac-labeled membranous puncta (n=15 cells); (*C*) confocal images of live

HeLa cell expressing hERG1b$_{GFP}$ with photobleaching of individual puncta imaged

before and after photobleaching; scale bar: 5 μm in large image and 1 μm in insets; (*D*)

quantification of fluorescence intensity of the photobleached puncta before and after

photobleaching; data are mean ± SEM; n=10 puncta from different cells. Data are from

a representative experiment repeated three times; (*E*) GFP tagged hERG1b amino-

terminal domain (1-64 residues) undergoes phase separation, with fluorescence

recovery after photobleaching of individual puncta; (*F*) quantification of normalized

fluorescence intensity after photobleaching; data are mean ± SEM; n= 13 puncta from

different cells in a representative experiment repeated three times; (*G*) droplet-

promoting regions in hERG1b identified by Fuzdrop, which includes the N-terminal

domain. (*H*) Confocal image of HeLa cell expressing $1bNT^{N15XN}_{GFP}$; scale bar: 10 µm in

large image and 1 µm in insets; (*I*) graph showing % cells with puncta in 1bNT-WT and

1bNT-$N_{15}$XN mutant protein-expressing cells; 1bNT-WT exemplar is shown in *E*; n= 3

independent wells per condition, with an average of 50 cells in each well, data are mean

± s.d., analyzed with a Student's t-test (*** indicating $p<0.001$).

**Fig. 6.** Sequestered hERG1b proteins can be rescued by newly expressed 1a. (*A*)

Experimental design; (*B*) mock-transfected cells showing hERG1b puncta; (*C*) 1a-

transfected cells showing hERG1b signal in or near plasma membrane in live cell

imaging; (*D*) quantification of the number of puncta per cell. Data are mean ± s.d., n=

35-42 cells per condition, and analyzed with a Student's t-test (**** indicating $p<0.0001$).

Scale bar: 10 µm in large image and 1 µm in inset.

**Fig. 1.**

**Fig. 2.**

**Fig. 3.**

A  1b_HA

B  1a-Y611H_HA

C  # of puncta

D  Transfection    Label & image    Image
   -24 hrs    0    2    4    6    8 hrs

E    1a_SNAP    1b_SNAP    1a-Y611H_SNAP    1b-Y271H_SNAP

0 hours

8 hours

F  Normalized CTFT vs Hours
   1b
   1a
   1a-Y611H  τ= 3.4 hr
   1b-Y271H  τ= 5.0 hr

G  # of puncta
   1b_SNAP    1b-Y271H_SNAP

**Fig. 4.**

**Fig. 5.**

A

Potomac | 1b_SNAP | Merge

ii

i

B

% Colocalization with Potomac spots

1b_SNAP

C

1b_GFP

Pre-bleach | Bleach | 1 min | 5 min

D

Normalized intensity

Time

E

1bNT_GFP

Pre-bleach | Bleach | 1 min | 3 min

F

Normalized intensity

Time (min)

G

Residue-based droplet-promoting probabilities (p_DP)

$p_{DP}$

1-23    235-267  287-297    516-634    687-699  732-762  779-819

H

1b-NT$^{N15XN}_{GFP}$

I

% of cells with puncta

***

1bNT    1bNT$^{N15XN}$

**Fig. 6.**

A

| Express 1b$_{SNAP}$ for 18 hours | → | Label with fluorescent ligand and block with non-fluorescent ligand | → | Transfect 1a mRNA | → | Image after 24 hours |

B

1b$_{SNAP}$

C

1a$_{GFP}$     1b$_{SNAP}$     Merge

1a transfected

D

Number of 1b puncta

****

Mock trans.    1a-mGFP trans.