Title: Conditional *Dystrophin* ablation causes profound effects on muscle development, neurobehavior, and extracellular matrix pathways

Muthukumar Karuppasamy[1], Katherine G. English[1], James R. Conner[2], Shelby N. Rorrer[1], Michael A. Lopez[1], David K. Crossman[3], Jodi R. Paul[4], Miguel A. Monreal-Gutierrez[5], Karen L. Gamble[4], Karyn A. Esser[4], Jeffrey J. Widrick[2], Louis M. Kunkel[2,6-9], and Matthew S. Alexander[1,10-12]

[1]Department of Pediatrics, Division of Neurology at the University of Alabama at Birmingham and Children's of Alabama, Birmingham, AL 35294

[2]Division of Genetics and Genomics, Boston Children's Hospital, Boston, MA 02115, USA

[3]Department of Genetics at the University of Alabama at Birmingham, Birmingham, AL 35294

[4]Department of Psychiatry and Behavioral Neurobiology, University of Alabama at Birmingham, Birmingham, AL 35294

[5]Department of Physiology and Aging, University of Florida, Gainesville, FL 32610, USA

[6]Department of Pediatrics, Harvard Medical School, Boston, MA 02115, USA

[7]The Stem Cell Program, Boston Children's Hospital, Boston, MA 02115, USA

[8]Harvard Stem Cell Institute, Cambridge, MA 02138, USA

[9]The Manton Center for Orphan Disease Research at Boston Children's Hospital, Boston, MA 02115, USA

[10]UAB Center for Exercise Medicine at the University of Alabama at Birmingham, Birmingham, AL, 35294

[11]UAB Civitan International Research Center (CIRC), at the University of Alabama at Birmingham, Birmingham, AL 35233

[12]UAB Center for Neurodegeneration and Experimental Therapeutics (CNET), Birmingham, AL 35294, USA

Word Count: 5,872
Character Count: 34,742

<u>**Correspondence should be directed to**</u>:
Matthew S. Alexander, PhD
University of Alabama at Birmingham and Children's of Alabama
1900 University Blvd. THT 929 Box 96
Birmingham, AL 35294
Email for correspondence: matthewalexander@uabmc.edu

ORCID IDs:

Muthukumar Karuppasamy: 0000-0003-0034-9950

Katherine G. English: 0000-0003-2757-5157

James R. Conner: 0000-0002-8886-1475

Michael A. Lopez: 0000-0002-3029-0761

David K. Crossman: 0000-0002-0981-169X

Jodi R. Paul: 0000-0002-5409-5944

Miguel A. Monreal-Gutierrez: 0000-0001-8601-4080

Karen L. Gamble: 0000-0003-3813-8577

Karyn A. Esser: 0000-0002-5791-1441

Jeffrey J. Widrick: 0000-0002-5597-1141

Louis M. Kunkel: 0000-0001-6126-0242

Matthew S. Alexander: 0000-0001-7406-5171

Author Contributions: M.K., K.G.E., J.R.C., J.R.P, M.A.M-G., S.N.R., D.K.C., and J.J.W. all performed experiments for the manuscript. M.A.L. and L.M.K. contributed key resources for the manuscript. M.K. M.A.L., K.L.G., K.A.E., J.J.W., and M.S.A. wrote the manuscript. All authors approved of the manuscript prior to its submission.

Competing Interest Statement: L.M.K is a consultant for Pfizer, Dyne Therapeutics, and Myofinity for muscle disease drug therapies. All other authors declare no conflicts of interest.

Classification: Medical Sciences

Keywords: Dystrophin, DMD, muscular dystrophy, skeletal muscle, Dystrophinopathy

Abstract

Dystrophin is an essential protein for regulating the transmission of intracellular force to the

extracellular matrix within the skeletal muscle. The mammalian dystrophin gene (*Dmd*) is X-linked

and has several isoforms with tissue-specific expression, including the large *Dp427m* muscle

transcript found in heart and skeletal muscle, and the *Dp427c* transcript that encodes the brain-

specific dystrophin cerebellar protein. Dystrophin-deficiency results in Duchenne muscular

dystrophy (also abbreviated DMD) in patients and understanding the pathways and tissues that

are affected by dystrophin loss will impact dystrophin replacement gene therapy and exon-

skipping DMD correction strategies. We generated conditional *Dystrophin* knockout mice by

targeting exon 52 of the mouse *Dystrophin* (*Dmd*^{flox52}) locus. We then combinatorially generated

dystrophin constitutive and inducible myofiber knockout (*Dmd* mKO; Dp427m) mice to evaluate

the tissue-specific function of the large skeletal muscle dystrophin isoform (Dp427m). Constitutive

embryonic deletion of the *Dystrophin* gene exclusively in skeletal myofibers resulted in a severe

skeletal muscle myopathy as demonstrated using histopathology, physiological force

assessments, and overall functional assessments compared to the *mdx* mouse model.

Transcriptomic analysis of skeletal myofibers of the *Dmd* mKO mice revealed the dysregulation

of key extracellular matrix and cytokine signaling pathways. Separately, we generated Purkinje

neuron cerebellar dystrophin knockout (*Dmd*:Pcp2 KO; Dp427c) mice that displayed

neurobehavioral deficits in social approach, social memory, and spatial navigation and working

memory. These studies reveal the essential requirement for dystrophin within muscle

development, growth, and regeneration in addition to normal neurobehavioral function.

Significance Statement

Duchenne muscular dystrophy is caused by the lack of a functional dystrophin protein in muscle.

The large dystrophin (Dp427m) isoform is expressed in skeletal, cardiac, and smooth muscle, but

its tissue-specific requirements remain unknown. We generated and characterized a conditional

skeletal muscle knockout mouse (*Dmd* mKO). Constitutive embryonic genetic ablation of skeletal

muscle *Dystrophin* resulted in muscle histopathologies similar to the *mdx* mouse, while postnatal

muscle *Dystrophin* ablation resulted in milder pathologies. Ablating cerebellar *Dystrophin* Dp427c

using a *Pcp2/L7*-Cre driver resulted in sociobehavioral defects. Transcriptomic analysis of the

Dmd mKO mice showed a severe reduction of extracellular matrix and cytokine signaling

pathways. Our study reveals an essential role for skeletal muscle dystrophin and identifies

essential pathways for modulation using dystrophin-replacement therapies.

Introduction

In Duchenne muscular dystrophy (DMD) patients there is a disruption of the dystrophin gene (also

abbreviated *DMD*) that is predominantly expressed in skeletal and cardiac muscle. DMD patients

develop progressive muscle weakness in their teenage years that results in severe muscle

degeneration and cardiorespiratory complications. The *DMD* gene is X-linked, thus DMD

predominantly affects males who lose the ability to walk by their early teenage years. Mammalian

dystrophin protein is an essential protein that is required for normal skeletal muscle function

through its key binding to the actin cytoskeleton and dystrophin-associated protein complex

(DAPC). When fully sequenced, the human *DMD* locus was found to be extremely large as it

spans 2.2 megabases (Mbs) and has multiple coding regions that are essential for the function of

the dystrophin protein(1). The regulation of the dystrophin protein is complex in that the major

protein (Dp427) is expressed in multiple tissues and organs including skeletal muscle, heart, lung,

intestine, and the brain(2). Further complexity of dystrophin transcriptional regulation exists in that

multiple isoforms and *DMD* promotors exist and the regulation of its expression on a tissue-

specific manner has yet to be fully elucidated.

The most widely used Duchenne mouse model is the *mdx* mouse, which harbors an *Dmd* exon

23 nonsense mutation (C>T) at nucleotide 3185 of the large *Dp427m* RNA transcript(3, 4). Several

other N-ethyl-N-nitrosourea (ENU) mutagenesis and targeted knockout mouse strains have been

generated since the identification of the *mdx* mouse, that all show varying muscle phenotypes

based on the DMD mutation type and *Dystrophin* exon(s) targeted(5). Additional DMD mouse and

other animal models have been generated and assessed for their dystrophin expression, mutation

location, muscle pathophysiologies, and their ability to respond to pre-clinical and clinical

corrective drugs. Attempts to generate conditional mouse models have had mixed outcomes due

the large *DMD* gene loci and difficulties in targeting the X chromosome using gene editing. One

study demonstrated that a dual *Dmd* exon 2 and 79 floxed mouse when mated with a tamoxifen-

inducible cardiac and skeletal muscle *MerCreMer* transgene resulted in 25-35% dystrophin gene

excision(6). A previously homologous recombination (HR)-established knockout mouse targeting

dystrophin exon 52 (a known DMD mutation "hot spot" region) resulted in the disruption of the

large Dp427, Dp260, and Dp140 dystrophin isoforms(7). This dystrophin exon 52 model (*mdx52*)

displayed severe muscle degeneration pathologies, in addition to cardiac and neurobehavioral

defects similar to those observed in the *mdx* (*Dmd* exon 23) mouse(7-10). Using this knowledge,

we generated a conditional-ready *Dmd* mouse by inserting loxP sites flanking *Dystrophin* exon 52

(*Dmd^{flox52}*) that allowed for the ablation of the large *Dp427* isoforms upon Cre-mediated excision.

Combinatorial mating with skeletal myofiber and brain transgenic lines allowed us to define the

tissue-specific requirements of the dystrophin protein in skeletal muscle in early and post-natal

muscle growth and regeneration.

Results

Generation and initial characterization of skeletal muscle Dystrophin knockout mice

We performed CRISPR-targeting of the mouse *Dystrophin* exon 52 locus and inserted two loxP

sequences in the intronic region immediately flanking the exon (**Figure 1A**). Upon Cre-mediated

excision of *Dmd* exon 52, the resulting DNA sequence contains an out-of-frame deletion that

impairs the stability of the Dp427, Dp260, and Dp140 transcripts as has been previously shown

in the *mdx52* mouse(11). We mated male *Dmd$^{flox52/Y}$* mice with the constitutive human skeletal

actin (*HSA*)-Cre to ablate expression of the large Dp427m protein within the skeletal myofibers

(**Figure 1B**). We compared our conditional dystrophin knockout mouse model to the *mdx^{5cv}*

mouse that has well-characterized dystrophic histological, molecular, and physiological muscle

symptoms(12). The resulting *Dmd$^{flox52/Y}$*:HSA-Cre+ (*Dmd*:HSA KO) mice have no overt

pathologies at birth, but similar to many *mdx* strains develop a progressive muscle degeneration

starting between 4-6 months of age (**Figure 1C**). Analysis of both fast twitch *Tibialis anterior* (TA)

and slow twitch (soleus) muscles from *Dmd*:HSA KO mice showed increased centralized

myonuclei, large areas of fibrosis, and increased collagen production throughout the myofibers

(**Figure 1C**). *Dmd*:HSA KO diaphragm muscle also revealed hallmark muscle degeneration,

increased centralized myonuclei, and the presence of collagen deposits compared to Cre-

controls (**Figure 1C**). Cross-sectional area (CSA) of *Dmd*:HSA KO TA myofibers showed an

increased frequency of smaller myofibers compared to Cre- controls and similar to those found in

our *mdx^{5cv}* mice (**Figures 1D and 1E**). Quantitative western blot assessments of Dp427m protein

levels in the *Dmd*:HSA KO TA and soleus muscles showed an average of <5% remaining Dp427m

protein in the TA and the soleus muscles (**Figure 1F**). This was further confirmed using

immunofluorescent antibody staining in these muscles showing severely reduced levels of the
large Dp427m protein (**Supplemental Figure S1**).

Skeletal myofiber loss of muscle dystrophin greatly impairs muscle physiological function
We performed a muscle functional analysis of the *Dmd*:HSA KO adult mice by performing an
assessment of muscle performance following TREAT-NMD standard operating procedures
(SOPs) for *mdx* mice(13). *Dmd*:HSA KO mice showed reduced rotarod latency to fall, open field
test (OFT) total distance travelled, and impaired forelimb grip strength (**Figures 2A-2C**). All of
these locomotor and muscle strength metrics were comparable to those of aged-matched *mdx*
mice. *Dystrophin*-deficient mice have impaired treadmill running performance due in part to the
lack of sarcolemma-associated neuronal NOS (nNOS) binding to Dp427 at the spectrin-like
repeats encoded by *Dmd* exons 45-48(14, 15). We performed a standardized treadmill downhill
running assessment of age-matched male WT, *mdx^{5cv}*, *Dmd$^{flox52/Y}$* (Cre negative), and *Dmd*:HSA
KO mice. The *Dmd*:HSA KO and *mdx^{5cv}* mice performed similarly in showing quicker times to
exhaustion, reduced overall distance travelled, and reduced velocity compared to internal cohort
control mice (**Figures 2D-2F**). These findings are consistent with the requirement of nNOS-
binding to skeletal muscle dystrophin as essential for skeletal muscle exercise and running
performance(16, 17). We assessed physiological force in EDL muscles from our *Dmd*:HSA KO
mice to determine if there were functional deficits in the peak force and resistance to high
mechanical strain. Despite their greater mass (**Supplemental Figure S2A**), EDL muscles from
the *Dmd*:HSA KO mice attained similar peak force as EDL's from the *Dmd$^{flox52/Y}$* Cre- controls
(**Supplemental Figure S2B**). This reduction in force per unit physiological cross-sectional area
of muscle (**Supplemental Figure S2C**) indicates a reduction in the quality of the *Dmd*:HSA KO
EDL. Eccentric contractions reduced force of muscles from both groups of mice but this response
was exacerbated in the *Dmd*:HSA KO mice (**Supplemental Figures S2D and S2E**). These results
suggest that skeletal muscle Dystrophin loss has a moderate reduction in myofiber force output
and resistance to mechanical strain in comparison with that of *mdx* mice.

Dystrophin **myofiber KO mice have impaired skeletal muscle regeneration following injury**
We performed a cardiotoxin (ctx)-induced injury to the TA muscles of the *Dmd*:HSA KO and control
Dmd$^{flox52/Y}$ mice to evaluate the regenerative capacity of the skeletal muscle in the absence of the
Dp427m protein. The *Dmd*:HSA KO mice showed a significant amount of fibrosis, collagen, and
necrotic regions within the muscle that was comparable to the *mdx^{5cv}* strain (**Supplemental
Figure S3A**). Immunofluorescent analysis of the myofibers from the *Dmd*:HSA KO mice showed

elevated levels of centralized myonuclei after 21 days post ctx-injury (**Supplemental Figure**
S3B). These results demonstrate the absolute requirement for the dystrophin protein in the
myofiber and suggest that dystrophin from the muscle satellite cells (MuSCs) cannot compensate
for the lack of myofiber dystrophin in the normal regenerative process.

Skeletal muscle-specific *Dystrophin* loss results in cytokine and ECM pathway disruption
We performed RNA-sequencing on the *Dmd*:HSA KO and Cre- control TA muscles to determine
what transcript and signaling pathways were disrupted upon the loss of skeletal muscle
Dystrophin (**Figure 3A**). Ingenuity Pathway Analysis of RNA transcripts revealed several
inflammatory cytokine transcripts that were disrupted as well as extracellular matrix (ECM)
including several *Collagen* isoforms (**Supplemental Figure S4**). Interestingly, several transcripts
known to be disrupted in DMD patients were also disrupted in the *Dmd* HSA KO mice including
an elevation of inducible nitric oxide synthase 2 *(iNos*; *Nos2)* (**Figure 3D**). We noted a strong
downregulation in key immune-responsive cytokines including *Tgfb1* which drives DMD muscle
pathologies in *Dmd* HSA KO mouse muscles that indicates disruption of key cell-cell signaling
that occurs between the skeletal myofiber and the immune cells that reside in the muscle (**Figures**
3B-3D). These transcriptomic analysis confirms and essential transcriptomic regulation of key
signaling pathways that are known regulators of dystrophin pathophysiology.

Postnatal skeletal myofiber *Dystrophin* loss impairs muscle maintenance despite
dystrophin protein stability
We next sought to evaluate the role of skeletal muscle dystrophin in the regulation of adult skeletal
muscle maintenance and regeneration. Subsequently, we mated our conditional *Dmd$^{flox52/Y}$* mice
to the *HSA*-MerCreMer tamoxifen-inducible skeletal myofiber Cre line to temporally ablate
Dp427m expression in the adult mice (**Figure 4A**). Two months after tamoxifen dosing, we
observed 25-30% reduction of total Dp427m protein within the TA skeletal muscles of *Dmd*:HSA-
MerCreMer+ (+ Tam) of *Dmd*:HSA-MerCreMer+ (-Tam) controls (**Figure 4B**). We observed a
similar 20-30% decrease of total Dp427m protein within the gastrocnemius and soleus muscles
in the *Dmd*:HSA-MerCreMer+ (+ Tam) mice (**Figure 4B**). These results are similar to the other
reported conditional *Dmd* mice that showed partial reduction of Dp427mprotein and highlights the
stability of the large dystrophin protein after 2 months of protein turnover(6). A histological
examination of the *Dmd*:HSA-MerCreMer+ (+ Tam) mice showed minimal myofiber degradation
and small pockets of regenerating myofibers compared to controls (**Figure 4C**). Overall myofiber
cross-sectional area (CSA) and minimal Feret diameter analysis revealed smaller clusters of

myofibers in the *Dmd*:HSA-MerCreMer+ (+ Tam) mice compared to no tamoxifen mock controls

(**Figures 4D and 4E**). These findings implicate the large amount of stability of the large dystrophin

Dp427m protein and implicate protein turnover as an essential part of dystrophin-restoration

therapies.

Loss of cerebellar dystrophin impairs neurobehavioral function and results in ASD-like

behaviors

Previous studies from our lab and others in various *mdx* strains demonstrated that the Dp427

cerebellar dystrophin isoform was essential for normal neurobehavior and social development(18-

21). We mated the conditional *Dmd*$^{flox52/Y}$ mice to *L7/Pcp2*-Cre (*Dmd*:Pcp2 KO) mice to ablate the

Dp427c protein in Purkinje neurons of the cerebellum (**Figure 5A**). Isolated cerebellums from

adult *Dmd*:Pcp2 KO mice showed loss of approximately 65% of cerebellar dystrophin protein

compared to Cre- controls (**Figure 5B**; **Supplemental Figure S5**). We performed a series of

neurobehavioral assessments in the *Dmd*:Pcp2 KO mice including social approach and social

novelty between novel and familiar objects and mice (**Figure 5C**). *Dmd*:Pcp2 KO showed

impaired social novelty in object recognition compared to *Dmd*$^{flox52/Y}$ Cre- control mice (**Figures**

5D and 5E). We performed a similar 3-chamber assessment (novel mouse versus familiar mouse)

and the *Dmd*:Pcp2 KO mice showed no preference for either mouse as compared to controls

suggesting that social behavior defects are linked to cerebellar dystrophin protein expression. We

performed both T- and Y-maze assessments in the *Dmd*:Pcp2 KO mice and found no differences

in T-maze alternations compared to Cre- control mice (**Figure 5F**). However, in the Y-maze

assessment, *Dmd*:Pcp2 KO mice had reduced alternations without a change in total arm visits

which together suggest reduced spatial working memory compared to Cre- controls (**Figure 5G**).

Memory tests were also conducted on the *Dmd*:Pcp2 KO mice that displayed impaired time to

find the platform in Morris water maze (**Figure 5H**). These findings support an essential role for

cerebellar Purkinje neuronal expression of *Dystrophin* and demonstrate a functional role for

Dp427 in the regulation of neurobehavior in DMD.

Discussion

The generation of novel dystrophinopathy animal models using CRISPR gene editing and other

next-generation genomic DNA editing tools highlights the growing need for comprehensive animal

models that accurately reflect disease symptoms and pathophysiologies(22). Our conditional

dystrophin knockout model allows for the tissue-specific study of the impact of both dystrophin

loss and when combined with dystrophin-replacement strategies, correct DMD strategies. Newer

dystrophinopathy mouse models have identified the impact of dystrophin loss on muscle disease
symptom progression. The *bmx* (Becker muscular dystrophy; *Dmd* exons 45-47 in-frame deletion)
mouse has been useful in elucidating the slower muscle disease progression that occurs in
patients(23).

Previous attempts to delineate skeletal muscle dystrophin protein expression using AAV-shRNAi
strategies have revealed dystrophin as essential for sarcomeric organization within the muscle
myofiber(24). A study of cultured *mdx* mouse myoblasts revealed a similar downregulation of
Collagen and histone transcripts along with impaired fusion and proliferation markers(25). Indeed,
there may be a significant amount of epigenetic remodeling that occurs which was shown in
embryonic stem (ES) cell cultures of *mdx* mice and human induced pluripotent stem cells
(hiPSCs)(26, 27). Given the alteration of myogenic determinant transcriptional pathways,
consideration of the impact of *Dystrophin* loss on epigenetic influences on myogenic
differentiation is warranted(28). Key autophagy and PTEN/AKT pathways have been shown to be
targets for therapeutic amelioration of dystrophic muscle pathologies(29-31). Questions remain
on the impact and influence of the large dystrophin muscle isoform on muscle satellite cells
(MuSCs) and its regulation of cellular polarity(32, 33). The Dp427 protein is expressed in muscle
satellite cells and it is currently unclear if restoration of dystrophin expression in the MuSC
population could compensate for myofiber dystrophin loss. Therapies to restore MuSC polarity in
DMD MuSCs may be beneficial for restoring the cellular kinetics and polarity defects observed in
DMD muscles(34, 35).

It is evident that the expression of dystrophin protein does occur in the brain and both large and
smaller dystrophin isoforms likely play independent and critical neuronal functions. Clinical studies
have demonstrated that loss of dystrophin is important for normal neurological and
neurobehavioral function as up to one-third of all DMD patients are diagnosed on the ASD
spectrum with symptoms ranging from mild cognitive impairment to severe developmental
delay(36-38). Evaluation of DMD mouse models have shown impaired neurobehavioral
responses in memory, learning, and overall electrophysiological neuronal function(20, 39-41). Our
Dmd:Pcp2 KO mice show similar defects to those in *mdx* mice, and highlight the essential role for
the large brain dystrophin protein expression. These results argue that newer platforms for exon-
skipping drug compounds that can more effectively penetrate the blood-brain-barrier (BBB) may
be required for the systemic restoration of dystrophin expression(42, 43). Our study demonstrates
the absolute requirement for both skeletal muscle and brain dystrophin and warrants further

evaluation of the tissue-specific pathways impacted by dystrophin loss to ultimately ameliorate

dystrophic disease pathologies.

Materials and Methods

Mice:

Wild type (*C57BL/6J*; strain #000664; Jackson Labs) and *mdx^{5cv}* (*C57BL/6J*; strain #002379;

Jackson Labs) used in all experiments were maintained in the Alexander lab colony under

pathogen-free conditions. Constitutive HSA-Cre (*Acta1*-Cre; strain# 006149) and tamoxifen-

inducible *HSA-MerCreMer* (*HSA*-MCM; strain# 031934) were purchased from Jackson Labs (Bar

Harbor, ME) and maintained on a congenic *C57BL/6J* background. The *L7/Pcp2*-Cre mice were

originally purchased from Jackson Labs and maintained on a congenic *C57BL/6J* background

that had been backcrossed greater than 6 generations (strain #006207; Jackson Labs). All mice

had a standard Teklad Global Rodent Diet (Inotiv; West Lafayette, IN) with *ad libitum* access to

food and water. All mouse strains and protocols were approved by the UAB Institutional Animal

Care and Use Committee (IACUC) under the protocol 21393.

Generation of conditional Dystrophin knockout mice:

Conditional *Dystrophin* knockout mice *(Dmd$^{flox52/Y}$*) were generated commercially (Cyagen; Santa

Clara, CA) via CRISPR guide editing of genomic DNA by inserting loxP sites in the intronic regions

flanking mouse *Dystrophin* exon 52 using a 12.2 kb homologous recombination vector. Briefly,

loxP sequences 5'-ATAACTTCGTATAGCATACATTATACGAAGTTAT-3' were inserted in the

intronic regions flanking the mouse *Dmd* exon 52 (*Dystrophin* NCBI transcript: NM_007868.6) in

C57Bl/6J mouse ES cells (mES). Positively-targeted mouse ES cells were screened for neomycin

cassette (later removed via FRT-mediated recombination) and diphtheria toxin subunit A (DTA)

selection. After PCR based confirmation of the presence of the loxP sites and removal of the

neomyocin cassette, the resultant ES cell line was injected into pseudopregnant females. The

subsequent F$_0$ offspring were genotyped for presence of the loxP sites at mouse *Dystrophin* exon

52, and then positively-targeted males were mated to WT (*C57Bl/6*) females to generate F$_1$

offspring. All positively-targeted F$_1$ mice were then maintained on the *C57Bl/6* strain for

subsequent matings. Mice were genotyped for the *Dmd$^{flox52/Y}$* allele using a forward primer "5-

TGACTGCATCTGCATACGTG-3" and reverse "5-GAAGCCTGGTAGGCACCAT-3". Using the

Roche Expand Long Template PCR System (MilliporeSigma; Burlington, MA; Cat# 11681842001)

the wild type band (2226 bp), targeted loxP conditional allele (2401 bp), and Cre-mediated

excised band (992 bp) were used for genotyping following the manufacturer's recommendations.

Later genotyping was performed commercially using Transnetyx probes (Cordova, TN). Isolation

of genomic DNA from TA muscles, cerebellum, and tail DNA were performed using Viagen

DirectPCR Lysis Reagent (Viagen Biotech; Los Angeles, CA; Cat# 102-T).

Histology and Immunofluorescent Imaging

Skeletal muscles were slowly cryofrozen as unfixed tissues in Scigen TissuePlus O.C.T.

Compound using an isopentane (FisherScientific; Cat# AC397221000) and liquid nitrogen bath

as unfixed tissues. Some muscle biopsies were placed in 30% sucrose (MilliporeSigma; Cat#

S0389) sink overnight at 4°C prior to the slow cryofreezing to preserve histological and protein

integrity. Muscles were sectioned on a Leica CM1950 cryotome at 7-10 μm thickness, and

specimens were placed on Fisherbrand Tissue Path Superfrost Plus Gold slides (Fisher Scientific;

Cat# FT4981gplus). H&E staining was performed using Hematoxylin (MilliporeSigma; Cat#

H9627) and Eosin Y (MilliporeSigma; Cat# 318906) solutions in a previously described

protocol(44). Masson's trichrome staining was performed using a Masson's Trichrome Stain Kit

(Polysciences, Inc; Warrington, PA; Cat# 25088-1) and picrosirius red staining was performed

using a Picrosirius Red Stain Kit (Polysciences, Inc; Cat# 24901-500). For immunofluorescent

staining, tissue slides were blocked for one hour in a solution of 10% goat serum/1xPBS and

incubated for one hour at room temperature using a M.O.M kit (Vector Labs; Cat# BMK-2202;

Newark, CA). Tissue slides were set using a VECTORSHIELD Antifade mounting medium with

DAPI kit (Vector Labs; Cat# H-1200-10),

Immunoblotting

Immunoblotting was performed on protein lysates using tissues that were snap frozen in liquid

nitrogen and were homogenized using a bead homogenizer and lysing matrix A beads (MP

Biomedicals; Cat# 1169100-CF) following the manufacturer's protocol. Samples were

homogenized in T-PER™ Tissue Protein Extraction Reagent (ThermoFisher Scientific; Cat#

78510) supplemented with a protease inhibitor cocktail (ThermoFisher Scientific; Cat# J61852-

XF). Protein lysates were quantified using a Pierce BCA Protein Assay Kit (ThermoFisher

Scientific; Cat# 23227), and unless otherwise stated, 50 μg of total whole cell lysate was resolved

on 4–20% polyacrylamide Mini-PROTEAN® TGX™ Precast Protein Gels (BioRad). Following an

overnight transfer onto PVDF membranes (BioRad; Cat# 1620175), samples were blocked with

5% BSA (RPI Corp.; Cat# A30075) and incubated with primary antibody overnight, washed 3

times, and incubated with secondary antibodies for 1 hour at room temperature. Membranes were

washed 4 times for 15 minutes with gentle rocking, before being incubated with SuperSignal™

West Dura Extended Duration Substrate (ThermoFisher Scientific; Cat# 34075). Membranes
were imaged on either a ChemiDoc MP Imaging System (BioRad) or a LI-COR Odyssey CLx
imager.
391

Antibodies

Primary antibodies used in these experiments were validated on dystrophin-deficient tissues or
appropriate control tissues for specificity. Primary antisera used for western immunobloting and
immunofluorescent labeling include: dystrophin all isoforms (Abcam; Cat# ab15277), dystrophin
Dys1 (Leica; Cat# NCL-DYS1), and vinculin (Cell Signaling Technology; Cat# 7076S). Secondary
antibodies used for western immunoblotting included anti-mouse IgG HRP (Cell Signaling
Technology; Cat# 7076S) and anti-rabbit IgG HRP (Cell Signaling Technology; Cat# 7074S).
Secondary fluorescent anti-mouse and anti-rabbit IgG fluorochrome-conjugated antibodies used
include: Alexa Fluor 488 (ThermoFisher Scientific; Cat# A-11001 and A-11008), Alexa Fluor 555
(ThermoFisher Scientific; Cat# A32727 and A32732), and DyLight 633 (ThermoFisher Scientific;
Cat# 35563). Fluorescently labeled sections were imaged on a Nikon A1R-HD25 Confocal
microscope using NIS Elements software, and later processed using Adobe Photoshop (2025
version).
405

Tamoxifen administration

We used a previously-established protocol for the *HSA*-MerCreMer strain of delivering (80 mg/kg
body weight) of (Z)-4-hydroxyl-tamoxifen (4-OHT) (MilliporeSigma; Cat# H7904) mixed with
sunflower seed oil (MilliporeSigma; Cat# S5007) via 5 daily intraperitoneal injections while
alternating left and right sides to avoid injection site reactions (45, 46). Control mice were given
equivalent IP injections of sunflower seed oil. Separate mouse cohorts were assessed at 30 day
intervals from days 0, 30, 60, 90, and 180 days post injection. Mice were monitored for
hepatotoxicity using cytochrome P450 assessments and injection site reactions after
administration.
415

Mice locomotor and functional analysis

Open field testing and mouse locomotor analysis was measured using a previously established 4
individual chamber sensor system protocol and data was analyzed using Ethovision XT software
platform (Noldus; Leesburg, VA)(47, 48). Mice were analyzed for functional analysis but were first
acclimated to the room and open-field chambers one day prior to activity and were given a five
minute additional adaptation period prior to activity recording. Mouse locomotor activity was

recorded for ten minutes with no external stimulation and the experimenter isolated in a separate

room.

Physiological force assessments

Mice were anesthetized with sodium pentobarbital (80 mg/kg) delivery intraperitoneal (IP)

injection. Isolated EDL muscles were suspended in a phosphate buffer equilibrated with 95% O_2,

5% CO_2 (35 °C). Muscle contractions were generated using a 150 ms, supramaximal stimulus

train (200 µs pulses) with the muscle held at its optimal length (L_o) for tetanic tension. Force was

normalized to physiological cross-sectional area assuming a fiber length to muscle length ratio of

0.44 as previously described(47). To evaluate the force-frequency relationship, each muscle was

stimulated at frequencies ranging from 30 to 300 Hz (peak force). Fixed-end force values were

expressed relative to maximal peak force and fit by a sigmoid curve. Changes in the relationships

were evaluated by differences in the inflection point (K, measured in Hz) and slope (H, unitless).

Muscles were then subjected to an active lengthening protocol consisting of the following

sequence: one fix-end contraction, 5 lengthening (eccentric) contractions, and two fixed-end

contractions. All contractions began with the muscle at L_o and were elicited with 150 ms trains at

300 Hz. The lengthening trials consisted of an initial 100 ms fixed-end contraction that allowed

the muscle to rise to peak force followed by a constant velocity stretch at four muscle fiber

lengths/s to a final length of 120% L_o. Force was evaluated at 95 ms of stimulation for both fixed-

end and lengthening trials.

Social neurobehavior and memory experiments

Social approach was determined by placing the test mouse in the middle chamber of a 3-chamber

system and determining if the mouse showed preference for a novel mouse or a novel object

(round ball). The social approach protocol was adapted from our previous protocol along with

other established mouse neurobehavioral assessments in autism spectrum disorder (ASD)

mouse models(18, 49). Mice were evaluated over 10 minute periods after a 10 minute acclimation

period in which the novel mouse and novel object were kept in blocked off chambers. Placement

of the novel mouse and novel object was switched between the two side chambers between

intervals, and the apparatus was cleaned between evaluations. Social novelty was determined by

placing the test mouse in the center chamber and evaluating the time spent in chambers

containing either an unfamiliar mouse (no previous exposure) or a familiar mouse (aged-matched

littermate). The mouse was monitored over a 10 min interval after the 10 minute acclimation

period, and its total movement, velocity, time spent in close interaction with familiar or unfamiliar

mouse were measured. T-maze and Y-maze (Maze Engineers; Skokie, IL) cohort assessments

were performed using previously described with video recordings captured after the mouse was

placed in the maze center(50, 51). Spatial memory and learning evaluation of the mouse cohorts

was performed using the Morris Water Maze (Noldus) from an adapted protocol previously used

in *mdx* mice(52, 53). Experimenter was double-blinded to treatment group and genotype of the

mice in each cage and de-blinded occurred after data analysis.

Skeletal muscle injury

A cardiotoxin-induced injury to the TA was performed by intramuscular injection of 40 µl of 10 µM

cardiotoxin (Millipore Sigma; Cat# 217503) given to adult, aged-matched male mice. The

contralateral TA muscle was used as a sham control and given an equivalent intramuscular

injection of phosphate-buffered saline (1x DPBS; Millipore Sigma; Cat# D8662) given as a control.

Mice were evaluated at day 0 (uninjured), 7, 14, and 21 days post cardiotoxin injury for histological

changes in muscle architecture and myofiber composition.

RNA-sequencing

Total RNA was extracted from TA skeletal muscles from aged-matched male mice using TRIzol

reagent (ThermoFisher Scientific; Cat# 15596026) for extraction following the manufacture's

protocol. RNA with an RNA Integrity Score (RIN) greater than 7.0 was selected for library

amplification. Poly A enrichment using a ribominus kit (ThermoFisher Scientific; Cat# K155002)

was performed on the samples. A total RNA library was amplified from approximately 200-300 ng

of RNA using a NovaSeq 6000 reagent kit v1.5 (Illumina; San Diego, CA; Cat# 20044417) before

being processed on an Illumina NovaSeq 6000 sequencer. Total reads were FASTQ files were

uploaded to the UAB High Performance Computer cluster for bioinformatics analysis with the

following custom pipeline built in the Snakemake workflow system (v5.2.2)(54). Following count

normalization, principal component analysis (PCA) was performed, and genes were defined as

differentially expressed genes (DEGs) if they passed a statistical cutoff containing an adjusted p-

value <0.05 (Benjamini-Hochberg False Discovery Rate (FDR) method) and if they contained an

absolute $\log_2$ fold change >=1. Functional annotation enrichment analysis was performed in the

NIH Database for Annotation, Visualization and Integrated Discovery (DAVID, v6.8) by separately

submitting upregulated and downregulated DEGs. Ingenuity pathway analysis (IPA) was

performed on the data as described(55). A p-value <0.05 cutoff was applied to identify gene

ontology (GO) terms and the FASTQ experimental files have been uploaded to NCBI's Gene

Expression Omnibus under accession number GSE284723.

Quantitative PCR analysis

Total RNA was extracted using TRIzol reagent (ThermoFisher Scientific) and the lysing matrix A beads (MP Biomedicals; Cat# 1169100-CF) as described above. Approximately 1 µg of RNA was then reverse transcribed using SuperScript IV First-Strand Synthesis System (ThermoFisher Scientific; Cat# 18091050). Approximately 0.5 µl of sample was then mixed with using TaqMan™ Fast Advanced Master Mix (ThermoFisher; Cat# 4444556) following the manufacturer's protocol and samples were analyzed on a QuantStudio 5 Real-Time PCR machine (ThermoFisher). The following Taqman real time PCR assays (ThermoFisher Scientific) were used to amplify specific transcripts: *Dmd* (exon 1, Mm03956078_s1; exons 16-17, Mm00464475_m1; exons 50-51, Mm01216958_m1; exons 64-65, Mm00464520_m1), and *Actb1* (Mm02619580_g1).

Statistical Analysis

For most dual cohort comparisons, either a two-tailed student's t-test or chi-squared test for normality was performed for all pairwise comparisons. For longitudinal assessments over time or across multiple cohorts, a one-way analysis of variance (ANOVA) with Least Significant Difference (LSD) was performed for all multiple comparisons. GraphPad Prism version 10 software (Graphpad Software; San Diego, CA) was used for all statistical analyses. An *a priori* hypothesis of $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$ was used for all reported data analyses. All graphs were represented as mean +/- SEM.

Acknowledgments and Funding

We would like to thank members of the Alexander lab and UAB Center for Exercise Medicine for their critical evaluation of our manuscript. We would also like to thank members of the Kunkel, Gamble, and Esser labs for reviewing our manuscript and assistance with data interpretation. M.S.A. is supported by NIH grants from the Office of the Director under award number (U54OD030167), and an R21 grant from National Institute of Neurological Disorders and Stroke (NINDS) (R21NS140497). Additional funding support was provided by the Hope4Gabe foundation. L.M.K. is supported by NIH National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS) grant R01AR064300, and the Bernard F. and Alva B. Gimbel Foundation. K.L.G. is funded by an NIH National Institute on Drug Abuse (NIDA) grants R01DA046096 and R01NS082413. K.A.E. is funded by NIH NIAMS grants (R01AR066082 and R01AR079220). M.A.L. is supported by an NIH NIAMS grant K08NS120812.

Figure Legends

Figure 1. Characterization of a myofiber-specific *Dmd* conditional knockout mouse. A. Schematic outlining the generation of myofiber-specific Dystrophin conditional knockout mice (*Dmd*:HSA KO) by breeding our $Dmd^{flox52/Y}$ mice to human skeletal actin (*HSA*)-Cre transgenic mice. Cre-mediated excision of *Dmd* exon 52 results in an out-of-frame disruption and instability of the Dp427 RNA transcript. **B.** Western immunoblotting for the large Dp427m protein in the TA and soleus muscles of WT (open), $Dmd^{flox52/Y}$ (blue), mdx^{5cv} (red), and *Dmd*:HSA KO (green) adult (6-month-old) male mice. **C.** Densitometry graphs WT, $Dmd^{flox52/Y}$, mdx^{5cv}, and *Dmd*:HSA KO immunoblots taken from TA and soleus muscle lysates and normalized to Vinculin protein. N = 5 separate mice per cohort. **D.** Histochemical staining of TA, soleus, and diaphragm muscles isolated from the WT, $Dmd^{flox52/Y}$, mdx^{5cv}, and *Dmd*:HSA KO male mice. Hematoxylin and eosin (H&E), picrosirius red, and Masson's trichrome histochemical staining performed. Scale bar = 100 μm. **E.** Percent fibrosis (%) in the TA muscles of the four mouse cohorts analyzed. **F.** Percentage (%) centralized myonuclei for the approximately 600 myofibers of the TA muscles of the four mouse cohorts analyzed. **G.** Cross-sectional area (CSA) displaying myofiber sizes (μm^2) shown for each of the four cohorts. Two-way ANOVA with Bonferroni correction performed. P-values of significance are shown: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, and **** $p < 0.001$.

Figure 2. Adult *Dmd* myofiber KO mice have a poor running capacity. A. Rotarod analysis shown for the four mouse cohorts: WT, $Dmd^{flox52/Y}$, mdx^{5cv}, and *Dmd*:HSA KO. **B.** Grip strength measured for mouse forelimbs shown normalized to grams (g) body weight. **C.** Open field testing (OFT) of the four cohorts of mice showing distance traveled in centimeters (cm). **D.** and **E.** Forced downhill treadmill running of the four cohorts of mice with average velocity and time to exhaustion shown. N = 6 mice/cohort. Two-way ANOVA with Bonferroni correction performed. P-values of significance are shown: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, and **** $p < 0.001$.

Figure 3. *Dmd* myofiber KO muscles have disrupted cytokine and ECM signaling pathways. **A.** Volcano plot of the approximately 6,200 dysregulated DEGs in the TA muscles of *Dmd*:HSA KO mice compared to $Dmd^{flox52/Y}$ (HSA-Cre-negative) aged-matched littermate controls. **B.** and **C.** Heatmaps of significantly dysregulated showing significantly elevated (> 2.0 fold) or downregulated (< 2.0 blue) DEGs with an emphasis on cytokine and ECM pathways. D. RNA-seq DeSeq2 FKPM counts of three candidate highly disrupted transcripts that are validated DMD

muscle biomarkers: *Nos2*, *Col4a4*, and *Tgfb1* are shown. Four mice of each cohort (*Dmd$^{flox52/Y}$*-

black; *Dmd*:HSA KO- red) were analyzed for differentially expressed genes (DEGs). Student's t-

test, two tailed used for comparisons. P-values of significance are shown: * $p < 0.05$, ** $p < 0.01$,

*** $p < 0.005$, and **** $p < 0.001$.

Figure 4. Postnatal loss of muscle dystrophin results in mild skeletal muscle pathologies.

A. Schematic for conditional ablation of postnatal dystrophin using a tamoxifen-inducible (± TAM)

strategy. *Dmd$^{flox52/Y}$* mice were mated to the *HSA*-MerCreMer (HSA-MCM) to conditionally ablate

dystrophin (+ TAM; red). *Dmd*:HSA (Cre+) mice given sunflower seed oil (- TAM) were used as

controls (black) After a one-week (5x per week) tamoxifen dosing (80 mg/kg body weight) in 8-

week-old male mice via intraperitoneal injection, mouse cohorts were evaluated at 0 (pre-dosing),

30, 60, 90, and 180 days (d) post deletion. N = 5 mice per cohort and time period for biopsy. **B.**

Densitometry western immunoblot measurements of the large dystrophin (Dp427m) protein

isoform measured in the TA, soleus, and diaphragm muscles. Note the robust persistence of the

large dystrophin protein and low protein turnover in the skeletal muscle tissues. **C.** H&E evaluation

of *Dmd$^{flox52/Y}$*:HSA-MCM (Cre+) mice (± TAM) evaluated at 180 days post ablation. Scale bar =

200 µm. **D.** and **E.** CSA and minimum Feret diameter measured at 180 days in the TA muscles

reveal smaller groupings of myofibers in the *md$^{flox52/Y}$*:HSA-MCM (Cre+) mice (+ TAM) mice

compared to (- TAM) controls. Student's t-test, two tailed used for comparisons. P-values of

significance are shown: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, and **** $p < 0.001$.

Figure 5. Loss of Purkinje neuron brain *Dystrophin* impairs neurobehaviors in mice. A.

Breeding schematic for the *Dmd$^{flox52/Y}$* line to the *L7/Pcp2*-Cre (cerebellar Cre driver) transgenic

line to generate Dmd:Pcp2 KO mice. **B.** Densitometry western immunoblot quantification of the

large dystrophin (Dp427) protein expression in the cerebellum, cortex, and hypothalamus regions

of the brain in the *Dmd$^{flox52/Y}$* (Cre-; open) and *Dmd*:Pcp2 KO (Cre+; grey) cohorts. N = 5 mice per

cohort were analyzed. **C.** Graphics showing the social approach (novel mouse versus novel

object) and social novelty (novel mouse versus familiar mouse) assays performed on the cohorts

of mice in a 3-chamber assay system. **D.** and **E.** Social approach and social novelty time in

chambers (1, 2, and 3) shown between the 3 analyzed mouse cohorts: the *Dmd$^{flox52/Y}$* (open bars),

Dmd:Pcp2 KO (grey bars), and *mdx^{5cv}* (red bars). Time in chamber and time investigating in

minutes (mins) are shown. N = 10 mice per cohort were analyzed with an acclimation period of

equivalent amounts. **F.** and **G.** T- and Y-maze assessments of the 3 mouse cohorts are shown

with spontaneous alternation and the number of total arm visits. **H.** Morris water maze (MWM)

assessments in the 3 cohorts are shown as distance to platform (cm) compared to trial day. Spatial

and reverse learning assessments were performed on separate days. N = 10 adult (6-month-old)

mice were analyzed for each cohort. Two-way ANOVA with Bonferroni correction performed. P-

values of significance are shown: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, and **** $p < 0.001$.

Supplementary Figure Legend

Supplemental Figure S1: Immunofluorescent confirmation of the loss of skeletal muscle

dystrophin. Immunofluorescent imaging of the TA skeletal muscle and the DYS1 (Dp427 large

isoform; purple) along with laminin (green) and DAPI (blue nuclei) co-labeling. Each of the four

mouse cohorts are represented: WT, $Dmd^{flox52/Y}$, mdx^{5cv}, and Dmd:HSA KO. Scale bar = 100 μm.

N = 5 adult (6-month-old) mice were analyzed for each cohort.

Supplemental Figure S2: Adult *Dmd* myofiber KO mice have physiological force deficits.

A. Muscle mass (mg) taken from muscle biopsies from Dmd^{flox52}/Y (Cre-) (black) and

$Dmd^{flox52/Y}$:HSA KO (Cre+) (red) mice shown. **B.** Absolute peak specific muscle force (mN). **C.**

Normalized peak specific muscle force normalized to CSA (kPa). **D.** Post-eccentric contraction

muscle force shown as relative force quantified over contraction number. **E.** Post eccentric relative

force (mN) and Force-Frequency H values are shown between the cohorts. N = 10 mice per

cohort. One way ANOVA with Bonferroni correction is shown. P-values of significance are shown:

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.005$, and **** $p < 0.001$; ns = non-significant.

Supplemental Figure S3: Conditional *Dmd* myofiber KO mice have impaired skeletal

muscle regeneration. A. Histochemical evaluation of the TA muscles 14 and 21 days post

cardiotoxin-induced skeletal muscle injury. H&E, picrosirius red, and Masson's trichrome staining

of the TA muscles showing elevated areas of fibrosis, myofiber degeneration, and necrosis in the

Dmd:HSA mKO mice at 14d and 21d post-ctx induced muscle injury compared to $Dmd^{flox52/Y}$

control mice. Scale bar = 100 μm. **B.** Laminin immunofluorescent imaging showing increased

centralized myonuclei and smaller myofibers in the 14d and 21d post-ctx injured Dmd:HSA mKO

mice. Scale bar = 80 μm. N = 5 mice per cohort.

Supplemental Figure S4: IPA pathway analysis of the conditional *Dmd* myofiber knockout

mice. IPA analysis of the top dysregulated signaling pathways disrupted in the adult TA muscles

of the *Dmd*:HSA mKO mice compared to *Dmd^flox52/Y^* control mice. -log p-values shown with

increased pathways (orange) and downregulated pathways (blue) shown.

Supplemental Figure S5: Confirmation of cerebellar dystrophin Dp427 protein knockdown

in Dmd:Pcp2 isolated cerebellums. Western immunoblot analysis showing knockdown of

cerebellar large dystrophin (Dp427c) in the isolated cerebellums of adult *Dmd*:Pcp2 KO mice. WT

(*C57BL/6J*), *Dmd^flox52/Y^*, and *Dmd*:Pcp2 KO mouse cohorts were analyzed. The Dp140 dystrophin

isoform was also reduced in the *Dmd*:Pcp2 KO mouse cohorts. GAPDH levels were used to

normalize protein lysates for comparative analyses.

References

1.   M. Koenig *et al.*, Complete cloning of the duchenne muscular dystrophy (DMD) cDNA
and preliminary genomic organization of the DMD gene in normal and affected
individuals. *Cell* **50**, 509-517 (1987).
2.   E. P. Hoffman, R. H. Brown, L. M. Kunkel, Dystrophin: the protein product of the
Duchenne muscular dystrophy locus. *Cell* **51**, 919 - 928 (1987).
3.   G. Bulfield, W. G. Siller, P. A. Wight, K. J. Moore, X chromosome-linked muscular
dystrophy (mdx) in the mouse. *Proceedings of the National Academy of Sciences* **81**,
1189-1192 (1984).
4.   P. Sicinski *et al.*, The molecular basis of muscular dystrophy in the mdx mouse: a point
mutation. *Science* **244**, 1578-1580 (1989).
5.   M. Rodrigues, Y. Echigoya, S.-i. Fukada, T. Yokota, Current Translational Research and
Murine Models For Duchenne Muscular Dystrophy. *J Neuromuscul Dis* **3**, 29-48 (2016).
6.   A. Bez Batti Angulski *et al.*, Investigations of an inducible intact dystrophin gene excision
system in cardiac and skeletal muscle in vivo. *Scientific Reports* **10**, 10967 (2020).
7.   E. Araki *et al.*, Targeted Disruption of Exon 52 in the Mouse Dystrophin Gene Induced
Muscle Degeneration Similar to That Observed in Duchenne Muscular Dystrophy.
Biochemical and Biophysical Research Communications **238**, 492-497 (1997).
8.   A. Saoudi *et al.*, Emotional behavior and brain anatomy of the mdx52 mouse model of
Duchenne muscular dystrophy. *Disease Models & Mechanisms* **14** (2021).
9.   S. Shibuya *et al.*, Muscle Plasma Membrane Changes in Dystrophin Gene Exon 52
Knockout Mouse. *Pathology - Research and Practice* **197**, 441-447 (2001).
10.   T. Oliver *et al.*, The glucocorticoid receptor acts locally to protect dystrophic muscle and
heart during disease. *Disease Models & Mechanisms* **17** (2024).
11.   Y. Hashimoto *et al.*, Brain Dp140 alters glutamatergic transmission and social behaviour
in the mdx52 mouse model of Duchenne muscular dystrophy. *Progress in Neurobiology*
216, 102288 (2022).
12.   N. Beastrom *et al.*, mdx5cv Mice Manifest More Severe Muscle Dysfunction and
Diaphragm Force Deficits than Do mdx Mice. *The American Journal of Pathology* **179**,
2464-2474 (2011).
13.   R. Willmann *et al.*, Enhancing translation: Guidelines for standard pre-clinical
experiments in mdx mice. *Neuromuscular Disorders* **22**, 43-49 (2012).

14. K. E. Wells *et al.*, Relocalization of neuronal nitric oxide synthase (nNOS) as a marker
for complete restoration of the dystrophin associated protein complex in skeletal muscle.
Neuromuscular Disorders **13**, 21-31 (2003).
15. Y. Lai *et al.*, Dystrophins carrying spectrin-like repeats 16 and 17 anchor nNOS to the
sarcolemma and enhance exercise performance in a mouse model of muscular
dystrophy. *The Journal of Clinical Investigation* **119**, 624-635 (2009).
16. J. A. Call, J. N. Mckeehen, S. A. Novotny, D. A. Lowe, Progressive resistance voluntary
wheel running in the mdx mouse. *Muscle & Nerve* **42**, 871-880 (2010).
17. K. E. Schill *et al.*, Muscle damage, metabolism, and oxidative stress in mdx mice: Impact
of aerobic running. *Muscle & nerve* **54**, 110-117 (2016).
18. M. S. Alexander *et al.*, Reversal of neurobehavioral social deficits in dystrophic mice
using inhibitors of phosphodiesterases PDE5A and PDE9A. *Translational Psychiatry* **6**,
e901-e901 (2016).
19. M. E. D. Stefano, V. Ferretti, C. Mozzetta, Synaptic alterations as a neurodevelopmental
trait of Duchenne muscular dystrophy. *Neurobiology of Disease* **168**, 105718 (2022).
20. C. Vaillend, R. Chaussenot, Relationships linking emotional, motor, cognitive and
GABAergic dysfunctions in dystrophin-deficient mdx mice. *Human Molecular Genetics*
26, 1041-1055 (2017).
21. A. Saoudi *et al.*, Impact of distinct dystrophin gene mutations on behavioral phenotypes
of Duchenne muscular dystrophy. *Disease Models & Mechanisms* **17** (2024).
22. J. W. McGreevy, C. H. Hakim, M. A. McIntosh, D. Duan, Animal models of Duchenne
muscular dystrophy: from basic mechanisms to gene therapy. *Disease Models &*
Mechanisms **8**, 195-213 (2015).
23. C. R. Heier *et al.*, The X-linked Becker muscular dystrophy (bmx) mouse models Becker
muscular dystrophy via deletion of murine dystrophin exons 45–47. *Journal of Cachexia,*
Sarcopenia and Muscle **14**, 940-954 (2023).
24. M. Ghahramani Seno *et al.*, Transcriptomic analysis of dystrophin RNAi knockdown
reveals a central role for dystrophin in muscle differentiation and contractile apparatus
organization. *BMC Genomics* **11**, 345 (2010).
25. M. R. F. Gosselin *et al.*, Loss of full-length dystrophin expression results in major cell-
autonomous abnormalities in proliferating myoblasts. *eLife* **11**, e75521 (2022).
26. In Y. Choi *et al.*, Concordant but Varied Phenotypes among Duchenne Muscular
Dystrophy Patient-Specific Myoblasts Derived using a Human iPSC-Based Model. *Cell*
Reports **15**, 2301-2312 (2016).
27. L. Caputo *et al.*, Acute conversion of patient-derived Duchenne muscular dystrophy
iPSC into myotubes reveals constitutive and inducible over-activation of TGFβ-
dependent pro-fibrotic signaling. *Skeletal Muscle* **10**, 13 (2020).
28. J. Massenet *et al.*, Epigenetic control of myogenic identity of human muscle stem cells in
Duchenne Muscular Dystrophy. *bioRxiv* 10.1101/2023.04.26.538414,
2023.2004.2026.538414 (2023).
29. Y. Liu *et al.*, PTPN1/2 inhibition promotes muscle stem cell differentiation in Duchenne
muscular dystrophy. *Life Science Alliance* **8**, e202402831 (2025).
30. F. Yue *et al.*, PTEN Inhibition Ameliorates Muscle Degeneration and Improves Muscle
Function in a Mouse Model of Duchenne Muscular Dystrophy. *Molecular Therapy* **29**,
132-148 (2021).
31. M. S. Alexander *et al.*, MicroRNA-486–dependent modulation of DOCK3/PTEN/AKT
signaling pathways improves muscular dystrophy–associated symptoms. *The Journal of*
Clinical Investigation **124**, 2651-2667 (2014).
32. N. A. Dumont *et al.*, Dystrophin expression in muscle stem cells regulates their polarity
and asymmetric division. *Nat Med* **21**, 1455-1463 (2015).

33.   Y. X. Wang *et al.*, EGFR-Aurka Signaling Rescues Polarity and Regeneration Defects in
Dystrophin-Deficient Muscle Stem Cells by Increasing Asymmetric Divisions. *Cell Stem*
Cell 10.1016/j.stem.2019.01.002 (2019).
34.   K. Kodippili, M. A. Rudnicki, Satellite cell contribution to disease pathology in Duchenne
muscular dystrophy. *Frontiers in Physiology* **14** (2023).
35.   U. Gurriaran-Rodriguez *et al.*, Wnt7a is Required for Regeneration of Dystrophic
Skeletal Muscle. *bioRxiv* 10.1101/2024.01.24.577041, 2024.2001.2024.577041 (2024).
36.   C. Pascual-Morena *et al.*, Dystrophin Genotype and Risk of Neuropsychiatric Disorders
in Dystrophinopathies: A Systematic Review and Meta-Analysis. *J Neuromuscul Dis* **10**,
159-172 (2023).
37.   M. Thangarajh *et al.*, Relationships between DMD mutations and neurodevelopment in
dystrophinopathy. *Neurology* **93**, e1597-e1604 (2019).
38.   M. Thangarajh *et al.*, Neurodevelopmental Needs in Young Boys with Duchenne
Muscular Dystrophy (DMD): Observations from the Cooperative International
Neuromuscular Research Group (CINRG) DMD Natural History Study (DNHS). *PLoS*
currents **10**, ecurrents.md.4cdeb6970e54034db54032bc54033dfa54054b54034d54987
(2018).
39.   W.-C. Wu, S. P. Bradley, J. M. Christie, J. R. Pugh, Mechanisms and Consequences of
Cerebellar Purkinje Cell Disinhibition in a Mouse Model of Duchenne Muscular
Dystrophy. *The Journal of Neuroscience* **42**, 2103-2115 (2022).
40.   T. L. Stay, L. N. Miterko, M. Arancillo, T. Lin, R. V. Sillitoe, In vivo cerebellar circuit
function is disrupted in an mdx mouse model of Duchenne muscular dystrophy. *Disease*
Models & Mechanisms 10.1242/dmm.040840, dmm.040840 (2019).
41.   R. Chaussenot *et al.*, Cognitive dysfunction in the dystrophin-deficient mouse model of
Duchenne muscular dystrophy: A reappraisal from sensory to executive processes.
Neurobiology of Learning and Memory **124**, 111-122 (2015).
42.   F. Zarrouki *et al.*, Partial Restoration of Brain Dystrophin and Behavioral Deficits by Exon
Skipping in the Muscular Dystrophy X-Linked (mdx) Mouse. *Annals of Neurology* **92**,
213-229 (2022).
43.   C. Vaillend *et al.*, Rescue of a Dystrophin-like Protein by Exon Skipping In Vivo Restores
GABAA-receptor Clustering in the Hippocampus of the mdx Mouse. *Mol Ther* **18**, 1683-
1688 (2010).
44.   C. Wang, F. Yue, S. Kuang, Muscle Histology Characterization Using H&E Staining and
Muscle Fiber Type Classification Using Immunofluorescence Staining. *Bio Protoc* **7**
(2017).
45.   L. A. Riley *et al.*, The skeletal muscle circadian clock regulates titin splicing through
RBM20. *eLife* **11**, e76478 (2022).
46.   B. D. Harfmann *et al.*, Muscle-specific loss of Bmal1 leads to disrupted tissue glucose
metabolism and systemic glucose homeostasis. *Skeletal Muscle* **6**, 12 (2016).
47.   A. L. Reid *et al.*, DOCK3 is a dosage-sensitive regulator of skeletal muscle and
Duchenne muscular dystrophy-associated pathologies. *Human Molecular Genetics* **29**,
2855-2871 (2020).
48.   Y. M. Kobayashi, E. P. Rader, R. W. Crawford, K. P. Campbell, Endpoint measures in the
mdx mouse relevant for muscular dystrophy pre-clinical studies. *Neuromuscular*
Disorders **22**, 34-42 (2012).
49.   M. Yang, J. L. Silverman, J. N. Crawley, "Automated Three-Chambered Social Approach
Task for Mice" in Current Protocols in Neuroscience. (John Wiley & Sons, Inc., 2011).
50.   C. Vaillend, A. Rendon, R. Misslin, A. Ungerer, Influence of dystrophin-gene mutation on
mdx mouse behavior. I. Retention deficits at long delays in spontaneous alternation and
bar-pressing tasks. *Behav Genet* **25**, 569-579 (1995).

51. S. Engelbeen, A. Aartsma-Rus, B. Koopmans, M. Loos, M. van Putten, Assessment of
Behavioral Characteristics With Procedures of Minimal Human Interference in the mdx
Mouse Model for Duchenne Muscular Dystrophy. *Frontiers in Behavioral Neuroscience*
14 (2021).
52. M. Verhaeg *et al.*, Learning, memory and blood–brain barrier pathology in Duchenne
muscular dystrophy mice lacking Dp427, or Dp427 and Dp140. *Genes, Brain and*
Behavior **23**, e12895 (2024).
53. C. V. Vorhees, M. T. Williams, Morris water maze: procedures for assessing spatial and
related forms of learning and memory. *Nature Protocols* **1**, 848-858 (2006).
54. J. Köster, S. Rahmann, Snakemake—a scalable bioinformatics workflow engine.
Bioinformatics **28**, 2520-2522 (2012).
55. A. Krämer, J. Green, J. Pollard, S. Tugendreich, Causal analysis approaches in
Ingenuity Pathway Analysis. *Bioinformatics* **30**, 523-530 (2014).

Figure 1. Characterization of a myofiber-specific *Dmd* conditional Dmd knockout mouse.

Figure 2: Adult *Dmd* myofiber KO mice have a poor running capacity.

Figure 3: *Dmd* myofiber KO muscles have disrupted cytokine and ECM signaling pathways.

Figure 4. Postnatal loss of muscle dystrophin results in a mild skeletal muscle pathologies.

Figure 5. Loss of Purkinje neuron brain *Dystrophin* impairs neurobehaviors in mice.