1
8 **Antiviral Activity of Rosemary Extract Against Zika Virus in Human Dermal Fibroblasts**
9

10 Jordan T. Masi[1], Robert W.E. Crozier[1], Jeremia M. Coish[1], Natalie J. Hicks[1], Carmela A.
11 Lowes[1], Evangelia Tsiani[1], Dustin Duncan[2], Edward I. Patterson[3,4] & Adam J. MacNeil[1*]
12
13
14 [1]Department of Health Sciences, Brock University, St Catharines, Ontario, L2S 3A1, Canada.
15
16 [2]Department of Chemistry, Brock University, St Catharines, Ontario, L2S 3A1, Canada.
17
18 [3]Department of Biological Sciences, Brock University, St Catharines, Ontario, L2S 3A1, Canada.
19
20 [4]Department of Biology, University of Louisiana at Lafayette, Lafayette, Louisiana 70504, USA.
21
22
23
[*]Correspondence: Dr. Adam J. MacNeil, Department of Health Sciences, Brock University, 1812
Sir Isaac Brock Way, Niagara Region, ON, Canada, L2S 3A1 amacneil@brocku.ca

Abbreviations
CC_{50}, 50% cytotoxic concentration
CA, Carnosic acid
CMC, Carboxymethylcellulose
CO, Carnosol
DENV, Dengue virus
DMEM, Dulbecco's modified Eagle's medium
DMSO, Dimethyl sulfoxide
EV71, Enterovirus 71
HFF-1, Human foreskin fibroblast cells
hRSV, Human respiratory syncytial virus
HSV, Herpes simplex virus
IC_{50}, 50% inhibitory concentration
JEV, Japanese encephalitis virus
MOI, Multiplicity of infection
NS, Non-structural
RT-qPCR, Reverse transcription-quantitative polymerase chain reaction
PBS, Phosphate-buffered saline
Pen/strep, penicillin/streptomycin
PFU, Plaque forming unit
RA, Rosmarinic acid
RD, Human rhabdomyosarcoma
RdRp, RNA-dependent RNA polymerase
RE, Rosemary extract
SI, Selectivity index
WHO, World Health Organization
WNV, West Nile virus
YFV, Yellow fever virus
ZIKV, Zika virus

Abstract

Flaviviruses have increasingly emerged and re-emerged in recent decades, infecting millions of people annually. Zika virus (ZIKV) is particularly concerning due to its associated pathological complications, including microcephaly in newborns and Guillain-Barré syndrome in adults, posing a significant threat to public health. Despite efforts made by the scientific community, no licensed drugs against flaviviruses have been developed. Medicinal plants show promise as a novel source of antiviral agents, as they possess a diverse array of biologically active secondary metabolites, making them potential candidates for therapeutic use. Here, we sought to investigate the antiviral potential of rosemary extract (RE) against ZIKV in human dermal fibroblasts (HFF-1), one of the earliest cellular targets of infection. ZIKV was treated with various concentrations of RE or its major polyphenols, including rosmarinic acid (RA), carnosic acid (CA), and carnosol (CO), and the infectivity of each sample was measured by plaque reduction assay. To evaluate the impact of RE on different stages of the ZIKV replication cycle, HFF-1 cells were treated before, during, and after infection, or the virus was treated before infection. RE exerted potent virucidal activity against ZIKV in both Vero and HFF-1 cells by directly acting on virus particles before infection. Importantly, RE significantly inhibited the later stages of the virus replication cycle by interfering with post-entry mechanisms within the host cell. Moreover, major RE-derived polyphenols CA and CO, but not RA, were shown to significantly reduce infectivity when ZIKV was pre-treated with each compound individually. Overall, RE significantly impairs ZIKV infection *in vitro* by directly interacting with virus particles prior to adsorption and interfering with post-entry processes of the viral replication cycle. This study highlights the antiviral potential of RE and its individual components, wFigurearranting further

77   investigation into the mechanisms underlying their activity and their effects on other medically

78   important flaviviruses.

Introduction

Zika virus (ZIKV) is a mosquito-borne virus that has received significant attention in recent years. ZIKV belongs to the *Flavivirus* genus within the *Flaviviridae* family, which includes dengue virus (DENV), yellow fever virus (YFV), Japanese encephalitis virus (JEV), West Nile virus (WNV), and other well-known human pathogens[1]. ZIKV was originally isolated from a Rhesus macaque monkey in Uganda when studying the ecology of YFV in 1947. After its discovery, ZIKV remained relatively unknown, sporadically causing minor outbreaks and mild febrile illness in few individuals[2,3]. The first major outbreak of ZIKV occurred on Yap Island in 2007, where close to 75% of the population was infected[4]. This was followed by an outbreak in French Polynesia from 2013 to 2014 and the largest recorded outbreak in Brazil in 2015, where approximately 1.3 million people were infected[3,5]. These outbreaks were linked to a significant increase in microcephaly in neonates of infected mothers and Guillain-Barré syndrome in adults, sparking a global panic[6]. Due to its rapid spread and associated neurological complications, ZIKV was declared a Public Health Emergency of International Concern by the World Health Organization (WHO) in February 2016[7].

ZIKV is a positive-sense single-stranded RNA virus with a genome of approximately 11 kb. The genome encodes a single polyprotein that is proteolytically cleaved into three structural proteins, including envelope, pre-membrane, and capsid proteins, along with seven non-structural (NS) proteins, including NS1, NS2A, NS2B, NS3, NS4A, NS4B, and NS5[3,4]. ZIKV is primarily transmitted by female *Aedes aegypti* mosquitoes, which inject the virus into the skin of the host during a blood meal[8]. Throughout this process, ZIKV encounters cells in the epidermal and dermal layers of the skin, including keratinocytes, fibroblasts, Langerhans cells, dendritic cells, and mast cells, representing the first line of defense against the virus[9,10]. Fibroblasts are the

most abundant cell type found in the dermis and are responsible for producing and maintaining

the extracellular matrix[11]. It has been well characterized that fibroblasts are highly permissive to

multiple strains of ZIKV and serve as a primary target for viral replication at the site of

inoculation. Moreover, multiple pro-inflammatory genes are upregulated in fibroblasts infected

with ZIKV, suggesting they contribute to the local inflammatory response in the early stages of

infection[10,12]. Fibroblasts are thus a compelling *in vitro* model for studying potential antiviral

agents against ZIKV.

As of May 2024, 92 countries and territories throughout five of the six WHO regions

have recorded autochthonous mosquito-borne transmission of ZIKV. Additionally, competent

populations of *Aedes aegypti* mosquitoes have been established in 60 countries and territories

globally, raising concerns that autochthonous transmission may be introduced in these areas[3,12].

Despite an extensive worldwide research effort, there are currently no vaccines approved to

prevent ZIKV infection and no specific antivirals available for any flavivirus, leaving supportive

care as the only option to manage symptoms. Given the severe complications associated with

ZIKV infection and the ongoing spread of flaviviruses to new regions, there is an urgent need to

develop effective antiviral agents to control future outbreaks[13–15].

An attractive approach to developing therapeutics is the use of medicinal plants, as they

possess a diverse array of bioactive secondary metabolites[14]. These compounds have been

shaped by millions of years of evolution to provide plants selective advantages, including

protection against oxidative stress, ultraviolet radiation, predators, and infection[16,17]. Historically,

humans have used plants as a source of medicines for treating various diseases, a practice that

continues today[18]. Plants have played a significant role in modern drug discovery, making up

approximately 6% of new molecular entities approved by the U.S. Food and Drug

Administration between the years 2000 and 2016[18,19]. In fact, many drugs currently available on

the market were isolated from plants, including the anti-inflammatory acetylsalicylic acid

(aspirin), the narcotic morphine, and the anti-glaucoma drug pilocarpine[14,20]. In the context of

mosquito-borne infections, quinine, derived from the bark of the cinchona tree, stands out as one

of the most important anti-malarial drugs in history and is listed on the WHO's Model List of

Essential Medicines for severe malaria[21]. Beyond their well-documented antioxidant and anti-

inflammatory properties, several plant-derived secondary metabolites have demonstrated

antiviral activity *in vitro* and *in vivo*, including quercetin and fisetin against DENV2, baicalein

against JEV and influenza virus, and epigallocatechin-3-gallate against herpes simplex virus

(HSV), Epstein Barr virus, and DENV, to name a few[22,23]. However, relatively few studies have

explored the antiviral potential of medicinal plants and their individual components against

ZIKV, highlighting a gap in the field.

Rosmarinus officinalis L. (rosemary) is a medicinal herb belonging to the *Lamiaceae*

family and is native to the Mediterranean region. Rosemary extract (RE) is widely used as a

natural preservative in food processing and has been extensively studied for its antioxidant, anti-

inflammatory, antibacterial, antitumor, neuroprotective, and other therapeutic properties[24–26].

These can be attributed to its rich polyphenolic composition, including phenolic acids like

rosmarinic (RA), ursolic, and caffeic acids, as well as phenolic diterpenoids like carnosic acid

(CA), carnosol (CO), and rosmanol[27]. Although the antibacterial activity of RE has been well

characterized[28], little is known about its antiviral activity. To the best of our knowledge, no

studies have explored the antiviral activity of RE in the context of flavivirus infection, although

it has been shown to be effective against human respiratory syncytial virus (hRSV)[29] and HSV[30–

33]. Additionally, RA, a major polyphenol in RE, was identified as a potential antiviral agent

against ZIKV and DENV *in silico*[34–36]. RA has also been shown to inhibit all four serotypes of

DENV *in vitro*[36] and significantly reduce mortality, viral loads, and pro-inflammatory cytokine

levels in mice infected with a lethal dose of JEV[37]. In the current study, we demonstrate that RE

potently inhibits ZIKV infection by directly interacting with virus particles before infection and

disrupting later stages of the viral replication cycle. Furthermore, we found that CA and CO, but

not RA, exert virucidal activity against ZIKV. These findings highlight the antiviral potential of

RE and its individual components, warranting further investigation into their mechanism of

action and activity against other flaviviruses.

Materials and Methods

Cells and virus

Human foreskin fibroblast cells (HFF-1; ATTC, SCRC-1041) and Vero E6 cells (African green monkey kidney cells; ATTC, CRL-1586) were maintained in Dulbecco's modified Eagle's medium (DMEM; Sigma, D6546) supplemented with 10% Serum Plus II (Sigma, 14009C) and 1% penicillin/streptomycin (pen/strep; Gibco, 15140-122) at 37°C with 5% CO_2. The PRAVBC59 strain of ZIKV (ATCC, VR1843) was propagated in Vero cells. Briefly, Vero monolayers were infected with ZIKV at a multiplicity of infection (MOI) of 0.01 and incubated at 37°C and 5% CO_2 for 1 h. The viral inoculum was removed, and cells were incubated for 5 days in culture medium containing 2% Serum Plus II and 1% pen/strep. Viral stocks were quantified by plaque assay, aliquoted, and stored at -80°C.

Reagent preparation

Rosemary extract was prepared by grinding and passing whole dried *Rosmarinus officinalis* L. leaves (purchased from Sobeys, Mississauga, ON, Canada; product of Turkey) through a mesh sieve. Extraction was performed following protocols established by the US National Cancer Institute. Ground leaves (5 g) were steeped overnight in 30 mL of dichloromethane-methanol (1:1). A slight vacuum was used to collect the filtrate, and a methanol extraction was performed for 30 min. The solvent was removed by rotary evaporation. For antiviral experiments, RE was dissolved in dimethyl sulfoxide (DMSO; Sigma, D2650) at a final concentration of 100 mg/mL and RA (Sigma, R4033), CA (Sigma, C0609), and CO (Sigma, C9617) were dissolved in DMSO at a final concentration of 100 mM. Aliquots were stored protected from light at -20°C until further use. For HPLC analysis, RE was dissolved in DMSO (Sigma, 34869) at 50 mg/mL and

authentic standards of RA (Cayman Chemicals, 70900), CA (Cayman Chemicals, 89820), and

CO (Cayman Chemicals, 89800) were dissolved in DMSO at 5 mg/mL immediately before use.

Cell viability assay

To determine the impact of RE treatment on cell viability, HFF-1 and Vero cells were

seeded in 96-well tissue culture plates and incubated at 37°C with 5% CO_2 until ~90% confluent.

Cells were treated in triplicate with DMEM containing 20 to 200 µg/mL of RE for 1 h at 37°C

with 5% CO_2, using 1% DMSO as a vehicle control. Subsequently, the medium was replaced

with fresh culture medium containing 10% Serum Plus II and 1% pen/strep, and cells were

incubated for either 96 h (HFF-1) or 120 h (Vero) at 37°C with 5% CO_2. These conditions

correspond to cells treated before or during infection or infected with pre-treated ZIKV.

Additionally, HFF-1 cells were incubated with 20 to 200 µg/mL of RE in DMEM with 10%

Serum Plus II and 1% pen/strep for 96 h, corresponding to cells treated after infection in the

time-of-addition analysis. Following incubation, culture medium was replaced with 100 µL of

fresh medium and 10 µL of WST-1 cell proliferation reagent (Roche, 5015944001). After 2 h,

absorbance was measured at 460 nm using a spectrophotometer (Bio-Tek, Synergy HT-1,

191356) and triplicates were averaged. The cytotoxic concentration of RE that reduced viability

by 50% (CC_{50}) was calculated by non-linear regression analysis (GraphPad Prism).

Plaque reduction assay

Plaque reduction assay was used to screen the antiviral activity of RE and three major

polyphenol constituents commonly reported in the literature[38–41], including RA, CA, and CO.

100 µL of ZIKV (290,000 PFU) was treated with 25 to 200 µg/mL of RE or 25 to 100 µM of

each compound individually for 1 h at 37°C with 5% CO_2, using 1% DMSO as a vehicle control.

The infectivity of each sample was then quantified by plaque assay. The inhibitory concentration

that reduced plaque numbers by 50% (IC_{50}) was calculated by non-linear regression analysis

(GraphPad Prism).

Time-of-addition analysis in HFF-1 cells

To determine which stages of infection are targeted by RE, time-of-addition analysis was

performed. Briefly, HFF-1 cells were seeded on 24-well plates and incubated at 37°C with 5%

CO_2 until ~90% confluent. Cells were infected with ZIKV at an MOI of 1 under four different

treatment conditions, with 1% DMSO included as a vehicle control. 1) Cells were pre-treated

with 20 to 100 μg/mL of RE for 1 h at 37°C. The supernatant was removed, and wells were

washed once with 1X phosphate-buffered saline (PBS; Gibco, 10010023) before viral

inoculation. 2) Cells were infected with ZIKV and simultaneously treated with 20 to 100 μg/mL

of RE for 1 h at 37°C. 3) Cells were infected with ZIKV for 1 h at 37°C, the inoculum was

removed, and culture medium containing 20 to 100 μg/mL of RE was added to each well for the

entire post-infection period. 4) ZIKV was incubated with 20 to 100 μg/mL of RE for 1 h at 37°C

before being added to the cells. In all treatment conditions, cells were washed once with PBS

after the 1 h adsorption period to remove unbound ZIKV. Next, 500 μL of culture medium with

10% Serum Plus II and 1% pen/strep was added to each well and cells were incubated at 37°C

and 5% CO_2. Supernatant was collected, aliquoted, and stored at -80°C after 0 and 96 h post-

infection. Viral titers were quantified by plaque assay and genome copies were quantified by

two-step reverse transcription-quantitative polymerase chain reaction (RT-qPCR). The inhibitory

concentration that reduced viral titers or genome copies by 50% (IC_{50}) was calculated by non-

linear regression analysis (GraphPad Prism).

Plaque assay

Vero E6 cells were seeded in 12- and 24-well plates and incubated at 37°C with 5% CO_2

until 90% confluent. Serial 1:10 dilutions were prepared for pre-treated ZIKV stocks (plaque

reduction assay) or supernatant harvested from HFF-1 cells (time-of-addition analysis) 96 h post-

infection, whereas a 1:2 dilution was prepared for samples collected 0 h post-infection. Cells

were infected in duplicate with 200 µL (1:10 dilutions) or 100 µL (1:2 dilutions) of each dilution

in 12- or 24-well plates, respectively. Following a 1 h adsorption period at 37°C, viral inoculum

was removed, and cells were overlaid with 1 mL (12-well) or 0.5 mL (24-well) of a 1:1 mixture

of carboxymethylcellulose (CMC) and DMEM with 2% Serum Plus II and 1% pen/strep. Plates

were incubated at 37°C with 5% CO_2 for 5 days for plaque development, the overlay was

discarded, and wells were washed twice with PBS. Cells were simultaneously fixed and stained

with 1% crystal violet dissolved in 8% formaldehyde and 30% ethanol for 30 min at room

temperature. Excess stain was removed using tap water and plaques were counted. Plaque

forming units per mL (PFU/mL) was calculated for each sample using the following equation:

236
$$\frac{PFU}{mL} = \frac{(\text{Average Number of Plaques})(\text{Dilution Factor})}{\text{Volume of Inoculum}}$$

RNA isolation and two-step RT-qPCR

Viral RNA was isolated from cell-free supernatant collected 96 h post-infection using the

QIAamp Viral RNA Kit (Qiagen, 52904) according to the manufacturer's instructions. cDNA

was generated by reverse transcription using RNA to cDNA EcoDry Premix kits (Takara Bio,

639549) in a SimpliAmp Thermal Cycler (Applied Biosystems, 228001548) and diluted 1:20 in

molecular grade water (Sigma, 693520). The StepOnePlus Real-Time PCR System (Applied

Biosystems, 272004476) was used to perform qPCR with KAPA SYBR Fast Master Mix

(KAPA Biosystems, KM4103) and amplification efficiency optimized primers at a final reaction

volume of 10 µL (1 µL cDNA + 9 µL master mix). The following forward and reverse primer set

was used: 5'-GCAAACGCGGTCGCAAACCT-3' and 5'-TGCTAACGCGAAGCCAGGGT-3'

(Integrated DNA Technologies)[42]. To determine the relative number of ZIKV copies in cell-free

supernatant, a standard curve was generated using ZIKV gBlock gene fragments (Integrated

DNA Technologies).

High-performance liquid chromatography (HPLC)

The polyphenolic composition of RE was analyzed using a Dionex UltiMate 3000

UHPLC system (Thermo Scientific) equipped with a Kinetex C18 column (100 mm × 4.6 mm ×

2.6 µm; Phenomenex, 00D-4462-E0) maintained at 30°C. The mobile phase consisted of (A)

water (Sigma, 270733) containing 1% (v/v) glacial acetic acid (Sigma, AX0074-6) and (B)

acetonitrile (VWR, BDH83639.400) containing 1% (v/v) glacial acetic acid. The following

gradient elution was performed: 0–25 min, 95–5% A; 25–28 min, 5% A; 28–30 min, 5–95% A;

30–35 min, 95% A. RE was dissolved in DMSO at 50 mg/mL and diluted ten-fold in 50:50 (v:v)

A:B to a final concentration of 5 mg/mL for analysis. Authentic standards of RA, CA, and CO

were prepared in DMSO at 5 mg/mL and diluted ten-fold in 50:50 (v:v) A:B to 500 µg/mL,

followed by serial two-fold dilutions. The injection volume was 5 µL, with a flow rate of 1.0

mL/min and detection wavelength of 280 nm. Peaks were identified by comparing retention

times with those of authentic standards. Quantification was performed using the external

standardization method in Chromeleon (v.7.1.1.1127), with standard curves generated by

plotting each concentration (31.75, 62.5, 125, 250, and 500 µg/mL) against the relative peak

area.

Statistical analysis

One-way ANOVA analysis was used to evaluate WST-1 viability assay and RT-qPCR

data, and two-way ANOVA analysis was used to evaluate plaque assay data. Multiple

comparisons were performed using Dunnett's *post-hoc* test where appropriate. CC_{50} and IC_{50}

values were determined by analyzing normalized data (% control) using non-linear regression

(variable slope). Results from all experiments were expressed as mean ± SEM and considered

significant if $p < 0.05$ between the vehicle control (DMSO) and RE or polyphenol treated groups.

Statistical analyses were conducted using GraphPad Prism (v.10.5.0).

Results

Cytotoxicity of RE in cell culture

WST-1 assay was used to evaluate the cytotoxic effects of RE treatment on HFF-1 and Vero cells. Both cell lines were exposed to increasing concentrations of RE (20–200 µg/mL) or an equivalent volume of DMSO (vehicle control). Incubation periods were selected to reflect the conditions used in the different functional assays. Results are expressed as a percentage of the vehicle control (set to 100% viability) for each independent experiment. No difference in viability was observed in HFF-1 cells treated with 20 to 80 µg/mL of RE for 1 h followed by a 96 h incubation compared to DMSO treated cells. However, there was a significant decrease in viability beginning at 100 µg/mL, with a CC_{50} of 120.2 µg/mL (Fig. 1a and 1c). Comparatively, HFF-1 cells treated for the entire 96 h incubation period showed a slight increase in cell viability at 40 µg/mL (p<0.01). There was a significant decrease in viability beginning at 80 µg/mL, with a CC_{50} of 107.3 µg/mL (Fig. 1b and 1d). Vero cells were unaffected by treatment with 20 µg/mL of RE for 1 h followed by a 120 h incubation (Fig. 2a).

RE dose-dependently inhibits ZIKV replication in Vero cells

To test the potential antiviral activity of RE, ZIKV was pre-treated with increasing concentrations of RE (25–200 µg/mL) or an equivalent volume of DMSO for 1 h. The viral titer of each sample was then quantified by plaque assay. Results demonstrated a significant dose-dependent decrease in infectious virions relative to the control (Fig. 2b, p<0.0001 for all tested concentrations). Concentrations of 25, 50, 75, 100, 125, 150, 175, and 200 µg/mL caused a 52.77 ± 5.66%, 69.06 ± 2.88%, 85.63 ± 1.49%, 92.83 ± 0.51%, 93.28 ± 0.63%, 96.74 ± 0.29%, 98.86 ± 0.06%, and 99.55 ± 0.10% reduction in ZIKV titers, respectively (Fig. 2c, IC_{50} = 25.06 µg/mL). Due to the nature of this assay, the selectivity index (SI) was not calculated. The highest

298   concentration of RE that ZIKV was pre-treated with was 200 μg/mL, but this was serially diluted

299   ten-fold before being added to Vero cells. As a result, Vero cells were exposed to a maximum

300   concentration of 20 μg/mL of RE, which was not cytotoxic (Fig. 2a).

301   *RE directly interacts with virus particles and inhibits post-entry mechanisms in fibroblasts*

302   To determine the antiviral potential of RE against ZIKV infection and its effect on

different stages of the viral replication cycle, HFF-1 cells were treated with RE before,

concurrent with, or after viral inoculation, or ZIKV was pre-treated with RE prior to infection.

Increasing concentrations of RE ranging from 20 to 100 μg/mL or an equivalent volume of

DMSO were used in all experiments. Based on what is frequently used in the literature[10,43–46],

cells were infected at an MOI of 1 and cell-free supernatant was collected 0 and 96 h post-

infection, which is when ZIKV titers peak in HFF-1 cells (Supplemental Fig. 1). The collected

supernatants were aliquoted and used for both plaque assay and RT-qPCR analysis. Percent

reductions were calculated relative to the DMSO control group within each respective treatment

condition. There were no statistically significant differences in viral titers between samples

collected 0 h post-infection for all treatment conditions (data not shown). No antiviral effects

were observed in HFF-1 cells treated with RE for 1 h prior to infection (Fig. 3a and 3b).

Treatment of HFF-1 cells during the adsorption period significantly enhanced ZIKV replication,

with viral titers increasing by $195.47 \pm 53.63\%$ ($p<0.0001$), $104.50 \pm 18.36\%$ ($p<0.01$), $98.68 \pm$

27.32% ($p<0.01$), $195.89 \pm 65.67\%$ ($p<0.0001$), and $88.65 \pm 41.33\%$ ($p<0.05$) and genome copy

numbers increasing by $320.97 \pm 36.65\%$ ($p<0.001$), $274.66 \pm 11.34\%$ ($p<0.01$), $247.84 \pm 61.02\%$

($p<0.01$), $303.73 \pm 79.16\%$ ($p<0.01$), and $264.92 \pm 68.04\%$ ($p<0.01$) at concentrations of 20, 40,

60, 80, and 100 μg/mL, respectively, relative to the control. However, these changes did not

follow a consistent trend across increasing concentrations of RE, as there was no clear dose-

response relationship (Fig. 4a and 4b). RE demonstrated the most potent antiviral activity in cells

treated after the 1 h infection period, as infectious virion production was significantly inhibited

by every concentration tested (Fig. 5a, p<0.0001). There was a $70.90 \pm 9.68\%$ and $99.99 \pm$

0.01% reduction in viral titers in cells treated with 20 and 40 μg/mL of RE and a $100 \pm 0.00\%$

reduction in viral titers below the detectable limit in cells treated with 60, 80, and 100 μg/mL of

RE (Fig. 5c, IC_{50} = 18.56 μg/mL, SI = 5.78). Genome copy numbers were also reduced by 95.87

$\pm 2.17\%$ (p<0.0001), $99.60 \pm 0.10\%$ (p<0.0001), $99.61 \pm 0.13\%$ (p<0.0001), and $99.95 \pm 0.03\%$

(p<0.0001) at concentrations of 40, 60, 80, and 100 μg/mL, respectively (Fig. 5b and 5d, IC_{50} =

23.12 μg/mL, SI = 4.64). However, concentrations of 80 μg/mL and above were cytotoxic in

HFF-1 cells treated with RE for 96 h (Fig. 1b). Lastly, pre-treatment of ZIKV with RE for 1 h

prior to infection significantly increased viral titers by $65.99 \pm 44.52\%$ (p<0.05) at 20 μg/mL and

reduced viral titers by $99.70 \pm 0.30\%$ (p<0.0001) at 60 μg/mL and $100 \pm 0.00\%$ (p<0.0001) at 80

and 100 μg/mL (Fig. 6a and 6c, IC_{50} = 56.25 μg/mL, SI = 2.14 μg/mL). Similarly, genome copy

numbers decreased by $55.14 \pm 22.42\%$ (p<0.01), $99.84 \pm 0.12\%$ (p<0.0001), $99.90 \pm 0.05\%$

(p<0.0001), and $99.91 \pm 0.06\%$ (p<0.0001) at concentrations of 40, 60, 80, and 100 μg/mL,

respectively (Fig. 6b and 6d, IC_{50} = 39.63 μg/mL, SI = 3.03 μg/mL). However, 100 μg/mL of RE

was cytotoxic when HFF-1 cells were treated for 1 h followed by a 96 h incubation (Fig. 1a).

CA and CO, but not RA, exhibit virucidal activity against ZIKV in Vero cells

To assess whether three main polyphenols in RE exert similar antiviral effects, ZIKV was

pre-treated with 25 to 100 μM of each compound individually or an equivalent volume of DMSO

for 1 h. The remaining infectivity of each sample was quantified by plaque assay. No effect on

viral replication was observed when ZIKV was pre-treated with RA (Fig. 7a). However, there

was a significant dose-dependent decrease in infectivity with CA (Fig. 8a) and CO (Fig. 9a).

Treatment with CA resulted in a $30.69 \pm 6.52\%$ (p<0.01), $69.50 \pm 5.23\%$ (p<0.0001), and 99.27

$\pm 0.12\%$ (p<0.0001) reduction in viral titers at concentrations of 25, 50, and 100 µM,

respectively, corresponding to an IC_{50} of 34.83 µM (Fig. 8b). Similarly, CO caused a $67.04 \pm$

1.97% (p<0.001), $93.81 \pm 1.64\%$ (p<0.0001), and $98.96 \pm 0.16\%$ (p<0.0001) reduction in viral

titers at concentrations of 25, 50, and 100 µM, respectively, with an IC_{50} of 19.55 µM (Fig. 9b).

HPLC analysis of RE

HPLC analysis was performed to identify and quantify CA, CO, and RA in the RE used

in this study. A representative chromatogram for the identified compounds is shown in Fig. 10,

and the corresponding results are presented in Table 1. The most abundant polyphenol per 1 mg

of RE was CA (78.41 ± 4.64 µg/mg), followed by RA (27.69 ± 0.23 µg/mg) and CO ($11.96 \pm$

0.32 µg/mg), with retention times of 17.62, 7.57, and 15.81 min, respectively. These amounts

correspond to concentrations of 235.86 ± 13.96 µM CA, 76.85 ± 0.64 µM RA, and 36.20 ± 0.97

µM CO when 1 mg of RE is dissolved in 1 mL. Accordingly, at the lowest concentration of RE

tested in the plaque reduction assay (25 µg/mL), ZIKV was exposed to approximately $5.90 \pm$

0.35 µM CA, 1.92 ± 0.02 µM RA, and 0.91 ± 0.02 µM CO.

360
Figure 1: Cell viability of HFF-1 cells treated with increasing concentrations of RE. (a)
HFF-1 cells were incubated with increasing concentrations of RE or a vehicle control (DMSO)
for 1 h followed by 96 h with fresh medium. (b) HFF-1 cells were incubated with increasing
concentrations of RE or a vehicle control (DMSO) for 96 h. Cell viability was determined by a
WST-1 assay. Data are expressed as mean % of control ± SEM for n = 3 independent
experiments. A one-way ANOVA and Dunnett's multiple comparisons test were used to
determine differences in cell viability following RE incubation. **$p<0.01$, ****$p<0.0001$ relative
to HFF-1 cells incubated with DMSO (0 µg/mL). CC_{50} values for HFF-1 cells treated for (c) 1 h
or (d) 96 h were calculated by non-linear regression analysis (inhibitor vs normalized response –
variable slope).

373
Figure 2: RE inhibits ZIKV replication in Vero cells in a dose-dependent manner. (a) Vero
cells were incubated with increasing concentrations of RE or a vehicle control (DMSO) for 1 h
followed by 120 h with fresh medium. Cell viability was determined by a WST-1 assay. Data are
expressed as % of control ± SEM for n = 3 independent experiments. Vero cells were exposed to
a maximum concentration of 20 µg/mL in the plaque reduction assay; thus, the corresponding
representative data is shown. A one-way ANOVA and Dunnett's multiple comparisons test were
used to determine differences in cell viability following RE incubation. (b) As an initial screen,
the virucidal activity of RE was evaluated by plaque reduction assay. ZIKV (290,000 PFU) was
incubated with increasing concentrations of RE or an equal volume of a vehicle control (DMSO)
for 1 h. The viral titer of each sample was then determined by plaque assay. Data are expressed
as mean PFU/mL ± SEM for n = 3 independent experiments, each performed in duplicate. A one-
way ANOVA and Dunnett's multiple comparisons test were used to determine differences in viral
titers following RE treatment. ****p<0.0001 relative to ZIKV incubated with DMSO (0 µg/mL)
for all concentrations tested. (c) The IC_{50} value was calculated by non-linear regression analysis
(inhibitor vs normalized response – variable slope). (d) Plaque assay images were taken using an
Amersham Typhoon Biomolecular Imager and represent wells from one independent experiment
in its entirety.

Figure 3: Treatment of HFF-1 cells with RE before infection does not impact ZIKV
replication. (a) HFF-1 cells were treated with increasing concentrations of RE or a vehicle
control (DMSO) for 1 h and subsequently infected with ZIKV (MOI = 1). Cell-free supernatants
were collected 0 and 96 h post-infection, and the viral titer were quantified by plaque assay. Data
are expressed as mean PFU/mL ± SEM for n = 3 independent experiments, each performed in
duplicate. A two-way ANOVA and Dunnett's multiple comparisons test were used to determine
differences in viral titers following RE treatment relative to HFF-1 cells treated with DMSO (0
μg/mL). (b) Viral RNA was extracted from cell-free supernatants collected 96 h post-infection
and quantified by RT-qPCR. Data are expressed as mean genome copies/mL ± SEM for n = 3
independent experiments, each performed in duplicate. A one-way ANOVA and Dunnett's
multiple comparisons test were used to determine differences in genome copy numbers following
RE treatment relative to HFF-1 cells treated with DMSO (0 μg/mL). (c) Plaque assay images
were taken using an Amersham Typhoon Biomolecular Imager and represent wells from one
independent experiment in its entirety.

Figure 4: Treatment of HFF-1 cells with RE during infection enhances ZIKV replication.
(a) HFF-1 cells were infected with ZIKV (MOI = 1) and simultaneously treated with increasing
concentrations of RE or a vehicle control (DMSO) for 1 h. Cell-free supernatants were collected
0 and 96 h post-infection, and the viral titer of each sample was quantified by plaque assay. Data
are expressed as mean PFU/mL ± SEM for n = 3 independent experiments, each performed in
duplicate. A two-way ANOVA and Dunnett's multiple comparisons test were used to determine
differences in viral titers following RE treatment. *p<0.05, **p<0.01, ****p<0.0001 relative to
HFF-1 cells treated with DMSO (0 µg/mL). (b) Viral RNA was extracted from cell-free
supernatants collected 96 h post-infection and quantified by RT-qPCR. Data are expressed as
mean genome copies/mL ± SEM for n = 3 independent experiments, each performed in
duplicate. A one-way ANOVA and Dunnett's multiple comparisons test were used to determine
differences in genome copy numbers following RE treatment. **p<0.01, ***p<0.001 relative to
HFF-1 cells treated with DMSO (0 µg/mL). (c) Plaque assay images were taken using an
Amersham Typhoon Biomolecular Imager and represent wells from one independent experiment
in its entirety.

Figure 5: Treatment of HFF-1 cells with RE after infection inhibits ZIKV replication. (a)
HFF-1 cells were infected with ZIKV (MOI = 1) for 1 h and subsequently treated with increasing
concentrations of RE or a vehicle control (DMSO) for the entire post-infection period. Cell-free
supernatants were collected 0 and 96 h post-infection and viral titers were quantified by plaque
assay. Data are expressed as mean PFU/mL ± SEM for n = 3 independent experiments, each
performed in duplicate. A two-way ANOVA and Dunnett's multiple comparisons test were used
to determine differences in viral titers following RE treatment. ****$p<0.0001$ relative to HFF-1
cells treated with DMSO (0 µg/mL). (b) Viral RNA was extracted from cell-free supernatants
collected 96 h post-infection and quantified by RT-qPCR. Data are expressed as mean genome
copies/mL ± SEM for n = 3 independent experiments, each performed in duplicate. A one-way
ANOVA and Dunnett's multiple comparisons test were used to determine differences in genome
copy numbers following RE treatment. ****$p<0.0001$ relative to HFF-1 cells treated with
DMSO (0 µg/mL). (c,d) IC_{50} values were calculated by non-linear regression analysis (inhibitor
vs normalized response – variable slope). (e) Plaque assay images were taken using an
Amersham Typhoon Biomolecular Imager and represent wells from one independent experiment
in its entirety.

Figure 6: RE exhibits virucidal activity against ZIKV in HFF-1 cells. (a) ZIKV was

incubated with increasing concentrations of RE or an equal volume of a vehicle control (DMSO)

for 1 h. Subsequently, HFF-1 cells were infected with each sample (MOI = 1) and cell-free

supernatants were collected 0 and 96 h post-infection. Viral titers were quantified by plaque

assay. Data are expressed as mean PFU/mL ± SEM for n = 3 independent experiments, each

performed in duplicate. A two-way ANOVA and Dunnett's multiple comparisons test were used

to determine differences in viral titers following RE treatment. *p<0.05, ****p<0.0001 relative

to HFF-1 cells infected with DMSO treated ZIKV (0 µg/mL). (b) Viral RNA was extracted from

cell-free supernatants collected 96 h post-infection and quantified by RT-qPCR. Data are

expressed as mean genome copies/mL ± SEM for n = 3 independent experiments, each

performed in duplicate. A one-way ANOVA and Dunnett's multiple comparisons test were used

to determine differences in genome copy numbers following RE treatment. **p<0.01,

****p<0.0001 relative to HFF-1 cells treated with DMSO (0 µg/mL). (c,d) IC$_{50}$ values were

calculated by non-linear regression analysis (inhibitor vs normalized response – variable slope).

(e) Plaque assay images were taken using an Amersham Typhoon Biomolecular Imager and

represent wells from one independent experiment in its entirety.

461
Figure 7: RA has no effect on ZIKV infection. (a) The virucidal activity of RA was evaluated
by plaque reduction assay. ZIKV (290,000 PFU) was incubated with increasing concentrations of
RA or an equal volume of a vehicle control (DMSO) for 1 h. Viral titers in each sample were
then determined by plaque assay. Data are expressed as mean PFU/mL ± SEM for n = 3
independent experiments, each performed in duplicate. A one-way ANOVA and Dunnett's
multiple comparisons test were used to determine differences in viral titers following RA
treatment relative to DMSO (0 µg/mL). (b) Plaque assay images were taken using an Amersham
Typhoon Biomolecular Imager and represent wells from one independent experiment in its
entirety.

471
472

Figure 8: CA inhibits ZIKV infection in a dose-dependent manner. (a) The virucidal activity
of CA was evaluated by plaque reduction assay. ZIKV (290,000 PFU) was incubated with
increasing concentrations of CA or an equal volume of a vehicle control (DMSO) for 1 h. Viral
titers in each sample were then determined by plaque assay. Data are expressed as mean PFU/mL
$\pm$ SEM for n = 3 independent experiments, each performed in duplicate. A one-way ANOVA and
Dunnett's multiple comparisons test were used to determine differences in viral titers following
CA treatment. **$p<0.01$, ****$p<0.0001$ relative to ZIKV incubated with DMSO (0 µg/mL). (b)
The IC_{50} value was calculated by non-linear regression analysis (inhibitor vs normalized
response – variable slope). (c) Plaque assay images were taken using an Amersham Typhoon
Biomolecular Imager and represent wells from one independent experiment in its entirety.

Figure 9: CO inhibits ZIKV infection in a dose-dependent manner. (a) The virucidal activity
of CO was evaluated by plaque reduction assay. ZIKV (290,000 PFU) was incubated with
increasing concentrations of CO or an equal volume of a vehicle control (DMSO) for 1 h. Viral
titers in each sample were then determined by plaque assay. Data are expressed as mean PFU/mL
± SEM for n = 3 independent experiments, each performed in duplicate. A one-way ANOVA and
Dunnett's multiple comparisons test were used to determine differences in viral titers following
CO treatment. ***$p<0.001$, ****$p<0.0001$ relative to ZIKV incubated with DMSO (0 µg/mL).
(b) The IC_{50} value was calculated by non-linear regression analysis (inhibitor vs normalized
response – variable slope). (c) Plaque assay images were taken using an Amersham Typhoon
Biomolecular Imager and represent wells from one independent experiment in its entirety.

497
Figure 10: Separation and identification of major polyphenols in RE. Representative HPLC
chromatograms of (a) RE and (b) a standard mixture of CA, CO, and RA. HPLC analysis was
performed using a Kinetex C18 column and chromatograms were recorded at 284 nm. Peaks
were identified by comparing retention times with authentic standards.

Table 1: Quantification of major polyphenols in RE. HPLC analysis was performed using a
Kinetex C18 column and detection at 284 nm. Concentrations of CA, CO, and RA were
determined using the external standardization method with authentic standards. Data are
expressed as mean ± SEM for n = 3 independent experiments.

Compound	Molecular weight (g/mol)	Retention time (min)	Concentration (µg/mg of RE)	Concentration (µM) in 1 mg/mL of RE	Concentration (µM) in 25 µg/mL of RE
CA	332.42	17.62 ± 0.01	78.41 ± 4.64	235.86 ± 13.96	5.90 ± 0.35
CO	330.42	15.82 ± 0.01	11.96 ± 0.32	36.20 ± 0.97	0.91 ± 0.02
RA	360.32	7.57 ± 0.01	27.69 ± 0.23	76.85 ± 0.64	1.92 ± 0.02

Discussion

Although cases of ZIKV have significantly declined since reaching their peak in 2016, the virus continues to circulate at low levels in tropical and subtropical regions, posing an ongoing threat to human health[15]. Predictive models estimate that by 2050, 1.3 billion people may be at risk of ZIKV infection, as rising global temperatures continue to expand the distribution of competent vectors and create conditions suitable for ZIKV transmission[47]. Despite extensive efforts by the global scientific community, there are no vaccines or antiviral treatments against ZIKV currently available, leaving us unprepared for potential future outbreaks[3]. This study aimed to evaluate the antiviral potential of RE against ZIKV in human dermal fibroblasts, one of the earliest cellular targets of infection. Here, we provide novel evidence demonstrating RE exerts potent *in vitro* antiviral effects against ZIKV. RE was found to significantly impair ZIKV infection in Vero cells when the virus was treated before infection (Fig. 2). To assess the effects of RE on the ZIKV replication cycle, HFF-1 cells were treated before, during, and after infection or ZIKV was treated before infection. No antiviral activity was observed when HFF-1 cells were treated before (Fig. 3) or during infection (Fig. 4), however, ZIKV replication unexpectedly increased in the latter condition. There was a significant decrease in viral titers and genome copies when ZIKV was pre-treated with RE, indicating RE may target the initial stages of the viral replication cycle in HFF-1 cells (Fig. 6). RE was most effective when HFF-1 cells were treated post-infection, indicating RE may also target the later stages of the viral replication cycle (Fig. 5). Of the major compounds tested, CA (Fig. 8) and CO (Fig. 9), but not RA (Fig. 7), possess antiviral activity against ZIKV. To the best of our knowledge, this is the first study describing the antiviral activity of RE in the context of flavivirus infection, specifically ZIKV.

To determine the antiviral potential of RE against ZIKV, we employed a plaque reduction assay. This approach is commonly used in the literature[36,48–50] to screen different compounds for

antiviral activity and involves directly treating the virus before infecting a susceptible cell line,

such as Vero. Here, we observed a significant dose-dependent decrease in viral replication with

all concentrations of RE, with the lowest concentration causing a 52.77% reduction in viral titers

(Fig. 2, $IC_{50} = 25.06$ µg/mL). Although very few studies have investigated the antiviral effects of

RE, comparable results have been shown against HSV-1[30,32] and HSV-2[30] in Vero cells.

However, Nolkemper et al. reported IC_{50} values of 0.646 µg/mL against HSV-1 and 1.055

µg/mL against HSV-2, which are much lower than those observed in the current study[30]. This

discrepancy can be potentially attributed to differences between the viruses and procedures used.

For example, Nolkemper et al. pre-treated 200 PFU of HSV before adding the entire inoculum to

RC-37 cells (African green monkey kidney cells), whereas we pre-treated 290,000 PFU of ZIKV

and infected Vero cells with serially diluted inoculum (10^{-1} to 10^{-3}). Furthermore, variations in

extraction solvents (water vs. methanol in our study) can substantially influence the chemical

composition and corresponding bioactivity of RE[51]. Overall, our results are in general agreement

with the literature, as several classes of natural products with anti-ZIKV activity exert a virucidal

effect by directly interacting with virus particles or exhibit biological effects during viral

adsorption and internalization[22].

To identify which stage(s) of the ZIKV replication cycle are affected by RE, we

performed four experiments using HFF-1 cells: 1) treatment of cells before infection, 2)

simultaneous treatment and infection of cells, 3) treatment of cells after infection, and 4)

treatment of ZIKV before infection. Fibroblasts are among the first cells that encounter ZIKV at

the site of inoculation, therefore, HFF-1 cells were selected as a relevant infection model. Results

show that treating cells before or during the adsorption phase does not impair ZIKV replication

(Fig. 3 and 4). Nolkemper et al. found similar results, with no effect on HSV-2 and a minimal

24% reduction in HSV-1 replication when RC-37 cells were treated with a non-cytotoxic

concentration of RE before infection. However, treatment of RC-37 cells during infection caused

a 36% and 67% reduction in HSV-1 and HSV-2 replication, respectively[30]. In another study, 20

µg/mL RE caused a 66-fold reduction in hRSV production in A549 cells treated before infection,

conflicting with our results, though a different virus and cell type[29]. Moreover, Lin $et\ al.$

demonstrated a significant 21.9% reduction in viral VP1 mRNA and an 88.6% reduction in

EV71 replication in human rhabdomyosarcoma (RD) cells treated with 10 µM of RA, one of the

main constituents found in RE, during infection. Similar to our findings, RA minimally inhibited

EV71 by 5.54% in RD cells treated before infection[52]. Surprisingly, we observed a statistically

significant increase in ZIKV replication in cells treated during infection with every concentration

of RE (Fig. 4), as well as in cells infected with ZIKV pre-treated with 20 µg/mL (Fig. 5). As far

as we know, this effect has not been reported in the literature. Considering that pH can have a

significant impact on ZIKV infectivity[53], we measured the pH of various concentrations of RE

prepared in DMEM. No major changes in pH were observed in comparison to DMSO, with all

values falling within a range of approximately 8.2 to 8.5 (data not shown). Therefore, the

increase in virus production may have occurred due to an experimental artefact, but further

investigation is required to determine if underlying mechanisms are at play.

Unlike the results above, RE was shown to exert potent antiviral activity when cells were

treated post-infection. There was a considerable 70.90% decrease in infectious virion production

(IC_{50} = 18.56 µg/mL, SI = 5.78) at the lowest concentration tested (20 µg/mL) and a 95.87%

decrease in genome copy numbers (IC_{50} = 23.12 µg/mL, SI = 4.64) at 40 µg/mL, with viral titers

falling below the detectable limit at 60 µg/mL and beyond (Fig. 5). This is in agreement with

Horvath et al., who found a 90.46% reduction in HSV-2 genome concentrations in Vero cells

treated with 0.47 mg/mL of RE after infection[33]. Similarly, Al-Megrin *et al.* reported that 30 and

40 µg/mL of RE reduces plaque formation of HSV-1 by 55% and HSV-2 by 65%, respectively,

when incorporated into the overlay[32]. In contrast to these findings, RE had no effect on HSV-1 or

HSV-2 plaque formation in RC-37 cells[30]. Of the main polyphenols found in RE, CA has been

shown to reduce HSV-2[33] and hRSV[29] genome concentrations in cells treated post-infection,

whereas EV71[52] mRNA and progeny virus production were reduced in RA treated cells, further

supporting our results. Based on an *in-silico* study, RE may interfere with the NS5 RNA-

dependent RNA polymerase (RdRp), which is essential for replicating the ZIKV genome. This is

due to inhibitory potential of CA (-6.8 kcal/mol), CO (-7.0 kcal/mol), RA (6.9 kcal/mol), and

several other compounds, which were predicted to have high binding energy against the RdRp,

although *in vitro* studies are required to validate these data[26]. Collectively, these data suggest that

RE may act on the later stages of the viral replication cycle, however, further research is

necessary to determine the underlying mechanisms, such as effects on gene expression or virion

assembly and release.

As seen with the plaque reduction assay, pre-treatment of ZIKV with RE significantly

inhibited infection in HFF-1 cells, resulting in a 99.70% reduction in progeny virions at a

concentration of 60 µg/mL and complete inhibition at 80 µg/mL (Fig. 6, IC_{50} = 56.25 µg/mL, SI

= 2.14). Viral RNA was also significantly reduced by 55.14%, 99.84%, 99.90%, and 99.91% at

concentrations of 40, 60, 80, and 100 µg/mL, respectively (IC_{50} = 39.63 µg/mL, SI = 3.03

µg/mL). Similar results were reported for HSV pre-treated with the maximum non-cytotoxic

concentration of RE (exact concentration not specified) in RC-37 cells, resulting in 97% and

99% reductions in HSV-1 and HSV-2, respectively[30]. Unlike our plaque reduction assay results,

no decrease in viral titers was observed when ZIKV was pre-treated with concentrations below

60 µg/mL, which may be due to the distinct species and tissue origin of each cell line, as well as

fundamental differences between the assays. For example, ZIKV was treated with RE for 1 h and

the infectivity of each sample was quantified by plaque reduction assay, whereas in the time-of-

addition analysis, plaque assay was used to quantify the release of infectious virions by HFF-1

cells after infection with pre-treated ZIKV. Since no antiviral effects were observed in HFF-1

cells treated during infection, these findings indicate that RE directly interacts with extracellular

ZIKV particles by potentially binding to viral proteins involved in mediating viral entry or

damaging virions, thereby impairing their ability to infect host cells.

To determine whether three major polyphenols in RE exert anti-ZIKV activity, each

compound was individually screened using a plaque reduction assay. Pre-treatment of ZIKV

with CA and CO significantly reduced ZIKV plaque formation in a dose-dependent manner, with

IC_{50} values of 34.83 and 19.55 µM, respectively (Fig. 8 and 9). Although both compounds are

phenolic diterpenoids bearing a catechol moiety, CA carries a carboxyl moiety, whereas CO

carries a lactone moiety[54]. These structural differences suggest that the lactone moiety may

contribute to the lower IC_{50} of CO. In comparison, RA had no effect on ZIKV infection at any

concentration tested (Fig. 7). This was somewhat unexpected, as RA is the only compound with

existing evidence of antiviral activity against flaviviruses. Pre-treatment of DENV with RA has

been shown to inhibit all four serotypes *in vitro*[36], as well as significantly reduce mortality, viral

mRNA, and proinflammatory cytokine levels in mice peritoneally treated with RA and infected

with a lethal dose of JEV[37]. There is also *in vitro* evidence demonstrating the efficacy of RA

against the influenza A virus H1N1[48], EV71[52,55], and hepatitis B virus[56], as well as *in vivo*

evidence against a lethal dose of EV71 in mice[57]. Based on the structural similarities among

flaviviruses, we anticipated that RA would exert antiviral activity against ZIKV. Its failure to do

so indicates that its effects may be specific to DENV and JEV. The antiviral activity of CA has

also been well characterized, as it was shown to be effective against HSV-2[33], hRSV[29], and

influenza A virus H1N1[48]. However, another study reported no effect against influenza A virus

H1N1[29]. Interestingly, we did not anticipate CO to have such a profound effect, as only two

groups have analyzed its antiviral properties with conflicting results[29,58]. HPLC analysis revealed

that the lowest concentration of RE tested by plaque reduction assay (25 µg/mL), which caused a

52.77% reduction in viral titers (Fig. 2), contained approximately 5.90 µM CA and 0.91 µM CO

(Table 1). These concentrations are substantially lower than the IC_{50} values determined for each

compound individually, suggesting that additional compounds within RE, or synergistic

interactions between compounds, underlie its antiviral activity and warrant further investigation.

There are several limitations worth considering. Many antiviral studies include a positive

control to validate assay performance and ensure the proper interpretation of results. Since no

clinically approved antiviral exists for ZIKV, we did not include a positive control, although a

repurposed drug with *in vitro* anti-ZIKV activity could theoretically serve this purpose.

Secondly, depending on soil characteristics, climate, light exposure, and other growing

conditions, the chemical composition of plant extracts may vary, limiting their reproducibility.

To address this, studies should report the composition of each extract, where possible, to

facilitate comparisons across the literature. Lastly, while *in vitro* studies provide valuable

insights and are necessary during the early stages of investigation, the observed effects may not

directly translate to humans. The human body consists of many diverse cell types interacting

within complex tissues and organ systems, and antiviral efficacy can be influenced by

metabolism, immune responses, bioavailability, and other factors. Therefore, it is important to

evaluate antivirals in small animal models, as these are more physiologically relevant. However,

a lack of immunocompetent animal models for ZIKV and other flaviviruses remains a challenge

that must be addressed moving forward.

As mosquitoes and flaviviruses continue to expand into new regions and increasingly

threaten human health, new antiviral agents are urgently needed. In this study, we demonstrate

that RE exhibits potent antiviral activity against ZIKV by directly interacting with virus particles

before infection in Vero and HFF-1 cells and interfering with post-entry processes of the ZIKV

replication cycle in HFF-1 cells. Screening three major polyphenols found in RE revealed that

CA and CO significantly reduce ZIKV infectivity and are promising candidates for follow-up

studies. Whether RE can also inhibit other medically important flaviviruses, such as DENV and

WNV, remains to be determined. Collectively, these findings provide novel evidence supporting

the antiviral potential of RE and its individual components, warranting further investigation into

their mechanisms of action and effects in additional cell and viral models.

Authorship Contributions

JTM, RWEC, JMC, EIP, and AJM conceived and designed the study. RWEC, JMC, and DD

assisted with experimental planning. JTM performed the experiments with assistance from

RWEC, NJH, and CAL. JTM and AJM analyzed the data and wrote the manuscript. All authors

contributed to discussion and editing of the manuscript.

Funding

This research has been supported and funded by grants and awards to AJM from the Natural

Sciences and Research Council of Canada (NSERC Discovery), Canada Foundation for

Innovation (CFI), the Ontario Research Fund (ORF) and Brock University. JTM was supported

by an NSERC Master's Canada Graduate Scholarship (CGS-M) and Queen Elizabeth II

Graduate Scholarship in Science and Technology (QEII-GSST); RWEC and JMC were

supported by an NSERC Doctoral Canada Graduate Scholarship (CGS-D) and NJH was

supported by an NSERC Doctoral Postgraduate Scholarship (PGS-D) and a QEII-GSST.

Conflicts of interests and disclosures

The authors declare no conflicts of interest.

References

1. Vázquez-Calvo, Á., Jiménez de Oya, N., Martín-Acebes, M. A., Garcia-Moruno, E. & Saiz, J.-C. Antiviral Properties of the Natural Polyphenols Delphinidin and Epigallocatechin Gallate against the Flaviviruses West Nile Virus, Zika Virus, and Dengue Virus. *Front. Microbiol.* **8**, (2017).

2. Noorbakhsh, F. *et al.* Zika Virus Infection, Basic and Clinical Aspects: A Review Article. *Iran. J. Public Health* **48**, 20–31 (2019).

3. Pielnaa, P. *et al.* Zika virus-spread, epidemiology, genome, transmission cycle, clinical manifestation, associated challenges, vaccine and antiviral drug development. *Virology* **543**, 34–42 (2020).

4. Duffy, M. R. *et al.* Zika Virus Outbreak on Yap Island, Federated States of Micronesia. *N. Engl. J. Med.* **360**, 2536–2543 (2009).

5. Zhao, R. *et al.* Flavivirus: From Structure to Therapeutics Development. *Life* **11**, 615 (2021).

6. Laureti, M., Narayanan, D., Fazakerley, J. K. & Kedzierski, L. Flavivirus Receptors: Diversity, Identity, and Cell Entry. *Front. Immunol.* **9**, (2018).

7. Gorshkov, K. *et al.* Zika Virus: Origins, Pathological Action, and Treatment Strategies. *Front. Microbiol.* **9**, 3252 (2019).

8. Bustos-Arriaga, J. *et al.* Soluble mediators produced by the crosstalk between microvascular endothelial cells and dengue-infected primary dermal fibroblasts inhibit dengue virus replication and increase leukocyte transmigration. *Immunol. Res.* **64**, 392–403 (2016).

9. Garcia, M., Wehbe, M., Lévêque, N. & Bodet, C. Skin innate immune response to flaviviral infection. *Eur. Cytokine Netw.* **28**, 41–51 (2017).

10. Hamel, R. *et al.* Biology of Zika Virus Infection in Human Skin Cells. *J. Virol.* **89**, 8880–8896 (2015).

11. Dick, M. K., Miao, J. H. & Limaiem, F. Histology, Fibroblast. *StatPearls* (2024).

12. Zika epidemiology update - May 2024. https://www.who.int/publications/m/item/zika-

epidemiology-update-may-2024 (2024).

13. Komarasamy, T. V., Adnan, N. A. A., James, W. & Balasubramaniam, V. R. Finding a chink

in the armor: Update, limitations, and challenges toward successful antivirals against

flaviviruses. *PLoS Negl. Trop. Dis.* **16**, e0010291 (2022).

14. Sadeer, N. B. *et al.* Secondary metabolites as potential drug candidates against Zika virus, an

emerging looming human threat: Current landscape, molecular mechanism and challenges

ahead. *J. Infect. Public Health* **16**, 754–770 (2023).

15. Musso Didier, Ko Albert I., & Baud David. Zika Virus Infection — After the Pandemic. *N.*

Engl. J. Med. **381**, 1444–1457 (2019).

16. Grigalunas, M., Brakmann, S. & Waldmann, H. Chemical Evolution of Natural Product

Structure. *J. Am. Chem. Soc.* **144**, 3314–3329 (2022).

17. Annunziata, G. *et al.* May Polyphenols Have a Role Against Coronavirus Infection? An

Overview of in vitro Evidence. *Front. Med.* **7**, (2020).

18. Dzobo, K. The Role of Natural Products as Sources of Therapeutic Agents for Innovative

Drug Discovery. *Compr. Pharmacol.* 408–422 (2022).

19. Patridge, E., Gareiss, P., Kinch, M. S. & Hoyer, D. An analysis of FDA-approved drugs:

natural products and their derivatives. *Drug Discov. Today* **21**, 204–207 (2016).

20. Dias, D. A., Urban, S. & Roessner, U. A Historical Overview of Natural Products in Drug

Discovery. *Metabolites* **2**, 303–336 (2012).

21. Zhou, B. & Yue, J.-M. Natural products are the treasure pool for antimalarial agents. *Natl.*

Sci. Rev. **9**, nwac112 (2022).

22. Pereira, R. S. *et al.* Natural Products and Derivatives as Potential Zika virus Inhibitors: A

Comprehensive Review. *Viruses* **15**, 1211 (2023).

23. Lee, J. L., Loe, M. W. C., Lee, R. C. H. & Chu, J. J. H. Antiviral activity of pinocembrin

against Zika virus replication. *Antiviral Res.* **167**, 13–24 (2019).

24. Shiravi, A. *et al.* Rosemary and its protective potencies against COVID-19 and other

cytokine storm associated infections: A molecular review. *Mediterr. J. Nutr. Metab.* **14**, 401–

416 (2021).

25. Harlina, P. W., Ma, M., Shahzad, R. & Khalifa, I. Effect of Rosemary Extract on Lipid

Oxidation, Fatty Acid Composition, Antioxidant Capacity, and Volatile Compounds of Salted

Duck Eggs. *Food Sci. Anim. Resour.* **42**, 689–711 (2022).

26. Activity of Phytochemical Constituents of Black Pepper and Rosemary Against Zika Virus:

An In silico Approach. *Lett. Appl. NanoBioScience* **11**, 3393–3404 (2021).

27. Zhang, L. & Lu, J. Rosemary (*Rosmarinus officinalis* L.) polyphenols and inflammatory

bowel diseases: Major phytochemicals, functional properties, and health effects. *Fitoterapia*

177, 106074 (2024).

28. Nieto, G., Ros, G. & Castillo, J. Antioxidant and Antimicrobial Properties of Rosemary

(Rosmarinus officinalis, L.): A Review. *Medicines* **5**, 98 (2018).

29. Shin, H.-B. *et al.* Antiviral activity of carnosic acid against respiratory syncytial virus. *Virol.*

J. **10**, 303 (2013).

30. Nolkemper, S., Reichling, J., Stintzing, F. C., Carle, R. & Schnitzler, P. Antiviral Effect of

Aqueous Extracts from Species of the Lamiaceae Family against Herpes simplex Virus Type

1 and Type 2 in vitro. *Planta Med.* **72**, 1378–1382 (2006).

31. Mancini, D. A. P., Torres, R. P., Pinto, J. R. & Mancini-Filho, J. Inhibition of DNA virus:

Herpes-1 (HSV-1) in cellular culture replication, through an antioxidant treatment extracted

from rosemary spice. *Braz. J. Pharm. Sci.* **45**, 127–133 (2009).

32. AL-Megrin, W. A. *et al.* Potential antiviral agents of Rosmarinus officinalis extract against

herpes viruses 1 and 2. *Biosci. Rep.* **40**, BSR20200992 (2020).

33. Horváth, G. *et al.* Carnosic Acid Inhibits Herpes Simplex Virus Replication by Suppressing

Cellular ATP Synthesis. *Int. J. Mol. Sci.* **25**, 4983 (2024).

34. Ahmed, S. R., Banik, A., Anni, S. M. & Chowdhury, M. M. H. Inhibitory potential of plant-

derived metabolites against Zika virus: a computational-aided approach. *Phytomedicine Plus*

1, 100129 (2021).

35. Samy, C. R. A. *et al.* (R)-(+)-Rosmarinic Acid as an Inhibitor of Herpes and Dengue Virus

Replication: an In Silico Assessment. *Rev. Bras. Farmacogn.* **33**, 543–550 (2023).

36. Panchal, R. *et al.* Antiviral Activity of Rosmarinic Acid Against Four Serotypes of Dengue

Virus. *Curr. Microbiol.* **79**, 203 (2022).

37. Swarup, V., Ghosh, J., Ghosh, S., Saxena, A. & Basu, A. Antiviral and Anti-Inflammatory

Effects of Rosmarinic Acid in an Experimental Murine Model of Japanese Encephalitis.

Antimicrob. Agents Chemother. **51**, 3367–3370 (2007).

38. Mena, P. *et al.* Phytochemical Profiling of Flavonoids, Phenolic Acids, Terpenoids, and

Volatile Fraction of a Rosemary (Rosmarinus officinalis L.) Extract. *Molecules* **21**, 1576

(2016).

39. Borrás-Linares, I. *et al.* Rosmarinus Officinalis Leaves as a Natural Source of Bioactive

Compounds. *Int. J. Mol. Sci.* **15**, 20585–20606 (2014).

40. Kheiria, H. *et al.* Total Phenolic Content and Polyphenolic Profile of Tunisian Rosemary

(*Rosmarinus officinalis* L.) Residues. *Natural Drugs from Plants* (2021).

41. Cedeño-Pinos, C., Martínez-Tomé, M., Murcia, M. A., Jordán, M. J. & Bañón, S.

Assessment of Rosemary (Rosmarinus officinalis L.) Extract as Antioxidant in Jelly Candies

Made with Fructan Fibres and Stevia. *Antioxidants* **9**, 1289 (2020).

42. Coish, J. M. *et al.* Zika Virus Replication in a Mast Cell Model is Augmented by Dengue

Virus Antibody-Dependent Enhancement and Features a Selective Immune Mediator

Secretory Profile. *Microbiol. Spectr.* **10**, e01772-22 (2022).

43. Gaudry, A. *et al.* The Flavonoid Isoquercitrin Precludes Initiation of Zika Virus Infection in

Human Cells. *Int. J. Mol. Sci.* **19**, 1093 (2018).

44. Cirne-Santos, C. C. *et al.* In vitro antiviral activity against Zika virus from a natural product

of the Brazilian red seaweed Bryothamnion triquetrum. *Acta Virol.* **65**, 402–410 (2022).

45. Saivish, M. V. *et al.* Antiviral Activity of Quercetin Hydrate against Zika Virus. *Int. J. Mol.*

Sci. **24**, 7504 (2023).

46. Vázquez-Calvo, Á., Jiménez de Oya, N., Martín-Acebes, M. A., Garcia-Moruno, E. & Saiz,

J.-C. Antiviral Properties of the Natural Polyphenols Delphinidin and Epigallocatechin Gallate

against the Flaviviruses West Nile Virus, Zika Virus, and Dengue Virus. *Front. Microbiol.* **8**,

1314 (2017).

47. Ryan, S. J. *et al.* Warming temperatures could expose more than 1.3 billion new people to

Zika virus risk by 2050. *Glob. Change Biol.* **27**, 84–93 (2021).

48. Elebeedy, D. *et al. In vitro* and computational insights revealing the potential inhibitory

effect of Tanshinone IIA against influenza A virus. *Comput. Biol. Med.* **141**, 105149 (2022).

49. Koch, C., Reichling, J., Schneele, J. & Schnitzler, P. Inhibitory effect of essential oils against

herpes simplex virus type 2. *Phytomedicine* **15**, 71–78 (2008).

50. Vázquez-Calvo, Á., Jiménez de Oya, N., Martín-Acebes, M. A., Garcia-Moruno, E. & Saiz,

J.-C. Antiviral Properties of the Natural Polyphenols Delphinidin and Epigallocatechin Gallate

against the Flaviviruses West Nile Virus, Zika Virus, and Dengue Virus. *Front. Microbiol.* **8**,

(2017).

51. Al-jaafreh, A. M. Evaluation of Antioxidant Activities of Rosemary (Rosmarinus officinalis

L.) Essential Oil and Different Types of Solvent Extractions. *Biomed. Pharmacol. J.* **17**, 323–

339 (2024).

52. Lin, W.-Y., Yu, Y.-J. & Jinn, T.-R. Evaluation of the virucidal effects of rosmarinic acid

against enterovirus 71 infection via in vitro and in vivo study. *Virol. J.* **16**, 94 (2019).

53. Müller, J. A. *et al.* Inactivation and Environmental Stability of Zika Virus. *Emerg. Infect.*

Dis. **22**, 1685–1687 (2016).

54. Mantzourani, C., Tarantilis, P. A. & Kokotou, M. G. Carnosic Acid and Carnosol: Analytical

Methods for Their Determination in Plants, Foods and Biological Samples. *Separations* **10**,

481 (2023).

55. Chung, Y.-C. *et al.* Magnesium lithospermate B and rosmarinic acid, two compounds present

in *Salvia miltiorrhiza,* have potent antiviral activity against enterovirus 71 infections. *Eur. J.*

Pharmacol. **755**, 127–133 (2015).

56. Tsukamoto, Y. *et al.* Rosmarinic acid is a novel inhibitor for Hepatitis B virus replication

targeting viral epsilon RNA-polymerase interaction. *PLOS ONE* **13**, e0197664 (2018).

57. Hsieh, C.-F. *et al.* Rosmarinic acid exhibits broad anti-enterovirus A71 activity by inhibiting

the interaction between the five-fold axis of capsid VP1 and cognate sulfated receptors.

Emerg. Microbes Infect. **9**, 1194–1205.

58. Aruoma, O. I. *et al.* An evaluation of the antioxidant and antiviral action of extracts of

rosemary and provençal herbs. *Food Chem. Toxicol.* **34**, 449–456 (1996).

Supplemental Material

816
Supplemental Figure 1: Infectious titers peak at 96 h in HFF-1 cells infected with ZIKV. (a)
HFF-1 cells were infected with ZIKV (MOI = 1) for 1 h at 37°C. Cell-free supernatants were
collected 0, 24, 48, 72, 96, and 120 h post infection and viral titers in each sample were
quantified by plaque assay. Data are expressed as PFU/mL ± SEM for n = 3 independent
experiments, each performed in duplicate. A two-way ANOVA and Dunnett's multiple
comparisons test were used to determine differences in viral titers between each timepoint.
****p<0.0001 relative to supernatant collected 96 h post-infection. (b) Plaque assay images were
taken using an Amersham Typhoon Biomolecular Imager and represent wells from one
independent experiment in its entirety.