VegVault database: linking global paleo-, and neo-vegetation data with functional traits and abiotic drivers

Authors

Ondřej Mottl[1,2], Franka Gaiser[3], Irena Šímová[1,4], Suzette G. A. Flantua[5,6]

Affiliations

1. Center for Theoretical Study, Charles University, Jilská 1, CZ-11000 Prague, Czech Republic
2. Department of Botany, Faculty of Science, Charles University, Benátská 2, CZ-12801 Prague, Czech Republic
3. Sport Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER) & Bayreuth Center of Sport Science (BaySpo), University of Bayreuth, 95447 Bayreuth, Germany Department of Ecology, Faculty of Science, Charles University, Viničná 7, CZ-12800 Prague, Czech Republic
4. Department of Biological Sciences, University of Bergen, PO Box 7803, N-5020 Bergen, Norway
5. Bjerknes Centre for Climate Research, Bergen, Norway

corresponding author(s): Ondřej Mottl (ondrej.mottl@gmail.com)

Abstract

Understanding the dynamics and persistence of biodiversity patterns over short (contemporary) and long (thousands of years) time scales is crucial for predicting ecosystem changes under global climate and land-use changes. A key challenge is integrating currently scattered ecological data to assess complex vegetation dynamics over time. Here, we present VegVault, an interdisciplinary SQLite database that uniquely integrates paleo- and neo-ecological plot-based vegetation data on a global and millennial scale, directly linking them with functional traits, soil, and climate information. VegVault currently comprises data from *BIEN*, *sPlotOpen*, *TRY*, *Neotoma*, *CHELSA*, and *WoSIS*, providing a comprehensive and ready-to-use resource for researchers across various fields to address questions about past and contemporary biodiversity patterns and their abiotic drivers. To further support the usability of the data, VegVault is complemented by the {*vaultkeepr*} R package, enabling streamlined data access, extraction, and manipulation. This study introduces the structure,

28 content, and diverse applications of VegVault, emphasizing its role in advancing ecological research

29 to improve predictions of biodiversity responses to global climate change.

Background & Summary

Biodiversity and ecosystem dynamics are currently profoundly influenced by global climate change,

creating an urgent need for robust predictive models capable of anticipating these impacts. Such

models are crucial in guiding conservation efforts during this global crisis. However, many existing

models rely on contemporary (present) ecological data alone, which limits their ability to capture the

long-term responses of species and ecosystems to environmental change. Incorporating

paleoecological data—"ecology of the past" as opposed to neo-ecological, or "ecology of the

present"—offers a vital temporal perspective that can enhance our understanding of biodiversity

dynamics under shifting climate and land-use[1–3].

In recent years, an increasing number of large-scale ecological datasets have become available,

providing unprecedented opportunities to investigate ecosystem dynamics across spatial and

temporal scales. These datasets encompass contemporary vegetation data (e.g., *Global Biodiversity*

Information Facility[4] ('*GBIF*' hereafter), *sPlot*[5], *Botanical Information and Ecology Network*[6] ('*BIEN*'

hereafter), paleo-vegetation data (e.g., *Neotoma Paleoecology Database*[7] ('*Neotoma*' hereafter)),

plant functional traits (e.g., *TRY plant trait database*[8] ('*TRY*' hereafter)), and abiotic drivers (e.g.,

CHELSA[9], *WorldClim*[10]), which are all critical for understanding the complex interactions that shape

ecosystems. Yet, despite the growing availability of such datasets, efforts to integrate these diverse

sources into a unified, open-access framework remain limited. An integrated database, built with

reproducible workflows, is a crucial step toward improving predictive models that estimate

biodiversity dynamics under changing climatic conditions. Addressing this gap requires

interdisciplinary solutions[11,12] and a commitment to compiling, processing, and linking data in a

systematic and transparent manner. Currently, no comprehensive framework exists to link global

vegetation data across temporal scales with functional traits and abiotic drivers.

There have been previous attempts to integrate vegetation data across extensive time scales and

link them with other data types, such as functional traits and abiotic data. For example,

BioDeepTime[13] compiles both contemporary and fossil biodiversity time series. However, it lacks

coverage of contemporary vegetation plots and, more importantly, omits direct links to functional

traits and abiotic data. GIFT[14] includes vegetation biodiversity linked with functional traits but only

holds taxonomic lists for specific regions, without data on relative abundances. Additionally, the

latter dataset lacks the long-term dimension of vegetation dynamics over time, derived from the

currently existing large databases containing paleoecological records, such as the Neotoma

database.

Our database—**VegVault**—addresses these limitations by offering a unique resource that compiles

extensive paleo- and neo-ecological vegetation data into a single, accessible SQLite database,

accompanied by functional traits and abiotic information (i.e., climate and soil). Its design bridges

paleo and contemporary data, enabling the study of long-term processes, and linking them to

today's patterns. Thereby, VegVault offers the possibility to specifically analyse the mechanisms of

ecosystem responses to environmental change which enhances our ability to study and ultimately

predict future biodiversity trajectories. Specifically, VegVault advances analyses across spatial,

temporal, and taxonomic scales in several ways:

• **Contemporary vegetation data**: VegVault incorporates plot-based data from *BIEN*[6] and

sPlotOpen[15] (open-access version of sPlot), including essential "true negative" information

about species' absence. These datasets are pre-cleaned for taxonomic and georeferencing

errors, ensuring high reliability for subsequent analyses. The focus on plot-based data rather

than occurrence data (e.g., *GBIF*) provides a closer match to paleoecological data formats,

facilitating robust comparisons across temporal scales.

• **Paleo-vegetation data**: VegVault employs standardised fossil pollen data processed using the

FOSSILPOL workflow[16] to integrate records globally from Neotoma. This workflow ensures

quality control and creates "ready-for-analysis" data through a reproducible and detailed

process for handling large fossil pollen dataset compilations (see **Methods** for details).

• **Interdisciplinary integration**: VegVault combines past and present vegetation data with

functional traits and abiotic information to support comprehensive ecological analyses. The

abiotic information includes soil data from the *World Soil Information Service*[17] ('*WoSIS*'

hereafter), and global temperature and precipitation datasets from *CHELSA* and *CHELSA-*

TRACE21[18,19], providing essential environmental context. Functional trait data from the *TRY*

database and *BIEN* enrich the database by linking species characteristics to ecological roles

and responses. This allows researchers to move beyond taxonomic diversity and explore

functional dimensions, offering deeper insights into ecosystem processes.

• **Taxonomic classification**: VegVault ensures consistent and standardised taxonomic
information by aligning all taxa present in vegetation data—both paleo and contemporary—with
the GBIF backbone taxonomy[20]. This process, implemented via the {*taxospace*} R package[21],
automates taxonomic classification by matching names to an existing taxonomy based on
similarity (e.g., "Ulmus Type" will be classified as genus *Ulmus*). While original taxonomic
names are preserved within the database, users can access data at varying taxonomic
resolutions (e.g., species, genus, family; if available) to suit specific analytical needs. The same
standardisation approach is applied to vegetation data and trait data, facilitating seamless
integration (e.g., enabling to calculation of genus-level average trait values, when species-level
identification is unavailable).

• **Accessibility**: VegVault is designed with user accessibility in mind. The database is supported
by the {*vaultkeepr*} R package[22], which simplifies the process of data extraction from the
database for project-specific needs. It ensures that researchers can efficiently retrieve and
subsequently manipulate data in a reproducible and transparent manner.

The comprehensive nature of VegVault enables diverse applications across scientific disciplines. Its
integration of fossil pollen data and functional traits allows researchers to investigate long-term
vegetation dynamics and their responses to past climate variations. Such integration provides
critical context for past ecological trends[e.g.,23,24], including trait information, which is currently
underrepresented in paleoecological analyses of vegetation dynamics[25]. While there are now
studies showing the relationship between functional traits and climate for global contemporary
data[26], VegVault allows testing the universality of such relationships across temporal scales. In
addition, VegVault provides the needed data for advanced modelling approaches, such as
comparing species distributions across temporal scales[e.g.,27], and analysing biodiversity patterns
using Joint Species Distribution Models[28] ('JSDM' hereafter). The database also facilitates studies of
functional ecology and biogeography by enabling assessments of shifts in ecosystem functioning
and climatic drivers over time, and by identifying the taxonomic and functional novelty of
communities undergoing climate-driven changes[e.g.,29,30]. Furthermore, the detailed climate data
(past and present), linked with vegetation records, provide a robust foundation for examining the
impacts of past and present climate change on ecosystems[e.g.,31]. These capabilities make VegVault
a valuable interdisciplinary resource for predicting biodiversity patterns and informing conservation
strategies and policy decisions aimed at mitigating climate change effects. While this is not an

exhaustive list of potential applications, VegVault offers a rich and accessible resource that can
significantly advance ecological research and global biodiversity conservation.

Methods

VegVault workflow and reproducibility

VegVault integrates data from multiple well-established sources to provide a comprehensive view
of vegetation dynamics. The current sources of data are summarised in **Table 1**.

The data processing and integration into VegVault follow a meticulous workflow to ensure
consistency and reliability. Each data source is first accessed and processed individually using a
specific pipeline. See **Figure 1** for the workflow process of accessing and processing all data sources.

In order to ensure the traceability of data, VegVault is designed as a versioned project. Each version
(including the current v1.0.0) is created using the git Tags system[32]. Each Tag is a snapshot of code
at certain points in time. Each of the individual data pipelines, coded in R programming language, is
publicly available and stored in a separate GitHub repository with a specific git Tag and published
with DOI (see legend of **Figure 1** for individual DOIs).

Finally, all processing pipelines with their corresponding Tags are then migrated into an SQLite
database using another dedicated GitHub repository, ensuring its reproducibility and transparency
of the development of the database.

In addition to the consolidation of all processed data into a unified SQLite database, the final
VegVault 'migration' repository performs three additional procedures: (i) taxon classification, (ii)
grouping of traits into *Trait Domains*, and (iii) creation of *gridpoints* for abiotic data.

(i) Taxon classification

As VegVault consists of data on taxa from various sources, the {*taxospace*} R package[21] is used to
classify the diverse taxa into a unifying backbone in the VegVault database. The {*taxospace*} tool
automatically aligns taxon names with the taxonomical backbone of the *GBIF*[20]. Specifically, we find
the best match of the raw names of taxa using *Global Names Resolver*[33], which is then aligned with
GBIF. Taxonomic information, detailed up to the family level, is stored for each taxon, ensuring
consistency and facilitating comparative analyses across different datasets. However, note that

taxonomic information down to the species level is not available for each taxon (e.g., some fossil
pollen types can only be identified to the genus or family level).

(ii) Grouping of *Traits* into *Trait Domains*

As there are differences in trait names across sources of data (e.g., "*Leaf nitrogen (N) content per*
leaf dry mass" and "*leaf nitrogen content per leaf dry mass*"), we added a new variable *Trait Domain*
that groups traits together following the trait selection of Diaz et al. (2016)[34]. For example, the trait
"*Plant height vegetative*" from *TRY* and the trait "*whole plant height*" from *BIEN* are both grouped
under the "*Plant height*" *Trait Domain*. See **Table 2** for the full list of traits and their *Trait Domain*.
Such grouping is aimed to serve as an efficient extraction of traits across various data sources.

(iii) Creation of *gridpoints* of abiotic data

The abiotic data in the VegVault database provide essential information on environmental factors
affecting vegetation distribution and traits, namely climate (past and present) and soil variables
(**Table 3**).

We developed a data structure that provides readily available environmental context for each
vegetation (and trait) record by creating spatio-temporal links between these records and abiotic
information. We first unified the resolution of the original rasters by resampling all abiotic data into
~ 25km resolution at 500-year time slices, reducing the amount of data. Next, as the rasters are not
suitable to be stored in an SQLite database, we created artificial points, called '*gridpoints*'**,** located
in the middle of each raster cell. This resulted in a uniform spatio-temporal matrix of gridpoints
holding the abiotic information (see **Figure 2B** or an example area). Next, we linked *gridpoints* and
other non-*gridpoint Samples*, namely *vegetation_plot*, *fossil_pollen_archive*, and *traits* (see **Internal**
Database Structure section), and calculated the spatial and temporal distances between them. We
discarded any *gridpoint Sample* beyond 50 km and/or 5000 years from any other non-*gridpoint*
Sample. See **Figure 2C** for an example of linking abiotic information at *gridpoints* and vegetation data.
In VegVault, users can select the information for each non-*gridpoint Sample* from the closest spatio-
temporal abiotic data or get the average from all surrounding *gridpoints*. By providing such a
comprehensive and well-linked structure between vegetation and abiotic data, VegVault enhances
the ability to study the interactions between vegetation and their environment, facilitating the
workflow for advanced ecological research and modelling efforts.

Data Records

The VegVault database v1.0.0 is an SQLite database of a size of ~ 110 GB stored in the *National Repository*[35] ('*REPO*'). It contains > 480,000 *Datasets*, > 13,000,000 *Samples*, > 100,000 *Taxa*, 6 vegetation *Traits*, > 11,000,000 *Trait* values, and 8 abiotic variables. The spatio-temporal overview of all vegetation data can be found in **Figure 3**.

Currently, the SQLite database consists of 29 interconnected tables with 87 fields (variables). However, the internal structure of VegVault may not be of relevance to a user, as the {*vaultkeepr*} R package processes all the data to output only the most relevant information in a "ready-to-analyse" format (see **Usage Notes** section).

Internal Database Structure

VegVault is organized into several sections to systematically manage the varying datasets it integrates. The main structure is the *Dataset*, which serves as the cornerstone of the database structure. *Datasets* are composed of *Samples*, representing individual data points within each dataset. Each *Dataset* will only contain several *Samples* if these differ in age. There are four types of *Datasets*, namely (i) contemporary vegetation plots, (ii) past vegetation (fossil pollen records), (iii) functional traits, and (iv) gridpoint (a dataset type created to hold abiotic data, see details in the **Methods** section).

For the contemporary (i) and past (ii) vegetation *Datasets*, the *Samples* hold information about *Taxa* as derived directly from the primary data sources. Trait information is organised in separate *Datasets* (as they are associated with unique information about their spatio-temporal location, references, etc) but linked to the same *Taxa* codes as those taxa in the vegetation *Datasets* (if present). Moreover, each *Taxa*, disregarding the source, is associated with classification information (i.e. species, genus and family name; see **Methods** section). While the vegetation data is the main focus of the database, the structure of separate *Datasets* ensures a comprehensive organization of the interlinked data. Note that there is more in-depth and updated documentation of database structure on the project website (*https://bit.ly/VegVault*).

Dataset

Here we will explain a selected number of features of the *Dataset*.

Dataset Type

The *Dataset Type* defines the most basic classification of each *Dataset*, ensuring that the vast amount of data is categorized systematically. Currently, VegVault contains the following types of *Datasets*:

- `vegetation_plot`: This type includes contemporary vegetation plot data, capturing contemporary vegetation characteristics and distributions.
- `fossil_pollen_archive`: This type encompasses past vegetation plot data derived from fossil pollen records, providing insights into past vegetation and climate dynamics.
- `traits`: This type contains functional trait data, detailing specific characteristics of plant species that influence their ecological roles.
- `gridpoints`: This type holds artificially created *Datasets* to manage abiotic data, here climate and soil information.

See **Figure 4A** for an overview of the number of *Datasets* in each *Dataset Type*.

Dataset Source-Type

Each *Dataset* is derived from a specific *Source-Type* including:

- **BIEN**[6] (Botanical Information and Ecology Network).
- **sPlotOpen**[15] (The open-access version of sPlot).
- **TRY**[8] (TRY Plant Trait Database).
- **Neotoma-FOSSILPOL** (FOSSILPOL[16]: Workflow for processing and standardizing global paleoecological pollen data from *Neotoma*[7]) Note that we specifically state *Neotoma-FOSSILPOL* and not just Neotoma, as *FOSSILPOL* not only provides the data acquisition but also alters it (e.g., creating new age-depth models). It also addresses major challenges in paleoecological data integration, such as age uncertainty, by incorporating probabilistic age-depth models and their associated uncertainty matrices. This enables the propagation of temporal uncertainty in subsequent analyses, a critical advancement for robust macroecological studies, previously flagged as a major issue with paleo-data[36].
- **Gridpoints** (artificially created *Datasets* for abiotic data).

See **Figure 4B** for an overview of the number of *Datasets* for each *Dataset Source-Type*. VegVault maintains detailed information on the source used to retrieve the original data, thereby enhancing the findability and referencing of primary data sources.

Sampling Method

Information on the *Sampling Method* is only available for both *vegetation_plot* and *fossil_pollen_archive Datasets*, and varies substantially reflecting the diverse nature of the data collected. Such information is crucial for understanding the context and limitations of each *Dataset Type*. For contemporary vegetation plots, sampling involves standardized plot inventories and surveys that capture detailed vegetation characteristics across various regions. Fossil pollen data are collected from sediment records from numerous different depositional environments[16] representing past vegetation and climate dynamics.

Dataset References

To support robust and transparent scientific research, each *Dataset* in VegVault can have multiple *References* at different levels as the raw data was pooled from other databases that provide data from various authors. The *Dataset Source-Type*, *Dataset Source*, and *Sampling Method* can all have their own *References*, providing detailed provenance and citation information. This multi-level referencing system enhances the traceability and validation of the data. The VegVault database v1.0.0 currently includes 706 sources of *Datasets* where each dataset can also have one or more direct references to specific data, further ensuring that users can accurately cite and verify the sources of their data.

Samples

Samples represent the main unit of data in the VegVault database, serving as the fundamental building blocks for all analyses. There are currently over 13 million *Samples* in the VegVault database v1.0.0 (of which ~ 1.6 million are *gridpoints* of abiotic data, see **Methods** section). Each *Sample* is linked to a specific *Dataset* via the *Dataset-Sample* table, which ensures that every sample is correctly associated with its corresponding *Dataset Type* (whether it is *vegetation_plot*, *fossil_pollen_archive*, *traits*, or *gridpoint*) and other *Dataset* properties (e.g., geographic location).

Vegetation plot size

The size of vegetation plots can vary substantially. This detail is crucial for ecological studies where plot size can influence species diversity and abundance metrics, thus impacting follow-up analyses and interpretations. To account for this variability, information about the plot size is recorded separately for each contemporary *Sample*. See **Figure 5** for an overview of the number of *Samples* in each size category.

 Sample age

VegVault encompasses both contemporary and paleo-data, necessitating accurate age information

for each *Sample*. Contemporary *Samples* are assigned an age of 0, while *Samples* from fossil pollen

records are in calibrated years before the present (cal yr BP). The "present" is here specified as 1950

AD. In addition, each *Sample* from the *fossil_pollen_archive Dataset* is also associated with an

uncertainty matrix. This matrix provides a range of potential ages derived from age-depth modelling,

reflecting the inherent uncertainty in dating paleoecological records. See an example of detailed age

uncertainty information for one fossil pollen record in **Figure 6**.

Sample reference

Each *Sample* in VegVault can have specific *References* in addition to those at the *Dataset*-level.

These individual *Sample References* provide detailed provenance and citation information, ensuring

that users can trace the origin and validation of each data point. This level of referencing enhances

the transparency and reliability of the data, especially when the database continues to be updated

in the future.

Taxa

The VegVault database records taxa names derived directly from the primary data sources. Each

individual *Taxon* is linked to corresponding *Samples* through the *SampleTaxa* table, ensuring

accurate and systematic association between species and their ecological data. Currently, the

VegVault database v1.0.0 holds over 100 thousand taxonomic names. Each taxonomic name

undergoes an automated classification (see **Methods** section). Through this procedure, only 1312

(1.2%) of all *Taxa* remained unclassified. See **Figure 7** for an overview of the number of species,

genus and family classification.

Traits

Functional traits of vegetation taxa follow the same structure of the *Dataset* and *Samples* obtained

directly from the *Dataset Source-Types*. Storing trait values in VegVault involves linking information

across *Datasets*, *Samples*, *Taxa*, and *Traits*. This comprehensive linkage ensures that each *Trait*

value is accurately associated with its relevant ecological, environmental and taxonomic context.

Yet, it is up to the user to decide how to further aggregate trait values if multiple trait samples of one

Trait Domain are available for the same environmental or taxonomic entity. To ensure clarity and

reproducibility, each *Trait* in the VegVault database can have additional *References* beyond the

general *Dataset* and *Sample References*. These *Trait*-specific *References* provide detailed

provenance and citation information, supporting rigorous scientific research and enabling users to

trace the origins and validation of each trait value.

Traits are further grouped into *Trait Domains* to allow for easier aggregation of *Traits* across data

sources (see **Methods** section). In total, six Trait domains are present: *Stem specific density*, *Leaf*

nitrogen content per unit mass, *Diaspore mass*, *Plant height*, *Leaf area*, *Leaf mass per area*, following

Diaz et al. (2016)[34]. See **Table 2** for an overview of *Traits* in various *Trait Domains* and the number of

Trait values.

Abiotic data

Currently, VegVault includes abiotic data from *CHELSA*[9], *CHELSA-TRACE21*[18,19], and *WoSIS*[17]. The

first two data sources provide high-resolution downscaled climatic data, while the latter offers

detailed soil information (only available for contemporary data). The abiotic variables included in the

database are summarized in **Table 3**. We opt for including the soil data, while only contemporary, as

they have been previously confirmed as important predictors for vegetation distribution[37–39]. Values

of abiotic data are stored in *gridpoints*, artificially created *Datasets* and *Samples* (see **Methods**),

with one *Dataset* holding more *Samples* only if they differ in age.

Technical Validation

The data validity of VegVault is driven by the completeness, accuracy, and reliability of the source

data. The licence of the primary data sources prohibits the changing of subsets of the data (e.g., *BIEN*

data has a *CC BY-NC-ND* licence). Nevertheless, several steps have been undertaken to ensure that

VegVault provides high-quality data for research applications.

For the fossil pollen records, strict data-quality rules have been applied by using the *FOSSILPOL*

workflow[16]. This includes the selection of depositional environments, ecological groups, chronology

control point types, and a minimum number of chronology control points to construct age-depth

models. Individual samples and records have been filtered by age limits, number of pollen grains,

maximum age interpolation, and number of valid levels. In addition, the accuracy of fossil pollen

data is increased by also re-estimation of all age-depth models using the Bayesian probabilistic

approach[40] and including the information about individual age uncertainty. For more information

about *FOSSILPOL* workflow, see https://bit.ly/FOSSILPOL. For exact values used for accessing,

processing, and filtration of fossil pollen data for VegVault v1.0.0, see

https://github.com/OndrejMottl/VegVault-FOSSILPOL/tree/v1.0.0.

Usage Notes

Guidelines for data reuse and update

Due to the version system of VegVault, the database is expected to be updated using new data

and/or changing the process of data migration and processing. Therefore, we recommend citing the

latest VegVault version with the corresponding DOI (v1.0.0 can be found here:

doi.org/10.48700/datst.m7ky4-4de51).

As the primary data sources are held under various licences, VegVault has a *CC BY-NC-ND 4.0*

Licence (https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en), allowing the usage of more

restricted data (i.e. *BIEN* data is under *CC BY-NC-ND*). We recommend users to carefully review the

restrictions associated with data usage. In addition, when citing the usage of VegVault, users should

always cite the primary sources of the data, as well as any individual datasets. For those purposes,

we created a function `*vaultkeepr::get_references()*` within the {*vaultkeepr*} R package (see below)

that will output a list of recommended references for the specific data compilation created by the

user.

Limitations and considerations for future research

Potential limitation lies in the completeness, accuracy, and reliability of the primary data sources.

For example, it has been suggested that improper data practices erode the quality of global

ecological databases such as *TRY*[41], which have been used in VegVault. We do not attempt to solve

these problems as VegVault's main purpose is the re-usability of the data. Note that for some cases

we are not permitted to make modifications based on the original data (e.g. *BIEN* data has a *CC BY-*

NC-ND licence).

{vaultkeepr}: R-package for accessing the VegVault database

In order to make usage of the VegVault database as easy as possible, we have developed an R-

package called {*vaultkeepr*}, providing a suite of functions to effectively interact with the database

directly from the R programming environment[42]. Functions include opening the database, extracting

datasets, filtering samples, and accessing specific taxa and traits. This package is a well-tested

interface   (>95%   code   coverage).   For   more   details,   see   {*vaultkeepr*}   website

(https://bit.ly/vaultkeepr), and see **Figure 8** for details about individual steps to retrieve the data from

VegVault.

Here we present 3 examples of potential projects and how to obtain data for such projects from

VegVault using {*vaultkeepr*} R-package. Note that we are specifically not trying to do any analysis,

only presenting a way to obtain data, which can be used for such follow-up projects.

1) Spatio-temporal patterns of the *Picea* genus across North America since the Last Glacial Maximum

359   The first example demonstrates how to retrieve data for the genus *Picea* across North America by

360   selecting both contemporary and fossil pollen plot datasets, filtering samples by geographic

361   boundaries and temporal range (0 to 15,000 yr BP), and harmonizing taxa to the genus level. The

362   resulting dataset allows users to study spatiotemporal patterns of *Picea* distribution over millennia

363   (**Figure 9A**). This can be accomplished by running the following code:

```r
data_na_plots_picea <-
  # Access VegVault
  vaultkeepr::open_vault(path = "<path_to_VegVault>") %>%
  # Start by adding dataset information
  vaultkeepr::get_datasets() %>%
  # Select both contemporary and paleo plot data
  vaultkeepr::select_dataset_by_type(
    sel_dataset_type = c(
      "vegetation_plot",
      "fossil_pollen_archive"
    )
  ) %>%
  # Limit data to North America
  vaultkeepr::select_dataset_by_geo(
    lat_lim = c(22, 60),
    long_lim = c(-135, -60)
  ) %>%
  # Add samples
  vaultkeepr::get_samples() %>%
  # Limit the samples by age
  vaultkeepr::select_samples_by_age(
    age_lim = c(0, 15e3)
  ) %>%
  # Add taxa & classify all data to a genus level
  vaultkeepr::get_taxa( classify_to = "genus") %>%
  # Extract only Picea data
  vaultkeepr::select_taxa_by_name(sel_taxa = "Picea") %>%
  vaultkeepr::extract_data()
```

393 ```

2) Joined-species Distribution Model for vegetation within the Czech Republic

395 In the second example, the project aims to do species distribution modelling for plant taxa in the

396 Czech Republic based on contemporary vegetation plot data and mean annual temperature. The

397 code includes selecting datasets and extracting relevant abiotic data (**Figure 9B**) as follows:

```r
data_cz_contemporary <-
  # Access VegVault file
  vaultkeepr::open_vault(path = "<path_to_VegVault>") %>%
  # Add the dataset information
  vaultkeepr::get_datasets() %>%
  # Select contemporary plot data and climate
  vaultkeepr::select_dataset_by_type(
    sel_dataset_type = c(
      "vegetation_plot",
      "gridpoints"
    )
  ) %>%
  # Limit data to Czech Republic
  vaultkeepr::select_dataset_by_geo(
    lat_lim = c(48.5, 51.1),
    long_lim = c(12, 18.9)
  ) %>%
  # Add samples
  vaultkeepr::get_samples() %>%
  # select only contemporary data
  vaultkeepr::select_samples_by_age(
    age_lim = c(0, 0)
  ) %>%
  # Add abiotic data
  vaultkeepr::get_abiotic_data() %>%
  # Select only Mean Annual Temperature (bio1)
  vaultkeepr::select_abiotic_var_by_name(sel_var_name = "bio1") %>%
  # add taxa
  vaultkeepr::get_taxa() %>%
  vaultkeepr::extract_data()
```

3) Patterns of community weighted mean of plant height for South and Central America between 6-12 ka cal yr BP

432 The third example focuses on obtaining data to be able to reconstruct plant height for South and

433 Central America between 6-12 ka cal yr BP (thousand calibrated years before present). This example

434 project showcases the integration of trait data with paleo-vegetation records to subsequently study

435 historical vegetation dynamics and functional composition of plant communities (**Figure 9C**):

```r
data_la_traits <-
```

14

```
438      # Access VegVault file
439      vaultkeepr::open_vault(path = "<path_to_VegVault>") %>%
440      # Add the dataset information
441      vaultkeepr::get_datasets() %>%
442      # Select contemporary plot data and climate
443      vaultkeepr::select_dataset_by_type(
444        sel_dataset_type = c(
445          "fossil_pollen_archive",
446          "traits"
447        )
448      ) %>%
449      # Limit data to South and Central America
450      vaultkeepr::select_dataset_by_geo(
451        lat_lim = c(-53, 28),
452        long_lim = c(-110, -38),
453        sel_dataset_type = c(
454          "fossil_pollen_archive",
455          "traits"
456        )
457      ) %>%
458      # Add samples
459      vaultkeepr::get_samples() %>%
460      # Limit to 6-12 ka yr BP
461      vaultkeepr::select_samples_by_age(
462        age_lim = c(6e3, 12e3)
463      ) %>%
464      # add taxa & classify all data to a genus level
465      vaultkeepr::get_taxa(classify_to = "genus") %>%
466      # add trait information & classify all data to a genus level
467      vaultkeepr::get_traits(classify_to = "genus") %>%
468      # Only select the plant height
469      vaultkeepr::select_traits_by_domain_name(sel_domain = "Plant height") %>%
470      vaultkeepr::extract_data()
471    ```
472
```

Code Availability

The VegVault database is built on SQLite, chosen for its robustness and compatibility with various data types. The database itself is under *CC BY-NC-ND 4.0* licence, due to the licences of underlying data (see **Usage Notes** section).

VegVault utilizes several tools and software to facilitate data management and access. All code used to generate the VegVault database is publicly available under MIT licence (see **Methods** section and **Figure 1** for details about individual GitHub repositories).

To enable easy access and manipulation of data, {*vaultkeepr*} R package has been developed as an open access software under MIT licence.

The code to create the figures in this manuscript is available under DOI:10.5281/zenodo.15201994.

Acknowledgements

OM is funded by the *Czech Science Foundation PIF* grant (GN23-06386I), by the *Charles University Research Centre program* (UNCE/24/SCI/006), and by the *Institutional Support for Science and Research* of the Ministry of Education, Youth and Sports of the Czech Republic. SGAF is funded by the *Trond Mohn Research Foundation* (TMF) and the University of Bergen by starting grant no. TMS2022STG03).

Author contributions

OM conceived the study and programmed all related code; SGAF, FG, and IS performed critical comments on the technical infrastructure of the database; OM & SGAF wrote the first draft of the manuscript; SGAF, FG, and IS provided several rounds of revisions. All authors approved of the final version of the manuscript.

Competing interests

The author(s) declare no competing interests.

Figures

Figure 1. Schematic workflow of accessing and processing all primary data sources into VegVault v1.0.0 database. 1) *Neotoma Paleoecological Database*[7]; **2)** *sPlotOpen*[15]; **3)** *Botanical Information and Ecology Network*[6]; **4)** *TRY Plant Trait Database*[8]; 5) *CHELSA*[9] and *CHELSA-TRACE21*[18,19]; **6)** *World Soil Information Service*[17]; **7)** *FOSSILPOL: Workflow for processing and standardizing global paleoecological pollen data*. A project-specific version with all the settings used for VegVault is stored as GitHub repository *VegVault-FOSSILPOL* (can be accessed as DOI: 10.5281/zenodo.14745060); **8)** The GitHub repository *VegVault-Vegetation_data* used to processes data for contemporary vegetation plots (can be accessed as DOI: 10.5281/zenodo.14745273); **9)** A GitHub repository *VegVault-Trait_data* used to processes data for plant vegetation traits (can be accessed as DOI: 10.5281/zenodo.14745086); **10)** The GitHub repository *VegVault-abiotic_data* used to process climate and soil abiotic data (can be accessed as DOI: 10.5281/zenodo.14745293); **11)** The GitHub repository *VegVault* used to integrate all data together and migrate them into SQLite database (can be accessed as DOI: 10.5281/zenodo.15201994). Additional steps such as the creation of *gridpoints*, grouping of *Traits* into *Trait Domains,* and classification of taxa are also present (for more details see **Methods**); **12)** VegVault - SQLite database (v1.0.0; DOI: 10.48700/datst.m7ky4-4de51); For details about primary data sources, see **Table 1**.

Figure 2. **Example of the resolution of the *gridpoint Datasets* extracting abiotic information.**
A: Map of Europe. The red line indicates country borders with the Czech Republic as the focus of the
map. **B**: Distribution of the individual *gridpoints* (grey squares) within the Czech Republic (red).
C: Linking vegetation plots and *gridpoints*. The green circles represent the contemporary vegetation
plots. The grey lines connect the vegetation points to their closest *gridpoint*. Note that the abiotic
data is provided to each vegetation plot from their closest *gridpoint* (i.e., the vegetation plots
connected to the same gridpoint will have the same abiotic data). *Gridpoints* without vegetation
plots in their spatio-temporal vicinity (here contemporary data) were excluded from the VegVault
database to reduce the database size. See **Methods** section for more details.

Figure 3. **Spatio-temporal distribution of vegetation Samples** (for both vegetation plots and fossil pollen archives). Note that this is an overview of the current state of VegVault (v1.0.0.). **A**: Spatial distribution with each point representing one plot. **B**: Temporal distribution of individual Samples.

Figure 4. **Visualisation of the number of *Datasets*.** **A**: Waffle plot coloured by *Dataset Type* with each square representing ~1000 *Datasets*. **B**: Number of *Datasets* by *Dataset Source-Type* grouped by *Dataset Type*. See **Internal Database Structure** section for more details.

Figure 5. **Distribution of *Samples* by vegetation plot sizes.** In VegVault, only *vegetation_plot*
Samples and *Datasets* have such information. See **Internal Database Structure** section for more
details.

Figure 6: **Example of age uncertainty of *Samples* for selected fossil pollen *Dataset* (ID: 91256).** Each point represents one *Sample*. Each horizontal line represents a range of all potential ages associated with that *Sample*. Potential ages are gathered from age-depth models and stored in the *SampleUncertainty* table. See **Internal Database Structure** section for more details about the data structure and *FOSSILPOL* workflow[16] for more details about age uncertainty from age-depth models.

551 **Figure 7**. **Total number of _Taxa_ and their classification present in VegVault**. See **Methods** section

552 for more information about the classification process.

Figure 8. **Schematic workflow of accessing and extracting data from VegVault database using {vaultkeepr} R package**. **1)** Download VegVault database (v1.0.0; DOI: 10.48700/datst.m7ky4-4de51); **2)** Download {vaultkeepr} R package from GitHub (current version 0.0.6; DOI: 10.5281/zenodo.14964737); **3)** In the R programming environment[42], the user has to select, which type(s) of datasets they would like to extracted. A minimum of one *Dataset Type (see* **Data Records** section for more details*)* must be specified when using the `vaultkeepr::select_dataset_by_type()` function; **4)** When the user is interested in data from a specific region, they can specific the geographical coordinates of a rectangular extent using the `vaultkeepr::select_dataset_by_geo()` function to only get *Samples* from the individual *Datasets* that were recorded within the region of interest. Additionally, the user can specify the temporal focus by filtering data within a specific period using the `vaultkeepr::select_samples_by_age()`; **5)** The user may also specify any further

attributes to be added to the data compilation. Specifically: `*get_taxa()*`—When extracting
contemporary or past vegetation (fossil pollen) records, the user most likely wishes to add
information about the abundance of individual *Taxa* in each *Sample*. To do so, the user can use the
`*vaultkeepr::get_taxa()*` function. In addition, the user can standardise the taxonomy, so that the
extracted *Taxa* can be compared. The parameter `*classify_to*` within the `*vaultkeepr::get_taxa()*`
function allows the user to specify a taxonomic level (species, genus or family) on which the data
should be comparable (see **Methods** about *Taxa* classification). Furthermore, the user can select
specific *Taxa* based on taxonomy by using the `*vaultkeepr::select_taxa_by_name()*` function.
`*get_traits()*`—When wishing to link the *Trait* data with the vegetation records, the user can use
`*vaultkeepr::get_traits()*` function to extract all *Trait Samples* of the earlier specified spatio-
temporal extent. Moreover, similarly to the `*vaultkeepr::get_taxa()*` function, the user can specify
the taxonomic level to which the data should be standardised by using the `*classify_to*` parameter.
Please note that the user has to decide in their further analysis how they want to aggregate the
measured traits per taxonomic group, e.g. by taking the mean. Further, the user can select a specific
Trait Domain (of the six available; see **Methods** about details) by using the
`*vaultkeepr::select_traits_by_domain_name()*` function. `*get_abiotic_data()*`—When wishing to
link vegetation or trait records with abiotic data, the abiotic data can be obtained by the
`*vaultkeepr::get_abiotic_data()*` function. The user can specify in which mode the abiotic data
should be linked, which can be either "*nearest*" (i.e. geographically closest records), "*mean*" or
"*median*" (summarising all abiotic records within a set geographical and/or temporal distance). This
can be further tweaked by its parameters `*limit_by_distance_km*` and `*limit_by_age_years*`.
Furthermore, specific abiotic variables can be chosen by the
`*vaultkeepr::select_abiotic_var_by_name()*` function; **6)** When defined all specifications for data
extraction, the user can execute the extraction using `*vaultkeepr::extract_data()*`. This will result in
a "ready-for-analyses" data compilation. Moreover, the user can use the
`*vaultkeepr::get_references()*` function to obtain all references required (and/or suggested) for such
compilation (see **Internal Database Structure** section for more details about hierarchical structure
of references). Finally, see the **Usage Notes** section for examples of data processing and extracting
using {vaultkeepr} R package.

Figure 9. **Visualisation of example data usage of VegVault using {*vaultkeepr*} package**. See **Usage Notes** section for code used to extract such data from VegVault and for details about the aim of each data compilation. **A**: Data for "*Spatio-temporal patterns of the Picea genus across North America since the Last Glacial Maximum*" example**.** The individual points represent vegetation data (*Samples* for fossil pollen or contemporary vegetation plots) within each age time slice. **B**: Data for "*Joined-species Distribution Model for vegetation within the Czech* Republic" example. Each point represents one contemporary vegetation plot. The colour represents the Mean Annual Temperature (MAT) associated with that plot. Note that several vegetation plots can have the same value due to sharing the same closest *gridpoint* (see **Figure 2**). **C**: Data for "*Patterns of community weighted mean of plant height for South and Central America between 6-12 ka cal* yr BP". The left figure represents the temporal length of fossil pollen *Datasets*, which are available for Central and South America between 6 and 12 ka BP (thousand years before present). The middle figure represents the distribution of various genera present in this data assembly. The right figure represents the distribution of average *Plant height Trait* per each genus present in the data assembly

Tables

Table 1: Main sources of data presented in VegVault v1.0.0.

Source	Type	Comment	References
Botanical Information and Ecology Network (*BIEN*)	Vegetation – contemporary data	Plot-based vegetation data	Botanical Information and Ecology Network (http://bien.nceas.ucsb.edu/bien/)[6]
sPlotOpen	Vegetation - contemporary data	The open-access version of *sPlot*[5]	Sabatini et al. (2021)[15]
FOSSILPOL: Workflow for processing and standardizing global paleoecological pollen data	Vegetation - Fossil pollen records	We specifically state FOSSILPOL and not Neotoma Paleoceologcal database[7], which is the primary source of the data, as FOSSILPOL not only accesses the data but also applies several selection criteria and adds new information (e.g., new age-depth models)	Flantua et al. (2023)[16]
TRY Plant Trait Database (*TRY*)	Functional traits of vegetation		Kattge et al. (2020)[8]
CHELSA	Abiotic data – climate	global downscaled climate data at 1km resolution	Karger et al. (2017)[9]
CHELSA-Trace21k	Abiotic data – past climate	global downscaled climate data at 1km resolution since the Last Glacial Maximum (21,000 yr BP)	Karger et al. (2020)[18] Karger et al. (2023)[19]
World Soil Information Service (*WoSIS*)	Abiotic data – soil types	global downscaled climate data at 1km resolution	Batjes et al. (2024)[17]

Table 2: Selected *Traits* presented in the VegVault v1.0.0. The *Traits* were selected based on similarity to Diaz et al. (2016)[34] and grouped into six main *Trait Domains* to simplify the navigation between different data sources.

Trait Domain	*Trait* name	*Trait* units	Source of *Datasets*	Number of *Datasets*
Stem specific density	*Stem specific density (SSD, stem dry mass per stem*	g/cm^3	*TRY*	8058

	fresh volume) or wood density			
Stem specific density	stem wood density	g/cm^3	BIEN	348
Leaf nitrogen content per unit mass	Leaf nitrogen (N) content per leaf dry mass	mg/g	TRY	11464
Leaf nitrogen content per unit mass	leaf nitrogen content per leaf dry mass	mg/g	BIEN	707
Diaspore mass	Seed dry mass	mg	TRY	1635
Diaspore mass	seed mass	mg	BIEN	48
Plant height	Plant height vegetative	m	TRY	11476
Plant height	whole plant height	m	BIEN	193492
Leaf area	Leaf area (in case of compound leaves: leaf, petiole excluded)	mm^2	TRY	1415
Leaf area	Leaf area (in case of compound leaves: leaf, petiole included)	mm^2	TRY	1374
Leaf area	Leaf area (in case of compound leaves: leaf, undefined if petiole in- or excluded)	mm^2	TRY	897
Leaf area	Leaf area (in case of compound leaves undefined if leaf or leaflet, undefined if petiole is in- or exluded)	mm^2	TRY	2202
Leaf area	Leaf area	mm^2	BIEN	902
Leaf mass per area	Leaf area per leaf dry mass (specific leaf area, SLA or 1/LMA): petiole included	mg/mm^2	TRY	1935
Leaf mass per area	Leaf area per leaf dry mass (specific leaf area, SLA or 1/LMA): undefined if petiole is in- or excluded)	mg/mm^2	TRY	8777

| Leaf mass per area | Leaf mass per area | mg/mm² | BIEN | 835 |

Table 3: Overview of abiotic variables present in VegVault v1.0.0.

Abiotic variable name	Abiotic variable description	Abiotic variable unit	Source of data
bio1	Annual Mean Temperature	°C	CHELSA & CHELSA-TRACE21
bio4	Temperature Seasonality	°C	CHELSA & CHELSA-TRACE21
bio6	Min Temperature of Coldest Month	°C	CHELSA & CHELSA-TRACE21
bio12	Annual Precipitation	kg m-2 year-1	CHELSA & CHELSA-TRACE21
bio15	Precipitation Seasonality	Unitless	CHELSA & CHELSA-TRACE21
bio18	Precipitation of Warmest Quarter	kg m-2 quarter-1	CHELSA & CHELSA-TRACE21
bio19	Precipitation of Coldest Quarter	kg m-2 quarter-1	CHELSA & CHELSA-TRACE21
HWSD2	Soil type	Unitless	WoSIS

References

1. Fordham, D. A. *et al.* Using paleo-archives to safeguard biodiversity under climate change. *Science* **369**, eabc5654 (2020).

2. Nieto-Lugilde, D. *et al.* Time to better integrate paleoecological research infrastructures with neoecology to improve understanding of biodiversity long-term dynamics and to inform future conservation. *Environ. Res. Lett.* **16**, 095005 (2021).

3. Reitalu, T., Kuneš, P. & Giesecke, T. Closing the gap between plant ecology and Quaternary palaeoecology. *Journal of Vegetation Science* **25**, 1188–1194 (2014).

4. GBIF.org. GBIF Home Page. https://www.gbif.org (2024).

5. Bruelheide, H. *et al.* sPlot – A new tool for global vegetation analyses. *Journal of Vegetation Science* **30**, 161–186 (2019).

6. Botanical Information and Ecology Network (BIEN). BIEN: Botanical Information and Ecology Network. http://bien.nceas.ucsb.edu/bien/ (n.d.).

7. Williams, J. W. *et al.* The Neotoma Paleoecology Database, a multiproxy, international, community-curated data resource. *Quat. res.* **89**, 156–177 (2018).

8. Kattge, J. *et al.* TRY plant trait database – enhanced coverage and open access. *Global Change Biology* **26**, 119–188 (2020).

9. Karger, D. N. *et al.* Climatologies at high resolution for the earth's land surface areas. *Sci Data* **4**, 170122 (2017).

10. Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* **37**, 4302–4315 (2017).

11. Schipper, E. L. F., Dubash, N. K. & Mulugetta, Y. Climate change research and the search for solutions: rethinking interdisciplinarity. *Clim Change* **168**, 18 (2021).

12. Intergovernmental Panel On Climate Change. *Climate Change and Land: IPCC Special Report*

on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food

Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems. (Cambridge University Press,

2022). doi:10.1017/9781009157988.

13. Smith, J. *et al.* BioDeepTime: A database of biodiversity time series for modern and fossil

assemblages. *Global Ecology and Biogeography* **32**, 1680–1689 (2023).

14. Weigelt, P., König, C. & Kreft, H. GIFT – A Global Inventory of Floras and Traits for macroecology

and biogeography. *Journal of Biogeography* **47**, 16–43 (2020).

15. Sabatini, F. M. *et al.* sPlotOpen – An environmentally balanced, open-access, global dataset of

vegetation plots. *Global Ecology and Biogeography* **30**, 1740–1764 (2021).

16. Flantua, S. G. A. *et al.* A guide to the processing and standardization of global palaeoecological

data for large-scale syntheses using fossil pollen. *Global Ecology and Biogeography* **32**, 1377–

1394 (2023).

17. Batjes, N. H., Calisto, L. & de Sousa, L. M. Providing quality-assessed and standardised soil data

to support global mapping and modelling (WoSIS snapshot 2023). *Earth System Science Data*

16, 4735–4765 (2024).

18. Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H. & Zimmermann, N. E. CHELSA-

TraCE21k: Downscaled transient temperature and precipitation data since the last glacial

maximum. (2020) doi:http://dx.doi.org/10.16904/envidat.211.

19. Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H. & Zimmermann, N. E. CHELSA-TraCE21k

– high-resolution (1km) downscaled transient temperature and precipitation data since the Last

Glacial Maximum. *Climate of the Past* **19**, 439–456 (2023).

20. GBIF Secretariat. GBIF Backbone Taxonomy. (2023).

21. Mottl, O. taxospace - v0.0.0.9003. https://doi.org/10.5281/zenodo.14699491 (2024).

22. Mottl, O. vaultkeepr - v0.0.6. https://doi.org/10.5281/zenodo.14964737 (2025).

23. Mottl, O. *et al*. Global acceleration in rates of vegetation change over the past 18,000 years.

Science **372,** 860–864 (2021).

24. Šímová, I., Ordonez, A. & Storch, D. The dynamics of the diversity–energy relationship during the

last 21,000 years. *Global Ecology and Biogeography* **32**, 707–718 (2023).

25. Brown, K. A. *et al*. Trait-based approaches as ecological time machines: Developing tools for

reconstructing long-term variation in ecosystems. *Functional Ecology* **37**, 2552–2569 (2023).

26. Li, J. & Prentice, I. C. Global patterns of plant functional traits and their relationships to climate.

Commun Biol **7**, 1–14 (2024).

27. Shipley, B., Dilkina, B. & McGuire, J. MegaSDM: Modelling Species Ranges in The Past And Future.

Bulletin of the Florida Museum of Natural History **60**, 115–115 (2023).

28. Pollock, L. J. *et al*. Understanding co-occurrence by modelling species simultaneously with a

Joint Species Distribution Model (JSDM). *Methods in Ecology and Evolution* **5**, 397–406 (2014).

29. Ordonez, A. & Gill, J. L. Unravelling the functional and phylogenetic dimensions of novel

ecosystem assemblages. *Philosophical Transactions of the Royal Society B: Biological Sciences*

379, 20230324 (2024).

30. Wieczynski, D. J. *et al*. Climate shapes and shifts functional biodiversity in forests worldwide.

Proceedings of the National Academy of Sciences **116**, 587–592 (2019).

31. Zurell, D., Fritz, S. A., Rönnfeldt, A. & Steinbauer, M. J. Predicting extinctions with species

distribution models. *Cambridge Prisms: Extinction* **1**, e8 (2023).

32. Chacon, S. & Straub, B. *Pro Git*. (Springer Nature, 2014).

33. Global Names Architecture. Global Names Resolver. https://resolver.globalnames.org/.

34. Díaz, S. *et al*. The global spectrum of plant form and function. *Nature* **529**, 167–171 (2016).

35. CESNET. National Repository. https://data.narodni-repozitar.cz/ (2024).

36. Dillon, E. M. *et al*. Challenges and directions in analytical paleobiology. *Paleobiology* **49**, 377–

393 (2023).

37. Thuiller, W. On the importance of edaphic variables to predict plant species distributions – limits

and prospects. *Journal of Vegetation Science* **24**, 591–592 (2013).

38. Zuquim, G., Costa, F. R. C., Tuomisto, H., Moulatlet, G. M. & Figueiredo, F. O. G. The importance

of soils in predicting the future of plant habitat suitability in a tropical forest. *Plant Soil* **450**, 151–

170 (2020).

39. Velazco, S. J. E., Galvão, F., Villalobos, F. & Júnior, P. D. M. Using worldwide edaphic data to

model plant species niches: An assessment at a continental extent. *PLOS ONE* **12**, e0186025

(2017).

40. Haslett, J. & Parnell, A. A Simple Monotone Process with Application to Radiocarbon-Dated

Depth Chronologies. *Journal of the Royal Statistical Society Series C: Applied Statistics* **57**, 399–

418 (2008).

41. Augustine, S. P., Bailey-Marren, I., Charton, K. T., Kiel, N. G. & Peyton, M. S. Improper data

practices erode the quality of global ecological databases and impede the progress of ecological

research. *Global Change Biology* **30**, e17116 (2024).

42. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for

Statistical Computing, Vienna, Austria, 2024).