VegVault dataset: linking global paleo-, and neo-vegetation data with functional traits and abiotic drivers

Authors

Ondřej Mottl[1,2], Franka Gaiser[3], Irena Šímová[1,4], Suzette G. A. Flantua[5,6]

Affiliations

1. Center for Theoretical Study, Charles University, Jilská 1, CZ-11000 Prague, Czech Republic
2. Department of Botany, Faculty of Science, Charles University, Benátská 2, CZ-12801 Prague, Czech Republic
3. Sport Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER) & Bayreuth Center of Sport Science (BaySpo), University of Bayreuth, 95447 Bayreuth, Germany
4. Department of Ecology, Faculty of Science, Charles University, Viničná 7, CZ-12800 Prague, Czech Republic
5. Department of Biological Sciences, University of Bergen, PO Box 7803, N-5020 Bergen, Norway
6. Bjerknes Centre for Climate Research, Bergen, Norway

corresponding author(s): Ondřej Mottl (ondrej.mottl@gmail.com)

Abstract

Understanding the dynamics and persistence of biodiversity patterns over short (contemporary) and long (thousands of years) time scales is crucial for predicting ecosystem changes under global climate and land-use changes. A key challenge is integrating currently scattered ecological data to assess complex vegetation dynamics over time. Here, we present VegVault, an interdisciplinary SQLite database that uniquely integrates paleo- and neo-ecological plot-based vegetation data on a global and millennial scale, directly linking them with functional traits, soil, and climate information. VegVault currently comprises data from *BIEN*, *sPlotOpen*, *TRY*, *Neotoma*, *CHELSA*, and *WoSIS*, providing a comprehensive and ready-to-use resource for researchers across various fields to address questions about past and contemporary biodiversity patterns and their abiotic drivers. To further support the usability of the data,

29 VegVault is complemented by the {*vaultkeepr*} R package, enabling streamlined data access,
30 extraction, and manipulation. This study introduces the structure, content, and diverse
31 applications of VegVault, emphasizing its potential role in advancing ecological research to
32 improve predictions of biodiversity responses to global climate change.

Background & Summary

Biodiversity and ecosystem dynamics are currently profoundly influenced by global climate
change, creating an urgent need for robust predictive models capable of anticipating these
impacts. Such models are crucial in guiding conservation efforts during this global crisis.
However, many existing models rely on contemporary (present day) ecological data alone,
which limits their ability to capture the long-term responses of species and ecosystems to
environmental change. Incorporating paleoecological data—"ecology of the past" as opposed to
neo-ecological, or "ecology of the present"—offers a vital temporal perspective that can
enhance our understanding of biodiversity dynamics under shifting climate and land-use[1–3].

In recent years, an increasing number of large-scale ecological datasets have become available,
providing unprecedented opportunities to investigate ecosystem dynamics across spatial and
temporal scales. These datasets encompass contemporary vegetation data (e.g., *Global*
Biodiversity Information Facility[4] ('*GBIF*' hereafter), *sPlot*[5], *Botanical Information and Ecology*
Network[6] ('*BIEN*' hereafter), paleo-vegetation data (e.g., *Neotoma Paleoecology Database*[7]
('*Neotoma*' hereafter)), plant functional traits (e.g., *TRY plant trait database*[8] ('*TRY*' hereafter)),
and abiotic drivers (e.g., *CHELSA*[9], *WorldClim*[10]), which are all critical for understanding the
complex interactions that shape ecosystems. Yet, despite the growing availability of such
datasets, efforts to integrate these diverse sources into a unified, open-access framework
remain limited. An integrated database, built with reproducible workflows, is a crucial step
toward improving predictive models that estimate biodiversity dynamics under changing climatic
conditions. Addressing this gap requires interdisciplinary solutions[11,12] and a commitment to
compiling, processing, and linking data in a systematic and transparent manner. Currently, no
comprehensive framework exists to link global vegetation data across temporal scales with
functional traits and abiotic drivers.

There have been previous attempts to integrate vegetation data across extensive time scales
and link them with other data types. For example, *BioDeepTime*[13] compiles both contemporary
and fossil biodiversity time series. However, it lacks coverage of contemporary vegetation plots

and, more importantly, omits direct links to functional traits and abiotic data. GIFT[14] includes

vegetation biodiversity linked with functional traits but only holds taxonomic lists for specific

regions, without data on relative abundances. Additionally, the latter database lacks the long-

term dimension of vegetation dynamics over time, derived from the currently existing large

databases containing paleoecological records, such as the Neotoma database.

VegVault addresses these limitations by offering a unique resource that compiles extensive

paleo- and neo-ecological vegetation data into a single, accessible SQLite database,

accompanied by functional traits and abiotic information (i.e., climate and soil). Its design

bridges paleo- and contemporary data, enabling the study of long-term processes, and linking

them to today's patterns. Thereby, VegVault offers the possibility to specifically analyse the

mechanisms of ecosystem responses to environmental change, which enhances our ability to

study and ultimately predict future biodiversity trajectories. See **Table 1** for the comparative

table that summarises the strengths and weaknesses of existing databases versus VegVault.

Specifically, VegVault advances analyses across spatial, temporal, and taxonomic scales in

several ways:

• **Contemporary vegetation data**: VegVault incorporates plot-based data from *BIEN*[6] and

sPlotOpen[15] (open-access version of sPlot), including essential "true negative" information

about species' absence. These datasets are pre-cleaned for taxonomic and georeferencing

errors, ensuring high reliability for subsequent analyses. The focus on plot-based data

rather than occurrence data (e.g., *GBIF*) provides a closer match to paleoecological data

formats, facilitating robust comparisons across temporal scales.

• **Paleo-vegetation data**: VegVault employs standardised fossil pollen data processed using

the *FOSSILPOL* workflow[16] to integrate records globally from Neotoma. This workflow

ensures quality control and creates "ready-for-analysis" data through a reproducible and

detailed process for handling large fossil pollen dataset compilations (see **Methods** for

details).

• **Interdisciplinary integration**: VegVault combines past and present vegetation data with

functional traits and abiotic information to support comprehensive ecological analyses. The

abiotic information includes soil data from the *WoSIS Soil Profile Database*[17] by *World Soil*

Information Service ('*WoSIS*' hereafter), and global temperature and precipitation datasets

from *CHELSA* and *CHELSA-TRACE21*[18,19], providing essential environmental context.

Functional trait data from the *TRY* database and *BIEN* enrich the database by linking

species characteristics to ecological roles and responses. This allows researchers to move

beyond taxonomic diversity and explore functional dimensions, offering deeper insights into
ecosystem processes.

• **Taxonomic classification**: VegVault ensures consistent and standardised taxonomic
information by aligning all taxa present in vegetation data—both paleo and contemporary—
with the GBIF backbone taxonomy[20]. This process, implemented via the {*taxospace*} R
package[21], automates taxonomic classification by matching names to an existing taxonomy
based on similarity (e.g., "Ulmus Type" will be classified as genus *Ulmus*). While original
taxonomic names are preserved within the VegVault, users can access data at varying
taxonomic resolutions (e.g., species, genus, family; if available) to suit specific analytical
needs. The same standardisation approach is applied to vegetation and trait data,
facilitating seamless integration between datasets. For instance, enabling the calculation of
genus-level average trait values, when species-level identification is unavailable.

• **Accessibility**: VegVault is designed with user accessibility in mind. The SQLite database is
supported by the {*vaultkeepr*} R package[22], which simplifies and streamlines the process of
data extraction for project-specific needs. It ensures that researchers can efficiently retrieve
and subsequently manipulate data in a reproducible and transparent manner.

The comprehensive nature of VegVault enables diverse applications across scientific disciplines.
Its integration of fossil pollen data and functional traits allows researchers to investigate long-
term vegetation dynamics and their responses to past climate variations. Such integration
provides critical context for past ecological trends[e.g.,23,24], including trait information, which is
currently underrepresented in paleoecological analyses of vegetation dynamics[25]. While there
are now studies showing the relationship between functional traits and climate for global
contemporary data[26], VegVault allows testing the universality of such relationships across
temporal scales. In addition, VegVault provides the needed data for advanced modelling
approaches, such as comparing species distributions across temporal scales[e.g.,27], and
analysing biodiversity patterns using Joint Species Distribution Models[28]. VegVault also
facilitates studies of functional ecology and biogeography by enabling assessments of shifts in
ecosystem functioning and climatic drivers over time, and by identifying the taxonomic and
functional novelty of communities undergoing climate-driven changes[e.g.,29,30]. Furthermore, the
detailed climate data (past and present) linked with vegetation records provide a robust
foundation for examining the impacts of past and present climate change on ecosystems[e.g.,31].
These capabilities make VegVault a valuable interdisciplinary resource for predicting
biodiversity patterns and informing conservation strategies and policy decisions aimed at

mitigating climate change effects. While this is not an exhaustive list of potential applications,
VegVault offers a rich and accessible resource that can significantly advance ecological
research and global biodiversity conservation.

Methods

VegVault workflow and reproducibility

VegVault integrates data from multiple well-established primary sources to provide a
comprehensive view of vegetation dynamics. The current sources of data are summarised in
Table 1.

The data processing and integration into VegVault follow a meticulous workflow to ensure
consistency and reliability. Each data source is first accessed and processed individually using
a specific pipeline. See **Figure 1** for the workflow process of accessing and processing all data
sources (1 to 6 in **Figure 1**).

Data extraction and pre-processing

139 To ensure the traceability of data, VegVault is designed as a versioned project. Each version
140 (including the current v1.0.0) is created using the git Tags system[32]. Each Tag is a snapshot of
141 code at certain points in time. Each of the individual data pipelines, coded in R programming
142 language, is publicly available and stored in a separate GitHub repository with a specific git Tag
143 and published with DOI (see legend of **Figure 1**).

Fossil pollen record (**Figure 1.7**)

Fossil pollen records have been downloaded from the *Neotoma Paleoecological Database*[7]
using their API on 26[th] June 2023. All data acquisition and processing have been done using the
FOSSILPOL: Workflow for processing and standardizing global paleoecological pollen data[16]
(version 1.0.0). This includes the selection of depositional environments, ecological groups,
chronology control point types, and a minimum number of chronology control points to construct
age-depth models. Individual samples and records have been filtered by age limits, number of
pollen grains, maximum age interpolation, and number of valid levels. In addition, the accuracy
of fossil pollen data is increased by re-estimating all age-depth models using the Bayesian
probabilistic approach[33] and including the information about individual age uncertainty. For field-

specific terminology, see Flantua et al.[16] and for more information about *FOSSILPOL* workflow,
see https://bit.ly/FOSSILPOL.

Individual steps in detail are as follows:

- 157 • Only records with type "pollen" have been requested with valid geological coordinates.
158 Specifically, longitude between -180 and 180, and latitude between -90 and 90.
- 159 • Records have been filtered by their depositional environments, where in this case we
160 only kept lakes, bogs, and mires. See **Table S1** for a list of selected depositional
161 environments.
- 162 • Only taxa in specific ecological groups have been accepted. See **Table S2** for a list of
163 selected ecological groups.
- 164 • All age-depth models have been recalculated using the {*Bchron*} package[33]. Only
165 records with at least 5 chronology control points of specific types have been retained.
166 See **Table S3** for a list of valid control types.
- 167 • To ensure the highest possible data quality and facilitate numerical comparisons across
168 multiple records, several filtering criteria were carefully chosen and applied:
  - 169 o Exclusion of samples with a total pollen count below a specified threshold. While
170 the preferred minimum number of pollen grains was initially set at 150, this
171 threshold resulted in a significant loss of records, particularly in regions with
172 limited data coverage. To balance data retention and quality, the threshold was
173 adjusted to 125, with the condition that less than 75% of the samples would have
174 a low pollen sum (i.e., lower than 125 pollen grains).
  - 175 o Exclusion of samples older than 3000 years of the last chronology control point.
  - 176 o Exclusion of samples beyond 20,000 cal yr BP.
  - 177 o Exclusion of records if the total number of samples was fewer than 5.

178 The VegVault-specific version of FOSSILPOL with all the individual R scripts and tables is
179 stored as a GitHub repository titled *VegVault-FOSSILPOL*
180 (https://github.com/OndrejMottl/VegVault-FOSSILPOL/tree/v1.0.0) and can be accessed as
181 DOI: 10.5281/zenodo.14745060.

Contemporary vegetation plots (**Figure 1**.8)

183 The primary sources of contemporary plot-based vegetation data are *BIEN*[6] (Botanical
184 Information and Ecology Network) and *sPlotOpen*[15] (the open-access version of sPlot[15]).

185 The GitHub repository titled *VegVault-Vegetation_data*
186 (*https://github.com/OndrejMottl/VegVault-Vegetation_data/tree/v1.1.0*)*,* used to process data for
187 contemporary vegetation plots, can be accessed as DOI: 10.5281/zenodo.16729978.

BIEN

The plot-based data have been downloaded using the {*RBIEN*} R package[34] (version 1.2.7) on
2[nd] August 2023 by using the function `BIEN::BIEN_plot_datasource()`. The following columns
have been retained: "datasource_id", "datasource", "plot_name", "sampling_protocol",
"methodology_reference", "methodology_description", "longitude", "latitude", "plot_area_ha",
"subplot", "individual_count", "family_matched", "name_matched", "name_matched_author",
"verbatim_family", "verbatim_scientific_name", "scrubbed_species_binomial",
"scrubbed_taxonomic_status", scrubbed_family", "scrubbed_author". All columns have been
renamed using the "snake case". Finally, rows containing NA values in "datasource_id,
"datasource", "plot_name", "longitude", "latitude", and/or "plot_area_ha" have been filtered out.

sPlotOpen

199 The sPlotOpen dataset[35] (version 2.0) have been downloaded on 26[th] September 2023. Tables
200 "DT2.oa" and "header.oa" have been linked via the "PlotObservationID" column, and rows
201 containing NA values in "PlotObservationID", "GIVD_ID", "Longitude", "Latitude", and/or
202 "Releve_area" have been filtered out. All columns have been renamed using the "snake case".

Functional traits of vegetation (**Figure 1.9**)

The primary sources of vegetation functional traits are *TRY Plant Trait Database*[8] and *BIEN*[6].

We followed Diaz et al. (2016)[36] and selected traits representing *Stem specific density*, *Leaf*
nitrogen content per unit mass, *Diaspore mass*, *Plant height*, *Leaf area*, and *Leaf mass per area*.

A GitHub repository titled *VegVault-Trait_data* (*https://github.com/OndrejMottl/VegVault-*
Trait_data/tree/v1.2.0)*,* used to process data for plant vegetation traits, can be accessed as
DOI: 10.5281/zenodo.16729756.

TRY

We made a data request from the *TRY Plant Trait Database* on the 29[th] of August 2023 (ID of
the request is 28498), requesting all values for traits with codes 3106, 4, 3108, 3110, 3112,

3114, 3116, 3117, 14, and 26 (as they are the closest to the Diaz et al. (2016)[36] description).

Next, we used the {*rtry*} R package[37] (version 1.1.0) to import the data into R.

For each trait ("TraitID"), we extracted all relevant observations ("ObservationID"), ensuring that
only observations unique to each trait are considered (i.e., filtering out overlapping observations
with other traits). Next, we identified all unique data ("DataID") associated with each trait.
Certain data were excluded ("DataID": 2221, 2222, 2223, 2224, 2225, 3646, 3647, 3698, 3699,
3727, 3728, 3730, 3731, 3849, 3850, 4029, and 4030) as not meaningful variation of the trait,
e.g. *height at 15 days.* We filtered the observations and extracted the actual trait values,
ensuring that only non-missing values were kept. Third, we extracted covariate information
(additional data associated with each observation but not directly a trait value stored in the
"DataName" column). Finally, all columns were renamed using the "snake case".

As there are differences in trait names across sources of data (e.g., "*Leaf nitrogen (N) content*
per leaf dry mass" and "*leaf nitrogen content per leaf dry mass*"), we added a new variable *Trait*
Domain that groups traits together following the trait selection by Diaz et al. (2016)[36]. For
example, the traits "*Leaf area (in case of compound leaves: leaf, petiole excluded)*" and "*Leaf*
area (in case of compound leaves: leaf, petiole included)" are grouped under the "Leaf area"
Trait Domain. See **Table 3** for the full list of traits and their *Trait Domain*. Such grouping is also
aimed to serve as an efficient extraction of traits across both *TRY* and *BIEN*.

BIEN

The trait data have been downloaded using the {*RBIEN*} R package[34] (version 1.2.7) on 15th
December 2023 by using the function `*BIEN::BIEN_trait_trait()*`. Specifically, we requested data
for traits "*whole plant height*", "stem wood density", "*leaf area*", "*leaf area per leaf dry mass*",
"*leaf nitrogen content per leaf dry mass*", and "*seed mass*". The following columns have been
retained: "trait_name", "trait_value", "unit", "id", "longitude", "latitude", "method", "url_source",
"source_citation", "project_pi", "scrubbed_species_binomial", and "access".

Abiotic data (**Figure 1.10**)

The primary sources of abiotic data are *CHELSA*[9], *CHELSA-TRACE21*[18,19], and *WoSIS Soil*
Profile Database[17]. The first two data sources provide high-resolution downscaled climatic data,
while the latter offers detailed soil information (only available for contemporary data). The abiotic
variables included in VegVault are summarised in **Table 4**.

The GitHub repository titled *VegVault-abiotic_data (https://github.com/OndrejMottl/VegVault-*
abiotic_data/tree/v1.1.0), used to process climate and soil abiotic data, can be accessed as
DOI: 10.5281/zenodo.14745293.

CHELSA

248 Climate data have been downloaded from *CHELSA*[9] (version 2.1) using the
249 {*ClimDatDownloadR*} R package[38] on the 8th of September 2023 with the function
250 `*ClimDatDownloadR::Chelsa.Clim.download()*`. The specific bio-variables selected are 1, 4, 6,
251 12, 15, 18, and 19. Next, we used the {*terra*} R package[39] to spatially aggregate the values
252 using the function `*terra::aggregate(factor = 25, fun = "median")*`.

CHELSA-TRACE21

254 Paleoclimate data have been downloaded from *CHELSA-TRACE21*[18,19] (version 1.0) on the 31st
255 December 2023. We only downloaded values for each 500-year time-slice between 0 and
256 18,000 years before present (BP) for the selected bio-variables (1, 4, 6, 12, 15, 18, and 19). For
257 each time slice, we also used the function `*terra::aggregate(factor = 25, fun = "median")*` to
258 spatially aggregate the values.

WoSIS Soil Profile Database

As soil information has been previously confirmed as important predictors for vegetation
distribution[40–42], we opted for including such data, though only representing the present day. We
downloaded the soil data from the *WoSIS Soil Profile Database*[17] on the 11th of September 2023
(both HWSD2_RASTER.zip and HWSD2_DB.zip). We extracted the soil type names (column
"HWSD2_SMU_ID") by combining tables "HWSD2_SMU" and "D_WRB4", and added this
information to the raster. Finally, we resampled the raster using the `*terra::resample(method =*
"near")` function to match the resolution of climate data.

Data integration (**Figure 1.11**)

268 All processing pipelines with their corresponding Tags are then migrated into an SQLite
269 database   using   the   GitHub   repository   titled   *VegVault*
270 (https://github.com/OndrejMottl/VegVault/tree/v1.0.0-website_rewiev)*,* which can be accessed
271 as DOI: 10.5281/zenodo.16987345).

Migrating sPlotOpen vegetation data

The Dataset name ("dataset_name") has been created from "plot_observation_id" as "splot_[plot_observation_id]". The original data source (e.g., sub-databases) from column "givd_id" has been stored in the *DatasetSourcesID* table.

The Sample name ("sample_name") has been created using column "plot_observation_id" as "splot_[plot_observation_id]". The Sample Size ("sample_size") has been created from the "releve_area" column. All samples are automatically assigned an age of 0.

Taxonomic names have been extracted from column "Species" and the abundances from "Original_abundance".

Migrating BIEN vegetation data

The Dataset name ("dataset_name") has been created as "bien_[row number]". The original data source (e.g., sub-databases) from column "datasource" has been stored in the *DatasetSourcesID* table. The Sampling method has been extracted from the "methodology_description" column, and its reference has been extracted from the "methodology_description" column.

The Sample name ("sample_name") has been created as "bien_[row number]". The Sample Size ("sample_size") has been created from the "plot_area_ha " column, and multiplied by 10,000 to be stored in square meters. All samples are automatically assigned an age of 0.

Taxonomic names have been extracted from column "name_matched" and the abundances from "taxon_name".

Migrating fossil pollen data

The Dataset name ("dataset_name") has been created from "dataset_id" (fossilpol_[dataset_id]). Note that the column "dataset_id" is from the primary source and does not match the "dataset_id" in VegVault. The original data source from column "source_of_data" has been stored in the *DatasetSourcesID* table. The Sampling method has been extracted from the "depositionalenvironment" column. The individual Dataset Reference have been extracted from the column "doi".

The Sample name ("sample_name") has been created using columns "dataset_id" and
"sample_id" as "fossilpol_[dataset_id]_[sample_id]". Ages have been extracted from column
"levels".

Age uncertainties from age-depth models have been extracted from "age_uncertainty" column.

Taxonomic names and their abundances have been extracted from the column
"counts_harmonised".

Migrating TRY functional traits

The Dataset name ("dataset_name") has been created as "try_[row number]". The original data
source (e.g., sub-databases) from column "dataset" has been stored in the *DatasetSourcesID*
table. The individual Dataset Reference have been extracted from the column
"reference_source".

The Sample name ("sample_name") has been created as "try_[row number]". The individual
Sample reference has been extracted from "dataset_reference_citation". All samples are
automatically assigned an age of 0.

Trait names have been extracted from "trait_full_name", taxonomic names from the column
"acc_species_name", and trait values from "trait_value".

Migrating BIEN functional traits

The Dataset name ("dataset_name") has been created as "bien_traits_[row number]". The
original data source (e.g., sub-databases) from column "project_pi" has been stored in the
DatasetSourcesID table. The individual Dataset Reference have been extracted from the
column "source_citation".

The Sample name ("sample_name") has been created using column "id" as "bien_traits_[id]". All
samples are automatically assigned an age of 0.

Trait names have been extracted from "trait_name", taxonomic names from the column
"scrubbed_species_binomial", and trait values from "trait_value". Trait "leaf mass per area" is
calculated from "leaf area per leaf dry mass" as 1/value.

Creation of *gridpoints* of abiotic data

We developed a data structure that provides readily available environmental context for each
vegetation (and trait) record by creating spatio-temporal links between these records and abiotic

information. As the rasters are not suitable to be stored in an SQLite database, we created
artificial points, called '*gridpoints*', located in the middle of each raster cell. This resulted in a
uniform spatio-temporal matrix of gridpoints holding the abiotic information (see **Figure 2B** for
an example area).

Gridpoints Dataset names ("dataset_name") have been created as "geo_[longitude]_[latitude]"
and gridpoints Sample names ("sample_name") have been created as
"geo_[longitude]_[latitude]_[age]". Next, we linked *gridpoints* and other non-*gridpoint Samples*,
namely *vegetation_plot*, *fossil_pollen_archive*, and *traits* (see **Internal Dataset Structure**
section), and calculated the spatial and temporal distances between them. We kept any
gridpoint Sample within 50 km and/or 5000 years from any other non-*gridpoint Sample* and
discarded the rest. See **Figure 2C** for an example of linking abiotic information at *gridpoints* and
vegetation data. In VegVault, users can select the information for each non-*gridpoint Sample*
from the closest spatio-temporal abiotic data or get the average from all surrounding *gridpoints*.
We hope that by providing such a comprehensive and well-linked structure between vegetation
and abiotic data, VegVault enhances the ability to study the interactions between vegetation and
their environment, facilitating the workflow for advanced ecological research and modelling
efforts.

Taxon classification

As VegVault consists of data on taxa from various sources, the {*taxospace*} R package[21] has
been used to classify the diverse taxa into a unifying backbone in the VegVault. The
{*taxospace*} tool automatically aligns taxon names with the taxonomical backbone of the *GBIF*[20].
Specifically, we find the best match of the raw names of taxa using the *Global Names*
Resolver[43], which is then aligned with *GBIF*. The resulting information of taxonomic
classification, detailed up to the family level, is stored for each taxon, ensuring consistency and
facilitating comparative analyses across different datasets. However, note that taxonomic
classification down to the species level is not available for each taxon (e.g., some fossil pollen
types can only be identified to the genus or family level). For several taxa, we were unable to
find a matching classification. Note that the taxon classification is additional information; the
original taxon name is always present and returned by default. Finally, we would like to stress to
the user that this classification is an automated process and may contain errors.

Data Records

The VegVault v1.0.0 is available at the *National Repository*[44] ('*REPO*') under DOI 10.48700/datst.t21cz-2nq07. It is an SQLite database of a size of ~ 110 GB, which contains > 480,000 *Datasets*, > 13,000,000 *Samples*, > 100,000 *Taxa*, 6 vegetation *Traits*, > 11,000,000 *Trait* values, and 8 abiotic variables. The spatio-temporal overview of all vegetation data can be found in **Figure 3**.

Internal Dataset Structure

Currently, the dataset consists of 31 interconnected tables with 87 fields (variables), which are described in detail below. See **Figure 4** for the structure (schema) of the SQLite database. However, the internal structure of VegVault may not be of relevance to a user, as the {*vaultkeepr*} R package processes all the data to output only the most relevant information in a "ready-to-analyse" format (see **Usage Notes** section). We will describe all tables and their inter-relationships in the following sections.

Metadata tables

There are currently several tables that are not directly linked to the data. First is the *Authors* table, which describes the authors and maintainers of VegVault, including the contact information. The next table is *version_control* table, which stores the information about database versions with a description of individual changes at a specific point in time. See **Table 5** and **Table 6** for the description of individual columns, respectively. Finally, there are *sqlite_stat1* and *sqlite_stat4* tables, which hold descriptive information about database indexes, used for improving the performance of data extractions. See **Table 7** and **Table 8** for the description of individual columns, respectively.

Datasets table

The table contains one *Dataset* per row, with each *Dataset* may contain: a unique *Dataset ID* ("dataset_id*"), *Dataset name* ("dataset_name"), geographic location ("coord_lat", "coord_long"), Dataset Type ("dataset_type_id*"), Dataset Source* ("data_source_id"), *Dataset Source Type* ("dataset_source_type_id"), and *Sampling Method* ("sampling_method_id"). See **Table 9** for the description of individual columns.

DatasetTypeID table

The table defines the most basic classification of each *Dataset*, ensuring that the vast amount of

data is categorised systematically (see **Table 10** for the description of individual columns).

Currently, VegVault contains the following *Dataset Types*:

- `*vegetation_plot*`: This type includes contemporary vegetation plot data, capturing
contemporary vegetation characteristics and distributions.

- `*fossil_pollen_archive*`: This type encompasses past vegetation plot data derived from
fossil pollen records, providing insights into past vegetation and climate dynamics.

- `*traits*`: This type contains functional trait data, detailing specific characteristics of plant
species that influence their ecological roles.

- `*gridpoints*`: This type holds artificially created *Datasets* to manage abiotic data, here
climate and soil information (a dataset type created to hold abiotic data, see details in
the **Methods** section).

See **Figure 5A** for an overview of the number of *Datasets* in each *Dataset Type.*

DatasetSourceTypeID table

The table contains information about the primary data source (see **Table 11** for the description

of individual columns). Therefore, each *Dataset* is derived from a specific *Source-Type*

including:

- **BIEN**[6] (Botanical Information and Ecology Network).

- **sPlotOpen**[15] (The open-access version of sPlot).

- **TRY**[8] (TRY Plant Trait Database).

- **Neotoma-FOSSILPOL** (FOSSILPOL[16]: Workflow for processing and standardizing
global paleoecological pollen data from *Neotoma*[7]) Note that we specifically state
Neotoma-FOSSILPOL and not just *Neotoma*, as *FOSSILPOL* not only provides the data
acquisition but also alters it (e.g., creating new age-depth models). It also addresses
major challenges in paleoecological data integration, such as age uncertainty, by
incorporating probabilistic age-depth models and their associated uncertainty matrices.
This enables the propagation of temporal uncertainty in subsequent analyses, a critical
advancement for robust macroecological studies, previously flagged as a major issue
with paleo-data[45].

- **gridpoints** (artificially created *Datasets* for abiotic data). See details in the **Methods**
section.

See **Figure 5B** for an overview of the number of *Datasets* for each *Dataset Source-Type.*

DatasetSourceTypeReference table

The table links the *Dataset Source-Types* to their specific *Reference.* Note that one *Dataset*
Source-Type may have several *References. See* **Table 12** for the description of individual
columns.

DatasetSourcesID *table*

The table contains information about the provider (*Dataset Source*) of the original data for the
primary data source. The VegVault v1.0.0 currently includes 706 sources of *Datasets,* where
each dataset can also have one or more direct references to specific data (see below), ensuring
that users can accurately cite and verify the sources of their data. See **Table 13** for the
description of individual columns.

DatasetSourcesReference table

The table links the *Dataset Sources* to their specific *Reference.* Note that one *Dataset Source*
may have several *References.* See **Table 14** for the description of individual columns.

SamplingMethodID table

The table contains information about specific *Sampling Method* in the form of an abbreviated ID,
which is only available for both *vegetation_plot* and *fossil_pollen_archive Datasets*, and varies
substantially, reflecting the diverse nature of the data collected. Such information is crucial for
understanding the context and limitations of each *Dataset Type.* For contemporary vegetation
plots, sampling involves standardised plot inventories and surveys that capture detailed
vegetation characteristics across various regions. Fossil pollen data are collected from sediment
records from numerous different depositional environments[16] representing past vegetation and
climate dynamics. See **Table 15** for the description of individual columns.

SamplingMethodReference table

The table links the *Dataset Sampling Methods* to their specific *Reference.* Note that one *Dataset*
Sampling Methods may have several *References.* See **Table 16** for the description of individual
columns.

DatasetReference table

This table links individual *Datasets* to their specific *References* (see **Table 17** for the description
of individual columns). Note that one *Dataset* may have several *References.* To support robust

and transparent scientific research, each *Dataset* in VegVault can have multiple *References* at
different levels as the raw data was pooled from other databases that provide data from various
authors. The *Dataset Source-Type*, *Dataset Source*, and *Sampling Method* can all have their
own *References*, providing detailed provenance and citation information. This multi-level
referencing system enhances the traceability and validation of the data.

Samples table

Samples represent the main unit of data in the *VegVault*, serving as the fundamental building
blocks for all analyses. There are currently over 13 million *Samples* in the VegVault v1.0.0 (of
which ~ 1.6 million are *gridpoints* of abiotic data, see **Methods** section). For contemporary and
fossil pollen records, each Sample is a representation of vegetation community at a specific
location and time. The *Samples* table contains one *Sample* per row, with each *Sample* may
contain: a unique *Sample ID* ("sample_id*"), *Sample name* ("sample_name"), temporal
information about *Sample* ("age*"), *Sample size* (size of the plot if available; "sample_size_id"*),*
and additional information about *Sample* ("sample_details"; this is currently not being used in
v1.0.0.). As VegVault encompasses both contemporary and paleo-data, accurate age
information for each *Sample* is needed. Contemporary *Samples* are assigned an age of 0, while
Samples from fossil pollen records are in calibrated years before the present (cal yr BP). The
'present' is here specified as 1950 AD. *See* **Table 18** for the description of individual columns.

DatasetSample table

467 Each *Sample* is linked to a specific *Dataset* via the *DatasetSample* table, which ensures that
468 every *Sample* is correctly associated with its corresponding *Dataset Type* (whether it is
469 *vegetation_plot*, *fossil_pollen_archive*, *traits*, or *gridpoint*) and other *Dataset* properties (e.g.,
470 geographic location). One *Dataset* contains several *Samples* only in cases where *Samples*
471 differ in time ("age*"). See **Table 19** for the description of individual columns.

SampleSizeID table

The size of vegetation plots can vary substantially. This detail is crucial for ecological studies
where plot size can influence species diversity and abundance metrics, thus impacting follow-up
analyses and interpretations[46]. To account for this variability, information about the plot size is
recorded separately for each contemporary *Sample*. See **Figure 6** for an overview of the
number of *Samples* in each size category and **Table 20** for the description of individual columns.

SampleUncertainty table

Each *Sample* from the *fossil_pollen_archive Dataset* is also associated with an uncertainty
matrix generated during the re-estimation of ages using FOSSILPOL workflow[16]. This matrix
provides a range of potential ages derived from age-depth modelling, reflecting the inherent
uncertainty in dating paleoecological records. See an example of detailed age uncertainty
information for one fossil pollen record in **Figure 7**, and the **Table 21** for the description of
individual columns

SampleReference table

Each *Sample* in VegVault can have specific *References* in addition to those at the *Dataset*-level.
These individual *Sample References* provide detailed provenance and citation information,
ensuring that users can trace the origin and validation of each data point. Note that a single
Sample can have several *References*. This level of referencing enhances the transparency and
reliability of the data, especially when the dataset continues to be updated in the future. See
Table 22 for the description of individual columns.

Taxa table

The VegVault database records the original taxonomic names derived directly from the primary
data sources, and currently, it holds over 100 thousand taxonomic names. See **Methods** for the
names of columns used from the primary sources. See **Table 19** for the description of individual
columns.

SampleTaxa table

Each individual *Taxon* is linked to corresponding *Samples* through the *SampleTaxa* table,
ensuring accurate and systematic association between species and their ecological data. Note
that the abundance information varies across the primary data sources (see **Methods** for the
names of columns used as abundance). Therefore, users have to be careful while processing
data from various sources. See **Table 24** for the description of individual columns.

TaxonClassification table

Each taxonomic name undergoes an automated classification (see **Methods** section), and
results are stored in the *TaxonClassification* table. Taxonomic classification for some *Taxa*
might be only available down to the genus or family level, while some most of the data is
classified to species level. See **Figure 8** for an overview of the number of species, genus and
family classification. However, 1312 (1.2%) of all *Taxa* currently do not contain classification,

and only their name from the primary source is available. The best possible name is returned for
each request, which means that *Taxa* without classification will return the name from the
primary source. See **Table 25** for the description of individual columns.

TaxonReference table

Each *Taxon* might get a reference. Currently, this is used to track the origin of the *Taxon* name
(i.e., which primary source was used first with this *Taxon*). Note that *Taxa*, generated from
taxonomic classification (see **Methods**), are associated with "taxospace" reference. See **Table**
26 for the description of individual columns.

Traits table

The *Traits* table contains the list of functional traits currently contained in VegVault. The table
contains one *Trait* per row, with each *Trait* may contain: a unique *Trait* ID ("trait_id"), original
Trait name from primary source ("trait_name"*;* see **Methods** for the details about the specific
columns used from primary sources), and *Trait Domain* ("trait_domain_id"). See **Table 20** for
the description of individual columns.

TraitsDomain table

Traits are grouped into *Trait Domains* to allow for easier aggregation of *Traits* across data
sources (see **Methods** section). In total, six *Trait Domains* are present: "Stem specific density",
"Leaf nitrogen content per unit mass", "Diaspore mass", "Plant height", "Leaf area", "Leaf mass
per area", following Diaz et al. (2016)[36]. See **Table 3** for an overview of *Traits* in various *Trait*
Domains and the number of *Trait* values. Yet, it is up to the user to decide how to further
aggregate trait values if multiple trait samples of one *Trait Domain* are available for the same
environmental or taxonomic entity. See **Table 28** for the description of individual columns.

TraitsValue table

In general, data of functional traits of vegetation taxa follow the same structure of the *Dataset*
and *Samples* obtained directly from the *Dataset Source-Types*. Therefore, the *TraitsValue* table
contains not only the actual measured value of *Trait* observation but also information about
linking information across *Datasets*, *Samples*, and *Taxa.* This comprehensive linkage ensures
that each *Trait* value is accurately associated with its relevant ecological, environmental and
taxonomic context. See **Table 29** for the description of individual columns.

 TraitsReference table

539 To ensure clarity and reproducibility, each *Trait* in the VegVault can have additional *References*

540 beyond the general *Dataset* and *Sample References*. These *Trait*-specific *References* provide

541 detailed provenance and citation information, supporting rigorous scientific research and

542 enabling users to trace the origins and validation of each trait value. See **Table 30** for the

543 description of individual columns.

AbioticVariable table

As the VegVault contains abiotic variables from several primary sources, the *AbioticVariable*

table contains descriptions of abiotic variables ("abiotic_variable_name"), their units

("abiotic_variable_unit"), and measurement details ("measure_details"). See **Table 4** for the

overview of abiotic variables and their primary source, and **Table 31** for the description of

individual columns.

AbioticData table

The table holds the actual values of abiotic variables (the units are the same for each *Abiotic*

Variable). See **Table 32** for the description of individual columns.

AbioticVariableReference table

Each *Abiotic Variable* can have a separate *Reference*, in addition to a *Dataset* and *Sample*. See

Table 33 for the description of individual columns.

AbioticDataReference table

As abiotic data is stored in a special *Dataset Type—gridpoints*, we have also pre-calculated the

spatio-temporal distance of *gridpoints* to other Datasets. See **Methods** for details about the

creation of *gridpoints*. There, the *AbioticDataReference* table contains information about specific

distances among *Samples* for both temporal (age in years) and geographic (distance in km).

See **Table 34** for the description of individual columns.

References table

This table contains all *References*, independent of the source of the reference and the type of

data. Each row contains a single *Reference* that is then linked to the type of data which is being

referenced. This allows to have a single *Reference* to be used across data types, but also one

data point having many different references.

Moreover, most primary sources of the data have a license, which requires correct attribution.
Therefore, each *Reference* has information on whether such a *Reference* needs to be cited
while using the specific data ("mandatory" column).

Technical Validation

571 The data validity of VegVault is driven by the completeness, accuracy, and reliability of the
572 source data. Several steps have been undertaken to ensure that VegVault provides high-quality
573 data for research applications. For the fossil pollen records, strict data-quality rules have been
574 applied by using the *FOSSILPOL* workflow[16]. The data preparation and execution are backed
575 up by open-source publicly available code versioned and controlled by a git Tag system.

Usage Notes

Guidelines for data reuse and update

Due to the version system of VegVault, the dataset is expected to be updated using new data
and/or changing the process of data migration and processing. Therefore, we recommend citing
the latest VegVault version with the corresponding DOI, see bit.ly/VegVault for the latest release
of the dataset (v1.0.0 can be found here: doi.org/10.48700/datst.t21cz-2nq07).

As the primary data sources are held under various licences, VegVault has a *CC BY 4.0*
Licence (https://creativecommons.org/licenses/by/4.0/), allowing the copying and redistribution
of the material as long as correct attribution is given. We recommend that users carefully review
the restrictions associated with data usage. In addition, when citing the usage of VegVault,
users should always cite the primary sources of the data, as well as any individual datasets. For
those purposes, we created a function `*vaultkeepr::get_references()*` within the {*vaultkeepr*} R
package (see below) that will output a list of recommended references for the specific data
compilation created by the user.

Limitations and considerations for future research

Potential limitation lies in the completeness, accuracy, and reliability of the primary data sources.
For example, it has been suggested that improper data practices erode the quality of global
ecological databases such as *TRY*[47], which have been used in VegVault. We do not attempt to
solve these problems as VegVault's main purpose is the re-usability of the data.

Next, as VegVault database contains more than 100GB of data, we advise the users to make
sure that their machine has enough space on their drives, as downloads can be challenging on
certain machines (e.g., a laptop).

Finally, to increase the accessibility of the data, we have also deposited the flat version of the
database (i.e. individual tables saved as CSV files), accessible under DOI: 10.48700/datst.thktf-
5rr35. However, note that many of those files are larger than 1GB and some machines do not
have enough memory (RAM) to fully load them.

{vaultkeepr}: R-package for accessing the VegVault database

In order to make usage of the VegVault as easy as possible and facilitate access to SQLite
databases, we have developed an R-package called {*vaultkeepr*}, providing a suite of functions
to effectively interact with the database directly from the R programming environment[48].

Functions include accessing the database, extracting datasets, filtering samples, and accessing
specific taxa and traits. Therefore, we suggest all users to use the SQLite database
accompanied by {vaultkeepr} R package to retrieve only project-specific data without the need
to load data into memory.

This package is a well-tested interface (>95% code coverage). See **Figure 9** for details about
individual steps to retrieve the data from VegVault. For more about the package, the function
documentation and examples of code usage, please see {*vaultkeepr*} website
(https://bit.ly/vaultkeepr). Here we present 3 examples of potential projects and how to obtain
data for such projects from VegVault using {*vaultkeepr*} R-package. Note that we are specifically
not trying to do any analysis, only presenting a way to obtain data, which can be used for such
follow-up projects:

Figure 10A: The first example demonstrates how to retrieve data for the genus *Picea*
across North America by selecting both contemporary and fossil pollen plot datasets,
filtering samples by geographic boundaries and temporal range (0 to 15,000 cal yr BP),
and harmonizing taxa to the genus level. The resulting dataset allows users to study
spatiotemporal patterns of *Picea* distribution over millennia.

Figure 10B: In the second example, the project aims to do species distribution modelling
for plant taxa in the Czech Republic based on contemporary vegetation plot data and
mean annual temperature. The project includes selecting datasets and extracting
relevant abiotic data.

Figure 10C: The third example focuses on obtaining data to be able to reconstruct plant
height for South and Central America between 6-12 cal kyr BP (thousand calibrated
years before present). This example project showcases the integration of trait data with
paleo-vegetation records to subsequently study historical vegetation dynamics and
functional composition of plant communities.

Data Availability

The VegVault v1.0.0 relational database (SQLite; ~110 GB) is publicly archived in the National Repository ('REPO') and available at https://doi.org/10.48700/datst.t21cz-2nq07. The deposit consists of a single SQLite file comprising 31 linked tables with 87 fields, containing >480,000 datasets, >13,000,000 samples, >100,000 taxa, six vegetation trait domains, >11,000,000 trait values, and eight abiotic variables. The database schema and field definitions are detailed in the manuscript (**Data Records**, **Figure 4**; **Table 5-35**). The database can be queried with any SQLite-compatible client or accessed programmatically via the companion RR package {vaultkeepr} (see **Usage Notes**). Version history and change logs are stored in the internal *version_control* table, and subsequent releases will be issued as versioned updates in the same repository.

Code Availability

The VegVault database is built on SQLite, chosen for its robustness and compatibility with various data types. The VegVault v1.0.0 is available at the *National Repository*[44] ('*REPO*') under DOI 10.48700/datst.t21cz-2nq07. The database itself is under *CC B 4.0* licence, see **Usage Notes** section.

VegVault utilizes several tools and software to facilitate data management and access. All code used to generate the VegVault dataset is publicly available under the MIT licence. The GitHub repository *VegVault* used to integrate all data together and migrate them into SQLite database (can be accessed as DOI: 10.5281/zenodo.16987345). See **Methods** section and **Figure 1** for details about individual GitHub repositories.

To enable easy access and manipulation of data, {*vaultkeepr*} R package has been developed as an open access software under MIT licence (https://bit.ly/vaultkeepr).

The code to create the figures in this manuscript is available under DOI:10.5281/zenodo.16987345.

Acknowledgements

OM is funded by the *Czech Science Foundation PIF* grant (GN23-06386I), by the *Charles University Research Centre program* (UNCE/24/SCI/006), and by the *Institutional Support for

662   *Science and Research* of the Ministry of Education, Youth and Sports of the Czech Republic.

663   SGAF is funded by the *Trond Mohn Research Foundation* (TMF) and the University of Bergen

664   by starting grant no. TMS2022STG03).

665   Author contributions

OM conceived the study and programmed all related code. SGAF provided strategic guidance

during the development of the project and manuscript, including decisions on data presentation

and structure. FG, IS, and SGAF provided critical feedback on the technical infrastructure and

database design. OM and SGAF wrote the first draft of the manuscript. SGAF, FG, and IS

contributed to multiple rounds of revision. All authors approved the final version of the

manuscript.

672   Competing interests

673   The author(s) declare no competing interests.

Figures

Figure 1. Schematic workflow of accessing and processing all primary data sources into VegVault v1.0.0 database. 1) *Neotoma Paleoecological Database*[7]; **2)** *sPlotOpen*[15]; **3)** *Botanical Information and Ecology Network*[6]; **4)** *TRY Plant Trait Database*[8]; 5) *CHELSA*[9] and *CHELSA-TRACE21*[18,19]; **6)** *World Soil Information Service*[17]; **7***) FOSSILPOL: Workflow for processing and standardizing global paleoecological pollen data*. A project-specific version with all the settings used for VegVault is stored as GitHub repository *VegVault-FOSSILPOL* (can be accessed as DOI: 10.5281/zenodo.14745060); **8)** The GitHub repository *VegVault-Vegetation_data* used to processes data for contemporary vegetation plots (can be accessed as DOI: 10.5281/zenodo.16729978); **9)** A GitHub repository *VegVault-Trait_data* used to processes data for plant vegetation traits (can be accessed as DOI: 10.5281/zenodo.16729756); **10)** The GitHub repository *VegVault-abiotic_data* used to process climate and soil abiotic data (can be accessed as DOI: 10.5281/zenodo.14745293); **11)** The GitHub repository *VegVault* used to integrate all data together and migrate them into SQLite database (can be accessed as DOI: 10.5281/zenodo.16987345). Additional steps such as the creation of *gridpoints*, grouping of *Traits* into *Trait Domains,* and classification of taxa are also present (for more details see **Methods**); **12)** VegVault - SQLite database (v1.0.0; DOI: 10.48700/datst.t21cz-2nq07); For details about primary data sources, see **Table 1**.

Figure 2. **Example of the resolution of the *gridpoint Datasets* extracting abiotic**
information. A: Map of Europe. The red line indicates country borders with the Czech Republic
as the focus of the map. **B**: Distribution of the individual *gridpoints* (grey squares) within the
Czech Republic (red). **C**: Linking vegetation plots and *gridpoints*. The green circles represent
the contemporary vegetation plots. The grey lines connect the vegetation points to their closest
gridpoint. Note that the abiotic data is provided to each vegetation plot from their closest
gridpoint (i.e., the vegetation plots connected to the same gridpoint will have the same abiotic
data). *Gridpoints* without vegetation plots in their spatio-temporal vicinity (here contemporary
data) were excluded from the VegVault to reduce the database size. See **Methods** section for
more details.

Figure 3. **Spatio-temporal distribution of vegetation Samples** (for both vegetation plots and

fossil pollen archives). Note that this is an overview of the current state of VegVault (v1.0.0.). **A**:

Spatial distribution with each point representing one plot. **B**: Temporal distribution of individual

Samples.

Figure 4: Visualisation of database structure (schema) created using https://dbdiagram.io/.

Each table contains its name, the names of individual columns, and the data type of each

column. In addition, primary key columns are indicated using a small key icon. Individual lines

indicate the connections

Figure 5. **Visualisation of the number of *Datasets*.** **A**: Waffle plot coloured by *Dataset Type*

with each square representing ~1000 *Datasets*. **B**: Number of *Datasets* by *Dataset Source-Type*

grouped by *Dataset Type.* See **Internal Dataset Structure** section for more details.

Figure 6. **Distribution of *Samples* by vegetation plot sizes.** In VegVault, only *vegetation_plot*

Samples and *Datasets* have such information. See **Internal Dataset Structure** section for more

details.

Figure 7: **Example of age uncertainty of *Samples* for selected fossil pollen *Dataset* (ID:**
91256). Each point represents one *Sample*. Each horizontal line represents a range of all
potential ages associated with that *Sample*. Potential ages are gathered from age-depth models
and stored in the *SampleUncertainty* table. See **Internal Dataset Structure** section for more
details about the data structure and *FOSSILPOL* workflow[16] for more details about age
uncertainty from age-depth models.

Figure 8. **Total number of *Taxa* and their classification present in VegVault**. See **Methods**

section for more information about the classification process.

Figure 9. **Schematic workflow of accessing and extracting data from VegVault database using {vaultkeepr} R package**. **1)** Download VegVault database (v1.0.0; DOI: 10.48700/datst.t21cz-2nq07); **2)** Download {vaultkeepr} R package from GitHub (current version 0.0.6; DOI: 10.5281/zenodo.14964737); **3)** In the R programming environment[48], the user has to select which type(s) of datasets they would like to extracted. A minimum of one *Dataset Type (see* **Data Records** section for more details*)* must be specified when using the `vaultkeepr::select_dataset_by_type()` function; **4)** When the user is interested in data from a specific region, they can specific the geographical coordinates of a rectangular extent using the `vaultkeepr::select_dataset_by_geo()` function to only get *Samples* from the individual *Datasets* that were recorded within the region of interest. Additionally, the user can specify the temporal focus by filtering data within a specific period using the `vaultkeepr::select_samples_by_age()`; **5)** The user may also specify any further attributes to be added to the data compilation. Specifically: `*get_taxa()*`—When extracting contemporary or past vegetation (fossil pollen) records, the user most likely wishes to add information about the abundance of individual *Taxa* in each *Sample*. To do so, the user can use the `vaultkeepr::get_taxa()` function. In addition, the user can standardise the taxonomy, so that the extracted *Taxa* can be compared. The parameter `*classify_to*` within the `vaultkeepr::get_taxa()` function allows the user to specify a taxonomic level (species, genus or family) on which the data should be comparable (see **Methods** about *Taxa* classification). Furthermore, the user can select specific *Taxa* based on taxonomy by using the `vaultkeepr::select_taxa_by_name()` function. `*get_traits()*`—When wishing to link the *Trait* data with the vegetation records, the user can use `vaultkeepr::get_traits()` function to extract all *Trait Samples* of the earlier specified spatio-temporal extent. Moreover, similarly to the `vaultkeepr::get_taxa()` function, the user can specify the taxonomic level to which the data should be standardised by using the `*classify_to*` parameter. Please note that the user has to decide in their further analysis how they want to aggregate the measured traits per taxonomic group, e.g. by taking the mean. Further, the user can select a specific *Trait Domain* (of the six available; see **Methods** about details) by using the `vaultkeepr::select_traits_by_domain_name()` function. `*get_abiotic_data()*`—When wishing to link vegetation or trait records with abiotic data, the abiotic data can be obtained by the `vaultkeepr::get_abiotic_data()` function. The user can specify in which mode the abiotic data should be linked, which can be either "*nearest*" (i.e., geographically closest records), "*mean*" or "*median*" (summarising all abiotic records within a set geographical and/or temporal distance). This can be further tweaked by its parameters `*limit_by_distance_km*` and `*limit_by_age_years*`. Furthermore, specific abiotic variables can be chosen by the

`*vaultkeepr::select_abiotic_var_by_name()*` function; **6)** When defined all specifications for data

extraction, the user can execute the extraction using `*vaultkeepr::extract_data()*`. This will result

in a "ready-for-analyses" data compilation. Moreover, the user can use the

`*vaultkeepr::get_references()*` function to obtain all references required (and/or suggested) for

such compilation (see **Internal Dataset Structure** section for more details about the

hierarchical structure of references). Finally, see the **Usage Notes** section for examples of data

processing and extracting using {vaultkeepr} R package.

Figure 10. **Visualisation of example data usage of VegVault using {*vaultkeepr*} package**.

See **Usage Notes** section for details about the aim of each data compilation. **A**: The individual

points represent vegetation data (*Samples* for fossil pollen or contemporary vegetation plots)

within each age time slice. **B**: Each point represents one contemporary vegetation plot. The

colour represents the Mean Annual Temperature (MAT) associated with that plot. Note that

several vegetation plots can have the same value due to sharing the same closest *gridpoint*

(see **Figure 2**). **C**: The left figure represents the temporal length of fossil pollen *Datasets*, which

are available for Central and South America between 6 and 12 ka BP (thousand years before

present). The middle figure represents the distribution of various genera present in this data

assembly. The right figure represents the distribution of average *Plant height Trait* per each

genus present in the data assembly. The code to reproduce these figures can be accessed from

the GitHub repository titled *VegVault* (https://github.com/OndrejMottl/VegVault/tree/v1.0.0-

website_rewiev), which can be accessed as DOI: 10.5281/zenodo.16987345.

Tables

Table 1: Overview comparison of selected databases and VegVault to highlight the latter's
novel contributions. Note that this comparison does not intend to be a comprehensive overview
of all ecological databases. We specify the presence of "Plot-based" vegetation data and
Species lists to flag the differences between stored data. The "Plot-based" databases contain a
full survey of the vegetation at specific locations and, therefore, contain true negatives.
Moreover, "Plot-based" databases contain some expression of species abundance.

Database name	Contains contemporary vegetation data?	Contains paleoecological data?	Contains abundance information?	Contains link to abiotic data?	Contains link to functional traits?
VegVault	⬜(Plot-based)	⬜(Plot-based)	⬜	⬜	⬜
BioDeepTime[13]	⬜(Plot-based; limited)	⬜(Plot-based)	⬜	⬜	⬜
GIFT[14]	⬜(Species lists)	⬜	⬜(Species lists)	⬜	⬜
Botanical Information and Ecology Network (BIEN)[6]	⬜(Plot-based)	⬜	⬜	⬜	⬜
sPlotOpen[15]	⬜(Plot-based)	⬜	⬜	⬜	⬜
Global Biodiversity Information Facility (GBIF)[4]	⬜(Occurrences)	⬜	⬜(Occurrences)	⬜	⬜
Neotoma Paleoecology	⬜	⬜(Plot-based)	⬜	⬜	⬜

Database (Neotoma)[7]					
TRY Plant Trait Database (TRY)[8]	▯	▯	▯	▯	▯

795 **Table 2**: Main sources of data presented in VegVault v1.0.0.

Source	Type	Time period	Comment	References
Botanical Information and Ecology Network (*BIEN*)	Vegetation	contemporary data	Plot-based vegetation data	Botanical Information and Ecology Network (*http://bien.nceas.ucsb.edu/bien/*)[6]
sPlotOpen	Vegetation	contemporary data	The open-access version of *sPlot*[5]	Sabatini et al. (2021)[15]
Neotoma Paleoecological database and *FOSSILPOL*: Workflow for processing and standardizing global paleoecological pollen data	Vegetation	paleo data	We specifically state both FOSSILPOL and Neotoma Paleoecological database, which is the primary source of the data, as FOSSILPOL not only accesses the data but also applies several selection criteria and adds new information (e.g., new age-depth models)	Flantua et al. (2023)[16] Williams, J. W. et al. (2018)[7]
TRY Plant Trait Database (*TRY*)	Functional traits of vegetation	contemporary data		Kattge et al. (2020)[8]
CHELSA	Abiotic data – climate	contemporary data	global downscaled climate data at 1km resolution	Karger et al. (2017)[9]
CHELSA-Trace21k	Abiotic data – climate	paleo data	global downscaled climate data at 1km resolution since the Last Glacial Maximum (21,000 yr BP)	Karger et al. (2020)[18] Karger et al. (2023)[19]
World Soil Information Service (*WoSIS*)	Abiotic data – soil types	contemporary data	global downscaled climate data at 1km resolution	Batjes et al. (2024)[17]

797 **Table 3:** Selected *Traits* presented in the VegVault v1.0.0. The *Traits* were selected based on
798 similarity to Diaz et al. (2016)[36] and grouped into six main *Trait Domains* to simplify the
799 navigation between different data sources.

Trait Domain	Trait name	Trait units	Source of Datasets	Number of Datasets
Stem specific density	*Stem specific density (SSD, stem dry mass per stem fresh volume) or wood density*	g/cm^3	*TRY*	8058
Stem specific density	*stem wood density*	g/cm^3	*BIEN*	348
Leaf nitrogen content per unit mass	*Leaf nitrogen (N) content per leaf dry mass*	mg/g	*TRY*	11464
Leaf nitrogen content per unit mass	*leaf nitrogen content per leaf dry mass*	mg/g	*BIEN*	707
Diaspore mass	*Seed dry mass*	mg	*TRY*	1635
Diaspore mass	*Seed mass*	mg	*BIEN*	48
Plant height	*Plant height vegetative*	m	*TRY*	11476
Plant height	*whole plant height*	m	*BIEN*	193492
Leaf area	*Leaf area (in case of compound leaves: leaf, petiole excluded)*	mm^2	*TRY*	1415
Leaf area	*Leaf area (in case of compound leaves: leaf, petiole included)*	mm^2	*TRY*	1374
Leaf area	*Leaf area (in case of compound leaves: leaf, undefined if petiole in- or excluded)*	mm^2	*TRY*	897
Leaf area	*Leaf area (in case of compound leaves undefined if leaf or leaflet, undefined if petiole is in- or excluded)*	mm^2	*TRY*	2202
Leaf area	*Leaf area*	mm^2	*BIEN*	902

Leaf mass per area	*Leaf area per leaf dry mass (specific leaf area, SLA or 1/LMA): petiole included*	mg/mm^2	*TRY*	1935
Leaf mass per area	*Leaf area per leaf dry mass (specific leaf area, SLA or 1/LMA): undefined if petiole is in- or excluded)*	mg/mm^2	*TRY*	8777
Leaf mass per area	*Leaf mass per area*	mg/mm^2	*BIEN*	835

801   **Table 4**: Overview of abiotic variables present in VegVault v1.0.0.

Abiotic variable name	Abiotic variable description	Abiotic variable unit	Time period	Source of data	Original spatial resolution	Original temporal resolution
bio1	Annual Mean Temperature	°C	contemporary and paleo data	*CHELSA & CHELSA-TRACE21*	*1 km (30 arc-second)*	*100 years for paleo records*
bio4	Temperature Seasonality	°C	contemporary and paleo data	*CHELSA & CHELSA-TRACE21*	*1 km (30 arc-second)*	*100 years for paleo records*
bio6	Min Temperature of Coldest Month	°C	contemporary and paleo data	*CHELSA & CHELSA-TRACE21*	*1 km (30 arc-second)*	*100 years for paleo records*
bio12	Annual Precipitation	kg m-2 year-1	contemporary and paleo data	*CHELSA & CHELSA-TRACE21*	*1 km (30 arc-second)*	*100 years for paleo records*
bio15	Precipitation Seasonality	Unitless	contemporary and paleo data	*CHELSA & CHELSA-TRACE21*	*1 km (30 arc-second)*	*100 years for paleo records*
bio18	Precipitation of Warmest Quarter	kg m-2 quarter-1	contemporary and paleo data	*CHELSA & CHELSA-TRACE21*	*1 km (30 arc-second)*	*100 years for paleo records*
bio19	Precipitation of Coldest Quarter	kg m-2 quarter-1	contemporary and paleo data	*CHELSA & CHELSA-TRACE21*	*1 km (30 arc-second)*	*100 years for paleo records*
HWSD2	Soil type	Unitless	contemporary data	*WoSIS*	*1 km (30 arc-second)*	*N/A*

803   **Table 5**: Column names and types for table *Authors*.

Column name	Data type	Description
author_id	integer	ID of an Author (unique)
author_fullname	character	Full name of an *Author*
author_email	character	Contact email of an *Author*
author_orcid	character	ORCID ID of an *Author*

805   **Table 6**: Column names and types for table *version_control.*

Column name	Data type	Description
id	integer	ID a database version (unique)
version	character	Version number
update_date	character	Date of the creation of that version
changelog	character	Text description of main changes in the database

807   **Table 7**: Column names and types for table *sqlite_stat1.*

Column name	Data type	Description
tbl	character	Name of a table
idx	character	Name of a database index
stat	character	Statistical information. Specifically, comma-separated list of numbers.

809 **Table 8**: Column names and types for table *sqlite_stat4.*

Column name	Data type	Description
tbl	character	Name of a table
idx	character	Name of a database index
neq	character	Number of entries in the index equal to the sample key
nlt	character	Number of entries less than the sample key
ndlt	character	Number of distinct values less than the sample key
sample	character	A sample value from the index (binary-encoded key values)

811   **Table 9**: Column names and types for table *Datasets*.

Column name	Data type	Description
dataset_id	integer	ID of a *Dataset* (unique)
dataset_name	character	Name of each *Dataset*
dataset_type_id	integer	ID of a *Dataset Type*
data_source_id	integer	ID of a Dataset Source
data_source_type_id	integer	ID of a Dataset Source-Type
coord_long	numeric	Geographical coordinates – longitude
coord_lat	numeric	Geographical coordinates – latitude
sampling_method_id	integer	ID of a *Sampling Method*

813 **Table 10**: Column names and types for table *DatasetTypeID.*

Column name	Data type	Description
dataset_type_id	integer	ID of a *Dataset Type* (unique)
dataset_type	character	Text description of individual *Dataset Types* (currently *vegetation_plot, fossil_pollen_archive, traits, gridpoints*)

815   **Table 11**: Column names and types for table *DatasetSourceTypeID*.

Column name	Data type	Description
data_source_type_id	integer	ID of a *Dataset Source-Type* (unique)
dataset_source_type	character	Text description of individual *Dataset Source-Type* (currently, *BIEN*, *sPlotOpen*, *TRY*, *Neotoma-FOSSILPOL*, *gridpoints*)

817   **Table 12**: Column names and types for table *DatasetSourceTypeReference*.

Column name	Data type	Description
data_source_type_id	integer	ID of a *Dataset Source-Type*
reference_id	integer	ID of a *Reference*

819   **Table 13**: Column names and types for table *DatasetSourcesID.*

Column name	Data type	Description
data_source_id	integer	ID of a *Dataset Source* (unique)
data_source_desc	character	Text description of individual *Dataset Sources* (e.g., name of the sub-database from the primary source)

821 **Table 14**: Column names and types for table *DatasetSourcesReference*.

Column name	Data type	Description
data_source_id	integer	ID of a *Dataset Source*
reference_id	integer	ID of a *Reference*

823   **Table 15**: Column names and types for table *SamplingMethodID.*

Column name	Data type	Description
sampling_method_id	integer	ID of a *Dataset Sampling Method* (unique)
sampling_method_details	character	Text description of individual *Dataset Sampling Method* (e.g., name of the sub-database from the primary source)

825 **Table 16**: Column names and types for table *SamplingMethodReference.*

Column name	Data type	Description
sampling_method_id	integer	ID of a *Dataset Sampling Method*
reference_id	integer	ID of a *Reference*

827 **Table 17**: Column names and types for table *DatasetReferences*.

Column name	Data type	Description
dataset_id	integer	ID of a *Dataset*
reference_id	integer	ID of a *Reference*

829 **Table 18**: Column names and types for table *Samples.*

Column name	Data type	Description
sample_id	integer	ID of a *Sample* (unique)
sample_name	character	Name of a Sample
sample_details	character	Specific description of a *Sample.* Currently not being used.
age	numeric	Age of sample. Mainly used for *fossil_pollen_archives*, where note the age of a *Sample* in calibrated years before present. Note that all contemporary *Samples*, have age of 0.
sample_size_id	integer	ID of a *Sample Size*

831 **Table 19**: Column names and types for table *DatasetSample.*

Column name	Data type	Description
dataset_id	integer	ID of a *Dataset*
sample_id	integer	ID of a *Sample*

833   **Table 20**: Column names and types for table *SampleSizeID*.

Column name	Data type	Description
sample_size_id	integer	ID of a size category (unique)
sample_size	numeric	Numeric expression of size
description	character	Mostly description of units in which the values are stored

835   **Table 21**: Column names and types for table *SampleUncertainty.*

Column name	Data type	Description
sample_id	integer	ID of a *Sample*
iteration	integer	ID of a iteration from age depth model. Currently, the is 1000 iteration per each Sample.
age	integer	Potential age of a Sample

837 **Table 22**: Column names and types for table *SampleReference*.

Column name	Data type	Description
sample_id	integer	ID of a *Sample*
reference_id	integer	ID of a *Reference*

839 **Table 23**: Column names and types for table *Taxa.*

Column name	Data type	Description
taxon_id	integer	ID of a *Taxon* (unique)
taxon_name	character	Name of a *Taxon* from primary source.

841 **Table 24**: Column names and types for table *SampleTaxa.*

Column name	Data type	Description
sample_id	integer	ID of a *Sample*
taxon_id	integer	ID of a *Taxon*
value	numeric	Abundance representation of a *Taxon* (the units may differ among primary sources, i.e., *Dataset Source-Types*)

843 **Table 25**: Column names and types for table *TaxonClassification.*

Column name	Data type	Description
taxon_id	integer	ID of a *Taxon* (unique)
taxon_species	integer	ID of a *Taxon,* which was assigned as species level
taxon_genus	integer	ID of a *Taxon,* which was assigned as genus level
taxon_family	integer	ID of a *Taxon,* which was assigned as family level

845 **Table 26**: Column names and types for table *TaxonReference.*

Column name	Data type	Description
taxon_id	integer	ID of a *Taxon*
reference_id	integer	ID of a Reference

847 **Table 27**: Column names and types for table *Traits.*

Column name	Data type	Description
trait_id	integer	ID of a *Trait* (unique)
trait_domain_id	integer	ID of a *Trait Domain*
trait_name	character	Name of the trait from the primary source. See **Methods** for the details about the specific columns used from primary sources.

849   **Table 28**: Column names and types for table *TraitsDomain.*

Column name	Data type	Description
trait_domain_id	integer	ID of a *Trait Domain* (unique)
trait_domain_name	character	Name of the *Trait Domain* from Diaz et al. (2016)[36]
trait_domanin_description	character	Additional information about the *Trait Domain*

851 **Table 29**: Column names and types for table *TraitsValue.*

Column name	Data type	Description
trait_id	integer	ID of a *Trait*
dataset_id	integer	ID of a *Dataset*
sample_id	integer	ID of a *Sample*
taxon_id	integer	ID of a T*axon*
trait_value	numeric	Value of specific measured observation of *Trait*. See **Methods** for the details about the specific columns used from primary sources.

853   **Table 30**: Column names and types for table *TraitsReference.*

Column name	Data type	Description
trait_id	integer	ID of a *Trait*
reference_id	integer	ID of a *Reference*

855   **Table 31**: Column names and types for table *AbioticVariable.*

Column name	Data type	Description
abiotic_variable_id	integer	ID of a *Abiotic Variable*
abiotic_variable_name	character	Name of a *Abiotic Variable* from primary source
abiotic_variable_unit	character	Unit of a *Abiotic Variable*
measure_details	character	Additonal details about *Abiotic Variable*

857   **Table 32**: Column names and types for table *AbioticData.*

Column name	Data type	Description
sample_id	integer	ID of a *Sample*
abiotic_variable_id	integer	ID of an *Abiotic Variable*
abiotic_value	numeric	The actual value of abiotic data

859 **Table 33**: Column names and types for table *AbioticVariableReference.*

Column name	Data type	Description
abiotic_variable_id	integer	ID of an *Abiotic Variable*
reference_id	integer	ID of a *Reference*

861   **Table 34**: Column names and types for table *AbioticDataReference.*

Column name	Data type	Description
sample_id	integer	ID of non-*gridpoint Sample*
sample_ref_id	integer	ID of *gridpoint Sample*
distance_in_km	integer	Distance among samples expressed in kilometres
distance_in_years	integer	Distance among samples expressed in years

863   **Table 35**: Column names and types for table *References.*

Column name	Data type	Description
reference_id	integer	ID of a *Reference* (unique)
reference_detail	character	Text of the reference (i.e., citation)
mandatory	Integer (Boolean)	Is the Reference mandatory to report while using the data? Most primary sources of the data have a license, which requires correct attribution.

References

1. Fordham, D. A. *et al.* Using paleo-archives to safeguard biodiversity under climate change.

Science **369**, eabc5654 (2020) https://doi.org/10.1126/science.abc5654.

2. Nieto-Lugilde, D. *et al.* Time to better integrate paleoecological research infrastructures with

neoecology to improve understanding of biodiversity long-term dynamics and to inform

future conservation. *Environ. Res. Lett.* **16**, 095005 (2021) https://doi.org/10.1088/1748-

9326/ac1b59.

3. Reitalu, T., Kuneš, P. & Giesecke, T. Closing the gap between plant ecology and

Quaternary palaeoecology. *Journal of Vegetation Science* **25**, 1188–1194 (2014)

https://doi.org/10.1111/jvs.12187.

4. GBIF.org. GBIF Home Page. https://www.gbif.org (2024) https://www.gbif.org.

5. Bruelheide, H. *et al.* sPlot – A new tool for global vegetation analyses. *Journal of Vegetation*

Science **30**, 161–186 (2019) https://doi.org/10.1111/jvs.12710.

6. Botanical Information and Ecology Network (BIEN). BIEN: Botanical Information and

Ecology Network. http://bien.nceas.ucsb.edu/bien/ (n.d.) http://bien.nceas.ucsb.edu/bien/.

7. Williams, J. W. *et al.* The Neotoma Paleoecology Database, a multiproxy, international,

community-curated data resource. *Quat. res.* **89**, 156–177 (2018)

https://doi.org/10.1017/qua.2017.105.

8. Kattge, J. *et al.* TRY plant trait database – enhanced coverage and open access. *Global*

Change Biology **26**, 119–188 (2020) https://doi.org/10.1111/gcb.14904.

9. Karger, D. N. *et al.* Climatologies at high resolution for the earth's land surface areas. *Sci*

Data **4**, 170122 (2017) https://doi.org/10.1038/sdata.2017.122.

10. Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces for

global land areas. *International Journal of Climatology* **37**, 4302–4315 (2017)

https://doi.org/10.1002/joc.5086.

11. Schipper, E. L. F., Dubash, N. K. & Mulugetta, Y. Climate change research and the search
for solutions: rethinking interdisciplinarity. *Clim Change* **168**, 18 (2021)
https://doi.org/10.1007/s10584-021-03237-3.

12. Intergovernmental Panel On Climate Change. *Climate Change and Land: IPCC Special*
Report on Climate Change, Desertification, Land Degradation, Sustainable Land
Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems.
(Cambridge University Press, 2022). https://doi.org/10.1017/9781009157988
https://www.cambridge.org/core/product/identifier/9781009157988/type/book.

13. Smith, J. *et al.* BioDeepTime: A database of biodiversity time series for modern and fossil
assemblages. *Global Ecology and Biogeography* **32**, 1680–1689 (2023)
https://doi.org/10.1111/geb.13735.

14. Weigelt, P., König, C. & Kreft, H. GIFT – A Global Inventory of Floras and Traits for
macroecology and biogeography. *Journal of Biogeography* **47**, 16–43 (2020)
https://doi.org/10.1111/jbi.13623.

15. Sabatini, F. M. *et al.* sPlotOpen – An environmentally balanced, open-access, global
dataset of vegetation plots. *Global Ecology and Biogeography* **30**, 1740–1764 (2021)
https://doi.org/10.1111/geb.13346.

16. Flantua, S. G. A. *et al.* A guide to the processing and standardization of global
palaeoecological data for large-scale syntheses using fossil pollen. *Global Ecology and*
Biogeography **32**, 1377–1394 (2023) https://doi.org/10.1111/geb.13693.

17. Batjes, N. H., Calisto, L. & de Sousa, L. M. Providing quality-assessed and standardised soil
data to support global mapping and modelling (WoSIS snapshot 2023). *Earth System*
Science Data **16**, 4735–4765 (2024) https://doi.org/10.5194/essd-16-4735-2024.

18. Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H. & Zimmermann, N. E. CHELSA-
TraCE21k: Downscaled transient temperature and precipitation data since the last glacial

maximum.   (2020)   http://dx.doi.org/10.16904/envidat.211

https://www.envidat.ch/dataset/chelsa_trace.

19. Karger, D. N., Nobis, M. P., Normand, S., Graham, C. H. & Zimmermann, N. E. CHELSA-

TraCE21k – high-resolution (1km) downscaled transient temperature and precipitation data

since the Last Glacial Maximum. *Climate of the Past* **19**, 439–456 (2023)

https://doi.org/10.5194/cp-19-439-2023.

20. GBIF Secretariat. GBIF Backbone Taxonomy. (2023).

21. Mottl, O. taxospace - v0.0.0.9003. https://doi.org/10.5281/zenodo.14699491 (2024)

https://doi.org/10.5281/zenodo.14699491 https://github.com/OndrejMottl/taxospace.

22. Mottl, O. vaultkeepr - v0.0.6. https://doi.org/10.5281/zenodo.14964737 (2025)

https://doi.org/10.5281/zenodo.14964737 https://github.com/OndrejMottl/vaultkeepr.

23. Mottl, O. *et al.* Global acceleration in rates of vegetation change over the past 18,000 years.

Science **372**, 860–864 (2021) https://doi.org/10.1126/science.abg1685.

24. Šímová, I., Ordonez, A. & Storch, D. The dynamics of the diversity–energy relationship

during the last 21,000 years. *Global Ecology and Biogeography* **32**, 707–718 (2023)

https://doi.org/10.1111/geb.13649.

25. Brown, K. A. *et al.* Trait-based approaches as ecological time machines: Developing tools

for reconstructing long-term variation in ecosystems. *Functional Ecology* **37**, 2552–2569

(2023) https://doi.org/10.1111/1365-2435.14415.

26. Li, J. & Prentice, I. C. Global patterns of plant functional traits and their relationships to

climate. *Commun Biol* **7**, 1–14 (2024) https://doi.org/10.1038/s42003-024-06777-3.

27. Shipley, B., Dilkina, B. & McGuire, J. MegaSDM: Modelling Species Ranges in The Past

And Future. *Bulletin of the Florida Museum of Natural History* **60**, 115–115 (2023)

https://doi.org/10.58782/flmnh.zwwl8127.

28. Pollock, L. J. *et al.* Understanding co-occurrence by modelling species simultaneously with

a Joint Species Distribution Model (JSDM). *Methods in Ecology and Evolution* **5**, 397–406

(2014) https://doi.org/10.1111/2041-210X.12180.

29. Ordonez, A. & Gill, J. L. Unravelling the functional and phylogenetic dimensions of novel

ecosystem assemblages. *Philosophical Transactions of the Royal Society B: Biological*

Sciences **379**, 20230324 (2024) https://doi.org/10.1098/rstb.2023.0324.

30. Wieczynski, D. J. *et al.* Climate shapes and shifts functional biodiversity in forests worldwide.

Proceedings of the National Academy of Sciences **116**, 587–592 (2019)

https://doi.org/10.1073/pnas.1813723116.

31. Zurell, D., Fritz, S. A., Rönnfeldt, A. & Steinbauer, M. J. Predicting extinctions with species

distribution models. *Cambridge Prisms: Extinction* **1**, e8 (2023)

https://doi.org/10.1017/ext.2023.5.

32. Chacon, S. & Straub, B. *Pro Git.* (Springer Nature, 2014).

33. Haslett, J. & Parnell, A. A Simple Monotone Process with Application to Radiocarbon-Dated

Depth Chronologies. *Journal of the Royal Statistical Society Series C: Applied Statistics* **57**,

399–418 (2008) https://doi.org/10.1111/j.1467-9876.2008.00623.x.

34. Maitner, B. S. *et al.* The bien r package: A tool to access the Botanical Information and

Ecology Network (BIEN) database. *Methods in Ecology and Evolution* **9**, 373–379 (2018)

https://doi.org/10.1111/2041-210X.12861.

35. Sabatini, F. M., Lenoir, J., Bruelheide, H. & sPlot Consortium. sPlotOpen – An

environmentally-balanced, open-access, global dataset of vegetation plots. iDiv Data

Repository   https://doi.org/10.25829/idiv.3474-bb7k72   (2021)

https://doi.org/10.25829/idiv.3474-bb7k72.

36. Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167–171 (2016)

https://doi.org/10.1038/nature16489.

37. Lam, O. H. Y. *et al.* 'rtry': An R package to support plant trait data preprocessing. *Ecology*

and Evolution **14**, e11292 (2024) https://doi.org/10.1002/ece3.11292.

38. Jentsch, helge, Weidinger, J. & Bobrowski, M. ClimDatDownloadR: Downloads Climate

Data from Chelsa and WorldClim. https://doi.org/10.5281/zenodo.7924343 (2025)

https://doi.org/10.5281/zenodo.7924343

https://github.com/HelgeJentsch/ClimDatDownloadR.

39. Hijmans, R. J. *Terra: Spatial Data Analysis*. (2025). https://rspatial.org/.

40. Thuiller, W. On the importance of edaphic variables to predict plant species distributions –

limits and prospects. *Journal of Vegetation Science* **24**, 591–592 (2013)

https://doi.org/10.1111/jvs.12076.

41. Zuquim, G., Costa, F. R. C., Tuomisto, H., Moulatlet, G. M. & Figueiredo, F. O. G. The

importance of soils in predicting the future of plant habitat suitability in a tropical forest. *Plant*

Soil **450**, 151–170 (2020) https://doi.org/10.1007/s11104-018-03915-9.

42. Velazco, S. J. E., Galvão, F., Villalobos, F. & Júnior, P. D. M. Using worldwide edaphic data

to model plant species niches: An assessment at a continental extent. *PLOS ONE* **12**,

e0186025 (2017) https://doi.org/10.1371/journal.pone.0186025.

43. Global Names Architecture. Global Names Resolver. https://resolver.globalnames.org/

https://resolver.globalnames.org/.

44. CESNET. National Repository. https://data.narodni-repozitar.cz/ (2024) https://data.narodni-

repozitar.cz/.

45. Dillon, E. M. *et al.* Challenges and directions in analytical paleobiology. *Paleobiology* **49**,

377–393 (2023) https://doi.org/10.1017/pab.2023.3.

46. Keil, P. *et al.* Patterns of beta diversity in Europe: the role of climate, land cover and

distance across scales. *Journal of Biogeography* **39**, 1473–1486 (2012)

https://doi.org/10.1111/j.1365-2699.2012.02701.x.

47. Augustine, S. P., Bailey-Marren, I., Charton, K. T., Kiel, N. G. & Peyton, M. S. Improper data

practices erode the quality of global ecological databases and impede the progress of

ecological   research.   *Global   Change   Biology*   **30**,   e17116   (2024)

https://doi.org/10.1111/gcb.17116.

48. R Core Team. *R: A Language and Environment for Statistical Computing.* (R Foundation for

Statistical Computing, Vienna, Austria, 2024). https://www.R-project.org/.

995
996

Supplementary Tables

Table S1: A list of selected depositional environments used during the processing of fossil

pollen data.

Depositional Environment	included
Abrocoma Midden	FALSE
Aeolian	FALSE
Alluvial fan	FALSE
Animal Midden	FALSE
Archaeological	FALSE
Archaeological Rock Shelter	FALSE
Arroyo	FALSE
Artificial Lake	FALSE
Ash Fall	FALSE
Backwater Lake	FALSE
Biological	FALSE
Blanket Bog	TRUE
Bog	TRUE
Caldera Lake	TRUE
Canal	FALSE
Cave	FALSE
Cave Dung Deposit	FALSE
Cirque Lake	TRUE
Coastal	FALSE
Coastal Bar	FALSE
Coastal lagoon	FALSE
Colluvium	FALSE
Deep Sea Benthic	FALSE
Deflation Basin Lake	TRUE
Delta	FALSE
Drained Lake	FALSE
Drift Filled Valley Lake	FALSE
Dune Dammed Lake	FALSE
Estuarine	FALSE
Explosion Crater Lake	TRUE
Fen	TRUE
Fjord	FALSE
Floating Mire	TRUE
Flood Scour Lake	TRUE

Floodplain	FALSE
Floodplain Basin	FALSE
Floodplain Mire	FALSE
Fluvial	FALSE
Fluvial Origin Lake	TRUE
Fluvial terrace lake	TRUE
Glacial	FALSE
Glacial Ice	FALSE
Glacial Meltwater Channel Lake	TRUE
Glacial Origin Lake	TRUE
Glacial Outburst Flood Lake	TRUE
Glacial Scour Lake	TRUE
Glacier Ice	FALSE
Glaciomarine	FALSE
Ice Thrust Lake	TRUE
Interdunal Lake	TRUE
Kettle Lake	FALSE
Lacustrine	TRUE
Lacustrine Beach	FALSE
Lake Dammed by Alluvial Fans	FALSE
Lake Marginal Fen	TRUE
Landslide Origin Lake	TRUE
Lava Flow Dammed Lake	TRUE
Loess	FALSE
Mangrove Swamp	TRUE
Marine	FALSE
Marine Beach	FALSE
Marine Wave-cut Platform	FALSE
Marsh	TRUE
Meander Belt	FALSE
Meteorite Impact Crater	TRUE
Mire	TRUE
Moist Soil	FALSE
Mor Humus	FALSE
Moraine Dammed Lake	TRUE
Natural Lake	TRUE
Natural Lake (Origin Unknown)	TRUE
Neotoma Midden	FALSE
Neotoma cinerea Midden	FALSE
Non-system tank	FALSE
Nonspecific Floodplain Lake	FALSE

Occupational Debris	FALSE
Open-water Transition Mire	TRUE
Other	FALSE
Oxbow Lake	FALSE
Paleosol	FALSE
Palustrine	TRUE
Periglacial Origin Lake	TRUE
Permafrost	FALSE
Petromus Midden	FALSE
Phyllotis Midden	FALSE
Playa	FALSE
Pond	TRUE
Poor Fen	FALSE
Procavia Midden	FALSE
Proglacial Lake	FALSE
Proglacial Lake	FALSE
Quarry	FALSE
Raised Bog	TRUE
Reservoir	FALSE
Rich Fen	TRUE
River Channel	FALSE
River Terrace	FALSE
Riverine	FALSE
Rock Shelter	FALSE
Salt Marsh	TRUE
Small Hollow	TRUE
Small Hollow Fen	TRUE
Soil	FALSE
Soligenous Mire	TRUE
Solution Origin Lake	TRUE
Spring	TRUE
Spring Mound	FALSE
Spring Pond	FALSE
Stream	FALSE
Stream-cut Exposure	FALSE
Swamp	TRUE
Tectonic Origin Lake	TRUE
Terrestrial	FALSE
Thaw Slump	FALSE
Thermokarst Lake	TRUE
Tidal Freshwater Forested	TRUE

Wetland	
Tidal Oligohaline Marsh	TRUE
Unknown	FALSE
Valley Glacier Lake	TRUE
Valley Mire	TRUE
Volcanic	FALSE
Volcanic Origin Lake	TRUE
Wind Origin Lake	TRUE
NA	FALSE

1000

1001   **Table S2**: A list of selected ecological groups used during the processing of fossil pollen data.

Ecological group	included
ACAR	FALSE
ACRI	FALSE
ALGA	FALSE
ANAC	FALSE
ANIM	FALSE
AQBR	FALSE
AQVP	FALSE
ARTH	FALSE
BIOM	FALSE
BRYZ	FALSE
CHAR	FALSE
CHRM	FALSE
CLAD	FALSE
COLE	FALSE
COPE	FALSE
DIAT	FALSE
DINO	FALSE
DIPT	FALSE
ELEM	FALSE
EMBR	FALSE
FLWO	FALSE
FORM	FALSE
FUNG	FALSE
GPOD	FALSE
INSE	FALSE
LABO	FALSE
LEPI	FALSE
MANG	TRUE

MNRL	FALSE
NEMA	FALSE
PALM	TRUE
PLNT	TRUE
PLOI	FALSE
PROT	FALSE
PSED	FALSE
PTER	FALSE
ROTI	FALSE
SEED	FALSE
SPNG	FALSE
SUCC	TRUE
TEAM	FALSE
TRSH	TRUE
UNID	FALSE
UPBR	FALSE
UPHE	FALSE
VACR	FALSE
VASC	TRUE
NA	FALSE

1002

1003 **Table S3**: A list of valid control types used during the age-depth modelling process during the
1004 processing of fossil pollen data.

Chronology control type	included	calibrated
Ambrosia rise	FALSE	FALSE
Annual laminations (varves)	TRUE	FALSE
Annual laminations (varves)/Sedimentation rate	TRUE	FALSE
Argon-argon	FALSE	FALSE
Biostratigraphic	FALSE	FALSE
Biostratigraphic, pollen	FALSE	FALSE
Biostratigraphic, pollen, Allerød/Younger Dryas	FALSE	FALSE
Biostratigraphic, pollen, Older Dryas/Allerød	FALSE	FALSE
Biostratigraphic, pollen, Oldest Dryas/Bølling	FALSE	FALSE
Biostratigraphic, pollen, Pleniglacial/Meiendorf	FALSE	FALSE
Biostratigraphic, pollen, Younger Dryas/Holocene	FALSE	FALSE
Biostratigraphic, pollen, regional	FALSE	FALSE

Collection date	TRUE	FALSE
Core bottom	FALSE	FALSE
Core top	TRUE	FALSE
Core top, estimated	TRUE	FALSE
Cultural	FALSE	FALSE
Deglaciation	FALSE	FALSE
European settlement horizon	FALSE	FALSE
Event stratigraphic	FALSE	FALSE
Extrapolated	FALSE	FALSE
Geochronologic	FALSE	FALSE
Geologic time scale	FALSE	FALSE
Geologic time-scale boundary	FALSE	FALSE
Guess	TRUE	FALSE
Heeb & Welten (1972) pollen boundary	FALSE	FALSE
Heinrich stadials	FALSE	FALSE
Historical documentation	TRUE	FALSE
Historical fire	TRUE	FALSE
IDW-2d Picea decline	FALSE	FALSE
IDW-2d Ulmus decline	FALSE	FALSE
Infrared Stimulated Luminescence (IRSL)	FALSE	FALSE
Interpolated	FALSE	FALSE
Introduced Pinus rise	FALSE	FALSE
Isotopic excursion	FALSE	FALSE
Lead-210	TRUE	FALSE
Marine isotope stage boundary	FALSE	FALSE
Marine isotope stage event	FALSE	FALSE
Marine isotope stages	FALSE	FALSE
Optically stimulated luminescence	TRUE	FALSE
Other absolute dating methods	FALSE	FALSE
Other dating methods	FALSE	FALSE
Oxygen-18	TRUE	FALSE
Palaeomagnetic	TRUE	FALSE
Picea decline	FALSE	FALSE
Pre-EuroAmerican settlement horizon	FALSE	FALSE
Quaternary event classification	FALSE	FALSE
Radiocarbon	TRUE	TRUE
Radiocarbon, average of two or more dates	TRUE	TRUE
Radiocarbon, calibrated	TRUE	FALSE
Radiocarbon, calibrated from calendar years	TRUE	FALSE
Radiocarbon, calibrated, Bayesian modelled	TRUE	FALSE
Radiocarbon, infinite	TRUE	FALSE

Radiocarbon, reservoir correction	TRUE	TRUE
Salsola rise	FALSE	FALSE
Section top	TRUE	FALSE
Sediment stratigraphic	FALSE	FALSE
Stratigraphic	FALSE	FALSE
Tephra	TRUE	FALSE
Tsuga decline	FALSE	FALSE
Uranium-series	TRUE	FALSE
W.O. van der Knaap pollen boundary	FALSE	FALSE
Welten (1982) pollen boundary	FALSE	FALSE

1005

one square is 1000 Datasets

1 get the **VegVault** datatabase

DOI: 10.48700/
datst.t21cz-2nq07

2 get the {vaultkeepr} R package

```
>library(remotes)
>install_github(
  "OndrejMottl/vaultkeepr"
)
```

3 Which type of dataset to use?

```
>select_dataset_by_type()
```
vegetation_plot   traits
fossil_pollen_archive gridpoints

4 For which spatio-temporal are you need the data?

```
>select_dataset_by_geo()
```
lat_lim
long_lim

```
>select_samples_by_age()
```
age_lim

5 Which atributes to include?

```
>get_taxa()
```
```
>get_abiotic_data()
```
```
>get_traits()
```

6 Extract data and references

```
>extract_data()
```
```
>get_references()
```