The core MICOS complex subunit Mic60 has been substituted by two cryptic mitofilin-

containing proteins in Euglenozoa

Shaghayegh Sheikh[1,2,†], Barbora Turpin Knotková[3], Corinna Benz[1], Marek Eliáš[4], Tomáš

Bílý[1,2], Alexey Bondar[2,5], Martina Tesařová[1], Eva Stříbrná[1], Jiří Heller[1], Michael Meinecke[3],

Hassan Hashimi[1,2*]

[1]Institute of Parasitology, Biology Centre, Czech Academy of Sciences, České Budějovice,

Czech Republic

[2]University of South Bohemia, Faculty of Science, České Budějovice, Czech Republic

[3]Heidelberg University Biochemistry Center (BZH), Heidelberg, Germany

[4]Department of Biology and Ecology, Faculty of Science, University of Ostrava, Ostrava,

Czech Republic

[5]Institute of Entomology, Biology Centre, Czech Academy of Sciences, České Budějovice,

Czech Republic

*corresponding author: hassan@paru.cas.cz

[†]current address: Department of Medical Biochemistry and Biophysics, Division of Molecular

Metabolism, Karolinska Institutet, Biomedicum, 171 65, Solna, Sweden

ABSTRACT

Cristae enclose respiratory chain complexes, making them the bioenergetic

subcompartments of mitochondria. The MICOS complex is among the inducers of

membrane curvature needed for crista formation. Resembling the respiratory chain

complexes, MICOS is organized around a core protein, the mitofilin-domain bearing Mic60,

that was inherited from the alphaproteobacterial progenitor of mitochondria. Extant

alphaproteobacteria express Mic60 to form their own bioenergetic subcompartments,

demonstrating the permeance of Mic60's form and function during prokaryotic and

eukaryotic evolution. Yet, unlike virtually all aerobic eukaryotes, Mic60 is not encoded within

the genomes of the multifarious protists that comprise the phylum Euglenozoa, including

trypanosomes. Here, we show that Mic60 has been replaced in euglenozoans by two cryptic

mitofilin domain-containing MICOS subunits, Mic34 and Mic40. Contrasting

alphaproteobacterial and mitochondrial Mic60, these are not integral membrane proteins.

Mic34 and Mic40 are as diverged from each other as both are to canonical Mic60. Reverse

genetics revealed they are intertwined with the oxidative protein folding pathway required for

mitochondrial–and crista–biogenesis, veiling a potential membrane remodeling role.

Nevertheless, Mic34 binds phospholipid bilayers *in vitro*. Mic34 and Mic40 heterologous

expression remodels gammaproteobacterial cytoplasmic membranes, like Mic60.

Unexpectedly, Mic34 overexpression elaborates the simplified tubular mitochondrion of a

Trypanosoma brucei life cycle stage with repressed oxidative phosphorylation. Furthermore,

this activity was ablated by mutations to Mic34's mitofilin domain that correspond to essential

motifs found in yeast Mic60's mitofilin domain. Thus, the mitofilin protein family is more

diverse than originally supposed, with two of its structurally most divergent members altering

the core of euglenozoan MICOS.

INTRODUCTION

Cristae are the bioenergetic subcompartments of mitochondria that house the molecular

machinery for cellular respiration (Pánek et al. 2020). Their biogenesis involves the extreme

bending of the mitochondrial inner membrane (IM) by dedicated–and usually multiprotein–

molecular machineries (Barbot and Meinecke 2016; Daumke and van der Laan 2025). Rows

of ATP synthase dimers are responsible for positive curvature at cristae rims (Kühlbrandt

2019) whereas the Mitochondrial contact site and Cristae Organizing System (MICOS)

complex forms both crista junctions, constricted zones of negative curvature (Colina-Tenorio

et al. 2020). Additionally, MICOS forms contact sites between the IM and outer membrane.

While ATP synthase dimerization appears to be a eukaryotic invention (Kühlbrandt 2019),

the keystone MICOS subunit Mic60 is of more ancient origin, being directly inherited from

the alphaproteobacterial symbiont that gave rise to mitochondria (Muñoz-Gómez et al. 2015;

Huynen et al. 2016). Mic60's function in the formation of bioenergetic subcompartments and

double membrane contact sites is conserved in extant alphaproteobacteria (Muñoz-Gómez

et al. 2023), supporting the hypothesis that the cenancestor of mitochondria and

alphaproteobacteria already possessed the requisite machinery to form cristae (Muñoz-

Gomez et al. 2017). Indeed, the shared properties of eukaryotic and prokaryotic Mic60 most

parsimoniously imply that their common ancestor had the capacity to remodel their inner

membranes into crista-like subcompartments (Tarasenko et al. 2017). Also, like the

respiratory chain complexes that are embedded in cristae, MICOS is also assembled around

a proteinaceous core inherited from the alphaproteobacterial symbiont (Roger et al. 2017).

Core architectures are necessarily more conserved than the surrounding shell of often

lineage-specific subunits to stabilize the functionality of these protein complexes

(Prokopchuk et al. 2023).

The canonical domain architecture of Mic60 (Muñoz-Gómez et al. 2015; Huynen et al. 2016)

forms a tertiary structure that allows the protein's homotetramerization, which has been

proposed to tubulate membranes (Bock-Bierbaum et al. 2022). A transmembrane domain

(TMD) embeds its N-proximal segment into the IM. A central coiled coil (CC) domain

provides an interaction interface bringing together two antiparallelly oriented Mic60

homodimers into a tetramer. The C-terminal mitofilin domain, made up of eight alpha helices,

directly interacts with phospholipids on the opposite side of a tubulated membrane. Via this

mechanism, Mic60 has been proposed to be the primary factor in crista junction constriction

(Stephan et al. 2020).

Given its central role in crista biogenesis, Mic60 is ubiquitously encoded in the genomes of

aerobic eukaryotes, with only a few exceptions (Muñoz-Gómez et al. 2015; Huynen et al.

2016). Flagellated protists belonging to the phylum Euglenozoa represent one such

anomaly. Euglenozoans are a sub-clade within the eukaryotic supergroup Discoba (Burki et

al. 2020). This phylum is named after Euglenida, a subdivision containing free-living, mostly

freshwater protists (Fig. 1A). Two sister clades branch next to Euglenida: Diplonemea, a

group of predominantly marine flagellates, and Kinetoplastea (or Kinetoplastida), named

after the kinetoplast, a massive network of mitochondrial DNA that comprises the organelle's

single genome (Kostygov et al. 2021). A species-poor group Symbiontida forms a fourth

euglenozoan lineage (Lax et al. 2021), but it is not considered further in this study due to

limited sequence resources.

Until it was isolated from the kinetoplastid *Trypanosoma brucei* (Kaurov et al. 2018), the

MICOS complex was only known from animals and yeast, grouped together in the diverse

Opisthokonta clade (Burki et al. 2020). *T. brucei* MICOS has some commonalities with

opisthokont MICOS, such as having the other central subunit, Mic10. While *T. brucei* MICOS

differs in having two Mic10 paralogs, one paralog associates with ATP synthase dimers

(Cadena et al. 2021), like in yeast (Rampelt et al. 2022). But *T. brucei* MICOS also has traits

not observed in opisthokont MICOS. One example is that half of the subunits assemble into

a soluble subcomplex within the intermembrane space (IMS; Fig. 2C) (Kaurov et al. 2018;

Eichenberger et al. 2019). Furthermore, it seems to be directly involved in a still unknown

aspect of the Mitochondrial Intermembrane Space Assembly (MIA) pathway (Kaurov et al.

2018; Kaurov et al. 2022), which facilitates the oxidative folding of proteins imported into the

IMS (Mordas and Tokatlidis 2015). Here, we address another trait that sets *T. brucei* MICOS

apart from MICOS in virtually all other eukaryotes. As predicted by the aforementioned

bioinformatic screen, the isolated *T. brucei* MICOS lacked a canonical Mic60. At the time of

discovery, a subunit bearing a superficially similar domain architecture, but lacking the

characteristic mitofilin domain, was annotated as a putative Mic60 (Kaurov et al. 2018).

However, here we present *in silico* and experimental evidence that two other MICOS

subunits named Mic34 (Uniprot ID: Q57ZZ1) and Mic40 (Q57YH2) truly represent cryptic

mitofilin family proteins. Thus, we henceforth will refer to the originally misannotated Mic60

(Q38DZ0) as Mic24. This discovery has expanded the mitofilin proteins to include members

that implement the eponymous domain unconventionally.

RESULTS

The soluble MICOS subunits Mic34 and Mic40 are paralogs that each contain a

mitofilin domain

A *bona fide* ortholog of Mic60 was not identified in the MICOS complex isolated from *T.*

brucei (Kaurov et al. 2018). However, when the largest *T. brucei* Mic34 and Mic40 MICOS

subunits were searched against the Pfam-A (v.37) database using HHpred (Söding et al.

2005), which implements sensitive protein homology detection based on pairwise

comparison of profile hidden Markov models (HMMs), both retrieved the family PF09731.14,

i.e. mitofilin as the best hit with convincing statistical significance: 98.08% probability and E-

value of 0.0056 for Mic34, and 96.77% probability and E-value of 2.2 for Mic40. We then

built a more representative profile HMM based on Mic34 and Mic40 multiple sequence

alignments (MSAs) of a phylogenetically comprehensive set of homologs retrieved from all

major Euglenozoa groups, i.e. Kinetoplastea, Diplonemea, and Euglenida (supplementary

dataset S1A, Supplementary Material online). The improved taxonomic coverage of the

query MSA increased the sensitivity of the homology detection by HHpred and reinforced the

credibility of the matches to mitofilin for Mic34 and Mic40 by increasing the probabilities to

99.16% and 98.03%, respectively, and lowering the E-values to 2.3e-7 and 0.021,

respectively (supplementary figs. S1 and S2, Supplementary Material online). In contrast, a

HHpred search with *T. brucei* Mic24 (formerly 'Mic60', see above) did not retrieve Mic60 or

the mitofilin domain even among all 250 best hits displayed by default. This did not change

when we used as queries custom alignments of a broader set of kinetoplastid or

euglenozoan Mic24 homologs (listed in supplementary dataset S1B, Supplementary Material

online). Comparison of profile HMMs derived from MSAs of the kinetoplastid and euglenid

Mic24 sequences revealed a region of homology that was restricted to a previously delimited

C-terminal α-helical region (Kaurov et al. 2018). We conclude that this represents a novel

Mic24-specific domain without homologs discernible by HHpred. Notably, in contrast to

Mic34 and Mic40 with ubiquitous occurrence in euglenozoans, no Mic24 homologs were

identified in diplonemids and the endosymbiotic kinetoplastids of the genus *Perkinsela*.

Inspection of the HHpred Mic34 and Mic40 alignments to the mitofilin protein family entry

showed that the Mic60 region detected to be homologous to one or the other euglenozoan

protein differs (Fig. 1A). For the Mic34-Mic60 comparison the match was restricted to a C-

terminal part of the former and a region of the latter exactly corresponding to the 'mitofilin

domain' as delineated by Benning et al. (2025). The match between Mic40 and Mic60

extended further upstream to include a larger part of the Mic60's central coiled-coil (CC)

region. This alignment covered almost the full Mic40 protein, except for a short N-terminal

region, which corresponds to a part of Mic40 not detected by mass spectroscopy in a

previous study, implying it potentially is not part of the mature protein (Kaurov et al. 2018).

To further illuminate the relationship between Mic60 and euglenozoan Mic34 and Mic40, we

focused on the mitofilin domain apparently shared by all three proteins. For this purpose, we

collected a reference set of Mic60 sequences enriched with proteins from eukaryotes most

closely related to Euglenozoa (i.e., other phyla of the supergroup Discoba; see

supplementary dataset S1C, Supplementary Material online) and created a custom

alignment of their mitofilin domain. Pairwise profile HMM-HMM comparisons of the different

mitofilin domains using HHpred indicated a higher sequence similarity in the case of Mic34-

Mic60 comparison than for the Mic40-Mic60 comparison (the respective alignment scores

being 93.09 and 72.52, respectively). Notably, an analogous comparison of the Mic34 and

Mic40 mitofilin domain HMMs yielded an alignment restricted to their C-terminal parts (and

hence with a much lower score, 47.38 or 49.55 depending on which domain is used as a

query). Phylogenetic analysis of the mitofilin domain sequences divided them into the three

expected groups, albeit with low bootstrap support for each clade, and confirmed the

mutually high divergence of Mic34 and Mic40 sequences (manifested by their branch length;

supplementary fig. S3, Supplementary Material online). The most parsimonious scenario

explaining the origin of Mic34 and Mic40 starts with a canonical *mic60* gene in the

euglenozoan ancestor followed by its duplication into the eventual Mic34 and Mic40

paralogs. Subsequent rapid evolution of the two paralogs diverged them considerably from

each other, both in the N-terminal region of the encoded protein as well as in the mitofilin

domain. Owing to the presence of both Mic34 and Mic40 orthologs in all three major

euglenozoan lineages, all these events must have happened before their divergence.

The mitofilin domain is the most conserved primary structure of prokaryotic and eukaryotic

Mic60 proteins (Muñoz-Gómez et al. 2015; Huynen et al. 2016). To bolster confidence that

Mic34 and Mic40 are cryptic mitofilin proteins, we employed AlphaFold2 predictions of the

mitofilin domains starting from its second alpha helix (α2) (Benning et al. 2025) found in the

three representatives from each euglenozoan group, namely *T. brucei* (Kinetoplastea),

Hemistasia phaeocysticola (Diplonemea), and *Euglena gracilis* (Euglenida) (Fig. 1B). These

tertiary structures were aligned to the mitofilin domain of discoban *Naegleria gruberi* Mic60,

which overlaps the theoretical *Saccharomyces cerevisiae* and the experimentally solved

Thermochaetoides thermophila (hereafter referred to by its previous name *Chaetomium*

thermophilum) (Bock-Bierbaum et al. 2022) mitofilin domains with a root-mean-square

distance (RMSD) of <1 Å; *N. gruberi*'s mitofilin aligned to each fungal one equivalently as the

two fungal structures did to each other (supplementary fig. S4A, Supplementary Material

online). Expectedly, the Mic34 mitofilin domains of the three euglenozoan representatives

are well aligned (supplementary fig. S4B, Supplementary Material online).

A linear map of Mic34 and Mic40's mitofilin domain alpha helices α2-α8 (Benning et al. 2025)

was derived from the high confidence predicted structures (Fig. 1C) and multiple sequence

alignments used to make the aforementioned profile HMM (Fig. 1A) and phylogenetic tree

(supplementary fig. S3, Supplementary Material online). The extended loop between α2 and

α3 in Mic60 *S. cerevisiae* (and to a lesser extent *N. gruberi*) is considerably reduced in

Mic34 and Mic40. Lipid binding site (LBS) 1, which is necessary for optimal lipid binding and

yeast cristae junction formation (Hessenberger et al. 2017), was present in all examined

Mic34 and Mic40. In almost all structural alignments, a highly conserved and functionally

important aromatic amino acid (Muñoz-Gómez et al. 2023) within LBS1 perfectly overlapped

(Fig. 1B; supplementary fig. S4A, Supplementary Material online). This phenylalanine is

staggered in *T. brucei* compared to all the other LBS1 structures examined (Fig. 1B;

supplementary fig. S4B, Supplementary Material online) because it is shifted by one amino

acid in the kinetoplastid primary structures (supplementary fig. S7F, Supplementary Material

online). Discoban LBS1 appears to be an amphipathic helix with the hydrophobic face

oriented like fungal LBS1; a notable exception is *T. brucei* Mic34, whose hydrophobic face is

shifted ~90º counterclockwise (supplementary fig. S4D, Supplementary Material online).

The adjacent LBS2 (α5) was much more variable, consistently being the least reliably

predicted motif that aligned poorly to the reference *N. gruberi* mitofilin domain

(supplementary fig. S4C, Supplementary Material online). Furthermore, it is missing in *H.*

phaeocysticola Mic34 and Mic40. The orientation of the last three alpha helices α6-8 is well

conserved in all three representative Mic34 proteins. In Mic40, this is a more malleable

feature, with *T. brucei* and *E. gracilis* having a shorter alpha helix downstream of α8, and α6

fused to LBS2 or missing, respectively (Fig. 1C). In summary, the mitofilin domain structure

of Mic34 appears to be more similar to canonical Mic60 than Mic40.

Mic34 and Mic40 are involved in the mitochondrial intermembrane space assembly

pathway that affects mitochondrial morphology

To verify *in silico* predictions that Mic34 and Mic40 are cryptic mitofilins, we examined how

their depletion in procyclic stage *T. brucei* (PCF), which bears mitochondria with mature

cristae that house an active electron chain (Bílý et al. 2021), affects the organelle's

morphology. These inducible-RNAi cell lines were previously characterized (Kaurov et al.

2018; Eichenberger et al. 2019). Here, we recapitulated the expected growth inhibition upon

RNAi silencing of Mic34 and Mic40 (supplementary fig. S5A, Supplementary Material

online). Qualitative (Fig. 2A; supplementary fig. S5B, Supplementary Material online) and

quantitative (Fig. 2B) assessment of mitochondrial complexity showed that branching was

reduced at 3- and 5-day time points (TP3 and TP5) after RNAi induction, with the organelle

exhibiting a tubular appearance in the latter. *S. cerevisiae mic60* null mutants also exhibit

alternations to the mitochondrial network (Rabl et al. 2009).

Interpretation of the observed morphological defects may be confounded by downregulation

of proteins imported into the IMS by the MIA pathway, which has been observed upon Mic34

silencing (Eichenberger et al. 2019). This role corresponds with the IMS localization of Mic34

and Mic40 (Kaurov et al. 2018).To see if Mic40 depletion has the same effect on IMS

proteins as Mic34, we first verified that the MIA substrate Tim9 (Peikert et al. 2017) was

downregulated during the time course of RNAi induction (supplementary fig. S5C,

Supplementary Material online). To gain a comprehensive view on the cohort of proteins that

may rely on Mic40 for their import, we performed a mitoproteome-wide analysis (*i.e.*

depletome) of Mic40 TP3 and TP5 relative to wild type *T. brucei* (Fig. 2C-D). The MICOS

subcomplex subunits residing completely in the IMS peripherally to the IM, were

downregulated prior to reduction of MIA substrates (Fig. 2C-D). A similar phenotype was

observed upon Mic34 silencing (Eichenberger et al. 2019). Indeed, the Mic40 depletome

overlapped with those of Mic34 (Fig. 2D, left) and Mic20 (Fig. 2D, middle), a thioredoxin-like

subunit of the peripheral, IMS-localized MICOS subcomplex (Kaurov et al. 2018; Kaurov et

al. 2022). Thus, Mic40 also plays an integral part in IMS protein import by MIA.

Returning to the question of whether IMS import defects may underlie the reduction in

mitochondrial complexity seen in Mic34 and Mic40 knockdown cells, we measured the

degree of branching in TP3 and TP5 Erv1 RNAi PCF (Fig. 2B; supplementary fig. S5B,

Supplementary Material online). Erv1 is a sulfhydryl oxidase that plays an essential role in

the MIA pathway (Mordas and Tokatlidis 2015; Peikert et al. 2017), and is not a component

of *T. brucei* MICOS (Kaurov et al. 2018; Turra et al. 2021). Erv1 depletion phenocopies those

of Mic34 and Mic40 to the same degree Erv1 depletion's effect on mitochondrial complexity

was previously reported based on qualitative observations (Niemann et al. 2013; Haindrich

et al. 2017). The Mic34 and Mic40 depletomes overlap with Erv1's (Peikert et al. 2017). It

should be noted that peripheral MICOS subunits were also downregulated upon Erv1

silencing (Peikert et al. 2017), possibly leading to the observed morphological alteration.

Consistent with this hypothesis, these subunits were by far the most affected at TP3 (Fig.

2D), when simplification of the mitochondrion was first observed during Mic40

downregulation (Fig. 2A-B). However, while these results do not rule out that Mic34 or Mic40

are directly involved in membrane remodeling, we still cannot decouple this function from its

established but still unexplained role in the MIA pathway.

Mic34 binds to phospholipids *in vitro*

First, we decided to test if Mic34 and Mic40 can deform liposomes *in vitro*, as has been

demonstrated for Mic60 (Hessenberger et al. 2017; Tarasenko et al. 2017). For this

experiment, recombinant *T. brucei* Mic34 and Mic40 bearing N-terminal His-tags (rMic34 and

rMic40, respectively) were expressed in *Escherichia coli*, where they sequestered into

inclusion bodies (supplementary fig. S6A-B, Supplementary Material online). After their

purification by sequential affinity and size exclusion chromatography, rMic34 and rMic40

were solubilized with varying concentrations of the mild detergent n-dodecyl β-d-maltoside

(DDM). While the majority of rMic34 proved to be soluble in 0.1% DDM, rMic40 was poorly

solubilized in any of the tested DDM concentrations (supplementary fig. S6C-D,

Supplementary Material online). Thus, we excluded rMic40 from further analysis.

Next, 0.1% DDM solubilized rMic34 was loaded onto large unilamellar vesicles (LUVs)

comprised of IM phospholipids (Barbot et al. 2015; Tarasenko et al. 2017), followed by DDM

removal. A fraction of rMic34 was carbonate extracted from the LUVs (Fig. 3A), which is the

expected behavior of the IMS-periphery localized Mic34 ((Kaurov et al. 2018) and also see

below). Surprisingly, most rMic34 acted as an integral protein in this biochemical assay. To

eliminate the possibility of DDM-mediated rMic34 insertion into the membrane, we performed

a Proteinase K protection assay to see if Mic34 is binding the exterior of the LUV (Fig. 3B).

The protease completely degraded rMic34 in the liposome sample even in the absence of

the membrane-disrupting detergent sodium dodecyl sulfate (SDS). This mimicked the

behavior of the *S. cerevisiae* rMic19, which serves as a negative control of phospholipid

binding due to an absence of this *in vitro* activity (Hessenberger et al. 2017), and thus

remains exposed to Proteinase K.

Next, we assayed whether rMic34 deformed the LUVs by flotation on Histodenz density

gradients (Fig. 3C). As previously reported (Barbot et al. 2015; Tarasenko et al. 2017), LUVs

alone float to the lightest fractions of the gradient. After incubation with rMic34, LUVs with or

without cardiolipin formed multiple bands in denser fractions of the gradient, reminiscent of

canonical Mic60 (Tarasenko et al. 2017). The presence of rMic34 in these fractions was

verified by immunoblotting (Fig. 3D), indicating that rMic34 binding is weighing down the

LUVs. The signal in the heaviest fraction corresponds to unbound rMic34, as seen in the

rMic19-incubated LUV negative control.

To see if the striated banding pattern we observed often corresponds to LUV deformation

(Barbot et al. 2015; Tarasenko et al. 2017), LUVs that flotated into lower density fractions

were isolated and observed by negative-stain transmission electron microscopy (TEM).

However, we could not reliably observe any obvious differences between the naked LUVs

and those incubated with rMic34, independently of the presence of cardiolipin in the LUV

membrane (Fig. 3E).

This lack of LUV deformation may be because rMic34 does not have *in vitro* membrane

deformation activity or this activity is inhibited by the 0.1% DDM treatment. To test the latter

hypothesis, we compared the liposome deformation activity of Epsin N-terminal Homology

Domain (ENTH) that underwent 0.1% DDM solubilization prior to incubation with Folch

liposomes spiked with 5% PI(4,5)P2 (DDM+) to that of untreated ENTH (Gleisner et al.

2016) (DDM-). DDM+ ENTH bound to the outside of liposomes (Fig. 3B-C). However,

liposome tubulation was only observed in the absence of the detergent treatment (Fig. 3F).

Thus, we are unable to address Mic34's putative membrane remodeling activity *in vitro* with

our current experimental set up. Nevertheless, we conclude that Mic34 binds phospholipids

in vitro in a cardiolipin-autonomous way like yeast Mic60 (Hessenberger et al. 2017).

Mic34 and Mic40 are capable of remodeling the cytoplasmic membrane of a

gammaproteobacterium upon export to the periplasm

After establishing that Mic34 is able to bind phospholipid membranes, we next investigated

whether Mic34 and Mic40 have membrane-bending activity *in vivo*. We employed an assay

that has been used to demonstrate that Mic60 has this capacity (Tarasenko et al. 2017).

Mic34 and Mic40 from *T. brucei* were heterologously expressed in the

gammaproteobacterium *E. coli*. To facilitate their export into the periplasm, which represents

the proteobacterial analog of the mitochondrial IMS, each MICOS subunit was fused to

maltose binding protein (MBP).

First, *E. coli* transformed with various MBP expression constructs were separated into

periplasm and spheroplast fractions (Fig. 4A). As expected, MBP alone was completely

exported into the periplasm. This and the retention of T7 polymerase within the spheroplasts

(supplementary fig. S7A, Supplementary Material online) verified successful fractionation.

While a minor portion of Mic34-MBP and Mic40-MBP was observed in the periplasm, they

were enriched in the spheroplast fraction (Fig. 4A). Carbonate extraction of the spheroplasts

was done to see whether Mic34-MBP and Mic40-MBP were embedded in the membrane

(Fig. 4A), like the integral subunit Mic24 fused to MBP (supplementary fig. S7B,

Supplementary Material online). While some Mic34-MBP and Mic40-MBP were observed in

the integral fraction–reminiscent of Mic34's behavior in LUVs (Fig. 3A)–most accumulated in

the soluble fraction (Fig. 4A). Our interpretation is that Mic34-MBP and Mic40-MBP tightly

associate with periplasm-facing part of the cytoplasmic membrane because (1) the protein is

partially observed in the periplasm fraction and (2) Mic34 and Mic40 are sorted into inclusion

bodies when expressed in the cytoplasm (supplementary fig. S6A-B, Supplementary

Material online).

Next, we looked if Mic34 and Mic40 remodeled *E. coli* membranes similarly to

heterologously expressed full-length *S. cerevisiae* Mic60 or just its IMS part (Tarasenko et al.

2017). After establishing by TEM that we can replicate this experiment (supplementary fig.

S7C, Supplementary Material online), including reproducing that MBP alone does not affect

E. coli ultrastructure (Fig. 4B), we observed Mic34-MBP and Mic40-MBP expressing *E. coli*

by electron tomography (Fig. 4C) and TEM (supplementary fig. S5D, Supplementary

Material online). Mic34-MBP and Mic40-MBP had respectively 5 and 3 times more vesicles

observed by TEM in the periphery of the cell as compared to the parental strain (Fig. 4D).

Immunogold labelling of the MBP moiety showed that both chimeras were also observed in

the periphery, associated with the majority of vesicles. Electron tomography allowed us to

better resolve these vesicles, which accumulated in the periplasm (Fig. 4C-D; supplementary

movies S1 and S2, Supplementary Material online). Furthermore, we used this method to

determine that the bulk of MBP-visualizing gold nanoparticles were associated with the

vesicles (supplementary fig. S7E, Supplementary Material online). Together these results

indicate that Mic34-MBP and Mic40-MBP are directly responsible for cytoplasmic membrane

remodeling when heterologously expressed in *E. coli*, phenocopying *S. cerevisiae* Mic60 in

this assay.

To test whether the predicted LBS helices of the slightly more conventional Mic34 (Fig. 1;

supplementary fig. S7F, Supplementary Material online) are important for membrane

remodeling, as has been demonstrated for *S. cerevisiae* Mic60 (Hessenberger et al. 2017),

we expressed a periplasm-targeted LBS deletion mutant (Mic34ΔLBS1+2). We first tested

whether the ~6 kDa smaller mutant was expressed (Fig. 4E) and targeted to the periplasmic

face of the *E. coli* cytoplasmic membrane (supplementary fig. S7A, Supplementary Material

online). As expected, this mutant failed to remodel the cytoplasmic membrane, having about

the same number of vesicles as the parental strain (Fig. 4D-E).

Wondering whether Mic34 and Mic40 can simultaneously remodel the cytoplasmic

membrane, we transformed *E. coli* with two plasmids respectively encoding Mic34-MBP or

Mic40-MBP. Co-expression was observed by gel staining, which revealed strong bands

corresponding to the theoretical molecular weights of each chimera (supplementary fig. S8A,

Supplementary Material online). This result was verified by immunoblotting with antibodies

specifically recognizing Mic34 and Mic40. These cells exhibited prominent vesicles within the

cytosol that were absent from the negative control when observed by electron tomography

(supplementary fig. S8B-F, Supplementary Material online). This result suggests that Mic34

and Mic40 may work in concert, evocative of how Mic60 self-interaction mediates membrane

remodeling activity (Bock-Bierbaum et al. 2022).

In conclusion, with an assay based on heterologous expression in *E. coli*, we were able to

show that (1) Mic34 and Mic40 have the capacity to remodel membranes, and (2) they may

act together to do so, and (3) Mic34's LBS is directly involved in this activity. Thus, we

provide empirical evidence that Mic34 and Mic40 are extremely diverged mitofilin family

proteins.

Overexpression of Mic34 causes reticulation of a morphologically simplified

trypanosomal mitochondrion

The long slender bloodstream form (BSF) *T. brucei* has a morphologically simplified

mitochondrion that runs as a tube along the cell body opposite the flagellum, reflecting the

cessation of cellular respiration in this life cycle stage (Jakob et al. 2016; Hughes et al. 2017;

Bílý et al. 2021). All MICOS complex subunits are downregulated in this stage compared to

PCF (Kaurov et al. 2018). Thus, expression of electron transport chain subunits is

necessarily triggered during BSF differentiation into PCF (Zíková 2022), corresponding with

mitochondrial reticulation.

Ectopic expression of V5-epitope tagged Mic34 (Mic34-V5$^+$) with 1000 ng/ml doxycycline in

wild type BSF was considerably higher than that of endogenous Mic34 (Fig. 5A).

Surprisingly, this overexpression reproducibly induced cytostasis after 24 hours, with

induction by even 10 ng/ml doxycycline effectively inhibiting growth (Fig. 5A; supplementary

fig. S9A-B, Supplementary Material online). Mic40-V5$^+$ ectopic expression only mildly

inhibited growth (supplementary fig. S9B, Supplementary Material online), perhaps due to its

lower expression (supplementary fig. S9A, Supplementary Material online). After verifying

that Mic34-V5$^+$is targeted to the mitochondrion (supplementary fig. S9C, Supplementary

Material online), we assayed how mitochondrial morphology was affected after 36 hours of

Mic34-V5$^+$ due to its cytostatic effect. Many Mic34-V5$^+$ cells exhibited profound reticulation of

their mitochondrion (Fig. 5C), with ~2.3 times more branching than in the wild type BSF

organelle (Fig. 5D). Apparent fragmentation of the organelle was also observed

(supplementary fig. S9D, Supplementary Material online). However, we did not notice any

obvious changes to the cristae that were sporadically observed by TEM (supplementary fig.

S9E, Supplementary Material online). This is consistent with the uninterrupted absence of

electron transport chain subunits and unaffected F_OF_1- ATP synthase expression in Mic34-

V5$^+$ cells (supplementary fig. S9F, Supplementary Material online). Furthermore, IMS protein

levels were not altered, unlike during Mic34 depletion in PCF (Eichenberger et al. 2019).

Thus, the observed morphological changes are likely due to Mic34-V5$^+$ acting on the

mitochondrion directly.

We were concerned that Mic34-V5$^+$ branching phenotype was an artifact of cell cycle arrest.

The mitochondrion becomes more branched during cell cycle progression to ensure

inheritance of the organelle after cytokinesis (Jakob et al. 2016; Hughes et al. 2017). Thus, a

defect to this process could secondarily cause a reticulated mitochondrion (Fig. 5E). To

address this issue, we counted the number of kinetoplasts and nuclei in Mic34-V5$^+$ cells

bearing a reticulated mitochondrion; branching starts during kinetoplast division, perceptively

increasing as the cell goes from having a single nucleus and kinetoplast (1N1K) to two

separate kinetoplasts (1N2K) and eventually two separate nuclei (2N2K) (Fig. 5E). The vast

majority of unsynchronized Mic34-V5$^+$ cells with a reticulated mitochondrion were 1N1K (Fig.

5C, F), at early cell cycle stages that normally have a tube-like mitochondrion. As expected,

wild type BSF exhibited branched mitochondria mostly in the later 2N2K cell cycle stage.

Thus, we conclude that Mic34-V5$^+$ overexpression directly affects mitochondrial morphology.

We hypothesized that the Mic34 LBS alpha helices may be important for the observed cell

cycle arrest due to mitochondrial remodeling. We overexpressed mutants in which both

(Mic34ΔLBS1+2$^+$) or just LBS1 (Mic34ΔLBS1$^+$) were deleted (supplementary fig. S7F,

Supplementary Material online). Like WT Mic34, these mutants were also targeted to the

mitochondrion (supplementary fig. S9C, Supplementary Material online). After 24 hours of

overexpression growth inhibition correlated with the extent of LBS deletion (Fig. 5A). The

mild growth inhibition in Mic34ΔLBS1+2$^+$ could be explained by its relatively lower

expression compared to others (Fig. 5B, *c.f.* lane 4 with 2 and 4). However, Mic34ΔLBS1+2$^+$

and Mic34-V5$^+$ induced with 10 ng/ml doxycycline (Fig. 5A, lane 4) are equivalent, with the

latter exhibiting more dramatic growth inhibition than the deletion mutant. Together with our

previous *E. coli* data, we conclude that the LBS of the mitofilin-like Mic34 plays an important

functional role as it does in canonical Mic60.

Tertiary (Fig. 1B-C) and primary structural alignments predicted that the conserved LBS1

phenylalanine of *T. brucei* LBS1 (F175) would be a critical residue for the observed Mic34

mitochondria remodeling given it and an upstream arginine (R) residue were shown to be

essential for yeast Mic60 remodeling activity (Hessenberger et al. 2017). We expressed a

double point mutant affecting these amino Mic34$^{R170D\_F174D}$, which was equivalently

expressed compared to Mic34-V5+)(Fig 5G; supplementary fig. S9C, Supplementary

Material online) and targeted to the mitochondrion (supplementary fig. S9C, Supplementary

Material online). The growth of Mic34$^{R170D\_F174D}$ was restored to almost the same rate as that

of the uninduced counterpart (Fig. 5G), which may be explained by a mitigation of apparent

fragmentation seen in Mic34$^{R170D\_F174D}$ compared to Mic34-V5$^+$ (Fig. 5H; supplementary S9D,

Supplementary Material online). Thus, these conserved amino acid residues are essential

for the dominant negative effect seen upon expression of Mic34-V5$^+$. We conclude that

Mic34 was directly responsible for the mitochondrial remodelling observed upon its

overexpression. Furthermore, the cytostatic effect of Mic34-V5$^+$ may be linked with the

apparent fragmentation that also occurs.

DISCUSSION

Here, we provide *in silico* and experimental evidence supporting the conclusion that

euglenozoan MICOS contains two soluble subunits that represent cryptic mitofilin family

proteins. In contrast to Mic60 from two domains of life, Mic34 and Mic40 do not contain TMD

that embeds them into a phospholipid bilayer, as supported by its repeated association with

the periphery of membranes in this and another (Kaurov et al. 2018) study. It has been

proposed before in passing (Hashimi 2019; Prokopchuk et al. 2023) that the ancestral Mic60

got split into two parts, one represented by the integral membrane Mic24 and the other by

the soluble Mic34 in the forebear of Euglenozoa. Our analyses did not provide any evidence

that Mic24 evolved from any part of the ancestral Mic60 gene, and its origin remains

obscure. However, our results corroborate that Mic34 indeed represents a cryptic mitofilin

domain-containing protein. Additionally, we demonstrate that the still hypothetical split of

integral and soluble components was followed by duplication of the latter, mitofilin domain-

containing part, resulting in the paralogous Mic34 and Mic40 pair. Admittedly, the

phylogenetic tree inferred from mitofilin domain sequences is not well resolved and does not

support this scenario, as it would need Mic34 and Mic40 to branch most closely to Mic60

from Heterolobosea (the sister group to Euglenozoa). However, this is not unexpected, given

the sequence divergence of the sampled mitofilin family proteins and the resulting

challenges for phylogenetic inference; note also that the branching order of Mic60 itself

conforms less to organismal phylogeny than that of Mic34 and Mic40.

The presence of cryptic mitofilin domains in Mic34 and Mic40 predicts that these are

membrane remodeling proteins like conventional Mic60 from prokaryotes and eukaryotes.

However, we were unable to verify Mic34 and Mic40's capacity to remodel phospholipid

bilayers *in vitro* here. Nevertheless, this assay did reveal Mic34 can bind liposomes. Thus,

we had to resort to assaying their putative role in membrane remodeling by other means.

Heterologous expression of Mic34 and Mic40 deformed *E. coli* cytosolic membranes, the

prokaryotic analog to the mitochondrial IM (Muñoz-Gomez et al. 2017; Muñoz-Gómez et al.

2023), as yeast and alphaproteobacterial Mic60 did (Tarasenko et al. 2017). However, a

difference should be noted here: Mic60 caused the cytoplasmic membrane to invaginate

inward into the cytoplasm whereas Mic34 and Mic40 promoted the accumulation of vesicles

within the IMS. Furthermore, Mic34 and Mic40 may work together as their co-expression in

E. coli results in conspicuous membrane remodeling. This suggests that Mic34 and Mic40

interact with each other *in vivo*, as conventional Mic60 acts on membranes as a

homooligomer (Bock-Bierbaum et al. 2022). Mic34 requires the LBS motifs it shares with

Mic60 (Hessenberger et al. 2017) for cytoplasmic membrane vesicularization, consistent

with the notion it is a cryptic member of the mitofilin protein family.

Mic34 ectopic expression in a *T. brucei* BSF with a reduced, tubular mitochondrion causes

elaboration into a branched organelle. This parallels the enhanced interconnectivity of

mitochondria upon Mic60 overexpression in primary and immortalized neuronal cell lines

(Van Laar et al. 2016). These assayed Mic34 membrane-deformation activities *in vivo* rely

on its predicted LBS motifs, further supporting Mic34 and its paralog Mic40 as diverged

mitofilin protein family members. LBS1 contains a conserved aromatic amino acid (Muñoz-

Gómez et al. 2023) that is essential for Mic60's *in vitro* and *in vivo* activities (Hessenberger

et al. 2017). This residue along with an upstream arginine residue is also essential for

Mic34-mediated BSF mitochondria remodeling, as predicted. The aromatic residue in *C.*

thermophila Mic60 is buried inside the hydrophobic interaction interface of the Mic60

homodimer (Bock-Bierbaum et al. 2022). Whether this residue plays a similar role in Mic34

homooligomerization and/or heterooligomerization remains an open question.

Mic34 overexpression did not alter cristae dramatically in BSF *T. brucei*, contradicting Mic60

overexpression phenotypes in yeast (Rabl et al. 2009) and alphaproteobacteria (Muñoz-

Gómez et al. 2023), manifested in the branching of their respective bioenergetic

subcompartments (*e.g.* cristae). However, this inconsistency can be explained by the lack of

an active electron transport chain in BSF (Bílý et al. 2021; Zíková 2022), which were present

when Mic60 was overexpressed in yeast and bacteria. Thus, an electron transport chain

may be necessary for populating and stabilizing elaborated crista membranes. Alternatively

and/or additionally, co-expression in *E. coli* implies Mic34 interacts stoichiometrically with

Mic40 and/or (an)other MICOS subunit(s) to shape crista.

One clear divergence from canonical Mic60 is that Mic34 (Eichenberger et al. 2019) and

Mic40 (this study) appear to be directly integrated into the MIA pathway for IMS protein

import in PCF, like the thioredoxin-like Mic20 subunit (Kaurov et al. 2018; Kaurov et al.

2022). In contrast, while yeast Mic60 augments the Mia40 sulfhydryl oxidase's mediation of

MIA (von der Malsburg et al. 2011), Mic60 deletion does not lead to depletion of MIA

substrates (Varabyova et al. 2013) like Mic34 and Mic40 silencing. The MIA defects seen

upon depletion of Mic34 and Mic40 complicate the interpretation of crista defects given that

putative respiratory chain subunits and assembly factors, plus the small Tim proteins, are

imported via MIA. Moreover, the latter are needed for import of IM proteins (Wenger et al.

2017; Smith et al. 2018), many of which are assembled into respiratory chain complexes

(Zíková 2022). Based on our results, the MIA pathway also is important for maintaining

reticulated mitochondria in PCF, as shown by Erv1 simplifying mitochondrial complexity to

the same degree as Mic34 and Mic40. However, it is possible that concomitant

downregulation of the soluble MICOS subunits Mic20 and Mic32–and by extension, perhaps

the rest of this subcomplex–may be responsible for the morphological phenotype of Erv1

RNAi after all (Peikert et al. 2017). Thus, at this time we cannot verify nor refute the

hypothesis that MICOS plays a role in maintaining a reticulated mitochondrial network *via* its

demonstrated membrane remodeling activity.

It is worth noting that this work has implications about the subcomplex architecture of

MICOS throughout eukaryotes. In opisthokonts, two adjacent integral membrane

subcomplexes are each assembled around their respective core subunit, Mic10 and Mic60

(Hashimi 2019; Colina-Tenorio et al. 2020). It is reasonable to expect this 'horizontal'

(relative to the membrane) configuration in the majority of aerobic eukaryotes because they

have both Mic10 and Mic60 (Muñoz-Gómez et al. 2015; Huynen et al. 2016). Uniquely,

trypanosomal–and by extension likely also euglenazoan–MICOS has a vertical architecture:

a membrane-embedded subcomplex containing two Mic10 paralogs and a peripheral

subcomplex comprised of soluble subunits located within the IMS (Eichenberger et al. 2019).

This divergent subcomplex architecture likely corresponds to the soluble nature of IMS

localized Mic34 and Mic40 (Kaurov et al. 2018). Significantly, the existence of the

Mic34/Mic40 subcomplex implies that a mitofilin domain represents the core of a MICOS

module regardless of whether or not it is directly IM anchored. Perhaps the Mic60

subcomplex is more predisposed to IM detachment than the Mic10 subcomplex. In

opisthokonts, the only soluble MICOS subunit Mic19 (plus its paralog Mic25 in vertebrates)

is part of the Mic60 complex (Muñoz-Gómez et al. 2015; Huynen et al. 2016; Hessenberger

et al. 2017; Bock-Bierbaum et al. 2022). In contrast, the persistence of the trypanosomal

Mic10 subcomplex within the IM suggests it is perpetually an integral IM MICOS module.

In the diverged euglenozoans and opisthokonts, a mitofilin protein represents the largest

subunit in the MICOS complex. This may reflect the mitofilin domain's role in bridging

juxtaposed membranes at contact sites and crista junctions as well as being an interaction

hub to facilitate coordination of MICOS with other fundamental pathways (Aaltonen et al.

2016; Michaud et al. 2016). Such a central role in MICOS is likely a product of the mitofilin

domain's ancient, pre-mitochondrial origin (Muñoz-Gómez et al. 2023). Unlike the rather

widespread loss of Mic60 in anaerobes without cristae, the transfiguration of Mic60 into

Mic34/Mic40 appears to be a singular event that occurred exclusively in Euglenozoa, joining

other unique mitochondrial apomorphies of this phylum (Cavalier-Smith 2016).

Our work here also shows that the mitofilin protein family is more diverse than originally

thought, given the conservation of Mic60 in alphaproteobacteria and mitochondria (Muñoz-

Gómez et al. 2023). It is unknown whether the duplication of the soluble mitofilin proteins is

an adaptation to loss of their respective TMDs, or perhaps the divergence of Mic34 and

Mic40's mitofilin domains necessitates loss of their membrane integration. Perhaps the

duplication of the mitofilin domain, resulting in the Mic34 and Mic40 paralogs during early

euglenozoan evolution, allows one of the mitofilin domains to be fine-tuned to the needs of

the subdivisions Euglenida, Diplonemea, and Kinetoplastea. This reveals surprising

variability of the mitofilin family of proteins found in the organismal diversity occurring in two

domains of life.

MATERIAL AND METHODS

Gathering of euglenozoan protein sequence datasets

Homologs of *T. brucei* Mic34 and Mic40 were identified by a combination of blast (Altschul et

al. 1997) and HMMER3 searches (Eddy 2011) in a variety of databases and sequence

resources, including the NCBI non-redundant protein sequences database, TriTryp (Alvarez-

Jarreta et al. 2024), EukProt v3 (Richter et al. 2022), and sequence assemblies and

annotations accompanying individual publications (details provided in supplementary dataset

S1, Supplementary Material online). Briefly, homologs were first collected with blast

(considering hits with e-value <1e-5) and aligned with MAFFT v7

(https://mafft.cbrc.jp/alignment/server/; (Katoh et al. 2019)). (Katoh et al. 2019)The HMMER3

package was then used for building a profile HMM for each MSA and searching the target

protein sequence sets (using the hmmbuild and hmmsearch command, respectively). Newly

identified hits missing in the original MSA and passing the program-defined inclusion

threshold were collected, added to the MSA and the whole search was repeated until no new

significant hits were retrieved or the search started to retrieve hits that were found not to

represent euglenozoan Mic34 or Mic40 proteins. In a few cases homologs were obtained by

tblastn searches against genome or transcriptome assemblies and the sequence of the

respective encoded proteins were deduced by conceptual translation.

Only a subset of identified sequences was retained for further analyses, taking into account

the degree of their completeness, whether both Mic34 and Mic40 homologs were identified

for the given species (some of the sequences resources, especially the various euglenid

single-cell transcriptomes, are very incomplete and did not yield both expected homologs),

and keeping the taxon sampling balanced (i.e. avoiding the inclusion of highly similar

sequences from closely related species). As a result, we obtained a set of Mic34 and Mic40

homologs from 44 euglenozoan taxa encompassing representatives of all three main

euglenozoan clades as well as their various sublineages, In a few cases the sequences

were attributed to a wrong organism in the given database, stemming from contamination of

the source protist cultures used for transcriptome sequences by a co-cultured euglenozoan,

whose identity was determined by investigating informative marker sequences present in the

same assembly, such as the 18S rRNA gene or elongation factor 2 (EF2) protein. All the

sequences were manually curated to achieve their accuracy (e.g. by removing artificial N-

terminal extensions frequently present in protein sequences automatically inferred from

transcriptome assemblies); in some cases the existing truncated sequences were extended

or even completed by manually recruiting original Illumina sequencing reads identified by

iterative blast searches against the respective sequencing runs in the Sequence Read

Archive (SRA; https://www.ncbi.nlm.nih.gov/sra).

Homologs of the *T. brucei* Mic24 protein (Tb927.9.10160) were searched and curated in a

manner analogous to that used for gathering the Mic34 and Mic40 sequence datasets. The

euglenid homologs retrieved with a profile HMM derived from a MSA of kinetoplastid Mic24

proteins included a protein from *Euglena gracilis* that was listed as a putative ortholog of *T.*

brucei Mic24 (=Mic60) in a previous study (Hammond et al. 2020). No candidates for Mic24

homologs were identified in any diplonemid and *Perkinsela* spp. even using sensitive

HMMER searches queried with profile HMMs of kinetoplastid, euglenid, or combined

kinetoplastid-euglenid Mic24 for MSAs, suggesting loss or extreme divergence of Mic24 in

these lineages. We likewise failed to identify a Mic24 homolog in the euglenid *Ploeotila vitrea*

(a species included in our selection of taxa for the Mic34 and Mic40 analysis), but

incompleteness of the available sequence data (a transcriptome assembly) cannot be

excluded in this case. All the Mic34, Mic40, and Mic24 sequences included in the final

datasets and the related metadata are provided in supplementary dataset S1,

Supplementary Material online.

HHpred searches

Homology of Mic34 and Mic40 to other proteins was rigorously examined with HHpred

(Söding et al. 2005) as available at the MPI Bioinformatics Toolkit server

(https://toolkit.tuebingen.mpg.de/tools/hhpred). The searches with queries representing

individual sequences, i.e. *T. brucei* Mic34 and Mic40, were done with the default parameters,

i.e. with initial three iterative HHblits searches against the UniRef30 database, which in both

cases retrieved a set of exclusively kinetoplastid homologs that were then included by the

program into the query MSA, used then for building a profile HMM and searching the target

database of profile HMMs based on HMM-HMMM comparisons. In the case of searches in

which the query was the custom Mic34 or Mic40 MSA (in each case computed with MAFFT

from the final set of 44 selected euglenozoan representatives), no initial HHblits searches

were allowed and the query MSA was directly used for creating the query profile HMM. For

reference we report results of HHpred searches against the Pfam-A_v37 database, which all

retrieved the Pfam database entry PF09731.14 as the only significant hit. Alternatively, we

also searched the COG_KOG_v.1 database and identified with both Mic34 and Mic40

(individually or as an MSA) the same entry as the best (and significant) hit, i.e. KOG1854,

which represent eukaryotic Mic60 proteins, reinforcing the results of the searches against

Pfam-A. The relative divergence of the mitofilin domains of the euglenozoan Mic34 and

Mic40 proteins and of Mic60 proteins from other eukaryotes was evaluated by carrying

pairwise profile HMM-HMM comparisons with HHpred, using as the query and the target the

respective custom MSAs (the same as used in the phylogenetic analysis; see the next

section).

Phylogenetic analysis of the mitofilin domain

A set of reference eukaryotic Mic60 sequences was gathered by selecting proteins from a

broader sequence set obtained by searching the EukProt v3 database with HMMER3 and a

profile HMM built from the seed alignment of the Pfam family PF09731 (mitofilin). Sequences

were selected to include a representative from different major eukaryote lineages, with a

denser sampling for the supergroup Discoba, owing to the presumed origin of the

euglenozoan Mic34 and Mic40 from an ancestral discoban Mic60. Additional discoban Mic60

sequences were identified by HMMER or blast searches in relevant sequence resources

from species not included in EukProt v3 and added to the dataset (details provided in

supplementary dataset S1, Supplementary Material online). The Mic60 sequences were

combined with the Mic34 and Mic40 sequences used in HHpred analyses (see above) and a

multiple alignment was built using MAFFT v7 and the L-INS-I iterative refinement method.

The alignment was restricted to the C-terminal region corresponding to the mitofilin domain

and used for tree inference performed the ETE3 3.1.3 pipeline (Huerta-Cepas et al. 2016) as

implemented on the GenomeNet server (https://www.genome.jp/tools-bin/ete). Specifically,

the alignment was trimmed with). Specifically, the alignment was trimmed with trimAl

v1.4.rev6 (Capella-Gutiérrez et al. 2009), removing columns with >20% of gaps, and the tree

was inferred using RAxML v8.2.11 (Stamatakis 2014), with the default parameters, the

substitution model PROTGAMMAJTT, and 100 bootstrap replicates. The resulting tree was

visualized in an arbitrarily rooted form using iTOL v6 (Letunic and Bork 2024) and further

processed using a graphical editor. v1.4.rev6 (Capella-Gutiérrez et al. 2009), removing

columns with >20% of gaps, and the tree was inferred using RAxML v8.2.11 (Stamatakis

2014), with the default parameters, the substitution model PROTGAMMAJTT, and 100

bootstrap replicates. The resulting tree was visualized in an arbitrarily rooted form using

iTOL v6 (Letunic and Bork 2024) and further processed using a graphical editor.

Structural predictions

The HHpred server was used to detect the mitofilin domain of Mic34 and Mic40 using hidden

Markov model profiles to interrogate the following manually set structural and domain

databases: PDB_mmCIF70_18_Jun; Pfam-A_v36; UniProt-SwissProt-viral70_3_Nov_2021;

TIGRFAMs_v15.0. E-values reported in Fig. 1A are for the top hit 'PF09731.12; Mitofilin'.

AlphaFold 2 predictions for *T. brucei* Mic34 and Mic40 and *N. gruberi* Mic60 were performed

according to a previously established pipeline catered to discoban proteins (Wheeler 2021).

Mitofilin domains were aligned in ChimeraX by restricting the matchmaker to the two

domains (Meng et al. 2023). The mitofilin alpha helices were mapped by manual inspection

of the aforementioned structural predictions and crosschecked with multiple sequence

alignments to identify deletions and fusions in the mitofilin domains.

***Trypanosoma brucei* culturing, transfection, fractionation and growth analysis**

The PCF Mic34, Mic40 and Erv1 inducible RNAi and SMOXp parental (Poon et al. 2012) cell

lines were generated with the same constructs plus strategy and then grown as previously

described (Haindrich et al. 2017; Kaurov et al. 2018). The pleomorphic AnTat1.1 BSF cell

line was cultivated in HMI-9 media (Hirumi and Hirumi 1989) at cell densities ≤8 × 10^5

cells/ml to maintain pleomorphism. Cells were transfected using a Amaxa Nucleofector (X-

001) with the pT7-3xV5 vector (Flaspohler John et al. 2010), in which wild type and mutated

mic34 and *mic40* was cloned to facilitate their inducible ectopic expression by addition of 1

µg/ml of doxycycline to the media. Cells were fractionated into cytoplasmic and organellar

fractions as previously described (Kaurov et al. 2018). Cell densities were measured on a

Beckham Coulter Z2 Cell and Particle Counter.

Mass spectroscopy analysis

For depletome data, mitochondria from 10^8 parental SMOXp, TP3 and TP5 Mic40i *T. brucei*

cells were crudely isolated as before (Kaurov et al. 2018) and snap frozen in liquid nitrogen

for storage at -80ºC prior to LC-MS/MS mass spectroscopy analysis, which was performed

and the data visualized by a previously established pipeline (Benz et al. 2022) with slight

modifications. Briefly, for sample prep, SP4 (Solvent Precipitation SP3) protocol without

beads was used (Johnston et al. 2022). Pellets of enriched mitochondria (50 µg of protein)

were solubilized by SDS (final concentration 1% [w/v]), reduced with TCEP [tris(2-

carboxyethyl)phosphine], alkylated with chloroacetamide and digested sequentially with Lys-

C and trypsin. Samples were desalted on Empore C18 StageTips, dried in a speedvac, and

dissolved in 0.1% TFA + 2% acetonitrile. About 0.5 µg of peptide digests were separated on

a 50 cm C18 column using 90 min elution gradient and analyzed in DIA mode on an

Orbitrap Exploris 480 (Thermo Fisher Scientific) mass spectrometer equipped with a FAIMS

unit.

Recombinant Mic34, Mic19 and ENTH expression and purification

The Mic34 open reading frame, omitting the region encoding the first 14 amino acid

residues, was cloned into the pSKB3 vector via the BamHI and HindIII restriction sites in

frame with the N-terminal His-tag sequence (Kaurov et al. 2022). The BL21(DE3)-derived C4

E. coli strain (Miroux and Walker 1996) was transformed with the plasmid. Upon reaching

0.6 OD_{600}, rMic34 production was induced with 1 mM isopropyl β-D-1-thiogalactopyranoside

(IPTG) for 3 h at 37ºC. Isolated inclusion bodies containing rMic34 were dissolved in 8 M

urea. This mixture was loaded into ÄKTAprime Plus liquid chromatography system

connected the 1 ml HisTrap column (GE Healthcare), followed by washing with lysis buffer (8

M urea, 2 mM Tris pH 7.2, 200 mM NaCl, 1mM EDTA) at a 1 ml/min flow rate. An increasing

gradient of elution buffer (lysis buffer with 400 mM imidazole) was introduced into the

column; rMic34 was eluted at 40% (v/v) elution buffer. Fractions in which rMic34 was

detected by Coomassie stained SDS-PAGE were pooled and loaded onto a HiLoad 16/600

Superdex 200 pg column (Cytiva) for size exclusion in 8 M urea, 150 mM NaCl, 10 mM Tris-

HCl, pH 8. Purest fractions were chosen based on SDS-PAGE analysis and the protein was

refolded through step-wise dialysis against refolding buffer (0.1% DDM, 150 mM NaCl, 20

mM HEPES, pH 7.4).

The *S. cerevisiae MIC19* coding sequence was cloned into the pGEX-6P-1 vector (Cytiva)

via the BamHI and SalI restriction sites with a C-terminal TEV cleavage site and His-tag, and

in frame with the N-terminal GST-tag. BL21(DE3) *E. coli* were transformed with the plasmid

and grown to an OD600 of 0.6 before induction with 1 mM IPTG for 20 h at 18°C. Cells were

lysed in lysis buffer (complete protease inhibitor cocktail without EDTA [Roche], 5% glycerol,

50 mM imidazole, 150 mM NaCl, 20 mM HEPES, pH 7.4), cleared by ultracentrifugation at

70000 × g for 25 min and loaded onto a His-Trap column (Cytiva) using an Äkta purifier

chromatography system. Unbound proteins were washed out with 10 column volumes (CV)

of lysis buffer and bound proteins eluted in 500 mM imidazole lysis buffer. Imidazole was

dialysed out in dialysis buffer (5% glycerol, 150 mM NaCl, 20 mM HEPES, pH 7.4). The GST

tag was removed via PreScission protease (Cytiva) and removal of GST and PreScission

protease on glutathione Sepharose (GE Healthcare) and the His-tag via incubation with TEV

protease in the presence of 14 mM β-mercaptoethanol and removal of His-tag and TEV

protease on nickel-nitrilotriacetic acid (NiNTA) agarose beads (Macherey-Nagel).

The *Rattus norvegicus* ENTH domain of Epsin was purified based on a previously published

protocol (Kroppen et al. 2021). In short, the protein was expressed with an N-terminal GST

tag from the pGEX-6-P2 vector in BL21(DE3) cells. Expression was induced in LB medium

with 1 mM IPTG for 3 h at 30°C. The cells were harvested and lysed in lysis buffer (complete

protease inhibitor cocktail without EDTA [Roche], 150 mM NaCl, 50 mM HEPES, 1 mM

EDTA, 1 mM DTT, pH 7.4). Lysed cleared by ultracentrifugation was loaded onto a GST-Trap

column (Cytiva) using an Äkta purifier chromatography system. Bound protein was eluted in

lysis buffer containing 10 mM reduced glutathione and peak fractions were concentrated.

The GST tag was cleaved with Precision Protease at 4°C overnight. To separate cleaved

GST, the buffer was then exchanged to low salt buffer (50 mM NaCl, 50 mM HEPES, pH 8)

using a Sephadex G-25 column (GE Healthcare) and the sample applied onto a HiTrap Q

column (Cytiva). The bound ENTH domain was eluted by steadily increasing the NaCl

concentration to 1 M. Concentrated peak fractions were further purified by size exclusion on

a HiLoad 16/600 Superdex 75 pg column (Cytiva) in 150 mM NaCl, 20 mM HEPES, pH 7.4,

and ENTH-containing fractions were pooled.

LUV preparation

Egg l-α-phosphatidylcholine (PC), egg l-α-phosphatidylethanolamine (PE), liver l-α-

phosphatidylinositol (PI), brain l-α-phosphatidylserine (PS), Heart cardiolipin (CL), and brain

L-α-phosphatidylinositol-4,5-bisphosphate (PI(4,5)P2) were purchased from Avanti Polar

Lipids, Inc. Folch bovine brain extract Fraction I was purchased from Sigma Aldrich. IMM

(48% PC/24%PE/16%PI/4%PS/8%CL), IMM-CL (56% PC/24%PE/16%PI/4%PS) and Folch

+ 5% PI(4,5)P2 lipid mixtures were dried under nitrogen flow to form thin lipid films. The IMM

phospholipid mixture used to make LUVs was informed by Horvath and Daum (2013). Then

they were dried in a desiccator for at least 3 h. Lipid films were hydrated in rehydration buffer

(150 mM NaCl and 20 mM HEPES buffer, pH 7.4). Resulting LUVs were subjected to

repeated freeze–thaw cycles and extruded through a 100-nm diameter polycarbonate filter

(Whatman). Protein was added to liposomes in 0.1% DDM rehydration buffer with a molar

protein/lipid ratio of 1:500.

Flotation assay, sodium carbonate extraction, proteinase K protection

Flotation assays were performed in nonionic Histodenz (Sigma-Aldrich) density gradients.

LUVs were placed on the bottom of the ultracentrifugation tubes, and a discontinuous

40/30/20/10/2% Histodenz density gradient was built up from high density to the lowest at

the top of the tube. LUVs were ultracentrifuged for 1 h at 273,000 × g and 4°C. Afterwards,

the gradient was dissected into nine equal fractions, precipitated with 10% TCA, and

analyzed by SDS-PAGE. For sodium carbonate extraction and proteinase K protection

assay, LUVs were flotated in a two-step density gradient consisting of 40% and 2%

Histodenz. LUVs and co-flotating proteins were collected from the interphase. For sodium

carbonate extraction, LUVs were incubated with 3 volumes of 100 mM $Na_2CO_{3'}$, pH 11, for

30 min at RT before ultracentrifugation at 167,000 × g for 30 min at 4°C to separate

membrane-bound from membrane-dissociated proteins. The supernatant containing lose

protein was TCA-precipitated before both fractions were analysed by SDS-PAGE. For

proteinase K protection assay, collected LUVs were washed in 8 volumes of rehydration

buffer (150 mM NaCl, 20 mM HEPES, pH 7.4), pelleted at 167,000 × g for 30 min at 4°C and

again resuspended in rehydration buffer before being incubated with 40 µg/ml proteinase K,

or buffer, in the presence or absence of 1% SDS for 15 min at RT. Then, 2 mM PMSF was

added and the samples incubated for 10 min at RT before being TCA-precipitated,

resuspended in SDS-loading dye containing 1 mM PMSF and analysed by SDS-PAGE. As a

control, proteins in absence of LUVs were treated in the same way.

Escherichia coli **culturing, transformation and subcellular fractionation**

Coding sequences (CDSs) of *T. brucei* Mic34, Mic40 and Mic24 (omitting the regions

encoding the first 14, 28, and 7 amino acid residues, respectively) were cloned into the

BamHI and HindIII restriction sites of the pMAL-P2X vector and transformed into BL21(DE3)

E. coli. The induction of MBP chimeras, which are targeted to the periplasm, was done as

previously (Tarasenko et al. 2017), except using LB broth. The subcellular fractionation of *E.*

coli was performed as before (Tarasenko et al. 2017), except the integral proteins remaining

in the cytoplasmic membrane of the spheroplasts were separated by 0.1M Na_2CO_3 pH 11.5

treatment like previously (Kaurov et al. 2018).

For co-expression, the same Mic40 CDS was cloned into pMAL-C2X, which bears

kanamycin resistance. BL21(DE3) *E. coli* were transformed concurrently with pMAL-P2X-

Mic34 and pMAL-C2X-Mic40 and *E. coli* hosting both were selected in LB supplemented

with 50 µg/ml kanamycin and 100 µg/ml ampicillin. *E.coli* lysates in were resolved on Mini-

PROTEAN TGX Stain-Free Precast gels, which allow visualization of abundant proteins by

UV illumination. Dual expression was verified by immunoblotting with antibody recognizing

Mic34 (Eichenberger et al. 2019) and Mic40. The latter rabbit antibody was produced by

Davids Biotechnologie GmbH (Regensburg, Germany) using denatured, His-tagged Mic40

recombinant protein as produced for LUV deformation assays.

Mic34 site directed mutagenesis and deletion of LBS motifs

A PCR reaction was performed on the pT7-3V5-Mic34 plasmid using oligos LUK0227 and

LUK0280, the former incorporating the mismatches necessary for introducing the desired

point mutations within the mitofilin domain of Mic34. ΔLBS1 and ΔLBS1+ 2 deletion mutants

were generated using either pT7-3V5-Mic34 or pMAL-p2X-Mic34 templates for a PCR

reaction with a primer downstream of LBS1 (ΔLBS1-R) and upstream of either LBS1

(ΔLBS1-F) or LBS2 (ΔLBS2-F), respectively. Primer sequences are:

LUK0280   5'-TATCGTTGAGGCGGGGAAAGCTCCCG-3'

LUK0227   5'-GCGCGTCCGAGATTATCGGCCGTTACATCCACAACAGCTGGTGA-3'

ΔLBS1-R   5'-TGAAGGGATGCTGTTTATGTACTTGAG-3'

ΔLBS1-F   5'-GCTCCCGCGGCTGAGCCCGTTC-3'

ΔLBS2-F   5'-TTCCGCACATCACTAAGCCCAACC-3'

These oligos containing 5'phosphates were ligated using T4 ligase (NEB) following

purification of the PCR product and DpnI digestion (to remove the original plasmid). The

resulting plasmid was verified by sequencing.

Immuno-electron tomography and transmission electron microscopy

The cell suspension was fixed in a mixture of 4% paraformaldehyde and 0.1%

glutaraldehyde in 0.1 M HEPES buffer for 1 h at RT. The cells were washed three times in

0.1 M HEPES buffer and embedded in 10% gelatine preheated at 37°C. The gelatine blocks

were immersed in 2.3 M sucrose for 72 h. The sample was placed on a stub and frozen in

liquid nitrogen, then cryo-sectioned into 80-100 nm sections, using a cryo-ultramicrotome

EM UC6 (Leica). The sections were placed on a formvar/carbon-coated nickel grid (300

Mesh) and allowed to thaw. The grids were then placed in 0.1 M HEPES buffer. After

treatment with blocking buffer (BB) composed of 1% fish skin gelatin dissolved in HEPES for

1 h, the grids were incubated with anti-MBP antibody diluted 1:20 in BB at RT for 30 min.

After washing in BB, the sections were incubated with the Protein A conjugated with 5 nm

gold nanoparticles (UMC, Utrecht) diluted 1:50 in BB and incubated for 45 min at RT.

Controls for nonspecific binding of the secondary antibody were performed by omitting the

primary antibody. Finally, the sections were washed in HEPES and dH_2O, stained with uranyl

acetate, embedded in methylcellulose, and inspected using a JEM-1400Flash transmission

electron microscope (JEOL).

Localization and analysis of immunogold labelling were performed utilizing electron

tomography. A JEM 2100F (JEOL) transmission electron microscope equipped with a direct

electron detector K2 Summit (Gatan) and controlled with the software package SerialEM

was used to acquire tilt series images in the range +57° to -63° with a tilt step of 1°. Electron

811 tomography reconstruction and filtering using nonlinear anisotropic diffusion were performed

812 in the software package IMOD (Mastronarde and Held 2017).

813 LUVs were imaged using a Talos L120C electron microscope equipped with a Ceta 16M

814 camera.

815 **Vesicle and nanoparticle counting**

816 Particle detection was performed by applying a density threshold to enhance the contrast

817 between the nanoparticles and the surrounding membrane. Particle counting was conducted

818 using the ImageJ software, where the occurrence of nanoparticles and vesicles was

819 analyzed with a precision limit of 5 nm. Vesicle counting was performed blind in 23 cells.

820 **Light microscopy and image analysis**

821 Images for 2D branching analysis were acquired on a BX63 widefield microscope (Olympus)

822 equipped with a 100× UPLSAPO NA 1.40 objective lens (Olympus), with an X-Cite excitation

823 light source (Excelitas), U-FUNA and U-FBWA fluorescence filter cubes (Olympus), and a

824 DP74 camera (Olympus) set to the 8-bit grayscale mode. Raw 8-bit grayscale images were

825 converted into the 32-bit format. Images were filtered using the difference of Gaussians s = 2

826 and 4. Background subtraction was done in Fiji (Schindelin et al. 2012) using the sliding

827 paraboloid and the radius of 20.0 pixels. Image thresholding was done using the triangle

828 method. Branching quantification and analysis was performed in Fiji using the 2D analysis

829 option of the Mitochondria Analyzer plugin (Chaudhry et al. 2019). We measured

830 mitochondria branching and length for 100 complete cells for each condition. All statistical

831 analyses were performed using the GraphPad Prism software, using an unpaired two-tailed

832 Student's t-test. A p-value of less than 0.05 was considered statistically significant.

833 Images for fragmentation analysis and 3D reconstruction were acquired on a FV3000 laser

834 scanning confocal microscope (Olympus). The microscope was equipped with a UAPON

835 100× TIRF NA 1.49 objective lens (Olympus), 405 nm and 488 nm lasers for excitation

836 (Coherent), quadband dichroic beamsplitter TRF89902 ET (Chroma), and high-sensitivity

GaAsP spectral detectors with detection windows set to 420-470 nm and 500-540 nm,

respectively. Z-stacks were acquired using Nyquist sampling and processed using the 3D

constrained iterative deconvolution with the maximum likelihood algorithm and the

theoretical point spread function in cellSens Dimension (Olympus). Creation of 3D models

was done in Imaris 10.0 (Oxford Instruments) using manual thresholding for surface creation

and removal of small objects. For fragmentation analysis maximum intensity projections

were taken from Z-stacks and processed in Fiji using the Mitochondria Analyzer plugin.

Single cells were cropped from the images and subjected to image thresholding using the

default parameters in Mitochondria Analyzer with the Block size of 1.85 micron and the C-

value of 3. The number of non-connected fragments for individual cells was counted from

binary images. We measured 35-50 cells for each condition. All statistical analyses were

performed using the GraphPad Prism software, using an unpaired two-tailed Mann-Whitney

test in GraphPad Prism. A p-value of less than 0.05 was considered statistically significant.

ACKOWLEDGEMENTS

We would like to thank André Schneider (University of Bern) and Alena Zíková (Biology

Centre, Czech Academy of Sciences) for antibodies (AS: Tim9, Mic34; AZ: Rieske, COXIV,

ATPβ, HSP70) plus Lawrence Rudy Cadena (Heinrich Heine University) for his input and

Julius Lukeš (BC CAS) for his support and Erv1 antibody. This work was supported by

Czech Science Foundation grant 23-07674S to HH. ME was supported by the European

Union under the LERCO project number CZ.10.03.01/00/22_003/0000003 via the

Operational Programme Just Transition. TB and MT were supported by the Czech

BioImaging grant from the ERD Funds of the Czech Ministry of Education. BTK was

supported by the Boehringer Ingelheim Fonds PhD Fellowship. BTK and MM were

supported by the Deutsche Forschungsgemeinschaft (SFB-1638/1 – 511488495 – P08 and

FOR-2848 – 401510699 - P05). The authors gratefully acknowledge Marek Vrbacký at the

Proteomics Service Laboratory at the Institute of Physiology (supported by RVO, ID

67985823) and Institute of Molecular Genetics (supported by RVO, ID 68378050) of the

Czech Academy of Sciences. The authors thank the Laboratory of Microscopy and Histology

at the Biology Centre, Czech Academy of Sciences for provision of instruments. We would

like to acknowledge access to the infrastructure and support provided by the Cryo-EM

Network at the Heidelberg University (HDcryoNet), which is funded and supported by the

German Research Founda1on (DFG), the Federal Ministry of Educa1on and Research

(BMBF) and the Ministry of Science Baden-Wu☐rOemberg, among others, within the

framework of the Excellence Strategy of the Federal and State Governments of Germany.

DATA AVAILABILITY

The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE partner repository with the dataset identifier PXD059459. Multiple

sequence alignments used in the analyses reported in the study are available in the

Figshare data repository (10.6084/m9.figshare.30001090).

REFERENCES

Aaltonen MJ, Friedman JR, Osman C, Salin B, di Rago JP, Nunnari J, Langer T, Tatsuta T. 2016. MICOS and phospholipid transfer by Ups2-Mdm35 organize membrane lipid synthesis in mitochondria. J. Cell Biol. 213:525-534.

Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucl. Acids Res. 25:3389-3402.

Alvarez-Jarreta J, Amos B, Aurrecoechea C, Bah S, Barba M, Barreto A, Basenko EY, Belnap R, Blevins A, Böhme U, et al. 2024. VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource center in 2023. Nucl. Acids Res. 52:D808-D816.

Barbot M, Jans DC, Schulz C, Denkert N, Kroppen B, Hoppert M, Jakobs S, Meinecke M. 2015. Mic10 oligomerizes to bend mitochondrial inner membranes at cristae junctions. Cell Metab. 21:756-763.

Barbot M, Meinecke M. 2016. Reconstitutions of mitochondrial inner membrane remodeling. J. Struct. Biol. 196:20-28.

Benning FMC, Bell TA, Nguyen TH, Syau D, Connell LB, Liao Y-T, Keating MP, Coughlin M, Nordstrom AEH, Ericsson M, et al. 2025. Ancestral sequence reconstruction of Mic60 reveals a residue signature supporting respiration in yeast. Protein Sci. 34:e70207.

Benz C, Müller N, Kaltenbrunner S, Váchová H, Vancová M, Lukeš J, Varga V, Hashimi H. 2022. Kinetoplastid-specific X2-family kinesins interact with a kinesin-like pleckstrin homology domain protein that localizes to the trypanosomal microtubule quartet. Mol. Microbiol. 118:155-174.

Bílý T, Sheikh S, Mallet A, Bastin P, Pérez-Morga D, Lukeš J, Hashimi H. 2021. Ultrastructural Changes of the Mitochondrion During the Life Cycle of *Trypanosoma brucei*. J. Eukaryot. Microbiol. 68:e12846.

Bock-Bierbaum T, Funck K, Wollweber F, Lisicki E, von der Malsburg K, von der Malsburg A, Laborenz J, Noel JK, Hessenberger M, Jungbluth S, et al. 2022. Structural insights into crista junction formation by the Mic60-Mic19 complex. Sci. Adv. 8:eabo4946.

Burki F, Roger AJ, Brown MW, Simpson AGB. 2020. The New Tree of Eukaryotes. Trends Ecol. Evol. 35:43-55.

Cadena LR, Gahura O, Panicucci B, Zíková A, Hashimi H, Blader Ira J. 2021. Mitochondrial Contact Site and Cristae Organization System and F1FO-ATP Synthase Crosstalk Is a Fundamental Property of Mitochondrial Cristae. mSphere 6:e00327-00321.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972-1973.

Cavalier-Smith T. 2016. Higher classification and phylogeny of Euglenozoa. Eur. J. Protistol. 56:250-276.

Chaudhry A, Shi R, Luciani DS. 2019. A pipeline for multidimensional confocal analysis of mitochondrial morphology, function, and dynamics in pancreatic β-cells. Am. J. Physiol. Endocrinol. Metab. 318:E87-E101.

Colina-Tenorio L, Horten P, Pfanner N, Rampelt H. 2020. Shaping the mitochondrial inner membrane in health and disease. J. Intern. Med. 287:645-664.

Daumke O, van der Laan M. 2025. Molecular machineries shaping the mitochondrial inner membrane. Nat. Rev. Mol. Cell Biol.:https://doi.org/10.1038/s41580-41025-00854-z.

Eddy SR. 2011. Accelerated Profile HMM Searches. PLoS Comput. Biol. 7:e1002195.

Eichenberger C, Oeljeklaus S, Bruggisser J, Mani J, Haenni B, Kaurov I, Niemann M, Zuber B, Lukeš J, Hashimi H, et al. 2019. The highly diverged trypanosomal MICOS complex is organized in a nonessential integral membrane and an essential peripheral module. Mol. Microbiol. 12:1731-1743.

Flaspohler John A, Jensen Bryan C, Saveria T, Kifer Charles T, Parsons M. 2010. A Novel Protein Kinase Localized to Lipid Droplets Is Required for Droplet Biogenesis in Trypanosomes. Eukaryot. Cell 9:1702-1710.

Gleisner M, Kroppen B, Fricke C, Teske N, Kliesch T-T, Janshoff A, Meinecke M, Steinem C.
2016. Epsin N-terminal Homology Domain (ENTH) Activity as a Function of Membrane
Tension. J. Biol. Chem. 291:19953-19961.
Haindrich AC, Boudová M, Vancová M, Peña-Diaz P, Horáková E, Lukeš J. 2017. The
intermembrane space protein Erv1 of *Trypanosoma brucei* is essential for mitochondrial Fe-
S cluster assembly and operates alone. Mol. Biochem. Parasitol. 214:47-51.
Hammond MJ, Nenarokova A, Butenko A, Zoltner M, Dobáková EL, Field MC, Lukeš J.
2020. A Uniquely Complex Mitochondrial Proteome from Euglena gracilis. Mol Biol Evol
37:2173-2191.
Hashimi H. 2019. A parasite's take on the evolutionary cell biology of MICOS. PLoS Pathog.
15:e1008166.
Hessenberger M, Zerbes RM, Rampelt H, Kunz S, Xavier AH, Purfurst B, Lilie H, Pfanner N,
van der Laan M, Daumke O. 2017. Regulated membrane remodeling by Mic60 controls
formation of mitochondrial crista junctions. Nat. Commun. 8:15258.
Hirumi H, Hirumi K. 1989. Continuous cultivation of Trypanosoma brucei blood stream forms
in a medium containing a low concentration of serum protein without feeder cell layers. J.
Parasitol. 75:985-989.
Horvath SE, Daum G. 2013. Lipids of mitochondria. Prog. Lipid Res. 52:590-614.
Huerta-Cepas J, Serra F, Bork P. 2016. ETE 3: Reconstruction, Analysis, and Visualization of
Phylogenomic Data. Mol. Biol. Evol. 33:1635-1638.
Hughes L, Borrett S, Towers K, Starborg T, Vaughan S. 2017. Patterns of organelle ontogeny
through a cell cycle revealed by whole-cell reconstructions using 3D electron microscopy. J.
Cell Sci. 130:637-647.
Huynen MA, Muhlmeister M, Gotthardt K, Guerrero-Castillo S, Brandt U. 2016. Evolution and
structural organization of the mitochondrial contact site (MICOS) complex and the
mitochondrial intermembrane space bridging (MIB) complex. Biochim. Biophys. Acta
1863:91-101.
Jakob M, Hoffmann A, Amodeo S, Peitsch C, Zuber B, Ochsenreiter T. 2016. Mitochondrial
growth during the cell cycle of *Trypanosoma brucei* bloodstream forms. Sci. Rep. 6:36265.
Johnston HE, Yadav K, Kirkpatrick JM, Biggs GS, Oxley D, Kramer HB, Samant RS. 2022.
Solvent Precipitation SP3 (SP4) Enhances Recovery for Proteomics Sample Preparation
without Magnetic Beads. Anal. Chem. 94:10320-10328.
Katoh K, Rozewicki J, Yamada KD. 2019. MAFFT online service: multiple sequence
alignment, interactive sequence choice and visualization. Brief. Bioinform. 20:1160-1166.
Kaurov I, Heller J, Deisenhammer S, Potěšil D, Zdráhal Z, Hashimi H. 2022. The essential
cysteines in the CIPC motif of the thioredoxin-like Trypanosoma brucei MICOS subunit
TbMic20 do not form an intramolecular disulfide bridge in vivo. Mol. Biochem. Parasitol.
248:111463.
Kaurov I, Vancová M, Schimanski B, Cadena LR, Heller J, Bílý T, Potěšil D, Eichenberger C,
Bruce H, Oeljeklaus S, et al. 2018. The diverged trypanosome MICOS complex as a hub for
mitochondrial cristae shaping and protein import. Curr. Biol. 28:3393-3407.
Kostygov AY, Karnkowska A, Votýpka J, Tashyreva D, Maciszewski K, Yurchenko V, Lukeš J.
2021. Euglenozoa: taxonomy, diversity and ecology, symbioses and viruses. Open Biol.
11:200407.
Kroppen B, Teske N, Yambire KF, Denkert N, Mukherjee I, Tarasenko D, Jaipuria G,
Zweckstetter M, Milosevic I, Steinem C, et al. 2021. Cooperativity of membrane-protein and
protein–protein interactions control membrane remodeling by epsin 1 and affects clathrin-
mediated endocytosis. Cell. Mol. Life Sci. 78:2355-2370.
Kühlbrandt W. 2019. Structure and mechanisms of F-type ATP synthases. Annu. Rev.
Biochem. 88:515-549.
Lax G, Kolisko M, Eglit Y, Lee WJ, Yubuki N, Karnkowska A, Leander BS, Burger G, Keeling
PJ, Simpson AGB. 2021. Multigene phylogenetics of euglenids based on single-cell
transcriptomics of diverse phagotrophs. Mol. Phylogen. Evol. 159:107088.
Letunic I, Bork P. 2024. Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic
tree display and annotation tool. Nucl. Acids Res. 52:W78-W82.

Mastronarde DN, Held SR. 2017. Automated tilt series alignment and tomographic
reconstruction in IMOD. J. Struct. Biol. 197:102-113.
Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023.
UCSF ChimeraX: Tools for structure building and analysis. Protein Sci. 32:e4792.
Michaud M, Gros V, Tardif M, Brugiere S, Ferro M, Prinz WA, Toulmay A, Mathur J, Wozny
M, Falconet D, et al. 2016. AtMic60 is involved in plant mitochondria lipid trafficking and is
part of a large complex. Curr. Biol. 26:627-639.
Miroux B, Walker JE. 1996. Over-production of Proteins in *Escherichia coli*: Mutant Hosts
that Allow Synthesis of some Membrane Proteins and Globular Proteins at High Levels. J.
Mol. Biol. 260:289-298.
Mordas A, Tokatlidis K. 2015. The MIA pathway: a key regulator of mitochondrial oxidative
protein folding and biogenesis. Acc. Chem. Res. 48:2191-2199.
Muñoz-Gómez SA, Cadena LR, Gardiner AT, Leger MM, Sheikh S, Connell LB, Bilý T,
Kopejtka K, Beatty JT, Koblížek M, et al. 2023. Intracytoplasmic-membrane development in
alphaproteobacteria involves the homolog of the mitochondrial crista-developing protein
Mic60. Curr. Biol. 33:1099-1111.e1096.
Muñoz-Gómez SA, Slamovits CH, Dacks JB, Baier KA, Spencer KD, Wideman JG. 2015.
Ancient homology of the mitochondrial contact site and cristae organizing system points to
an endosymbiotic origin of mitochondrial cristae. Curr. Biol. 25:1489-1495.
Muñoz-Gomez SA, Wideman JG, Roger AJ, Slamovits CH. 2017. The origin of mitochondrial
cristae from alphaproteobacteria. Mol. Biol. Evol. 34:943-956.
Niemann M, Wiese S, Mani J, Chanfon A, Jackson C, Meisinger C, Warscheid B, Schneider
A. 2013. Mitochondrial outer membrane proteome of Trypanosoma brucei reveals novel
factors required to maintain mitochondrial morphology. Mol. Cell. Proteom. 12:515-528.
Pánek T, Eliáš M, Vancová M, Lukeš J, Hashimi H. 2020. Returning to the Fold for Lessons
in Mitochondrial Crista Diversity and Evolution. Curr. Biol. 30:R575-R588.
Peikert CD, Mani J, Morgenstern M, Kaser S, Knapp B, Wenger C, Harsman A, Oeljeklaus
S, Schneider A, Warscheid B. 2017. Charting organellar importomes by quantitative mass
spectrometry. Nat. Commun. 8:15272.
Poon SK, Peacock L, Gibson W, Gull K, Kelly S. 2012. A modular and optimized single
marker system for generating *Trypanosoma brucei* cell lines expressing T7 RNA polymerase
and the tetracycline repressor. Open Biol. 2:110037.
Prokopchuk G, Butenko A, Dacks JB, Speijer D, Field MC, Lukeš J. 2023. Lessons from the
deep: mechanisms behind diversification of eukaryotic protein complexes. Biol. Rev. Camb.
Philos. Soc. 98:1910-1927.
Rabl R, Soubannier V, Scholz R, Vogel F, Mendl N, Vasiljev-Neumeyer A, Korner C, Jagasia
R, Keil T, Baumeister W, et al. 2009. Formation of cristae and crista junctions in mitochondria
depends on antagonism between Fcj1 and Su e/g. J. Cell Biol. 185:1047-1063.
Rampelt H, Wollweber F, Licheva M, de Boer R, Perschil I, Steidle L, Becker T, Bohnert M,
van der Klei I, Kraft C, et al. 2022. Dual role of Mic10 in mitochondrial cristae organization
and ATP synthase-linked metabolic adaptation and respiratory growth. Cell Rep. 38:110290.
Richter DJ, Berney C, Strassert JFH, Poh Y-P, Herman EK, Muñoz-Gómez SA, Wideman
JG, Burki F, de Vargas C. 2022. EukProt: A database of genome-scale predicted proteins
across the diversity of eukaryotes. Peer Community J. 2:e56.
Roger AJ, Munoz-Gomez SA, Kamikawa R. 2017. The origin and diversification of
mitochondria. Curr. Biol. 27:1177-1192.
Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S,
Rueden C, Saalfeld S, Schmid B, et al. 2012. Fiji: an open-source platform for biological-
image analysis. Nat. Methods 9:676-682.
Smith JT, Singha UK, Misra S, Chaudhuri M. 2018. Divergent Small Tim Homologues Are
Associated with TbTim17 and Critical for the Biogenesis of TbTim17 Protein Complexes in
Trypanosoma brucei. mSphere 3:e00204-00218.
Söding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein homology
detection and structure prediction. Nucl. Acids Res. 33:W244-W248.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. Bioinformatics 30:1312-1313.
Stephan T, Brüser C, Deckers M, Steyer AM, Balzarotti F, Barbot M, Behr TS, Heim G,
Hübner W, Ilgen P, et al. 2020. MICOS assembly controls mitochondrial inner membrane
remodeling and crista junction redistribution to mediate cristae formation. EMBO J.
39:e104105.
Tarasenko D, Barbot M, Jans DC, Kroppen B, Sadowski B, Heim G, Mobius W, Jakobs S,
Meinecke M. 2017. The MICOS component Mic60 displays a conserved membrane-bending
activity that is necessary for normal cristae morphology. J. Cell Biol. 216:889-899.
Turra GL, Liedgens L, Sommer F, Schneider L, Zimmer D, Vilurbina Perez J, Koncarevic S,
Schroda M, Muhlhaus T, Deponte M. 2021. In Vivo Structure-Function Analysis and Redox
Interactomes of Leishmania tarentolae Erv. Microbiol. Spectr. 9:e0080921.
Van Laar VS, Berman SB, Hastings TG. 2016. Mic60/mitofilin overexpression alters
mitochondrial dynamics and attenuates vulnerability of dopaminergic cells to dopamine and
rotenone. Neurobiol. Dis. 91:247-261.
Varabyova A, Topf U, Kwiatkowska P, Wrobel L, Kaus-Drobek M, Chacinska A. 2013. Mia40
and MINOS act in parallel with Ccs1 in the biogenesis of mitochondrial Sod1. FEBS J.
280:4943-4959.
von der Malsburg K, Muller JM, Bohnert M, Oeljeklaus S, Kwiatkowska P, Becker T,
Loniewska-Lwowska A, Wiese S, Rao S, Milenkovic D, et al. 2011. Dual role of mitofilin in
mitochondrial membrane organization and protein biogenesis. Dev. Cell 21:694-707.
Wenger C, Oeljeklaus S, Warscheid B, Schneider A, Harsman A. 2017. A trypanosomal
orthologue of an intermembrane space chaperone has a non-canonical function in
biogenesis of the single mitochondrial inner membrane protein translocase. PLoS Pathog.
13:e1006550.
Wheeler RJ. 2021. A resource for improved predictions of Trypanosoma and Leishmania
protein three-dimensional structure. PLoS One 16:e0259871.
Zíková A. 2022. Mitochondrial adaptations throughout the Trypanosoma brucei life cycle. J.
Eukaryot. Microbiol. 69:e12911.

1068

1069

1070

Fig. 1. The N-termini of Mic34 and Mic40 retain a mitofilin domain.

(A) Schematic representation of HHpred results using representative profile HMMs. The

searches detected statistically significant matches (the respective values indicated in the

figure) between both Mic34 and Mic40 and the Pfam family PF09731.14 (*i.e.*, Mic60), with

the specific matching regions delimited by the grey zones. The three proteins are drawn to

scale according to the reference proteins for each HMM (arbitrarily Mic34 and Mic40 from *T.*

brucei and *Dictyostelium discoideum* Mic60). For Mic60 (PF09731.14) three different

structural/functional regions are delimited, including a TMD (TM), a coiled-coil region

(boundaries as reported (Hessenberger et al. 2017) for *C. thermophila* Mic60), and the

mitofilin domain as recently redefined (Benning et al. 2025).

(B) The AlphaFold2 predicted structures of the putative mitofilin domains of Mic34 and Mic40

(light purple) aligned to the predicted mitofilin domain of discoban *N. gruberi* Mic60 (alpha

helices colored as in B). The root-mean-square distance (RMSD) of the alignments is given

in red below the model. Scale bar, 10 Å. Dashed-boxed insets show alignment of conserved

LBS1 aromatic amino acid. Scale bar, 1 Å. Major euglenozoan clades depicted as in

(Kostygov et al. 2021): Eugl., Euglenida; Dipl., Diplonemea; Kin., Kinetoplastea.

(C) Map of the mitofilin alpha-helical motifs from yeast and *N. gruberi* (Ng) Mic60, plus

representative Mic34 and Mic40 from the three clades in A. Mitofilin alpha helices are color

coded and correspond to names on top. The conserved aromatic amino acid residue of

LBS1 is color coded based on the key below the uppermost LBS1 (F, phenylalanine; W,

tryptophan; Y, tyrosine). Abbreviations: α, alpha helix; Ct, *C. thermophilum*; Eg, *E. gracilis*;

Hp, *H. phaeocysticola*; LBS, lipid binding site; Sc, *S cerevisiae*; Tb, *T. brucei*. The PDB

accession number for the solved CtMic60 mitofilin domain structure (Bock-Bierbaum et al.

2022), in which LBS2 was necessarily deleted, is given in parentheses.

1096

Fig. 2. Mitochondrial remodeling by Mic34 and Mic40 seems to be coupled with their involvement in the mitochondrial intermembrane space assembly pathway.

(A) Representative 3D reconstruction of mitochondria in wild type (WT) and Mic34 knockdown *T. brucei* cells at time points (TPs) 3 and 5 days after Dox induction. Mitochondria labeled by an anti-HSP70 antibody are in green; the nucleus and kinetoplast DNA (arrowhead) are in blue. Scale bar, 1 µm.

(B) Scatter plot showing the number of mitochondrial branches per cell (n=100) in WT plus Mic34, Mic40, and ERV1 RNAi at TP 3 and 5 of doxycycline induction. Data are presented as individual values with their mean and standard deviation indicated by bar and whiskers. Statistical significance: **p = 0.01 and ****p < 0.0001.

(C) Schematic representation of the MICOS complex and ERV1. Peripheral MICOS subunits whose depletomes were determined are colored in yellow. Integral membrane components are shown in gray. The lipid bilayer is represented as a double gray line. The ERV1 dimer is depicted in red.

(D) Volcano plots of Mic40 depletomes measured at TP3 (top row) and TP5 (bottom) of RNAi induction. Each row shows the same Mic40 TP3 and TP5 depletome data colored to highlight overlap with the depletomes of (left to right) Mic34 (Eichenberger et al. 2019), Mic20 (Kaurov et al. 2018; Kaurov et al. 2022), and Erv1 (Peikert et al. 2017). Points corresponding to MICOS subunits are individually labelled as in C. X-axis, $\log_2$-transformed fold change of Mic40 RNAi over WT; y-axis, -log of P-value of measurement from triplicates.

Fig. 3. Mic34 binds to phospholipid bilayers *in vitro*.

(A) Carbonate extraction of Mic34 loaded large unilamellar vesicles (LUVs) with 25 or 75 mM sodium carbonate. Supernatant, S; pellet, P.

(B) Proteinase K (PK) protection assay of LUVs loaded with Mic34 or Mic19 and Folch liposomes loaded with ENTH visualized with Coomassie Brilliant Blue staining after SDS-PAGE.

(C) Flotated LUVs with and without (-CL) cardiolipin separated on aa noncontinuous

Histodenz density gradient (schema on left) loaded with recombinant Mic34. Compare

banding pattern with LUVs flotated alone (right).

(D) Histodenz density gradient fractions resolved by SDS-PAGE and stained with

Coomassie Brilliant Blue or transferred to a membrane for incubation with an anti-His

antibody (Mic34 with and without CL).

(E) Negatively stained LUVs loaded with Mic34. Scale bar, 100 nm.

(F) Negatively stained Folch + 5% PI(4,5)P2 liposomes incubated with the membrane

remodeling protein ENTH that did (+DDM) or did not (DDM-) undergo prior solubilization with

0.1% DDM. Scale bar, 200 nm.

1134

Fig. 4. Heterologously expressed Mic34 and Mic40 remodels the cytoplasmic

membrane of *E. coli* in an LBS-dependent manner upon periplasm export.

(A) Schema (left) of periplasting by osmotic shock to separate spheroplast (*i.e. the*

compartment bound by the cytoplasmic membrane) and periplasm fractions. The former was

fractioned by carbonate extraction to separate integral and peripheral membrane proteins.

Western blots on right show the localization of MBP-Mic34, MBP-Mic40, and MBP alone in

E. coli, as detected with an anti-MBP antibody.

(B) Representative transmission electron micrographs of *E. coli* when MBP expression is not

(left) and is (right) induced. Scale bar, 200 µm.

(C) Representative electron tomograms of immunogold-labeled MBP-Mic34 and MBP-Mic40

using an anti-MBP antibody. Full cells are shown in the main image with dashed boxes

corresponding to magnified inset images that depict either unmarked or traced images. In

the latter, pink line traces the cytoplasmic membrane, red the outer membrane, and orange

the vesicles; yellow circles highlight electron-dense gold nanoparticles. Scale bars: main

image, indicated above bar; inset, 50 nm. Insets report the depth in the Z-dimension of the

depicted plane of the tomogram when applicable.

(D) Bar graph reporting the total number of vesicles observed in TEM images (n=22) of cells

expressing MBP tagged Mic34, Mic40, Mic34ΔLBS1+2, and of WT cells. The number of

vesicles arising in Mic34 and Mic40 expressing cells. Vesicles directly associated with

nanoparticles are shaded in light purple with their proportion shown.

(E) Western blot probed using an anti-MBP antibody to show MBP-Mic34 and MBP-

Mic34ΔLBS1+2 expression (left) and a representative TEM image of *E. coli* expressing

MBP-Mic34ΔLBS1+2. Scale bar= 200 nm).

1158

Fig. 5. **Mic34 overexpression induces reticulation of the tubular mitochondrion of**

bloodstream form *T. brucei*.

(A) Growth of Mic34 overexpression (O/E) as measured by cell density plotted

logarithmically under different doxycycline (Dox) concentrations (left). Solid line with circles,

1000 ng/ml; triangle,100 ng/mL; square,10 ng/ml. Dashed line, 0 ng/ml Dox. Standard

deviation at each time point (n = 3) is not visible at this scale. Right, western blots probed

with anti-Mic34 (detecting both endogenous Mic34 and V5-tagged Mic34) and anti-V5 tag

antibodies, with fluorescently detected proteins used as a loading control (load).

(B) Left, growth of cells overexpressing Mic34 WT (black circle), ΔLBS1+2 (purple square),

ΔLBS1 (red triangle) by induction with 1000 ng/ml Dox (dotted line) compared to uninduced

cells (solid lines). Standard deviation at each time point (n = 3) is not visible at this scale.

Right, western blot probed with anti-V5 to show expression of Mic34 WT, ΔLBS1+2, and

ΔLBS1 with (+) and without (–) 100 ng/ml Dox induction.

(C) Representative 3D reconstructions of mitochondria in parental and WT Mic34 O/E cells.

Mitochondria labeled by anti-HSP70 antibody are in green; the DAPI-stained nucleus and

kinetoplast DNA (arrowhead) are in blue. Scale bar, 1 μm. Note all cells are 1N1K cells.

(D) Quantification of mitochondrial branching in 100 parental and Mic34 WT O/E (36 hours

after Dox induction) cells. Statistical significance: ****, $p < 0.0001$.

(E) Schema of the *T. brucei* bloodstream cell cycle under normal conditions (yellow

background with black arrows) and upon Dox-induced Mic34 overexpression (↑Mic34 O/E)

(red arrow). Mitochondria are in blue and the number of nuclei (N) and kinetoplast DNA (K) is

given below each cell. Dashed arrows indicate a pathway to defective 1N2K cells under

normal Mic34 expression and overexpression.

(F) Bar graph showing the percentage of cells with reticulated mitochondria from different

cell cycle stages depicted in E in parental and WT Mic34 overexpression cells.

(G) Growth of *T. brucei* overexpressing Mic34^{R170DF174D} with 1 μg/ml of doxycycline (Dox+)

compared to uninduced (Dox-) cell lines. Standard deviation whiskers at each time point (n =

3) are faintly visible at 3 days in this scale. This overlays the growth 1 μg/ml of doxycycline-

induced Mic34 WT and its uninduced counterpart from A (grey). Right, western blots probed

with anti-V5 tag antibody (top) and loading control (bottom; load) as in A.

(H) Quantification of mitochondrial fragmentation in Mic34 WT and Mic34^{R170DF174D}

overexpressing (+, 36 hours after Dox induction) and non-induced (-) cells. Statistical

significance: *, $p < 0.05$. Number of cells counted: WT-, 42; WT+, 37; Mic34^{R170DF174D} (both +

and -), 47.

1193

1194

1195

A

KDa

	Peri	Sphero	Integral	Periph
76					Mic34
82					Mic40
42					MBP

B

Non-induced   MBP-induced

C   MBP-Mic34

Z = - 2.2 nm
Z = - 2.8 nm
Z = - 8.4 nm
Z = - 17.9 nm
Z = - 7.8 nm
Z = - 4.0 nm
Z = - 7.8 nm

500 nm

MBP-Mic40

Z = - 4.5 nm   Z = - 6.2 nm   Z = + 5 nm

200 nm

D

Vesicle number

- Direct association
- No association

61.1%   67.7%

Mic34   Mic40   ΔLBS1+2   WT

E

	Mic34	ΔLBS1+2
76		
60			MBP

MBP-ΔLBS1+2