Title: Functional genomics of trypanotolerant and trypanosusceptible cattle infected with

Trypanosoma congolense across multiple time points and tissues

Short title: Functional genomics of trypanotolerant and trypanosusceptible cattle

Authors and Affiliations: Gillian P. McHugo[1], James A. Ward[1], John A. Browne[1], Grace M.

O'Gorman[2], Kieran G. Meade[1,3], Emmeline W. Hill[1], Thomas J. Hall[1], David E.

MacHugh[1,3,4*]

[1] UCD School of Agriculture and Food Science, University College Dublin, Belfield, Dublin,

Ireland.

[2] UK Agri-Tech Centre, Innovation Centre, York Science Park, York, UK.

[3] UCD Conway Institute of Biomolecular and Biomedical Research, University College

Dublin, Dublin, Ireland.

[4] UCD One Health Centre, University College Dublin, Dublin, Ireland.

* Corresponding author

Email: david.machugh@ucd.ie

Abstract

Human African trypanosomiasis (HAT), or sleeping sickness, is a neglected tropical disease caused by infection with trypanosome parasites (*Trypanosoma* spp.). These are transmitted by infected tsetse flies (*Glossina* spp.) and cause a similar disease in animals, known as African animal trypanosomiasis (AAT). AAT is one of the largest constraints to livestock production in sub-Saharan Africa and causes a financial burden of approximately $4.5 billion annually. Some African *Bos taurus* cattle populations have an important evolutionary adaptation known as trypanotolerance, a genetically determined tolerance of infection by trypanosome parasites (*Trypanosoma* spp.). Trypanotolerant African *B. taurus* N'Dama and trypanosusceptible *Bos indicus* Boran cattle responded in largely similar ways during trypanosome infection when gene expression was examined using blood, liver, lymph node, and spleen samples with peaks and troughs of gene expression differences following the cyclic pattern of parasitaemia exhibited during trypanosome infection. However, differences in response to infection between the two breeds were reflected in differential expression of genes related to the immune system such as those encoding antimicrobial peptides and cytokines, including, for example, the antimicrobial peptide encoding genes *LEAP2*, *CATHL3*, *DEFB4A*, and *S100A7* and the cytokine genes *CCL20*, *CXCL11*, *CXCL13*, *CXCL16*, *CXCL17*, *IL33*, and *TNFSF13B*. In addition, transcriptional profiling of peripheral blood identified expression differences in genes relating to coagulation and iron homeostasis, which supports the hypothesis that the dual control of parasitaemia and the anaemia resulting from the innate immune response to trypanosome parasites is key to trypanotolerance and provide new insights into the molecular mechanisms underlying this phenomenon.

Author Summary

Trypanosome parasites are transmitted by infected tsetse flies and cause the neglected tropical disease human African trypanosomiasis (HAT) and the similar African animal trypanosomiasis (AAT), which is one of the largest impediments to livestock production in sub-Saharan Africa. Taurine (*Bos taurus*) and indicine (*Bos indicus*) cattle shared a common ancestor more than 150,000 years ago, and in the intervening period significant genomic differences have evolved between the two groups. Importantly, several African *B. taurus* populations have evolved an adaptation known as trypanotolerance, a genetically determined tolerance of infection by trypanosome parasites. Trypanotolerant African taurine and trypanosusceptible indicine cattle responded in largely similar ways during trypanosome infection when gene expression was examined using blood, liver, lymph node, and spleen samples with peaks and troughs of gene expression differences following the cyclic pattern of the number of trypanosome parasites in the blood. Transcriptional profiling of these tissues highlighted genes related to the multiple facets of the immune system; notably, for peripheral blood, differences observed for genes relating to coagulation and iron homeostasis support the existing hypothesis that control of both parasite number and anaemia is an important feature of the trypanotolerance trait and provide new insights into the molecular mechanisms underlying this phenomenon.

Introduction

Trypanosomiasis is prevalent in humid and semi-humid regions of Africa and is a wasting disease caused by parasitic protozoa of the genus *Trypanosoma* which are transmitted by biting insect vectors such as tsetse flies (*Glossina* spp.) [1]. Human African trypanosomiasis (HAT), caused by *T. brucei gambiense* or *T. brucei rhodesiense* infection, is also known as sleeping sickness and is classed as a neglected tropical disease (NTD) by the World Health Organisation [2]. Trypanosomes also infect animals and cause particularly severe disease in domesticated animals—African animal trypanosomiasis (AAT)—for which the symptoms, as in HAT, include fever, severe weight loss, and anaemia [1, 3, 4]. As the disease progresses, animals weaken and become paralysed and unfit for work [1]. AAT is caused by a wider range of trypanosome parasites, most commonly *T. congolense, T. vivax*, and *T. brucei* ssp., although the prevalence and distribution of trypanosome species varies and co-infections are possible [5, 6]. In addition, domestic animals can also be infected with the human-infective trypanosome species, for which they act as a reservoir [5, 7].

Animal models, including mice, have long been used to study human-infective trypanosomes leading to important insights into host-trypanosome interactions [8-10]. The usefulness of these animal models has been improved by the differential host tolerance of, or susceptibility to, trypanosome infection that has been found between different cattle breeds, mouse strains, and human populations, although the underlying genes differ between host species [11-14]. Conversely, while it is much less common, humans may also be infected with trypanosome species more frequently found in animals, resulting in atypical human trypanosomiasis [5, 15]. With climate change and increasing globalisation there is growing concern that this atypical human trypanosomiasis may represent an emerging neglected zoonotic disease that underscores the importance of a One Health approach to both human and animal trypanosomiasis [16].Trypanosomiasis has a limiting effect on sub-Saharan livestock production, because even with application of trypanocidal drugs, the susceptibility

of most cattle to trypanosomiasis makes production economically unsustainable in many

regions on the African continent [17].

Cattle are significant components of rural economies and livelihoods in Africa. They

provide milk, meat, fertiliser and traction and represent mobile assets that serve as a financial

buffer for poor families, particularly for pastoralists and women [18, 19]. There are

approximately 150 breeds of indigenous cattle in sub-Saharan Africa and African cattle

represent a complex mosaic of African *Bos taurus*, European *B. taurus*, *B. indicus*, and

various hybrid populations [20-22]. African cattle form a gradient of *B. taurus* and *B. indicus*

ancestry across the continent [20]. This is a result of multiple domestications and the

subsequent spread of distinct populations of cattle across Africa [23]. The spread of

agriculture and livestock herding has also shaped African human genetics and linguistic

variation, illustrating the importance of domesticated species in influencing recent human

evolution [24, 25]. Zebu or indicine breeds, which have primarily *B. indicus* ancestry, are

favoured by many farmers due to their larger size and higher production yields [26]. However,

cattle populations in West Africa tend to have higher level of taurine (*B. taurus*) ancestry [20,

27]. This is because some indigenous *B. taurus* breeds have an advantage in western sub-

Saharan Africa due to their tolerance of trypanosomes, a trait termed "trypanotolerance" [26,

28]. These cattle exhibit a greater ability to control parasitaemia and anaemia, making them

more productive than *B. indicus* cattle or other *B. taurus* breeds in areas infested with tsetse

flies and trypanosomes [26, 29]. These trypanotolerant breeds, which include the longhorn

N'Dama and shorthorn Baoule, Lagune, and Somba breeds, are therefore an important genetic

resource as they are uniquely suited to livestock production in these areas [30, 31].

Trypanotolerant breeds only make up 6% of the total cattle population of Africa and but

represent 17% of the cattle in the tsetse infested areas [32]. Although trypanotolerance is a

valuable trait in these areas, trypanotolerant breeds such as the N'Dama, are not more widely

used because of their relatively inferior production characteristics, unpredictable temperament

and smaller size, which make them unsuitable for draft purposes [32, 33].

Trypanotolerance has been shown to be a heritable trait, although there is variability in

tolerance between animals within breeds [13, 29, 34, 35]. Trypanotolerant breeds are also less

susceptible to other infectious diseases such as helminthiasis, ticks and tick-borne-diseases

and genes associated with the immune response have been found to be under selection in west

African cattle populations [36, 37]. Trypanotolerant and trypanosusceptible breeds have been

shown to vary in their immune responses and exhibit gene expression differences in various

tissues at key points during infection after infection with trypanosomes [33, 38-42]. They also

show differences in ability to control anaemia over two years [43]. There are also *B. taurus*/*B.*

indicus hybrid breeds that have been observed to be trypanotolerant [44]. However,

trypanotolerant breeds with high levels of African *B. taurus* ancestry are better able to control

anaemia while hybrid animals exhibit intermediate levels of control when compared to

susceptible *B. indicus* breeds [45]. The genes, genomic regulatory elements (GREs), and

genomic regulatory networks (GRNs) underpinning trypanotolerance remain largely

unknown, although several candidate genes and regulatory mechanisms have been suggested

based on population genomics and functional genomics studies [33, 40, 42, 43, 46-49]. It has

been proposed that a better understanding of the trypanotolerance trait will facilitate genome-

informed breeding programmes that could also leverage transgenesis to enhance the trait,

thereby increasing the productivity of livestock production in sub-Saharan Africa [26, 33].

In this study we analysed Affymetrix® Bovine Genome Array gene expression data

across multiple tissues from trypanotolerant N'Dama and trypanosusceptible Boran cattle that

had been infected with *T. congolense* across an infection time course [33, 40, 41]. We used

previously published data from liver, lymph node, and spleen tissue samples and combined

these data with new peripheral blood mononuclear cell (PBMC) data from the same animals.

Differentially expressed genes (DEGs) were detected across the infection time course for each

breed and between the two breeds for each time point. The sets of DEGs were then used for

functional enrichment analyses focused on gene ontology (GO) to catalogue and interrogate

biological processes associated with disease pathogenesis and trypanotolerance.

Materials and methods

Data sources

New microarray gene expression data

Affymetrix® Bovine Genome Array data sets were generated for 40 peripheral blood mononuclear cell (PBMC) samples from 10 African cattle (5 trypanotolerant N'Dama and 5 trypanosusceptible Boran) collected at four time points for a *T. congolense* infection experiment performed in 2003 [39-41]. The cattle, all female and aged between 19–28 months, were reared together at the International Livestock Research Institute (ILRI) ranch at Kapiti Plains Estate, Kenya which is located in an area free from tsetse flies and trypanosomiasis [41]. The cattle were experimentally infected with the *T. congolense* clone IL1180 [50, 51] delivered via the bites of eight infected tsetse flies (*Glossina morsitans morsitans*) [52, 53] at the ILRI laboratories in Nairobi, Kenya. The flies were allowed to feed on the shaved flanks of the animals until engorgement. The infections were carried out in three batches which took place two weeks apart and were confirmed by microscopy. Parasitaemia and anaemia measured by packed cell volume (PCV) were monitored throughout the course of the infection [41].

Peripheral blood (200 ml) was collected in heparinised syringes before infection and at 14, 25, and 34 days post-infection (DPI), these time points were selected to align with the first wave of parasitaemia [41]. This animal work was completed prior to the requirement for formal Institutional Permission in Ireland, which is based on European Union Directive 2010/63/EU; however, all animal procedures were approved by the Institutional Animal Use and Care (IAUC) Committee of ILRI in accordance with the UK Animals (Scientific Procedures) Act 1986 and all efforts were made to ensure high welfare standards and ethical handling of animal subjects.

Peripheral blood mononuclear cell (PBMC) samples were isolated from the blood

samples with Percoll™ gradients (GE Healthcare, Buckinghamshire, UK) and total RNA was

extracted from the PBMCs in TriReagent (Molecular Research Center, Inc., Cincinnati, OH,

USA) and the RNA samples were then DNase-treated and purified using an RNeasy mini kit

(Qiagen Ltd., Crawley, UK). RNA quality and quantity was assessed using the 18S/28S ratio

and RNA integrity number (RIN) on an Agilent Bioanalyzer with the RNA 6000 Nano

LabChip® kit (Agilent Technologies, Inc., Santa Clara, CA, USA). Following this, cDNA

labelling, hybridisation and scanning for the microarray experiments were performed by

Almac Diagnostic Services (Craigavon, Northern Ireland) using a one-cycle

amplification/labelling protocol. Gene expression data in the form of cell intensity files

(.CEL) were generated using the Affymetrix GeneChip® Operating Software (GCOS)

package. The Affymetrix® GeneChip® Bovine Genome Array data sets generated for this

study have been deposited in the European Molecular Biology Laboratory European

Bioinformatics Institute ArrayExpress data repository under accession number E-MTAB-

14517 (temporary manuscript reviewer link:

https://www.ebi.ac.uk/biostudies/arrayexpress/studies/E-MTAB-14517?key=b9fd37fb-644d-

442b-b443-53506a43a8df).

Previously published microarray gene expression data

Additional Affymetrix® Bovine Genome Array data sets were obtained from a published

study using solid tissues samples from the same animal infection time course [33] resulting in

a total of 220 biological samples across 12 infection time points and four tissues (for both

N'Dama and Boran) before filtering (**Table 1**). **Fig 1** illustrates the experimental design and

study workflow. The computer code required to repeat and reproduce the analyses is available

at doi.org/10.5281/zenodo.11502109.

Table 1. Days post infection, tissues, numbers of N'Dama (NDAM) and Boran (BORA)
samples, and sources of microarray data used in this study before and after sample filtering.

Days post infection	Tissue	No. NDAM samples pre-filtering	No. BORA samples pre-filtering	No. NDAM samples post-filtering	No. BORA samples post-filtering	Source*
0	Blood (PBMC)	5	5	5	5	a
0	Liver	20	20	19	18	b
0	Lymph node	5	5	5	4	b
0	Spleen	5	5	5	4	b
12	Liver	5	5	5	5	b
14	Blood (PBMC)	5	5	5	5	a
15	Liver	5	5	5	5	b
18	Liver	5	5	5	5	b
21	Liver	5	5	5	5	b
21	Lymph node	5	5	4	5	b
21	Spleen	5	5	5	5	b
25	Blood (PBMC)	5	5	5	5	a
26	Liver	5	5	3	5	b
29	Liver	5	5	5	5	b
32	Liver	5	5	5	5	b
34	Blood (PBMC)	5	5	5	5	a
35	Liver	5	5	5	5	b
35	Lymph node	5	5	4	5	b
35	Spleen	5	5	5	5	b
Population total		110	110	105	106	
Total		220		211		

185 *a this study, b [33].

Fig 1. Diagram showing the experimental design and study workflow. Trypanosome image by Matus Valach and cattle images by Tracy A. Heath and T. Michael Keesey via phylopic.org and tissue images via healthicons.org. Colours were generated from the khroma (v. 1.10.0) [54] and viridis (v. 0.6.3) [55] R packages.

Data filtering and normalisation

Quality control with log transformation was performed using affy (v. 1.80.0) [56], Biobase (v. 2.62.0) [57], and arrayQualityMetrics (v. 3.58.0) [58] with R (v. 4.3.2) [59]. As in previous microarray studies, any individual samples that were identified as outliers in two or more quality control tests were discarded from the analysis [60]. Normalisation was performed using farms (v. 1.25.0) [61] with R (v. 4.3.2).

The intensity of expression for each sample was visualised both with the raw data and after normalisation using affy (v. 1.80.0), Biobase (v. 2.62.0), dplyr (v. 1.1.2) [62], ggh4x (v. 0.2.4) [63], ggplot2 (v. 3.4.2) [64], ggtext (v. 0.1.2) [65], magrittr (v. 2.0.3) [66], readr (v. 2.1.4) [67], reshape2 (v. 1.4.4) [68], and stringr (v. 1.5.0) [69] with R (v. 4.3.2). Colours were generated from khroma (v. 1.10.0) [54] and viridis (v. 0.6.3) [55]. Informative probe sets were identified and extracted using farms (v. 1.25.0) with R (v. 4.3.2).

Principal component analysis

Principal component analysis (PCA) was performed using limma (v. 3.58.1) [70] with R (v. 4.3.2). The results were visualised using dplyr (v. 1.1.2), ggplot2 (v. 3.4.2), patchwork (v. 1.2.0) [71], and stringr (v. 1.5.0), with R (v. 4.3.2). Colours were generated from khroma (v. 1.10.0) and viridis (v. 0.6.3).

Differential expression analysis

Differential expression analysis was performed using limma (v. 3.58.1) [70] with R (v. 4.3.2). The correlation between samples from the same animal was estimated and included in the linear model [70, 72]. The contrast matrix contained 64 contrasts across the two populations, four tissues and 12 time points which included direct (DIRE) contrasts to identify changes in expression in the N'Dama samples relative to the Boran samples ($DIRE_i = N_i - B_i$, where N represents the N'Dama population and B represents the Boran population), N'Dama (NDAM) and Boran (BORA) contrasts to identify changes in expression within the

populations relative to time 0 (NDAM$_i$ = N$_i$ – N$_0$; BORA$_i$ = B$_i$ – B$_0$) and response (RESP)

contrasts to identify changes in expression over time in the N'Dama samples relative to the

Boran (RESP$_i$ = [N$_i$ – N$_0$] – [B$_i$ – B$_0$]) (**Fig 1**, **Table S1**) [33]. Moderated *t*-statistics,

moderated *F*-statistic, and log-odds of differential expression were calculated for the

informative probe sets by empirical Bayes moderation of the standard errors towards a global

value [70, 73]. Global Benjamini-Hochberg correction for multiple testing was applied across

all contrasts [70, 74]. Probe sets with an adjusted *P*-value of ≤ 0.05 (B-H $P_{adj.}$ ≤ 0.05) were

determined to be significantly differentially expressed [70]. The microarray probe sets were

converted to genes using gprofiler2 (v. 0.2.2) [75] with R (v. 4.3.2).

The results of the differential expression analysis were visualised using ComplexUpset

(v. 1.3.3) [76], dplyr (v. 1.1.2), ggh4x (v. 0.2.4), ggplot2 (v. 3.4.2), ggrepel (v. 0.9.3) [77],

patchwork (v. 1.2.0), readr (v. 2.1.4), rlang (v. 1.1.1) [78], stringr (v. 1.5.0), and tidyr (v 1.3.0)

[79] with R (v. 4.3.2). Colours were generated from khroma (v. 1.10.0) and viridis (v. 0.6.3).

228   Functional enrichment analysis

Functional enrichment was performed and visualised using dplyr (v. 1.1.2), ggh4x (v.

0.2.4), ggplot2 (v. 3.4.2), ggrepel (v. 0.9.3), gprofiler2 (v. 0.2.2), magick (v. 2.8.1) [80] with

ImageMagick (v. 6.9.12.96) [81], magrittr (v. 2.0.3), purrr (v. 1.0.1) [82], readr (v. 2.1.4),

rlang (v. 1.1.1), scales (v. 1.2.1) [83], and stringr (v. 1.5.0) with R (v. 4.3.2). Colours were

generated from khroma (v. 1.10.0) and viridis (v. 0.6.3). The background set was the set of

informative probe sets with valid Ensembl IDs. The query sets included those that were

significantly differentially expression (adjusted *P*-value ≤ 0.05) for each of the 64 contrasts.

The analysis was restricted to Gene Ontology (GO) terms and driver GO terms were

highlighted. Generic Enrichment Map (GEM) files were generated and used as input for

EnrichmentMap (v. 3.3.6) [84] with Cytoscape (v. 3.8.0) [75, 85, 86]. AutoAnotate (v. 1.3.5)

[87] was used to create clusters of GO terms with genes in common with clusterMaker2 (v.

2.0) [88] and annotate them with appropriate names using WordCloud (v. 3.1.4) [89]. The

yFiles Layout Algorithms (v. 1.1.3) method [90] was used to remove overlaps in the networks.

Results

Data filtering and normalisation

Nine samples were indicated as outliers by two or more of the quality control tests and were therefore removed, leaving 211 samples for analysis (**Fig S1**, **Table 1**). After normalisation the data were filtered to retain only informative probe sets, which resulted in a data set of 11,050 probe sets or 45.89% of the initial 24,128 probe sets.

Principal component analysis

Based on the top ten principal components (PCs), the first PC explained 82.86% of the total variation in the data for PC1–10 and separated the liver from the blood, lymph node, and spleen samples (**Fig 2**). The second PC explained a further 9.10% of the total variation for PC1–10 and separated the blood from the lymph node and spleen samples (**Fig 2**). The third and fourth PCs explained 2.24% and 1.78% of the total variation for PC1–10, respectively, and separated the samples according to days post infection (dpi) while the fifth PC, which explained 1.32% of the total variation in the data for PC1–10 separated the lymph node and spleen samples (**Fig S2 S4**).

Fig 2. A. Principal component analysis (PCA) of the microarray data set with samples coloured according to tissue and with the shape indicating the
population. The first two principal components (PC1 and PC2) are shown, and **B.** bar chart of proportion of variance of the top ten PCs.

Differential expression analysis

The results of the differential expression analysis are detailed in **Table S2** and the number of significant (B-H $P_{adj.} \leq 0.05$) differentially expressed genes (DEGs) generally increased over time across all contrast types and tissues until 21 dpi when they began to decrease slightly before rising again to their peak at 34 or 35 dpi (**Fig 3**, **S5**☐**S8**). The highest numbers of significant DEGs were found in the N'Dama and Boran contrasts, followed by the direct contrasts (**Fig 3**). Within the response contrasts the highest number of significant DEGs were found in the lymph node samples at 35 dpi (**Fig 3**, **S5**) while the blood samples at 34 dpi had the highest number of significant DEGs in the direct contrasts (**Fig 3**, **S6**). The highest number of significant DEGs in the N'Dama and Boran contrasts were at 35 dpi in the lymph node and liver samples, respectively (**Fig 3**, **S7**, **S8**). The numbers of significant DEGs were generally higher in the N'Dama contrasts than in the Boran contrasts early in the time course (**Fig** 3).

Fig 3. Bar chart showing the numbers of significant differentially expressed genes (DEGs) for all 64 contrasts (B-H $P_{adj.}$ ≤ 0.05). The extent of the bar above and below 0 on the *y*-axis indicates the numbers of significant DEGs with increased and decreased expression, respectively. The position on the *x*-axis indicates the number of days post infection (dpi). The bar colours represent the contrast type while the bar outline colours represent the tissue.

The N'Dama and Boran contrasts also had a higher number of significant DEGs in common between them when overlaps among all 64 contrasts were examined (**Fig S9**). Within the response and direct contrasts any overlaps in significant DEGs were generally confined to those between samples from the same tissue type at different time points (**Fig S10**, **S11**) while the N'Dama and Boran contrasts included more overlaps between tissues and time points (**Fig S12**, **S13**).

There were also many overlaps when the top 10 most significant genes with increased and decreased expression were examined for each of the contrasts (**Table 2**, **S3□S5**, **Fig 4**, **S14□S27**). The set of 1,267 genes identified as being in the top 10 most significant for increased or decreased expression for each of the 64 contrasts contained just 602 unique genes (**Table 2**, **S3□S5**). Of these, 328 were in the top 10 most significant for only one contrast, with the remaining 274 in the top 10 most significant for an average of 3.43 contrasts (**Table 2**, **S3□S5**). The response contrasts had the most unique genes in the top 10 most significant with 129 of the 328 unique top genes, while the direct, N'Dama and Boran contrasts had 75, 58 and 66 unique top genes, respectively (**Table 2**, **S3□S5**). The most common genes in the top 10 most significant differentially expressed for all the contrasts included *TMSB10,* which was in the top 10 most significant genes for 16 of the 64 contrasts, followed by *CYRIB*, *PTPRC*, *SPI1*, and *TTLL1*, which were in the top 10 most significant genes for 14 of the 64 contrasts (**Table 2**, **S3□S5**). The *TMSB10* gene was in the top 10 most significant genes with increased expression for the liver samples at all time points for the N'Dama and Boran contrasts (**Table S4**, **S5**). The *CYRIB*, *PTPRC*, and *SPI1* genes showed similar patterns with the exclusion of 12 dpi (**Table S4**, **S5**). *TTLL1* was in the top 10 most significant genes with increased expression in the blood samples at 25 and 34 dpi and the liver samples at 18 dpi for the response contrasts (**Table 2**). It was also in the top 10 most significant genes with increased expression for the blood samples at 0, 14, 25, and 34 dpi, the liver samples at 18

303   dpi, and the lymph node and spleen samples at 0, 21, and 35 dpi for the direct contrasts (**Table**

304   **S3**).

Fig 4. Volcano plot showing the results of the RESP contrast for the peripheral blood mononuclear cell (PBMC) samples at 34 days post infection (dpi). Each data point represents a gene with the position on the *x*- and *y*-axes indicating the log$_2$ fold change and -log$_{10}P_{adj.}$, respectively. Genes above the horizontal dashed line are significantly differentially expressed with the colours representing the change in expression. The top 10 most significant genes for increased and decreased expression with gene symbols are labelled.

The most common genes in the top 10 most significant differentially expressed for the

response contrasts included *CYP4B1* and *RAB31,* which were in the top 10 most significant

genes for four of the 15 response contrasts, followed by *OLFM4*, *TMEM45B*, and *TTLL1,*

which were in the top 10 most significant genes for three of the 15 response contrasts (**Table**

2). For the response contrasts, *CYP4B1* was in the top 10 most significant genes with

increased expression for the spleen samples at 21 and 35 dpi and the top 10 most significant

genes with decreased expression for the lymph node samples at the same time point contrasts

(**Table 2**). It was also in the top 10 most significant genes with increased expression for the

spleen samples at 0 dpi and the top 10 most significant genes with decreased expression for

the lymph node samples at the same time point for the direct contrasts (**Table S3**). *RAB31*

was only in the top 10 most significant genes with decreased expression for the liver samples

at 26, 29, 32, and 35 dpi for the response contrasts (**Table 2**). For the response contrasts,

OLFM4 was in the top 10 most significant genes with decreased expression for the blood

samples at 14, 25, and 34 dpi (**Fig 4**, **Table 2**). It was also in the top 10 most significant genes

with decreased expression for the blood samples at 0 dpi for the direct contrasts, and at 14,

25, and 34 dpi for the N'Dama contrasts (**Table S3**, **S4**). For the response contrasts,

TMEM45B was in the top 10 most significant genes with increased expression for the liver

samples at 21, 26, and 29 dpi (**Table 2**). It was also in the top 10 most significant genes with

decreased expression for the liver samples at 0, 12, and 15 dpi for the direct contrasts (**Table**

S3).

Table 2. Tissue, days post infection (dpi) and the top 10 most significant genes with
increased and decreased expression with valid gene symbols for the response contrasts.

Tissue	Days post infection	Most significant genes with increased expression
BL	14	SRSF11, SNORD50B, CCAR1, RESF1, CLECL1P, KTN1, SENP6, KRR1, CCNL1, TRMT13
BL	25	RESF1, TTLL1, ADAMDEC1, COX6C, KLF5, S100A7, TLR6, CHIC2, ZNF318, RUM1
BL	34	CHIC2, TTLL1, SPDEF, PLAU, CDADC1, SNORD26, SNORD27, MRPL57, POLB, ZC3H12A
LI	12	IL18BP, FOXA3, CXCL11, DVL1, REG3A, CD40, BIRC3, SRGN, MAPK12, WARS1
LI	15	SGCB, RNF39, ITIH2, FBXO34, GNPAT, TCF20, bta-mir-2442, C8H9orf64, SCRN2, STAG2
LI	18	PCF11, TTLL1, PER1, NRG2, CIPC, MKNK2, BAG3, RPL22, ACTR8
LI	21	TMEM45B, RFTN1, UPF2, TCEA3, LTB, CPQ, PLIN5, ABCC5, OSGIN1, SMARCA2
LI	26	TMEM45B, CIDEA, ECT2L, CCDC191, RETSAT, PMVK, CA5A, ANP32A
LI	29	MAGI3, TMEM45B, NGFR, PPP2R5A, CBS, DPYS, MTMR4, UGT2B10, COLEC11, GSTA4
LI	32	SFRP2, APLNR, PLAC8B, CYP3A28, AAMDC, ST7L, MSMO1, SQLE, CES1, TCEA3
LI	35	CES1, RETSAT, CYP3A28, MTMR4, KCNK17, ABHD14B, ACP6, SYBU, ACY1, GCDH
LN	21	ALB, CXCL13, ATP6V0D2, EAF2, SUCLG1, ORC1, SNORA73, ECRG4, SKA2, TNFSF13B
LN	35	CCL20, RGS13, DEFB4A, MYB, ELL3, SNORA73, LXN, PLEKHF2, EED, SPAG5
SP	21	TRMT10B, CYP4B1, NXPH1, KCNA3, MDK, KIRREL2, PTGDS, FAM83D, PRR5L, LTB
SP	35	CYP4B1, MAB21L1, FAM83D, IKZF2, SNCA, PRR5L, RNASE4, RGS13, ZCCHC17, TARP
Tissue	**Days post infection**	**Most significant genes with decreased expression**
BL	14	OLFM4, PGA5, CDC42EP3, ORAI3, KAT7, VAT1L, PMF1, OASL, RAVER1, ICAM3
BL	25	SLC6A4, OLFM4, PGA5, SMOX, PRDM1, SLC9A6, PFKFB4, CCR7, MEIS1, F11R
BL	34	CCR7, ENDOD1, SLC6A4, PIM2, OLFM4, SMOX, TUBB1, ITGA2, DSTN, SLC40A1
LI	12	NUDT12, APOB, PLD1, MYO1B, IQGAP2, CRYZ, EPS8, MASP1, APC, CYP4V2
LI	15	MAPK6, TCIRG1, CXCL17, CUL1, XRN2
LI	18	MEP1B, SLC1A3, TJP2, OLR1, ACTG1, bta-mir-3533, IGFALS, NAE1, IMPA1, SNX6
LI	21	LEAP2, C15H11orf52, SLC25A25, ABHD10, ANGPTL8, LPAR6, KBTBD6, MTRF1, MFSD2A, ISG12(B)
LI	26	PAMR1, RAB31, VLDLR, GPLD1

332 **Table 2 continued.**

Tissue	Days post infection	Most significant genes with decreased expression
LI	29	PEG3, RAB31, JCHAIN, WWOX, OTULINL, DNMT1, OLR1, ACTG1, bta-mir-3533, PLTP
LI	32	MEP1B, SLC1A3, JCHAIN, RPSA, SNORA62, XRCC5, RAB31, LDHA, WWOX, CORO1B
LI	35	ANGPTL8, RAB31, LEAP2, ANGPTL4, VLDLR, PLOD2, BNIP3, GNL2, MTMR11, S100A8
LN	21	CLDN11, STAB1, CYP4B1, BRB, P2RX4, PROS1, KANK3, HYAL2, ELOVL7, COLEC12
LN	35	BRB, CYP4B1, STAB1, CLDN11, PROS1, HEPH, P2RX4, MAN1C1, C1QTNF5, SCARA5
SP	21	BMP2, FABP3, GRO1, THBS1, BCL10, F2RL2, CLDN1, SAMSN1, TLR4, ARL14EP
SP	35	NELL2, CD14, ITGA8, MUSTN1, RBP1, F2RL2, CXCL16, IL33, CATHL3, BMP2

Functional enrichment analysis

GO terms related to regulation of the mitotic cell cycle were significantly enriched for

the genes with significantly increased expression in the blood and lymph node samples for

the response contrasts (**Fig 5**). The top driver GO terms for the blood samples included

GO:0006396 RNA processing and *GO:0005730 nucleolus* (**Fig S28**) while the top driver GO

terms for the lymph node samples included *GO:0007059 chromosome segregation*,

GO:0005694 chromosome, *GO:0006259 DNA metabolic process*, *GO:0006260 DNA*

replication, *GO:0051301 cell division*, *GO:0003677 DNA binding*, and *GO:0008017*

microtubule binding (**Fig S30**). GO terms related to steroid metabolic processes

(*GO:0008202 steroid metabolic process*) and oxidoreductase activity (*GO:0016705*

oxidoreductase activity, acting on paired donors, with incorporation or reduction of

molecular oxygen, *GO:0016712 oxidoreductase activity, acting on paired donors, with*

incorporation or reduction of molecular oxygen, reduced flavin or flavoprotein as one donor,

and incorporation of one atom of oxygen) were significantly enriched for the genes with

significantly increased expression in the liver samples for the response contrasts (**Fig 5**, **S29**).

No GO terms were significantly enriched for genes with significantly increased expression in

the spleen samples for the response contrasts (**Fig 5**).

GO terms related to blood coagulation (*GO:0007596 blood coagulation*) and the cell-

substrate junction (*GO:0030055 cell-substrate junction*) were significantly enriched for the

genes with significantly decreased expression in the blood samples for the response contrasts

(**Fig 5**, **S28**). GO terms related to receptor-mediated endocytosis were significantly enriched

for the genes with significantly decreased expression in the liver and lymph node samples for

the response contrasts (**Fig 5**). The top driver GO terms included *GO:0003823 antigen*

binding for the liver samples (**Fig S29**), and *GO:0071944 cell periphery*, *GO:0009986 cell*

surface, *GO:0007155 cell adhesion*, and *GO:0006897 endocytosis* for the lymph node

samples (**Fig S30**). GO terms related to multicellular development and morphogenesis

including the driver GO terms *GO:0007167 enzyme-linked receptor protein signaling*

pathway, *GO:0032501 multicellular organismal process*, and *GO:0071495 cellular response*

to endogenous stimulus were also significantly enriched for the genes with significantly

decreased expression in the lymph node samples for the response contrasts (**Fig 5**, **S30**). GO

terms related to cell migration and motility (*GO:0016477 cell migration*, *GO:0048870 cell*

motility) and cargo receptor activity (*GO:0038024 cargo receptor activity*) were also enriched

for these genes (**Fig 5**, **S30**). GO terms related to the extracellular region (*GO:0005576*

extracellular region) were significantly enriched for the genes with significantly decreased

expression in both the lymph node and spleen samples for the response contrasts (**Fig 5**, **S30**,

S31), while GO terms related to lipopolysaccharide binding (*GO:0001530*

lipopolysaccharide binding) were also significantly enriched for the genes with significantly

decreased expression in the spleen samples for the response contrasts (**Fig 5**, **S31**).

For the direct contrasts, GO terms related to organelle metabolic processes were

significantly enriched for the genes with significantly increased expression in the blood, liver,

and lymph node samples (**Fig S32**). The GO term for leukocyte proliferation was

significantly enriched for genes with significantly increased expression in the blood samples,

while GO terms related to scavenger receptor activity and cell periphery were also

significantly enriched for genes with significantly increased expression in the lymph node

samples (**Fig S32**). GO terms related to receptor signalling pathways were significantly

enriched for genes with significantly decreased expression in the liver and lymph node

samples, while GO terms related to coagulation were significantly enriched for genes with

significantly decreased expression in the blood samples (**Fig S32**). GO terms related to actin

binding and cytoskeleton, immune response and antigen binding, small molecule and

pyruvate metabolic processes, and the cytosolic ribosome were also significantly enriched for

genes with significantly decreased expression in the liver samples (**Fig S32**). GO terms

related to tube development, cell migration, extracellular region, cell adhesion, and calcium

ion binding were significantly enriched for genes with significantly decreased expression in

the lymph node samples (**Fig S32**). GO terms related to metaphase chromosome alignment,

the chromosome centromeric region, and the microtubule cytoskeleton were significantly

enriched for genes with significantly decreased expression in the spleen samples (**Fig S32**).

For the N'Dama contrasts, GO terms related to cell process regulation were

significantly enriched for genes with significantly increased expression in all tissues (**Fig

S33**). GO terms related to GTPase activator activity, phagocytosis, and protein kinase binding

were also significantly enriched for genes with significantly increased expression in the liver

samples (**Fig S33**). GO terms related to cell development and regulation were significantly

enriched for genes with significantly decreased expression in all tissues (**Fig S33**). GO terms

related to metabolic processes, peroxisome organization, electron transfer activity,

oxidoreductase activity, and the endoplasmic reticulum membrane were enriched for genes

with significantly decreased expression in the liver samples (**Fig S33**). GO terms related to

actin cytoskeleton organization, extracellular matrix, cell projection, and vesicle-mediated

transport were significantly enriched for genes with significantly decreased expression in the

lymph node and spleen samples (**Fig S33**). GO terms related to the extracellular region were

also significantly enriched for genes with significantly decreased expression in the lymph

node samples, while GO terms related to extracellular matrix organization were significantly

enriched for genes with significantly decreased expression in the spleen samples (**Fig S33**).

For the Boran contrasts, GO terms related to cell process regulation were significantly

enriched for genes with significantly increased expression in all tissues and GO terms related

to cell response regulation were significantly enriched for genes with significantly decreased

expression in all tissues, while GO terms related to metabolic processes were also

significantly enriched for genes with significantly decreased expression in the liver and

spleen samples (**Fig S34**).

Fig 5. EnrichmentMap network of significantly enriched gene ontology (GO) terms identified from g:Profiler functional enrichment of significantly differentially expressed genes (DEGs) for the response contrasts. Each node represents a GO term with the colour of the node representing the tissue and the size representing the number of genes in the GO term. The edges indicate overlap between the GO terms with the width of the edges representing the similarity coefficient for the connected GO terms. The GO terms are clustered by AutoAnotate with the background colour of the clusters representing the direction of expression. The clusters are labelled with the size of the label scaling with the number of GO terms in the cluster.

Discussion

The data filtering and normalisation steps resulted in a similar number of samples and probe sets for analysis as previous studies that used Affymetrix® Bovine Genome Array data sets to examine host responses to trypanosomiasis and other infectious diseases of cattle, including the previous analysis of the liver, lymph node, and spleen samples [33, 91-93]. The clear separation of the liver samples from the other tissues that is evident for PC1, which also explained the majority of the total variation in the data set for PC1–10, may be due to several factors (**Fig 2**). The first is biological differentiation between the liver samples and those of the other tissues. This is supported by a previous gene expression microarray study using rat (*Rattus norvegicus*) tissues, which also observed that PC1 in a PCA separated liver samples from those taken from blood, lymph node, spleen, and other tissues [94]. This biological difference may have been compounded by the higher number of liver samples in the data set from Noyes and colleagues [33] due to the experimental design, which encompassed a larger number of time points for the liver samples as well as a greater number of animals from which the liver samples were taken to prevent the collection of multiple liver biopsies from the same animal at consecutive time points [33]. This experimental design consideration also resulted in more control liver samples for the pre-infection time point [33]. In total, 115 of the 211 samples that passed the quality control filters or 54.50% of the filtered data set were liver samples (**Table 1**). Because an unequal number of samples is known to increase the distance between groups with larger sample sizes in a PCA, it is possible that the disproportionately large number of liver samples in the present study has given rise to the effect seen in PC1 [95-97]. It is therefore unsurprising that PC2, which explained a further 9.10% of the total variation in the data set for PC1–10, separated the blood samples from the lymph node and spleen samples (**Fig 2**) as the blood samples were the second most numerous in the data set with 40 samples or 18.96% of the filtered data set (**Table 1**). It is not until PC5, which

explained 1.32% of the total variation in the data set for PC1–10, that the lymph node and

spleen samples are separated (**Fig S4**). The samples from these tissues are the least numerous

in the data set with 29 spleen and 27 lymph node samples, making up 13.74% and 12.80% of

the filtered data set, respectively (**Table 1**). The spleen and lymph nodes are also both key

components of the lymphatic and immune systems and their functional and morphological

similarities as hemopoietic organs that filter bodily fluids have been long established [98-

100]. It is logical, therefore, that these tissues would show similar patterns of gene expression

both before and after infection. The third and fourth PCs (PC3 and PC4), which explained

2.24% and 1.78% of the total variation for PC1–10, respectively, separated the samples

temporally in order of dpi (**Fig S2**, **S3**). This illustrates the overall similarity in response to

infection across the populations and is in agreement with RNA-seq data from whole blood

samples taken during a similar trypanosome infection time course experiment [42]. It also

shows that the systemic response to trypanosome infection in peripheral blood is distinct

from that in the liver and other tissues and that blood sampling alone may not be sufficient to

capture all the tissue specific responses to infection.

Gene expression in bovine tissues can be measured using a variety of methods, one of

which is the Affymetrix® Bovine Genome Array based on GeneChip® technology [101] and

which contains more than 24,000 probe sets representing over 23,000 gene transcripts. The

Affymetrix® Bovine Genome Array platform has been previously used to study infectious

disease in cattle, including trypanosomiasis [33], and for mycobacterial infections that cause

tuberculosis and paratuberculosis [91-93]. Importantly, although expression microarrays have

been largely supplanted by RNA sequencing (RNA-seq), when the same biological samples

were analysed using both the Bovine Genome Array and RNA-seq, gene expression data

obtained using the two platforms were extremely well correlated in peripheral blood and

alveolar macrophages [102-104]. In addition, it has recently been shown that peripheral blood

gene expression data generated using RNA-seq, a different microarray technology (long

oligonucleotide microarrays), and reverse transcription quantitative real-time PCR (RT-

qPCR) were well correlated in independent studies of trypanosome infection in cattle [42].

The pattern of an increase in the numbers of significant DEGs until 21 days post

infection followed by a decrease before rising to a final peak at 35 days post infection across

contrast types and tissues (**Fig 3**, **S5**☐**S8**) is consistent with known information about the

cyclic nature of parasitaemia during trypanosome infection, which has been shown to have an

initial peak at 20 dpi [38]. It is also in agreement with the measures of parasitaemia in the

blood samples during this infection time course experiment, which showed peaks of

parasitaemia from 15 to 22 dpi [41]. The final peak of significant DEGs at the end of the

infection time course is consistent with analysis of the same blood samples using a gene

expression platform with less gene content—a bovine long oligonucleotide (BLO) array—

which detected the highest number of significant DEGs between the two populations at 34

dpi [40]. It is also in agreement with a study using RNA-seq data from whole blood samples

that found the highest number of DEGs in N'Dama samples at the end of a similar

trypanosome infection time course experiment [42]. The higher numbers of significant DEGs

in the N'Dama contrasts when compared to the Boran contrasts is in agreement with previous

results   and   supports   the   hypothesis   that,   compared   to   trypanosusceptible   Boran,

trypanotolerant N'Dama cattle exhibit an earlier proinflammatory response, which is evident

in both PBMC and whole peripheral blood [40-42]. That the response contrasts had the

lowest number of significant DEGs is to be expected given that these contrasts are examining

the data set for differences in the rate of change of gene expression between the populations

[33]. In addition, the numbers of significant DEGs for the response contrasts were similar to

those found in the original analysis by Noyes and colleagues [33], which employed a similar

approach. The numbers of genes in common between the contrasts is to be expected due to

the shared underlying data set while the numbers of genes in common within the contrasts is

likely due to similar patterns of gene expression found across the populations, tissues, and

time points during the infection (**Fig S9□S13**). This is in agreement with RNA-seq data that

showed large overlaps in DEGs in response to trypanosome infection between five different

cattle populations with varying levels of trypanotolerance [42].

The similarities in gene expression are also illustrated by the high level of overlap in

the top 10 most significant DEGs with increased and decreased expression between the

contrasts (**Table 2**, **S3□S5**). The most common genes in the top 10 most significant DEGs for

all the contrasts included those related to the immune system, such as the most common

significant DEG—the thymosin beta 10 gene (*TMSB10*), which has been identified as a hub

gene in the response to Rift Valley fever (RVF), an important viral disease in African cattle

[105]. Other immune-related genes in the set of the most common genes in the top 10 most

significant DEGs for all the contrasts included the CYFIP related Rac1 interactor B gene

(*CYRIB*), which encodes a regulator of phagocytosis [106], and the protein tyrosine

phosphatase receptor type C gene (*PTPRC*), which is an essential regulator of T and B cell

antigen receptor-mediated activation and has been highlighted by multiple studies of

trypanosome infection in mice [107-112]. Similarly, the Spi-1 proto-oncogene gene (*SPI1*),

which encodes a transcription factor that activates gene expression during myeloid and B-

lymphoid cell development, has also been implicated as an important host gene in

trypanosome infection in the mouse [111, 113]. The presence of immune genes in the top 10

most significant DEGs is to be expected given the nature of the infection time course

experiment and is in agreement with previous results from this and other infection

experiments, illustrating the similarity in immune responses between the N'Dama and Boran

cattle during trypanosome infection [33, 40, 42].

The most common genes in the top 10 most significant DEGs for the response contrasts

included genes which are involved in in inflammation and immunity such as the olfactomedin

4 gene (*OLFM4*) and the transmembrane protein 45B gene (*TMEM45B*) [114, 115]. Other

common genes in the top 10 most significant DEGs for the response contrasts are more

varied in function and included the cytochrome P450 family 4 subfamily B member 1 gene

(*CYP4B1*), which is involved in drug metabolism [116]; the RAB31, member RAS oncogene

family gene (*RAB31*), which is a small GTPase [117]; and the TTL family tubulin

polyglutamylase complex subunit L1 gene (*TTLL1*), which is involved in microtubule

cytoskeleton organisation [118].

The top 10 most significant DEGs for the response contrasts also included genes

encoding antimicrobial peptides (AMPs), which are key components of the innate immune

system that have huge therapeutic potential and are important for the healthy function of a

variety of bovine tissues [119-122]. This is in agreement with previous RT-qPCR results

using blood samples from the same time course infection experiment; in this regard, it has

previously been hypothesised that trypanotolerance may be partly due to inherited regulatory

sequence variation in genes encoding antimicrobial peptides [39]. For example, the liver

enriched antimicrobial peptide 2 gene (*LEAP2*) [123], which was in the top 10 most

significant DEGs with decreased expression for the liver samples at 21 and 35 dpi for the

response contrasts (**Table 2**). The previous RT-qPCR results showed that *LEAP2* exhibited

significantly decreased expression in N'Dama samples at 34 dpi relative to day 0 while the

Boran animals showed no such change [39]. Other AMP genes in the top 10 most significant

DEGs for the response contrasts which examine the N'Dama samples relative to the Boran

samples included the cathelicidin antimicrobial peptide gene (*CATHL3*) [124] with decreased

expression in the spleen samples at 35 dpi, which is notable since cathelicidins were found to

be the most effective of the three classes of AMP at killing both insect and bloodstream forms

of *T. brucei* in mice [125]; the defensin beta 4A gene (*DEFB4A*) [126] with increased

expression in the lymph node samples at 35 dpi; and S100 calcium binding protein A7 gene

(*S100A7*) [127] with increased expressed in peripheral blood at 25 dpi (**Table 2**) [39, 124,

126, 127].

Another group of genes represented in the top 10 most significant DEGs for the

response contrasts were cytokine genes (**Table 2**). This is consistent with both RT-qPCR and

BLO microarray results from the same infection time course experiment and previous studies

using the same animals [33, 40, 41]. This observation agrees with more recent

transcriptomics studies from trypanotolerant and trypanosusceptible cattle and studies of

trypanosome infection in the mouse [42, 128]. These genes included the C-C motif

chemokine ligand 20 gene (*CCL20*) with increased expression in the lymph node samples at

35 dpi; the C-X-C motif chemokine ligand 11 (*CXCL11*) with increased expression in the

liver samples at 12 dpi; the C-X-C motif chemokine ligand 13 gene (*CXCL13*) with increased

expression in the lymph node samples at 21 dpi; the C-X-C motif chemokine ligand 16 gene

(*CXCL16*) with decreased expression in the spleen samples 35 dpi; the C-X-C motif

chemokine ligand 17 gene (*CXCL17*) with decreased expression in the liver samples at 15

dpi; the interleukin 33 gene (*IL33*) with decreased expression in the lymph node samples at

35 dpi; and the TNF superfamily member 13b gene (*TNFSF13B*) with increased expression

in the lymph node samples at 21 dpi (**Table 2**).

Related genes such as cytokine receptors and inhibitors were also in the top 10 most

significant DEGs for the response contrasts (**Table 2**). These included the C-C motif

chemokine receptor 7 gene (*CCR7*) with decreased expression in the blood samples at 25 and

34 dpi; the C1q and TNF related 5 gene (*C1QTNF5*) with decreased expression in lymph

node samples at 35 dpi; and the interleukin 18 binding protein gene (*IL18BP*) with increased

expression in the liver samples at 12 dpi (**Table 2**). Additionally, mitogen-activated protein

kinase (MAPK) genes were present in the top 10 most significant DEGs, which is in

agreement with previous work using the same animals and also a more recent study of

trypanosome infection in cattle [33, 40, 42]. In this regard, it is notable that MAPK signalling

pathways are involved in the response to proinflammatory cytokines and are known to be

subverted by *T. cruzi* and other trypanosomatid parasites to evade the host immune response

[129-131].

A final notable group of genes present in the top 10 most significant DEGs were those

encoding apolipoproteins, including apolipoprotein B (*APOB*) with decreased expression for

the response contrast in the liver samples at 12 dpi (**Table 2**); apolipoprotein L3 (*APOL3*)

with increased expression for the direct contrast in the blood samples at 25 dpi (**Table S3**);

and apolipoprotein M (*APOM*) with decreased expression for the Boran contrast in the liver

samples at 15 dpi (**Table S5**). These genes are related to apolipoprotein 1 (*APOL1*) which

encodes the protein that makes up the trypanosome lytic factor (TLF) that is present in human

and western lowland gorilla (*Gorilla gorilla gorilla*) serum [132]. TLF is taken up by

susceptible trypanosomes where it interferes with their lysosomes and mitochondria, thereby

conferring host resistance to most trypanosome species [132]. It is therefore interesting that

related genes are among the top 10 most differentially expressed in this study.

Gene ontology terms related to regulation of the mitotic cell cycle, steroid metabolic

processes and oxidoreductase activity (**Fig 5**) were significantly enriched for the genes with

significantly increased expression for the response contrasts, which is in agreement with

results from a previous study using the same animals [33]. The overlap and grouping of the

GO terms between the blood and lymph node samples illustrates the similarity of responses to

infection for these tissues. The lack of GO terms significantly enriched for genes with

significantly increased expression for the response contrasts in the spleen samples (**Fig 5**) can

be explained by the lower numbers of significant DEGs with increased expression for these

samples (**Fig 3**, **S5**). A larger range of GO terms were significantly enriched for the genes

with significantly decreased expression for the response contrasts (**Fig 5**). This is likely due

to the higher number of significant DEGs with decreased expression, which would allow

more GO terms to be significantly enriched for these genes (**Fig 3**). The GO terms

significantly enriched for the genes with significantly decreased expression for the response

contrasts included those related to multicellular development and morphogenesis, receptor-

mediated endocytosis, cell migration and motility, cargo receptor activity, extracellular

region, and lipopolysaccharide binding (**Fig 5**). These observations are also in agreement

with results of the previous analysis using the same animals [33].

It is logical that the GO terms that are significantly enriched for genes with

significantly decreased expression for the response contrasts are related to blood coagulation

in the PBMC samples (**Fig 5**, **S28**). The effects of blood-borne trypanosome parasites on the

blood of the host have been long studied in both cattle and humans [133-135]. Notably,

anaemia is the main cause of death due trypanosomiasis and the ability to control this

anaemia is consider critical to trypanotolerance in cattle [136, 137]. While trypanosomes do

obtain iron from their environment in the blood of the host and that iron homeostasis and

metabolism in the trypanosome parasite are known to be essential for infection, it has been

observed that anaemia does not correlate with parasitaemia [138-141]. Additionally,

trypanosomes require much less iron than a mammalian cell [142, 143]. The anaemia caused

by trypanosome infection is therefore considered to be an immune response, which may be

driven by cytokines [137, 144, 145].

During trypanosome infection in cattle, control of anaemia and parasitaemia are key components of trypanotolerance but are considered to represent separate genetically determined traits [146]. This is exemplified by genes related to iron ion homeostasis such as the solute carrier family 40 member 1 gene (*SLC40A1*) [147], which was noted to be among the most divergent between trypanotolerant N'Dama and trypanosusceptible Boran as a result of a marked reduction in expression in the N'Dama population over the course of the infection experiment relative to the pre-infection level in PBMC [40]. *SLC40A1* was also in the top 10 most significant DEGs with decreased expression in the blood samples at 34 dpi for the response contrast in this study, highlighting the divergent nature of the expression of this gene between the N'Dama and Boran populations during trypanosome infection (**Table 2**). The reduction of cellular iron export by the SLC40A1 protein is thought to be a component of an innate immune-driven strategy to prevent bloodstream pathogens from accessing iron, with anaemia as a side effect for the host [137, 140, 144]. A related gene, the solute carrier family 11 member 1 gene (*SLC11A1*), which regulates iron homeostasis in macrophages was found to be differentially expressed between N'Dama and African indicine cattle during trypanosome infection using RNA-seq data [42] and variants of this gene are associated with susceptibility to several infectious diseases, including tuberculosis in cattle [148, 149]. Finally, it is noteworthy that the previous work by Noyes and colleagues [33], which used a different functional enrichment approach also highlighted biological pathways related to iron ion homeostasis.

The significantly enriched GO terms showed similar patterns for the direct contrasts with those enriched for the genes with significantly increased expression related to organelle metabolic processes for multiple tissues, while those enriched for the genes with significantly decreased expression showed a greater range of processes (**Fig S32**). Gene ontology terms related to coagulation were also significantly enriched for genes with significantly decreased

expression in the PBMC samples (**Fig S32**). For the N'Dama and Boran contrasts, the

significantly enriched GO terms were related to cell regulation across all tissues, time points,

and direction of expression (**Fig S33**, **S34**). This is likely a result of the higher numbers of

significant DEGs for these contrasts, which led to higher numbers of significantly enriched

GO terms that were too varied for the EnrichmentMap method to summarise effectively (**Fig**

3) [84].

In conclusion, trypanotolerant N'Dama and trypanosusceptible Boran cattle responded

in largely similar ways during trypanosome infection when gene expression was examined

using PBMC, liver, lymph node, and spleen samples with peaks and troughs of gene

expression following the cyclic pattern of parasitaemia exhibited during trypanosome

infection. Differences in response to infection between the two populations include genes

related to the immune system such as those encoding antimicrobial peptides and cytokines.

Within the PBMC samples, differences in genes relating to coagulation and iron homeostasis

support the hypothesis that the dual abilities to control both parasitaemia and the anaemia

resulting from the innate immune response to trypanosome parasites are key to

trypanotolerance. This work adds to our understanding of host-trypanosome interactions

across multiple tissues and time points as well as to our knowledge of how genetic variation

underlying the host response can lead to differential host tolerance of, or susceptibility to,

trypanosome infection. This improves our general understanding of mammalian trypanosome

infection which may also be applicable to human African trypanosomiasis.

Acknowledgements

We thank Morris Agaba, Olivier Hanotte, Stephen J. Kemp, and Stephen V. Gordon for assistance with sample resources and for useful scientific discussion.

References

References

1.   Steverding D. The history of African trypanosomiasis. Parasit Vectors. 2008;1(1):3.

Epub 20080212. https://doi.org/10.1186/1756-3305-1-3. PubMed PMID: 18275594.

2.   Brun R, Blum J, Chappuis F, Burri C. Human African trypanosomiasis. Lancet.

2010;375(9709):148-59. Epub 20091014. https://doi.org/10.1016/S0140-6736(09)60829-1.

PubMed PMID: 19833383.

3.   Taylor KA. Immune responses of cattle to African trypanosomes: protective or

pathogenic? Int J Parasitol. 1998;28(2):219-40. https://doi.org/10.1016/s0020-

7519(97)00154-9. PubMed PMID: 9512986.

4.   Taylor KA, Mertens B. Immune response of cattle infected with African

trypanosomes. Mem Inst Oswaldo Cruz. 1999;94(2):239-44. https://doi.org/10.1590/s0074-

02761999000200022. PubMed PMID: 10224536.

5.   Ibrahim MAM, Weber JS, Ngomtcho SCH, Signaboubo D, Berger P, Hassane HM, et

al. Diversity of trypanosomes in humans and cattle in the HAT foci Mandoul and Maro,

Southern Chad-A matter of concern for zoonotic potential? PLoS Negl Trop Dis.

2021;15(6):e0009323. Epub 20210609. https://doi.org/10.1371/journal.pntd.0009323.

PubMed PMID: 34106914.

6.   Hounyèmè RE, Kaboré J, Gimonneau G, Somda MB, Salou E, Missihoun AA, et al.

Molecular epidemiology of Animal African Trypanosomosis in southwest Burkina Faso.

PLoS Negl Trop Dis. 2022;16(8):e0010106. Epub 20220822.

https://doi.org/10.1371/journal.pntd.0010106. PubMed PMID: 35994491.

7.   N'Djetchi MK, Ilboudo H, Koffi M, Kaboré J, Kaboré JW, Kaba D, et al. The study of

trypanosome species circulating in domestic animals in two human African trypanosomiasis

foci of Côte d'Ivoire identifies pigs and cattle as potential reservoirs of Trypanosoma brucei

gambiense. PLoS Negl Trop Dis. 2017;11(10):e0005993. Epub 20171018.

https://doi.org/10.1371/journal.pntd.0005993. PubMed PMID: 29045405.

8.   Antoine-Moussiaux N, Magez S, Desmecht D. Contributions of experimental mouse

models to the understanding of African trypanosomiasis. Trends Parasitol. 2008;24(9):411-8.

Epub 20080804. https://doi.org/10.1016/j.pt.2008.05.010. PubMed PMID: 18684669.

9.   Magez S, Caljon G. Mouse models for pathogenic African trypanosomes: unravelling
the immunology of host-parasite-vector interactions. Parasite Immunol. 2011;33(8):423-9.
https://doi.org/10.1111/j.1365-3024.2011.01293.x. PubMed PMID: 21480934.

10.   Moulton JE, Stevens DR. Animal model of human disease: trypanosomiasis, sleeping
sickness. Am J Pathol. 1978;91(3):693-6. PubMed PMID: 655266.

11.   Kao WH, Klag MJ, Meoni LA, Reich D, Berthier-Schaad Y, Li M, et al. *MYH9* is
associated with nondiabetic end-stage renal disease in African Americans. Nat Genet.
2008;40(10):1185-92. Epub 2008/09/17. https://doi.org/10.1038/ng.232. PubMed PMID:
18794854.

12.   Genovese G, Friedman DJ, Ross MD, Lecordier L, Uzureau P, Freedman BI, et al.
Association of trypanolytic ApoL1 variants with kidney disease in African Americans.
Science. 2010;329(5993):841-5. Epub 2010/07/22. https://doi.org/10.1126/science.1193032.
PubMed PMID: 20647424.

13.   Berthier D, Breniere SF, Bras-Goncalves R, Lemesre JL, Jamonneau V, Solano P, et
al. Tolerance to trypanosomatids: a threat, or a key for disease elimination? Trends Parasitol.
2016;32(2):157-68.   Epub   20151128.   https://doi.org/10.1016/j.pt.2015.11.001.   PubMed
PMID: 26643519.

14.   Kemp SJ, Teale AJ. Genetic basis of trypanotolerance in cattle and mice. Parasitol
Today. 1998;14(11):450-4. https://doi.org/10.1016/s0169-4758(98)01334-9. PubMed PMID:
17040846.

15.   Truc P, Büscher P, Cuny G, Gonzatti MI, Jannin J, Joshi P, et al. Atypical human
infections by animal trypanosomes. PLoS Negl Trop Dis. 2013;7(9):e2256. Epub 20130912.
https://doi.org/10.1371/journal.pntd.0002256. PubMed PMID: 24069464.

16.   Kumar R, Gupta S, Bhutia WD, Vaid RK, Kumar S. Atypical human trypanosomosis:
Potentially emerging disease with lack of understanding. Zoonoses Public Health.
2022;69(4):259-76. Epub 20220330. https://doi.org/10.1111/zph.12945. PubMed PMID:
35355422.

17.   Okello I, Mafie E, Eastwood G, Nzalawahe J, Mboera LEG. African animal
trypanosomiasis: a systematic review on prevalence, risk factors and drug resistance in sub-

Saharan Africa. J Med Entomol. 2022;59(4):1099-143. https://doi.org/10.1093/jme/tjac018.
PubMed PMID: 35579072.

18.   Herrero M, Thornton PK. Livestock and global change: emerging issues for
sustainable food systems. Proc Natl Acad Sci U S A. 2013;110(52):20878-81. Epub
2013/12/18. https://doi.org/10.1073/pnas.1321844111. PubMed PMID: 24344313.

19.   Thornton PK. Livestock production: recent trends, future prospects. Philos Trans R
Soc   Lond   B   Biol   Sci.   2010;365(1554):2853-67.   Epub   2010/08/18.
https://doi.org/10.1098/rstb.2010.0134. PubMed PMID: 20713389.

20.   Hanotte O, Bradley DG, Ochieng JW, Verjee Y, Hill EW, Rege JE. African
pastoralism: genetic imprints of origins and migrations. Science. 2002;296(5566):336-9.
PubMed PMID: 11951043.

21.   MacHugh DE, Shriver MD, Loftus RT, Cunningham P, Bradley DG. Microsatellite
DNA variation and the evolution, domestication and phylogeography of taurine and zebu
cattle (Bos taurus and Bos indicus). Genetics. 1997;146(3):1071-86. Epub 1997/07/01.
https://doi.org/10.1093/genetics/146.3.1071. PubMed PMID: 9215909.

22.   Kim J, Hanotte O, Mwai OA, Dessie T, Bashir S, Diallo B, et al. The genome
landscape of indigenous African cattle. Genome Biol. 2017;18(1):34. Epub 20170220.
https://doi.org/10.1186/s13059-017-1153-y. PubMed PMID: 28219390.

23.   Mwai O, Hanotte O, Kwon YJ, Cho S. African indigenous cattle: unique genetic
resources in a rapidly changing world. Asian-Australas J Anim Sci. 2015;28(7):911-21. Epub
2015/06/25. https://doi.org/10.5713/ajas.15.0002R. PubMed PMID: 26104394.

24.   Holden CJ. Bantu language trees reflect the spread of farming across sub-Saharan
Africa: a   maximum-parsimony   analysis.   Proc   Biol   Sci.   2002;269(1493):793-9.
https://doi.org/10.1098/rspb.2002.1955. PubMed PMID: 11958710.

25.   Tishkoff SA, Reed FA, Friedlaender FR, Ehret C, Ranciaro A, Froment A, et al. The
genetic   structure   and   history   of   Africans   and   African   Americans.   Science.
2009;324(5930):1035-44. Epub 20090430. https://doi.org/10.1126/science.1172257. PubMed
PMID: 19407144.

26.   Yaro M, Munyard KA, Stear MJ, Groth DM. Combatting African animal
trypanosomiasis (AAT) in livestock: the potential role of trypanotolerance. Vet Parasitol.
2016;225:43-52. Epub 20160504. https://doi.org/10.1016/j.vetpar.2016.05.003. PubMed
PMID: 27369574.

27.   Freeman AR, Meghen CM, MacHugh DE, Loftus RT, Achukwi MD, Bado A, et al.
Admixture and diversity in West African cattle populations. Mol Ecol. 2004;13(11):3477-87.
https://doi.org/10.1111/j.1365-294X.2004.02311.x. PubMed PMID: 15488005.

28.   Murray M, Trail JC, D'Ieteren GD. Trypanotolerance in cattle and prospects for the
control of trypanosomiasis by selective breeding. Rev Sci Tech. 1990;9(2):369-86.
https://doi.org/10.20506/rst.9.2.506. PubMed PMID: 2132686.

29.   Murray M, Black SJ. African trypanosomiasis in cattle: working with nature's
solution. Vet Parasitol. 1985;18(2):167-82. https://doi.org/10.1016/0304-4017(85)90065-2.
PubMed PMID: 3898544.

30.   d'Ieteren GD, Authié E, Wissocq N, Murray M. Trypanotolerance, an option for
sustainable livestock production in areas at risk from trypanosomosis. Rev Sci Tech.
1998;17(1):154-75. https://doi.org/10.20506/rst.17.1.1088. PubMed PMID: 9638808.

31.   Trail JC, D'Ieteren GD, Teale AJ. Trypanotolerance and the value of conserving
livestock genetic resources. Genome. 1989;31(2):805-12. https://doi.org/10.1139/g89-142.
PubMed PMID: 2534387.

32.   Smetko A, Soudre A, Silbermayr K, Muller S, Brem G, Hanotte O, et al.
Trypanosomosis: potential driver of selection in African cattle. Front Genet. 2015;6:137.
Epub 20150421. https://doi.org/10.3389/fgene.2015.00137. PubMed PMID: 25964796.

33.   Noyes H, Brass A, Obara I, Anderson S, Archibald AL, Bradley DG, et al. Genetic
and expression analysis of cattle identifies candidate genes in pathways responding to
Trypanosoma congolense infection. Proc Natl Acad Sci U S A. 2011;108(22):9304-9. Epub
20110518. https://doi.org/10.1073/pnas.1013486108. PubMed PMID: 21593421.

34.   Naessens J, Teale AJ, Sileghem M. Identification of mechanisms of natural resistance
to African trypanosomiasis in cattle. Vet Immunol Immunopathol. 2002;87(3-4):187-94.
https://doi.org/10.1016/s0165-2427(02)00070-3. PubMed PMID: 12072233.

35.   Trail JC, d'Ieteren GD, Maille JC, Yangari G. Genetic aspects of control of anaemia

development in trypanotolerant N'Dama cattle. Acta Trop. 1991;48(4):285-91.

https://doi.org/10.1016/0001-706x(91)90016-d. PubMed PMID: 1674402.

36.   Gautier M, Flori L, Riebler A, Jaffrezic F, Laloe D, Gut I, et al. A whole genome

Bayesian scan for adaptive genetic divergence in West African cattle. BMC Genomics.

2009;10:550. Epub 20091121. https://doi.org/10.1186/1471-2164-10-550. PubMed PMID:

19930592.

37.   Kim SJ, Ka S, Ha JW, Kim J, Yoo D, Kim K, et al. Cattle genome-wide analysis

reveals genetic signatures in trypanotolerant N'Dama. BMC Genomics. 2017;18(1):371. Epub

20170512. https://doi.org/10.1186/s12864-017-3742-2. PubMed PMID: 28499406.

38.   Hill EW, O'Gorman GM, Agaba M, Gibson JP, Hanotte O, Kemp SJ, et al.

Understanding bovine trypanosomiasis and trypanotolerance: the promise of functional

genomics.   Vet   Immunol   Immunopathol.   2005;105(3-4):247-58.

https://doi.org/10.1016/j.vetimm.2005.02.004. PubMed PMID: 15808304.

39.   Meade KG, O'Gorman GM, Hill EW, Narciandi F, Agaba M, Kemp SJ, et al.

Divergent antimicrobial peptide (AMP) and acute phase protein (APP) responses to

Trypanosoma congolense infection in trypanotolerant and trypanosusceptible cattle. Mol

Immunol.   2009;47(2-3):196-204.   Epub   20091103.

https://doi.org/10.1016/j.molimm.2009.09.042. PubMed PMID: 19889461.

40.   O'Gorman GM, Park SD, Hill EW, Meade KG, Coussens PM, Agaba M, et al.

Transcriptional profiling of cattle infected with *Trypanosoma congolense* highlights gene

expression signatures underlying trypanotolerance and trypanosusceptibility. BMC

Genomics. 2009;10:207. Epub 20090501. https://doi.org/10.1186/1471-2164-10-207.

PubMed PMID: 19409086.

41.   O'Gorman GM, Park SD, Hill EW, Meade KG, Mitchell LC, Agaba M, et al. Cytokine

mRNA profiling of peripheral blood mononuclear cells from trypanotolerant and

trypanosusceptible cattle infected with *Trypanosoma congolense*. Physiol Genomics.

2006;28(1):53-61. Epub 20060919. https://doi.org/10.1152/physiolgenomics.00100.2006.

PubMed PMID: 16985010.

42.   Peylhard M, Berthier D, Dayo G-K, Chantal I, Sylla S, Nidelet S, et al. Whole blood

transcriptome profiles of trypanotolerant and trypanosusceptible cattle highlight a differential

modulation of metabolism and immune response during infection by *Trypanosoma*

congolense. Peer Community J. 2023;3. https://doi.org/10.24072/pcjournal.239.

43.   Dayo GK, Gautier M, Berthier D, Poivey JP, Sidibe I, Bengaly Z, et al. Association

studies in QTL regions linked to bovine trypanotolerance in a West African crossbred

population. Anim Genet. 2012;43(2):123-32. Epub 20110722. https://doi.org/10.1111/j.1365-

2052.2011.02227.x. PubMed PMID: 22404348.

44.   Bahbahani H, Afana A, Wragg D. Genomic signatures of adaptive introgression and

environmental adaptation in the Sheko cattle of southwest Ethiopia. PLoS ONE.

2018;13(8):e0202479.   Epub   20180816.   https://doi.org/10.1371/journal.pone.0202479.

PubMed PMID: 30114214.

45.   Berthier D, Peylhard M, Dayo GK, Flori L, Sylla S, Bolly S, et al. A comparison of

phenotypic traits related to trypanotolerance in five west african cattle breeds highlights the

value of shorthorn taurine breeds. PLoS ONE. 2015;10(5):e0126498. Epub 2015/05/09.

https://doi.org/10.1371/journal.pone.0126498. PubMed PMID: 25954819.

46.   Alvarez I, Perez-Pardal L, Traore A, Fernandez I, Goyache F. Lack of haplotype

structuring for two candidate genes for trypanotolerance in cattle. J Anim Breed Genet.

2016;133(2):105-14. Epub 20150914. https://doi.org/10.1111/jbg.12181. PubMed PMID:

26365013.

47.   Goyache F, Perez-Pardal L, Fernandez I, Traore A, Menendez-Arias NA, Alvarez I.

Ancient autozygous segments subject to positive selection suggest adaptive immune

responses   in   West   African   cattle.   Gene.   2021;803:145899.   Epub   20210813.

https://doi.org/10.1016/j.gene.2021.145899. PubMed PMID: 34400278.

48.   Rajavel A, Heinrich F, Schmitt AO, Gultas M. Identifying cattle breed-specific partner

choice of transcription factors during the African trypanosomiasis disease progression using

bioinformatics   analysis.   Vaccines   (Basel).   2020;8(2).   Epub   20200523.

https://doi.org/10.3390/vaccines8020246. PubMed PMID: 32456126.

49.   Rajavel A, Klees S, Hui Y, Schmitt AO, Gultas M. Deciphering the molecular
mechanism underlying African animal trypanosomiasis by means of the 1000 Bull Genomes
Project genomic dataset. Biology (Basel). 2022;11(5). Epub 20220513.
https://doi.org/10.3390/biology11050742. PubMed PMID: 35625470.

50.   Geigy R, Kauffmann M. Sleeping sickness survey in the Serengeti area (Tanzania)
1971. I. Examination of large mammals for trypanosomes. Acta Trop. 1973;30(1):12-23.
PubMed PMID: 4144952.

51.   Nantulya VM, Musoke AJ, Rurangirwa FR, Moloo SK. Resistance of cattle to tsetse-
transmitted challenge with *Trypanosoma brucei* or *Trypanosoma congolense* after
spontaneous recovery from syringe-passaged infections. Infect Immun. 1984;43(2):735-8.
https://doi.org/10.1128/iai.43.2.735-738.1984. PubMed PMID: 6693173.

52.   Akol GW, Murray M. Early events following challenge of cattle with tsetse infected
with *Trypanosoma congolense*: development of the local skin reaction. Vet Rec.
1982;110(13):295-302. https://doi.org/10.1136/vr.110.13.295. PubMed PMID: 7072104.

53.   Dwinger RH, Murray M, Moloo SK. Potential value of localized skin reactions
(chancres) induced by *Trypanosoma congolense* transmitted by *Glossina morsitans centralis*
for the analysis of metacyclic trypanosome populations. Parasite Immunol. 1987;9(3):353-62.
https://doi.org/10.1111/j.1365-3024.1987.tb00513.x. PubMed PMID: 3601446.

54.   Frerebeau N. khroma: Colour Schemes for Scientific Data Visualization. manual.
2023 2023. Report No.

55.   Garnier, Simon, Ross, Noam, Rudis, Robert, et al. viridis(Lite) - Colorblind-Friendly
Color Maps for R. manual. 2023 2023. Report No.

56.   Gautier L, Cope L, Bolstad BM, Irizarry RA. *affy*—analysis of Affymetrix GeneChip
data at the probe level. Bioinformatics. 2004;20(3):307-15.
https://doi.org/10.1093/bioinformatics/btg405. PubMed PMID: 14960456.

57.   Huber W, Carey VJ, Gentleman R, Anders S, Carlson M, Carvalho BS, et al.
Orchestrating high-throughput genomic analysis with Bioconductor. Nat Methods.
2015;12(2):115-21. https://doi.org/10.1038/nmeth.3252. PubMed PMID: 25633503.

58.   Kauffmann A, Gentleman R, Huber W. arrayQualityMetrics—a bioconductor package
for quality assessment of microarray data. Bioinformatics. 2009;25(3):415-6. Epub
20081223. https://doi.org/10.1093/bioinformatics/btn647. PubMed PMID: 19106121.

59.   R Core Team. R: A Language and Environment for Statistical Computing. manual.
Vienna, Austria: R Foundation for Statistical Computing, 2023 2023. Report No.

60.   Rue-Albrecht K, Magee DA, Killick KE, Nalpas NC, Gordon SV, MacHugh DE.
Comparative functional genomics and the bovine macrophage response to strains of the
Mycobacterium genus. Front Immunol. 2014;5:536. Epub 20141105.
https://doi.org/10.3389/fimmu.2014.00536. PubMed PMID: 25414700.

61.   Hochreiter S, Clevert DA, Obermayer K. A new summarization method for
Affymetrix probe level data. Bioinformatics. 2006;22(8):943-9. Epub 20060210.
https://doi.org/10.1093/bioinformatics/btl033. PubMed PMID: 16473874.

62.   Wickham H, François R, Henry L, Müller K, Vaughan D. dplyr: A Grammar of Data
Manipulation. manual. 2023 2023. Report No.

63.   van den Brand T. ggh4x: Hacks for 'ggplot2'. manual. 2023 2023. Report No.

64.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer; 2009.

65.   Wilke CO, Wiernik BM. ggtext: Improved Text Rendering Support for 'ggplot2'.
manual. 2022.

66.   Bache SM, Wickham H. magrittr: A Forward-Pipe Operator for R. manual. 2022
2022. Report No.

67.   Wickham H, Hester J, Bryan J. readr: Read Rectangular Text Data. manual. 2023
2023. Report No.

68.   Wickham H. Reshaping data with the reshape package. J Stat Softw. 2007;21(12):1 -
20. https://doi.org/10.18637/jss.v021.i12.

69.   Wickham H. stringr: Simple, Consistent Wrappers for Common String Operations.
manual. 2023 2023. Report No.

70.   Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers

differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids

Res. 2015;43(7):e47. Epub 20150120. https://doi.org/10.1093/nar/gkv007. PubMed PMID:

25605792.

71.   Pedersen TL. patchwork: The Composer of Plots. manual. 2023 2023. Report No.

72.   Smyth GK, Michaud J, Scott HS. Use of within-array replicate spots for assessing

differential expression in microarray experiments. Bioinformatics. 2005;21(9):2067-75. Epub

20050118. https://doi.org/10.1093/bioinformatics/bti270. PubMed PMID: 15657102.

73.   Smyth GK. Linear models and empirical bayes methods for assessing differential

expression in microarray experiments. Stat Appl Genet Mol Biol. 2004;3:Article3. Epub

20040212. https://doi.org/10.2202/1544-6115.1027. PubMed PMID: 16646809.

74.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. J Roy Stat Soc B Met. 1995;57(1):289-300.

75.   Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. g:Profiler-

interoperable web service for functional enrichment analysis and gene identifier mapping

(2023   update).   Nucleic   Acids   Res.   2023;51(W1):W207-W12.

https://doi.org/10.1093/nar/gkad347. PubMed PMID: 37144459.

76.   Krassowski M. ComplexUpset. manual. 2020 2020. Report No.

77.   Slowikowski K. ggrepel: Automatically Position Non-Overlapping Text Labels with

'ggplot2'. manual. 2023 2023. Report No.

78.   Henry L, Wickham H. rlang: Functions for Base Types and Core R and 'tidyverse'

Features. manual. 2024 2024. Report No.

79.   Wickham H, Vaughan D, Girlich M. tidyr: Tidy Messy Data. manual. 2023 2023.

Report No.

80.   Ooms J. magick: Advanced Graphics and Image-Processing in R. manual. 2023 2023.

Report No.

81.   ImageMagick Studio LLC. ImageMagick. 2023.

82.   Wickham H, Henry L. purrr: Functional Programming Tools. manual. 2023 2023.
Report No.

83.   Wickham H, Pedersen TL, Seidel D. scales: Scale Functions for Visualization.
manual. 2023 2023. Report No.

84.   Merico D, Isserlin R, Stueker O, Emili A, Bader GD. Enrichment map: a network-
based method for gene-set enrichment visualization and interpretation. PLoS ONE.
2010;5(11):e13984. Epub 20101115. https://doi.org/10.1371/journal.pone.0013984. PubMed
PMID: 21085593.

85.   Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a
software environment for integrated models of biomolecular interaction networks. Genome
Res. 2003;13(11):2498-504. https://doi.org/10.1101/gr.1239303. PubMed PMID: 14597658.

86.   Reimand J, Isserlin R, Voisin V, Kucera M, Tannus-Lopes C, Rostamianfar A, et al.
Pathway enrichment analysis and visualization of omics data using g:Profiler, GSEA,
Cytoscape   and   EnrichmentMap.   Nat   Protoc.   2019;14(2):482-517.
https://doi.org/10.1038/s41596-018-0103-9. PubMed PMID: 30664679.

87.   Kucera M, Isserlin R, Arkhangorodsky A, Bader GD. AutoAnnotate: A Cytoscape app
for summarizing networks with semantic annotations. F1000Res. 2016;5:1717. Epub
20160715. https://doi.org/10.12688/f1000research.9090.1. PubMed PMID: 27830058.

88.   Morris JH, Apeltsin L, Newman AM, Baumbach J, Wittkop T, Su G, et al.
clusterMaker: a multi-algorithm clustering plugin for Cytoscape. BMC Bioinformatics.
2011;12:436. Epub 20111109. https://doi.org/10.1186/1471-2105-12-436. PubMed PMID:
22070249.

89.   Oesper L, Merico D, Isserlin R, Bader GD. WordCloud: a Cytoscape plugin to create
a visual semantic summary of networks. Source Code Biol Med. 2011;6:7. Epub 20110407.
https://doi.org/10.1186/1751-0473-6-7. PubMed PMID: 21473782.

90.   Becker MY, Rojas I. A graph layout algorithm for drawing metabolic pathways.
Bioinformatics. 2001;17(5):461-7. https://doi.org/10.1093/bioinformatics/17.5.461. PubMed
PMID: 11331241.

91.   Magee DA, Taraktsoglou M, Killick KE, Nalpas NC, Browne JA, Park SD, et al.
Global gene expression and systems biology analysis of bovine monocyte-derived
macrophages in response to in vitro challenge with *Mycobacterium bovis*. PLoS ONE.
2012;7(2):e32034. Epub 20120222. https://doi.org/10.1371/journal.pone.0032034. PubMed
PMID: 22384131.

92.   MacHugh DE, Taraktsoglou M, Killick KE, Nalpas NC, Browne JA, E. PSD, et al.
Pan-genomic analysis of bovine monocyte-derived macrophage gene expression in response
to *in vitro* infection with *Mycobacterium avium* subspecies *paratuberculosis*. Vet Res.
2012;43(1):25. Epub 20120328. https://doi.org/10.1186/1297-9716-43-25. PubMed PMID:
22455317.

93.   Killick KE, Browne JA, Park SD, Magee DA, Martin I, Meade KG, et al. Genome-
wide transcriptional profiling of peripheral blood leukocytes from cattle infected with
Mycobacterium bovis reveals suppression of host immune genes. BMC Genomics.
2011;12:611. Epub 20111219. https://doi.org/10.1186/1471-2164-12-611. PubMed PMID:
22182502.

94.   Ezendam J, Staedtler F, Pennings J, Vandebriel RJ, Pieters R, Harleman JH, et al.
Toxicogenomics of subchronic hexachlorobenzene exposure in Brown Norway rats. Environ
Health Perspect. 2004;112(7):782-91. https://doi.org/10.1289/ehp.112-1241993. PubMed
PMID: 15159207.

95.   Kriegel H-P, Kröger P, Schubert E, Zimek A, editors. A general framework for
increasing the robustness of PCA-based correlation clustering algorithms2008; Berlin,
Heidelberg: Springer Berlin Heidelberg.

96.   McVean G. A genealogical interpretation of principal components analysis. PLoS
Genet. 2009;5(10):e1000686. Epub 20091016. https://doi.org/10.1371/journal.pgen.1000686.
PubMed PMID: 19834557.

97.   Novembre J, Stephens M. Interpreting principal component analyses of spatial
population genetic variation. Nat Genet. 2008;40(5):646-9. Epub 20080420.
https://doi.org/10.1038/ng.139. PubMed PMID: 18425127.

98.   Lewis SM, Williams A, Eisenbarth SC. Structure and function of the immune system

in the spleen. Sci Immunol. 2019;4(33). https://doi.org/10.1126/sciimmunol.aau6085.

PubMed PMID: 30824527.

99.   Buettner M, Bode U. Lymph node dissection – understanding the immunological

function of lymph nodes. Clin Exp Immunol. 2012;169(3):205-12.

https://doi.org/10.1111/j.1365-2249.2012.04602.x. PubMed PMID: 22861359.

100.   Jønsson V. Comparison and definition of spleen and lymph node: a phylogenetic

analysis. J Theor Biol. 1985;117(4):691-9. https://doi.org/10.1016/s0022-5193(85)80247-2.

PubMed PMID: 4094460.

101.   Dalma-Weiszhausz DD, Warrington J, Tanimoto EY, Miyada CG. The Affymetrix

GeneChip platform: an overview. Methods Enzymol. 2006;410:3-28.

https://doi.org/10.1016/S0076-6879(06)10001-4. PubMed PMID: 16938544.

102.   Casey ME, Meade KG, Nalpas NC, Taraktsoglou M, Browne JA, Killick KE, et al.

Analysis of the bovine monocyte-derived macrophage response to *Mycobacterium avium*

subspecies *paratuberculosis* infection using RNA-seq. Front Immunol. 2015;6:23. Epub

20150204. https://doi.org/10.3389/fimmu.2015.00023. PubMed PMID: 25699042.

103.   McLoughlin KE, Nalpas NC, Rue-Albrecht K, Browne JA, Magee DA, Killick KE, et

al. RNA-seq transcriptional profiling of peripheral blood leukocytes from cattle infected with

Mycobacterium bovis. Front Immunol. 2014;5:396. Epub 20140826.

https://doi.org/10.3389/fimmu.2014.00396. PubMed PMID: 25206354.

104.   Nalpas NC, Park SD, Magee DA, Taraktsoglou M, Browne JA, Conlon KM, et al.

Whole-transcriptome, high-throughput RNA sequence analysis of the bovine macrophage

response to *Mycobacterium bovis* infection in vitro. BMC Genomics. 2013;14:230. Epub

20130408. https://doi.org/10.1186/1471-2164-14-230. PubMed PMID: 23565803.

105.   Gitau JK, Macharia RW, Mwangi KW, Ongeso N, Murungi E. Gene co-expression

network identifies critical genes, pathways and regulatory motifs mediating the progression

of Rift Valley Fgorever in *Bos taurus*. Heliyon. 2023;9(7):e18175. Epub 20230714.

https://doi.org/10.1016/j.heliyon.2023.e18175. PubMed PMID: 37519716.

106. Kuiper JWP, Krause J, Potgeter L, Adrian J, Hauck CR. A genome-wide genetic screen identifies CYRI-B as a negative regulator of CEACAM3-mediated phagocytosis. J Cell Sci. 2023;136(11). Epub 20230602. https://doi.org/10.1242/jcs.260771. PubMed PMID: 37264948.

107. Al Barashdi MA, Ali A, McMullin MF, Mills K. Protein tyrosine phosphatase receptor type C (PTPRC or CD45). J Clin Pathol. 2021;74(9):548-52. Epub 20210526. https://doi.org/10.1136/jclinpath-2020-206927. PubMed PMID: 34039664.

108. Goodhead I, Archibald A, Amwayi P, Brass A, Gibson J, Hall N, et al. A comprehensive genetic analysis of candidate genes regulating response to *Trypanosoma congolense* infection in mice. PLoS Negl Trop Dis. 2010;4(11):e880. Epub 20101109. https://doi.org/10.1371/journal.pntd.0000880. PubMed PMID: 21085469.

109. Noyes HA, Alimohammadian MH, Agaba M, Brass A, Fuchs H, Gailus-Durner V, et al. Mechanisms controlling anaemia in *Trypanosoma congolense* infected mice. PLoS ONE. 2009;4(4):e5170. Epub 20090413. https://doi.org/10.1371/journal.pone.0005170. PubMed PMID: 19365556.

110. Guerrero NA, Camacho M, Vila L, Iniguez MA, Chillon-Marinas C, Cuervo H, et al. Cyclooxygenase-2 and prostaglandin E2 signaling through prostaglandin receptor EP-2 favor the development of myocarditis during acute *Trypanosoma cruzi* infection. PLoS Negl Trop Dis. 2015;9(8):e0004025. Epub 20150825. https://doi.org/10.1371/journal.pntd.0004025. PubMed PMID: 26305786.

111. Pham HTT, Magez S, Choi B, Baatar B, Jung J, Radwanska M. Neutrophil metalloproteinase driven spleen damage hampers infection control of trypanosomiasis. Nat Commun. 2023;14(1):5418. Epub 20230905. https://doi.org/10.1038/s41467-023-41089-w. PubMed PMID: 37669943.

112. Antoine-Moussiaux N, Cornet A, Cornet F, Glineur S, Dermine M, Desmecht D. A non-cytosolic protein of *Trypanosoma evansi* induces CD45-dependent lymphocyte death. PLoS ONE. 2009;4(5):e5728. Epub 20090529. https://doi.org/10.1371/journal.pone.0005728. PubMed PMID: 19478957.

113.   Le Coz C, Nguyen DN, Su C, Nolan BE, Albrecht AV, Xhani S, et al. Constrained
chromatin accessibility in PU.1-mutated agammaglobulinemia patients. J Exp Med.
2021;218(7). Epub 20210505. https://doi.org/10.1084/jem.20201750. PubMed PMID:
33951726.

114.   Liu W, Rodgers GP. Olfactomedin 4 expression and functions in innate immunity,
inflammation, and cancer. Cancer Metastasis Rev. 2016;35(2):201-12.
https://doi.org/10.1007/s10555-016-9624-2. PubMed PMID: 27178440.

115.   Tanioku T, Nishibata M, Tokinaga Y, Konno K, Watanabe M, Hemmi H, et al.
Tmem45b is essential for inflammation- and tissue injury-induced mechanical pain
hypersensitivity. Proc Natl Acad Sci U S A. 2022;119(45):e2121989119. Epub 20221102.
https://doi.org/10.1073/pnas.2121989119. PubMed PMID: 36322717.

116.   Röder A, Hüsken S, Hutter MC, Rettie AE, Hanenberg H, Wiek C, et al. Spotlight on
CYP4B1. Int J Mol Sci. 2023;24(3). Epub 20230120. https://doi.org/10.3390/ijms24032038.
PubMed PMID: 36768362.

117.   Banworth MJ, Liang Z, Li G. A novel membrane targeting domain mediates the
endosomal or Golgi localization specificity of small GTPases Rab22 and Rab31. J Biol
Chem. 2022;298(9):102281. Epub 20220719. https://doi.org/10.1016/j.jbc.2022.102281.
PubMed PMID: 35863437.

118.   Jentzsch J, Wunderlich H, Thein M, Bechthold J, Brehm L, Krauss SW, et al.
Microtubule polyglutamylation is an essential regulator of cytoskeletal integrity in
Trypanosoma brucei. J Cell Sci. 2024;137(3). Epub 20240212.
https://doi.org/10.1242/jcs.261740. PubMed PMID: 38205672.

119.   Kumar R, Ali SA, Singh SK, Bhushan V, Mathur M, Jamwal S, et al. Antimicrobial
peptides in farm animals: an updated review on its diversity, function, modes of action and
therapeutic prospects. Vet Sci. 2020;7(4). Epub 20201218.
https://doi.org/10.3390/vetsci7040206. PubMed PMID: 33352919.

120.   Davies D, Meade KG, Herath S, Eckersall PD, Gonzalez D, White JO, et al. Toll-like
receptor and antimicrobial peptide expression in the bovine endometrium. Reprod Biol

Endocrinol. 2008;6:53. Epub 20081118. https://doi.org/10.1186/1477-7827-6-53. PubMed PMID: 19017375.

121. Tetens J, Friedrich JJ, Hartmann A, Schwerin M, Kalm E, Thaller G. The spatial expression pattern of antimicrobial peptides across the healthy bovine udder. J Dairy Sci. 2010;93(2):775-83. https://doi.org/10.3168/jds.2009-2729. PubMed PMID: 20105549.

122. Alva-Murillo N, Tellez-Perez AD, Sagrero-Cisneros E, Lopez-Meza JE, Ochoa-Zarzosa A. Expression of antimicrobial peptides by bovine endothelial cells. Cell Immunol. 2012;280(1):108-12. Epub 20121219. https://doi.org/10.1016/j.cellimm.2012.11.016. PubMed PMID: 23298865.

123. Krause A, Sillard R, Kleemeier B, Kluver E, Maronde E, Conejo-Garcia JR, et al. Isolation and biochemical characterization of LEAP-2, a novel blood peptide expressed in the liver. Protein Sci. 2003;12(1):143-52. https://doi.org/10.1110/ps.0213603. PubMed PMID: 12493837.

124. Whelehan CJ, Barry-Reidy A, Meade KG, Eckersall PD, Chapwanya A, Narciandi F, et al. Characterisation and expression profile of the bovine cathelicidin gene repertoire in mammary tissue. BMC Genomics. 2014;15:128. Epub 20140213. https://doi.org/10.1186/1471-2164-15-128. PubMed PMID: 24524771.

125. McGwire BS, Olson CL, Tack BF, Engman DM. Killing of African trypanosomes by antimicrobial peptides. J Infect Dis. 2003;188(1):146-52. Epub 20030616. https://doi.org/10.1086/375747. PubMed PMID: 12825184.

126. Selsted ME, Tang YQ, Morris WL, McGuire PA, Novotny MJ, Smith W, et al. Purification, primary structures, and antibacterial activities of beta-defensins, a new family of antimicrobial peptides from bovine neutrophils. J Biol Chem. 1993;268(9):6641-8. PubMed PMID: 8454635.

127. Regenhard P, Leippe M, Schubert S, Podschun R, Kalm E, Grotzinger J, et al. Antimicrobial activity of bovine psoriasin. Vet Microbiol. 2009;136(3-4):335-40. Epub 20081206. https://doi.org/10.1016/j.vetmic.2008.12.001. PubMed PMID: 19167844.

128. Uzonna JE, Kaushik RS, Gordon JR, Tabel H. Cytokines and antibody responses during *Trypanosoma congolense* infections in two inbred mouse strains that differ in

resistance. Parasite Immunol. 1999;21(2):57-71. https://doi.org/10.1046/j.1365-3024.1999.00202.x. PubMed PMID: 10101716.

129.   Soares-Silva M, Diniz FF, Gomes GN, Bahia D. The mitogen-activated protein kinase (MAPK) pathway: role in immune evasion by trypanosomatids. Front Microbiol. 2016;7:183. Epub 20160224. https://doi.org/10.3389/fmicb.2016.00183. PubMed PMID: 26941717.

130.   Kaur P, Goyal N. Pathogenic role of mitogen activated protein kinases in protozoan parasites. Biochimie. 2022;193:78-89. Epub 20211024. https://doi.org/10.1016/j.biochi.2021.10.012. PubMed PMID: 34706251.

131.   Kuriakose S, Onyilagha C, Singh R, Olayinka-Adefemi F, Jia P, Uzonna JE. TLR-2 and MyD88-dependent activation of MAPK and STAT proteins regulates proinflammatory cytokine response and immunity to experimental *Trypanosoma congolense* infection. Front Immunol. 2019;10:2673. Epub 20191122. https://doi.org/10.3389/fimmu.2019.02673. PubMed PMID: 31824484.

132.   Pays E, Vanhollebeke B, Vanhamme L, Paturiaux-Hanocq F, Nolan DP, Pérez-Morga D. The trypanolytic factor of human serum. Nat Rev Microbiol. 2006;4(6):477-86. https://doi.org/10.1038/nrmicro1428.

133.   Wellde BT, Chumo DA, Onyango FK, Reardon MJ, Roberts LM, Njogu AR, et al. *Trypanosoma vivax*: disseminated intravascular coagulation in cattle. Ann Trop Med Parasitol. 1989;83 Suppl 1:177-83. https://doi.org/10.1080/00034983.1989.11812422. PubMed PMID: 2619392.

134.   Greenwood BM, Whittle HC. Coagulation studies in Gambian trypanosomiasis. Am J Trop Med Hyg. 1976;25(3):390-4. https://doi.org/10.4269/ajtmh.1976.25.390. PubMed PMID: 937630.

135.   Choudhuri S, Garg NJ. Platelets, macrophages, and thromboinflammation in Chagas disease. J Inflamm Res. 2022;15:5689-706. Epub 20221004. https://doi.org/10.2147/JIR.S380896. PubMed PMID: 36217453.

136.   Naessens J. Bovine trypanotolerance: A natural ability to prevent severe anaemia and haemophagocytic syndrome? Int J Parasitol. 2006;36(5):521-8. Epub 20060313. https://doi.org/10.1016/j.ijpara.2006.02.012. PubMed PMID: 16678182.

137. Stijlemans B, Beschin A, Magez S, Van Ginderachter JA, De Baetselier P. Iron
homeostasis and *Trypanosoma brucei* associated immunopathogenicity development: a
battle/quest for iron. Biomed Res Int. 2015;2015:819389. Epub 20150518.
https://doi.org/10.1155/2015/819389. PubMed PMID: 26090446.

138. Magez S, Schwegmann A, Atkinson R, Claes F, Drennan M, De Baetselier P, et al.
The role of B-cells and IgM antibodies in parasitemia, anemia, and VSG switching in
Trypanosoma brucei-infected mice. PLoS Pathog. 2008;4(8):e1000122. Epub 20080808.
https://doi.org/10.1371/journal.ppat.1000122. PubMed PMID: 18688274.

139. Gilabert Carbajo C, Cornell LJ, Madbouly Y, Lai Z, Yates PA, Tinti M, et al. Novel
aspects of iron homeostasis in pathogenic bloodstream form *Trypanosoma brucei*. PLoS
Pathog. 2021;17(6):e1009696. Epub 20210623. https://doi.org/10.1371/journal.ppat.1009696.
PubMed PMID: 34161395.

140. Taylor MC, Kelly JM. Iron metabolism in trypanosomatids, and its crucial role in
infection. Parasitology. 2010;137(6):899-917. Epub 20100215.
https://doi.org/10.1017/S0031182009991880. PubMed PMID: 20152063.

141. Dick CF, Alcantara CL, Carvalho-Kelly LF, Lacerda-Abreu MA, Cunha ESNL,
Meyer-Fernandes JR, et al. Iron uptake controls *Trypanosoma cruzi* metabolic shift and cell
proliferation. Antioxidants (Basel). 2023;12(5). Epub 20230422.
https://doi.org/10.3390/antiox12050984. PubMed PMID: 37237850.

142. Basu S, Horakova E, Lukes J. Iron-associated biology of *Trypanosoma brucei*.
Biochim Biophys Acta. 2016;1860(2):363-70. Epub 20151030.
https://doi.org/10.1016/j.bbagen.2015.10.027. PubMed PMID: 26523873.

143. Steverding D. Bloodstream forms of *Trypanosoma brucei* require only small amounts
of iron for growth. Parasitol Res. 1998;84(1):59-62. https://doi.org/10.1007/s004360050357.
PubMed PMID: 9491428.

144. Stijlemans B, Vankrunkelsven A, Brys L, Magez S, De Baetselier P. Role of iron
homeostasis in trypanosomiasis-associated anemia. Immunobiology. 2008;213(9-10):823-35.
Epub 20080907. https://doi.org/10.1016/j.imbio.2008.07.023. PubMed PMID: 18926297.

145.   Stijlemans B, Vankrunkelsven A, Caljon G, Bockstal V, Guilliams M, Bosschaerts T,

et al. The central role of macrophages in trypanosomiasis-associated anemia: rationale for

therapeutical approaches. Endocr Metab Immune Disord Drug Targets. 2010;10(1):71-82.

https://doi.org/10.2174/187153010790827966. PubMed PMID: 20158497.

146.   Naessens J, Leak SG, Kennedy DJ, Kemp SJ, Teale AJ. Responses of bovine

chimaeras combining trypanosomosis resistant and susceptible genotypes to experimental

infection with Trypanosoma congolense. Vet Parasitol. 2003;111(2-3):125-42.

https://doi.org/10.1016/s0304-4017(02)00360-6. PubMed PMID: 12531289.

147.   Donovan A, Lima CA, Pinkus JL, Pinkus GS, Zon LI, Robine S, et al. The iron

exporter ferroportin/Slc40a1 is essential for iron homeostasis. Cell Metab. 2005;1(3):191-

200. https://doi.org/10.1016/j.cmet.2005.01.003. PubMed PMID: 16054062.

148.   Holder A, Garty R, Elder C, Mesnard P, Laquerbe C, Bartens MC, et al. Analysis of

genetic variation in the bovine *SLC11A1* gene, its influence on the expression of *NRAMP1*

and potential association with resistance to bovine tuberculosis. Front Microbiol.

2020;11:1420. Epub 20200630. https://doi.org/10.3389/fmicb.2020.01420. PubMed PMID:

32714308.

149.   Archer NS, Nassif NT, O'Brien BA. Genetic variants of SLC11A1 are associated with

both autoimmune and infectious diseases: systematic review and meta-analysis. Genes

Immun. 2015;16(4):275-83. Epub 20150409. https://doi.org/10.1038/gene.2015.8. PubMed

PMID: 25856512.

1163

Supporting information captions

Table S1. Contrast ID, contrast type, tissue, days post infection (dpi), and formula for each of the 64 contrasts.

Table S2. Differential expression analysis results including the probe ID, gene ID, $\log_2$ fold change, and $P_{adj.}$ for each of the 64 contrasts.

Table S3. Tissue, days post infection (dpi), and the top 10 most significant genes with increased and decreased expression with valid gene symbols for the direct contrasts.

Table S4. Tissue, days post infection (dpi), and the top 10 most significant genes with increased and decreased expression with valid gene symbols for the N'Dama contrasts.

Table S5. Tissue, days post infection (dpi), and the top 10 most significant genes with increased and decreased expression with valid gene symbols for the Boran contrasts.

Fig S1. Boxplots showing the $\log_2$ expression intensity of the probe sets for each sample of the **A.** raw and **B.** normalised data after quality control filtering separated into NDAM and BORA populations and coloured according to tissue. The position along the horizontal axis indicates the days post infection. Outlier probe sets are shown as grey dots.

Fig S2. A. Principal component analysis (PCA) of the microarray data set with samples coloured according to days post infection (dpi) with the outer colour representing the tissue and shape indicating the population showing the first and third principal components (PC1 and PC3), and **B.** bar chart of proportion of variance of the top ten PCs.

Fig S3. A. Principal component analysis (PCA) of the microarray data set with samples coloured according to days post infection (dpi) with the outer colour representing the tissue and shape indicating the population showing the first and fourth principal components (PC1 and PC4), and **B.** bar chart of proportion of variance of the top ten PCs.

Fig S4. A. Principal component analysis (PCA) of the microarray data set with samples coloured according to tissue and shape indicating the population showing the first and fifth principal components (PC1 and PC5), and **B.** bar chart of proportion of variance of the top ten PCs.

Fig S5. Bar chart showing the numbers of significantly differentially expressed genes for the RESP contrasts. The extent of the bar above and below 0 on the vertical axis indicates the numbers of significantly differentially expressed genes (DEGs) with increased and decreased expression, respectively. The position on the horizontal indicates the number of days post infection (dpi) and the colour of the bars represents the tissue.

Fig S6. Bar chart showing the numbers of significantly differentially expressed genes for the
DIRE contrasts. The extent of the bar above and below 0 on the vertical axis indicates the
numbers of significantly differentially expressed genes (DEGs) with increased and decreased
expression, respectively. The position on the horizontal indicates the number of days post
infection (dpi) and the colour of the bars represents the tissue.

Fig S7. Bar chart showing the numbers of significantly differentially expressed genes for the
NDAM contrasts. The extent of the bar above and below 0 on the vertical axis indicates the
numbers of significantly differentially expressed genes (DEGs) with increased and decreased
expression, respectively. The position on the horizontal indicates the number of days post
infection (dpi) and the colour of the bars represents the tissue.

Fig S8. Bar chart showing the numbers of significantly differentially expressed genes for the
BORA contrasts. The extent of the bar above and below 0 on the vertical axis indicates the
numbers of significantly differentially expressed genes (DEGs) with increased and decreased
expression, respectively. The position on the horizontal indicates the number of days post
infection (dpi) and the colour of the bars represents the tissue.

Fig S9. UpSet plot showing the top 20 intersections among all 64 contrasts. The horizontal
bars indicate the total number of significant differentially expressed genes (DEGs) for each
contrast while the vertical bars indicate the number of significant DEGs in common between
the contrasts annotated with black dots connected by lines in the intersection matrix. The
background colour of the stripes in the intersection matrix and horizontal bars represents
the tissue. The colour of the bars represents the contrast type with black bars indicating an
overlap between different contrast types. The outline colour of the bars also represents the
tissue with no outline representing an overlap between different tissues.

Fig S10. UpSet plot showing the top 20 intersections among the RESP contrasts. The
horizontal bars indicate the total number of significant differentially expressed genes (DEGs)
for each contrast while the vertical bars indicate the number of significant DEGs in common
between the contrasts annotated with black dots connected by lines in the intersection
matrix. The background colour of the stripes in the intersection matrix and horizontal bars
represents the tissue. The colour of the bars represents the days post infection (dpi) with
black bars indicating an overlap between different timepoints. The outline colour of the bars
also represents the tissue with no outline representing an overlap between different tissues.

Fig S11. UpSet plot showing the top 20 intersections among the DIRE contrasts. The
horizontal bars indicate the total number of significant differentially expressed genes (DEGs)
for each contrast while the vertical bars indicate the number of significant DEGs in common
between the contrasts annotated with black dots connected by lines in the intersection
matrix. The background colour of the stripes in the intersection matrix and horizontal bars
represents the tissue. The colour of the bars represents the days post infection (dpi) with

black bars indicating an overlap between different timepoints. The outline colour of the bars
also represents the tissue with no outline representing an overlap between different tissues.

Fig S12. UpSet plot showing the top 20 intersections among the NDAM contrasts. The
horizontal bars indicate the total number of significant differentially expressed genes (DEGs)
for each contrast while the vertical bars indicate the number of significant DEGs in common
between the contrasts annotated with black dots connected by lines in the intersection
matrix. The background colour of the stripes in the intersection matrix and horizontal bars
represents the tissue. The colour of the bars represents the days post infection (dpi) with
black bars indicating an overlap between different timepoints. The outline colour of the bars
also represents the tissue with no outline representing an overlap between different tissues.

Fig S13. UpSet plot showing the top 20 intersections among the BORA contrasts. The
horizontal bars indicate the total number of significant differentially expressed genes (DEGs)
for each contrast while the vertical bars indicate the number of significant DEGs in common
between the contrasts annotated with black dots connected by lines in the intersection
matrix. The background colour of the stripes in the intersection matrix and horizontal bars
represents the tissue. The colour of the bars represents the days post infection (dpi) with
black bars indicating an overlap between different timepoints. The outline colour of the bars
also represents the tissue with no outline representing an overlap between different tissues.

Fig S14. Volcano plot showing the results of the RESP contrast for the peripheral blood
mononuclear cell (PBMC) samples at 14 days post infection (dpi). Each data point represents
a gene with the position on the *x*- and *y*-axes indicating the $\log_2$ fold change and $-\log_{10}P_{adj.}$,
respectively. Genes above the horizontal dashed line are significantly differentially expressed
with the colours representing the change in expression. The top 10 most significant genes
for increased and decreased expression with gene symbols are labelled.

Fig S15. Volcano plot showing the results of the RESP contrast for the peripheral blood
mononuclear cell (PBMC) samples at 25 days post infection (dpi). Each data point represents
a gene with the position on the *x*- and *y*-axes indicating the $\log_2$ fold change and $-\log_{10}P_{adj.}$,
respectively. Genes above the horizontal dashed line are significantly differentially expressed
with the colours representing the change in expression. The top 10 most significant genes
for increased and decreased expression with gene symbols are labelled.

Fig S16. Volcano plot showing the results of the RESP contrast for the liver samples at 12
days post infection (dpi). Each data point represents a gene with the position on the *x*- and
y-axes indicating the $\log_2$ fold change and $-\log_{10}P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S17. Volcano plot showing the results of the RESP contrast for the liver samples at 15
days post infection (dpi). Each data point represents a gene with the position on the *x*- and
y-axes indicating the $\log_2$ fold change and $-\log_{10} P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S18. Volcano plot showing the results of the RESP contrast for the liver samples at 18
days post infection (dpi). Each data point represents a gene with the position on the *x*- and
y-axes indicating the $\log_2$ fold change and $-\log_{10} P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S19. Volcano plot showing the results of the RESP contrast for the liver samples at 21
days post infection (dpi). Each data point represents a gene with the position on the *x*- and
y-axes indicating the $\log_2$ fold change and $-\log_{10} P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S20. Volcano plot showing the results of the RESP contrast for the liver samples at 26
days post infection (dpi). Each data point represents a gene with the position on the *x*- and
y-axes indicating the $\log_2$ fold change and $-\log_{10} P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S21. Volcano plot showing the results of the RESP contrast for the liver samples at 29
days post infection (dpi). Each data point represents a gene with the position on the *x*- and
y-axes indicating the $\log_2$ fold change and $-\log_{10} P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S22. Volcano plot showing the results of the RESP contrast for the liver samples at 32
days post infection (dpi). Each data point represents a gene with the position on the *x*- and
y-axes indicating the $\log_2$ fold change and $-\log_{10} P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S23. Volcano plot showing the results of the RESP contrast for the liver samples at 35
days post infection (dpi). Each data point represents a gene with the position on the x- and
y-axes indicating the $\log_2$ fold change and $-\log_{10}P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S24. Volcano plot showing the results of the RESP contrast for the lymph node samples at
21 days post infection (dpi). Each data point represents a gene with the position on the x-
and y-axes indicating the $\log_2$ fold change and $-\log_{10}P_{adj.}$, respectively. Genes above the
horizontal dashed line are significantly differentially expressed with the colours representing
the change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S25. Volcano plot showing the results of the RESP contrast for the lymph node samples at
35 days post infection (dpi). Each data point represents a gene with the position on the x-
and y-axes indicating the $\log_2$ fold change and $-\log_{10}P_{adj.}$, respectively. Genes above the
horizontal dashed line are significantly differentially expressed with the colours representing
the change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S26. Volcano plot showing the results of the RESP contrast for the spleen samples at 21
days post infection (dpi). Each data point represents a gene with the position on the x- and
y-axes indicating the $\log_2$ fold change and $-\log_{10}P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.

Fig S27. Volcano plot showing the results of the RESP contrast for the spleen samples at 35
days post infection (dpi). Each data point represents a gene with the position on the x- and
y-axes indicating the $\log_2$ fold change and $-\log_{10}P_{adj.}$, respectively. Genes above the horizontal
dashed line are significantly differentially expressed with the colours representing the
change in expression. The top 10 most significant genes for increased and decreased
expression with gene symbols are labelled.**Fig S28.** g:Profiler functional enrichment of
significantly differentially expressed genes (DEGs) in the peripheral blood mononuclear cell
(PBMC) sample RESP contrasts. Each circle represents a significantly enriched GO term with
the size indicating the ratio of the intersection between the term and the DEGs. The vertical
axis shows the $-\log_{10}P_{adj.}$ value and the vertical panels and colours indicate the direction of
change in expression. The horizontal panels indicate the source of the term and the position
within the panels groups terms from the same GO subtree. The top driver GO terms (up to a
maximum of 10) are indicated with a black outline and label.

Fig S29. g:Profiler functional enrichment of significantly differentially expressed genes
(DEGs) in the liver sample RESP contrasts. Each circle represents a significantly enriched GO
term with the size indicating the ratio of the intersection between the term and the DEGs.
The vertical axis shows the -$\log_{10}P_{adj.}$ value and the vertical panels and colours indicate the
direction of change in expression. The horizontal panels indicate the source of the term and
the position within the panels groups terms from the same GO subtree. The top driver GO
terms (up to a maximum of 10) are indicated with a black outline and label.

Fig S30. g:Profiler functional enrichment of significantly differentially expressed genes
(DEGs) in the lymph node sample RESP contrasts. Each circle represents a significantly
enriched GO term with the size indicating the ratio of the intersection between the term and
the DEGs. The vertical axis shows the -$\log_{10}P_{adj.}$ value and the vertical panels and colours
indicate the direction of change in expression. The horizontal panels indicate the source of
the term and the position within the panels groups terms from the same GO subtree. The
top driver GO terms (up to a maximum of 10) are indicated with a black outline and label.

Fig S31. g:Profiler functional enrichment of significantly differentially expressed genes
(DEGs) in the spleen sample RESP contrasts. Each circle represents a significantly enriched
GO term with the size indicating the ratio of the intersection between the term and the
DEGs. The vertical axis shows the -$\log_{10}P_{adj.}$ value and the vertical panels and colours indicate
the direction of change in expression. The horizontal panels indicate the source of the term
and the position within the panels groups terms from the same GO subtree. The top driver
GO terms (up to a maximum of 10) are indicated with a black outline and label.

Fig S32. EnrichmentMap network of significantly enriched GO terms identified from
g:Profiler functional enrichment of significant differentially expressed genes (DEGs) for the
DIRE contrasts.Each node represents a GO term with the colour of the node representing the
tissue and the size representing the number of genes in the GO term. The edges indicate
overlap between the GO terms with the width of the edges representing the similarity
coefficient for the connected GO terms. The GO terms are clustered by AutoAnotate with
the background colour of the clusters representing the direction of expression. The clusters
are labelled with the size of the label scaling with the number of GO terms in the cluster.

Fig S33. EnrichmentMap network of significantly enriched GO terms identified from
g:Profiler functional enrichment of significant differentially expressed genes (DEGs) for the
NDAM contrasts. Each node represents a GO term with the colour of the node representing
the tissue and the size representing the number of genes in the GO term. The edges indicate
overlap between the GO terms with the width of the edges representing the similarity
coefficient for the connected GO terms. The GO terms are clustered by AutoAnotate with
the background colour of the clusters representing the direction of expression. The clusters
are labelled with the size of the label scaling with the number of GO terms in the cluster.

Fig S34. EnrichmentMap network of significantly enriched GO terms identified from
g:Profiler functional enrichment of significant differentially expressed genes (DEGs) for the
BORA contrasts. Each node represents a GO term with the colour of the node representing
the tissue and the size representing the number of genes in the GO term. The edges indicate
overlap between the GO terms with the width of the edges representing the similarity
coefficient for the connected GO terms. The GO terms are clustered by AutoAnotate with
the background colour of the clusters representing the direction of expression. The clusters
are labelled with the size of the label scaling with the number of GO terms in the cluster.