Two cowpea Rubisco activase isoforms for crop thermotolerance

Armida Gjindali; Rhiannon Page; Catherine J. Ashton; Ingrid Robertson; Mike T. Page;

Duncan Bloemers; Peter D. Gould; Dawn Worrall; Douglas J. Orr; Elizabete Carmo-Silva

Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

Corresponding author email: e.carmosilva@lancaster.ac.uk

Submission date: 31st January 2025

Number of figures: 4 (all figures in colour)

Number of tables: 2

Number of Supporting Information figures: 15 (all figures in colour)

Number of Supporting Information tables: 10

Word count: 9520 (Introduction 1574, M&M 4096, Results 2126, Discussion 1724)

Summary

• A vital crop for sub-Saharan Africa, cowpea productivity is threatened by climate
change, including increased heatwave intensity, duration, and frequency. Rubisco
activase (Rca) is a key molecular chaperone that maintains Rubisco activity and due
to its thermal sensitivity is a key target for improving crop climate resilience.

• We identified and characterised four Rca isoforms in *Vigna unguiculata* (L.) Walp. cv.
IT97K-499-35 *in vitro*: Rca1β, Rca8α, Rca10α and Rca10β. Cowpea leaf and plant
growth traits were also investigated during a 5-day +10 °C heatwave that included
assessment of Rubisco and Rca activity in leaf extracts, and relative changes in the
abundance of the four Rca isoforms.

• Cowpea Rca10α and Rca10β had higher thermal maxima, broader thermal optima
and higher rates of ATP hydrolysis and Rubisco reactivation *in vitro*. In the absence
of water deficit, the heatwave caused only mild effects, including increases in leaf
temperature and expression of *Rca10*, small decreases in Rubisco activity and
activation state, and unaltered temperature response of Rubisco activation by Rca.

• The superior Rca10α and Rca10β isoforms offer the prospect of enhancing the
thermotolerance of cowpea and other crops in anticipation of more extreme future
heatwaves.

Keywords: Activase, cowpea, heat stress, photosynthesis, Rca, Rubisco, temperature
response, *Vigna unguiculata*.

36   Introduction

Changes in global temperatures including heatwave events pose significant threats to global

food security (Moore *et al.*, 2021), making increased crop resilience against heat stress

imperative (Song *et al.*, 2023). Regions closer to the equator, particularly sub-Saharan

Africa, are the most vulnerable to rising temperatures (Harboure, 2024). Elevated

temperatures can lead to decreased photosynthesis and plant growth (Hatfield & Prueger,

2011; Bagati *et al.*, 2021; Zahra *et al.*, 2023; Becker *et al.*, 2023). Cowpea (*Vigna*

unguiculata (L.) Walp) is a major source of protein for populations in sub-Saharan Africa

and, despite being adapted to a warm climate, it is still vulnerable to heat stress – even a 1

°C increase in night temperature can result in significant yield loss (Warrag & Hall, 1984;

Gonçalves *et al.*, 2016; Mohammed *et al.*, 2024). Therefore, it is imperative to identify

physiological and biochemical traits that confer thermotolerance in cowpea to develop heat

resilient crops (Barros *et al.*, 2021).

Moderate heat stress, defined as temperatures up to 40 °C (Crafts-Brandner & Salvucci,

2000; Kumar *et al.*, 2009; Ara *et al.*, 2013), has a profound impact on plant photosynthesis,

adversely affecting both the electron transport chain and the Calvin-Benson-Bassham (CBB)

cycle (Chen *et al.*, 2022). Rubisco and chlorophyll leaf content also decline under moderate

heat stress (Mathur *et al.*, 2014). Chlorophyll biosynthesis enzymes exhibit susceptibility to

heat stress; however, the reduction in Rubisco content cannot be attributed to denaturation

of Rubisco, as alterations in secondary structure are observed only at temperatures

exceeding 45 °C (Li *et al.*, 2002; Mathur *et al.*, 2014). Increased temperatures enhance

Rubisco carboxylation activity; but also lead to reduced Rubisco specificity and increased O_2

diffusion, both of which result in higher oxygenation over carboxylation and increased

photorespiration (Hermida-Carrera *et al.*, 2016; Galmés *et al.*, 2016). Importantly,

photorespiration does not fully account for the decline in carbon assimilation during

moderate heat stress, as the Rubisco activation state has been demonstrated to decrease

under both photorespiratory and non-photorespiratory conditions during mild heat stress

(Feller *et al.*, 1998; Crafts-Brandner & Law, 2000; Galmés *et al.*, 2016).

Numerous studies have demonstrated that Rubisco inactivation is one of the primary causes

of reduced photosynthetic efficiency during moderate heat stress (Feller *et al.*, 1998; Crafts-

Brandner & Law, 2000; Portis, 2003; Ristic *et al.*, 2009; Carmo-Silva *et al.*, 2015; Perdomo

et al., 2017; Wijewardene *et al.*, 2021; Scafaro *et al.*, 2023). To catalyse carboxylation or

oxygenation, Rubisco catalytic sites require binding of CO_2 and Mg^{2+} as cofactors to form the

active carbamylated Rubisco prior to binding the substrate ribulose 1,5-biphosphate (RuBP).

Rubisco can become inactivated by the binding of inhibitory sugar phosphates to

carbamylated catalytic sites or by misfire products that cannot dissociate and maintain

catalytic sites in their closed, inactive conformation (Orr *et al.*, 2023). Inactivation can also

occur when RuBP binds to an uncarbamylated Rubisco catalytic site. Inhibited Rubisco

requires its molecular chaperone Rubisco activase (Rca) to restore its activity (Carmo-Silva

& Salvucci, 2013). Rca is a nuclear encoded protein that belongs to the AAA+ ATPases

superfamily (Hazra *et al.*, 2015) and, in its active multimeric form, exhibits two activities: ATP

hydrolysis and Rubisco reactivation (Bhat *et al.*, 2017; Serban *et al.*, 2018). Rca docks on

Rubisco's large subunit and, using ATP, performs conformational remodelling to release the

inhibitory compound, thus modulating Rubisco activity by facilitating the dissociation of

bound inhibitors. Elevated temperatures increase the frequency of Rubisco inactivation due

to higher concentrations of some inhibitory substrates (Salvucci & Crafts-Brandner, 2004, p.

20). Furthermore, although Rubisco is a relatively thermostable enzyme, Rca is thermolabile

and even a moderate increase in temperature results in loss of its activity (Crafts-Brandner &

Salvucci, 2000; Salvucci *et al.*, 2001).

Rca isoforms within a single species vary in their regulatory properties (Zhang & Portis,

1999; Carmo-Silva & Salvucci, 2013; Perdomo *et al.*, 2019; Scafaro *et al.*, 2019b). Many

species express two types of Rca isoforms: longer α isoforms (43-47 kDa) and shorter β

isoforms (41-43 kDa) that can be encoded either by individual genes or the same gene via

alternative splicing (Carmo-Silva *et al.*, 2015). Rca α isoforms contain a C-terminal extension

with two cysteine residues shown to form a disulfide bond which is regulated by thioredoxin-f

in *Arabidopsis* (Zhang & Portis, 1999). This regulatory mechanism links Rubisco activity with

alterations in the redox potential induced by environmental stimuli, such as high light (Zhang

et al., 2002; Perdomo *et al.*, 2021). The redox regulation is species-specific, with

Solanaceae species such as tobacco (*Nicotiana tabacum* L.) and tomato (*Solanum*

lycopersicum L.) containing only the β Rca isoform that lacks the cysteine residues and the

redox regulation of the enzyme (Carmo-Silva & Salvucci, 2013). In species containing only

redox insensitive Rca, light regulation appears to be mediated by changes in the ADP/ATP

ratio of the chloroplast (Carmo-Silva & Salvucci, 2013).

In addition to variation in the number of genes and the isoforms they produce, the

expression and relative abundance of Rca isoforms vary among species and as a response

to temperature (Law & Crafts-Brandner, 2001; Wang *et al.*, 2010; Degen *et al.*, 2020;

Perdomo *et al.*, 2021). In rice (*Oryza sativa* L.) two Rca isoforms are produced via

alternative splicing and while Rca β is more abundant under optimal steady state conditions

(Wang *et al.*, 2010), heat stress induces upregulation of Rca α (Scafaro *et al.*, 2016). In

maize (*Zea mays* L.) the two isoforms derive from two genes *ZmRcaα* and *ZmRcaβ*

encoding an α (43 kDa) and a β (41 kDa) isoform respectively (Yin *et al.*, 2014). Under

optimal temperature conditions, the ZmRca β protein isoform is expressed in higher levels

than ZmRca α and both exhibit comparable diurnal variation in expression levels (Ristic *et*
al., 2009). Jiménez *et al.* reported an increase in ZmRca α in seedlings exposed to higher
temperatures, however in more mature heat-treated plants this effect was not consistent
(Ristic *et al.*, 2009; Stainbrook *et al.*, 2024). In wheat (*Triticum aestivum* L.) there are two
Rca genes, one of which produces a β isoform while the other (*TaRca2*) is alternatively
spliced and produces both an α and a β isoform (Nagarajan & Gill, 2018). The Rca α/β ratio
changes when wheat is exposed to heat stress, suggesting that the relative abundance of
the different Rca isoforms is linked with the physiological response to heat stress (Warrag &
Hall, 1984; Crafts-Brandner & Salvucci, 2000). By contrast, in *Arabidopsis* the Rca isoform
ratio is not impacted by heat, with the α and β isoforms originating from alternative splicing
continuing to be expressed at a 1:1 ratio (Zhang & Portis, 1999). These species-specific
responses of Rca and Rubisco regulation to heat stress illustrate the need to investigate this
system directly in the crop of interest.

While the exact mechanism of Rca deactivation at high temperatures is still not fully
understood, it is evident that the optimum temperature range for Rca is species-specific and
in tune with the photosynthetic optimum (Scafaro *et al.*, 2016). Therefore, the temperature at
which Rca loses its activity depends on environmental acclimation; i.e., plants that grow at
high temperatures, such as cotton and agave, maintain Rca activity at higher temperatures
(Salvucci & Crafts-Brandner, 2004; Carmo-Silva & Salvucci, 2011; Scafaro *et al.*, 2016;
Shivhare & Mueller-Cajar, 2017). Rca thermotolerance also differs between closely related
species and among the native Rca isoforms within a plant species. The wild rice species
Oryza australiensis Domin and *Oryza meridionalis* Ng. contain thermostable Rca isoforms
that enable sustained high Rubisco activity and photosynthetic rate even at 45 °C (Scafaro
et al., 2016, 2018). In rice, spinach and cotton, the α isoform exhibits higher thermostability
whereas in wheat, it is one of the two shorter β (TaRca1β) isoforms that is most
thermostable (Kurek *et al.*, 2007; Wang *et al.*, 2010; Keown & Pearce, 2014; Degen *et al.*,
2020). Increasing the thermostability of Rca is considered one of the most promising
strategies for increasing Rubisco activity and enhancing photosynthesis under elevated
temperature (Carmo-Silva *et al.*, 2015; Wijewardene *et al.*, 2021).

In this study we identified four Rca isoforms in cowpea encoded by three genes: *Rca1*, *Rca8*
and *Rca10*, with the latter producing two isoforms via alternative splicing. Characterisation of
their ATPase and Rubisco reactivation activities across a range of temperatures *in vitro*
revealed that the two Rca10 protein isoforms have higher thermal maxima and broader
thermal optima for both activities. We then examined the effect of moderate heat stress
during a 5-day heatwave at 38 °C (+10°C compared to control) on the expression and
abundance of Rca isoforms and subsequent impacts on Rubisco activation in the leaves of

young cowpea plants. Heat treatment resulted in decreased Rubisco activation state and

upregulation of *Rca8a* and *Rca10*, however *Rca10* levels were consistently the lowest.

Despite upregulation of the more thermotolerant isoforms, the temperature profile of Rubisco

reactivation by Rca in leaf extracts remained unaltered in heat-treated plants. Leaf

temperature increased by 6 °C during the heatwave, reaching 32 °C, which is within the

temperature range where Rca activity is maintained above 70% of the maximum (T_{opt}) for all

isoforms. We therefore conclude that warm-adapted cowpea plants can cope with the mild

heat stress induced by a 5-day heatwave at 38 °C in the absence of water deficit, which did

not impair Rca function. Overexpression of the most thermotolerant, yet least abundant, Rca

isoforms in cowpea will enable adaptation to leaf temperatures resulting from heatwaves that

are more intense or accompanied by water deficit, which are likely to occur in sub-Saharan

Africa.

Materials and Methods

Gene identification

Cowpea *Rca* genes were identified by searching the cowpea genome (Phytozome 12, cowpea genome v1.1) using keywords and known Rca protein sequences from *Arabidopsis thaliana*, *Nicotiana tabacum*, and *Triticum aestivum* (Supporting Information Table S1)(Goodstein *et al.*, 2012; Lonardi *et al.*, 2019). This identified 3 genes for *Rca* in cowpea: *Vigun01g219300* (*Rca1*), *Vigun08g150700* (*Rca8*), and *Vigun10g051600* (*Rca10*). Alignment with known sequences and key conserved features of canonical Rcas allowed identification of the products of each gene. Transit peptide sequences for each protein were identified by alignment to known Rca sequences (with known transit peptide locations). The analysis was repeated using the updated Phytozome 13 cowpea genome v1.2 and yielded identical Rca isoforms (Supporting Information Fig.S1). Analysis of transcript sequences showed that *Rca1* encodes a 384 amino acid (aa) mature protein consistent with a β isoform. *Rca8* encodes a 419 aa α isoform and a 295 aa isoform that is shorter than a regular β isoform and unlikely to be functional. *Rca10* encodes both a 419 aa α and a 382 aa β isoform by alternative splicing.

The cowpea *HSP20* (*Vigun06g052200*) gene, used as a heat stress marker, was identified as a homolog of the soybean *HSP20* (*Glyma.18g094600*) shown to be upregulated in two soybean cultivars exposed to heat stress (Song *et al.*, 2017). The Arabidopsis homolog (*At4G27670*) is also upregulated in heat stress (eFP Browser Stress Series, https://bar.utoronto.ca/)(Kilian *et al.*, 2007; Winter *et al.*, 2007). The *Glyma.18g094600* coding sequence was downloaded from SoyBase (https://legacy.soybase.org) and used to search the cowpea genome as above (Brown *et al.*, 2021).

Cloning of cowpea Rca and construction of *E. coli* expression vectors

Coding regions for the mature proteins of cowpea Rca1β, Rca8α, Rca10α and Rca10β were codon optimised for *E. coli, de novo* synthesised and subcloned into pUC57 (Kan[R]) by Aruru Molecular Ltd. (Dundee, UK). A Golden Gate compatible *E.coli* expression vector was created from the pET His TEV LIC plasmid (a gift from Scott Gradia, Addgene plasmid #29653; http://n2t.net/addgene:29653; RRID:Addgene_29653). Ligation independent cloning was used to add a Golden Gate cassette taken from pICH47742 (part of the MoClo Toolkit that was a gift from Sylvestre Marillonnet, Addgene kit # 1000000044) into pET His TEV LIC, generating the new Golden Gate compatible pET His TEV GG plasmid (Weber *et al.*, 2011; Werner *et al.*, 2012). Cloning a coding region into this vector allows production of a target protein fused to an N-terminal histidine (His) tag that can be removed with TEV protease. PCR was used to add Golden Gate overhangs to the four optimised *Rca* sequences (see

Supporting Information Table S2 for primers used). Golden Gate reactions (final volume 15

µL) were performed with a 2:1 molar ratio of insert:plasmid and the following reagents (final

concentration): 1× T4 DNA Ligase buffer, 0.1 mg mL^{-1} BSA, 1 mM ATP, 40 KU/mL T4 DNA

Ligase, 2 KU/mL *Bsa* I. All reagents from NEB, except ATP (Promega). Reaction conditions

were 30 cycles of [37 °C for 2 min followed by 16 °C for 3 min], 37 °C for 5 min, 80 °C for 10

min, and 16 °C for 1 min. 5 µL of the reaction mix was used to transform *E. coli* DH5α cells.

Resulting colonies were used to generate plasmid DNA for sequencing with primers T7F and

T7R (Source Bioscience) to verify correct sequence including in-frame fusion of the coding

region to the His tag.

Recombinant Rca expression and purification

Expression constructs were used to transform BL21(DE3)pLysS *E. coli* competent cells

(Invitrogen). PCR and sequencing were used to confirm the presence of the correct

constructs in the BL21(DE3)plysS *E. coli* cells. Several cell lines were tested for each

construct to identify the cell strains with the highest expression levels for the target proteins.

Expression experiments were then scaled up for protein purification.

Starter cultures (10 mL M9ZB medium, containing kanamycin and chloramphenicol, both at

50 µg/mL) were inoculated from glycerol stocks and grown overnight at 37 °C with 225 rpm

orbital shaking. Dynamite medium (containing kanamycin and chloramphenicol, at 50 µg/mL)

was seeded with the overnight starter culture (10 mL starter culture in 1 L Dynamite

medium). This culture was grown for 3-4 h at 37 °C with shaking at 220 rpm until OD_{595}=0.6-

0.8 and then IPTG (0.5 mM final concentration) added to initiate protein expression. Cultures

were then transferred to 24 °C with 220 rpm shaking for 20 h. Cells were pelleted by

centrifugation (5,000 × *g*, 4 °C, 20 min), washed with 40 mL buffer (50 mM HEPES pH 7.2,

300 mM NaCl, 5 mM $MgCl_2$), transferred to 50 mL screw capped tubes before finally

collecting by centrifugation (5,000 × *g*, 4 °C, 20 min). The pellet was flash frozen in liquid

nitrogen and stored at -80 °C.

Cell pellets were thawed on ice and resuspended in Lysis Buffer (50 mM HEPES pH 7.2, 5

mM $MgCl_2$, 300 mM NaCl, 50 mM Imidazole, 0.1% Triton X-100 (v/v), 10 µM leupeptin, 1

mM PMSF), using 7 mL buffer for every 1 g of pellet. Cells were lysed on ice using a probe

sonicator and 5 cycles of 30 s on 30 s off until the solution was homogeneous. Cell debris

was removed by centrifugation (23,000 × *g*, 4 °C, 30 min) and the soluble fraction pushed

through a 0.45 µm syringe filter prior to chromatography.

Purification of the Rca protein was achieved using an NGC chromatography system (Bio-

Rad, UK), equipped with a nickel-charged IMAC column (1 mL HisTrap Fast Flow, Cytiva)

and a desalting column (10 mL Bio-Scale Mini Bio-Gel P6 Desalting Cartridge, Bio-Rad).

IMAC and desalting columns were equilibrated with Equilibration Buffer (5 mL: 50 mM

HEPES pH 8, 300 mM NaCl, 5 mM $MgCl_2$, 50 mM imidazole) and Final Buffer (13 mL: 50

mM HEPES pH 8, 5 mM NaCl, 5% glycerol), respectively. The sample was applied to the

IMAC column, followed by Wash Buffer (15 mL: 50 mM HEPES pH 8, 300 mM NaCl, 5 mM

$MgCl_2$, 125 mM imidazole). Bound protein was eluted using a higher concentration of

imidazole (2.1 mL: 50 mM HEPES pH 8, 300 mM NaCl, 5 mM $MgCl_2$, 300 mM imidazole)

and directly applied to the desalting column. Buffer was exchanged with 15 mL Final Buffer

and 1 mL fractions collected. Fractions containing protein were identified using a UV/Vis

detector reading at 280 nm. Fractions containing the highest protein concentration

(determined using Bradford reagent, BIO-RAD) were pooled, ATP and DTT (2 mM and 5

mM final concentration respectively) added and 50-100 µL aliquots frozen in liquid nitrogen

before storage at -80 °C.

In vitro characterization of Rca thermal optima

The rate of ATP hydrolysis by Rca isoforms was assayed using the method described by

Chifflet *et al.* (1988), with modifications as described within Degen *et al.* (2020). ATP

hydrolysis was measured at a range of temperatures (20-55 °C) using 5-minute assays, with

reactions initiated through addition of Rca (final concentration 1 µM Rca monomer) and

quenched via addition of SDS. After quenching, all reactions and standards were kept at 4

°C until inorganic phosphate (P_i) determination. The P_i released during ATP hydrolysis was

determined by measuring the absorbance of molybdenum blue at 850 nm in a

spectrophotometer (SpectroStar Nano, BMG Labtech, UK).

Rubisco reactivation assays were conducted at a range of temperatures (20-45 °C) following

the procedures of Barta *et al.* (2011) and Degen *et al.* (2020). Briefly, Rubisco purified as per

Amaral *et al.* (2024) was used to prepare inhibited Rubisco complexes (ER) through

incubation with RuBP overnight at 4 °C. The two-stage reactivation assay comprises a first

stage assay where Rca is added to inhibited Rubisco (ER) at a test temperature (20-45 °C),

with aliquots from this assay taken at time intervals to initiate the second stage assay that

measures Rubisco activity. This Rubisco activity is a function of the Rca, time allowed for

reactivation, and the temperature of the first stage (Supporting Information Fig. S2). With

known quantities of Rca and Rubisco, and appropriate controls as detailed within Degen *et*

al. (2020), Rca reactivation of Rubisco can be established. Maximum Rubisco activity was

determined at every temperature and was comparable among assays with each of the four

Rca isoforms.

Independent purifications of each isoform were used as biological replicates. For ATPase

activity, three biological replicates were measured for Rca8α and four for the each of the rest

of the isoforms. For Rubisco reactivation, four biological replicates were measured for
temperatures 20-32 °C and three for temperatures 38-45 °C. To estimate maximum ATP
hydrolysis and Rubisco activation activities and the corresponding temperatures at which the
maximum values are attained, second- to third-order polynomials and generalized additive
models (GAM) were fitted to the experimental data using the gam function from the mgcv
1.8–24 package in R (Wood, 2017). The model that best fit the experimental data was
selected based on the Akaike information criterion using the AIC function (Akaike, 1974)
(Supporting Information Table S4). The selected model was fitted to each set of the
biological replicates (Table 1) as well as the whole dataset (Supporting Information Table
S3). T_{max} was calculated as the temperature corresponding to maximum activity. T_{opt} was
calculated as the range where activity was maintained above 70% of maximum value. $T_{0.5}$
was calculated as the temperature where activity dropped below 50% of maximum value.

Plant growth and sampling

Vigna unguiculata (L.) Walp. cv. IT97K-499-35 seeds (kindly provided by B.B. Singh) were
germinated in 0.6 L Deepots (D40H, Stuewe & Sons) containing 1:1 (v:v) mixture of nutrient-
rich compost (Petersfield Growing Mediums) and silver sand (horticultural grade, Royal
Horticultural Society). Plants were grown in eight growth cabinets (MC1000, Snijders Labs)
and watered as needed to soil saturation. Cabinets were maintained at day/night
temperatures of 28/18 °C, 70% relative humidity and 400 ppm CO_2 concentration (control
conditions). A 12 h photoperiod of constant light of 850 µmol m^{-2} s^{-1} at canopy level was
maintained via LED lighting (NS1 spectrum, Valoya). Four cabinets were randomly selected
to apply a heat treatment starting at 13 days after sowing (das) including an intermediate day
with day/night temperatures of 34/24 °C followed by 5 days of 38/28 °C (Supporting
Information Fig. S3a). Data loggers (OM-EL_USB-2, Omega Engineering Ltd.) were placed
at canopy level in each cabinet to record temperature and relative humidity every 15 minutes
(Supporting Information Fig. S3b).

Twenty cowpea plants were grown in each cabinet and sampled or measured for different
analyses following a randomised design (Supporting Information Fig. S4). All samples and
measurements were taken from the central leaflet of the first trifoliate leaf. Leaf disc samples
were taken 4-5 h into the light period, after which plants were cut above the unifoliate leaves
to reduce shading of neighbouring plants. Separate plants were used for sample collection in
subsequent time points. For non-destructive measurements, such as plant growth traits and
chlorophyll content, the same individuals were measured at all time points.

Growth measurements were obtained at 15, 17 and 22 das (days 2 and 4 of the heatwave
and on day 4 of the post-heatwave recovery period, Supporting Information Fig. S6). Leaf

chlorophyll content was estimated using a chlorophyll concentration meter (MC-100,

Apogee) by measuring 3 positions on the central leaflet for each plant. Plant height was

measured using a ruler from the soil level to the apical meristem. Leaf length and width

(widest part perpendicular to the length) were also measured with a ruler for the central

leaflet only. Leaf thickness was measured with a digitronic calliper 110-DBL (Moore &

Wright). Total above-ground biomass was determined on a separate set of plants by drying

leaves and stems at 60 °C until constant weight.

Chlorophyll fluorescence (method below) measurements were conducted every day of the

heatwave. Three 0.38 cm^2 leaf discs were excised and quickly placed with the adaxial

surface upwards on a single layer of wet filter paper on a thin plastic tray (<0.5 mm) and

immediately transported to the fluorescence imager.

A thermal imaging camera (CAT S60 smartphone, Caterpillar) was used prior to sampling to

assess T_{leaf}. Each image was later analysed with FLIR Thermal Studio software (Teledyne

FLIR LLC), providing the mean temperature across an area of the leaf surface (Supporting

Information Fig. S5).

For RNA-seq analyses (method below), four 0.55 cm^2 leaf discs were collected from 4

independent biological replicates per treatment per timepoint (days 1, 3 and 5 of heatwave)

using a cork borer, then immediately snap frozen in liquid nitrogen and stored at -80 °C.

For protein and Rubisco analyses, four 0.55 cm^2 leaf discs were excised as above and

quickly placed with the adaxial surface upwards on a single layer of wet filter paper to

maintain turgidity. The filter paper was contained in a thin plastic tray and positioned on the

surface of the water in a circulating water bath in the artificial sunlight simulation rig (Taylor

et al., 2022). PPFD at leaf-disc-level was 850 µmol m^{-2} s^{-1}. Water bath temperature was

maintained at 26 °C or 32 °C (±0.1 °C) for control or heat-treated plants, respectively, to

mimic leaf temperatures in the growth cabinet. After 1 h incubation leaf discs were quickly

blotted on absorbent paper to remove excess moisture, then immediately frozen in liquid

nitrogen and stored at -80 °C.

For Rubisco reactivation in leaf extracts (LE), eight 0.55 cm^2 leaf discs were sampled on day

5 of the heatwave. A different plant was used for each of the seven temperatures of the

temperature response curve (20, 25, 32, 38, 40, 45 °C). Samples were quickly frozen in

liquid nitrogen, then stored at -80 °C until further analysis.

Chlorophyll fluorescence

Samples were analysed using a chlorophyll fluorescence imager (closed 800C FluorCam,

Photon System Instruments). The imager contained a customized temperature-controlled

sample surface area which comprised an aluminium plate with imbedded copper piping

connected to a circulating temperature-controlled water bath (Optima TX150, Grant

Instruments). The filter paper was in contact with sponges which acted as water reservoirs to

ensure the paper remained wet during analysis (Supporting Information Fig. S7a). The

temperature of the sample surface area and leaf discs were recorded using type k

thermocouples and a data logger (RDXL4SD, Omega). Additional thermocouple wires were

placed in contact with the sample surface area, and extra leaf discs, to record temperature

throughout the analysis (Supporting Information Fig. S7a). Samples were first dark adapted

for 1 h at 26 °C. After dark adaptation, the Fv/Fm protocol in the software (FluorCam 7,

Photon System Instruments) was used to make measurements of the maximum quantum

efficiency of photosystem II over a range of temperatures. The temperature of the leaf discs

was used to assess when the measurement temperature had been reached and was stable.

Once stable, the temperature was held for 3 min before the Fv/Fm measurement was

taken.

The data for each set of three leaf discs from the same plant were averaged and plateauing

Fv/Fm points at higher temperatures were removed before the remaining data were used to

calculate T_{crit} (Supporting Information Fig. S7b) with the segmented function from the

segmented package(Muggeo, 2017) in R. This function uses a previously fitted linear model

to identify a breakpoint and fits two linear models before and after the breakpoint. The

breakpoint in the Fv/Fm curve was taken as T_{crit} while the slopes of the two linear models

were taken as m1 & m2 (Supporting Information Fig. S7b, c, d).

Protein extraction, Rubisco activity and content

Proteins were extracted from leaf disc samples and Rubisco activity assays were performed

in a randomised order, as described previously (Taylor *et al.*, 2022; Ashton *et al.*, 2024). The

activation state of Rubisco was calculated as the ratio of initial to total activity. In short,

aliquots of the leaf extract supernatant were added to an assay mix containing $^{14}CO_2$ and

RuBP to initiate assays that gave an estimate of Rubisco initial activity, whereas Rubisco

total activity was measured after allowing a 3-minute incubation period of the leaf extract in

the assay mix without RuBP to allow full carbamylation of Rubisco.

Aliquots of leaf extract supernatant were also used for determining Rubisco content via ^{14}C-

CABP (2-carboxy-D-arabinitol 1,5-bisphosphate) binding, as previously described (Ruuska

et al., 1998; Ashton *et al.*, 2024) and total soluble protein content, using the Bradford method

(Bradford, 1976).

Gene expressions analysis

Frozen leaf disc samples were ground to a fine powder in liquid nitrogen using a pestle and mortar. Total RNA was extracted from 20-30 mg tissue using a NucleoSpin™ RNA Plant Kit (Macherey-Nagel) including a DNase treatment, according to the manufacturer's instructions. RNA concentration and quality were determined by spectrophotometry using a microplate reader and LVis plate (SPECTROstar Nano, BMG Labtech) (Supporting Information Table S5). Further QC analysis of RNA samples was performed by Novogene using a Qubit fluorometer (Invitrogen) and an Agilent 5400 fragment analyzer, followed by mRNA purification (using poly-T oligo-attached magnetic beads), library construction and 2 x 150 bp paired-end (PE150) sequencing using an Illumina NovaSeq 6000 (read depth of 30 million reads).

An initial filtering was applied to the raw data to remove reads containing adapters, containing N > 10% (N represents bases that cannot be determined) and low-quality bases (Phred < 5). Further read quality assessment and trimming was performed using the FastQC tool, version 0.12.1 (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and cutadapt, version 4.2 (Martin, 2011). Low quality bases (Phred < 30) were trimmed from 3' ends of each read, flanking N bases were removed and trimmed reads shorter than 50 bp were discarded (for sequencing and alignment statistics see Supporting Information Table S6).

Cowpea IT97K-499-35 reference transcriptome v1.2 (Lonardi *et al.*, 2019; Liang *et al.*, 2024) was downloaded from Phytozome 13 (Goodstein *et al.*, 2012). Salmon, version 1.10.1 (Patro *et al.*, 2017) was first used to index the reference transcriptome and then quantify expression of clean reads at the transcript level. To identify changes in the cowpea Rca transcripts TPM values were extracted from the Salmon Quant.sf output files for Rca, RbcS and HSP20 transcripts. Three transcripts are present for Rca1β (the first two result in the same protein, the third is missing some of the N domain) and TPM values for all three were combined. Three transcripts exist for Rca8α (transcripts 2 and 3 are non-functional and have low expression) so here we only present data for the first transcript. For Rca10 transcripts 1, 3 and 4 are combined to give total expression for the alpha isoform and transcript 2 alone represents the beta isoform. The three transcripts for RbcS were also combined.

Global gene expression analysis was carried out in R version 4.4.1 (2024-06-14, Race for Your Life). Raw counts from Salmon were loaded into EdgeR 4.2.1 using tximport 1.32.0 and rtracklayer 1.64.0. Transcript data for each gene was combined at this stage so that all global analysis was performed at the gene-level. PCA plot was made using logCPM the PCAtools 2.16.0 to check that biological replicates group together and to look for outlying

samples. Here we used the counts per million (CPM) to remove the variability due to library
depth. The EdgeR trimmed mean of M-values (TMM) method was used for normalisation of
read counts. The data was filtered using a CPM threshold to remove genes with zero or near
zero expression. A generalised linear model was then fitted to the normalised and filtered
data for each gene. Empirical Bayes quasi-likelihood F-tests were used to test whether
genes were significantly differentially expressed between heat-treated and control plants on
each day of the heatwave. To control for multiple testing, p-values were calculated using the
false discovery rate (FDR) of Benjamini-Hochberg. Differentially expressed genes (DEGs)
were then identified using the threshold values of a false discovery rate (FDR) of < 0.05 and
a $\log2$ (|fold change|) of >1.5.

Rca gene expression changes were validated via RT-qPCR (Supporting Information Fig.
S11). For sampling and detailed methodology see Supporting Information Tables S7 & 8.

Rca quantification via western blotting

To quantify the abundance of different Rca protein isoforms, an aliquot of supernatant
resulting from Rubisco analysis was mixed with SDS-PAGE loading buffer and separated as
previously described on Perdomo *et al.* (2018) on a 12% TGX Stain-Free gel (Bio-Rad). Gels
were imaged to assess total protein before transfer to a nitrocellulose membrane using a dry
blotting system (iBlot2, Thermo Fisher Scientific). Membranes were stained with REVERT
Total Protein Stain for normalisation (Li-Cor Biosciences) and imaged prior to application of
antibodies. To compare the abundance of α and β isoforms of Rca, an antibody with broad
specificity for both isoforms was used (Feller *et al.*, 1998,a gift from Michael Salvucci), with
detection via a secondary fluor-tagged antibody (IRDye800CW, Li-Cor Biosciences,
RRID:AB_1660973). A dilution series of a pooled sample was run on every gel alongside the
samples to verify antibody detection was within the linear range. To quantify Rca8α, a more
specific antibody was used (Abcalis) in place of the broad specificity antibody. The anti-
Rca8α specific antibody was generated using phage display with the peptide selected
targeted against KRGAFYGKAAQQINVP (amino acid residue 376-391), as described in
Bloemers & Carmo-Silva (2024). All blot images were obtained using an Odyssey FC (Li-Cor
Biosciences). Empiria studio software (version 2.3.0.154, Li-Cor Biosciences) was used for
image data analysis and normalisation with the total protein stain, with values determined as
signal intensity.

Characterization of Rca thermal optima in leaf extracts

Rubisco reactivation by Rca in leaf extracts was determined as described in Carmo-Silva &
Salvucci (2011). Eight leaf discs per plant were sampled from the youngest fully expanded
leaf on day 5 of the heatwave. Separate samples were used for assaying each of the seven

temperatures of the temperature response curve shown in Figure 4, and eight plant sample

replicates were used per temperature. Samples were stored at -80 °C until further analysis.

One replicate of samples from one cabinet at control and one cabinet at high temperature

was used in each day of Rca assays, with samples allocated randomly to each assay

temperature. Supporting Information Fig. S14 illustrates the method which uses leaf extracts

as the source of both Rubisco and Rca, incubates sub-aliquots with different assays mixes,

and then carries out a two-stage assay very similar to the assay with the purified proteins.

This enables determination of the capacity of a population of Rca, drawn from plants grown

in different conditions, to reactivate Rubisco. Model fitting and temperature parameter (T_{max},

T_{opt} and $T_{0.5}$) calculations were conducted as described for the *in vitro* Rca activity assays

above. From the same samples, aliquots of leaf homogenate were used for leaf chlorophyll

content determination as described previously in Sales *et al.* (2022) and Taylor *et al.* (2020),

according to equations by Lichtenthaler (1987). Aliquots of the desalted extract were used

for total soluble protein (TSP) content determination using the Bradford method (Bradford,

1976). Results are presented in Supporting Information Table S3.

Data analyses

Geneious 2024.0.7 was used to generate protein alignments depicted in Figure 1a. Data

was processed using R 4.3.1 and RStudio 2024.09.0. Graphs were prepared using the

ggplot 2 (Wickham, 2017), EnhancedVolcano (Blighe *et al.*, 2018), VennDiagramm (Chen,

2022), ggpubr (Kassambara, 2023) and patchwork (Lin Pedersen, 2024) R packages.

Outliers were detected with the Outliers package using Tukey's fences method, where

outliers are defined as extreme values that are 1.5 times the interquartile range (1.5 IQR)

below the first quartile or 1.5 IQR above the third quartile. Box plots show medians and the

first and third quartiles (25[th] and 75[th] percentiles), and whiskers extend from the hinge to the

largest or smallest value. Symbols represent individual data points (technical or biological

replicates defined for each plot either on methods or figure legend). Bar plots show means

and error bars represent standard error. For statistical analysis, normality of data was

assessed using the Shapiro-Wilk test and equality of variances using the Levene test. For

measurements on multiple days, two-way ANOVA tests were applied to determine if there

were significant day, treatment or interactive effects, followed by post-hoc Tukey tests for

multiple comparisons with Bonferroni correction. Compact letters display (cld) were

generated after the post-hoc pairwise comparisons using packages mutlicomp (Bretz *et al.*,

2016) and multicompView (Graves *et al.*, 2024). In cases where there was a slight violation

of normality or homogeneity of variance when data between different days was compared,

two-way ANOVA was still chosen as the most robust method. If normality and variance were

severely violated, data was log transformed, and two-way ANOVA was applied to the

transformed data provided assumptions were not severely violated in the transformed data.

For measurements taken at a single timepoint (growth parameters), t-test was used to

determine statistical significance between treatments. Statistical analysis results and full

datasets are provided in the Source Data file.

Results

Cowpea contains four Rubisco activase isoforms

The cowpea genome contains three genes that encode Rca. The gene on chromosome 1

(*Vigun01g219300)* encodes an Rca β, the gene on chromosome 8 (*Vigun08g150700)* an

Rca α, and the gene on chromosome 10 (*Vigun10g051600)* encodes both Rca α and β

isoforms via alternative splicing (Figure 1a). The gene on chromosome 8 also encodes a

truncated version of Rca, however this would not produce a canonical, fully functional Rca

as it has a premature stop codon before the α-helical domain which is necessary for the

recognition of Rubisco by Rca (Portis *et al.*, 2008). Rca isoforms in cowpea exhibit minor

sequence differences, primarily in the N-terminal region (the flexible domain critical for

Rubisco binding) (Stotz *et al.*, 2011; Shivhare *et al.*, 2019, p. 20) and the α-helical domain

(Supporting Information Fig. S1). These variations may contribute to functional diversity of

Rca in cowpea, potentially allowing for fine-tuning of responses to different environmental

conditions.

Cowpea Rca isoforms differ in thermal optima of activity *in vitro*

Previous studies have shown the potential for intraspecies variation of Rca thermal optima

(Degen *et al.*, 2020). With four distinct isoforms in cowpea and its known ability to grow in

warm (20-37 °C) temperatures (Barros *et al.*, 2021), recombinant versions of each isoform

were produced for *in vitro* characterisation of cowpea Rca thermal optima (Figure 1b). Both

enzyme activities were assessed, i.e., the ability to reactivate inhibited cowpea Rubisco and

the ATPase activity that drives reactivation (Robinson & Portis, 1989). Rubisco reactivation

by Rca was calculated after accounting for the spontaneous reactivation of Rubisco,

expressed as a percentage of maximum Rubisco activity. The spontaneous reactivation of

Rubisco, although insignificant at lower temperatures, increases with temperature

(Supporting Information Fig. S2) as the Rubisco catalytic site becomes more flexible. For all

the Rca isoforms, the temperature corresponding to the peak of activity (T_{max}), the range of

temperatures at which activity was maintained above 70% of the maximum value (T_{opt}) and

the temperature at which activity drops below 50% of the maximum ($T_{0.5}$) were consistently

higher for ATPase activity than for Rubisco reactivation activity (Table 1). A higher sensitivity

of Rubisco reactivation compared to ATP hydrolysis has been previously reported and may

be due to the interaction with Rubisco being the key step impacted by elevated temperatures

in Rca (Degen *et al.*, 2020). It has also been shown that while ATP hydrolysis can be

catalysed by Rca oligomers consisting of three subunits, Rubisco reactivation requires

oligomers containing more than three subunits (Keown *et al.*, 2013; Hazra *et al.*, 2015).

Figure 1. Temperature response of cowpea Rca isoforms in vitro. a) Amino acid residue
alignment of the four cowpea Rca isoforms encoded by genes located on chromosomes 1, 8
and 10. Black vertical lines indicate differences in sequence among the genes. The following
corresponding regions of the protein are shown: N-domain, α/β subdomain, α-helical
domain, C-domain, and C-terminal extension. b) Temperature response of ATP hydrolysis
and Rubisco activation by cowpea Rubisco activase isoforms *in vitro*. Rate of ATP hydrolysis
and Rubisco reactivation by recombinantly produced purified cowpea Rca isoforms Rca1β,
Rca8α, Rca10α and Rca10β. Assays were performed at the indicated temperatures using 1
µM purified Rca and 5 mM ATP. Activation of pre-inhibited Rubisco used 5 µM Rubisco
active sites ($R_{A.S.}$) in the ER form (1:5 Rca:$R_{A.S}$). Rubisco activation is presented relative to
maximum Rubisco activity, determined as the Rubisco activity when Rca is allowed to
reactivate inhibited Rubisco for 5 min. Values represent biological replicates (independent
purifications; n=3-4 biological replicates). Lines represent the best fit for each enzyme
(selected by the Akaike information criterion) and shaded areas denote the 95% confidence
interval for each fit (Supporting Information Table S4). Full sample data are provided in the
source data file.

Table 1. Temperature response of *in vitro* Rubisco activase (Rca) activity. Maximum rate of ATP hydrolysis and Rubisco activation and corresponding temperature (T_{max}), optimum temperature range (T_{opt}, above 70% activity) and temperature above the optimum at which 50% of the maximum activity remains ($T_{0.5}$). Values were estimated from best-fit models that describe the *in vitro* temperature response of each Rca isoform (Supporting Information Table S4). The best-fit model was applied to each of three biological replicates with complete temperature response curves shown in Figure 1b. The first biological replicate was excluded from the fitting of the individual replicates because the temperature response of Rca activity was incomplete (determined only up to 32°C); this replicate is included in the model fitting to the complete temperature response data set (Supporting Information Table S3). Values show here are the mean ± standard error of the mean of values determined for individual replicates of temperature response curves (n=3).

Rca Isoform	ATPase_{max} (mol min^{-1} mol Rca^{-1})	T_{max} (°C)	T_{opt} (°C)	$T_{0.5}$ (°C)
1β	28.0 ± 5.3	36.7 ± 0.5	29.9 – 43.4	43.4 ± 1.1
8α	35.9 ± 8.1	35.9 ± 0.7	28.4 – 43.1	45.6 ± 0.2
10α	179.0 ± 36.8	42.6 ± 1.1	30.9 – 51.6	53.6 ± 0.7
10β	183.4 ± 58.5	40.9 ± 1.1	28.5 – 52.1	53.5 ± 0.3
Rca Isoform	Rubisco reactivation$_{max}$ (mol R$_{A.S.}$ min^{-1} mol Rca^{-1})	T_{max} (°C)	T_{opt} (°C)	$T_{0.5}$ (°C)
1β	0.10 ± 0.03	29.5 ± 1.2	27.7 – 34.4	36.1 ± 1.5
8α	0.09 ± 0.02	26.0 ± 2.9	22.1 – 36.0	41.0 ± 2.9
10α	0.14 ± 0.01	34.5 ± 1.0	26.5 – 40.8	42.4 ± 0.2
10β	0.19 ± 0.01	32.5 ± 0.4	24.0 – 40.9	43.3 ± 0.8

The two Rca isoforms encoded by *Rca10* were faster at hydrolysing ATP and reactivating

Rubisco, had higher thermal maxima (T_{max}), broader thermal optima (T_{opt}), and showed

thermal sensitivity ($T_{0.5}$) at warmer temperatures compared to the other two isoforms (Figure

1b; Table 1). T_{max}, T_{opt}, and $T_{0.5}$ calculated for both ATP hydrolysis and Rubisco reactivation

were generally similar for Rca1β and Rca 8α (Table 1), but Rca1β exhibited the lowest $T_{0.5}$

for reactivation activity (36.1 ± 1.5 °C). Up to 34 °C, all Rca isoforms maintained at least

70% of maximum Rubisco reactivation activity. Rca10α exhibited the highest temperature

corresponding to the peak of activity for both ATP hydrolysis (T_{max} = 42.6 ± 1.1 °C) and

Rubisco reactivation (T_{max} = 34.5 ± 1.0 °C). T_{max} values for Rca10α and Rca10β were at

least 3 °C higher for both ATP hydrolysis and Rubisco reactivation relative to Rca1β and

Rca8α (Table 1). Combined, the results show that the cowpea Rca10α and Rca10β isoforms

have faster rates, higher thermal maxima, broader thermal optima, and are less sensitive to

warmer temperatures than Rca1β and Rca8α.

A 5-day heatwave of +10 °C causes mild heat stress in cowpea

To investigate heat-induced changes in cowpea Rca *in planta*, young plants were exposed

to a 5-day heatwave consisting of an increase of 10 °C during both day and night. Plants

were cultivated for two weeks at 28/18 °C day/night, then in half of the controlled

environment cabinets the temperature was increased to 34/24 °C day/night for one day and

subsequently to 38/28 °C day/night for five days (Figure 2a, Supporting Information Fig. S3,

4). The intermediate day aimed to resemble a gradual increase in temperature in field

conditions. Plants were well watered throughout the heatwave as the focus was on the

impact of heat on Rca and we wanted to avoid confounding effects of water deficit on CO_2

diffusion and assimilation. Cabinet and leaf temperatures were monitored throughout the

experiment to ensure the plants were under heat stress conditions, with heat-treated plants

showing a 6 °C increase in leaf temperature to 32 °C (Figure 2a, Supporting Information Fig.

S5). Thus, plants experiencing the heatwave had leaf temperatures within the T_{opt} of Rca

activity.

Plant growth traits were determined four days after the heatwave, after plants were allowed

to recover, and showed that the 5-day heatwave of +10 °C had only a mild impact on

cowpea (Figure 2b). Exposure to elevated temperature resulted in decreased leaf thickness

and increased plant height; however, no effect on aboveground dry biomass was observed.

Total chlorophyll also decreased in leaves that had been exposed to heat when determined

four days after the heatwave (Figure 2b). Measurement of leaf and plant traits during the

heatwave suggested an initial increase in the rate of cowpea leaf and plant growth (day 2,

Supporting Information Fig. S6), but only minimal differences between control and heat at

the end of and after the heatwave.

Figure 2. Cowpea response to a heatwave in controlled environment. a) Experimental design, heatwave and sampling times. Plants were grown at 28/18 °C day/night for 13 days, then exposed to an intermediate heat treatment for a day at 34/24 °C day/night before being exposed to a 38/28 °C day/night heatwave for five days. Sampling: (F) photosystem II maximum efficiency (Fv/Fm) was measured daily during the heatwave; (R) leaf discs for RNA-seq analysis, total soluble protein (TSP), Rubisco activity and amount were sampled on days 1, 3 and 5 of the heatwave; (A) samples for Rca temperature response in leaf extracts were taken on day 5 of heatwave; (B) growth parameters and biomass were measured on

day 22 of plant growth, four days after the end of the heatwave. Cabinet and leaf

temperature were monitored daily. b) Leaf and whole plant growth parameters were

measured four days after the end of the heatwave (n=15). Relative chlorophyll content was

determined using a concentration meter based on leaf reflectance. c) T_{crit} was calculated for

each day of the heatwave as the breaking point where the two Fv/Fm slopes meet

(Supporting Information Fig. S7). d) Rubisco activation state (n=7-8), (b) initial (n=7-8) and

(c) total activity per leaf area (n=6-8), Rubisco amount (n=7-8) and total soluble protein

(TSP; n=6-8) were measured in leaf extracts on days 1, 3 and 5 of the heatwave. P-values

were calculated using two-way ANOVA followed by Tukey post-hoc tests. Full sample data

are provided in the source data file.

As a proxy for photosynthetic activity, photosystem II (PSII) maximum efficiency (Fv/Fm) was

monitored throughout the heatwave (Supporting Information Fig. S7). Changes in Fv/Fm

indicate modifications in the functionality of PSII reaction centres, with a decrease indicating

photooxidative damage and reduced photochemistry quantum yield (Murchie & Lawson,

2013). Fv/Fm data collected from plants throughout the heatwave enabled determination of

T_{crit} (Figure 2c), which corresponds to the temperature at which chlorophyll *a* fluorescence

rises rapidly in response to temperature due to thermal damage in PSII, and is used as an

indicator of photosynthetic thermal stability (Perez & Feeley, 2020; Posch *et al.*, 2022). In

heat-treated cowpea, Fv/Fm did not decline compared to control plants (Supporting

Information Fig. S7e), indicating that no irreversible damage was caused to PSII reaction

centres due to heat stress. In fact, heat-treated plants showed a progressive increase in the

temperature at which Fv/Fm decreased, resulting in a significantly higher T_{crit} after 3 days,

denoting increased PSII thermal stability (p < 0.001, Figure 2c). These results indicate a

dynamic response that allows for PSII efficiency to adapt to warmer conditions.

Heat stress induces a decline in Rubisco activation state

In contrast to T_{crit}, the activation state of Rubisco decreased with heat treatment (Figure 2d).

Importantly, a decline in Rubisco activation was observed from the beginning to the end of

the experiment as leaves aged. This decline was observed earlier in heat-treated plants (day

3) compared to control plants (day 5), indicating that heat stress accelerated this decline in

Rubisco function. Rubisco activation state, defined as the ratio of initial to total Rubisco

activity, reflects the enzyme's operational efficiency under different conditions. Initial activity

represents the enzyme's immediate functionality upon extraction, while total activity reflects

its potential after full carbamylation (Ashton *et al.,* 2024). Although Rubisco activation state

was similar on day 5, the initial and total activity was significantly lower in heat-treated

compared to control plants (p < 0.001, Figure 2d). This decrease in Rubisco activity on day

5 of the heatwave was associated with decreased Rubisco amount and TSP (Figure 2d), as

no significant differences were detected in Rubisco specific activity, expressed per quantity

of Rubisco protein (Supporting Information Fig. S8).

Heat stress induces the expression of thermotolerant Rca isoforms

Potential heat-induced changes in gene expression of *Rca* isoforms were assessed through

RNA sequencing analysis (RNA-seq) on days 1, 3 and 5 of the heatwave. A principal

component analysis (PCA) distinguished gene expression changes caused by heat stress

and those resulting from leaf aging (Figure 3a, Supporting Information Fig. S9). Heat-

induced differential gene expression occurred from the first day of the heatwave, with both

up and down-regulation of genes evident in heat-treated plants compared to control

conditions throughout the duration of the heatwave (Supporting Information Fig. S9). Heat

stress was confirmed by upregulation of the heat shock protein *HSP20* (Song *et al.*, 2017)

from day 1 of the heatwave (Figure 3b). The heatwave also decreased expression of the

Rubisco small subunit (*RbcS*), which is likely associated with the lower Rubisco amounts

(Figure 3c).

Under control temperature conditions, *Rca1β* was found to be the most abundant Rca

transcript, followed by *Rca8α*, while *Rca10α* and *Rca10β* were expressed at much lower

levels (Figure 3c). Of the hundreds of up and downregulated genes (Supporting Information

Fig. S9), volcano plots showed that *HSP20* and *Rca10* were both upregulated on days 1, 3,

and 5 of the heatwave (Supporting Information Fig. S10). Heat-treated plants showed a

(non-significant) tendency for lower *Rca1β* expression, whereas *Rca8α*, *Rca10α* and

Rca10β were all upregulated during the heatwave (Figure 3c). It is noteworthy that despite

the upregulation of both *Rca10* transcripts, their expression levels remained much lower

than *Rca1β* and *Rca8α* (Figure 3c). Changes in *HSP20* and the four *Rca* transcript levels

were also confirmed via RT-qPCR (Supporting Information Fig. S11). Moreover, promoter

analysis revealed multiple stress responsive elements in *Rca1β and Rca8α*, while *Rca10*

contains two heat stress response elements and has fewer other stress responsive elements

(Supporting Information Fig. S12).

Figure 3. Differential gene expression and Rca protein abundance in cowpea during a heatwave. a) Principal components (PCA) plot of gene expression in leaves of heat-treated and control cowpea plants on days 1, 3 and 5 of the heatwave. b) Expression levels of the

heat shock protein 20 (HSP20) and the small Rubisco subunits (RbcS) were used as

reference. Bar plots represent means and error bars standard error. c) Expression levels of

the four cowpea Rca isoforms in control and heat-treated plant on days 1, 3 and 5 of the

heatwave (n=4). d) Protein abundance of Rca isoforms in leaves of control and heat-treated

plants on days 1, 3 and 5 of the heatwave (relative units). Rca α and β isoforms were

detected using an anti-Rca antibody and normalised to total protein. The size difference

between the α (45 kDa) and β (41 kDa) isoforms enabled the identification as separate

bands, and the estimation of the α/β ratio. P-values were calculated using two-way ANOVA

followed by Tukey post-hoc tests. Samples were run in separate gels and normalised to total

protein via reversible staining of the membrane. Blot images and full sample data are

provided in the source data file.

To investigate if changes in gene expression are reflected at protein level, the abundance of

leaf Rca protein was quantified via immunoblotting (Figure 3d). A pan-Rca antibody that

reacts with all isoforms of Rca and enables quantification of Rca α and Rca β isoforms, as

these run separately in gel electrophoresis due to the difference in molecular weight, was

used to identify changes in abundance between α and β isoforms (Perdomo *et al.*, 2018).

Whilst differential gene expression was observed, changes in Rca protein were less

pronounced. As *Rca10β* transcript levels are almost negligible compared to *Rca1β*, it is

reasonable to infer that the β isoform protein is predominantly attributable to *Rca1β*

expression. Expression of *Rca8α* is also likely to produce most of the α isoform protein. Rca

β remained unchanged during the heatwave, while the α isoform decreased from day 1 to

day 5 but was not significantly different between control and heat-treated plants (Figure 3d).

Protein abundance of Rca8α was also determined using a specific antibody that only reacts

with this isoform (Bloemers & Carmo-Silva, 2024). Rca8α showed a near identical pattern to

the pan-Rca antibody for Rca α with a non-significant tendency for increased abundance in

heat treated plants compared to the control (Supporting Information Fig. S13a). A significant

increase in the α/β ratio was however detected in day 3 of the heatwave, which may indicate

differences in the relative abundance of Rca isoforms (Figure 3d). Due to their very low

abundance, Rca10α & Rca10β protein isoforms were undetected in leaf extracts with Rca10

specific antibodies (Supporting Information Fig. S13b).

Heat-treated plants maintain Rca rate and thermal optimum

To test the hypothesis that heat-induced changes in the relative abundance of Rca isoforms

are reflected in activity, we characterised the temperature response of Rubisco reactivation

by Rca in protein extracts from leaves sampled on day 5 of the heatwave (Figure 4). Leaf

extracts were desalted to produce the free Rubisco apoenzyme (uncarbamylated form),

which was subsequently inhibited through incubation with RuBP (Carmo-Silva & Salvucci,

2011). The inhibited Rubisco in the leaf extract was then used to determine the rate of

reactivation by the Rca holoenzyme via removal of RuBP from uncarbamylated Rubisco

catalytic sites, at a range of temperatures (Supporting Information Fig. S14). Concurrent

measurement of spontaneous reactivation of Rubisco in the absence of ATP, which is

required for Rca activity, enabled calculation of the spontaneous (Supporting Information

Fig. S15) and Rca-mediated rate of Rubisco reactivation (Figure 4).

Figure 4. Temperature response of Rca activity in leaf extracts of control and heat-

treated cowpea plants. Leaf discs were sampled on day 5 of the heatwave. The Rubisco

and Rca contained in the leaves of each plant were extracted, Rubisco was inhibited by

binding of RuBP to decarbamylated catalytic sites in the absence of CO_2 and Mg^{2+}, then

reactivation by Rca was measured at varying temperatures by comparing an assay in

presence of ATP and ATP-regenerating system with the negative control assay in the

absence of ATP. The rate of Rubisco reactivation was calculated from measurements of

Rubisco activity at 30 and 90 s after the start of the assay and expressed relative to

maximum Rubisco activity in the same leaf extracts (as described in Supporting Information

Fig. S14). Symbols represent biological replicates (leaf extracts from independent plant

samples; n=8). Lines represent the best fit for each group of plants (selected by the Akaike

information criterion) and coloured areas denote the 95% confidence interval for each fit

(Supporting Information Table S9). Full sample data are provided in the source data file.

Rubisco reactivation by Rca tended to be faster in heat-treated compared to control plants,

however this increase was not statistically significant (*p > 0.05*, Figure 4). Heat-treated plants

exhibited similar thermal optima of Rubisco reactivation by Rca to control plants at

approximately 38 °C (Figure 4). T_{opt} and $T_{0.5}$ were also similar between control and heat-

treated plants (Table 2), indicating that the large fold-change in *Rca10* transcript abundance

(Figure 3c) and the small change in heat-induced protein Rca isoform ratio (Figure 3d) did

not alter the temperature profile of Rubisco reactivation by Rca in the leaves of heat-treated

plants (Figure 4). Interestingly, the T_{max}, T_{opt} and, to a lesser extent, $T_{0.5}$ of Rubisco

reactivation by Rca were higher in leaf extracts (Table 2) than the respective values

measured for each individual Rca isoform purified after recombinant expression (Table 1).

Table 2. Temperature response of Rubisco activase (Rca) activity in leaf extracts.

Maximum rate of Rubisco activation and corresponding temperature (T_{max}), optimum

temperature range (T_{opt}, above 70% activity) and temperature above the optimum at which

50% of the maximum activity remains ($T_{0.5}$). Values were estimated from best-fit models that

describe the temperature response of the pool of Rca isoforms extracted from leaves of control

and heat-treated cowpea on day 5 of the heatwave (Supporting Information Table S9). The

best-fit model for each treatment was applied to combined replicates shown in Figure 4 (n=6-

8).

Treatment	Rubisco reactivation$_{max}$ (mol R$_{A.S.}$ min^{-1})	T_{max} (°C)	T_{opt} (°C)	$T_{0.5}$ (°C)
Control	7.3	38.4	31.9 – 42.3	43.1
Heat	9.0	37.8	28.2 – 42.2	43.3

Discussion

Cowpea Rca isoforms differ in thermal optima

We investigated the temperature response of Rubisco activase in cowpea, a vital crop for sub-Saharan Africa that is adapted to warm temperatures, yet susceptible to the increasing pressures of global warming. Several studies demonstrated that Rca thermolability limits photosynthesis at elevated temperatures (Feller *et al.*, 1998; Crafts-Brandner & Law, 2000; Salvucci *et al.*, 2006; Perdomo *et al.*, 2017), making it a target for enhancing crop yield under elevated temperatures. In this study, we identified and characterised four Rca isoforms in cowpea: Rca1β, Rca8α, Rca10α and Rca10β. *In vitro* analysis across a range of temperatures revealed a higher thermal optimum for Rca10α and Rca10β for both ATPase and Rubisco reactivation activities (Figure 1b). The most abundant Rca1β and Rca8α isoforms in cowpea leaves (Figure 3) both exhibited lower thermal optima and thermal maxima, as well as slower rates of catalysis for both activities compared to the isoforms encoded by *Rca10* (Table 1).

Differences in thermal optima among isoforms have been reported for other species. In rice, the α isoform is more thermotolerant while in wheat one of the two β isoforms exhibits a higher thermal optima (Scafaro *et al.*, 2018; Shivhare *et al.*, 2019; Degen *et al.*, 2020). For spinach, the β isoform is more sensitive to heat stress, with Rca β homo-hexamers displaying a lower thermal midpoint compared to Rca α homo-hexamers or hetero-hexamers comprising both α and β isoforms (Keown & Pearce, 2014). Notably, for all Rca isoforms, the T_{max} for ATPase activity is higher than for Rubisco reactivation, potentially due to differential oligomeric requirements for each activity. The proposed mechanism for Rca-mediated Rubisco reactivation involves threading Rubisco through the AAA+ pore, akin to the method employed by AAA+ ATPases that adjust the conformation of misfolded proteins (Bhat *et al.*, 2017; Houry *et al.*, 2017; Waheeda *et al.*, 2023). This mechanism requires hexameric Rca for Rubisco reactivation, whilst ATP hydrolysis can occur with three Rca subunits forming an oligomer (Keown *et al.*, 2013; Hazra *et al.*, 2015).

The maximum rates of both ATP hydrolysis and Rubisco reactivation were higher for the Rca10 isoforms compared to Rca1β and Rca8α (Table 1). Similar differences among Rca isoforms were observed in rice and wheat but only for ATP hydrolysis, with the more thermostable Rca isoforms exhibiting lower Rubisco reactivation maxima compared to the less thermostable isoforms (Scafaro *et al.*, 2016, 2019a; Shivhare & Mueller-Cajar, 2017; Degen *et al.*, 2020). In cowpea, we show that the increased thermostability of Rca10 isoforms occurs without a penalty in Rubisco reactivation rate. The broader T_{opt} of Rca10α and Rca10β spans and extends above the T_{opt} of Rca1β and Rca8α (Table 1). Thus,

Rca10α and Rca10β appear to be ideal candidates to maintain CO_2 assimilation by Rubisco

both at optimal and supra-optimal temperatures. Curiously, the promoter regions of both

Rca1β and Rca8α contain multiple stress responsive elements while Rca10α and Rca10β

have only two heat response elements (Supporting Information Fig. S12), suggesting an

engineering approach may be required to alter the Rca isoforms predominantly involved in

maintaining Rubisco activity *in planta.*

Cowpea responds dynamically to a 5-day 38/28 °C heatwave

A 5-day heatwave at the vegetative growth stage (10 °C increase to 38/28 °C day/night) is

considered moderate heat stress for most cowpea cultivars (Lonardi *et al.*, 2019; Barros *et*

al., 2021). Plants were well-watered to avoid confounding effects of water deficit limiting the

diffusion of CO_2 to Rubisco sites (Carmo-Silva *et al.*, 2012). The leaf temperature only

increased to 32 °C (Figure 2a), plant growth parameters were only mildly impacted, and the

aboveground dry biomass remained unaffected (Figure 2bc). Notably, it has been shown that

a heatwave during the vegetative stage can have a delayed impact on cowpea yield

(Mohammed *et al.*, 2024), which would not necessarily be expressed as reduced biomass at

the end of the heatwave. T_{crit}, an indicator of thermal lability of PSII, was higher on day 5 of

the 38/28 °C heatwave (Figure 2c) suggesting some capacity for PSII thermal acclimation

(Perez & Feeley, 2020). In wheat, increased T_{crit} after exposure to elevated temperature in

field and controlled environment experiments correlated with a dynamic response of PSII to

heat stress (Posch *et al.*, 2022). On day 5 of the 38/28 °C heatwave, cowpea plants had

decreased Rubisco activities and amount, lower leaf thickness, total chlorophyll and TSP

(Figure 2b), suggesting accelerated leaf senescence. Similar results were observed in heat-

treated wheat (Qu *et al.*, 2023), which could have a delayed impact on biomass production.

Thermotolerant Rca isoforms increase but remain low under heat

At optimal temperatures, *Rca1β* was the isoform most abundantly expressed, with *Rca8α*

following, while *Rca10α* and *Rca10β* were only marginally expressed (Figure 3c). Under

heat stress, *Rca8α*, *Rca10α* and *Rca10β* expression was upregulated, however *Rca10*

transcript levels remained very low compared to *Rca1β* and *Rca8α*. The corresponding

protein abundance was only marginally higher for Rca8α and remained undetectable for

Rca10α and Rca10β (Supporting Information Fig. S13; Figure 3d). The Rca α/β ratio

decreased as the leaves aged, and on day 3 of the heatwave heat-treated plants maintained

higher α/β ratio than control pants (Figure 3d). Variation in Rca α/β ratio under optimal and

supra-optimal temperatures may reflect differences in thermostability and regulatory

mechanisms (Zhang & Portis, 1999; Wang *et al.*, 2010; Degen *et al.*, 2021). While Rca α is

subject to redox regulation via thioredoxin-f (Zhang & Portis, 1999; Shivhare *et al.*, 2019;

Kim *et al.*, 2020) and tends to be more sensitive to inhibition by ADP (Zhang & Portis, 1999;

Scafaro *et al.*, 2019b; Amaral *et al.*, 2024a), variation also exists in the properties of Rca β

isoforms (Scafaro *et al.*, 2019a; Perdomo *et al.*, 2019).

Changes in relative abundance of Rca isoforms under heat stress are species-specific

(Yamori & von Caemmerer, 2009). In *Arabidopsis*, the two isoforms are comparable in

thermotolerance and both increase with heat maintaining a 1:1 ratio (Ristic *et al.*, 2009).

Conversely, in rice and wheat, Rca isoforms differ in temperature response and heat induces

upregulation of the thermotolerant variants (Scafaro *et al.*, 2018, 2019a; Degen *et al.*, 2021).

While the same was observed for cowpea in the present study, the protein abundance of the

thermostable Rca isoforms remained below detection levels in plants exposed to the 5-day

38/28 °C heatwave. Overexpression of thermostable Rca via genetic engineering led to

enhanced photosynthesis and yield (Kurek *et al.*, 2007; Shivhare & Mueller-Cajar, 2017;

Scafaro *et al.*, 2018; Degen *et al.*, 2020). Increasing Rca thermostability, and to a lesser

extent upregulation of its quantity, have been shown to ameliorate the reduction in Rubisco

activation state caused by elevated temperatures (Kurek *et al.*, 2007; Scafaro *et al.*, 2018,

2019a; Kim *et al.*, 2020). The results presented here suggest that increasing Rca10α and

Rca10β abundance in cowpea, while possible in longer or hotter heat conditions, would

likely require an engineering approach.

Rubisco reactivation by Rca in cowpea remains unaltered under heat

The temperature response of Rubisco reactivation in leaf extracts of plants exposed to the 5-

day 38/28 °C heatwave showed a similar pattern and comparable thermal optima to control

plants (Figure 4). Scafaro *et al.* (2019a) reported increases Rubisco reactivation rate and $T_{0.5}$

of Rca in leaf extracts from heat-treated wheat. In cowpea, the minor increase in rate was

statistically insignificant and T_{opt} and $T_{0.5}$ were comparable between control and heat-treated

plants, with a slight increase of the T_{opt} range in heat-treated plants (Table 2). These results

indicate that the heatwave in the absence of water deficit caused only mild heat stress,

allowing plants to respond by adjusting isoform ratios in the leaf while preserving the Rca

rate and temperature response.

Interestingly, the T_{opt} upper bound determined in leaf extracts was 42 °C for control and

heat-treated plants, much higher than the T_{leaf} in heat-treated plants (Table 2, Figure 2a).

The maximum Rca rate in leaf extracts (Table 2) was also higher than the Rubisco

reactivation rate of each isoform measured *in vitro* (Table 1). Similar results were observed

in rice and wheat but only for the more thermostable isoforms (Scafaro *et al.*, 2016, 2019a).

Higher Rca rate in leaf extracts may be attributed to several possible factors: differences in

structural conformation of recombinant Rca, post-translational modifications that regulate

Rca activity in the leaf and are absent in recombinant proteins, hetero-hexamers comprising

different isoforms *in planta* exhibiting higher thermal optima, or interaction of Rca with other

components in the chloroplast. Chen *et al.* (2010) reported dynamic and reversible binding of

rice Rca to the thylakoid membrane while it is active in remodelling the conformation of

inhibitor-bound Rubisco. Binding of Rca to the thylakoid membrane has also been reported

during heat stress although its binding to the chloroplast ribosomes was proposed to have a

protective role (Rokka *et al.*, 2001). Moreover, heat stress has been shown to induce

association of Rca with β-subunit of chaperonin-60 (cpn60β), a potential heat shock

molecular chaperone (Salvucci, 2008). Thus, while *in vitro* analyses are useful to compare

the properties of the individual isoforms, *in vivo* analyses as shown here for cowpea enable

assessment of the overall Rca response within the chloroplast stroma.

Conclusion

Cowpea is a vital source of protein for sub-Saharan Africa, and depite being warm-adapted,

it is threatened by future heatwaves, which are likely to increase in intensity, duration, and

frequency. Here, we identified two isoforms of Rca that have higher thermal maxima,

broader thermal optima, and faster rates of ATP hydrolysis and Rubisco reactivation than the

predominant Rca isoforms present in cowpea leaves. We showed that a 5-day 38/28 °C

heatwave in the absence of water deficit caused leaf temperature to increase to 32 °C. Since

this temperature is below the temperature at which Rca activity decreased below 70% of

maximum for each Rca isoform, the impact of such a heatwave on Rubisco function and

biomass production was only mild. However, cowpea plants growing in smallholder farmer

fields in sub-Saharan Africa are likely to experience water deficit alongside warmer

heatwaves (Almazroui *et al.*, 2020; Song *et al.*, 2023). This combination can impact the

ability of the plant to cool its leaves leading to higher leaf temperatures (Carmo-Silva *et al.*,

2012; Fahad *et al.*, 2017; Sato *et al.*, 2024). Increasing the abundance of the superior

Rca10α and Rca10β isoforms characterised here provides an exciting opportunity to

enhance the resilience of cowpea and other crops to future extreme climates.

Acknowledgements:

We thank Dr Sam Taylor (Lancaster University) for input and discussions around

experimental design and calculations for determining T_{crit}. Ben Davies (Lancaster University)

supported plant maintenance, measurements and sampling. We thank Dr Rachel Baxter

(Lancaster University) for assistance with data analysis, and Dr Joana Amaral for useful

discussions on data interpretation.

Competing interests:

The authors declare no conflicts of interest.

Author contributions:

E.C.S conceived and supervised the research. A.G., R.P., I.R., D.J.O., E.C.S. designed

experiments. M.T.P. designed and executed the cloning strategy. D.B., P.D.G., D.W.

produced and purified recombinant proteins. D.W. and C.J.A. purified Rubisco. D.B. and

A.G. performed ATP hydrolysis measurements. A.G. performed Rubisco reactivation

measurements. R.P. and I.R. performed heatwave experiments. R.P. performed RNAseq,

RT-qPCR and plant growth measurements. R.P. and P.D.G. performed promoter analyses.

I.R. performed fluorescence measurements. A.G., R.P. and C.J.A. determined activity and

amount of Rubisco and Rubisco activase in leaf extracts. A.G., R.P., C.J.A. and I.R.

analysed the data. A.G. and E.C.S. wrote the manuscript with help from all authors.

Data availability:

Data presented are available in the accompanying supporting information and source data.

The RNA-seq data reported in this article has been deposited in DDBJ's Sequence Read

Archive (DRA) and are accessible through DRR BioProject accession number PRJDB19806.

Processed RNA-seq data are submitted in the DDBJ's Genomic Expression Archive (GEA)

and are accessible through the Experiment accession number (E-GEAD-889). Other data

associated with the study are available from the corresponding author.

Funding:

This work was supported by the project Realizing Increased Photosynthetic Efficiency

(RIPE), that is funded by Gates Agricultural Innovations grant investment 57248, awarded to

Lancaster University by the University of Illinois, USA. D.B. was supported by a PhD

scholarship from Lancaster University.

References

Akaike H. **1974**. A new look at the statistical model identification. *IEEE Transactions on Automatic Control* **19**: 716–723.

Almazroui M, Saeed F, Saeed S, Nazrul Islam M, Ismail M, Klutse NAB, Siddiqui MH. **2020**. Projected change in temperature and precipitation over Africa from CMIP6. *Earth Systems and Environment* **4**: 455–475.

Amaral J, Lobo AKM, Carmo-Silva E. **2024a**. Regulation of Rubisco activity in crops. *New Phytologist* **241**: 35–51.

Amaral J, Lobo AKM, Carmo-Silva E, Orr DJ. **2024b**. Purification of Rubisco from Leaves. In: Covshoff S, ed. Photosynthesis : Methods and Protocols. New York, NY: Springer US, 417–426.

Ara N, Nakkanong K, Lv W, Yang J, Hu Z, Zhang M. **2013**. Antioxidant enzymatic activities and gene expression associated with heat tolerance in the stems and roots of two Cucurbit species ('*Cucurbita maxima*' and '*Cucurbita moschata*') and their interspecific inbred line "*Maxchata*". *International Journal of Molecular Sciences* **14**: 24008.

Ashton CJ, Page R, Lobo AKM, Amaral J, Siqueira JA, Orr DJ, Carmo-Silva E. **2024**. Radiometric determination of Rubisco activation state and quantity in leaves. In: Methods in Enzymology. Academic Press.

Bagati S, Rai GK, Bhadwal D, Berwal MK. **2021**. Biotechnological approaches for the development of heat stress tolerance in crop plants. In: Abiotic Stress Tolerance Mechanisms in Plants. CRC Press.

Barros JRA, Guimarães MJM, Silva RM e, Rêgo MTC, de Melo NF, de Melo Chaves AR, Angelotti F. **2021**. Selection of cowpea cultivars for high temperature tolerance: physiological, biochemical and yield aspects. *Physiology and Molecular Biology of Plants* **27**: 29–38.

Barta C, Carmo-Silva AE, Salvucci ME. **2011**. Purification of Rubisco activase from leaves or after expression in *Escherichia coli*. *Methods in Molecular Biology (Clifton, N.J.)* **684**: 363–374.

Becker R, Schüth C, Merz R, Khaliq T, Usman M, Beek T aus der, Kumar R, Schulz S. **2023**. Increased heat stress reduces future yields of three major crops in Pakistan's Punjab region despite intensification of irrigation. *Agricultural Water Management* **281**: 108243.

Bhat JY, Thieulin-Pardo G, Hartl FU, Wendler P, Mili Ci C G, Bracher A, Maxwell A, Ciniawsky S, Mueller-Cajar O, Engen JR, *et al.* 2017. Mechanism of enzyme repair by the AAA + chaperone Rubisco activase. *Molecular Cell* **67**: 744–756.

Blighe, K, S Rana, M Lewis. **2018**. EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling.

Bloemers D, Carmo-Silva E. **2024**. Antibody design for the quantification of photosynthetic proteins and their isoforms. In: Covshoff S, ed. Photosynthesis : Methods and Protocols. New York, NY: Springer US, 405–416.

Bradford MM. **1976**. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* **72**: 248–254.

Bretz F, Hothorn T, Westfall P. **2016**. *Multiple Comparisons Using R*. CRC Press.

Brown AV, Conners SI, Huang W, Wilkey AP, Grant D, Weeks NT, Cannon SB, Graham
MA, Nelson RT. **2021**. A new decade and new data at SoyBase, the USDA-ARS soybean
genetics and genomics database. *Nucleic Acids Research* **49**: D1496–D1501.

Carmo-Silva AE, Gore MA, Andrade-Sanchez P, French AN, Hunsaker DJ, Salvucci ME.
2012. Decreased CO_2 availability and inactivation of Rubisco limit photosynthesis in cotton
plants under heat and drought stress in the field. *Environmental and Experimental Botany* **83**:
1–11.

Carmo-Silva AE, Salvucci ME. **2011**. The activity of Rubisco's molecular chaperone,
Rubisco activase, in leaf extracts. *Photosynthesis Research* **108**: 143–155.

Carmo-Silva AE, Salvucci ME. **2013**. The Regulatory Properties of Rubisco Activase Differ
among Species and Affect Photosynthetic Induction during Light Transitions 1. *Plant*
Physiology Ò **161**: 1645–1655.

Carmo-Silva E, Scales JC, Madgwick PJ, Parry MAJ. **2015**. Optimizing Rubisco and its
regulation for greater resource use efficiency. *Plant, Cell & Environment* **38**: 1817–1832.

Chen H. **2022**. VennDiagram: Generate High-Resolution Venn and Euler Plots.

Chen J-H, Tang M, Jin X-Q, Li H, Chen L-S, Wang Q-L, Sun A-Z, Yi Y, Guo F-Q. **2022**.
Regulation of Calvin–Benson cycle enzymes under high temperature stress. *aBIOTECH* **3**:
65.

Chen J, Wang P, Mi H-L, Chen G-Y, Xu D-Q. **2010**. Reversible association of ribulose-1, 5-
bisphosphate carboxylase/oxygenase activase with the thylakoid membrane depends upon
the ATP level and pH in rice without heat stress. *Journal of Experimental Botany* **61**: 2939–
2950.

Chifflet S, Torriglia A, Chiesa R, Tolosa S. **1988**. A method for the determination of inorganic
phosphate in the presence of labile organic phosphate and high concentrations of protein:
Application to lens ATPases. *Analytical Biochemistry* **168**: 1–4.

Crafts-Brandner SJ, Law RD. **2000**. Effect of heat stress on the inhibition and recovery of
the ribulose-1,5-bisphosphate carboxylase/oxygenase activation state. *Planta* **212**: 67–74.

Crafts-Brandner SJ, Salvucci ME. **2000**. Rubisco activase constrains the photosynthetic
potential of leaves at high temperature and CO_2. *Proceedings of the National Academy of*
Sciences of the United States of America **97**: 13430–13435.

Degen GE, Orr DJ, Carmo-Silva E. **2021**. Heat-induced changes in the abundance of wheat
Rubisco activase isoforms. *New Phytologist* **229**: 1298–1311.

Degen GE, Worrall D, Carmo-Silva E. **2020**. An isoleucine residue acts as a thermal and
regulatory switch in wheat Rubisco activase. *Plant Journal* **103**: 742–751.

Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A, Sadia S, Nasim W, Adkins
S, Saud S, *et al.* 2017. Crop production under drought and heat stress: Plant responses and
management options. *Frontiers in Plant Science* **8**.

Feller U, Crafts-Brandner SJ, Salvucci ME. **1998**. Moderately high temperatures inhibit ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) activase-mediated activation of Rubisco. *Plant Physiol* **116**: 539–546.

Galmés J, Hermida-Carrera C, Laanisto L, Niinemets Ü. **2016**. A compendium of temperature responses of Rubisco kinetic traits: variability among and within photosynthetic groups and impacts on photosynthesis modeling. *Journal of Experimental Botany* **67**: 5067–5091.

Gonçalves A, Goufo P, Barros A, Domínguez-Perles R, Trindade H, Rosa EAS, Ferreira L, Rodrigues M. **2016**. Cowpea (*Vigna unguiculata* L. Walp), a renewed multipurpose crop for a more sustainable agri-food system: nutritional advantages and constraints. *Journal of the Science of Food and Agriculture* **96**: 2941–2951.

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U, Putnam N, *et al.* 2012. Phytozome: a comparative platform for green plant genomics. *Nucleic Acids Research* **40**: D1178–D1186.

Graves S, Piepho H-P, Selzer L, Dorai-Raj S. **2024**. multcompView: Visualizations of Paired Comparisons.

Harboure P. **2024**. WMO Global Annual to Decadal Climate Update.

Hatfield JL, Prueger JH. **2011**. Agroecology: Implications for Plant Response to Climate Change. In: Crop Adaptation to Climate Change. Wiley-Blackwell.

Hazra S, Henderson JN, Liles K, Hilton MT, Wachter RM. **2015**. Regulation of ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) activase. *The Journal of Biological Chemistry* **290**: 24222–24236.

Hermida-Carrera C, Kapralov MV, Galmés J. **2016**. Rubisco catalytic properties and temperature response in crops. *Plant Physiology* **171**: 2549.

Houry WA, Wachter R, Prahlad V, Hayer-Hartl M, Bhat JY, Thieulin-Pardo G, Hartl FU. **2017**. Rubisco activases: AAA+ chaperones adapted to enzyme repair. *Frontiers in Molecular Biosciences* **4**: 20.

Kassambara A. **2023**. ggpubr: 'ggplot2' Based Publication Ready Plots.

Keown JR, Griffin MDW, Mertens HDT, Pearce FG. **2013**. Small oligomers of ribulose-bisphosphate carboxylase/oxygenase (Rubisco) activase are required for biological activity. *The Journal of Biological Chemistry* **288**: 20607–20615.

Keown JR, Pearce FG. **2014**. Characterization of spinach ribulose-1,5-bisphosphate carboxylase/ oxygenase activase isoforms reveals hexameric assemblies with increased thermal stability. *Biochem. J* **464**: 413–423.

Kilian J, Whitehead D, Horak J, Wanke D, Weinl S, Batistic O, D'Angelo C, Bornberg-Bauer E, Kudla J, Harter K. **2007**. The AtGenExpress global stress expression data set: protocols, evaluation and model data analysis of UV-B light, drought and cold stress responses. *The Plant Journal* **50**: 347–363.

Kim SY, Stessman DJ, Wright DA, Spalding MH, Huber SC, Ort DR. **2020**. *Arabidopsis* plants expressing only the redox-regulated Rca-α isoform have constrained photosynthesis and plant growth. *Plant Journal* **103**: 2250–2262.

Kumar A, Li C, Portis AR. **2009**. *Arabidopsis thaliana* expressing a thermostable chimeric Rubisco activase exhibits enhanced growth and higher rates of photosynthesis at moderately high temperatures. *Photosynthesis Research* **100**: 143–153.

Kurek I, Chang TK, Bertain SM, Madrigal A, Liu L, Lassner MW, Zhu G. **2007**. Enhanced thermostability of *Arabidopsis* Rubisco activase improves photosynthesis and growth rates under moderate heat stress. *The Plant Cell* **19**: 3230–3241.

Law RD, Crafts-Brandner SJ. **2001**. High temperature stress increases the expression of wheat leaf ribulose-1,5-bisphosphate carboxylase/oxygenase activase protein. *Archives of Biochemistry and Biophysics* **386**: 261–267.

Li G, Mao H, Ruan X, Xu Q, Gong Y, Zhang X, Zhao N. **2002**. Association of heat-induced conformational change with activity loss of Rubisco. *Biochemical and Biophysical Research Communications* **290**: 1128–1132.

Liang Q, Muñoz-Amatriaín M, Shu S, Lo S, Wu X, Carlson JW, Davidson P, Goodstein DM, Phillips J, Janis NM, *et al.* 2024. A view of the pan-genome of domesticated cowpea (*Vigna unguiculata* [L.] Walp.). *The Plant Genome* **17**: e20319.

Lichtenthaler HK. **1987**. [34] Chlorophylls and carotenoids: Pigments of photosynthetic biomembranes. In: Plant Cell Membranes. Methods in Enzymology. Academic Press, 350–382.

Lin Pedersen T. **2024**. patchwork: The Composer of Plots.

Lonardi S, Ia Mu~ Noz-Amatria In M, Liang Q, Shu S, Wanamaker SI, Lo S, Tanskanen J, Schulman AH, Zhu T, Luo M-C, *et al.* 2019. The genome of cowpea (*Vigna unguiculata* [L.] Walp.). *The Plant Journal* **98**: 767–782.

Martin M. **2011**. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**: 10.

Mathur S, Agrawal D, Jajoo A. **2014**. Photosynthesis: Response to high temperature stress. *Journal of Photochemistry and Photobiology B: Biology* **137**: 116–126.

Mohammed SB, Ongom PO, Togola A, Boukar O. **2024**. Enhancing Cowpea tolerance to elevated temperature: Achievements, challenges and future directions. *Agronomy* **14**: 513.

Moore CE, Meacham-Hensold K, Lemonnier P, Slattery RA, Benjamin C, Bernacchi CJ, Lawson T, Cavanagh AP. **2021**. The effect of increasing temperature on crop photosynthesis: from enzymes to ecosystems. *Journal of Experimental Botany* **72**: 2822–2844.

Muggeo VMR. **2017**. Interval estimation for the breakpoint in segmented regression: a smoothed score-based approach. *Australian & New Zealand Journal of Statistics* **59**: 311–322.

Murchie EH, Lawson T. **2013**. Chlorophyll fluorescence analysis: a guide to good practice and understanding some new applications. *Journal of Experimental Botany* **64**: 3983–3998.

Nagarajan R, Gill KS. **2018**. Evolution of Rubisco activase gene in plants. *Plant Molecular Biology* **96**: 69–87.

Orr DJ, Robijns AKJ, Baker CR, Niyogi KK, Carmo-Silva E. **2023**. Dynamics of Rubisco regulation by sugar phosphate derivatives and their phosphatases. *Journal of Experimental Botany* **74**: 581–590.

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. **2017**. Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods* **14**: 417–419.

Perdomo JA, Buchner P, Carmo-Silva E. **2021**. The relative abundance of wheat Rubisco activase isoforms is post-transcriptionally regulated. *Photosynthesis Research* **148**: 47–56.

Perdomo JA, Capó-Bauçà S, Carmo-Silva E, Galmés J. **2017**. Rubisco and Rubisco activase play an important role in the biochemical limitations of photosynthesis in rice, wheat, and maize under high temperature and water deficit. *Frontiers in Plant Science* **8**: 490.

Perdomo JA, Degen GE, Worrall D, Carmo-Silva E. **2019**. Rubisco activation by wheat Rubisco activase isoform 2β is insensitive to inhibition by ADP. *Biochemical Journal* **476**: 2595–2606.

Perdomo JA, Sales CRG, Carmo-Silva E. **2018**. Quantification of photosynthetic enzymes in leaf extracts by immunoblotting. *Methods in Molecular Biology (Clifton, N.J.)* **1770**: 215–227.

Perez TM, Feeley KJ. **2020**. Photosynthetic heat tolerances and extreme leaf temperatures. *Functional Ecology* **34**: 2236–2245.

Portis AR. **2003**. Rubisco activase - Rubisco's catalytic chaperone. *Photosynthesis Research* **75**: 11–27.

Portis AR, Li C, Wang D, Salvucci ME. **2008**. Regulation of Rubisco activase and its interaction with Rubisco. *Journal of Experimental Botany* **59**: 1597–1604.

Posch BC, Hammer J, Atkin OK, Bramley H, Ruan Y-L, Trethowan R, Coast O. **2022**. Wheat photosystem II heat tolerance responds dynamically to short- and long-term warming. *Journal of Experimental Botany* **73**: 3268–3282.

Qu L, Gu X, Li J, Guo J, Lu D. **2023**. Leaf photosynthetic characteristics of waxy maize in response to different degrees of heat stress during grain filling. *BMC Plant Biology* **23**: 469.

Ristic Z, Momčilović I, Bukovnik U, Prasad PVV, Fu J, DeRidder BP, Elthon TE, Mladenov N. **2009**. Rubisco activase and wheat productivity under heat-stress conditions. *Journal of Experimental Botany* **60**: 4003–4014.

Robinson SP, Portis AR. **1989**. Adenosine triphosphate hydrolysis by purified Rubisco activase. *Archives of Biochemistry and Biophysics* **268**: 93–99.

Rokka A, Zhang L, Aro EM. **2001**. Rubisco activase: An enzyme with a temperature-dependent dual function? *Plant Journal* **25**: 463–471.

Ruuska S, Andrews TJ, Badger MR, Hudson GS, Laisk A, Price GD, Caemmerer S von. **1998**. The interplay between limiting processes in C3 photosynthesis studied by rapid-response gas exchange using transgenic tobacco impaired in photosynthesis. *Functional Plant Biology* **25**: 859–870.

Sales CRG, Molero G, Evans JR, Taylor SH, Joynson R, Furbank RT, Hall A, Carmo-Silva E. **2022**. Phenotypic variation in photosynthetic traits in wheat grown under field versus glasshouse conditions. *Journal of Experimental Botany* **73**: 3221–3237.

Salvucci ME. **2008**. Association of Rubisco activase with chaperonin-60b: a possible mechanism for protecting photosynthesis during heat stress. *Journal of Experimental Botany* **59**: 1923–1933.

Salvucci ME, Crafts-Brandner SJ. **2004**. Relationship between the heat tolerance of photosynthesis and the thermal stability of Rubisco activase in plants from contrasting thermal environments. **134**: 1460–1470.

Salvucci ME, DeRidder BP, Portis AR. **2006**. Effect of activase level and isoform on the thermotolerance of photosynthesis in Arabidopsis. *Journal of Experimental Botany* **57**: 3793–3799.

Salvucci ME, Osteryoung KW, Crafts-Brandner SJ, Vierling E. **2001**. Exceptional Sensitivity of Rubisco Activase to Thermal Denaturation in Vitro and in Vivo. *Plant Physiology* **127**: 1053–1064.

Sato H, Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K. **2024**. Complex plant responses to drought and heat stress under climate change. *The Plant Journal* **117**: 1873–1892.

Scafaro AP, Atwell BJ, Muylaert S, Reusel BV, Ruiz GA, Rie JV, Gallé A. **2018**. A thermotolerant variant of Rubisco activase from a wild relative improves growth and seed yield in rice under heat stress. *Frontiers in Plant Science* **9**: 1663.

Scafaro AP, Bautsoens N, den Boer B, Van Rie J, Gallé A. **2019a**. A Conserved sequence from heat-adapted species improves Rubisco activase thermostability in wheat. *Plant Physiology* **181**: 43–54.

Scafaro AP, Gallé A, Van Rie J, Carmo-Silva E, Salvucci ME, Atwell BJ. **2016**. Heat tolerance in a wild *Oryza* species is attributed to maintenance of Rubisco activation by a thermally stable Rubisco activase ortholog. *New Phytologist* **211**: 899–911.

Scafaro AP, Posch BC, Evans JR, Farquhar GD, Atkin OK. **2023**. Rubisco deactivation and chloroplast electron transport rates co-limit photosynthesis above optimal leaf temperature in terrestrial plants. *Nature Communications* **14**: 2820.

Scafaro AP, Vleesschauwer DD, Bautsoens N, Hannah MA, Boer B den, Gallé A, Rie JV. **2019b**. A single point mutation in the C-terminal extension of wheat Rubisco activase dramatically reduces ADP inhibition via enhanced ATP binding affinity. *The Journal of Biological Chemistry* **294**: 17931.

Serban AJ, Breen IL, Bui HQ, Levitus M, Rebekka Wachter XM, Jez JM. **2018**. Assembly–disassembly is coupled to the ATPase cycle of tobacco Rubisco activase. *J. Biol. Chem* **293**: 19451–19465.

Shivhare D, Mueller-Cajar O. **2017**. *In vitro* characterization of thermostable CAM Rubisco activase reveals a Rubisco interacting surface loop. **174**: 1505–1516.

Shivhare D, Ng J, Tsai Y-CC, Mueller-Cajar O. **2019**. Probing the rice Rubisco-Rubisco activase interaction via subunit heterooligomerization. *Proceedings of the National Academy of Sciences of the United States of America* **116**: 24041–24048.

Song J, Tong G, Chao J, Chung J, Zhang M, Lin W, Zhang T, Bentler PM, Zhu W. **2023**. Data driven pathway analysis and forecast of global warming and sea level rise. *Scientific Reports* **13**: 5536.

Song K, Yim WC, Lee B-M. **2017**. Expression of heat shock proteins by heat stress in soybean. *Plant Breeding and Biotechnology* **5**: 344–353.

Stainbrook SC, Aubuchon LN, Chen A, Johnson E, Si A, Walton L, Ahrendt AJ, Strenkert D, Jez JM. **2024**. C4 grasses employ distinct strategies to acclimate Rubisco activase to heat stress. *Bioscience Reports* **44**.

Stotz M, Mueller-Cajar O, Ciniawsky S, Wendler P, Hartl FU, Bracher A, Hayer-Hartl M. **2011**. Structure of green-type Rubisco activase from tobacco. *Nature Structural & Molecular Biology* **18**: 1366–1370.

Taylor SH, Gonzalez-Escobar E, Page R, Parry MAJ, Long SP, Carmo-Silva E. **2022**. Faster than expected Rubisco deactivation in shade reduces cowpea photosynthetic potential in variable light conditions. *Nature Plants* **8**: 118–124.

Taylor SH, Orr DJ, Carmo-Silva E, Long SP. **2020**. During photosynthetic induction, biochemical and stomatal limitations differ between Brassica crops. *Plant, Cell & Environment* **43**: 2623–2636.

Waheeda K, Kitchel H, Wang Q, Chiu P-L. **2023**. Molecular mechanism of Rubisco activase: Dynamic assembly and Rubisco remodeling. *Frontiers in Molecular Biosciences* **10**: 1125922.

Wang D, Li X-F, Zhou Z-J, Feng X-P, Yang W-J, Jiang D-A. **2010**. Two Rubisco activase isoforms may play different roles in photosynthetic heat acclimation in the rice plant. *Physiologia Plantarum* **139**: 55–67.

Warrag MOA, Hall AE. **1984**. Reproductive responses of cowpea (*Vigna unguiculata* (L.) Walp.) to heat stress. II. Responses to night air temperature. *Field Crops Research* **8**: 17–33.

Weber E, Engler C, Gruetzner R, Werner S, Marillonnet S. **2011**. A modular cloning system for standardized assembly of multigene constructs. *PLOS ONE* **6**: e16765.

Werner S, Engler C, Weber E, Gruetzner R, Marillonnet S. **2012**. Fast track assembly of multigene constructs using Golden Gate cloning and the MoClo system. *Bioengineered Bugs* **3**: 38–43.

Wickham H. **2017**. Tidyverse: Easily install and load the 'tidyverse'. *R package version* **1**: 2017.

Wijewardene I, Shen G, Zhang H. **2021**. Enhancing crop yield by using Rubisco activase to improve photosynthesis under elevated temperatures. **1**: 2.

Winter D, Vinegar B, Nahal H, Ammar R, Wilson GV, Provart NJ. **2007**. An "Electronic fluorescent pictograph" browser for exploring and analyzing large-scale biological data sets. *PLOS ONE* **2**: e718.

Wood SN. **2017**. *Generalized Additive Models: An Introduction with R, Second Edition*. New York: Chapman and Hall/CRC.

Yamori W, von Caemmerer S. **2009**. Effect of Rubisco activase deficiency on the temperature response of CO_2 assimilation rate and Rubisco activation state: Insights from

transgenic tobacco with reduced amounts of Rubisco activase. *Plant Physiology* **151**: 2073–
2082.

Yin Z, Zhang Z, Deng D, Chao M, Gao Q, Wang Y, Yang Z, Bian Y, Hao D, Xu C. **2014**.
Characterization of Rubisco activase genes in maize: An α-isoform gene functions alongside
a β-isoform gene. *Plant Physiology* **164**: 2096–2106.

Zahra N, Hafeez MB, Wahid A, Al Masruri MH, Ullah A, Siddique KHM, Farooq M. **2023**.
Impact of climate change on wheat grain composition and quality. *Journal of the Science of*
Food and Agriculture **103**: 2745–2751.

Zhang N, Kallis RP, Ewy RG, Portis AR. **2002**. Light modulation of Rubisco in *Arabidopsis*
requires a capacity for redox regulation of the larger Rubisco activase isoform. *Proceedings*
of the National Academy of Sciences of the United States of America **99**: 3330–3334.

Zhang N, Portis AR. **1999**. Mechanism of light regulation of Rubisco: A specific role for the
larger Rubisco activase isoform involving reductive activation by thioredoxin-f. *Plant Biology*
96: 9438–9443.

1184

Supporting Information

Supporting Information Figures

Fig. S1. Amino acid residue alignment of the four cowpea Rca isoforms.

Fig. S2. *In vitro* Rubisco reactivation by the four cowpea Rca isoforms.

Fig. S3. Heatwave experimental design and cabinet temperature monitoring.

Fig. S4. Cabinet plant layout design.

Fig. S5. Leaf temperature monitoring over the heatwave experiment.

Fig. S6. Growth parameters and chlorophyll content during and after heat stress.

Fig. S7. T_{crit} calculation from temperature response of chlorophyll fluorescence-derived Fv/Fm.

Fig. S8. Specific Rubisco activities of control and heat-treated plants.

Fig. S9. Differential gene expression in heat-treated *versus* control cowpea plants.

Fig. S10. Volcano plots depicting differential gene expression based on log change across the
days of heatwave.

Fig. S11. Gene expression of heat shock protein 20 (HSP20) and cowpea Rca transcripts.

Fig. S12. Potential cis-acting regulatory elements in cowpea Rca promoter regions.

Fig. S13. Protein abundance of Rca isoforms in leaves of control and heat-treated plants.

Fig. S14. Protocol outline for determination of Rca activity in leaf extracts (LE).

Fig. S15. Rubisco reactivation by the pool of cowpea Rca isoforms in leaf extracts (LE).

Supporting Information Tables

Table S1. Reference protein sequences used to identify cowpea Rca genes.

Table S2. Primer sequences for adding Golden Gate overhangs to Rca coding regions.

Table S3. Optimum temperature of in vitro Rubisco activase (Rca) activity.

Table S4. Modelling of the *in vitro* temperature response of ATP hydrolysis and Rubisco
activation by cowpea Rca isoforms.

Table S5. RNA sample QC analysis prior to RNA sequencing.

Table S6. Sequencing and alignment statistics.

Table S7. MIQE checklist for RT-qPCR.

Table S8. RT-qPCR primers.

Table S9. Modelling of the temperature response of Rubisco activation by cowpea Rca in leaf
extracts of control and heat-treat plants.

Table S10. Leaf total soluble protein (TSP) and chlorophyll content.

References