Under the cover of darkness: a transcriptomic exploration of clubroot during the night.

Andrea Garvetto[1], Susann Auer[2], Freia Benade[2], Michaela Hittorf[1,3], Jutta Ludwig-Müller[2], Sigrid Neuhauser[1]

Corresponding authors: Andrea Garvetto (andrea.garvetto@uibk.ac.at), Sigrid Neuhauser (sigrid.neuhauser@uibk.ac.at)
[1]Department of Microbiology, Universität Innsbruck, Technikerstraße 25, 6020 Innsbruck, Tirol, Austria.
[2]Faculty of Biology, Technische Universität Dresden, 01062 Dresden, Germany.
[3]Department of Botany, Universität Innsbruck, Sternwartestraße 15, 6020 Innsbruck, Tirol, Austria.

Summary

Plasmodiophora brassicae (Phytomyxea, Rhizaria) is the etiological agent of clubroot disease, one of the most important diseases of Brassicaceae crops. Alteration of metabolism and hormone homeostasis leads to the formation of tumour-like galls in the roots of affected plants. Host plant energy metabolism, defence and developmental processes are under strong temporal control and the very same processes are affected by clubroot. For the first time, this study uses time-resolved transcriptome analyses to explore how *P. brassicae* affects *Arabidopsis thaliana* in the night during intermediate (14 days after inoculation; DAI) and late (21 DAI) infection. Day-night differences in gene expression were more pronounced in younger rather than older plants in our differential gene expression (DGE) analysis. Consequently, intermediate phases of infection showed more day-night differences than later ones. Clustering of differentially expressed genes (DEGs) in functional categories highlighted how some of the typical processes known to be disrupted by clubroot infection are more significantly affected in the night and also uncovered some disrupted exclusively in the night. RNA modification stood out as the most unambiguously upregulated process in infected *Arabidopsis* roots in the night. Analysis of the interaction between clubroot infection and diel oscillations in gene expression detected modifications in the rhythmicity of central circadian clock components during the infection. We discuss our findings in the context of manipulation of plant defence and metabolism, identifying targets for experimental validation and highlighting potential new lines of investigation of our time-resolved datasets to better understand the interaction between *P. brassicae* and its host.

Significance statement: *Plasmodiophora brassicae* impacts physiological processes under strong temporal control in Brassicaceae hosts: e.g., metabolism, hormone homeostasis and defence. Here, for the first time, we performed a time-resolved transcriptomic exploration of clubroot disease at intermediate and mature stages of infection. We identify a previously unrecognised role for nocturnal manipulation of organellar RNA editing and disruption of rhythmicity in circadian clock components. We provide a dataset enabling further exploration of the impact of clubroot on plant circadian processes.

Keywords: Biotrophic parasite, Circadian, Host-pathogen interactions, *Plasmodiophora brassicae,* Protist, Time-resolved transcriptome, *Arabidopsis thaliana*.

Introduction

Responsible for around 10 % of harvest losses worldwide, *Plasmodiophora brassicae* Woronin is one of the most impactful pathogens of Brassicaceae crops (Dixon, 2009). This obligate biotrophic parasitic protist belonging to the class Phytomyxea (Rhizaria; Sierra et al., 2016) has a complex life cycle featuring a short-lasting, symptomless primary phase occurring in the root hairs and epidermis of a broad array of plant hosts; and a secondary, long-lasting and symptomatic phase, specifically occurring in hosts within the family Brassicaceae where durable resting spores are formed (Kageyama & Asano, 2009; Liu et al., 2020). The secondary phase of the infection occurs in the root cortex and stele, where it impacts the host physiology and anatomy, causing successive hyperplasia and hypertrophy in the affected tissues (Olszak et al., 2019), leading to the typical gall formation from which the common name "clubroot disease" derives. This modification of the host anatomy and physiology is required for the parasite to generate a strong local nutrient sink at the site of infection, inducing a systemic redirection of the host photosynthates (Malinowski et al., 2019). Clubroot impact on susceptible plants is heavy, with infected hosts featuring stunted growth, increased susceptibility to wilting and early onset of senescence (Javed et al., 2023). Molecular investigations have been proven paramount in understanding the influence of *P. brassicae* on its host physiology (Agarwal et al., 2011; Ciaghi et al., 2019; Irani et al., 2018; Siemens et al., 2006), highlighting highly impacted processes deserving more extensive investigation, such as: cell wall expansion, remodelling and lignification (Stefanowicz et al., 2021; Tu et al., 2024); cell cycle activation and modification (Olszak et al., 2019) and sugar homeostasis (Siemens et al., 2011). Pathogen-induced disruption of sugar homeostasis is crucial to the trophic interaction between *P. brassicae* and its host, whereby the parasite generates a sugar sink in the gall tissue by increasing symplastic sugar delivery via extracellular invertases (Siemens et al., 2011) and by promoting phloem differentiation and local accumulation of sugar transporters at the site of infection (Walerowski et al., 2018).

Because of its central role in plant physiology, the disruption of sugar metabolism likely mediates

systemic reactions at the organism level in clubroot disease, but the substantial anatomical changes

observed are also the result of its impact on hormone homeostasis (Devos et al., 2006; Ludwig-Müller et

al., 2009). Hormones involved in plant defence such as salicylic acid (SA) and jasmonic acid (JA) have

been broadly linked to heightened resistance against *P. brassicae* (Gravot et al., 2012; Lemarié et al.,

2015; Lovelock et al., 2013, 2016); and the parasite seems to directly disrupt both (Ludwig-Müller et al.,

2015; Smolko et al., 2024). *P. brassicae* also induces accumulation of the stress-related hormone abscisic

acid (ABA) as a consequence of the impaired vascular transport and water supply caused by gall

formation (Ludwig-Müller et al., 2009). *Arabidopsis* mutants impaired in the perception of the stress-

related hormone ethylene (ET) showed increased resistance only when the mutation targeted the

crosstalk between ethylene and auxin transport (Ludwig-Müller et al., 2009; Vandenbussche et al.,

2003). Indeed, auxins have been found to play an important role in clubroot disease, gradually increasing

in the galls with the development of the infection and mutants impaired in indole acetic acid (IAA)

biosynthesis were found to produce smaller or developmentally delayed galls (Grsic-Rausch et al., 2000;

Neuhaus et al., 2000). Investigations highlighted how the *P. brassicae* protein PbGH3 can conjugate

auxins (Smolko et al., 2024) and the recently identified gene *SYNERGISTIC ON AUXIN AND CYTOKININ 1*

(*SYAC1*), a cross-talk component between the auxin and cytokinin pathways in *A. thaliana* roots, made

plants more susceptible to infection and exacerbated the symptoms of clubroot when constitutively

expressed; supporting a pivotal role of the auxin-cytokinin balance in this interaction (Hurný et al., 2020).

In biotrophic interactions between plants and pathogens, the role of cytokinins in establishing local

nutrient sinks and in suppressing cell death has long been recognized (Walters & McRoberts, 2006),

nonetheless the overall impact of *P. brassicae* on host cytokinins is ambiguous. Studies have reported

both suppression and increase in metabolism and levels of cytokinins (Dekhuijzen, 1981; Dekhuijzen &

Overeem, 1971; Malinowski et al., 2016), but decreased cytokinin degradation on the host side (Siemens

et al., 2006) coupled with evidence of endogenous cytokinin production by the pathogen (Müller &

Hilgenberg, 1986; Rolfe et al., 2016) seem to support the need for cytokinins in the local establishment

of a *de novo* meristematic sink area in the initial phases of infection (Devos et al., 2005, 2006).

The above listed processes impacted by *P. brassicae* are known to be strongly influenced by diel (i.e., day

and night) and circadian oscillations in plants. While most living organisms have evolved a way to

synchronise their physiology to the environmental rhythms in order to maximize their fitness, plants are

particularly affected by these rhythms as they rely for growth and development on light, arguably one of

the most rhythmic timers/*zeitgebers* on the planet (Henriques et al., 2018; Sanchez & Kay, 2016).

Changes in light and other oscillating environmental parameters such as temperature, entrain the core

circadian clock, a transcriptional/translational feedback loop central to the adjustment of rhythmic

physiological responses in plants (S. Wang et al., 2022). Moreover, the same rhythmic physiological

processes influenced by environmental factors via the circadian clock also act back on it as endogenous

zeitgebers such as in the case of sugars derived from photosynthetic carbon fixation, whose correlation

with the availability of diurnal light has long been ascertained (Graf et al., 2010; Mora-García et al.,

2017).

Besides being under strong temporal control, the availability of photosynthates is also spatially resolved.

For example, a sucrose transporter involved in phloem unloading has been found to be specifically

upregulated in roots during the night, matching a steady increase in nocturnal root growth that peaks

shortly after dawn (Durand et al., 2018; Yazdanbakhsh et al., 2011; Yazdanbakhsh & Fisahn, 2010). The

synthesis and accumulation of defence hormones has also been reported to be under the influence of

diel oscillations in gene expression, in particular the synthesis and accumulation of competing JA and SA;

with JA-mediated resistance to herbivores and necrotrophic pathogens being upregulated in the day,

whilst SA-mediated resistance to biotrophic parasites heightened in the night (Goodspeed et al., 2012).

As already mentioned, the majority of the processes under circadian control also provide retroactive

feedback to inform and entrain the circadian clock itself (S. Wang et al., 2022). Since pathogens are

known to manipulate hormones and to strongly affect energy homeostasis, it is possible that these

processes are central to mediate pathogen-induced modifications on the host internal clock (Lu et al.,

2017).

Many biotrophic parasites have been shown to possess intrinsic transcriptionally-mediated rhythms

(Rijo-Ferreira et al., 2017, 2020) and in well-investigated pathosystems, such as *Plasmodium* spp. in

vertebrates, the parasite contemporarily synchronizes with (Motta et al., 2023) and disrupts (Prior et al.,

2019) the circadian rhythmicity of the host.

We reasoned that, as an obligate biotrophic intracellular parasite, *P. brassicae* is likely to have evolved

similar ways to influence the rhythmic gene expression of its host in order to maximize its benefits and

hinder defence; especially considering its strong influence on host diel processes such as for example

carbohydrates allocation to the root (Graf et al., 2010; Malinowski et al., 2019; Yazdanbakhsh et al.,

2011). Nevertheless, until now only the diurnal phase of clubroot infection has been explored. This study

aims at providing the first time-resolved RNA-seq dataset, covering both day and night, for intermediate

(14 days post-inoculation) and mature (21 days post-inoculation) clubroot disease in *Arabidopsis*

thaliana. We provide a first exploration of the data generated in order to highlight processes of the host

that are influenced by the parasite specifically during the night and further to understand the interplay

between diurnal/nocturnal changes in gene expression and clubroot disease.

Results

Overview of gene expression levels in infected and healthy plants in the day and in the night

The major factor influencing the clustering of DEGs of *A. thaliana* in PCAs is their response to *P. brassicae*

(Fig. 1 A & C). In the intermediate phases of infection (14 DAI) the first two principal components (PCs)

explain a cumulative 78% of the variation (Fig. 1 A). Infected and healthy plants strongly separate along

the PC1, explaining more than half of the variation in gene expression (63%). Noticeably, the PC2

(explaining 15% of the variation) mostly separated the samples according to night or day (factor "Time

point"). Samples collected at 21 DAI seem to be less affected by the time of sampling, with PC1

explaining 83% of the variation and being clearly linked to the infected or healthy status of the plant

samples, and PC2 explaining only 5% of the variation (Fig. 1 C) without clear separation of the samples

along the factor "Time point". A PCA of the gene expression levels in uninfected plants resulted in a

similar pattern, whereby older plants did not show a time-dependent response (Additional File 3).

Befittingly, in plants 14 DAI more than one third (36,75%) of the differentially expressed genes (DEGs)

are specific to the night phase and 16,40% to the day (Fig. 1 B), whilst the proportion of DEGs that are

exclusively detected during day or night is decreasing at 21 DAI indicating a reduced influence of the

factor "Time point" on gene expression from the intermediate to the late stages of infection (21,48% in

the night, 17,52% in the day and 61% shared; Fig. 1 D).

Regulation of enriched GO categories in infected plants during day and night

Differentially expressed genes (DEGs) were largely downregulated in infected plants across all sampling

times and DAI, as shown in Additional Files 4 and 5. This overall trend was further confirmed through an

analysis of average expression levels within overrepresented Gene Ontology (GO) categories, which

revealed a consistent predominance of downregulated over upregulated categories.

At 14 DAI during the day, this imbalance was particularly evident: 63 GO categories showed

downregulation compared to just 25 that were upregulated, while one category— "response to molecule

of fungal origin"—exhibited an equal mix of both (Fig. 2A and Additional File 6).

The ten most significantly upregulated categories during the day were associated with responses to

environmental stress, development, and biosynthesis, with highly significant enrichment seen in

processes such as response to toxic substances, anatomical structure morphogenesis, pigment

biosynthesis, and photosynthetic activity. In contrast, the most significantly downregulated categories

were related to core metabolic and defense responses, including hypoxia response, carbohydrate

metabolism, salicylic acid signaling, and floral development (Fig. 2 A and Additional File 6).

A similar expression pattern was observed during the night at 14 DAI, with 67 GO categories

downregulated and 30 upregulated, alongside five showing no overall change (Fig. 2B and Additional File

6). Upregulated categories at night included many processes related to stress-response and signaling

such as response to toxic substance, RNA modification, ABC-type transporter activity, and response to

reactive oxygen species. Meanwhile, downregulated categories again pointed to a suppression of key

metabolic and defense functions, including phenylpropanoid metabolism, oxygen response, glycoside

metabolism, and response to salicylic acid.

By 21 DAI, the contrast between down- and upregulated categories became even more pronounced,

consistent with the higher number of DEGs observed at this later stage of infection (Fig. 3, Additional

Files 5 and 9). During the day, 94 categories were downregulated compared to just 15 upregulated.

These upregulated categories continued to reflect photosynthetic and light-responsive processes, such

as plastid organization, pigment biosynthesis, circadian rhythm regulation, and peroxisomal activity.

Conversely, the most downregulated categories included pathways related to toxic substance response,

carbohydrate metabolism, extracellular signaling, and salicylic acid response, indicating a persistent

suppression of metabolic and defense processes.

At night, the upregulated GO categories again included several also enriched during the day, such as

photosynthesis and plastid organization. Additional night-specific upregulated processes included

responses to light intensity, chlorophyll biosynthesis, and abiotic stress. Downregulated categories at

night often mirrored those observed during the day, particularly in areas such as carbohydrate

metabolism, tetrapyrrole binding, and external encapsulating structure organization. However, some

processes—such as inorganic ion transport and secondary metabolite biosynthesis—were uniquely or

more strongly suppressed during the night phase.

All GO terms and their associated gene lists used prior to plotting in Figures 2 and 3 are provided in

tabular format in Additional File 10.

Effect of the interaction between clubroot and time of the day on A. thaliana gene expression

The variables "Condition" and "Time Point" are not fully independent, both influence gene expression at

the same time. To identify the genes that are affected by both of these variables we used a statistical

"interaction effect" model (IE; Duda et al., 2023). This identified 217 functionally annotated genes at 14

DAI which were strongly impacted by both variables (Fig. 4). These 217 DEGs can be grouped in eight

clusters according to their expression patterns. Enriched GO terms could be detected in 6 of these

clusters (cluster 1, 3, 4, 5, 6 and 8) hinting at how infection and time of the day are affecting processes in

the host (Additional File 8 GO terms overrepresentation analysis of clusters of DEGs driven by the

interaction between infection and time of the day 14 DAI8).

Clusters 1 and 3 are characterised by a loss of diel oscillation in infected plants, where the genes are

always active regardless of the time of the day. Cluster 1 is enriched in the terms related to senescence,

circadian rhythm and cell signalling, where control plants show upregulation during the day. Cluster 3 is

characterised by the terms "response to nitrate" (particularly enriched in genes encoding glutaredoxins

or GRXS; Ehrary et al., 2020) and terms related to the chloroplast membrane (enriched in genes encoding

chlorophyllide a oxygenase, chloro-ribosomal subunits and FKBP16-2), terms showing upregulation in the

night in the control. Cluster 7 also loses diel oscillation, but the changes in this cluster are characterised

by a downregulation of genes in infected plants which are upregulated during day in control plants.

In contrast cluster 2 acquires a diel oscillation in infected plants with genes being upregulated in the day

and downregulated in the night, an oscillation absent in the control plants. Genes in cluster 6 show an

inversion of the normal pattern of expression, whereby nocturnal upregulation in the control is

overridden in infected plants by a nocturnal downregulation (and vice versa). Cluster 6 is enriched in

term "response to fructose". Cluster 4, enriched in terms related to glycosylation of proteins, shows upregulation by the infection with a stronger effect during the day. Clusters 5 and 8 show no distinct pattern. Cluster 5 is associated with the terms "phosphatidylinositol phosphate kinase activity", "phosphotransferase activity, phosphate group as acceptor" and "nucleoside transmembrane transporter activity" and cluster 8 with terms related to cellular response to abiotic stress and protein localization to the nucleus.

Fewer genes were impacted by the interaction between time of day and infection at 21 days post-infection (DAI) compared to 14 DAI (96 genes with functional annotations). The clustering of the samples cannot discriminate day and night but it maintains a clear separation along infected and control plants (Fig. 5). The pattern of gene expression could divide the DEGs in five clusters, three of which contain enriched GO terms (Additional File 9). Genes in cluster 1 and cluster 2 show a general upregulation in infected plants and in cluster 2 terms related to response to hypoxia and sugars were enriched. Cluster 3 shows an opposite pattern, with genes mainly downregulated by the infection enriching categories related to the development of anatomical structures and cytoplasmic stress granules. The main expression pattern identified in cluster 4 is the upregulation of genes normally downregulated during the night. This cluster is described by GO terms related to circadian/rhythmic processes, chloroplast membranes, histone methylation and transport of organic acids. Cluster 5 shows a more complex expression pattern, were most of the genes upregulated in the night in control conditions are upregulated in the night in infected conditions and vice versa, but no enriched GO terms were found for this cluster.

Discussion

This study explores the effects of *P. brassicae* infection on transcriptional diel oscillations in *A. thaliana*, identifying significant effects on gene expression dynamics. Intermediate infection stages (14 DAI) show strong diel influence, aligning with peak metabolic activity of the parasite and initiation of gall formation

(Devos et al., 2005; Liu et al., 2020), whereas later stages (21 DAI) reveal reduced rhythmicity, likely

driven by plant senescence rather than the pathogen. Nonetheless, overall the strongest driver of gene

expression changes is the infection itself, overriding time-of-day effects, and highlighting a complex

interaction between host circadian mechanisms and pathogen strategies. Processes such as

phenylpropanoid metabolism and RNA modification are disproportionately affected at night and we link

these to disruptions of organellar dynamics and defense in infected roots.

Time-resolved RNA-seq analysis of clubroot disease in *A. thaliana* shows an increasing number of DEGs

as the infection advances, reflecting the progressive cell-by-cell colonisation of roots by *P. brassicae*

(Kunkel, 1918) and its systemic and diversified impact on the plant, where each cell in infected tissues

may contain parasitic plasmodia at different stages of development. Existing transcriptomic datasets

featuring a developmental dimension support this (Irani et al., 2018; Schuller et al., 2014). The effect of

diel changes in gene expression in infected plants appear to be stronger at 14 DAI as compared to 21 DAI

(Fig.1) and a decreased influence of day and night on gene expression is also evident in control plants

(Additional File 3) indicating a general circadian dampening with plant age. The relationship between

senescence and circadian rhythm in ageing plants has been studied in leaves, revealing a similar pattern

of reduced amplitude and an increase in frequency of circadian oscillations with age (Kim & Hong, 2019;

Lee et al., 2021; Song et al., 2018). A specific circadian clock or 'root clock' has been shown to integrate

exogenous cues such as light, water availability and temperature in order to adjust root gene oscillations

with the external environment (Bustillo-Avendaño et al., 2022). The root clock is closely linked to shoot

apical meristem circadian oscillations (Davis et al., 2022; Mortada et al., 2024; Perez-Garcia et al., 2022)

and likely influenced by senescence in above ground tissues, although specific work on the root

developmental clock has not yet addressed its interplay with root senescence itself (Siqueira et al.,

2022). At 21 DAI clubroot infected *A. thaliana* plants are four weeks old and therefore approaching the

end of their six-week life-cycle (Meinke et al., 1998). This advanced senescence would explain the

reduced influence of diel rhythmicity observed at 21 DAI (Fig. 1 C&D, 3 and 5). In contrast, a clear

influence of night and day has been observed at 14 DAI (14 DAI; Fig. 1 A&B, 2 and 4).

The infection is the strongest driver of changes in gene expression (Fig. 1 A), but the genes affected differ

between day and night especially in the intermediate phases (i.e., 14 DAI; Fig. 2 and 3; Additional Files 6

and 7). GO categories enriched by infection-related DEGs during the day are in line with previous,

comparable studies (Ciaghi et al., 2019; Irani et al., 2018). Nonetheless, interpreting these results

requires careful consideration, since each enriched GO category contains upregulated and

downregulated genes and since the z-score reflects only the count of genes up/downregulated but not

the magnitude of changes. For example, specific genes involved in low oxygen responses during clubroot

such as PYRUVATE DECARBOXYLASE 1 (PDC1) and ALCOHOL DEHYDROGENASE 1 (ADH1) have been

found to be upregulated in previous studies (Gravot et al., 2016). These specific genes are also

upregulated in our dataset, even though low oxygen response categories are overall downregulated (Fig.

2). Similarly, cell wall-related categories are consistently downregulated in our dataset, though individual

genes like expansins and pectin methylesterases show contrasting gene-specific expression patterns,

which is in line with previous findings (Ciaghi et al., 2019; Irani et al., 2018; Stefanowicz et al., 2021).

The GO term "RNA modification" was the most striking night-exclusive category in terms of differential

expression, gene counts and significance at 14 DAI. The majority of genes (90/101) are assigned to

pentatricopeptide repeat-containing proteins (PPR proteins), with most of them being upregulated

(86/101) and part of the PLS-class PPR proteins (67/101). Land plant genomes harbour an expanded

number of PPR protein encoding genes, specifically PLS (Barkan & Small, 2014), and many of them are

involved in the RNA stabilisation, splicing and editing in chloroplasts and mitochondria, necessary for the

correct development and functioning of the target organelle (Barkan & Small, 2014; X. Wang et al.,

2021). These data thus suggest a pivotal role for nocturnal manipulation of mitochondria and plastids by

P. brassicae.

On the one hand, the upregulation of genes potentially involved in organelle dynamics in infected *A.*

thaliana would fit the well-documented phenomenon of accumulation of amyloplasts in the root cortex

of plants affected by clubroot (Garvetto et al., 2023; Ma et al., 2022). Disruption of transcription is

known to inhibit the differentiation of plastids into amyloplasts (Enami et al., 2011; Liebers et al., 2017),

suggesting a potential involvement of post-transcriptional editing in the transition. Furthermore,

cytokinin concentration, another factor necessary for the transition to amyloplast (Enami et al., 2011), is

increased locally by *P. brassicae* (Müller & Hilgenberg, 1986; Siemens et al., 2006) arguing for its

importance in clubroot development. Amyloplast induction driven by alterations in the cytokinins-to-

auxins ratio would enable the storage of sugars directed toward the rapidly growing root in the night

(Graf et al., 2010; Yazdanbakhsh et al., 2011; Yazdanbakhsh & Fisahn, 2010), where the parasite

generates an even stronger physiological sink (Keen & Williams, 1969; Malinowski et al., 2019;

Walerowski et al., 2018). Nevertheless, on the other hand, plastidial RNA editing has been found to be

more effective in green tissues as compared to non-photosynthetic ones (Peeters & Hanson, 2002) and a

high proportion of PPR in our data seem to target the mitochondrion. Therefore, a hypothesis

encompassing organelle-mediated defences might be more parsimonious.

RNA editing is required for the correct functioning of chloroplasts and mitochondria and mutations of

PPR proteins often lead to defective growth or lethal phenotypes, in particular by stimulating the

production of reactive oxygen species (ROS) due to defects in the post-transcriptional modifications of

genes targeting these two organelles (Y. Wang & Tan, 2025). Evidence exists of plants exploiting

increased ROS levels caused by the disruption of the electron transport chain in chloroplasts and

mitochondria as a signal to promote defence or to initiate oxidative burst and fend off pathogens. In *A.*

thaliana chloroplasts, reduced editing of *ndhB* mRNA (encoding the B subunit of the chloroplast NADH

dehydrogenase-like complex - NDH) caused by the disruption of PPR proteins CRR21 and CRR2 (both

upregulated in our dataset) led to substantially enhanced ROS production and subsequent resistance to

fungal pathogens (García-Andrade et al., 2013). A recent study found that lack of editing and splicing of

mitochondria-targeting mRNAs *nad1*, *nad3*, *nad6* and *nad7* (encoding subunits of mitochondrial Complex

I) caused by downregulation of the PPR protein RESISTANCE TO PHYTOPHTHORA PARASITICA 7 (RTP7)

results in ROS overproduction, oxidative burst and incompatible interaction with the pathogenic

oomycete *Phytophthora parasitica*(Yang et al., 2022) Although not a PPR, multiple organellar RNA-

editing factor 8 (MORF8) participates in RNA editing in both chloroplasts and mitochondria, and has been

shown to decrease resistance to *P. parasitica* when targeting the mitochondrion, putatively by stabilizing

the respiratory chain thereby avoiding ROS production, SA signalling and oxidative burst (Yang et al.,

2020). This is particularly interesting as *morf8* is among the genes assigned to "RNA modification"

upregulated in the night in our analysis. Therefore, the increased RNA editing observed in our dataset

can be interpreted as an attempt from the side of *P. brassicae* to maintain functional mitochondria and

plastids, subsequently dampening ROS production and defence. The hormone salicylic acid (SA)

stimulates mitochondrial ROS production (J. Wang et al., 2022) and is involved in a positive feedback

loop (Mencia et al., 2020), suppressing mitochondrial RNA editing and being induced by the ensuing ROS

production as observed in (Yang et al., 2020). SA is also under strong diel control and peaks in the middle

of the night (Zheng et al., 2015; Lu et al., 2017; Karapetyan & Dong, 2018), therefore disruptions in SA

signalling by *P. brassicae* in the moment of its normal peak concentration might help to explain the

enhanced nocturnal upregulation of "RNA modification" in our analyses.

Indeed, SA-mediated defences are known to be affected in the interaction between *P. brassicae* and its

host (Galindo-González et al., 2020; Mencia et al., 2022), as exemplified by the parasite's effector protein

PbBSMT methylating SA to methyl-salicylate (MeSA) in order to enhance its transport away from the site

of infection, thereby reducing its local concentration and dampening the immune response (Djavaheri et

al., 2019; Ludwig-Müller et al., 2015). Accordingly, an overall downregulation of the category "response

to salicylic acid" was observed in our analyses in the intermediate and mature infection regardless of the

time of the day (Fig. 2 and 3; Additional File 10).

A putative targeting of nocturnal SA signalling by *P. brassicae* might also explain why the biosynthesis of

phenylpropanoids, enhanced by SA (J. Y. Chen et al., 2006), is more severely impacted in the night (64

genes enriching the category "phenylpropanoid metabolic process" in the night as compared to 49 in the

day; Additional File 10; Bauters et al., 2021). Phenylpropanoids are known to be involved in plant

mechanical and chemical defence, especially via the induction of lignin deposition in secondary cell walls

and the accumulation of flavonoids (Dong & Lin, 2021). The pivotal importance of lignification of

secondary cell walls in preventing the colonization and spread of *P. brassicae* is known, with the parasite

actively interfering in this process in susceptible *B. napus* varieties (Tu et al., 2024) and downregulation

of lignin biosynthesis has been observed in clubroot infected *A. thaliana* (Agarwal et al., 2011). Our data

suggest that the manipulation of the nocturnally gated SA response by *P. brassicae* is a key driver of the

changes observed in defence and organellar dynamics, arguing in favour of the exploration of nocturnal

phases of clubroot to better understand this disease. Furthermore, the recently established link between

SA acid and circadian clock(Fraser et al., 2024) advocates for the identification of genes whose

oscillations are equally affected by temporal and disease dynamics.

The IE model allowed for the detection of this small and interesting subset of DEGs whose expression

pattern is simultaneously affected by time of the day and infection (Duda et al., 2023). The overall

reduction of DEGs at 21 DAI as compared to 14 DAI (i.e., 96 against 217) is likely due to the already

observed decreased effect of diel oscillations in older plants and explains the absence of day/night

grouping of the samples at 21 DAI (Fig. 5). At 14 DAI, when the effect of time on gene expression is

stronger, DEGs appropriately divide samples into day/night and infected/control groups (Fig. 4). At 14

DAI, row clustering reveals varying effects of infection on the diel gene expression patterns, the most

notable being the suppression of rhythmicity through upregulation of genes that are normally

downregulated at night (i.e., cluster 1 in Fig. 4). Interestingly, the GO term enrichment analysis of the

DEGs in cluster 1 features the terms "Circadian rhythm" and "Rhythmic process" (Additional File 8). A

closer look at the list of genes contained in cluster 1 revealed seven genes of acknowledged importance

in rhythmic and circadian processes: *GIGANTEA* (GI), *PSEUDO-RESPONSE REGULATOR* 3 (PRR3), 7 (PRR7)

and 9 (PRR9); *PHYTOCHROME INTERACTING FACTOR 4* (PIF4), *NIGHT LIGHT-INDUCIBLE AND CLOCK-*

REGULATED 1 (LNK1) and 4 (LNK4). Members of the core circadian clock such as GI and the PRRs are

known to be directly involved in the transcriptional/translational feedback loop central to circadian

oscillations (Sanchez & Kay, 2016), but also to integrate endogenous and exogenous stimuli to inform

and entrain it (S. Wang et al., 2022), a role they share with PIF4 and LNK1/4 (Kidokoro et al., 2023; Zhu et

al., 2016). Whilst LNK1 and LNK4 seems to mainly integrate temperature stimuli in the circadian

oscillator (Kidokoro et al., 2023), all other genes detected above are variously involved in the integration

of information about temperature, light, sugar and hormone levels within the circadian clock (S. Wang et

al., 2022). While temperature and light are unlikely to be involved in mediating the pathogen influence

on the diel rhythm of the host, *P. brassicae* influence on hormones (Jayasinghege et al., 2023; Wei et al.,

2021) and sugar levels (Keen & Williams, 1969; Ma et al., 2022) is documented and a more plausible

mediator. Considering the suppression of the diel cycle observed in cluster 1 and the fact that the

majority of the rhythmic genes we identified within it are inhibitors of the central oscillator CIRCADIAN

CLOCK ASSOCIATED 1 / LATE ELONGATED HYPOCOTYL (CCA1/LHY), either directly such as PRR9, PRR7

(Farré & Liu, 2013) or indirectly such as PRR3 (via TOC1; Para et al., 2007) and LNK1 (via RVE8 and PRR5;

Kidokoro et al., 2023), it is possible that these genes are mediating the influence of the pathogen on the

circadian clock.

In the shoot PRR7 is activated in condition of low sugars by a mechanism involving the activation of the

sugar sensor SUCROSE NON FERMENTING RELATED KINASE 1 (SnRK1), itself phosphorylating the

transcription factor bZIP63 which activates PRR7 transcription (Frank et al., 2018). Interestingly, SnRK1

has been found to be the direct target of the recently described *P. brassicae*-specific effector protein

PBZF1 (W. Chen et al., 2021). SnRK1 is a central sensor of nutrient stress in plants (Han et al., 2024), and

the fact that this gene seems to be at the center of direct (i.e., via effectors) and indirect (i.e., via

changes in sugar levels) manipulation by *P. brassicae* argues for its pivotal role in the interaction. Recent

studies have shown that sucrose has an even higher importance in conveying circadian information in

the root, where sucrose pulses from the shoot regulate diel oscillations of PRR7 (and PRR9; Uemoto et

al., 2023). Unlike sucrose and trehalose-6-phosphate levels, plants do not regulate starch turnover based

on its abundance (Seki et al., 2017). As a result, the parasites direct consumption of sucrose and/or the

induction of amyloplastic storage could create a continuous flow of photosynthates towards the

infection-induced sink, disrupting the normal rhythmic pattern of sugar delivery to the root. This

scenario of a nutrient deprived tissue with consistent expression of PRR7 is further supported in our

study by the high expression of the "starvation gene" SKIP20 (a.k.a. KMD4 or At3g59940; Graf et al.,

2010) in infected roots.

Despite showing expression patterns like the one discussed above, GI and PIF4 are less clearly influenced

by sugar metabolism. Sucrose pulses in sugar-depleted condition in the dark are known to be able to re-

establish the oscillations of the circadian clock by post-transcriptionally stabilizing GI but they do not

change its expression levels (Dalchau et al., 2011; Haydon et al., 2017); whilst sucrose influences the

expression pattern of various PIFs, but not of PIF4 (Shor et al., 2017). GI is a plant specific circadian

protein promoting the transcription of thousands of genes and whose competitive interaction with PIF4

integrates endogenous oscillations with light and temperature in order to optimize growth (Nohales et

al., 2019; Park et al., 2020). Among the many processes influenced by GI, senescence is particularly

interesting, since it is also found in our GO term enrichment analysis of cluster 1 ("Plant organ

senescence"; "Leaf senescence"; Additional File 8) and prior studies have shown delayed senescence

when *GI* expression is suppressed (Thiruvengadam et al., 2015).

In our dataset, the established connection between circadian rhythms and senescence processes may be

linked to the continuous expression of *GI* during the night, explaining the premature senescence and the

yellowing of leaves observed as typical clubroot symptoms. GI nocturnal breakdown also regulates

gibberellin mediated growth by allowing DELLA degradation and enabling the expression of downstream

growth promoting genes (Nohales & Kay, 2019). The upregulation of *GI* in the night in our dataset might

therefore be involved in disrupting the root growth (especially lateral root formation and main root

elongation) by making cells insensitive to gibberellin (Shtin et al., 2022). Intriguingly, GI is also involved in

stabilization of REPRESSOR OF GA (RGA; Park et al., 2020), itself inhibiting the gibberellin-induced

transcription of PIF4 (another promoter of growth genes), which is instead upregulated in the night in

infected plants. These results highlight the complexity of the influence of the circadian oscillator on

downstream processes acting at the post-transcriptional level.

Conclusions

Overall, our exploration highlights a previously unrecognized nocturnal impact of *P. brassicae* on

organellar dynamics and plant defences, and suggests a disruption of the host circadian processes. We

provide target genes amenable to experimental validation, the necessary next step in order to answer

questions on the time-specific cross-talk between *P. brassicae* and its host. Nocturnal manipulation of

host organellar RNA editing by the parasite is likely to be linked to the diel oscillations in SA signalling

and seemingly mediated by a subset of transcripts enriched in PPR encoding genes. The question

remains whether PRR genes are a direct target of *P. brassicae* manipulation or whether the influence of

the pathogen on their expression pattern is mediated by manipulation of SA signalling. Ongoing analyses

of the diel changes in gene expression on the side of the parasite might help identifying time-specific

effectors and help clarifying their targets. Furthermore, our investigation already revealed that circadian

oscillations in gene expression (i.e., oscillations at a finer scale than the day and night) are altered by *P.*

brassicae. In line with recent findings (Fu et al., 2025), the application of specific tools for the analysis of

differential rhythmicity to our dataset may further identify effectors allowing *P. brassicae* to hijack the

circadian clock components to disrupt the physiology of the host plant.

Experimental procedures

Experimental setup & biological material

After germination, seedlings of *Arabidopsis thaliana* ecotype Col-0 obtained from the Seed Stock Centre

(stock N1092, Nottingham, UK) were grown on a soil substrate (Einheitserde Type P, pH 5.8; Hermina

Samen, Germany) mixed with sand (Sahara Spielsand; Hornbach Germany) in the ratio 4:1. This soil

mixture was steam-sterilized prior to use for 120 minutes at 100 °C. All plants were grown in a Sanyo

MLR-350H climate chamber at a temperature of 23°C during the day and 18°C during the night. The

light/dark cycle was set to 12:12 hours, with a light intensity of 100 μmol. Genome-sequenced

Plasmodiophora brassicae isolate e3 (Schwelm et al., 2015) obtained from Dr. Siemens (Fähling et al.,

2004) was propagated on *B. rapa* var. *pekinensis* (ISP International Seed Processing GmbH, Quedlinburg,

Germany) and pure spore suspensions were obtained as described in (Smolko et al., 2024). One week

after germination, *A. thaliana* Col-0 plants were individually inoculated with 1 mL of *P. brassicae* isolate

e3 spore suspension at a concentration of 10^6 spores/mL in 50 mM KH_2PO_4 buffer (pH 5.5, adjusted with

K_2HPO_4) or with a mock inoculation of an equal amount of the same buffer. Healthy controls and infected

plants were harvested 14 and 21 days after inoculation (DAI, Additional File 1 A and B), in each occasion

at four different time points: two points during the "day" (5 hr 50' and 10 hr 40' after artificial dawn) and

two points during the "night" (15 hr 30' and 19 hr 20'; i.e., 3 hr 30' and 7 hr 20' after artificial dusk). The

time point 14 DAI was chosen as it marks the establishment of the secondary infection phase with

colonization of cortical tissue by metabolically active plasmodia and onset of visible symptoms in the

root (Liu et al., 2020). The second time point (i.e., 21 DAI) has been chosen as it matches the shift from

actively feeding plasmodia to resting spore production (Liu et al., 2020). Eight to twelve plants were

grown per each time point / treatment and harvested by pooling four plant roots in 1 mL ice-cold

RNAlater (Ambion, Austin, TX, USA) in 1.5 mL Eppendorf tubes. Samples were then left to rest overnight,

before removing and refreshing the RNAlater and storing them at -80°C until RNA extraction.

RNA extraction & sequencing

Samples in RNAlater were thawed on ice and one or two plant roots (when biomass from a single root

specimen was judged insufficient for RNA extraction) per each tube were transferred to ethanol-

sterilised and RNAseAway treated tin foil squares with equally treated steel tweezers (Additional File 1 C

and D). A root sample as described above constitute one replicate, and triplicate were selected and

processed per each time point, for infected and mock inoculated control plants at 14 DAI and 21 DAI,

resulting in a total of 48 processed roots (Additional File 2). Roots were carefully wrapped and snap-

frozen in liquid nitrogen before transferring them in pre-mixed bead matrix tubes (D1034-MX; Biozym,

Hessisch Oldendorf, Germany) amended with 200 μL IB buffer (SPLIT RNA Extraction Kit, Lexogen,

Vienna, Austria) to decrease the risk of RNA degradation and immediately homogenized with a FastPrep

bead mill (MP Biomedicals, Santa Ana, CA, USA) for 40 s at 60 ms^{-1} three times. After tissue disruption,

another 200 μL of ice cold IB buffer was added to the samples. The RNA extraction proceeded following

the protocol SPLIT RNA extraction of total RNA (SPLIT RNA Extraction Kit, Lexogen, Vienna, Austria) as

suggested by the manufacturer without further modifications. Assessment of the RNA extracts quality

and quantity was performed on an Agilent Bioanalyzer 2100 (Agilent Technologies, Palo Alto, CA, USA)

and Quantus fluorometer (Promega Corporation, Fitchburg, WI, USA). Strand-specific library preparation

was performed on poly-A-selected mRNA using a NEBNext Ultra II Directional RNA Library Prep Kit with

NEBNext Poly(A) mRNA Magnetic Isolation Module and NEBNext Multiplex Oligos for Illumina (New

England Biolabs, Ipswich, MA, USA). The quality and concentration of cDNA in the resulting libraries was

assessed on an Agilent Bioanalyzer 2100 and Quantus fluorometer, before paired-end sequencing (2x150

bp on a NovaSeq S1; Illumina, San Diego, CA, USA) at the VBCF NGS Unit (Vienna, Austria).

Sequencing data processing

Read quality was checked using FastQC (Andrews, 2010) and quality reports visualized in MultiQC v1.6

(Ewels et al., 2016) before and after read filtering and adapter trimming with Trim Galore v0.6.2 (quality

score threshold 30; https://zenodo.org/doi/10.5281/zenodo.5127898). To assign reads to the host plant

and the parasite, Bowtie2 v2.3.5.1 (Langmead & Salzberg, 2012) was used to map them against the

concatenated genomes of *Arabidopsis thaliana* (TAIR10; Lamesch et al., 2011) and *Plasmodiophora*

brassicae (Pldbra_eH_r1) with a "Place-to-go-based" approach as defined in (O'Keeffe & Jones, 2019).

Obtained read pools (separately for the host and the parasite) were assembled with Trinity v2.10.0 (in

genome guided mode; Grabherr et al., 2011)) and resulting transcripts annotated with Trinotate v3.2.1

(Bryant et al., 2017). Expression estimates were obtained by pre-aligning the reads to the transcripts

using STAR v2.7.1a (Dobin et al., 2013) and assigned to assembled transcript with Salmon (Patro et al.,

2017). Raw NGS sequences of the dataset obtained in this study are deposited in NCBI Sequence Read

Archive (SRA) under the BioProject accession number PRJNA1212907.

Differential gene expression

DESeq2 Bioconductor package (Love et al., 2014) in R (R Core Team, 2021) was used to estimate

differential gene expression. In order to detect differences in the response of plants to infection by *P.*

brassicae separately during the day and during the night, differentially expressed genes (DEGs) were

calculated on pooled counts. Briefly, read counts from all time points sampled during the day (i.e., 5 hr

50' and 10 hr 40' after artificial dawn) and during the night (15 hr 30' and 19 hr 20' after artificial dawn)

were pooled before further analyses.

This resulted in six replicates for each of the four conditions, as defined by the two factors "Condition"

(Infected or Control) and "Time point" (Day or Night; Additional file 2). An experimental design with two

factors and interaction was used for estimating the DEGs separately for 14 DAI and 21 DAI, thus resulting

in five comparisons: infected vs control in the day, infected vs control in the night, control in the night vs

in the day, infected in the night vs in the day and interaction. Genes were considered differentially

expressed when the adjusted p-value was ≤ 0.05 (Wald test p-value adjusted for multiple testing with

Benjamini–Hochberg correction) and a $\log_2$-fold change was ≤ –1 or ≥ 1.

Principal component analysis and Venn diagram

Principal component analysis (PCA) was calculated in the DESeq2 Bioconductor package in R (Love et al.,

2014) and further modified in ggplot2 (Wickham, 2010). The PCA was performed on variance stabilizing

transformed (vst) count data from the DESeqDataSet object, without taking into account the sample

information (i.e., blind=TRUE), as suggested by the DESeq2 manual. Venn diagrams on DEGs were

calculated in R with the Package VennDiagram (H. Chen, 2022).

GO term over representation analyses

ClusterProfiler (Wu et al., 2021) was used within R to perform gene ontology (GO) over-representation

tests using lists of DEGs in the comparisons between infected and healthy plants in the day and in the

night. Significance of the enriched GO categories was assessed by Fisher's exact test controlled for False

discovery rate (FDR) with Benjamini–Hochberg correction and the q-value cutoff was set to 0.05.

Because the RNA-seq strategy adopted was designed to measure all expressed mRNAs (total RNA

extraction followed by poly-A tail selection), the background gene list (or "universe") against which over-

representation was performed, was composed of a list of all protein coding genes from *A. thaliana*

TAIR10 genome release. The package GOplot (Walter et al., 2015) was used to combine expression data

with functional categories. Using the function "circle_dat", the package was used to summarise and

append to each enriched GO category an overall expression value (i.e., a z-score) calculated as follows:

$$z\ score = \frac{(n° \text{ upregulated genes in GO } - n° \text{ downregulated genes in GO })}{\sqrt{Total\ n° \ of\ genes\ in\ GO}}$$

The function "reduce_overlap" within the same package was used to eliminate all but one GO category

sharing more than 25% of the genes before plotting the results with the function "GObubble".

Heatmap of gene expression levels

Genes whose differential expression levels were driven by the interaction between the factor "Time

point" and the factor "Condition" were detected and their expression levels were visualised with

heatmaps as implemented in the package pheatmap (Kolde, 2019) in R, on row-scaled, log-normalised

read counts and with default "Euclidean" clustering method. All figures were further refined and

finalized in Inkscape (Inkscape Project, 2020).

Author contributions

523   Experimental concept and design: SN, JLM. Wet lab work: SA, FB, AG, MH. Bioinformatics and statistical

524   analysis: AG. Analysis of Results: AG, SN, SA, JLM, MH. Manuscript writing: AG, SN, SA, JLM. Figures and

525   tables: AG. All authors read and approved the final manuscript.

Acknowledgements

527   We are grateful to Bettina Schneidhofer for her support in the wet lab. We would also like to thank the

528   VBCF NGS Unit (www.vbcf.ac.at) in particular Carmen Czepe for her support during the sequencing and

529   Nadine Tatto for her paramount contribution and continuous support with the bioinformatic analyses.

Conflict of interests

The authors declare that they have no competing interests. This article does not contain any studies with

human or animal participants

Data availability statement

Raw NGS sequences of the dataset generated and analysed in the current study are available in NCBI

Sequence Read Archive (SRA) under the BioProject accession number PRJNA1212907 or are available

from the corresponding author on request.

Funding

This research was funded in whole by the Austrian Science Fund (FWF) [grant DOI:10.55776/Y801]. For

open access purposes, the authors have applied a CC BY public copyright license to any author accepted

manuscript version arising from this submission.

Supporting Information

Additional file 1. Above and below ground phenotypical traits of infected and control plants at 14 and 21 DAI.

Additional file 2. Detailed overview of the samples sequenced in this study.

Additional file 3. PCA on control *A. thaliana* plants, showing the decreased influence of the temporal factor with plant age.

Additional file 4. Volcano plot showing the number of DEGs in the day and night in infected *A. thaliana* plants 14 DAI.

Additional file 5. Volcano plot showing the number of DEGs in the day and night in infected *A. thaliana* plants 21 DAI.

Additional file 6. GO categories enriched in DEGs in *A. thaliana* roots 14 DAI infected by *P. brassicae.*

Additional file 7. GO categories enriched in DEGs in *A. thaliana* roots 21 DAI infected by *P. brassicae.*

Additional file 8. GO terms overrepresentation analysis of clusters of DEGs driven by the interaction between infection and time of the day 14 DAI.

Additional file 9. GO terms overrepresentation analysis of clusters of DEGs driven by the interaction between infection and time of the day 21 DAI.

557 **Additional file 10.** Enriched GO terms and corresponding genes underpinning figure 2 and 3 before data
558 reduction.

561 *Fig. 1 Day/night-informed overview of gene expression in samples of A. thaliana roots 14 DAI (A, B) and 21 DAI (C, D) with P.*
562 *brassicae. A and C, PCA of A. thaliana root samples structured by gene expression estimates (i.e., vst counts). B and D,*
563 *Proportions of DE genes between infected and healthy A. thaliana roots assigned to the day, night or both.*

Fig. 2 GO bubble chart of GO categories enriched in DEGs in infected A. thaliana roots during the day (A, yellow) and night (B, blue) 14 DAI by P. brassicae. The bubbles are arranged according to their overall expression level or z-score on the x axis and according to the negative logarithmic scale of the adjusted p-value on the y axis. The bubble size is proportional to the gene counts within each category. The significance threshold of adjusted p-value < 0,05 (1,3 in -log10 scale) is represented by coloured lines parallel to the x axis (orange in the day and turquoise in the night).

Fig. 3 GO bubble chart of GO categories enriched in DEGs in infected A. thaliana roots during the day (top, yellow) and night (blue, bottom) 21 DAI by P. brassicae. The bubbles are arranged according to their overall expression level or z-score on the x axis and according to the negative logarithmic scale of the adjusted p-value on the y axis. The bubble size is proportional to the gene counts within each category. The significance threshold of adjusted p-value < 0,05 (1,3 in -log10 scale) is represented by coloured lines parallel to the x axis (orange in the day and turquoise in the night).

586 *Fig. 4 Heatmap highlighting DEGs explained by the interaction between infection and time of the day as detected in infected A*
587 *.thaliana roots 14 DAI The row-normalised and log transformed gene counts are shown for each sample separately. The status of*
588 *each sample is described by a combination of green (control plant) or purple (infected plant) and yellow (day) or (blue) bars.*
589 *DEGs with similar patterns of expression have been grouped in 8 clusters (numbers in white boxes).*

Fig. 5: Heatmap highlighting DEGs explained by the interaction between infection and time of the day as detected in infected A .thaliana roots 21 DAI The row-normalised and log transformed gene counts are shown for each sample separately. The status of each sample is described by a combination of green (control plant) or purple (infected plant) and yellow (day) or (blue) bars. DEGs with similar patterns of expression have been grouped in 5 clusters (numbers in white boxes).

591
592
593
594

References

Agarwal, A., Kaul, V., Faggian, R., Rookes, J. E., Ludwig-Müller, J., & Cahill, D. M. (2011). Analysis of global host gene expression during the primary phase of the Arabidopsis thalianaPlasmodiophora brassicae interaction. *Functional Plant Biology*, *38*(6), 462–478. https://doi.org/10.1071/FP11026

Andrews, S. (2010). *FastQC A Quality Control tool for High Throughput Sequence Data*. Http://Www.Bioinformatics.Babraham.Ac.Uk/Projects/Fastqc/.

Barkan, A., & Small, I. (2014). Pentatricopeptide repeat proteins in plants. *Annual Review of Plant Biology*, *65*, 415–442. https://doi.org/10.1146/annurev-arplant-050213-040159

Bauters, L., Stojilković, B., & Gheysen, G. (2021). Pathogens pulling the strings: Effectors manipulating salicylic acid and phenylpropanoid biosynthesis in plants. In *Molecular Plant Pathology* (Vol. 22, Issue 11, pp. 1436–1448). John Wiley and Sons Inc. https://doi.org/10.1111/mpp.13123

Bryant, D. M., Johnson, K., DiTommaso, T., Tickle, T., Couger, M. B., Payzin-Dogru, D., Lee, T. J., Leigh, N. D., Kuo, T. H., Davis, F. G., Bateman, J., Bryant, S., Guzikowski, A. R., Tsai, S. L., Coyne, S., Ye, W. W., Freeman, R. M., Peshkin, L., Tabin, C. J., … Whited, J. L. (2017). A Tissue-Mapped Axolotl De Novo Transcriptome Enables Identification of Limb Regeneration Factors. *Cell Reports*, *18*(3), 762–776. https://doi.org/10.1016/j.celrep.2016.12.063

Bustillo-Avendaño, E., Serrano-Ron, L., & Moreno-Risueno, M. A. (2022). The Root Clock as a Signal Integrator System: Ensuring Balance for Survival. *Frontiers in Plant Science*, *13*(May). https://doi.org/10.3389/fpls.2022.886700

Chen, H. (2022). *VennDiagram: Generate High-Resolution Venn and Euler Plots*.

Chen, J. Y., Wen, P. F., Kong, W. F., Pan, Q. H., Zhan, J. C., Li, J. M., Wan, S. B., & Huang, W. D. (2006). Effect of salicylic acid on phenylpropanoids and phenylalanine ammonia-lyase in harvested grape berries. *Postharvest Biology and Technology*, *40*(1), 64–72. https://doi.org/10.1016/j.postharvbio.2005.12.017

Chen, W., Li, Y., Yan, R., Ren, L., Liu, F., Zeng, L., Sun, S., Yang, H., Chen, K., Xu, L., Liu, L., Fang, X., & Liu, S. (2021). SnRK1.1-mediated resistance of Arabidopsis thaliana to clubroot disease is inhibited by the novel Plasmodiophora brassicae effector PBZF1. *Molecular Plant Pathology*, *22*(9), 1057–1069. https://doi.org/10.1111/mpp.13095

Ciaghi, S., Schwelm, A., & Neuhauser, S. (2019). Transcriptomic response in symptomless roots of clubroot infected kohlrabi (Brassica oleracea var. gongylodes) mirrors resistant plants. *BMC Plant Biology*, *19*(1), 1–14. https://doi.org/10.1186/s12870-019-1902-z

Dalchau, N., Baek, S. J., Briggs, H. M., Robertson, F. C., Dodd, A. N., Gardner, M. J., Stancombe, M. A., Haydon, M. J., Stan, G. B., Gonçalves, J. M., & Webb, A. A. R. (2011). The circadian oscillator gene GIGANTEA mediates a long-term response of the Arabidopsis thaliana circadian clock to sucrose. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(12), 5104–5109. https://doi.org/10.1073/pnas.1015452108

Davis, W., Endo, M., & Locke, J. C. W. (2022). Spatially specific mechanisms and functions of the plant
circadian clock. *Plant Physiology*, *190*(2), 938–951. https://doi.org/10.1093/plphys/kiac236

Dekhuijzen, H. M. (1981). The Occurrence of Free and Bound Cytokinins in Plasmodia of Plasmodiophora
brassicae Isolated from Tissue Cultures of Clubroots. *Plant Cell Reports*, *213*, 29–30.

Dekhuijzen, H. M., & Overeem, J. C. (1971). The role of cytokinins in clubroot formation. *Physiological*
Plant Pathology, *1*(2), 151–161. https://doi.org/10.1016/0048-4059(71)90024-5

Devos, S., Laukens, K., Deckers, P., Van Der Straeten, D., Beeckman, T., Inzé, D., Van Onckelen, H.,
Witters, E., & Prinsen, E. (2006). A hormone and proteome approach to picturing the initial
metabolic events during Plasmodiophora brassicae infection on Arabidopsis. *Molecular Plant-*
Microbe Interactions, *19*(12), 1431–1443. https://doi.org/10.1094/MPMI-19-1431

Devos, S., Vissenberg, K., Verbelen, J. P., & Prinsen, E. (2005). Infection of Chinese cabbage by
Plasmodiophora brassicae leads to a stimulation of plant growth: Impacts on cell wall metabolism
and hormone balance. *New Phytologist*, *166*(1), 241–250. https://doi.org/10.1111/j.1469-
8137.2004.01304.x

Dixon, G. R. (2009). The occurrence and economic impact of plasmodiophora brassicae and clubroot
disease. *Journal of Plant Growth Regulation*, *28*(3), 194–202. https://doi.org/10.1007/s00344-009-
9090-y

Djavaheri, M., Ma, L., Klessig, D. F., Mithöfer, A., Gropp, G., & Borhan, H. (2019). Mimicking the host
regulation of salicylic acid: A virulence strategy by the clubroot pathogen plasmodiophora
brassicae. *Molecular Plant-Microbe Interactions*, *32*(3), 296–305. https://doi.org/10.1094/MPMI-
07-18-0192-R

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras,
T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21.
https://doi.org/10.1093/bioinformatics/bts635

Dong, N. Q., & Lin, H. X. (2021). Contribution of phenylpropanoid metabolism to plant development and
plant–environment interactions. *Journal of Integrative Plant Biology*, *63*(1), 180–209.
https://doi.org/10.1111/jipb.13054

Duda, J. C., Drenda, C., Kästel, H., Rahnenführer, J., & Kappenberg, F. (2023). Benefit of using interaction
effects for the analysis of high-dimensional time-response or dose-response data for two-group
comparisons. *Scientific Reports*, *13*(1), 1–12. https://doi.org/10.1038/s41598-023-47057-0

Durand, M., Mainson, D., Porcheron, B., Maurousset, L., Lemoine, R., & Pourtau, N. (2018). Carbon
source–sink relationship in Arabidopsis thaliana: the role of sucrose transporters. *Planta*, *247*(3),
587–611. https://doi.org/10.1007/s00425-017-2807-4

Ehrary, A., Rosas, M., Carpinelli, S., Davalos, O., Cowling, C., Fernandez, F., & Escobar, M. (2020).
Glutaredoxin AtGRXS8 represses transcriptional and developmental responses to nitrate in
Arabidopsis thaliana roots. *Plant Direct*, *4*(6). https://doi.org/10.1002/pld3.227

Enami, K., Ozawa, T., Motohashi, N., Nakamura, M., Tanaka, K., & Hanaoka, M. (2011). Plastid-to-nucleus
retrograde signals are essential for the expression of nuclear starch biosynthesis genes during

amyloplast differentiation in tobacco BY-2 cultured cells. *Plant Physiology*, *157*(1), 518–530.
https://doi.org/10.1104/pp.111.178897

Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–3048.
https://doi.org/10.1093/bioinformatics/btw354

Fähling, M., Graf, H., & Siemens, J. (2004). Characterization of a Single-Spore Isolate Population of
Plasmodiophora brassicae Resulting from a Single Club. *Journal of Phytopathology*, *152*, 438–444.

Farré, E. M., & Liu, T. (2013). The PRR family of transcriptional regulators reflects the complexity and
evolution of plant circadian clocks. *Current Opinion in Plant Biology*, *16*(5), 621–629.
https://doi.org/10.1016/j.pbi.2013.06.015

Frank, A., Matiolli, C. C., Viana, A. J. C., Hearn, T. J., Kusakina, J., Belbin, F. E., Wells Newman, D.,
Yochikawa, A., Cano-Ramirez, D. L., Chembath, A., Cragg-Barber, K., Haydon, M. J., Hotta, C. T.,
Vincentz, M., Webb, A. A. R., & Dodd, A. N. (2018). Circadian Entrainment in Arabidopsis by the
Sugar-Responsive Transcription Factor bZIP63. *Current Biology*, *28*(16), 2597-2606.e6.
https://doi.org/10.1016/j.cub.2018.05.092

Fraser, O. J. P., Cargill, S. J., Spoel, S. H., & van Ooijen, G. (2024). Crosstalk between salicylic acid
signalling and the circadian clock promotes an effective immune response in plants. *Npj Biological*
Timing and Sleep, *1*(1). https://doi.org/10.1038/s44323-024-00006-0

Fu, M., Zhou, Y., Zhang, X., Yang, K., Xu, Y., Wang, X., Chen, Z., Wang, Y., Shi, Y., Ma, L., Liu, H., Deng, Y.,
Cheng, S., Chu, J., Song, J., Sun, T., Wang, Y., Wang, W., & Zhou, M. (2025). A pathogen effector
HaRxL10 hijacks the circadian clock component CHE to perturb both plant development and
immunity. *Nature Communications* , *16*(1). https://doi.org/10.1038/s41467-025-56787-w

Galindo-González, L., Manolii, V., Hwang, S. F., & Strelkov, S. E. (2020). Response of Brassica napus to
Plasmodiophora brassicae Involves Salicylic Acid-Mediated Immunity: An RNA-Seq-Based Study.
Frontiers in Plant Science, *11*(July), 1–20. https://doi.org/10.3389/fpls.2020.01025

García-Andrade, J., Ramírez, V., López, A., & Vera, P. (2013). Mediated Plastid RNA Editing in Plant
Immunity. *PLoS Pathogens*, *9*(10). https://doi.org/10.1371/journal.ppat.1003713

Garvetto, A., Murúa, P., Kirchmair, M., Salvenmoser, W., Hittorf, M., Ciaghi, S., Harikrishnan, S. L.,
Gachon, C. M. M., Burns, J. A., & Neuhauser, S. (2023). Phagocytosis underpins the biotrophic
lifestyle of intracellular parasites in the class Phytomyxea (Rhizaria). *New Phytologist*, *238*(5), 2130–
2143. https://doi.org/10.1111/nph.18828

Goodspeed, D., Chehab, E. W., Min-Venditti, A., Braam, J., & Covington, M. F. (2012). Arabidopsis
synchronizes jasmonate-mediated defense with insect circadian behavior. *Proceedings of the*
National Academy of Sciences of the United States of America, *109*(12), 4674–4677.
https://doi.org/10.1073/pnas.1116368109

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L.,
Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F.,
Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev, A. (2011). Full-length transcriptome

assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, *29*(7), 644–652.
https://doi.org/10.1038/nbt.1883

Graf, A., Schlereth, A., Stitt, M., & Smith, A. M. (2010). Circadian control of carbohydrate availability for
growth in Arabidopsis plants at night. *Proceedings of the National Academy of Sciences of the*
United States of America, *107*(20), 9458–9463. https://doi.org/10.1073/pnas.0914299107

Gravot, A., Deleu, C., Wagner, G., Lariagon, C., Lugan, R., Todd, C., Wendehenne, D., Delourme, R.,
Bouchereau, A., & Manzanares-Dauleux, M. J. (2012). Arginase induction represses gall
development during clubroot infection in arabidopsis. *Plant and Cell Physiology*, *53*(5), 901–911.
https://doi.org/10.1093/pcp/pcs037

Gravot, A., Richard, G., Lime, T., Lemarié, S., Jubault, M., Lariagon, C., Lemoine, J., Vicente, J., Robert-
Seilaniantz, A., Holdsworth, M. J., & Manzanares-Dauleux, M. J. (2016). Hypoxia response in
Arabidopsis roots infected by Plasmodiophora brassicae supports the development of clubroot.
BMC Plant Biology, *16*(1), 1–10. https://doi.org/10.1186/s12870-016-0941-y

Grsic-Rausch, S., Kobelt, P., Siemens, J. M., Bischoff, M., & Ludwig-Müller, J. (2000). Expression and
localization of nitrilase during symptom development of the clubroot disease in Arabidopsis. *Plant*
Physiology, *122*(2), 369–378. https://doi.org/10.1104/pp.122.2.369

Han, C., Wang, H., Shi, W., & Bai, M.-Y. (2024). The molecular associations between the SnRK1 complex
and carbon/nitrogen metabolism in plants. *New Crops*, *1*(December 2023), 100008.
https://doi.org/10.1016/j.ncrops.2023.12.003

Haydon, M. J., Mielczarek, O., Frank, A., Román, Á., & Webb, A. A. R. (2017). Sucrose and ethylene
signaling interact to modulate the circadian clock. *Plant Physiology*, *175*(2), 947–958.
https://doi.org/10.1104/pp.17.00592

Henriques, R., Papdi, C., Ahmad, Z., & Bögre, L. (2018). Circadian Regulation of Plant Growth. *Annual*
Plant Reviews Online, *1*, 675–704. https://doi.org/10.1002/9781119312994.apr0655

Hurný, A., Cuesta, C., Cavallari, N., Ötvös, K., Duclercq, J., Dokládal, L., Montesinos, J. C., Gallemí, M.,
Semerádová, H., Rauter, T., Stenzel, I., Persiau, G., Benade, F., Bhalearo, R., Sýkorová, E., Gorzsás,
A., Sechet, J., Mouille, G., Heilmann, I., … Benková, E. (2020). SYNERGISTIC ON AUXIN AND
CYTOKININ 1 positively regulates growth and attenuates soil pathogen resistance. *Nature*
Communications, *11*(1). https://doi.org/10.1038/s41467-020-15895-5

Inkscape Project. (2020). *Inkscape*. Retrieved from https://inkscape.org.

Irani, S., Trost, B., Waldner, M., Nayidu, N., Tu, J., Kusalik, A. J., Todd, C. D., Wei, Y., & Bonham-Smith, P.
C. (2018). Transcriptome analysis of response to Plasmodiophora brassicae infection in the
Arabidopsis shoot and root. *BMC Genomics*, *19*(1), 23. https://doi.org/10.1186/s12864-017-4426-7

Javed, M. A., Schwelm, A., Zamani-Noor, N., Salih, R., Silvestre Vañó, M., Wu, J., González García, M.,
Heick, T. M., Luo, C., Prakash, P., & Pérez-López, E. (2023). The clubroot pathogen Plasmodiophora
brassicae: A profile update. *Molecular Plant Pathology*, *24*(2), 89–106.
https://doi.org/10.1111/mpp.13283

Jayasinghege, C. P. A., Ozga, J. A., Manolii, V. P., Hwang, S. F., & Strelkov, S. E. (2023). Impact of
Susceptibility on Plant Hormonal Composition during Clubroot Disease Development in Canola
(Brassica napus). *Plants*, *12*(16). https://doi.org/10.3390/plants12162899

Kageyama, K., & Asano, T. (2009). Life cycle of plasmodiophora brassicae. *Journal of Plant Growth*
Regulation, *28*(3), 203–211. https://doi.org/10.1007/s00344-009-9101-z

Karapetyan, S., & Dong, X. (2018). Redox and the circadian clock in plant immunity: A balancing act. In
Free Radical Biology and Medicine (Vol. 119, pp. 56–61). Elsevier Inc.
https://doi.org/10.1016/j.freeradbiomed.2017.12.024

Keen, N. T., & Williams, P. H. (1969). Translocation of Sugars into Infected Cabbage Tissues during
Clubroot Development. *Plant Physiology*, *44*(5), 748–754.

Kidokoro, S., Konoura, I., Soma, F., Suzuki, T., Miyakawa, M., Shinozaki, K., & Yamaguchi-Shinozaki, K.
(2023). Clock-regulated coactivators selectively control gene expression in response to different
temperature stress conditions in Arabidopsis. *Proceedings of the National Academy of Sciences*,
120(16), 1–11. https://doi.org/10.1073/pnas

Kim, H., & Hong, S. (2019). Role of the circadian clock in fine-tuning the process of leaf senescence in
plants. *Translational Medicine of Aging*, *3*(2019), 26–30. https://doi.org/10.1016/j.tma.2018.12.001

Kolde, R. (2019). *pheatmap: Pretty Heatmaps*.

Kunkel, L. (1918). Tissue invasion by Plasmodiophora brassicae. *Journal of Agricultural Research*, *12*(543),
543–572.

Lamesch, P., Berardini, T. Z., Li, D., Swarbreck, D., Wilks, C., Sasidharan, R., Muller, R., Dreher, K.,
Alexander, D. L., Garcia-Hernandez, M., Karthikeyan, A. S., Lee, C. H., Nelson, W. D., Ploetz, L.,
Singh, S., Wensel, A., & Huala, E. (2011). The Arabidopsis Information Resource (TAIR): Improved
gene annotation and new tools. *Nucleic Acids Research*, *40*(D1), 1202–1210.
https://doi.org/10.1093/nar/gkr1090

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*, *9*(4),
357–359. https://doi.org/10.1038/nmeth.1923

Lee, J., Kang, M. H., Kim, J. Y., & Lim, P. O. (2021). The Role of Light and Circadian Clock in Regulation of
Leaf Senescence. *Frontiers in Plant Science*, *12*(April). https://doi.org/10.3389/fpls.2021.669170

Lemarié, S., Robert-Seilaniantz, A., Lariagon, C., Lemoine, J., Marnet, N., Jubault, M., Manzanares-
Dauleux, M. J., & Gravot, A. (2015). Both the Jasmonic Acid and the Salicylic Acid Pathways
Contribute to Resistance to the Biotrophic Clubroot Agent Plasmodiophora brassicae in
Arabidopsis. *Plant and Cell Physiology*, *56*(11), 2158–2168. https://doi.org/10.1093/pcp/pcv127

Liebers, M., Grübler, B., Chevalier, F., Lerbs-Mache, S., Merendino, L., Blanvillain, R., & Pfannschmidt, T.
(2017). Regulatory shifts in plastid transcription play a key role in morphological conversions of
plastids during plant development. *Frontiers in Plant Science*, *8*.
https://doi.org/10.3389/fpls.2017.00023

Liu, L., Qin, L., Zhou, Z., Hendriks, W. G. H. M., Liu, S., & Wei, Y. (2020). Refining the life cycle of
Plasmodiophora brassicae. *Phytopathology*, *110*(10), 1704–1712. https://doi.org/10.1094/PHYTO-
02-20-0029-R

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biology*, *15*(12), 1–21. https://doi.org/10.1186/s13059-014-0550-8

Lovelock, D. A., Donald, C. E., Conlan, X. A., & Cahill, D. M. (2013). Salicylic acid suppression of clubroot in
broccoli (Brassicae oleracea var. italica) caused by the obligate biotroph Plasmodiophora brassicae.
Australasian Plant Pathology, *42*(2), 141–153. https://doi.org/10.1007/s13313-012-0167-x

Lovelock, D. A., Šola, I., Marschollek, S., Donald, C. E., Rusak, G., van Pée, K. H., Ludwig-Müller, J., &
Cahill, D. M. (2016). Analysis of salicylic acid-dependent pathways in Arabidopsis thaliana following
infection with Plasmodiophora brassicae and the influence of salicylic acid on disease. *Molecular*
Plant Pathology, *17*(8), 1237–1251. https://doi.org/10.1111/mpp.12361

Lu, H., McClung, C. R., & Zhang, C. (2017). Tick Tock: Circadian Regulation of Plant Innate Immunity.
Annual Review of Phytopathology, *55*, 287–311. https://doi.org/10.1146/annurev-phyto-080516-
035451

Ludwig-Müller, J., Jülke, S., Geiß, K., Richter, F., Mithöfer, A., Šola, I., Rusak, G., Keenan, S., & Bulman, S.
(2015). A novel methyltransferase from the intracellular pathogen Plasmodiophora brassicae
methylates salicylic acid. *Molecular Plant Pathology*, *16*(4), 349–364.
https://doi.org/10.1111/mpp.12185

Ludwig-Müller, J., Prinsen, E., Rolfe, S. A., & Scholes, J. D. (2009). Metabolism and plant hormone action
during clubroot disease. *Journal of Plant Growth Regulation*, *28*(3), 229–244.
https://doi.org/10.1007/s00344-009-9089-4

Ma, Y., Choi, S. R., Wang, Y., Chhapekar, S. S., Zhang, X., Wang, Y., Zhang, X., Zhu, M., Liu, D., Zuo, Z., Yan,
X., Gan, C., Zhao, D., Liang, Y., Pang, W., & Lim, Y. P. (2022). Starch content changes and
metabolism-related gene regulation of Chinese cabbage synergistically induced by Plasmodiophora
brassicae infection. *Horticulture Research*.

Malinowski, R., Novák, O., Borhan, M. H., Spíchal, L., Strnad, M., & Rolfe, S. A. (2016). The role of
cytokinins in clubroot disease. *European Journal of Plant Pathology*, *145*(3), 543–557.
https://doi.org/10.1007/s10658-015-0845-y

Malinowski, R., Truman, W., & Blicharz, S. (2019). Genius architect or clever thief — How
Plasmodiophora brassicae reprograms host development to establish a pathogen-oriented
physiological sink. *Molecular Plant-Microbe Interactions*, *32*(10), 1259–1266.
https://doi.org/10.1094/MPMI-03-19-0069-CR

Meinke, D. W., Cherry, J. M., Dean, C., Rounsley, S. D., & Koornneef, M. (1998). Arabidopsis thaliana: A
model plant for genome analysis. *Science*, *282*(5389).
https://doi.org/10.1126/science.282.5389.662

Mencia, R., Céccoli, G., Fabro, G., Torti, P., Colombatti, F., Ludwig-Müller, J., Alvarez, M. E., & Welchen, E.
(2020). OXR2 increases plant defense against a hemibiotrophic pathogen via the salicylic acid
pathway. *Plant Physiology*, *184*(2), 1112–1127. https://doi.org/10.1104/pp.19.01351

Mencia, R., Welchen, E., Auer, S., & Ludwig-Müller, J. (2022). A Novel Target (Oxidation Resistant 2) in
Arabidopsis thaliana to Reduce Clubroot Disease Symptoms via the Salicylic Acid Pathway without
Growth Penalties. *Horticulturae*, *8*(1). https://doi.org/10.3390/horticulturae8010009

Mora-García, S., de Leone, M. J., & Yanovsky, M. (2017). Time to grow: circadian regulation of growth
and metabolism in photosynthetic organisms. *Current Opinion in Plant Biology*, *35*, 84–90.
https://doi.org/10.1016/j.pbi.2016.11.009

Mortada, M., Xiong, L., & Mas, P. (2024). Dissecting the complexity of local and systemic circadian
communication in plants. *Npj Biological Timing and Sleep*, *1*(1), 1–7.
https://doi.org/10.1038/s44323-024-00003-3

Motta, F. C., McGoff, K., Moseley, R. C., Cho, C. Y., Kelliher, C. M., Smith, L. M., Ortiz, M. S., Leman, A. R.,
Campione, S. A., Devos, N., Chaorattanakawee, S., Uthaimongkol, N., Kuntawunginn, W.,
Thongpiam, C., Thamnurak, C., Arsanok, M., Wojnarski, M., Vanchayangkul, P., Boonyalai, N., …
Haase, S. B. (2023). The parasite intraerythrocytic cycle and human circadian cycle are coupled
during malaria infection. *Proceedings of the National Academy of Sciences of the United States of*
America, *120*(24). https://doi.org/10.1073/pnas.2216522120

Müller, P., & Hilgenberg, W. (1986). Isomers of zeatin and zeatin riboside in clubroot tissue: evidence for
trans-zeatin biosynthesis by Plasmodiophora brassicae. *Physiologia Plantarum*, *66*(2), 245–250.
https://doi.org/10.1111/j.1399-3054.1986.tb02415.x

Neuhaus, K., Grsic-Rausch, S., Sauerteig, S., & Ludwig-Müller, J. (2000). Arabidopsis plants transformed
with nitrilase 1 or 2 in antisense direction are delayed in clubroot development. *Journal of Plant*
Physiology, *156*(5–6), 756–761. https://doi.org/10.1016/S0176-1617(00)80243-6

Nohales, M. A., & Kay, S. A. (2019). GIGANTEA gates gibberellin signaling through stabilization of the
DELLA proteins in Arabidopsis. *Proceedings of the National Academy of Sciences of the United*
States of America, *116*(43), 21893–21899. https://doi.org/10.1073/pnas.1913532116

Nohales, M. A., Liu, W., Duffy, T., Nozue, K., Sawa, M., Pruneda-Paz, J. L., Maloof, J. N., Jacobsen, S. E., &
Kay, S. A. (2019). Multi-level Modulation of Light Signaling by GIGANTEA Regulates Both the Output
and Pace of the Circadian Clock. *Developmental Cell*, *49*(6), 840-851.e8.
https://doi.org/10.1016/j.devcel.2019.04.030

O'Keeffe, K. R., & Jones, C. D. (2019). Challenges and solutions for analysing dual RNA-seq data for non-
model host–pathogen systems. *Methods in Ecology and Evolution*, *10*(3), 401–414.
https://doi.org/10.1111/2041-210X.13135

Olszak, M., Truman, W., Stefanowicz, K., Sliwinska, E., Ito, M., Walerowski, P., Rolfe, S., & Malinowski, R.
(2019). Transcriptional profiling identifies critical steps of cell cycle reprogramming necessary for
Plasmodiophora brassicae-driven gall formation in Arabidopsis. *Plant Journal*, *97*(4), 715–729.
https://doi.org/10.1111/tpj.14156

Para, A., Farré, E. M., Imaizumi, T., Pruneda-Paz, J. L., Harmon, F. G., & Kay, S. A. (2007). PRR3 is a
vascular regulator of TOC1 stability in the Arabidopsis circadian clock. *Plant Cell*, *19*(11), 3462–
3473. https://doi.org/10.1105/tpc.107.054775

Park, Y. J., Kim, J. Y., Lee, J. H., Lee, B. D., Paek, N. C., & Park, C. M. (2020). GIGANTEA Shapes the
Photoperiodic Rhythms of Thermomorphogenic Growth in Arabidopsis. *Molecular Plant*, *13*(3),
459–470. https://doi.org/10.1016/j.molp.2020.01.003

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast and bias-
aware quantification of transcript expression. *Nature Methods*, *14*(4), 417–419.
https://doi.org/10.1038/nmeth.4197

Peeters, N. M., & Hanson, M. R. (2002). Transcript abundance supercedes editing efficiency as a factor in
developmental variation of chloroplast gene expression. *RNA*, *8*(4), 497–511.
https://doi.org/10.1017/S1355838202029424

Perez-Garcia, P., Serrano-Ron, L., & Moreno-Risueno, M. A. (2022). The nature of the root clock at single
cell resolution: Principles of communication and similarities with plant and animal pulsatile and
circadian mechanisms. *Current Opinion in Cell Biology*, *77*, 102102.
https://doi.org/10.1016/j.ceb.2022.102102

Prior, K. F., O'Donnell, A. J., Rund, S. S. C., Savill, N. J., van der Veen, D. R., & Reece, S. E. (2019). Host
circadian rhythms are disrupted during malaria infection in parasite genotype-specific manners.
Scientific Reports, *9*(1). https://doi.org/10.1038/s41598-019-47191-8

R Core Team. (2021). *R: A Language and Environment for Statistical Computing*.

Rijo-Ferreira, F., Acosta-Rodriguez, V. A., Abel, J. H., Kornblum, I., Bento, I., Kilaru, G., Klerman, E. B.,
Mota, M. M., & Takahashi, J. S. (2020). The malaria parasite has an intrinsic clock. *Science*, *368*,
746–753. http://science.sciencemag.org/

Rijo-Ferreira, F., Takahashi, J. S., & Figueiredo, L. M. (2017). Circadian rhythms in parasites. In *PLoS*
Pathogens (Vol. 13, Issue 10). Public Library of Science.
https://doi.org/10.1371/journal.ppat.1006590

Rolfe, S. A., Strelkov, S. E., Links, M. G., Clarke, W. E., Robinson, S. J., Djavaheri, M., Malinowski, R.,
Haddadi, P., Kagale, S., Parkin, I. A. P., Taheri, A., & Borhan, M. H. (2016). The compact genome of
the plant pathogen Plasmodiophora brassicae is adapted to intracellular interactions with host
Brassica spp. *BMC Genomics*, *17*(1), 1–15. https://doi.org/10.1186/s12864-016-2597-2

Sanchez, S. E., & Kay, S. A. (2016). The plant circadian clock: From a simple timekeeper to a complex
developmental manager. *Cold Spring Harbor Perspectives in Biology*, *8*(12), 1–16.
https://doi.org/10.1101/cshperspect.a027748

Schuller, A., Kehr, J., & Ludwig-Müller, J. (2014). Laser microdissection coupled to transcriptional profiling
of arabidopsis roots inoculated by plasmodiophora brassicae indicates a role for brassinosteroids in
clubroot formation. *Plant and Cell Physiology*, *55*(2), 392–411. https://doi.org/10.1093/pcp/pct174

Schwelm, A., Fogelqvist, J., Knaust, A., Jülke, S., Lilja, T., Bonilla-Rosso, G., Karlsson, M., Shevchenko, A.,
Dhandapani, V., Choi, S. R., Kim, H. G., Park, J. Y., Lim, Y. P., Ludwig-Müller, J., & Dixelius, C. (2015).
The Plasmodiophora brassicae genome reveals insights in its life cycle and ancestry of chitin
synthases. *Scientific Reports*, *5*(May), 1–12. https://doi.org/10.1038/srep11153

Seki, M., Ohara, T., Hearn, T. J., Frank, A., Da Silva, V. C. H., Caldana, C., Webb, A. A. R., & Satake, A.
(2017). Adjustment of the Arabidopsis circadian oscillator by sugar signalling dictates the regulation
of starch metabolism. *Scientific Reports*, *7*(1), 1–11. https://doi.org/10.1038/s41598-017-08325-y

Shor, E., Paik, I., Kangisser, S., Green, R., & Huq, E. (2017). PHYTOCHROME INTERACTING FACTORS
mediate metabolic control of the circadian system in Arabidopsis. *New Phytologist*, *215*(1), 217–
228. https://doi.org/10.1111/nph.14579

Shtin, M., Dello Ioio, R., & Del Bianco, M. (2022). It's Time for a Change: The Role of Gibberellin in Root
Meristem Development. *Frontiers in Plant Science*, *13*(May), 1–7.
https://doi.org/10.3389/fpls.2022.882517

Siemens, J., González, M. C., Wolf, S., Hofmann, C., Greiner, S., Du, Y., Rausch, T., Roitsch, T., & Ludwig-
Müller, J. (2011). Extracellular invertase is involved in the regulation of clubroot disease in
Arabidopsis thaliana. *Molecular Plant Pathology*, *12*(3), 247–262. https://doi.org/10.1111/j.1364-
3703.2010.00667.x

Siemens, J., Keller, I., Sarx, J., Kunz, S., Schuller, A., Nagel, W., Schmülling, T., Parniske, M., & Ludwig-
Müller, J. (2006). Transcriptome analysis of Arabidopsis clubroots indicate a key role for cytokinins
in disease development. *Molecular Plant-Microbe Interactions*, *19*(5), 480–494.
https://doi.org/10.1094/MPMI-19-0480

Sierra, R., Canas-Duarte, S. J., Burki, F., Schwelm, A., Fogelqvist, J., Dixelius, C., González-García, L. N.,
Gile, G. H., Slamovits, C. H., Klopp, C., Restrepo, S., Arzul, I., & Pawlowski, J. (2016). Evolutionary
origins of rhizarian parasites. *Molecular Biology and Evolution*, *33*(4), 980–983.
https://doi.org/10.1093/molbev/msv340

Siqueira, J. A., Otoni, W. C., & Araújo, W. L. (2022). The hidden half comes into the spotlight: Peeking
inside the black box of root developmental phases. *Plant Communications*, *3*(1), 1–17.
https://doi.org/10.1016/j.xplc.2021.100246

Smolko, A., Jülke, S., Benade, F., Široká, J., Pěnčík, A., Vuk, T., Bauer, N., Salopek-Sondi, B., & Ludwig-
Müller, J. (2024). The GH3 Protein from the Clubroot Pathogen Plasmodiophora brassicae Causes
Hormone-Related Phenotypes in Arabidopsis thaliana. *Journal of Plant Growth Regulation*, *43*(8),
2715–2738. https://doi.org/10.1007/s00344-024-11300-3

Song, Y., Jiang, Y., Kuai, B., & Li, L. (2018). Circadian clock-associated 1 inhibits leaf senescence in
arabidopsis. *Frontiers in Plant Science*, *9*(March), 1–9. https://doi.org/10.3389/fpls.2018.00280

Stefanowicz, K., Szymanska-Chargot, M., Truman, W., Walerowski, P., Olszak, M., Augustyniak, A.,
Kosmala, A., Zdunek, A., & Malinowski, R. (2021). Plasmodiophora brassicae-Triggered Cell
Enlargement and Loss of Cellular Integrity in Root Systems Are Mediated by Pectin Demethylation.
Frontiers in Plant Science, *12*(July). https://doi.org/10.3389/fpls.2021.711838

Thiruvengadam, M., Shih, C. F., & Yang, C. H. (2015). Expression of An Antisense Brassica oleracea
GIGANTEA (BoGI) Gene in Transgenic Broccoli Causes Delayed Flowering, Leaf Senescence, and
Post-Harvest Yellowing Retardation. *Plant Molecular Biology Reporter*, *33*(5), 1499–1509.
https://doi.org/10.1007/s11105-015-0852-3

Tu, J., Qin, L., Karunakaran, C., Wei, Y., & Peng, G. (2024). Lignin accumulation in cell wall plays a role in
clubroot resistance. *Frontiers in Plant Science*, *15*(July), 1–16.
https://doi.org/10.3389/fpls.2024.1401265

Uemoto, K., Mori, F., Yamauchi, S., Kubota, A., Takahashi, N., Egashira, H., Kunimoto, Y., Araki, T.,
Takemiya, A., Ito, H., & Endo, M. (2023). Root PRR7 Improves the Accuracy of the Shoot Circadian
Clock through Nutrient Transport. *Plant and Cell Physiology*, *64*(3), 352–362.
https://doi.org/10.1093/pcp/pcad003

Vandenbussche, F., Smalle, J., Le, J., Saibo, N. J. M., De Paepe, A., Chaerle, L., Tietz, O., Smets, R.,
Laarhoven, L. J. J., Harren, F. J. M., Van Onckelen, H., Palme, K., Verbelen, J. P., & Van Der Straeten,
D. (2003). The Arabidopsis mutant alh1 illustrates a cross talk between ethylene and auxin. *Plant*
Physiology, *131*(3), 1228–1238. https://doi.org/10.1104/pp.010850

Walerowski, P., Gündel, A., Yahaya, N., Truman, W., Sobczak, M., Olszak, M., Rolfe, S., Borisjuk, L., &
Malinowski, R. (2018). Clubroot disease stimulates early steps of phloem differentiation and
recruits SWEET sucrose transporters within developing galls. *Plant Cell*, *30*(12), 3058–3073.
https://doi.org/10.1105/tpc.18.00283

Walter, W., Sánchez-Cabo, F., & Ricote, M. (2015). GOplot: An R package for visually combining
expression data with functional analysis. *Bioinformatics*, *31*(17), 2912–2914.
https://doi.org/10.1093/bioinformatics/btv300

Walters, D. R., & McRoberts, N. (2006). Plants and biotrophs: a pivotal role for cytokinins? *Trends in Plant*
Science, *11*(12), 581–586. https://doi.org/10.1016/j.tplants.2006.10.003

Wang, J., Xu, G., Ning, Y., Wang, X., & Wang, G. L. (2022). Mitochondrial functions in plant immunity. In
Trends in Plant Science (Vol. 27, Issue 10, pp. 1063–1076). Elsevier Ltd.
https://doi.org/10.1016/j.tplants.2022.04.007

Wang, S., Steed, G., & Webb, A. A. R. (2022). Circadian entrainment in Arabidopsis. *Plant Physiology*,
190(2), 981–993. https://doi.org/10.1093/plphys/kiac204

Wang, X., An, Y., Xu, P., & Xiao, J. (2021). Functioning of PPR Proteins in Organelle RNA Metabolism and
Chloroplast Biogenesis. *Frontiers in Plant Science*, *12*(February), 1–8.
https://doi.org/10.3389/fpls.2021.627501

Wang, Y., & Tan, B. C. (2025). Pentatricopeptide repeat proteins in plants: Cellular functions, action
mechanisms, and potential applications. In *Plant Communications* (Vol. 6, Issue 2). Cell Press.
https://doi.org/10.1016/j.xplc.2024.101203

Wei, X., Zhang, Y., Zhao, Y., Xie, Z., Hossain, M. R., Yang, S., Shi, G., Lv, Y., Wang, Z., Tian, B., Su, H., Wei,
F., Zhang, X., & Yuan, Y. (2021). Root Transcriptome and Metabolome Profiling Reveal Key
Phytohormone-Related Genes and Pathways Involved Clubroot Resistance in Brassica rapa L.
Frontiers in Plant Science, *12*(December), 1–17. https://doi.org/10.3389/fpls.2021.759623

Wickham, H. (2010). ggplot2: Elegant Graphics for Data Analysis. *Journal of Statistical Software*, *1*(1),
128–129. https://doi.org/10.1002/wics.10

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., Fu, X., Liu, S., Bo, X.,
& Yu, G. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
Innovation, *2*(3), 100141. https://doi.org/10.1016/j.xinn.2021.100141

Yang, Y., Fan, G., Zhao, Y., Wen, Q., Wu, P., Meng, Y., & Shan, W. (2020). Cytidine-to-uridine RNA editing
factor NbMORF8 negatively regulates plant immunity to phytophthora pathogens. *Plant Physiology*,
184(4), 2182–2198. https://doi.org/10.1104/pp.20.00458

Yang, Y., Zhao, Y., Zhang, Y., Niu, L., Li, W., Lu, W., Li, J., Schäfer, P., Meng, Y., & Shan, W. (2022). A
mitochondrial RNA processing protein mediates plant immunity to a broad spectrum of pathogens
by modulating the mitochondrial oxidative burst. *Plant Cell*, *34*(6), 2343–2363.
https://doi.org/10.1093/plcell/koac082

Yazdanbakhsh, N., & Fisahn, J. (2010). Analysis of Arabidopsis thaliana root growth kinetics with high
temporal and spatial resolution. *Annals of Botany*, *105*(5), 783–791.
https://doi.org/10.1093/aob/mcq048

Yazdanbakhsh, N., Sulpice, R., Graf, A., Stitt, M., & Fisahn, J. (2011). Circadian control of root elongation
and C partitioning in Arabidopsis thaliana. *Plant, Cell and Environment*, *34*(6), 877–894.
https://doi.org/10.1111/j.1365-3040.2011.02286.x

Zheng, X. Y., Zhou, M., Yoo, H., Pruneda-Paz, J. L., Spivey, N. W., Kay, S. A., & Dong, X. (2015). Spatial and
temporal regulation of biosynthesis of the plant immune signal salicylic acid. *Proceedings of the*
National Academy of Sciences of the United States of America, *112*(30), 9166–9173.
https://doi.org/10.1073/pnas.1511182112

Zhu, J. Y., Oh, E., Wang, T., & Wang, Z. Y. (2016). TOC1-PIF4 interaction mediates the circadian gating of
thermoresponsive growth in Arabidopsis. *Nature Communications*, *7*, 1–10.
https://doi.org/10.1038/ncomms13692

1001

Supporting information

1004

1005 *Additional File 1 Phenotypical traits of rosettes (A and B) and root samples (C and D) for specimen of healthy and infected A.*
1006 *thaliana Col-0 harvested at 14 d.a.i. (top figure plate) and 21 d.a.i. (bottom figure plate). Scale bars in C and D = 5 mm.*

1007

1008

1009

1010 *Additional File 2 Detailed overview of the samples sequenced in this study*

Condition	Age (days after infection)	Hours after artificial dawn	Replicate	Raw reads

			code	
Control	14	5 hr 50' (day)	127925	41193025
		5 hr 50' (day)	127935	50545664
		5 hr 50' (day)	127936	35205326
		10 hr 40' (day)	127906	56725609
		10 hr 40' (day)	127912	54273366
		10 hr 40' (day)	127918	38720139
		15 hr 30' (night)	127926	41309013
		15 hr 30' (night)	127931	55027843
		15 hr 30' (night)	127937	38004169
		19 hr 20' (night)	127907	39958547
		19 hr 20' (night)	127913	37101449
		19 hr 20' (night)	127919	40693049
Infected	14	5 hr 50' (day)	127964	48508205
		5 hr 50' (day)	127970	41810661
		5 hr 50' (day)	127976	54739923
		10 hr 40' (day)	127946	42594195
		10 hr 40' (day)	127952	41155738
		10 hr 40' (day)	127958	58101982
		15 hr 30' (night)	127965	43247574
		15 hr 30' (night)	127971	46509846
		15 hr 30' (night)	127977	57602784
		19 hr 20' (night)	127947	40263406
		19 hr 20' (night)	127953	41398461
		19 hr 20' (night)	127959	50627500
Control	21	5 hr 50' (day)	127908	52533389
		5 hr 50' (day)	127914	36118824
		5 hr 50' (day)	127941	64871718
		10 hr 40' (day)	127928	41179331
		10 hr 40' (day)	127933	45662748
		10 hr 40' (day)	127939	54002775
		15 hr 30' (night)	127909	43278660
		15 hr 30' (night)	127915	43201375
		15 hr 30' (night)	127920	42466381
		19 hr 20' (night)	127929	43287455
		19 hr 20' (night)	127934	46270564
		19 hr 20' (night)	127940	53657896
Infected	21	5 hr 50' (day)	127948	59286277
		5 hr 50' (day)	127954	62740880
		5 hr 50' (day)	127960	46195940
		10 hr 40' (day)	127967	48074903
		10 hr 40' (day)	127973	61491275
		10 hr 40' (day)	127979	62580301

15 hr 30' (night)	127949	55537283
15 hr 30' (night)	127955	65514599
15 hr 30' (night)	127961	54889653
19 hr 20' (night)	127968	85469436
19 hr 20' (night)	127974	71731948
19 hr 20' (night)	127980	58536488

1011

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

1022   *Additional File 3 PCA on control A. thaliana plants, showing the decreased influence of the temporal factor (i.e., day and night)*
1023   *on the overall gene expression levels with the ageing of the plants.*

1024

1025

1026 *Additional File 4 Volcano plot showing the number of DEGs in the day (top figure) and night (bottom figure) in infected A.*
1027 *thaliana plants 14 DAI*

1028

1029

1030

Additional File 5 Volcano plot showing the number of DEGs in the day (top figure) and night (bottom figure) in infected A.
1031
1032   *thaliana plants 21 DAI*

1033

1034

1035
1036

GO term categories affected by *P.brassicae* infection in *A. thaliana* 14 d.a.i.							
Day				Night			
Upregulated							
ID	term	count	adj_pval	zscore	ID	term	count	adj_pval	zscore
GO:00 09636	response to toxic substance	55	8,92	3,64	GO:00 09636	response to toxic substance	68	11,73	3,40
GO:00 46906	tetrapyrrole binding	89	7,47	1,38	GO:00 09451	RNA modification	101	7,86	7,06
GO:00 48646	anatomical structure formation involved in morphogenesis	68	5,68	1,21	GO:01 40359	ABC-type transporter activity	41	6,98	0,47
GO:00 08299	isoprenoid biosynthetic process	54	4,74	0,27	GO:00 90332	stomatal closure	26	5,69	0,39
GO:00 48580	regulation of post-embryonic development	91	4,41	0,31	GO:00 00302	response to reactive oxygen species	50	5,36	0,28
GO:00 46148	pigment biosynthetic process	52	4,38	1,94	GO:00 02213	defense response to insect	32	4,97	1,06
GO:00 38023	signaling receptor activity	36	4,31	0,67	GO:00 48580	regulation of post-embryonic development	108	4,70	0,38
GO:00 48046	apoplast	70	4,22	1,20	GO:00 09739	response to gibberellin	40	4,41	0,32
GO:00 10287	plastoglobule	21	3,75	2,40	GO:00 48544	recognition of pollen	20	3,82	3,13
GO:00 09767	photosynthetic electron transport chain	19	3,41	3,44	GO:00 09991	response to extracellular stimulus	79	3,60	0,11
GO:00 42651	thylakoid membrane	73	3,14	5,97	GO:00 45165	cell fate commitment	30	2,96	0,73
GO:00 16829	lyase activity	65	3,06	0,12	GO:00 99503	secretory vesicle	46	2,90	0,59
GO:00 09739	response to gibberellin	31	2,94	1,26	GO:00 71214	cellular response to abiotic stimulus	63	2,81	0,13
GO:00 09644	response to high light intensity	19	2,87	1,15	GO:00 07166	cell surface receptor signaling pathway	19	2,70	0,69
GO:00 15267	channel activity	41	2,86	0,47	GO:20 00377	regulation of reactive oxygen species metabolic process	21	2,63	1,53
GO:01 02483	scopolin beta-glucosidase activity	14	2,39	0,53	GO:00 09723	response to ethylene	60	2,45	0,26
GO:00 10168	ER body	7	2,35	2,65	GO:19 02679	negative regulation of RNA biosynthetic process	53	2,45	0,41
GO:00 16054	organic acid catabolic process	35	2,09	0,85	GO:00 08171	O-methyltransferase activity	16	2,38	0,50
GO:20 00377	regulation of reactive oxygen species metabolic process	17	2,02	1,21	GO:00 48825	cotyledon development	18	2,32	1,41
GO:00 48544	recognition of pollen	14	1,96	0,53	GO:00 00712	resolution of meiotic recombination intermediates	7	2,21	2,65
GO:00 00373	Group II intron splicing	9	1,82	0,33	GO:00 42440	pigment metabolic process	82	2,01	0,66
GO:00 09630	gravitropism	21	1,66	0,65	GO:00 08283	cell population proliferation	27	1,90	0,96
GO:00 08283	cell population proliferation	22	1,51	0,43	GO:00 48235	pollen sperm cell differentiation	12	1,87	0,58
GO:19 00865	chloroplast RNA modification	6	1,45	1,63	GO:00 00079	regulation of cyclin-dependent protein serine/threonine	15	1,84	0,26

ID	term	count	adj_pval	zscore	ID	term	count	adj_pval	zscore
						kinase activity			
GO:00 06638	neutral lipid metabolic process	10	1,41	0,63	GO:00 42793	plastid transcription	7	1,84	1,89
					GO:00 02832	negative regulation of response to biotic stimulus	15	1,78	0,77
					GO:00 10114	response to red light	25	1,64	0,20
					GO:00 10090	trichome morphogenesis	19	1,63	0,23
					GO:00 19432	triglyceride biosynthetic process	9	1,47	0,33
					GO:19 01681	sulfur compound binding	11	1,40	0,30
Zero z-score									
ID	term	count	adj_pval	zscore	ID	term	count	adj_pval	zscore
GO:00 02238	response to molecule of fungal origin	10	3,43	0,00	GO:00 08299	isoprenoid biosynthetic process	74	8,81	0,00
					GO:00 48437	floral organ development	72	4,62	0,00
					GO:19 01698	response to nitrogen compound	64	3,43	0,00
					GO:00 09608	response to symbiont	6	1,75	0,00
					GO:00 09629	response to gravity	26	1,66	0,00
Downregulated									
ID	term	count	adj_pval	zscore	ID	term	count	adj_pval	zscore
GO:00 71456	cellular response to hypoxia	86	18,78	-3,23	GO:00 44262	cellular carbohydrate metabolic process	144	17,31	-5,33
GO:00 44262	cellular carbohydrate metabolic process	125	17,57	-4,74	GO:00 03953	NAD+ nucleosidase activity	27	16,14	-4,81
GO:00 44550	secondary metabolite biosynthetic process	107	14,08	-3,77	GO:00 09698	phenylpropanoid metabolic process	64	14,42	-3,25
GO:00 03953	NAD+ nucleosidase activity	24	14,04	-3,27	GO:00 36294	cellular response to decreased oxygen levels	87	13,71	-3,11
GO:00 09751	response to salicylic acid	107	11,64	-4,54	GO:00 16798	hydrolase activity acting on glycosyl bonds	127	11,94	-5,06
GO:00 12501	programmed cell death	59	9,99	-2,21	GO:00 20037	heme binding	104	9,74	-0,98
GO:00 48584	positive regulation of response to stimulus	72	7,78	-1,89	GO:00 09751	response to salicylic acid	117	9,70	-2,68
GO:00 42537	benzene-containing compound metabolic process	29	7,18	-2,79	GO:00 45229	external encapsulating structure organization	107	9,36	-3,96
GO:00 42802	identical protein binding	89	6,82	-0,74	GO:00 16143	S-glycoside metabolic process	82	9,34	-1,77
GO:00 48437	floral organ development	69	6,71	-0,12	GO:00 48646	anatomical structure formation involved in morphogenesis	87	8,42	-1,61
GO:00 00302	response to reactive oxygen species	47	6,58	-0,44	GO:00 48584	positive regulation of response to stimulus	84	8,32	-0,44
GO:00 62012	regulation of small molecule metabolic process	54	6,54	-4,08	GO:00 10087	phloem or xylem histogenesis	55	7,08	-4,45
GO:00 45229	external encapsulating structure organization	81	5,53	-3,22	GO:20 00652	regulation of secondary cell wall biogenesis	28	6,87	-4,54
GO:00 90332	stomatal closure	23	5,38	-1,46	GO:00 42537	benzene-containing compound metabolic process	31	6,56	-3,05
GO:19	regulation of cell wall	25	5,14	-4,60	GO:00	organic acid transport	46	5,92	-1,18

GO ID	Description	N	Val1	Val2	GO ID	Description	N	Val1	Val2
03338	organization or biogenesis				15849				
GO:00 09743	response to carbohydrate	48	4,95	-0,87	GO:00 10150	leaf senescence	83	5,86	-0,77
GO:00 16747	acyltransferase activity transferring groups other than amino-acyl groups	69	4,63	-0,60	GO:00 10648	negative regulation of cell communication	41	5,85	-0,78
GO:00 48585	negative regulation of response to stimulus	56	4,49	-0,80	GO:00 06811	monoatomic ion transport	94	5,78	-0,83
GO:00 10087	phloem or xylem histogenesis	41	4,16	-3,90	GO:00 15698	inorganic anion transport	40	5,54	-1,58
GO:00 15698	inorganic anion transport	32	4,09	-0,71	GO:00 10143	cutin biosynthetic process	16	5,41	-0,50
GO:00 09733	response to auxin	83	3,77	-0,99	GO:00 08219	cell death	68	5,30	-2,43
GO:00 10150	leaf senescence	64	3,75	-0,75	GO:00 00981	DNA-binding transcription factor activity RNA polymerase II-specific	79	5,30	-0,56
GO:00 46777	protein autophosphorylation	40	3,70	-2,85	GO:00 07623	circadian rhythm	49	4,99	-0,43
GO:00 42594	response to starvation	48	3,64	-0,29	GO:00 90627	plant epidermal cell differentiation	62	4,92	-2,29
GO:01 40359	ABC-type transporter activity	29	3,49	-0,56	GO:00 71365	cellular response to auxin stimulus	42	4,84	-0,31
GO:00 15849	organic acid transport	34	3,43	-0,34	GO:00 42802	identical protein binding	94	4,33	-0,62
GO:00 06816	calcium ion transport	23	3,38	-1,46	GO:00 09735	response to cytokinin	39	3,62	-1,12
GO:00 48235	pollen sperm cell differentiation	13	3,02	-0,28	GO:00 09308	amine metabolic process	43	3,48	-2,29
GO:00 70417	cellular response to cold	13	3,02	-0,28	GO:00 43621	protein self-association	35	3,32	-0,51
GO:00 43621	protein self-association	29	2,79	-0,56	GO:00 34285	response to disaccharide	25	3,22	-1,80
GO:00 06631	fatty acid metabolic process	61	2,77	-0,38	GO:00 19899	enzyme binding	81	3,01	-2,33
GO:00 09308	amine metabolic process	35	2,76	-2,87	GO:00 22603	regulation of anatomical structure morphogenesis	50	2,84	-1,70
GO:00 09624	response to nematode	22	2,76	-1,28	GO:00 46777	protein autophosphorylation	43	2,83	-0,15
GO:19 01698	response to nitrogen compound	52	2,72	-0,28	GO:00 09741	response to brassinosteroid	34	2,65	-2,74
GO:00 42887	amide transmembrane transporter activity	18	2,64	-1,41	GO:00 09705	plant-type vacuole membrane	36	2,58	-1,33
GO:00 48767	root hair elongation	21	2,46	-1,53	GO:00 15295	solute:proton symporter activity	13	2,57	-0,83
GO:00 09408	response to heat	70	2,46	-0,96	GO:00 06880	intracellular sequestering of iron ion	8	2,47	-0,71
GO:00 19900	kinase binding	29	2,43	-3,53	GO:00 09826	unidimensional cell growth	96	2,45	-2,04
GO:00 10073	meristem maintenance	51	2,41	-0,42	GO:00 48046	apoplast	72	2,44	-1,18
GO:00 09625	response to insect	13	2,38	-0,28	GO:00 09624	response to nematode	24	2,42	-0,82
GO:00 00981	DNA-binding transcription factor activity RNA polymerase II-specific	56	2,32	-0,53	GO:00 06833	water transport	14	2,34	-1,07
GO:00 51644	plastid localization	9	2,27	-0,33	GO:00 08236	serine-type peptidase activity	43	2,33	-1,98
GO:00	nucleoside	11	2,24	-2,11	GO:00	meristem maintenance	59	2,26	-0,13

05337	transmembrane transporter activity				10073				
GO:00 42578	phosphoric ester hydrolase activity	66	2,24	-2,95	GO:00 51119	sugar transmembrane transporter activity	19	2,18	-1,61
GO:00 71215	cellular response to abscisic acid stimulus	59	2,19	-3,78	GO:00 42886	amide transport	15	2,18	-1,29
GO:00 07389	pattern specification process	60	2,11	-1,03	GO:00 10038	response to metal ion	51	2,05	-2,66
GO:00 09705	plant-type vacuole membrane	29	2,06	-1,30	GO:00 08374	O-acyltransferase activity	30	2,02	-2,92
GO:00 09531	secondary cell wall	9	2,06	-3,00	GO:00 09411	response to UV	31	2,00	-0,90
GO:00 05507	copper ion binding	37	2,01	-0,16	GO:00 30570	pectate lyase activity	10	1,96	-0,63
GO:00 55062	phosphate ion homeostasis	9	1,91	-1,00	GO:00 70417	cellular response to cold	12	1,87	-0,58
GO:00 09269	response to desiccation	9	1,91	-1,67	GO:00 35556	intracellular signal transduction	90	1,87	-1,26
GO:00 09826	unidimensional cell growth	77	1,82	-1,71	GO:00 10214	seed coat development	17	1,87	-1,70
GO:00 04620	phospholipase activity	15	1,76	-1,81	GO:00 51049	regulation of transport	30	1,85	-1,46
GO:00 71214	cellular response to abiotic stimulus	48	1,64	-0,29	GO:00 50660	flavin adenine dinucleotide binding	37	1,84	-0,82
GO:00 09735	response to cytokinin	27	1,58	-0,96	GO:00 04551	dinucleotide phosphatase activity	6	1,84	-1,63
GO:00 08171	O-methyltransferase activity	12	1,53	-0,58	GO:00 09615	response to virus	25	1,83	-1,00
GO:00 30145	manganese ion binding	13	1,52	-3,05	GO:00 06631	fatty acid metabolic process	65	1,67	-1,86
GO:00 50660	flavin adenine dinucleotide binding	30	1,51	-0,37	GO:00 09850	auxin metabolic process	19	1,63	-0,23
GO:00 40034	regulation of development heterochronic	13	1,47	-0,28	GO:00 09408	response to heat	77	1,63	-1,48
GO:00 45926	negative regulation of growth	10	1,47	-0,63	GO:00 16829	lyase activity	68	1,60	-0,73
GO:00 71367	cellular response to brassinosteroid stimulus	18	1,45	-2,36	GO:00 51923	sulfation	7	1,55	-2,65
GO:00 16307	phosphatidylinositol phosphate kinase activity	6	1,43	-0,82	GO:00 30145	manganese ion binding	15	1,54	-3,36
GO:00 15081	sodium ion transmembrane transporter activity	11	1,32	-1,51	GO:00 07010	cytoskeleton organization	65	1,49	-3,35
					GO:00 09269	response to desiccation	9	1,47	-1,67
					GO:00 90158	endoplasmic reticulum membrane organization	6	1,42	-1,63
					GO:00 15851	nucleobase transport	9	1,38	-1,00
					GO:00 09631	cold acclimation	15	1,31	-2,32

1037

1038

1039

1040

1041

1042

1043
1044

GO term categories affected by *P.brassicae* infection in *A. thaliana* 21 d.a.i.								
Day				Night				
Upregulated								
ID	term	count	adj_pval	zscore	ID	term	count	adj_pval	zscore
GO:0015979	photosynthesis	123	14,82	8,75	GO:0015979	photosynthesis	132	16,94	9,05
GO:0009657	plastid organization	121	11,02	5,36	GO:0009991	response to extracellular stimulus	148	14,73	1,97
GO:0016054	organic acid catabolic process	76	8,94	0,92	GO:0009657	plastid organization	131	12,99	6,55
GO:0046148	pigment biosynthetic process	91	8,43	3,25	GO:0003953	NAD+ nucleosidase activity	27	11,87	0,19
GO:0009646	response to absence of light	33	8,32	0,87	GO:0000302	response to reactive oxygen species	81	11,34	0,56
GO:0007623	circadian rhythm	68	6,98	0,97	GO:0042802	identical protein binding	151	9,49	0,08
GO:0009639	response to red or far red light	134	5,25	0,69	GO:0044282	small molecule catabolic process	99	9,20	0,90
GO:0002213	defense response to insect	37	3,95	0,49	GO:0009642	response to light intensity	143	9,17	5,44
GO:0005777	peroxisome	94	2,63	0,41	GO:0015995	chlorophyll biosynthetic process	51	8,01	4,90
GO:0000373	Group II intron splicing	13	2,17	1,39	GO:0071214	cellular response to abiotic stimulus	107	7,63	1,45
GO:0007166	cell surface receptor signaling pathway	21	1,81	2,40	GO:0007623	circadian rhythm	72	7,53	2,59
GO:0006641	triglyceride metabolic process	14	1,79	0,53	GO:0009743	response to carbohydrate	81	7,52	1,00
GO:1901361	organic cyclic compound catabolic process	90	1,55	0,84	GO:0010150	leaf senescence	115	6,56	1,40
GO:0015749	monosaccharide transmembrane transport	12	1,49	1,15	GO:0051093	negative regulation of developmental process	73	5,26	0,35
GO:0005315	inorganic phosphate transmembrane transporter activity	9	1,35	1,00	GO:0071949	FAD binding	37	4,80	1,81
					GO:0043621	protein self-association	50	4,58	0,28
					GO:0015144	carbohydrate transmembrane transporter activity	39	4,41	0,48
					GO:0051253	negative regulation of RNA metabolic	81	3,19	0,56

ID	term	count	adj_pval	zscore	ID	term	count	adj_pval	zscore
						process			
					GO:0002213	defense response to insect	36	3,10	0,33
					GO:0008271	secondary active sulfate transmembrane transporter activity	9	3,00	0,33
					GO:0100014	meristem initiation	21	2,49	0,22
					GO:0022411	cellular component disassembly	40	2,33	0,63
					GO:0009704	de-etiolation	10	2,29	0,63
					GO:0040034	regulation of development heterochronic	22	2,28	0,43
					GO:0009739	response to gibberellin	45	2,28	0,45
					GO:0000373	Group II intron splicing	13	2,01	1,39
					GO:1900865	chloroplast RNA modification	9	1,92	2,33
					GO:0010181	FMN binding	13	1,57	0,83
					GO:0045454	cell redox homeostasis	16	1,48	2,50
					GO:0031406	carboxylic acid binding	37	1,33	0,82
					GO:0019432	triglyceride biosynthetic process	11	1,32	0,90
Zero z-score									
ID	term	count	adj_pval	zscore	ID	term	count	adj_pval	zscore
GO:0080027	response to herbivore	18	6,49	0	GO:0009636	response to toxic substance	94	15,27	0
GO:1901699	cellular response to nitrogen compound	32	4,90	0	GO:0010118	stomatal movement	78	9,44	0
GO:0009624	response to nematode	36	4,40	0					
GO:0004033	aldo-keto reductase (NADP) activity	12	2,49	0					
Downregulated									
ID	term	count	adj_pval	zscore	ID	term	count	adj_pval	zscore
GO:0044262	cellular carbohydrate metabolic process	196	23,13	-6,00	GO:0044262	cellular carbohydrate metabolic process	208	25,39	-5,82
GO:0009698	phenylpropanoid metabolic process	84	18,71	-4,80	GO:0044550	secondary metabolite biosynthetic process	171	17,41	-8,34

GO ID	Description				GO ID	Description			
GO:0009636	response to toxic substance	94	16,84	-1,24	GO:0045229	external encapsulating structure organization	156	13,46	-7,37
GO:0003953	NAD+ nucleosidase activity	29	15,42	-0,93	GO:0046906	tetrapyrrole binding	158	12,94	-1,91
GO:0046906	tetrapyrrole binding	158	15,42	-1,75	GO:0098660	inorganic ion transmembrane transport	107	12,43	-1,64
GO:0016143	S-glycoside metabolic process	117	14,45	-6,19	GO:0006721	terpenoid metabolic process	120	12,11	-0,55
GO:0036294	cellular response to decreased oxygen levels	109	14,30	-3,93	GO:0036294	cellular response to decreased oxygen levels	107	11,74	-3,96
GO:0009991	response to extracellular stimulus	139	13,31	-0,93	GO:0090698	post-embryonic plant morphogenesis	99	11,65	-1,51
GO:0045229	external encapsulating structure organization	148	12,76	-7,07	GO:0140359	ABC-type transporter activity	60	11,13	-1,29
GO:0009751	response to salicylic acid	159	12,21	-5,63	GO:0009751	response to salicylic acid	162	11,09	-5,34
GO:0048046	apoplast	126	11,47	-1,07	GO:0006631	fatty acid metabolic process	132	11,01	-1,04
GO:0000302	response to reactive oxygen species	76	10,23	-0,46	GO:1901615	organic hydroxy compound metabolic process	158	10,82	-2,23
GO:0098660	inorganic ion transmembrane transport	94	8,68	-1,24	GO:0048046	apoplast	127	10,03	-0,44
GO:0010016	shoot system morphogenesis	104	8,66	-3,14	GO:0090627	plant epidermal cell differentiation	93	7,81	-4,87
GO:0006721	terpenoid metabolic process	106	8,62	-2,33	GO:0009966	regulation of signal transduction	155	7,60	-4,10
GO:0042537	benzene-containing compound metabolic process	41	8,59	-2,97	GO:0042446	hormone biosynthetic process	40	7,56	-3,48
GO:0006631	fatty acid metabolic process	119	8,38	-1,19	GO:0006955	immune response	114	7,50	-0,75
GO:0090696	post-embryonic plant organ development	83	7,98	-1,43	GO:0015849	organic acid transport	63	7,27	-1,64
GO:0016798	hydrolase activity acting on glycosyl bonds	150	7,89	-5,88	GO:0016829	lyase activity	123	6,77	-3,70
GO:0090627	plant epidermal cell differentiation	89	7,64	-6,04	GO:0016746	acyltransferase activity	127	6,66	-2,40
GO:0006955	immune response	110	7,60	-3,05	GO:0009624	response to nematode	42	6,55	-0,31
GO:0016747	acyltransferase activity transferring groups other than amino-acyl groups	116	7,43	-2,04	GO:0009225	nucleotide-sugar metabolic process	30	6,10	-3,65
GO:0010817	regulation of hormone levels	107	7,40	-5,32	GO:2000280	regulation of root development	39	5,74	-2,08

GO ID					GO ID				
GO:0015849	organic acid transport	61	7,34	-2,94	GO:0010038	response to metal ion	87	5,69	-2,47
GO:0140359	ABC-type transporter activity	51	7,24	-1,82	GO:0010143	cutin biosynthetic process	19	5,47	-1,61
GO:0016682	oxidoreductase activity acting on diphenols and related substances as donors oxygen as acceptor	21	7,08	-3,71	GO:0015267	channel activity	74	5,30	-1,86
GO:1901615	organic hydroxy compound metabolic process	138	6,97	-2,04	GO:0009741	response to brassinosteroid	54	5,25	-2,72
GO:0010118	stomatal movement	69	6,91	-1,81	GO:0009733	response to auxin	144	5,24	-2,17
GO:0010087	phloem or xylem histogenesis	68	6,77	-5,82	GO:0048226	Casparian strip	13	4,87	-3,61
GO:0042802	identical protein binding	134	6,56	-1,21	GO:0022626	cytosolic ribosome	106	4,85	-6,60
GO:0009741	response to brassinosteroid	55	6,42	-3,10	GO:0005856	cytoskeleton	124	4,80	-5,75
GO:0048226	Casparian strip	14	6,39	-3,74	GO:0035556	intracellular signal transduction	146	4,40	-1,99
GO:0022626	cytosolic ribosome	107	6,39	-6,48	GO:0004620	phospholipase activity	29	4,35	-2,41
GO:0010038	response to metal ion	85	6,15	-2,71	GO:0102483	scopolin beta-glucosidase activity	24	4,31	-0,41
GO:0048437	floral organ development	99	5,92	-2,11	GO:0010087	phloem or xylem histogenesis	63	4,25	-4,66
GO:0010646	regulation of cell communication	146	5,90	-5,96	GO:0007389	pattern specification process	112	4,23	-3,59
GO:0016829	lyase activity	114	5,84	-3,56	GO:0009826	unidimensional cell growth	148	4,20	-3,45
GO:0009733	response to auxin	140	5,77	-3,21	GO:0008283	cell population proliferation	44	4,07	-0,30
GO:0090693	plant organ senescence	120	5,68	-1,10	GO:0048438	floral whorl development	77	3,89	-1,03
GO:0010143	cutin biosynthetic process	18	5,06	-1,89	GO:1904680	peptide transmembrane transporter activity	21	3,83	-1,09
GO:0015267	channel activity	70	5,00	-2,15	GO:0030029	actin filament-based process	49	3,81	-3,00
GO:0009743	response to carbohydrate	70	4,90	-0,48	GO:0051923	sulfation	12	3,67	-2,31
GO:0006833	water transport	22	4,90	-2,13	GO:0009615	response to virus	40	3,63	-0,95
GO:0019900	kinase binding	51	4,87	-4,90	GO:0090351	seedling development	75	3,61	-1,27

GO ID	Term	Count			GO ID	Term	Count		
GO:0005507	copper ion binding	68	4,66	-2,67	GO:0006972	hyperosmotic response	32	3,54	-0,71
GO:0070417	cellular response to cold	20	4,55	-1,79	GO:0006576	biogenic amine metabolic process	38	3,47	-3,57
GO:0006972	hyperosmotic response	33	4,45	-1,22	GO:0005509	calcium ion binding	85	3,36	-3,80
GO:0051094	positive regulation of developmental process	54	4,42	-2,18	GO:0009408	response to heat	122	3,23	-1,99
GO:0009723	response to ethylene	89	4,37	-2,23	GO:0030145	manganese ion binding	24	3,16	-2,45
GO:0051093	negative regulation of developmental process	67	4,33	-0,61	GO:0055086	nucleobase-containing small molecule metabolic process	124	3,10	-0,54
GO:0008171	O-methyltransferase activity	23	3,98	-1,04	GO:0009629	response to gravity	41	3,09	-1,72
GO:0005774	vacuolar membrane	69	3,92	-3,49	GO:0099503	secretory vesicle	60	3,03	-2,84
GO:1903338	regulation of cell wall organization or biogenesis	31	3,86	-4,49	GO:0009631	cold acclimation	25	2,94	-0,20
GO:0038023	signaling receptor activity	50	3,68	-1,13	GO:0006833	water transport	19	2,90	-2,06
GO:0035556	intracellular signal transduction	135	3,64	-3,87	GO:0035266	meristem growth	37	2,87	-0,16
GO:1901568	fatty acid derivative metabolic process	25	3,43	-1,40	GO:0016614	oxidoreductase activity acting on CH-OH group of donors	69	2,82	-0,12
GO:0090351	seedling development	71	3,43	-1,54	GO:0000981	DNA-binding transcription factor activity RNA polymerase II-specific	95	2,78	-0,31
GO:0050660	flavin adenine dinucleotide binding	55	3,34	-0,13	GO:0008171	O-methyltransferase activity	21	2,73	-1,09
GO:0008283	cell population proliferation	40	3,29	-2,21	GO:0044273	sulfur compound catabolic process	13	2,73	-2,50
GO:0006576	biogenic amine metabolic process	36	3,28	-4,00	GO:0005507	copper ion binding	63	2,68	-3,15
GO:0009826	unidimensional cell growth	135	3,14	-5,25	GO:1901698	response to nitrogen compound	84	2,64	-0,87
GO:0051923	sulfation	11	3,13	-2,71	GO:0009838	abscission	19	2,59	-2,06
GO:0005509	calcium ion binding	80	3,10	-4,70	GO:0032504	multicellular organism reproduction	81	2,58	-2,11
GO:0010073	meristem maintenance	83	3,03	-2,52	GO:0008236	serine-type peptidase activity	59	2,47	-1,95

GO ID	Description	n	v1	v2	GO ID	Description	n	v1	v2
GO:0009739	response to gibberellin	46	3,00	-0,59	GO:0042886	amide transport	20	2,47	-0,89
GO:0050829	defense response to Gram-negative bacterium	13	2,94	-1,94	GO:0008378	galactosyltransferase activity	20	2,47	-2,24
GO:0009735	response to cytokinin	47	2,83	-2,19	GO:0005615	extracellular space	46	2,38	-1,77
GO:0000981	DNA-binding transcription factor activity RNA polymerase II-specific	91	2,81	-3,04	GO:0010223	secondary shoot formation	17	2,30	-1,21
GO:0042578	phosphoric ester hydrolase activity	109	2,81	-3,54	GO:0000278	mitotic cell cycle	85	2,16	-4,23
GO:0008236	serine-type peptidase activity	58	2,79	-2,63	GO:0016311	dephosphorylation	70	2,10	-2,63
GO:0008378	galactosyltransferase activity	20	2,72	-2,24	GO:1901607	alpha-amino acid biosynthetic process	56	2,08	-1,60
GO:0051253	negative regulation of RNA metabolic process	75	2,68	-0,81	GO:0019199	transmembrane receptor protein kinase activity	18	2,05	-0,47
GO:0032504	multicellular organism reproduction	78	2,68	-4,08	GO:0044087	regulation of cellular component biogenesis	47	2,04	-3,94
GO:0099503	secretory vesicle	56	2,63	-2,94	GO:0015851	nucleobase transport	13	2,01	-0,83
GO:0030029	actin filament-based process	43	2,62	-3,81	GO:0006644	phospholipid metabolic process	67	1,94	-1,83
GO:0051302	regulation of cell division	30	2,42	-1,10	GO:0003008	system process	11	1,87	-1,51
GO:0009411	response to UV	42	2,36	-1,54	GO:0032879	regulation of localization	47	1,68	-1,60
GO:0042886	amide transport	19	2,33	-2,52	GO:0045927	positive regulation of growth	17	1,51	-0,24
GO:0030145	manganese ion binding	21	2,27	-2,40	GO:0009555	pollen development	104	1,43	-2,16
GO:0000079	regulation of cyclin-dependent protein serine/threonine kinase activity	20	2,25	-1,79					
GO:0015294	solute:monoatomic cation symporter activity	17	2,24	-0,73					
GO:0043094	cellular metabolic compound salvage	29	2,20	-0,56					
GO:0043621	protein self-association	40	2,16	-0,63					
GO:0009631	cold acclimation	22	2,12	-1,28					
GO:0009408	response to heat	109	2,08	-1,25					
GO:0008970	phospholipase A1 activity	9	2,02	-3,00					
GO:0003008	system process	11	2,00	-1,51					
GO:0000278	mitotic cell cycle	80	1,96	-5,81					

GO:009 9080	supramolecular complex	113	1,87	-4,80						
GO:000 5615	extracellular space	42	1,87	-3,09						
GO:190 1607	alpha-amino acid biosynthetic process	52	1,77	-1,11						
GO:001 0214	seed coat development	20	1,44	-2,24						
GO:000 9606	tropism	49	1,32	-2,14						
GO:005 5086	nucleobase-containing small molecule metabolic process	105	1,31	-0,88						

1045

1046

1047

1048

1049

1050

1051 *Additional File 8 GO terms overrepresentation analysis of clusters of DEGs driven by the interaction between infection and time*
1052 *of the day 14 DAI*

1053

1054

1055 *Additional File 9 GO terms overrepresentation analysis of clusters of DEGs driven by the interaction between infection and time*
1056 *of the day 21 DAI*

1057

1058 https://fileshare.uibk.ac.at/f/03d6ddfe1e6145bca1a6/?dl=1

1059 *Additional File 10 Enriched GO terms and corresponding genes underpinning figure 2 and 3 before data reduction.*