Readiness potential in human functional magnetic resonance imaging motor task data.

Marika Strindberg[1,*], Yu Tian[2] & Peter Fransson[1]

[1]Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden.

[2] Center for Systems Biology, Dresden, Germany.

* Corresponding author
Marika Strindberg
Department of Clinical Neuroscience
Nobels väg 9
SE-171 77 Stockholm, SWEDEN
E-mail: marika.strindberg@ki.se

Themes: Phase synchronization, readiness potential, anticipation, attention, fMRI.

Significance statement

The readiness potential (RP) is observed in EEG recordings as a slow build-up of negative electrical potential in the supplementary motor area (SMA) starting up to two seconds before self-initiated motor movements. It is believed to play a causal role in motor preparation. Given that it is only detectable by averaging 30-40 epochs, its existence as a real signal has been challenged by new modeling approaches which suggests the early RP-component is merely a summation of stochastic noise. In this study, we demonstrate that the equivalent of the RP is present in functional magnetic resonance imaging (fMRI) data in a distributed network known as the "action mode network". The RP-equivalent is present in single subjects and single epochs.

Abstract

The readiness potential (RP), also known as the Bereichtschaftspotential, is commonly observed in EEG recordings as a slow build-up of negative electrical potential prior to self-directed motor movements (Kornhuber et al. 1965). In this study, we analyzed motor task-based functional magnetic resonance (fMRI) data acquired from 262 human participants, focusing on the degree of signal phase synchronization across different frequency spans and brain regions. A new method that clusters brain regions based on instantaneous phase allowed for time resolved estimation of cross-subject phase synchronization. We show that during rest periods that precede movement, an fMRI equivalent of the RP is gradually established in a network that encompasses the bilateral supplementary motor area and parts of the cingulo-opercular network, recently described as the brain's Action Mode network (Dosenbach et al. 2025). Importantly, an anticipatory negative shift in the fMRI signal was observed in both single subjects and single epochs. Tongue movement elicited strong synchronization between the orbitofrontal, ventromedial, and temporal pole cortices. Our findings suggest a link between fMRI and electrophysiological recordings of anticipatory motor events in the brain. The method presented here grades brain synchronization along an in-phase and anti-phase continuum and has applications in clinical settings, as well as for cognitive brain mapping that goes beyond anticipation.

Introduction.

The readiness potential (RP) is characterized by a gradual increase in negative potential, observable in time-locked averages of neurophysiological recordings of the brain preceding self-initiated motor movements[1-3]. The RP comprises early and late components. The early component commences up to 2 s prior to motor movement and predominantly reflects bilateral activity in the supplementary motor area (SMA), pre-SMA, and anterior mid-cingulate cortex. The late component initiates approximately 400 ms before movement onset and is lateralized to the hemisphere opposite the motor

movement, corresponding to activity in the motor cortex [4]. By contrast, self-reports and

empirical evidence indicate that the intention to act typically arises 200 ms before a

motor action [5].

Since its initial observation in the 1960s[3], the readiness potential (RP) has been

extensively investigated[1,6–8]. However, its functional significance and existence as a

genuine phenomenon in the brain remain debatable. The discrepancy in timing between

self-reports of the decision to act and the onset of the RP has even been employed as an

argument against the concept of free will[9]. The RP is most readily observed prior to self-

initiated movement, wherein participants are instructed to move when they feel the urge

to do so. This contrasts with cued movements, in which participants moved only after

receiving a cue. Although the RP is also observed in the latter context, it manifests with

reduced amplitude[6]. Common to these scenarios is the perceived imperative to move,

with the distinction that timing is less constrained in the self-paced scenario.

The classical interpretation posits that the readiness potential (RP) plays a causal role in

motor preparation. However, RP has also been observed in instances where participants

decide to move but do not execute the movement[6]. This challenges the strict classical

interpretation and suggests that RP may reflect a more general decision-making process.

Consequently, it remains uncertain whether the RP is indeed distinct from other negative

potentials observed prior to cognitive tasks in cued paradigms that do not necessarily

involve a motor component, such as the contingent negative variation (CNV) and

stimulus-preceding negativity (SPN)[2]. The CNV and SPN are generally considered to

reflect anticipation and attention, engaging brain areas essential for upcoming tasks [2].

A more recent explanatory framework, supported by modelling approaches and empirical

data, has provided a radical revaluation of the RP [10]. The stochastic accumulator model

proposes that the RP may not be a genuine signal but rather an artifact resulting from

averaging time-locked stochastic noise. According to this model, the decision to move

occurs when the accumulation of noise reaches a certain (undefined) threshold. Thus,

averaging the epochs of movement introduces a selection bias where the pre-threshold

data are inherently lower. Consequently, this model suggests that the post-threshold or

late RP component reflects a decision process, whereas the pre-threshold build-up
represents stochastic noise.

Recent empirical data suggest that slow cortical potentials (SCP) could account for the
RP phenomena[11]. SCPs are slow fluctuations (0.01–1 Hz) in the cortical potential in
neurophysiological recordings that are spontaneously present[6,11]. There is strong
evidence of a correlation between SCPs and slow fluctuations in BOLD fMRI [12]. A negative
shift in SCPs has been linked to increased responsiveness and cognitive performance [11].
Experimental findings indicate that both the subjective urge to move and the likelihood of
movement are greater during a negative shift than during a positive shift in SCP. The SCP
sampling hypothesis posits that the RP represents the subjective urge to move [6,11]. From
this perspective, the early component of the RP is the summation of the predominantly
negative shifts in the SCP preceding movement. It has also been demonstrated that it is
possible to modulate the SCP by intention and delay movement onset, even when a
strong negative shift, indicative of an urge, is present[11,13].

In summary, three principal explanatory frameworks for the RP have been proposed: 1) a
direct causal role in preparing motor action, rendering it necessary for motor movement;
2) a result of stochastic fluctuations that trigger a decision to move upon reaching a
certain threshold, thereby rendering the early component of the RP an artifact; and 3) a
reflection of the likelihood of ongoing fluctuations in cortical potential, where negative
shifts create an urge to move.
A central issue contributing to the varied interpretations of the readiness potential (RP) is
the low signal-to-noise ratio in electroencephalography (EEG), which requires averaging
across 30-40 time-locked epochs or several electrodes to observe the RP[1,2,14]. A crucial
step towards a more comprehensive understanding of the RP involves unequivocally
determining whether the RP exists within single epochs or is merely an artifact of
averaging.

In this study, we leverage the enhanced signal-to-noise ratio and spatial precision of
functional magnetic resonance imaging (fMRI) to address this question. We employed a
new time-resolved data analysis method using a substantial sample of the Human

Connectome Project dataset[15,16], focusing on fMRI motor task data involving hand, foot, and tongue movements across two distinct BOLD signal frequency domains, 0.066-0.189 Hz and 0.030-0.010 Hz. Our findings show that within the lower frequency domain, the equivalent of the readiness potential (RP) emerges towards the end of fixation periods between task blocks. It is observed bilaterally in the supplementary motor area (SMA) and most other regions of the cingulo-opercular action mode networks (CO-AMN). Notably, we demonstrate that this gradual and increasingly negative shift in the blood-oxygen-level-dependent (BOLD) signal is readily observable in single epochs and individual subjects.

Results

Intrinsic mode signal extraction and phase community assignment.

Functional imaging data from the two motor tasks in the fMRI HCP dataset (282 participants) were parcellated using a standardized parcellation scheme[17] (results from the final run are presented here, with differences between runs noted when relevant). Bandpass-filtered intrinsic mode signals from each parcel were extracted using the complete ensemble empirical mode decomposition (CEEMD) algorithm[18] (Suppl. Fig. S1). The analysis was restricted to two intrinsic modes that represented signal changes in a lower (0.010-0.030 Hz) and upper (0.066-0.189 Hz) frequency band. Instantaneous phase was computed at each time point and intrinsic mode time series. A phase-clustering algorithm was developed to form communities of parcels that consistently spanned a limited phase range across time and subjects (Fig. 1). The algorithm was designed such that any pairwise relationship between parcels within the same community would have an instantaneous phase difference that did not exceed a predefined threshold. The results were six major and two minor communities for both frequency bands (referred to as COMs). The major COMs were similar in size, with each covering a limited section of the unit circle (Fig. 1A), with some overlaps (Fig. 1B). Consequently, each community consistently represented approximately the same instantaneous phase, that is, sections of positive and negative shifts (deflections) of the time series (Figs. 1B, C). This provided

a simple way to compare time series across parcels, time points, and subjects. The major COMs were numbered sequentially (COM1 – COM6) as they appear in the unit circle in Fig. 1A. To assess the statistical significance of the results, the time series were phase-shuffled, and the same analysis was applied to the phase-shuffled data.

Two types of task-locked synchronization.

Our analysis identified two types of motor task-locked synchronizations across subjects for both frequency bands. The first type involved regular fluctuations in the average COM size and was observed for all the COMs. These fluctuations were time-locked to the paradigm and were independent of the specific motor task performed. The oscillatory peaks in size corresponded to the switching order of the respective COMs, reflecting the progression of the signal phase (see Figs. 1D-E). In both frequency bands, the size of COM5, which aligned approximately with the peaks of the negative shifts in the time series (Figs. 1B-C) peaked at each cue (and at the time point where the cue would have been for the fixation blocks). The remaining five major COMs exhibited peaks in size in progressive order, thereby completing a phase cycle within the period of a task block (cue + task) in both the upper and lower frequency bands.

The second type comprised brain-network-specific patterns of phase synchronization across subjects. Cross-subject phase synchronization was defined for each parcel as the percentage of subjects sharing the same community assignment at a given time, thus providing a measure of the degree to which the phase of each parcel time series was synchronized across subjects. For the upper frequency band (Fig. 2), peaks of synchronization were observed for the visual (VIS) and attention networks (ventral attention, VAN, and dorsal attention, DAN), suggesting engagement independent of the specificity of motor tasks and instead mirroring the structure and timing of the block design. To a lesser extent, this phenomenon was also observed in segments of the frontoparietal network (FPN), basal ganglia (BG), thalamus (Thal), and for a few parcels in the default mode network (DMN) (Fig. 2). Marked synchronization in COM5 aligned with the onset of the cues was followed by a sequence of phase progression delineated by

peaks in synchronization in COM6, COM1, COM2, COM4, and COM4. Thus, a complete cycle of the signal phase was achieved within the duration of a single motor task block (i.e., 3s cue + 12s task). This corresponds to a distinct peak in the power spectrum at 0.066 Hz, representing the 15s block design (Suppl. Fig. S1). Most parcels in the visual network exhibited cross-subject desynchronization during the fixation blocks (Suppl. Fig. S2A-B), indicating that the phase of the parcel time series was no longer synchronized across subjects. This desynchronization phenomenon was also observed in a few parcels of the attention networks.

In the lower-frequency band (Fig. 3), COM membership patterns were unique to each of the five motor tasks (left/right foot, left/right hand, and tongue movement). Notably, specific subcohorts of parcels within the somatomotor network (SOM) were selectively synchronized, depending on the specific body part being moved (Fig. 3). Additionally, peaks in cross-subject phase synchronization were present during fixation periods in regions associated with anticipation and planning of motor movements, a result that will be elaborated later. While the block design-induced responses resulted in cycles of cross-subject phase synchronization that were completed within a task block for the upper frequency band, the period of task-related cross-subject phase synchronization in the lower frequency band persisted over approximately two and a half blocks (33 s) (Fig. 3, Suppl. Fig. S1).

Task-locked synchronization in the somatomotor network.

Fig. 4A shows the intrinsic mode signals (lower frequency band) for five parcels in the

motor cortex directly involved in the execution of hand, foot, and tongue movements. The

corresponding progression of the cross-subject phase synchronization for each of the six

communities is shown in Fig. 4B. Importantly, the cross-subject phase synchronization in

Fig. 4B adds time-resolved information on the statistical significance of the averaged

intrinsic mode signal in Fig. 4A in terms of the percentage of the total number of subjects

aligned in phase. Statistical significance thresholds of cross-subject phase

synchronization were calculated from the phase-shuffled data (Suppl. Fig. S3). There was

a strong positive relationship between intrinsic mode signal amplitude and the maximum

(across the six communities) cross-subject phase synchronization (Suppl. Fig. S4). The

whole-brain maps shown in Fig. 4C depict the degree to which other brain areas were

correlated in terms of synchronization (across subjects) with the five seed parcels. This

shows that several regions beyond the five parcels in the motor cortex, such as the

associated somatosensory cortex and neighboring dorsal regions in the dorsal attention

network, were synchronized during motor task execution. Notably, the highest degree of

cross-subject phase synchronization was observed in the hand somatosensory cortex.

Tongue movements induced nearly equal-sized synchronizations in the associated motor

regions in both hemispheres. It also induced significant synchronization with the

orbitofrontal cortex and the temporal poles (Suppl. Fig. S7,8). Hand activation was

unilateral, whereas motor parcels of the feet exhibited weak synchronization when the

contralateral foot moved.

Patterns of whole-brain task-locked synchronization.

A whole-brain perspective of community phase synchronization during an fMRI motor

task is shown in Fig. 5. The primary visual cortex (V1) and segments of the ventral visual

stream, along with the primary somatosensory and motor areas of the tongue (more

pronounced on the right side), demonstrated the highest degree of synchronization within

the upper frequency band (Fig. 5). The somatosensory and motor regions were associated

with the maximum degree of synchronization in the lower frequency band (Fig. 5B). We computed the average correlation between the parcel community synchronization time series (mean across the six main communities and all subjects) and sorted them according to network (Figs. 5C, D).

In the upper-frequency band (Fig. 5C), the DMN and limbic networks were decoupled (more so in the left hemisphere than in the right), whereas most parcels in the remaining networks were highly correlated in terms of cross-subject synchronization. An inspection of the mean pattern of synchronization per parcel (that is, by taking the mean of the rows in Figs. 5C and 5D) revealed a distinct pattern of integration (high average correlation) between the VIS, DAN, VAN, Thal, BG, DMN (right hemisphere) networks, and most of the FPN network for the upper frequency band (Fig. 5E). Additionally, limbic and predominantly left-hemispheric DMN regions were segregated in relation to other networks.

In the lower frequency band, the cross-subject synchronization time series for the VIS, attentional (VAN, DAN), FPN, Thal, and BG networks exhibited relatively high intra-network correlations (Fig. 5D). Additionally, the between-network correlation for the Thal and BG networks was strong. Moreover, the SOM network displayed interhemispheric decoupling (Fig. 4), except for the tongue motor areas, which were strongly synchronized (Fig. 5D). The motor and somatosensory cortices, as well as portions of the DMN and visual poles, appeared decoupled from the rest of the brain, as measured by the average correlation of the cross-subject synchronization time series (Figs. 5D, F). The maximum degree of cross-subject phase synchronization exhibited a near-oscillatory pattern, with three peaks per block (Suppl. Fig. S5B).

Activity of the SMA and cingulo-opercular action mode network.

It is well known from the literature that the supplementary motor cortex (SMA), pre-SMA, anterior mid-cingulate cortex (aMCC), insula, supramarginal gyrus, putamen, and

thalamus are activated before the motor cortex [4,18]. In instances where movement is self-
paced or coupled with a cognitively demanding task, the prefrontal cortex is also engaged
[2,6]. These regions collectively constitute the cingulo-opercular action mode network (CO-
AMN), which encompasses the middle frontoparietal operculum (inferior precentral
sulcus), middle precentral sulcus, supramarginal gyrus, anterior prefrontal cortex (aPFC),
putamen, and thalamus[19]. This network is also known as a timing network in different
contexts[20].

Our analysis revealed a distinct pattern of cross-subject phase synchronization in the
bilateral SMA, right precentral sulcus, left inferior precentral sulcus, and right
supramarginal gyrus (SMG) (Fig. 6A, See also Suppl. Figs. S9-10 for regions exhibiting the
most pronounced synchronized negative shift across subjects). At the onset of the motor
task paradigm and at the conclusion of the two fixation blocks, which were interspersed
with the task blocks at irregular intervals, significant peaks in cross-subject phase
synchronization (i.e., COM5 and COM6) were observed (Fig. 6B). These corresponded to
peaks in the negative shifts in the averaged intrinsic mode signal time series (lower-
frequency band) (Fig. 6C). Notably, these negative shifts were also present in the intrinsic
mode signal time series of the individual subjects for each fixation period (Fig. 6D).
Importantly, the negative shifts in the intrinsic mode signals of the individual subejcts had
their equivalent in the raw BOLD time series (Fig. 6E), albeit with a temporal shift, such
that the maximum negative deflection was observed slightly before that of the raw BOLD
time series (Fig. 6D-E, see also Suppl. Fig. S11). The results were very similar when
investigating a subcohort (N = 70) with minimal head movement (Supp. Fig. S12).

313 Cross-subject synchronization and inter-effector regions.

315 Emerging evidence indicates that the motor cortex can be divided into effector and inter-
316 effector (IE) regions[21]. The effector regions are responsible for the execution of motor
317 movement, whereas the IE regions have been proposed to facilitate whole-body

integration and motor planning, displaying non-preferential activation during motor movements. The IE regions exhibit extensive connectivity with each other and with CO-AMN. In the parcellation utilized in this study[17], only the middle and inferior IE regions were identifiable; specifically, one parcel lateral to the tongue and one situated between the tongue and hand motor regions. Consistent with the literature, we observed that the cross-subject synchronization time series of the IEs were generally highly correlated with each another (Pearson's r range [0.59, 0.87]) (Suppl. Fig. S13). Contrary to the findings of the Gordon study[21], which examined individual subjects, our results indicated that IE regions exhibit a preference for specific motor activities. The cross-subject synchronization time series demonstrated a strong positive average correlation with tongue movement (r = 0.75, SD = 0.07), positive average correlation with hand movements (r = 0.49, SD = 0.19), and negative average correlation with foot movements (r = -0.28, SD = 0.12) (average calculated across the four effector regions and two motor regions). There was a stronger average correlation between the IE regions and the SMA network (r = 0.59, SD = 0.10) compared to the average correlation between the IE regions and the rest of the brain (r = 0.18, SD = 0.30).

Discussion

We employed a new time-resolved fMRI method to demonstrate that the equivalent of the readiness potential is present in a substantial sample of Human Connectome fMRI motor task data[15,16]. During rest (fixation) intervals between motor tasks, a slow and increasingly negative shift in the intrinsic mode BOLD signal was observed bilaterally in the SMA prior to a of forthcoming motor actions within a cued paradigm (block design). Although the negative shift was most pronounced in the SMA, it was also observed in regions largely overlapping with the cingulo-opercular action mode network (CO-AMN) (Fig. 6A, Suppl. Fig. S9), and disappeared after the cue for the upcoming task was presented. CO-AMN is associated with heightened arousal, externally focused attention, cue-specific processing, planning, and processing of interoceptive signals[19]. The SMA is part of the ventral attention network for the parcellation schema used in this study, further attesting to its involvement in attentional processes [17]. Importantly also, SMA and anterior insula

are consistently involved in both reward and loss anticipation in monetary incentive delay

tasks[22]. Regions in the CO-AMN have also been shown to be activated by processes

involving timing in the brain; therefore, they are referred to as the brain timing network[20,23].

A large meta-analysis of timing studies using fMRI showed that a right hemispheric

dominant version of CO-AMN consistently activates during timing tasks involving either

perceptual or motor components, or both[23]. This network has a topology very similar to

that of the SMA - centred network, which synchronizes during the fixation periods in our

results (Suppl. Fig. S9). The minor differences observed in the network boundaries

between the CO-AMN, the timing network, and the SMA-centred network reported here

are likely a reflection of the flexible engagement of these regions during different task

demands, mirroring real-world complexity, which recruits similar, partly overlapping

networks. For example, the PFC is not engaged in the motor tasks, which can be attributed

to the basic paradigm that lacks significant cognitive complexity [19].

CO-AMN has also been found to be strongly correlated with the recently described IE

regions[21]. We found similar, but more attenuated, correlations (Suppl. Fig. S13). This

discrepancy could be explained by methodological differences between precision

mapping in a smaller number of densely sampled individuals and the parcellated large

sample used here. Task-specific activation of some IE regions observed in our study is

likely to have the same cause.

Importantly, the negative signal shift in the CO-AMN (Fig. 6) was present in single subjects

and for single epochs, with some minor differences in timing. These results suggest that

the negative shift serves as a general indicator of active anticipation within the CO-AMN

for an upcoming motor task. It is also likely to include some degree of motor planning.

However, because the exact nature of the upcoming task is unknown during the fixation

period (which limb, which side), this planning must be general in scope; thus, no

lateralization is seen. Recently, results from an electrophysiological study showed that

the RP amplitude increased as a function of learning the optimal time to move. No

increase was observed when the timing was random, and learning was impossible. The

authors concluded that the RP reflects expectations more strongly than uncertainty [24].

Our results support the notion that learning the timing of motor movements is reflected

in the activity of the SMA and CO-AMN when motor movements are predictable. During the first of the two back-to-back fMRI motor runs (Suppl. Fig. S14), fixation periods were interspersed with three consecutive task blocks. A peak in the negative shift in cross-subject synchronization was observed towards the end of the fixation blocks. It was also observed where the third fixation block would have been placed if it had followed the same order between tasks, which it did not (see Online Methods and Suppl. Fig S17). In the second run, there were no signs of this type of expectation (Fig. 6B). This finding suggests a rapid learning effect (within one run), that fixation periods in this paradigm occurred at irregular intervals. The rapid learning of timing is well documented, as is the involvement of the SMA in timing [20,23]. In addition, in both runs, the SMA showed a clear negative shift right at the start of the task runs in the same way as at the end of the fixation period. In both runs, there was a positive shift in synchronization at the end of the last task block. We interpret this as a relaxation response, as opposed to a negative shift, indicating anticipation.

While the cross-subject phase-synchronization algorithm closely captured both positive and negative shifts in the intrinsic mode signals, in some instances, the negative shifts were an artifact of the intrinsic mode decomposition, with no clear correspondence in the raw BOLD signal. An example is the pre-cue negative shift in motor movement-specific parcels (Fig. 4, Suppl. Fig. S6). Importantly however, the negative shifts in the intrinsic mode signals of the SMA and CO-AMN during fixation periods had a clear equivalent in the raw BOLD time series, albeit slightly shifted in time (Fig. 6, Suppl. Fig. S9-12).

Our results partially reconcile previous interpretations of the RP in so far that they demonstrate that RP is a genuine phenomenon, observable not only in electrophysiological recordings but also in fMRI data, and detectable at the level of individual subjects and single epochs (Fig. 6). Its bilateral distribution suggests that it is indistinguishable from contingent negative variation (CNV) when lateralization of the upcoming motor task is unknown, supporting the hypothesis that early CNV and RP components largely represent the same type of signal manifesting in slightly different contexts[2,11]. However, the design of the task paradigm examined here does not permit

fine-tuned discrimination of potential differences in the late components of CNV and RP signals or conclusions regarding its role in the decisional process[1,10]. While the relationship between the RP and fMRI BOLD response has been investigated in a few studies, those studies have not identified the RP per se in the BOLD signal. Instead, they observed BOLD activations (positive deflections) associated with the negative potentials in the EEG [4,25].

The negative shift in the sluggish BOLD signal is naturally more protracted in time than the RP and other negative potentials observed in the neurophysiological recordings with better temporal resolution. Importantly, it appeared in the slower of the two main frequency intervals that were the focus of our analysis. This was clearly differentiated from another type of recruitment of the ventral and dorsal attention networks, which together with the visual network, exhibited task-locked synchronization in the higher-frequency band (Fig. 2). It is well known from neurophysiological recordings that periodic sensory stimulation can induce a rhythmic brain response that mirrors the stimulation frequency [26]. It has been proposed that slow cortical potentials (SCPs) that are recruited [6,26]. Given the slow nature of both SCPs and fMRI BOLD signals, they are likely to reflect the same or similar processes, albeit with a time lag[11,12]. In contrast to the regular synchronization patterns in the CO-AMN generated by the block design in the upper frequency band (Fig. 2), the slow negative shift in the CO-AMN in the lower frequency domain (Fig. 6) appears to be driven by an active anticipation process. This means that internally generated and externally presented information is processed in different frequency bands in the same region. This is in line with previous fMRI studies, which have shown differentiated processing in different frequency bands[27–29]. Our results suggest that this processing is sequential and not simultaneous for the SMA, because the peaks in cross-subject synchronization in the lower frequency band are accompanied by desynchronization in the upper frequency band (Suppl. Fig. S15). Finally, we noted previously unreported complex co-activation of the orbitofrontal cortex (OFC), ventromedial prefrontal cortex (vmPFC), and temporal poles for tongue movements involving both positive and negative shifts in the BOLD signal (Suppl. Figs. S7-8).

Taken together, the results of our time-resolved analysis of BOLD signal time courses,

based on their instantaneous phases and the corresponding phase-ruled community

membership, contribute new knowledge on the role of negative shifts in fMRI signals.

Negative BOLD signals are conventionally interpreted as "deactivations", hence "not

active." Given the results presented here, this notion seems too simplistic[30]. Instead, in

analogy with the closely related phenomenon of SCPs, that likely reflect cyclic

modulations in overall cortical excitability [31], negative shifts in the BOLD-signal does not

necessarily correspond to a passive, "non-use" process, but rather an active process of

inhibition that significantly impacts the brains' state-specific responsiveness and it is

amenable to subjective intention [30–33]. De-activation responses, notably deactivation of

the DMN during task paradigms, were noted early in the era of fMRI research [30,34,35]. These

anti-correlations between network activity were by some initially thought to be artifacts[36].

More recently, whole-brain approaches have begun to harness information present in the

entire brain simultaneously, moving away from the bias of activation versus deactivation

[32,37–40]. Our new method demonstrated cross-subject synchronization through cycles of

both positive and negative shifts in the BOLD signal, further highlighting the importance

of the whole-brain perspective. It also provides a methodological approach for

investigating anticipation and attention in a wide variety of fMRI paradigms. Given the

importance of anticipation in perception and the creation of subjective reality, as well as

its close association with prediction, it provides an exciting opportunity to investigate

these phenomena further [41–44].

Material and methods

Data

fMRI data from 381 young and healthy participants of both sexes from the HCP 1200

subject release were initially included in this study [15,45]. The two motor fMRI task runs

(block design, scanned back-to-back) used here were acquired at the end of a close to an

one-hour fMRI recording session[16]. In the first run (run A), phase-encoding was in the right-

left direction and the second (run B) phase encoding was in the left-right direction.

Subjects with head movement with a peak frame-wise displacement root mean square

(fdrms) of > 0.5 mm were excluded from each of the two runs (See Fig. 1 for fdrms time
series, run B). In the first run, 266 subjects remained, and in the second run, 262 subjects
remained. Because the cues of most motor tasks were associated with very minor
average head movements in this group, we also performed the same analysis on a subset
of subjects with very low motion (max fdrms <0.2mm) (Suppl. Fig. S16).

The second run (run B) was the focus of the main analysis. When appropriate, to highlight
important differences, results from the first run (run A) will also be included. The motor
task consisted of five movements (finger-tapping left, finger-tapping right, toe squeezing
right, toe squeezing left, and tongue movement)[16]. The duration of each block was 12 s
and was preceded by a 3 second cue. Each type of movement/block was repeated once.
Additionally, three fixation blocks (duration: 15 s) were interspersed between motor task
blocks. The ordering of movements in runs A and B was different, but some of the basic
structure was preserved (Suppl. Fig. S17).

Preprocessing and parcellation

Minimally pre-processed volume-based data provided by the HCP were used; no further
pre-processing of the data was conducted[46]. The pre-processing pipeline included
gradient unwarping, motion correction in six directions, distortion correction using a field
map, nonlinear registration to the MNI reference space using the FSL tool FNIRT, and
intensity normalization[16,46]. For cortical parcellation, the Schaefer 200 brain area
parcellation (seven networks) was used[17]. It is important to note that the parcels in the
Schaefer parcellation are not symmetrical across hemispheres. Furthermore, the
subcortical section of the Human Brainnetome Atlas[47] was used to delineate the
thalamus, basal ganglia, hippocampus, and amygdala. In total, the parcellation
comprised of 236 brain parcels. The amygdala and hippocampus parcels were assigned
to the "limbic network" in the Schaefer parcellation scheme, whereas the thalamus and
basal ganglia were treated as separate networks. The analysis therefore considered nine
separate networks: Visual (VIS), Somato-motor (SOM), Dorsal attention (DAN), Ventral
attention (VAN), Limbic (Limb), Frontoparietal (FPN), Default Mode (DMN), Thalamus
(Thal), Basal Ganglia (BG). This was the same parcellation used in previous studies [39,40].

Complex weights, instantaneous phase, and Complete Ensemble Empirical mode decomposition (CEEMD)

Time series analysis is an example in which complex numbers are ubiquitous. A real signal S can be represented in analytical (i.e. complex-valued) form using polar coordinates $S = A \times e^{i\theta}$, where A is the amplitude of the signal, theta is the angle and i the imaginary unit. The same signal can be expressed in Cartesian coordinates as $S = A(\cos(\theta) + i \times sin(\theta))$. The cosine term represents the real part of the phase angle and the sine component represents the imaginary part.

Signal phase is one of the classic measures used to assess co-activation between regions of the brain [48,49]. It has mostly been used in MEG and EEG studies and to a lesser extent in fMRI data analysis [39,40,49–52]. Phase synchronization is the alignment of the phases of two signals. This can refer to the relationship between two signal time series. In the idealized case, the phase difference between the two time series is constant over time. It can also refer to a single time point by estimating the instantaneous phase using the Hilbert transformation[53]. The assumption is that a small difference in phase translates into information exchange between regions[48]. Thus, the smaller the difference, the stronger the cooperation. An important methodological strength of phase as a measure of interregional relationships in the brain is that it quantifies relationships on a continuous spectrum of positive and negative values. This is conceptually similar to correlation, which also spans positive and negative dimensions. The instantaneous phase provides the added advantage of being time-resolved. Previous studies have shown high similarity between time-resolved changes in connectivity between BOLD-signals based on correlation and instantaneous phase measures respectively[39,54,55]. Importantly, the instantaneous phase extracted using Hilbert transform must be unambiguous. This implies that the time series cannot contain riding waves, which are typically present in real-world signals. Narrow-band pass filtering is the most common method for making a time series smooth with unambiguous phase representation at each time point. The frequency range of the BOLD-signal that contain neuronally associated information remains an open question. Some argue for a very narrow range, while there is growing evidence pointing to a wide range of meaningful frequencies within the BOLD-spectrum[51].

In fact, there seems to be task-specific segregation of network activation in different

frequency bands, which is also present to some extent during rest [27,28,56–58]. Thus, a narrow

filter is likely resulting in information loss, which may be relevant. Also, the choice of

frequency cut-off is often arbitrary.

Considering these factors, we opted to use an adaption of the empirical mode

decomposition (EMD) algorithm called Complete Ensemble Empirical Mode

Decomposition (CEEMD). The EMD is a heuristic sifting algorithm that decomposes a

signal into oscillatory modes called intrinsic mode functions [59]. In each mode the total

number of maxima and zero crossings differ by at the most one. This means that there will

be no riding waves and no ambiguity regarding phase. The waveforms are locally

approximately symmetric [59]. This, by design, forces a negative deflection prior to and after

a positive deflection (albeit it could be very minor) whether it is present in the raw signal

or not. Because the combined modes reconstruct the original signal, an imposed

deflection is compensated in the subsequent modes. To interpret the results correctly, it

is necessary to examine the original time series. To reduce the downstream effects of this

process, we first applied a broad bandpass filter ([0.003,0.3] Hz) to reduce the noise

because the (CE)EMD extracts the highest frequencies first. This also substantially

reduced mode mixing. The CEEMD is a noise-assisted development of the EMD which

results in a complete decomposition of the signal with only a small error and improved

separation of frequencies, that is, reduced mode mixing [60]. Due to edge effects, 10 time

points at the beginning and end of each time series were removed.

Phase clustering algorithm

As a starting point, we define the dense complex weighted matrix M_t where t is the time.

Dense here means that all pairwise relationships between parcels are included without

threshold.   Each entry $M_t(x,y)$ is an edge that describes the instantaneous phase

difference $\theta_{x,y}$ between parcel (node) x and y at time t. Complex weighted networks are

also an emerging research topic in the network science community, and we refer

interested readers to, for example, Böttcher and Porter 2024 and Tian and Lambiotte 2024, for more theoretical support[61,62]. This algorithm is greedy and has a hierarchical structure. It optimizes two main criteria: integration (criterion one) and segregation (criterion two). Integration in this context means that *all* pairwise relationships within the same community have a maximum phase difference defined by threshold one, $thr1 = \cos(\theta_1)$. We investigated six different choices for θ_1: $\frac{5\pi}{6}, \frac{\pi}{2}, \frac{\pi}{3}, \frac{\pi}{6}, \frac{\pi}{9}, \frac{\pi}{18}$. In this context, an integrated community refers to a community that exhibits *phase consistency*, meaning that the maximum phase difference between any pair of nodes within the community is less than θ_1. Segregated means that *all* pairwise relationships between nodes in two different communities have a *minimum phase difference* defined by threshold two, $thr2 = \cos(\theta_2)$, where $\theta_2 = -\theta_1$. There are also two secondary, more generous thresholds, to account for scenarios where lesser degrees of segregation are present: $thr2b = \cos(\theta_{2b})$, where $\theta_{2b} = \frac{\pi}{2}$ and $thr2c = thr1$. $Thr2b$ is only relevant when $\theta_1 < \theta_{2b}$ or in situations where no cluster is formed based on $thr2$. $Thr2c$ is applied only when segregation occurred within this range $[\theta_1, \theta_{2b}]$. Thresholds beyond $thr2c$ are superfluous because it equals a scenario when there is only a single phase-consistent cluster based on $thr1$. In each iteration, the algorithm aims to identify a pair of maximally segregated (criterion two), phase-consistent (integrated) communities (criterion one) if a minimum segregation of *at least* $Thr2c (= thr1)$ exists in the data. In each iteration, to harmonize communities across time points, the first community is based on the parcel with the smallest instantaneous phase value among the unassigned parcels, that is $-\pi$ in the idealized case. The second community is defined based on the parcel most segregated from the seed parcel of the first community. The first community is subsequently expanded by finding groups of parcels that fulfil both criteria one and two. If multiple such groups exist, the group with the smallest mean phase difference with the seed parcel is chosen. The second community was expanded in the same manner. Communities formed beyond the first iteration were sorted based on their leading and lagging relations with the communities in the first iteration. The imaginary component (*sin*) of the phase difference between two communities provides information on which community is leading, and which is lagging. Let ϑ_1 be the instantaneous phase of *Com1* and ϑ_2 that of *ComX*. Then, if $\sin(\vartheta_x - \vartheta_1) > 0$, *ComX* is leading in relation to *Com1*.

However, this is a truth with modification since $\vartheta_1 = \vartheta_1 + 2\pi$ in which case Com1 would

be ahead of *ComX*. If a leading community already exists, a secondary leading community

is formed. If instead $\sin(\vartheta_x - \vartheta_1) < 0$, *Com X* lags in relation to *Com1*.

The results for $\theta_1 = \frac{\pi}{3}$ are presented in the main text. The algorithm was designed for

community comparison between multiple iterations of a complex weighted network,

where the distribution of phases was assumed to be similar across iterations. In the case

of time series, the iterations are discrete time points. The algorithm can be adapted to

situations in which the phase distribution varies widely between the iterations. This is

done using the iteration with the largest distribution as the reference.

Phase shuffled data

To test whether the results were not merely driven by chance, the algorithm was applied

to surrogate data. Because the key property of the investigation was phase, phase

shuffled data were used as surrogates[63]. The shuffling of the phases preserves the power

spectrum of a time course while destroying the temporal order of the phase. The intrinsic

mode time series of the second run (run B) was shuffled. Thus, the shuffled data provided

statistical thresholds for significant results. Because of edge effects resulting from the

phase shuffling process, 30 time points were excluded at the beginning and end of each

time series, instead of only 10, which were excluded from the real data.

622
623
624
625

References

1. Schurger, A., Hu, P. 'Ben', Pak, J. & Roskies, A. L. What Is the Readiness Potential?

Trends Cogn. Sci. **25**, 558–570 (2021).

2. Brunia, C. H. M., Van Boxtel, G. J. M. & Böcker, K. B. E. *Negative Slow Waves as Indices*

of Anticipation: The Bereitschaftspotential, the Contingent Negative Variation, and

the Stimulus-Preceding Negativity. (Oxford University Press, 2011).

doi:10.1093/oxfordhb/9780195374148.013.0108.

3. KORNHUBER, H. H. & DEECKE, L. [CHANGES IN THE BRAIN POTENTIAL IN

VOLUNTARY MOVEMENTS AND PASSIVE MOVEMENTS IN MAN: READINESS

POTENTIAL AND REAFFERENT POTENTIALS]. *Pflugers Arch. Gesamte Physiol.*

Menschen Tiere **284**, 1–17 (1965).

4. Cunnington, R., Windischberger, C. & Moser, E. Premovement activity of the pre-

supplementary motor area and the readiness for action: Studies of time-resolved

event-related functional MRI. *Hum. Mov. Sci.* **24**, 644–656 (2005).

5. Schultze-Kraft, M. *et al*. The point of no return in vetoing self-initiated movements.

Proc. Natl. Acad. Sci. **113**, 1080–1085 (2016).

6. Armstrong, S., Sale, M. V. & Cunnington, R. Neural Oscillations and the Initiation of

Voluntary Movement. *Front. Psychol.* **9**, 2509 (2018).

7. Di Russo, F. *et al*. Beyond the "Bereitschaftspotential": Action preparation behind

cognitive functions. *Neurosci. Biobehav. Rev.* **78**, 57–81 (2017).

8. Shibasaki, H. & Hallett, M. What is the Bereitschaftspotential? *Clin. Neurophysiol.*

117, 2341–2356 (2006).

9. Libet, B., Gleason, C. A. & Wright, E. W. TIME OF CONSCIOUS INTENTION TO ACT IN

RELATION TO ONSET OF CEREBRAL ACTIVITY (READINESS-POTENTIAL).

10. Schurger, A., Sitt, J. D. & Dehaene, S. An accumulator model for spontaneous neural

activity prior to self-initiated movement. *Proc. Natl. Acad. Sci.* **109**, (2012).

11. Schmidt, S., Jo, H.-G., Wittmann, M. & Hinterberger, T. 'Catching the waves' – slow

cortical potentials as moderator of voluntary action. *Neurosci. Biobehav. Rev.* **68**,

639–650 (2016).

12. He, B. J. & Raichle, M. E. The fMRI signal, slow cortical potential and consciousness.

Trends Cogn. Sci. **13**, 302–309 (2009).

13. Jo, H.-G., Wittmann, M., Borghardt, T. L., Hinterberger, T. & Schmidt, S. First-person

approaches in neuroscience of consciousness: Brain dynamics correlate with the

intention to act. *Conscious. Cogn.* **26**, 105–116 (2014).

14. Jo, H.-G., Hinterberger, T., Wittmann, M., Borghardt, T. L. & Schmidt, S. Spontaneous

EEG fluctuations determine the readiness potential: is preconscious brain activation

a preparation process to move? *Exp. Brain Res.* **231**, 495–500 (2013).

15. Glasser, M. F. *et al.* The Human Connectome Project's neuroimaging approach. *Nat.*

Neurosci. **19**, 1175–1187 (2016).

16. Barch, D. M. *et al.* Function in the human connectome: Task-fMRI and individual

differences in behavior. *NeuroImage* **80**, 169–189 (2013).

17. Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from

Intrinsic Functional Connectivity MRI. *Cereb. Cortex* **28**, 3095–3114 (2018).

18. Deiber, M.-P., Honda, M., Ibañez, V., Sadato, N. & Hallett, M. Mesial Motor Areas in

Self-Initiated Versus Externally Triggered Movements Examined With fMRI: Effect of

Movement Type and Rate. *J. Neurophysiol.* **81**, 3065–3077 (1999).

19. Dosenbach, N. U. F., Raichle, M. E. & Gordon, E. M. The brain's action-mode

network. *Nat. Rev. Neurosci.* **26**, 158–168 (2025).

20. Bader, F. & Wiener, M. Neuroimaging Signatures of Metacognitive Improvement in

Sensorimotor Timing. *J. Neurosci.* JN-RM-1789-22 (2023)

doi:10.1523/JNEUROSCI.1789-22.2023.

21. Gordon, E. M. *et al.* A somato-cognitive action network alternates with effector

regions in motor cortex. *Nature* **617**, 351–359 (2023).

22. Oldham, S. *et al.* The anticipation and outcome phases of reward and loss

processing: A neuroimaging meta-analysis of the monetary incentive delay task.

Hum. Brain Mapp. **39**, 3398–3418 (2018).

23. Naghibi, N. *et al.* Embodying Time in the Brain: A Multi-Dimensional Neuroimaging

Meta-Analysis of 95 Duration Processing Studies. *Neuropsychol. Rev.* **34**, 277–298

(2024).

24. Travers, E., Friedemann, M. & Haggard, P. The Readiness Potential reflects planning-

based expectation, not uncertainty, in the timing of action. *Cogn. Neurosci.* **12**, 14–

27 (2021).

25. Nguyen, V. T., Breakspear, M. & Cunnington, R. Reciprocal Interactions of the SMA

and Cingulate Cortex Sustain Premovement Activity for Voluntary Actions. *J.*

Neurosci. **34**, 16397–16407 (2014).

26. Duecker, K. *et al.* Challenges and Approaches in the Study of Neural Entrainment. *J.*

Neurosci. **44**, e1234242024 (2024).

27. Chang, C. & Glover, G. H. Time–frequency dynamics of resting-state brain

connectivity measured with fMRI. *NeuroImage* **50**, 81–98 (2010).

28. Thompson, W. H. & Fransson, P. The frequency dimension of fMRI dynamic

connectivity: Network connectivity, functional hubs and integration in the resting

brain. *NeuroImage* **121**, 227–242 (2015).

29. Hipp, J. F. & Siegel, M. BOLD fMRI Correlation Reflects Frequency-Specific Neuronal

Correlation. *Curr. Biol.* **25**, 1368–1374 (2015).

30. He, B. J. Spontaneous and Task-Evoked Brain Activity Negatively Interact. *J. Neurosci.*

33, 4672–4682 (2013).

31. Vanhatalo, S. *et al.* Infraslow oscillations modulate excitability and interictal

epileptic activity in the human cortex during sleep. *Proc. Natl. Acad. Sci.* **101**, 5053–

5057 (2004).

32. Buonomano, D. V. & Maass, W. State-dependent computations: spatiotemporal

processing in cortical networks. *Nat. Rev. Neurosci.* **10**, 113–125 (2009).

33. Garrison, K. A. *et al.* Effortless awareness: using real time neurofeedback to

investigate correlates of posterior cingulate cortex activity in meditators' self-report.

Front. Hum. Neurosci. **7**, (2013).

34. Fransson, P. Spontaneous low-frequency BOLD signal fluctuations: An fMRI

investigation of the resting-state default mode of brain function hypothesis. *Hum.*

Brain Mapp. **26**, 15–29 (2005).

35. Fox, M. D. *et al.* The human brain is intrinsically organized into dynamic,

anticorrelated functional networks. *Proc. Natl. Acad. Sci.* **102**, 9673–9678 (2005).

36. Chai, X. J., Castañón, A. N., Öngür, D. & Whitfield-Gabrieli, S. Anticorrelations in

resting state networks without global signal regression. *NeuroImage* **59**, 1420–1428

(2012).

37. Gonzalez-Castillo, J. *et al.* Whole-brain, time-locked activation with simple tasks

revealed using massive averaging and model-free analysis. *Proc. Natl. Acad. Sci.*

109, 5487–5492 (2012).

38. Deco, G., Tononi, G., Boly, M. & Kringelbach, M. L. Rethinking segregation and

integration: contributions of whole-brain modelling. *Nat. Rev. Neurosci.* **16**, 430–439

(2015).

39. Strindberg, M., Fransson, P., Cabral, J. & Ådén, U. Spatiotemporally flexible

subnetworks reveal the quasi-cyclic nature of integration and segregation in the

human brain. *NeuroImage* **239**, 118287 (2021).

40. Fransson, P. & Strindberg, M. Brain network integration, segregation and quasi-

periodic activation and deactivation during tasks and rest. *NeuroImage* **268**, 119890

(2023).

41. Clark, A. Whatever next? Predictive brains, situated agents, and the future of

cognitive science. *Behav. Brain Sci.* **36**, 181–204 (2013).

42. Buzsáki, G., Peyrache, A. & Kubie, J. Emergence of Cognition from Action. *Cold*

Spring Harb. Symp. Quant. Biol. **79**, 41–50 (2014).

43. De Lange, F. P., Heilbron, M. & Kok, P. How Do Expectations Shape Perception?

Trends Cogn. Sci. **22**, 764–779 (2018).

44. Lee, C. S., Aly, M. & Baldassano, C. Anticipation of temporally structured events in

the brain. *eLife* **10**, (2021).

45. Uğurbil, K. *et al.* Pushing spatial and temporal resolution for functional and diffusion

MRI in the Human Connectome Project. *NeuroImage* **80**, 80–104 (2013).

46. Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human

Connectome Project. *NeuroImage* **80**, 105–124 (2013).

47. Fan, L. *et al.* The Human Brainnetome Atlas: A New Brain Atlas Based on

Connectional Architecture. *Cereb. Cortex* **26**, 3508–3526 (2016).

48. Varela, F., Lachaux, J.-P., Rodriguez, E. & Martinerie, J. The brainweb: Phase

synchronization and large-scale integration. *Nat. Rev. Neurosci.* **2**, 229–239 (2001).

49. Glerean, E., Salmi, J., Lahnakoski, J. M., Jääskeläinen, I. P. & Sams, M. Functional

Magnetic Resonance Imaging Phase Synchronization as a Measure of Dynamic

Functional Connectivity. *Brain Connect.* **2**, 91–101 (2012).

50. Cabral, J. *et al*. Cognitive performance in healthy older adults relates to spontaneous

switching between states of functional connectivity during rest. *Sci. Rep.* **7**, 5135

(2017).

51. Niazy, R. K., Xie, J., Miller, K., Beckmann, C. F. & Smith, S. M. Spectral characteristics

of resting state networks. in *Progress in Brain Research* vol. 193 259–276 (Elsevier,

2011).

52. Ponce-Alvarez, A. *et al*. Resting-State Temporal Synchronization Networks Emerge

from Connectivity Topology and Heterogeneity. *PLOS Comput. Biol.* **11**, e1004100

(2015).

53. Honari, H., Choe, A. S. & Lindquist, M. A. Evaluating phase synchronization methods

in fMRI: A comparison study and new approaches. *NeuroImage* **228**, 117704 (2021).

54. Pedersen, M., Omidvarnia, A., Zalesky, A. & Jackson, G. D. On the relationship

between instantaneous phase synchrony and correlation-based sliding windows for

time-resolved fMRI connectivity analysis. *NeuroImage* **181**, 85–94 (2018).

55. Preti, M. G., Bolton, T. A. & Van De Ville, D. The dynamic functional connectome:

State-of-the-art and perspectives. *NeuroImage* **160**, 41–54 (2017).

56. Billings, J. C. W. *et al*. Instantaneous brain dynamics mapped to a continuous state

space. *NeuroImage* **162**, 344–352 (2017).

57. Yaesoubi, M., Allen, E. A., Miller, R. L. & Calhoun, V. D. Dynamic coherence analysis

of resting fMRI data to jointly capture state-based phase, frequency, and time-

domain information. *NeuroImage* **120**, 133–142 (2015).

58. He, B. J., Zempel, J. M., Snyder, A. Z. & Raichle, M. E. The Temporal Structures and

Functional Significance of Scale-free Brain Activity. *Neuron* **66**, 353–369 (2010).

59. Huang, N. E. *et al.* The empirical mode decomposition and the Hilbert spectrum for

nonlinear and non-stationary time series analysis. *Proc. R. Soc. Lond. Ser. Math.*

Phys. Eng. Sci. **454**, 903–995 (1998).

60. Torres, M. E., Colominas, M. A., Schlotthauer, G. & Flandrin, P. A complete ensemble

empirical mode decomposition with adaptive noise. in *2011 IEEE International*

Conference on Acoustics, Speech and Signal Processing (ICASSP) 4144–4147 (IEEE,

Prague, Czech Republic, 2011). doi:10.1109/ICASSP.2011.5947265.

61. Böttcher, L. & Porter, M. A. Complex networks with complex weights. *Phys. Rev. E*

109, 024314 (2024).

62. Tian, Y. & Lambiotte, R. Spreading and Structural Balance on Signed Networks. *SIAM*

J. Appl. Dyn. Syst. **23**, 50–80 (2024).

63. Lancaster, Iatsenko, Pidde, Ticcinelli, & Stefanovska. Surrogate data for hypothesis

testing in physical systems. *Phys. Rep.* **2018**, (2018).

790   Figures

791
792

Fig. 1. *Community assignment based on the instantaneous phase of motor task fMRI*

data. A) Color wheel in which each instantaneous signal phase corresponds to a color. B)

Distribution of the instantaneous phase (radians) across subjects, parcels, and time

points) per community (plot is cut at π). For color coding, the median value of the phase

distribution in each community was used based on (A). C) Correspondence between an

intrinsic mode signal time course (lower frequency band) and community membership

(COM1-6) for a single subject and parcel. For example, COM5 approximately represents

the maximum points of negative deflection (or signal shift) over time. The average fraction

of parcels (out of 236) for each community across time was computed for the upper

(0.066-0.189 Hz) (D) and lower (0.030-0.010 Hz) frequency bands (E).

Fig. 2. *Network-specific cross-subject phase synchronization (n=282) in six communities (COM1-6) during an fMRI motor task (results shown here refer to signal changes in the upper frequency range of: 0.066-0.189 Hz). The color bar reflects the percentage of the total number of subjects (282 subjects) that were synchronized in each community at each point in time.*

Fig. 3. *Network-specific cross-subject phase synchronization (n=282) in six communities (COM1-6) during an fMRI motor task (results shown here refer to signal changes in the lower-frequency range: 0.030-0.010 Hz). The color bar reflects the percentage of the total number of subjects (282 subjects) that were synchronized in each community at each point in time.*

Fig. 4. *Averaged (n=282) intrinsic mode signals (lower frequency band, shaded area = ± 1 SD) **(A)** and the fraction of subjects that were phase-synchronized into COM1-6 across time for the fMRI motor task **(B)** for five parcels in the motor cortex (SOM network) responsible for movement execution: hands (LH_SomMot_6, RH_SomMot_14), feet (LH_SomMot_12, RH_SomMot_14), and tongue (RH_SomMot_18). The correlation maps depicted in **(C)** show the average degree of correlation (Pearson's r × 100) between the seed parcels in the motor cortex and the rest of the brain in terms of the percentage of synchronized subjects (correlation analysis performed separately for each COM and then averaged). The y-axis in (B) represents the fraction (in percentage) of subjects being synchronized. The horizontal black lines denote the mean degree of cross-subject*

synchronization (16.67%) for the phase-shuffled version of the data (dotted lines = ± 2 SD).

Only the left hemisphere motor cortex representation of the tongue is shown (right side

strongly correlated, Pearsons' r = 0.97). See Suppl. Fig. S6 for the averaged raw BOLD

signals. The amygdala, hippocampus, and basal ganglia parcels are not shown here.

Fig. 5*. Panels **A** (upper frequency band) and **B** (lower frequency band) show the maximum*

degree of cross-subject synchronization across time and communities for all the parcels

(282 subjects). The network (with left hemispheric parcels initially grouped, followed by

right hemispheric parcels) and parcel pair-specific presentations of the mean correlation

of community synchronization (averaged separately for COM1-6 and all subjects) over

*time are shown in panels **C** and **D**, respectively. In panels **E** and **F**, the mean (i.e., by*

*averaging across rows of the correlation matrices shown in panels **C** and **D**) correlation of*

synchronization for all parcels (Pearson's r × 100) is shown. The amygdala, hippocampus,

and basal ganglia parcels are not shown.

Fig. 6. *Cross-subject phase synchronization and subject-specific intrinsic mode signals*

for the left supplementary motor area (LH_SalVentAttn_Med_3) during the fMRI motor

*task. Panel **A** depicts the mean degree of correlation of cross-subject phase*

synchronization between the left hemisphere SMA parcel and all other brain regions

*(Pearson's r × 100), showing substantial overlap with the CO-AMN network[19]. Panel **B***

shows the cross-subject phase synchronization time series (COM1-6; black line

represents the mean of the results based on phase-scrambled data, dotted lines = ± 2

SD). The average intrinsic mode signal time-course (lower frequency band) is shown in

*panel **C**, shaded area = ± 1SD) and the subject-specific time series are shown in panel **D**.*

*For comparison, the raw BOLD signal intensity time-courses are shown in panel **E**.*

856
857

Supplement

864 ***Suppl. Fig. S1.*** *Power-spectrum for four intrinsic mode functions (IMFs) (final run). IMF1*

865 *[0.1318, 0.4613] Hz,* ***IMF2 (upper frequency band) [0.0659, 0.1876] Hz,*** *IMF 3 **(lower***

866 ***frequency band) [0.0304, 0.0963] Hz***, *IMF4 [0.0152, 0.1064] Hz. Only results from upper*

867 *and lower frequency bands are presented.*

885
886

A.   **B.**

890
891 **C.**
892

893 ■ RIGHT HAND ■ RIGHT FOOT ■ LEFT HAND ■ LEFT FOOT ■ TONGUE ■ FIXATION ■ CUE
894

895 ***Suppl. Fig. S2. A)*** *and* ***B)*** *example of a visual network parcels cross-subject*

896 *synchronization, upper frequency band desynchronizes during fixation blocks (grey),*

897 *second run.* ***C)*** *Visual parcel that doesn't show significant cross-subject synchronization.*

898 *X-axis is time in minutes. Y-axis is percent of total subjects.*

901
902

903
904 **A.** **B.**

905 ***Suppl. Fig. S3***. *Distribution of cross-subject synchronization across all six communities,*

906 *time-points and subjects (count) based on the phase scrambled data. Due to edge effects*

907 *of the phase scrambling 15 timepoints were cropped at the start and finish.* ***A)*** *upper*

908 *frequency domain,* ***B)*** *lower frequency band*.

909
910
911
912

913
914
915 **A.** **B.**
916

917 ***Suppl. Fig. S4.*** *Scatterplots showing the relationship between intrinsic mode signal*

918 *amplitude (absolute value) and maximum (among the six communities) cross subject*

919 *phase synchronization for all of the 236 parcels and 264 time points in the time-series.* ***A)***

920 *Upper frequency band, Pearsons' r = 0.60,* ***B)*** *Lower frequency band, Pearsons' r = 0.66.*

922
A.   **B.**

■ RIGHT HAND ■ RIGHT FOOT ■ LEFT HAND ■ LEFT FOOT ■ TONGUE ■ FIXATION ■ CUE

Suppl. Fig. S5. Maximum cross-subject phase synchronization across the six main
communities as a function of time in the first (Run A) and second (Run B) run A) Upper
frequency band, B) lower frequency band. Synchronization is generally higher in the lower
compared to the upper frequency band. The lower frequency band exhibit a near
oscillatory pattern with three peaks per block.

932
933 *A. Right hand (LH_SomMot_12)*   *B. Left hand (RH_SomMot_14)*

935 *C. Right foot (LH_SomMot_15)*   *D. Left foot (RH_SomMot_18)*

E. *Right foot (RH_SomMot_7)*   **F.** *Left foot (LH_SomMot_6)*

■ RIGHT HAND ■ RIGHT FOOT ■ LEFT HAND ■ LEFT FOOT ■ TONGUE ■ FIXATION ■ CUE

Suppl. Fig. S6. *Averaged (n=282) raw BOLD time-series for the six movement specific parcels in the motor cortex (shaded area = ± 1 SD)). Tongue motor movement parcels from both hemisphere motor cortex are included (E, F). Comparing the averaged raw BOLD with the lower frequency intrinsic modes in Fig. 4 suggests that the pronounced negative deflections seen in the latter prior to task execution in this case are artefacts of the CEEMD-algorithm (See Methods). X-axis is time in minutes. Y-axis is arbitrary units.*

A. First Run   **B. Second Run**

Suppl. Fig. S7. *Cross-subject phase synchronization correlation maps for tongue movement, lower frequency band.   A) First Run. B) Second Run (main results). Tongue movements induced synchronization in the temporal pole, orbitofrontal cortex (OFC), ventromedial prefrontal cortex (vmPFC), and specific regions of the basal ganglia, notably in the left ventral caudate nucleus. These synchronizations were antiphase, or*

segregated, relative to the tongue motor areas, except for the left temporal pole in the second run. During the two experimental runs, tongue movements engaged the cerebral hemispheres in a partially opposite manner. In the initial run, in-phase synchronization was observed in the right temporal pole rather than in the left, and above-baseline synchronization with the basal ganglia was absent. Note the opposite correlation pattern in the ventral temporal lobes and left lateral OFC between the two runs.

Suppl. Fig. S8. *Example from to parcels with opposite signal changes during tongue movement (second run). Left column: Left medial temporal pole (LH_Limbic_TempPole_3). Right column: Right caudal orbito-frontal cortex (RH_Limbic_OFC_2). First row: mean raw BOLD signal (+/- 1 SD in grey). Second row: Mean intrinsic mode signal (+/- 1 SD in grey). Third row: Cross-subject synchronization time-series.*

Suppl. Fig. S9. *The parcels most strongly connected with the left hemispheric SMA: right SMA, right middle and inferior precentral sulcus, left inferior precentral sulcus and right supramarginal gyrus (SMG). Strong connection was defined as the combination of a) mean correlation of synchronization with SMA r > 0.75 and b) minimum peak in cross-subject synchronization > 40% of total subjects.*

Suppl. Fig. S10. *Cross-subject synchronization time-series for the parcels in Suppl. Fig S9, i.e. the parcels with the strongest connection with the left hemispheric SMA.* **A1.** *LH_SomMot_5,* **A2.** *LH_SomMot_14,* **A3.** *LH_SalVentAttn_ParOper_2,* **A4.** *LH_SalVentAttn_FrOperIns_4,* **B1.** *RH_SalVentAttn_TempOccPar_1,* **B2.** *RH_SalVentAttn_PrC_1,* **B3.** *RH_SalVentAttn_FrOperIns_4,* **B4.** *RH_SalVentAttn_Med_3.*

RIGHT HAND **RIGHT FOOT** **LEFT HAND** **LEFT FOOT** **TONGUE** **FIXATION** **CUE**

Suppl. Fig. S11. *Left SMA (LH_SalVentAttn_Med_3) raw averaged (across subjects)*

BOLD time-series. X-axis is time in minutes. Y-axis is arbitrary units.

A. **B.**

RIGHT HAND **RIGHT FOOT** **LEFT HAND** **LEFT FOOT** **TONGUE** **FIXATION** **CUE**

Suppl. Fig. S12. *Low motion subjects (N = 70, maximum frame-wise displacement root*

mean square (fdrms) < 0.2 mm), individual left SMA time series, A) Intrinsic mode time-

series lower frequency band, B) Raw BOLD time-series

1076

A. **B.**

C. **D.**

Suppl. Fig S13. *Inter-effector regions average correlation of the six coss-subject phase synchronization timeseries with the rest of the brain (Range is Pearson's correlation x100).*

A) Left inferior IE, B) Right inferior IE, C) Left middle IE, D) Right middle IE.

Suppl. Fig. S14. *First run (A), Cross-subject phase synchronization time-series for left SMA (LH_SalVentAttn_Med_3). Note the blue peak at the second last task. This is the location the last fixation block would have been following the pattern of the first two fixation blocks. X-axis is time in minutes. Y-axis is percent of total subjects.*

Suppl. Fig. S15. *Left SMA (LH_SalVentAttn_Med_3), upper frequency band. De-synchronizations are noted at the end of the two first (i.e. full-length) fixation blocks. The last fixation block is cut due to edge artefacts of the band-pass filtering and intrinsic mode decomposition (see Methods). X-axis is time in minutes. Y-axis is percent of total subjects.*

1113

A.   **B.**

■ RIGHT HAND ■ RIGHT FOOT ■ LEFT HAND ■ LEFT FOOT ■ TONGUE ■ FIXATION ■ CUE

Suppl. Fig. S16. Mean (across subjects) fdrms time series for the second run (run B). A) thr < 0.5mm (N= 262), B) thr < 0.2 mm (N = 70). ± 1 SD in gray shades.

■ RIGHT HAND ■ RIGHT FOOT ■ LEFT HAND ■ LEFT FOOT ■ TONGUE ■ FIXATION ■ CUE

Suppl. Fig. S17. Block design for the two runs: first run (A) and second run (B).