AMPK signaling is dysregulated in human tendinopathy and loss of AMPKα1 leads to cell, matrix and mechanical dysfunction in mouse Achilles tendon

LeeAnn A. Hold[1,2], Nicole Migotsky[2], Syeda N. Lamia[2], Steph S. Steltzer[1,2], Sydney Grossman[2], Jessica Chen[2], Seung-Ho Bae[2], Paige Cordts[2], Tessa Phillips[2,3], Matthew J. O'Meara[4,5], Carol Davis[1], Susan V. Brooks[1,6], Moeed Akbar[7], Neal Millar[7], Megan L. Killian[1,2], Adam C. Abraham[2,8*]

[1] Molecular and Integrative Physiology, University of Michigan, Ann Arbor, MI, United States
[2] Orthopedic Surgery, University of Michigan, Ann Arbor, MI, United States
[3] University of Toledo College of Medicine, Toledo, Ohio, United States
[4] Gilbert S. Omenn Department of Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, United States
[5] Medicinal Chemistry, University of Michigan, Ann Arbor, MI, United States
[6] Biomedical Engineering, University of Michigan, Ann Arbor, MI, United States
[7] Institute of Infection, Immunity and Inflammation, University of Glasgow, Glasgow, UK
[8] Lead contact
*Correspondence: adaabrah@med.umich.edu

SUMMARY

Tendinopathy is a tendon disorder that is caused by the failure to self-repair and has many pathological characteristics such as disorganized ECM and decreased cell viability. We have identified a possible target to combat these changes, AMPK, an energy stress sensor that was shown to maintain intracellular homeostasis. Through bulk RNA-sequencing of healthy and tendinopathic tendons from humans we have identified a novel finding of downregulation of AMPK signaling in the tendinopathic samples which suggests AMPK plays a role in tendon homeostasis. Our studies utilizing a conditional knock-out of Prkaa1 in tendon in mice showed that loss of AMPK results in degenerative ECM, impaired biomechanical properties and increased cellular senescence in Achilles tendon through the lifespan. Additionally, we found that exercise can delay senescence onset independent of AMPK. These findings highlight the importance of energy metabolism in tendon health which will assist in understanding the onset and progression of tendinopathy.

KEYWORDS

Extracellular matrix, AMPKa1, Prkaa1, senescence, biomechanics, exercise

INTRODUCTION

Tendons are poorly vascularized tissues rich in extracellular matrix (ECM) that transmit force from muscle to bone. The mechanical demand placed on these tissues requires a precise balance of metabolic processes to maintain functional homeostasis[1,2]. Tendinopathy, is a tendon disorder and while the precise causes and characteristics remain to be elucidated, it is associated with results in pain, swelling, and diminished biomechanical properties. is a clinical problem that affects ~3.5 million people in the US[3]. Risk factors for tendinopathy include age, metabolic disorders and limited exercise[3]. Tendinopathy is caused by failure of tendon to self-repair and is characterized by degenerative ECM, decreased cell viability, and poor biomechanical function[4]. Currently, treatment for tendinopathy has limited non-surgical treatment options[3], therefore elucidating targets for a druggable therapy will improve current clinical limitations. Previously, Wu et. al established a promising molecular mechanism for treatment of diabetic induced tendinopathy through the AMP-activated protein kinase (AMPK) pathway[5]. The AMPK pathway is an intracellular stress response activated in times of nutrient deficiency, and exercise to regulate energy balance and maintain homeostasis[6]. In response to a decrease in the ATP:AMP ratio, AMPK is

phosphorylated and redirects metabolism to stimulate catabolic processes while simultaneously
decreasing anabolic processes[6]. While AMPK is a critical regulator of homeostasis in other tissues[7,8] as
well as a potential regulator of ECM remodeling in musculoskeletal disease[9], little is known about how
AMPK regulates energy balance and homeostasis in tendon. A key pathological characteristic of
tendinopathy includes dysregulation of homeostasis, such as matrix disorganization and decreased cell
viability[3]. Additionally, regulation of metabolic mechanisms play a role in tendon healing and
degeneration[10,11] establishing the potential for metabolic modulators as therapeutic targets for
tendinopathy. Therefore, the potential in targeting AMPK motivates the need to define the role of AMPK in
tendon homeostasis.

The basal metabolic rate of tendon is relatively low compared to other musculoskeletal tissues, possibly
related to the high number of quiescent cells present in tendon during homeostasis[12]. The quiescent
phenotype presents during times of limited nutrients and is coupled with a reversible cell cycle arrest. It is
believed that tendon fibroblasts (TFs) become predominantly quiescent in the early stages of
maturation[13]. Grinstein et. al showed in mice that after post-natal day 30 mitotic activity declines
significantly and few cells continue to divide through the remainder of life[13]. Regulation of quiescence
through AMPK has been shown to be age dependent in muscle stem cells[14,15].With age, AMPK activity is
dampened resulting in a push towards senescence through nuclear translocation of cyclin inhibitors[14].
Furthermore, activation of AMPK restored cell survival, through upregulation of autophagy and reduced
nuclear translocation of cyclin inhibitors, in aged skeletal muscle stem cells[14]. These trends may be
consistent in tendon, and in a study by Dai et. al, it was shown that that activation of the AMPK signaling
cascade through metformin reduced the effects of aging *in vivo* and *in vitro*[16]. Using tendon progenitor
cells *in vitro*, metformin reduced senescence markers p16 and p21[16]. Additionally, they showed metformin
reduced the aging phenotype by inhibiting senescence marker p16 and the formation of heterotopic
ossification in 10 month-old rats [16]. Taken together, these data implicate AMPK as a possible target to
regulate cellular fate in tendon and prevent age-associated functional decline.

AMPK is a heterotrimeric complex composed of α, β and γ subunits[6]. There are two α subunits, α1 and
α2: AMPKα1 is ubiquitously expressed in all tissues while AMPKα2 is dominant in skeletal muscle, heart,
and liver[17,18]. Reduction of AMPKα1 has been associated with musculoskeletal diseases such as
osteoarthritis (OA)[19]. For example, loss of AMPKα1 leads to accelerated progression and increased
severity of OA[9,19]. However, while dysregulation of AMPKα1 in some degenerative musculoskeletal
diseases is known, the mechanisms that underlie its role in tendinopathy remain unclear.

In this study, we identified that the AMPK signaling pathway is transcriptionally suppressed in patients
with tendinopathy. Additionally, we found that AMPK is necessary for tendon homeostasis by mediating
expression of *Prkaa1* in TFs. We used Cre-lox to target deletion of *Prkaa1*, the gene that encodes
AMPKα1, in tendon progenitors in mice. This targeted deletion led to normal development of tendons;
however, we saw dysregulation of cell cycle and ECM biological processes in the transcriptome at 1
month of age (1M). By 3 months of age (3M), we found that tendons lacking *Prkaa1* had impaired
mechanical properties and upregulation of senescence associated marker, p21. With increased age,
tendons lacking *Prkaa1* developed expansive ectopic calcification and had elevated markers of
senescence associated p16 and p21. Voluntary wheel running was beneficial in tendons of *Prkaa1-*
deficient mice and led to increased ECM organization and suppressed p16 and p21. Our results
demonstrate AMPKα1 is necessary for tendon homeostasis and exercise, an energy demanding process,
which can delay senescence onset independent of AMPK α1 suggesting energy balance is crucial for
cellular viability in tendon.

RESULTS

Metabolic pathways and AMPK signaling are dysregulated in tendinopathy

Using bulk RNA-sequencing, we compared healthy hamstring tendon to diseased rotator cuff tendons
from human donors and found a downregulation of multiple genes associated with the mitochondrial

electron transport chain and energy production, suggesting an overall dysregulation in metabolism (Fig. 1,
Table S1 for patient demographics). Enriched biological processes were related to energy homeostasis,
response to nutrients, and cytokine signaling (Fig. 1A). Additionally, enriched molecular functions (Fig 1B)
and KEGG pathways were also related to AMPK activity and signaling (Fig. 1C), respectively, with KEGG
pathways also enriched for regulation of energy metabolism (i.e. glycolysis and oxidative
phosphorylation), metabolic pathways and cytokine signaling (Fig. 1C). When analyzing the differentially
expressed genes (DEGs) produced from the KEGG pathway results, the AMPK signaling pathway was
primarily driven by six DEGs, of which 5 were downregulated in tendinopathic tendons (Fig. 1D,E), with
phosphofructokinase (*PFKP*) as the only upregulated gene (Fig. 1D,E). The top 10 hits for DEGs
associated with metabolic pathways were also downregulated with tendinopathy, including genes
associated with the mitochondrial electron transport chain (Fig. 1D,E). Specifically, tendinopathy samples
had downregulated mitochondrial encoded NADH dehydrogenase 1 (*MT-ND1*; Fig. 1D,E), which
regulates mitochondrial complex one and the NAD^+/NADH ratio[20]. A decrease in the NAD^+/NADH ratio
has been associated with decreased AMPK activation and increased production of the senescence
associated secretory phenotype (SASP)[21]. Additionally, DEGs associated with the cytokine-cytokine
receptor interaction pathway included activation of multiple interleukins (e.g. *IL1*, *IL13*, and *IL36*; Fig.
1D,E), another characteristic of the SASP[22]. Considering these findings, we sought to identify the role on
AMPK in the mouse Achilles tendon in an age-dependent manner.

Figure 1: Metabolic pathways and AMPK signaling are dysregulated in human tendinopathy.
(A-C) Enrichment quantification of differentially expressed genes (DEGs) from bulk RNA sequencing of
human tendinopathic samples compared to healthy control tendons. (A. Enriched GO biological
processes; B. Enriched GO molecular functions; C. Enriched Kegg pathways.)
(D) Signaling diagram from individual DEGs from KEGG pathways (Gray: DEGs in AMPK signaling
pathway; Pink: DEGs from metabolic pathways; Purple: DEGs in cytokine-cytokine receptor interaction
pathway; blue arrow representative of downregulated DEGs, orange arrow representative of upregulated
DEGs)

123 (E) Relative expression of DEGs from signaling diagram (log2 fold change) from human tendinopathic
124 samples compared to healthy control tendons. n= 7 samples/group. Adjusted p-value: p<0.05,
125 log2FC±1.5
126
Prkaa1^{Scx-Cre} cKO mouse tendons showed dysregulation of cell cycle processes, increased
calcification and impaired biomechanical properties.

In the Achilles tendons of mice at 1M of age, we identified 621 upregulated and 1008 downregulated
genes in Prkaa1Scx-Cre cKO females compared to age-matched WT females using bulk RNA sequencing,
yet transcriptional differences were not observed in male mice at 1M or 3M of age. Prkaa1Scx-Cre cKO
tendons from female mice at 1M had enriched biological processes related to cell cycle regulation, matrix
organization and response to nutrients compared to WT females (Fig. 2A). When comparing cell cycle
DEGs, we found a robust upregulation of cell cycle arrest markers in tendons from female Prkaa1Scx-Cre
cKO mice such as *Ccngd3,* the gene that encodes cyclin D3*,* and *Cdkn1a*, the gene that encodes p21
(Fig. 2B). Finally, when comparing metabolism DEGs we identified a significant upregulation of *Bop1*, a
known inhibitor of cell cycle progression*,* in cKO tendons compared to WT at 1M[23] (Fig.2B). Cellular
senescence is defined as terminal growth arrest in cells even if stimulated by mitogens, however they are
metabolically and synthetically active[24]. Although there is no clear pathway to senescence, early markers
of senescence include changes in glucose metabolism and increased cyclin kinase activity. These
findings suggest that Achilles tendons in Prkaa1Scx-Cre cKO mice exhibit a transcriptional profile that
supports the onset of senescence[25,26]. Senescent cells are also known to contribute to ECM remodeling
[27]. When comparing tendon and matrix related DEGs, we found *Scx* was significantly upregulated in
female Prkaa1Scx-Cre cKO tendons, while *Lox*, *Loxl3* and both *Col1a1* and *Col1a2* were downregulated
compared to WT female tendons (Fig. 2C). A decrease in *Scx* expression in tendon indicates a shift
towards a more mature tendon state[28]. Our results showing an increase in Scx mRNA expression in the
cKO cells suggest while the WT cells are transitioning into a more mature tendon the cKO cells
transcriptionally resemble a progenitor tendon cell. At the protein level, LOX catalyzes the formation of
crosslinks in collagen and is essential for the formation of collagen fibril, decreased *Lox* gene expression
has been associated shown to reduce the ability to withstand load in *in vitro* tendon constructs [29]. We
found the Prkaa1Scx-Cre cKO Achilles tendons had impaired mechanical function compared to WT controls.
First, we found the maximum failure load of Prkaa1Scx-Cre cKO Achilles tendons trended downward at 1M
(p = 0.0509) and was significantly reduced at 3M compared to WT controls (Fig. 2D,E), however the
maximum load equilibrated by 9M of age (Fig. 2D). Prkaa1Scx-Cre cKO Achilles tendons were significantly
larger (cross-sectional area, CSA) compared to WT controls at 9M (Fig. 2F). When normalizing the load
to CSA we found peak mechanical stress of Prkaa1Scx-Cre cKO Achilles tendons was also significantly
reduced at 3M and 9M of age compared to WT controls (Fig. 2G). When comparing at 3% stain the
tangent modulus of Prkaa1Scx-Cre cKO Achilles tendons were significantly reduced at 3M of age (Table
S2). Finally, we saw no changes to stiffness between WT and cKO tendons at any age we investigated
(Table S2). The functional changes we observed were likely not due to impaired muscle function, as we
saw no differences in muscle force production in Prkaa1Scx-Cre cKO mice at 3M (Tables S3-9). We next
analyzed the Prkaa1Scx-Cre cKO Achilles tendons histologically and identified positive silver nitrate-stained
areas we will report as lesions (Fig.2H), Quantification of silver nitrate-stained lesions at 9M of age
showed that Prkaa1Scx-Cre cKO Achilles tendons had a significantly greater percent of positive lesion area
compared to WT controls (Fig. 2J) and X-ray confirmed these regions appeared as radio-opaque ectopic
calcification (Fig. 2I). Cells contributing to ectopic calcification can exhibit nuclear swelling due to
increased stress caused by aberrant calcium levels[30]. In line with this, we found the cell nuclei in lesions
at 9M were rounder compared to non-lesion "healthy" tendon within Prkaa1Scx-Cre cKO Achilles tendons as
well as in WT tendons (Fig. 2K). Finally, we found that collagen organization, measured using quantitative
polarized light imaging (qPLI) of fiber birefringence on histological sections, was not different between
Prkaa1Scx-Cre cKO tendons and WT controls at 1, 3 and 9 months of age (Fig. S1A,B). In summary we
identified dysregulation of matrix interaction signaling as well as markers of pre-mature senescence
therefore, we sought to identify how loss of *Prkaa1* effects ECM specific adhesion and if Prkaa1Scx-Cre
cKO tendons exhibited additional senescence markers with age.

A. Biological Processes

Response to stress
Response to glucose
Glucose metabolic process
Glucose homeostasis
ECM organization
Collagen catabolic process
Cellular response to oxidative stress
Cell adhesion mediated by integrin

Enrichment

0 1 2 3 4 5 6

#DEGs
10 20 30

B. Cell Cycle Diagram

Cdkn1a → Senescence

Ccnd3

Cell Cycle Progression

Bop1

C. Tendon Maturation Diagram

Mesenchymal Stem Cell

Osteoblast Progenitor — *Runx2*

Osteoblast — *Sparc*

Tendon Progenitor — *Scx* *Sox9*

Immature tendon cells — *Mkx*

Mature tendon cells — *Col1a1* *Lox* *Dcn* *Col1a2*

D. Max load (N) — 1M, 3M (****), 9M

E. Force (N) vs Displacement (mm) — 1M, 3M, WT, cKO

F. CSA (mm²) — 1M, 3M, 9M (*)

G. Max Stress (MPa) — 1M, 3M (**), 9M (**), WT, cKO

H. WT, cKO — Male, Female

I. WT, cKO

J. Lesion Area at 9M (%) — WT, cKO (*)

K. Nuclear Circularity — WT, cKO (****, ****)

Figure 2: Phenotypic characterization of Prkaa1[Scx-Cre] cKO Achilles tendons.
(A) Quantification of GO biological processes found during bulk RNA sequencing in 1M female Prkaa1[Scx-
Cre] cKO mice compared to WT controls.
(B) Signaling diagram created from cell cycle related DEGs in Prkaa1[Scx-Cre] cKO mice compared WT
controls. (blue arrow representative of downregulated DEGs, orange arrow representative of upregulated
DEGs)
(C) Signaling diagram created from tendon maturation related DEGs in Prkaa1[Scx-Cre] cKO mice compared
WT controls. (blue arrow representative of downregulated DEGs, orange arrow representative of
upregulated DEGs, gray bay representative of no significant change)
n=3 samples/group. Adjusted p-value: Adjusted p-value: $p<.05$, log2FC±1.5
(D-G) Biomechanical outcomes from 1, 3 and 9M WT and Prkaa1[Scx-Cre] cKO Achilles tendons (D.
Maximum Load before failure; E. Representative Force vs Displacement curve for 1M and 3M WT and
Prkaa1[Scx-Cre] cKO Achilles tendons F. Cross-sectional area (CSA); G. Maximum stress).
n =6 - 10/group, * $p<.05$, ** $p<0.01$; two-way ANOVA with Sidak correction.
(H) Representative X-ray images of dystrophic calcification, highlighted by white arrows in the 9M WT and
cKO Achilles tendons.
(I) Representative silver nitrate-stained images in 9M WT and Prkaa1[Scx-Cre] cKO Achilles tendons. Scale
bar = 200µm.
(J) Quantification of positive silver nitrate-stained lesions in 9M WT and Prkaa1[Scx-Cre] cKO Achilles
tendons. n =6/group, * $p<0.05$.
(K) Quantification of nuclear circularity 9M WT and Prkaa1[Scx-Cre] cKO Achilles tendons. ((H) = healthy, (L)
= lesion; n =3/group, ****$p<.0001$; by paired t-test.
198
199   ***Prkaa1[Scx-Cre] cKO Achilles tendon fibroblasts have altered ECM specific adhesion and show an***
200   ***increase in senescence associated markers.***
201
To compare cellular adhesion between Prkaa1[Scx-Cre] cKO and WT TFs we plated cells on an array of 36
matrix spots comprised of individual and mixtures of 9 ECM component proteins and measured nuclear
counts after 24h as a proxy for adhesion. Across all conditions, spot counts ranged from 0 to 408 with a
median of 95. To analyze the drivers of this variation, we fit a Bayesian general linear regression models
with genotype and matrix proteins as covariates and using weakly informative priors and compared the
posterior distributions over regression weights. As a baseline model, we assumed no statistical interaction
between genotype and the matrix proteins and since the outcomes are counts, we used a Poisson link
function. We found that Prkaa1[Scx-Cre] cKO TFs were less adherent than WT TFs (Fig. 3A). Additionally,
collagen (COL) 1, COL6, fibronectin and vitronectin were more permissive while COL3 was less
permissive for adhesion regardless of genotype (Fig. 3A). Considering an interaction model where
genotype and matrix interact suggested there was a preference of Prkaa1[Scx-Cre] cKO TF adhesion for Col5
and tropoelastin. Additionally, Prkaa1[Scx-Cre] cKO TFs showed a decreased preference for COL1 and
laminin (Fig. 3B). There is increasing evidence showing the composition of the ECM potentiates cellular
senescence[31]. We identified senescence in Prkaa1[Scx-Cre] cKO tail TFs using Beta-Galactosidase staining
(Fig. 4A). Two cell-cycle inhibitors that are commonly regarded as markers of senescent cells are the
cyclin-dependent kinase inhibitors p21 and p16[32]. Here, we used immunohistochemistry of the Achilles
tendon and identified a significant increase in p21 and p16 in Prkaa1[Scx-Cre] cKO mice compared to WT
controls at 9M of age (Fig. 4B-E). It is important to note that even in the 9M WT samples we saw >10% of
the tendon cell population expressed both p16/p21. Cells expressing senescent markers can play critical
roles in tissue regeneration and homeostasis[33], suggesting that while we saw a significant increase in
senescent markers in Prkaa1[Scx-Cre] cKO Achilles tendons, senescence may be a normal process in
tendon homeostasis. Moreover, given the ability to remove senescent cells through exercise, we decided
to use voluntary wheel running to determine the effects of exercise on senescence in the Achilles tendon
independent of AMPK. However, given our cells ability to respond to nutrient depletion is impaired we
were unsure how these tendons would respond to exercise. Therefore, we gave WT and Prkaa1[Scx-Cre]
cKO access to voluntary running wheels.

Figure 3: Tendon fibroblasts from Prkaa1^{Scx-Cre} cKO mice have impaired ECM specific adhesion.

(A) Representative images of Hoechst staining used for COLI and COL III.

(B-C) Posterior distributions with median and 80% confidence intervals of Bayesian regression models quantified from nuclear counts (B. Baseline; C. Interaction; Grey bar represents median number of all nuclear counts regardless of genotype) n = 3/group.

Figure 4: Prkaa1$^{Scx-Cre}$ cKO tendon fibroblasts had increased expression of senescence markers in vitro and in vivo.
(A) Representative Beta-Galactosidase stain of WT and Prkaa1$^{Scx-Cre}$ cKO tail TFs *in vitro* after five days in culture. Scale bar = 200µm. n=3/group.
(B and C) Representative IHC images in 9M WT and Prkaa1$^{Scx-Cre}$ cKO Achilles tendons of (B) p21 and (C) p16 staining using DAB.
(D and E) Quantification of (D) positive p21 cells and (E) positive p16 cells in 9M WT and Prkaa1$^{Scx-Cre}$ cKO Achilles tendons. Scale bar = 500µm. n =6/group (pink = female; blue = male), ***p<0.001, ****p<0.0001; two-way ANOVA with Sidak correction.

Exercise increases organization and reduces the onset of senescence in Prkaa1$^{Scx-Cre}$ cKO Achilles tendons.

We identified a significant linear relationship between distance ran and days with wheels in the WT mice (r^2=0.7295) during voluntary wheel running (WR) but not the Prkaa1$^{Scx-Cre}$ cKO mice (r^2=0.02202) showing the WT mice ability to adapt (Fig. 5A). The body weight of Prkaa1$^{Scx-Cre}$ cKO mice was significantly lower than WT mice in cage activity (CA) group but was rescued with WR (Fig. 5B). Additionally, fasting blood glucose (FBG) was significantly elevated in the Prkaa1$^{Scx-Cre}$ cKO WR mice compared to CA mice at 3M of age (Fig. 5C). Voluntary wheel running can influence the metabolic activity in rats where *Gapdh* was shown to be significantly downregulated as well as an increase in overall food intake compared to controls[34]. Using bulk RNA sequencing of Achilles tendons, we found 109 upregulated and 78 downregulated genes with WR compared to CA in WT tendons. In WT tendons, enriched biological processes were related to protein phosphorylation, cell differentiation, and positive regulation of gene expression and angiogenesis (Fig. 5D). When comparing DEGs from cell differentiation processes, we found upregulation of *Csrp3* (Fig. 5E). Additionally, when comparing DEGs related to phosphorylation, we saw upregulation of *Sbk3* and *Sbk2* and downregulation of *Bmpr1b*. Protein phosphorylation, the most common post-translational modification, utilizes ATP and is representative of good energy balance within a cell[35] suggesting voluntary wheel running promotes energy homeostasis in tendons. In Prkaa1$^{Scx-Cre}$ cKO tendons, we found 169 upregulated and 272 downregulated genes with WR compared to CA. Enriched biological processes included positive regulation of gene expression, inflammatory response, and apoptosis (Fig. 5F). When comparing DEGs from inflammatory response we again found upregulation of *Csrp3* as well as *Cul3* (Fig.5G). When looking at the positive regulation of gene expression we found upregulation of *Itgb3*, *Itga3* and *Bmpr1b* (Fig.5G). Integrin expression is upregulated with increased mechanical signaling, thought to be necessary to transmit mechanical stimuli to the cells during loading[36]. When looking at the structure of the tendon we found a significant increase in tendon organization in the WR Prkaa1$^{Scx-Cre}$ cKO mice when comparing to the WT WR and a trend upward when comparing to the Prkaa1$^{Scx-Cre}$ cKO CA group (p = 0.0505; Fig. 5H) However, we found no significant change in tendon alignment between the groups (Fig. 5I). Immunohistochemistry of Achilles tendons revealed a significant decrease in senescence-associated p21 with WR compared to CA in Prkaa1$^{Scx-Cre}$ cKO mice (Fig. 5J). Finally, we found WR significantly suppressed senescence-associated p16 regardless of genotype (Fig. 5J).

Figure 5: Effects of exercise on Prkaa1$^{Scx-Cre}$ cKO mice.
(A) Quantification of distance run for WT and Prkaa1$^{Scx-Cre}$ cKO mice.
(B) Quantification of body weight for WT and Prkaa1$^{Scx-Cre}$ cKO WR and CA mice.
(C) Quantification of fasting blood glucose (FBG) mg/dL for wheel running groups for WT and Prkaa1$^{Scx-Cre}$ cKO WR and CA mice. n =6-12/group, females = pink; males = blue. * $p<0.05$, ** $p<0.01$; two-way ANOVA with Sidak correction.

(D) Quantification of GO biological processes found during bulk RNA sequencing in WT WR mice
compared to CA controls.
(E) Quantification of the log2 fold change of DEGs from WT mice from biological processes WR mice
compared to CA controls (Cell differentiation; Phosphorylation).
(F) Quantification of GO biological processes found during bulk RNA sequencing in Prkaa1[Scx-Cre] cKO
mice WR mice compared to CA controls.
(G) Quantification of the log2 fold change of DEGs in WR Prkaa1[Scx-Cre] cKO mice from biological
processes (Inflammatory response; Positive regulation of gene expression). n= 3-6 samples/group.
Adjusted p-value: Adjusted p-value: $p < .05$, log2FC±1.5
(H and I) Quantification of alignment for WT and Prkaa1[Scx-Cre] cKO WR and CA mice (H. Degree of linear
polarization (DoLP); I. Standard Deviation Angle of Polarization (AOP)), females = pink; males = blue
(J) Representative IHC images of p16 and p21 in CA and WR Prkaa1[Scx-Cre] cKO Achilles tendons. Scale
bar = 500µm.Quantification of p16 positive cells. Quantification of positive p21 cells. n =6/group, * $p < 0.05$,
** $p < 0.001$, ***$p < 0.0001$; two-way ANOVA with Sidak correction.
(K) Quantification of positive p16 cells. n =6/group, females = pink; males = blue. * $p < 0.05$, ** $p < 0.01$,
***$p < 0.001$; two-way ANOVA with Sidak correction.
(L) Quantification of positive p21 cells. n =6/group, females = pink; males = blue. * $p < 0.05$, ** $p < 0.01$,
***$p < 0.001$; two-way ANOVA with Sidak correction.
300

DISCUSSION

Defining the role of energy metabolism in the context of tendon health and disease is critical for the advancement of non-surgical therapeutics for patients with tendinopathy. While the role AMPK has been established as a regulator of ECM organization and homeostasis in other musculoskeletal tissues[7–9,19], the role of AMPK in tendon homeostasis and tendinopathy has previously been unclear. Here we have demonstrated that AMPK signaling is dysregulated in human tendinopathy patients and *Prkaa1* is necessary to maintain homeostasis in adult mouse Achilles tendon. These data demonstrated that loss of *Prkaa1* resulted in premature senescence as well as accelerated onset of ectopic calcification. Additionally, we found that exercise delays the onset of senescence in a *Prkaa1*-independent manner suggesting energy usage is essential for the maintenance of tendon health.

To the best of our knowledge, we are the first to identify dysregulation of the AMPK signaling pathway in tendinopathy patients. We identified upregulation of genes related to SASP production in tendinopathy patients and these trends are consistent when evaluating other data sets from tendinopathic patients[37]. In a study by Zhu et. al, screening for biomarkers of tendinopathy revealed common processes dysregulated in tendinopathy were related to the cell cycle, mitochondria function and pro-inflammatory responses as well as mTOR and NF-κB signaling pathways[37]. Additionally, other musculoskeletal diseases can be driven by senescence and the SASP[38–40]. In osteoarthritis, it is speculated that synovial macrophages, fibroblasts and osteoclasts promote SASP production, leading to an inflammatory response that enhances disease progression[39,40]. Taken together, our findings, along with previous insights into musculoskeletal disease, suggest progression of tendinopathic disease may be driven by the presence of senescent fibroblasts and maintained through an increased proinflammatory response.

We identified the formation of ectopic calcification in Prkaa1[Scx-Cre] cKO Achilles tendons, especially males. Surprisingly, our bulk RNA sequencing results showed an upregulation of both *Runx2* and *Dlx5* in female, but not male, Prkaa1[Scx-Cre] cKO tendons. The relationship between AMPK and RUNX2 can vary depending on cell type[41–43]. In osteoblasts, AMPK can regulate Runx2 post-transcriptionally and promote osteogenesis[41]. However, in vascular mesenchymal stem cells, AMPK can downregulate Runx2 expression, and loss of AMPKα1 can promote osteoblastic differentiation[43]. Spatially, we noticed the presence of calcification near the peritenon, suggesting the possibility of cell migration into the tendon core before the onset of calcification. Peritenon cells have been identified as being more stem-like than tendon core cells, expressing several stem cell markers[44]. Additionally, peritenon cells are capable of undergoing osteogenic differentiation *in vitro*[44]. Taken together, TFs and peritenon cells may respond similarly to mesenchymal stem cells with loss of AMPK, pushing them toward an osteogenic state. Future directions will explore the role of the peritenon in the formation of ectopic calcification and how migration and trans-differentiation of these cells may be induced by SASP. The formation of calcification within soft tissues has been linked to increase of senescent cells[16,45,46]. Here we identified loss of *Prkaa1* results in both premature senescence and ectopic calcification in the mouse Achilles tendon. Our data supports proposed mechanisms in which an increase p21 responds to a replicative related process (e.g DNA damage) which precedes the increase in p16, while both together correspond to late senescent cells and the presence of CDK inhibition and cell cycle arrest[47]. Our data also supports previous findings in tendon suggesting that modulation of the AMPK/mTOR signaling cascade via metformin delays the accumulation of senescent cells, cell morphological changes, and calcification[16,48]. Surprisingly, we found that while we had no sex differences in our senescence associated marker expression, we had distinct differences in calcification area between our male and female mice. Future directions will explore estrogen as a protector of calcification in the female tendon as well as identify the potential of the SASP to induce calcification and additional therapeutics to delay onset of senescence.

Integrins regulate the onset of senescence in fibroblasts[49] AMPK is a known regulator of β-1 integrin[50]. Consequently, loss of AMPK in fibroblasts can increase cell spreading as well as fibronectin fiber formation[50]. Here, we found transcriptional changes of multiple integrins in both the 1M timepoint as well as with exercise in Prkaa1[Scx-Cre] cKO tendons. Additionally, we found that TFs from Prkaa1[Scx-Cre] cKO tails

have impaired ECM substrate-specific adhesion. Different ECM substrates have unique integrin binding properties, and taken together these data suggest that AMPK activity is responsive to different ECM substrates and integrin accessibility.

The reduction of senescent cells through exercise has been seen in fibrotic disease models[51,52]. Here we found that voluntary wheel running delayed the onset of senescence markers, p16 and p21, as well as increased cell cycle progression markers (i.e. Regulator of cell cycle progression (*Rgcc*)) in Prkaa1[Scx-Cre] cKO tendons. In addition to regulating cell cycle progression, *Rgcc* may have protective effects against fibrosis, and stimulation of *Rgcc* correlated with enrichment of ECM organization pathways[53]. These trends correlate with our findings that exercise induced a significant increase in *Rgcc* gene expression as well as increased collagen alignment and integrin expression in Prkaa1[Scx-Cre] cKO tendons.

In summary, these data show that AMPK, which is dysregulated in tendinopathy, is necessary for the maintenance of homeostasis and mechanical function in the mouse Achilles tendon. Additionally, we found exercise has a protective effect against senescence independent of AMPK and promotes energy homeostasis in tendon. These findings highlight the importance of energy balance metabolism in tendon health and overall improves our understanding of the onset and progression of tendinopathy as well as identification of potential treatment options for tendon disease.

Limitations of the study

Although our data parallels other bulk-RNA sequencing of human Achilles tendinopathy, our cohort of patients was small with only seven per group focusing on the rotator cuff. Additionally, despite conducting bulk-RNA sequencing at 1M and 3M on female and male mice, we only observed large transcriptional differences in our female 1M mice. We speculate either our timepoint was too late to capture similar differences in the male mice or the phenotype was driven by post translational or other modifications. Future work with proteomics could assist in divulging these differences.

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Adam Abraham (adaabrah@med.umich.edu).

Materials availability
This study did not generate new unique reagents.

Data and code availability
- Bulk-seq data have been deposited at GEO at GEO: accession number and are publicly available as of the date of publication.
- All raw data will be deposited to Deep Blue Data

ACKNOWLEDGMENTS
University of Michigan Center for Cellular Plasticity and Organ Design Emerging Scholars program (ACA); National Science Foundation Graduate Research Fellowship DGE2241144 to LH; National Institutes of Health R01AR079367 to MLK; Michigan Integrative Musculoskeletal Health Core Center P30 AR069620 to MLK; University of Michigan Rackham Graduate Student Research Grant to LH; University of Michigan Integrative Musculoskeletal Health Core Pilot and Feasibility Grant to ACA; National Science Foundation CAREER 144448 to MLK; Library preparation and next-generation sequencing was carried out in the Advanced Genomics Core at the University of Michigan. We are grateful for histological support from Emma Snyder-White and Carol Whitinger (through the Michigan Integrative Musculoskeletal Health Center Core). We are grateful for thoughtful discussions about tendon biomechanics with Sarah Calve at the University of Colorado.

AUTHOR CONTRIBUTIONS

397   Conceptualization L.A.H, M.L.K, A.C.A; methodology, L.A.H., M.O., S.L., S.V.B., M.L.K, A.C.A.
398   Investigation, L.A.H., N.M., J.C., S.S.S., P.C., C.D., S-H.B., S.V.B., S.G., S.L., T.P., M.O., M.A., N.A.;
399   writing—original draft, L.A.H., N.M., M.L.K., A.C.A.; writing—review & editing, L.A.H., N.M., J.C., S.S.S.,
400   P.C., S-H.B., C.D., S.G.,S.L., T.P., M.O., S.V.B., M.A., N.M., M.L.K., A.C.A; funding acquisition, L.A.H,
401   M.L.K., and A.C.A..; resources, M.L.K. and A.C.A; supervision, M.L.K. and A.C.A.

SUPPLEMENTAL INFORMATION

Document S1. Figures S1 and Table S1 -S2

Figure S1. Quantification of alignment for WT and Prkaa1[Scx-Cre] cKO WR and CA animals, A) degree of linear polarization (DoLP); B) standard deviation of the angle of polarization (AOP).

Table S1. Patient Demographics for Bulk-RNA Sequencing

Table S2. Biomechanical Properties of Murine Achilles Tendons in wildtype (WT) and *Prkaa1[Scx-Cre]mice*

Table S3. EDL muscle function in 3M wildtype (WT) and *Prkaa1[Scx-Cre]mice*

Table S4. Soleus muscle function in 3M wildtype (WT) and *Prkaa1[Scx-Cre]mice*

Table S5. Gastrocnemius muscle function in 3M wildtype (WT) and *Prkaa1[Scx-Cre]mice*

Table S6. EDL muscle response (frequency sweep) of 3M wildtype (WT) and *Prkaa1[Scx-Cre]mice*

Table S7. Soleus muscle response (frequency sweep) in 3M wildtype (WT) and *Prkaa1[Scx-Cre]mice*

Table S8. Gastrocnemius muscle response to nerve stimulation (frequency sweep) in 3M wildtype (WT) and *Prkaa1[Scx-Cre]mice*

Table S9. Gastrocnemius muscle response with direct muscle stimulation (frequency sweep) in 3M wildtype (WT) and *Prkaa1[Scx-Cre]mice*

REFERENCES

1. Winnicki, K., Ochała-Kłos, A., Rutowicz, B., Pękala, P.A., and Tomaszewski, K.A. (2020). Functional anatomy, histology and biomechanics of the human Achilles tendon — A comprehensive review. Annals of Anatomy - Anatomischer Anzeiger *229*, 151461. https://doi.org/10.1016/j.aanat.2020.151461.

2. Montagna, C., Svensson, R.B., Bayer, M.L., Rizza, S., Maiani, E., Yeung, C.-Y.C., Filomeni, G., and Kjær, M. (2022). Autophagy guards tendon homeostasis. Cell Death Dis *13*, 402. https://doi.org/10.1038/s41419-022-04824-7.

3. Millar, N.L., Silbernagel, K.G., Thorborg, K., Kirwan, P.D., Galatz, L.M., Abrams, G.D., Murrell, G.A.C., McInnes, I.B., and Rodeo, S.A. (2021). Tendinopathy. Nat Rev Dis Primers *7*, 1. https://doi.org/10.1038/s41572-020-00234-1.

4. Fouda, M.B., Thankam, F.G., Dilisio, M.F., and Agrawal, D.K. (2017). Alterations in tendon microenvironment in response to mechanical load: potential molecular targets for treatment strategies. Am J Transl Res *9*, 4341–4360.

5. Wu, Y.-F., Wang, H.-K., Chang, H.-W., Sun, J., Sun, J.-S., and Chao, Y.-H. (2017). High glucose alters tendon homeostasis through downregulation of the AMPK/Egr1 pathway. Sci Rep *7*, 44199. https://doi.org/10.1038/srep44199.

6. Herzig, S., and Shaw, R.J. (2018). AMPK: guardian of metabolism and mitochondrial homeostasis. Nat Rev Mol Cell Biol *19*, 121–135. https://doi.org/10.1038/nrm.2017.95.

7. Zong, H., Ren, J.M., Young, L.H., Pypaert, M., Mu, J., Birnbaum, M.J., and Shulman, G.I. (2002). AMP kinase is required for mitochondrial biogenesis in skeletal muscle in response to chronic energy deprivation. Proc. Natl. Acad. Sci. U.S.A. *99*, 15983–15987. https://doi.org/10.1073/pnas.252625599.

8. Myers, R.W., Guan, H.-P., Ehrhart, J., Petrov, A., Prahalada, S., Tozzo, E., Yang, X., Kurtz, M.M., Trujillo, M., Gonzalez Trotter, D., et al. (2017). Systemic pan-AMPK activator MK-8722 improves glucose homeostasis but induces cardiac hypertrophy. Science *357*, 507–511. https://doi.org/10.1126/science.aah5582.

9. Zhou, S., Lu, W., Chen, L., Ge, Q., Chen, D., Xu, Z., Shi, D., Dai, J., Li, J., Ju, H., et al. (2017). AMPK deficiency in chondrocytes accelerated the progression of instability-induced and ageing-associated osteoarthritis in adult mice. Sci Rep *7*, 43245. https://doi.org/10.1038/srep43245.

10. Izumi, S., Otsuru, S., Adachi, N., Akabudike, N., and Enomoto-Iwamoto, M. (2019). Control of glucose metabolism is important in tenogenic differentiation of progenitors derived from human injured tendons. PLoS ONE *14*, e0213912. https://doi.org/10.1371/journal.pone.0213912.

11. Patel, S.H., Yue, F., Saw, S.K., Foguth, R., Cannon, J.R., Shannahan, J.H., Kuang, S., Sabbaghi, A., and Carroll, C.C. (2019). Advanced Glycation End-Products Suppress Mitochondrial Function and Proliferative Capacity of Achilles Tendon-Derived Fibroblasts. Sci Rep *9*, 12614. https://doi.org/10.1038/s41598-019-49062-8.

12. Ackerman, J.E., Best, K.T., Muscat, S.N., and Loiselle, A.E. (2021). Metabolic Regulation of
Tendon Inflammation and Healing Following Injury. Curr Rheumatol Rep *23*, 15.
https://doi.org/10.1007/s11926-021-00981-4.

13. Grinstein, M., Dingwall, H.L., O'Connor, L.D., Zou, K., Capellini, T.D., and Galloway, J.L.
(2019). A distinct transition from cell growth to physiological homeostasis in the tendon.
eLife *8*, e48689. https://doi.org/10.7554/eLife.48689.

14. White, J.P., Billin, A.N., Campbell, M.E., Russell, A.J., Huffman, K.M., and Kraus, W.E.
(2018). The AMPK/p27Kip1 Axis Regulates Autophagy/Apoptosis Decisions in Aged
Skeletal Muscle Stem Cells. Stem Cell Reports *11*, 425–439.
https://doi.org/10.1016/j.stemcr.2018.06.014.

15. McKay, L.K., and White, J.P. (2021). The AMPK/p27Kip1 Pathway as a Novel Target to
Promote Autophagy and Resilience in Aged Cells. Cells *10*, 1430.
https://doi.org/10.3390/cells10061430.

16. Dai, G., Li, Y., Zhang, M., Lu, P., Zhang, Y., Wang, H., Shi, L., Cao, M., Shen, R., and Rui,
Y. (2023). The Regulation of the AMPK/mTOR Axis Mitigates Tendon Stem/Progenitor Cell
Senescence and Delays Tendon Aging. Stem Cell Rev and Rep *19*, 1492–1506.
https://doi.org/10.1007/s12015-023-10526-0.

17. Stapleton, D., Mitchelhill, K.I., Gao, G., Widmer, J., Michell, B.J., Teh, T., House, C.M.,
Fernandez, C.S., Cox, T., Witters, L.A., et al. (1996). Mammalian AMP-activated Protein
Kinase Subfamily. Journal of Biological Chemistry *271*, 611–614.
https://doi.org/10.1074/jbc.271.2.611.

18. Woods, A., Azzout-Marniche, D., Foretz, M., Stein, S.C., Lemarchand, P., Ferré, P.,
Foufelle, F., and Carling, D. (2000). Characterization of the Role of AMP-Activated Protein
Kinase in the Regulation of Glucose-Activated Gene Expression Using Constitutively Active
and Dominant Negative Forms of the Kinase. Molecular and Cellular Biology *20*, 6704–
6711. https://doi.org/10.1128/MCB.20.18.6704-6711.2000.

19. June, R.K., Liu-Bryan, R., Long, F., and Griffin, T.M. (2016). Emerging role of metabolic
signaling in synovial joint remodeling and osteoarthritis. Journal Orthopaedic Research *34*,
2048–2058. https://doi.org/10.1002/jor.23420.

20. Lin, X., Zhou, Y., and Xue, L. (2024). Mitochondrial complex I subunit MT-ND1 mutations
affect disease progression. Heliyon *10*, e28808.
https://doi.org/10.1016/j.heliyon.2024.e28808.

21. Nacarelli, T., Lau, L., Fukumoto, T., Zundell, J., Fatkhutdinov, N., Wu, S., Aird, K.M.,
Iwasaki, O., Kossenkov, A.V., Schultz, D., et al. (2019). NAD+ metabolism governs the
proinflammatory senescence-associated secretome. Nat Cell Biol *21*, 397–407.
https://doi.org/10.1038/s41556-019-0287-4.

22. Kumari, R., and Jat, P. (2021). Mechanisms of Cellular Senescence: Cell Cycle Arrest and
Senescence Associated Secretory Phenotype. Front. Cell Dev. Biol. *9*, 645593.
https://doi.org/10.3389/fcell.2021.645593.

23. Strezoska, Ž., Pestov, D.G., and Lau, L.F. (2002). Functional Inactivation of the Mouse
Nucleolar Protein Bop1 Inhibits Multiple Steps in Pre-rRNA Processing and Blocks Cell
Cycle Progression. Journal of Biological Chemistry *277*, 29617–29625.
https://doi.org/10.1074/jbc.M204381200.

24. Shtutman, M., Chang, B.-D., Schools, G.P., and Broude, E.V. (2017). Cellular Model of p21-
Induced Senescence. Methods Mol Biol *1534*, 31–39. https://doi.org/10.1007/978-1-4939-
6670-7_3.

25. Ogrodnik, M. (2021). Cellular aging beyond cellular senescence: Markers of senescence
prior to cell cycle arrest *in vitro* and *in vivo*. Aging Cell *20*, e13338.
https://doi.org/10.1111/acel.13338.

26. Gorgoulis, V., Adams, P.D., Alimonti, A., Bennett, D.C., Bischof, O., Bishop, C., Campisi, J.,
Collado, M., Evangelou, K., Ferbeyre, G., et al. (2019). Cellular Senescence: Defining a
Path Forward. Cell *179*, 813–827. https://doi.org/10.1016/j.cell.2019.10.005.

27. Mavrogonatou, E., Pratsinis, H., Papadopoulou, A., Karamanos, N.K., and Kletsas, D.
(2019). Extracellular matrix alterations in senescent cells and their significance in tissue
homeostasis. Matrix Biology *75–76*, 27–42. https://doi.org/10.1016/j.matbio.2017.10.004.

28. Yoshimoto, Y., Takimoto, A., Watanabe, H., Hiraki, Y., Kondoh, G., and Shukunami, C.
(2017). Scleraxis is required for maturation of tissue domains for proper integration of the
musculoskeletal system. Sci Rep *7*, 45010. https://doi.org/10.1038/srep45010.

29. Herchenhan, A., Uhlenbrock, F., Eliasson, P., Weis, M., Eyre, D., Kadler, K.E., Magnusson,
S.P., and Kjaer, M. (2015). Lysyl Oxidase Activity Is Required for Ordered Collagen
Fibrillogenesis by Tendon Cells. Journal of Biological Chemistry *290*, 16440–16450.
https://doi.org/10.1074/jbc.M115.641670.

30. Giachelli, C.M. (1999). Ectopic Calcification. The American Journal of Pathology *154*, 671–
675. https://doi.org/10.1016/S0002-9440(10)65313-8.

31. Blokland, K.E.C., Pouwels, S.D., Schuliga, M., Knight, D.A., and Burgess, J.K. (2020).
Regulation of cellular senescence by extracellular matrix during chronic fibrotic diseases.
Clinical Science *134*, 2681–2706. https://doi.org/10.1042/CS20190893.

32. Campisi, J., and d'Adda Di Fagagna, F. (2007). Cellular senescence: when bad things
happen to good cells. Nat Rev Mol Cell Biol *8*, 729–740. https://doi.org/10.1038/nrm2233.

33. De Magalhães, J.P. (2024). Cellular senescence in normal physiology. Science *384*, 1300–
1301. https://doi.org/10.1126/science.adj7050.

34. Legerlotz, K., Schjerling, P., Langberg, H., Brüggemann, G.-P., and Niehoff, A. (2007). The
effect of running, strength, and vibration strength training on the mechanical, morphological,
and biochemical properties of the Achilles tendon in rats. Journal of Applied Physiology *102*,
564–572. https://doi.org/10.1152/japplphysiol.00767.2006.

35. Ardito, F., Giuliani, M., Perrone, D., Troiano, G., and Muzio, L.L. (2017). The crucial role of
protein phosphorylation in cell signaling and its use as targeted therapy (Review).

International Journal of Molecular Medicine *40*, 271–280.
https://doi.org/10.3892/ijmm.2017.3036.

36. Popov, C., Burggraf, M., Kreja, L., Ignatius, A., Schieker, M., and Docheva, D. (2015).
Mechanical stimulation of human tendon stem/progenitor cells results in upregulation of
matrix proteins, integrins and MMPs, and activation of p38 and ERK1/2 kinases. BMC
Molecular Biol *16*, 6. https://doi.org/10.1186/s12867-015-0036-6.

37. Zhu, Y.X., Huang, J.Q., Ming, Y.Y., Zhuang, Z., and Xia, H. (2021). Screening of key
biomarkers of tendinopathy based on bioinformatics and machine learning algorithms. PLoS
ONE *16*, e0259475. https://doi.org/10.1371/journal.pone.0259475.

38. Falvino, A., Gasperini, B., Cariati, I., Bonanni, R., Chiavoghilefu, A., Gasbarra, E., Botta, A.,
Tancredi, V., and Tarantino, U. (2024). Cellular Senescence: The Driving Force of
Musculoskeletal Diseases. Biomedicines *12*, 1948.
https://doi.org/10.3390/biomedicines12091948.

39. Zeng, N., Yan, Z.-P., Chen, X.-Y., and Ni, G.-X. (2020). Infrapatellar Fat Pad and Knee
Osteoarthritis. Aging and disease *11*, 1317. https://doi.org/10.14336/AD.2019.1116.

40. Jeon, O.H., David, N., Campisi, J., and Elisseeff, J.H. (2018). Senescent cells and
osteoarthritis: a painful connection. Journal of Clinical Investigation *128*, 1229–1237.
https://doi.org/10.1172/JCI95147.

41. Wei, J., Shimazu, J., Makinistoglu, M.P., Maurizi, A., Kajimura, D., Zong, H., Takarada, T.,
Iezaki, T., Pessin, J.E., Hinoi, E., et al. (2015). Glucose Uptake and Runx2 Synergize to
Orchestrate Osteoblast Differentiation and Bone Formation. Cell *161*, 1576–1591.
https://doi.org/10.1016/j.cell.2015.05.029.

42. Chava, S., Chennakesavulu, S., Gayatri, B.M., and Reddy, A.B.M. (2018). A novel
phosphorylation by AMP-activated kinase regulates RUNX2 from ubiquitination in
osteogenesis over adipogenesis. Cell Death Dis *9*, 754. https://doi.org/10.1038/s41419-018-
0791-7.

43. Lu, Y., Yuan, T., Min, X., Yuan, Z., and Cai, Z. (2021). AMPK: Potential Therapeutic Target
for Vascular Calcification. Front. Cardiovasc. Med. *8*, 670222.
https://doi.org/10.3389/fcvm.2021.670222.

44. Cadby, J.A., Buehler, E., Godbout, C., Van Weeren, P.R., and Snedeker, J.G. (2014).
Differences between the Cell Populations from the Peritenon and the Tendon Core with
Regard to Their Potential Implication in Tendon Repair. PLoS ONE *9*, e92474.
https://doi.org/10.1371/journal.pone.0092474.

45. Dai, G.-C., Wang, H., Ming, Z., Lu, P.-P., Li, Y.-J., Gao, Y.-C., Shi, L., Cheng, Z., Liu, X.-Y.,
and Rui, Y.-F. (2024). Heterotopic mineralization (ossification or calcification) in aged
musculoskeletal soft tissues: A new candidate marker for aging. Ageing Research Reviews
95, 102215. https://doi.org/10.1016/j.arr.2024.102215.

46. Arefin, S., Buchanan, S., Hobson, S., Steinmetz, J., Alsalhi, S., Shiels, P.G., Kublickiene, K.,
and Stenvinkel, P. (2020). Nrf2 in early vascular ageing: Calcification, senescence and
therapy. Clinica Chimica Acta *505*, 108–118. https://doi.org/10.1016/j.cca.2020.02.026.

47. Stein, G.H., Drullinger, L.F., Soulard, A., and Dulić, V. (1999). Differential roles for cyclin-
dependent kinase inhibitors p21 and p16 in the mechanisms of senescence and
differentiation in human fibroblasts. Mol Cell Biol *19*, 2109–2117.
https://doi.org/10.1128/MCB.19.3.2109.

48. Zhang, J., Li, F., Nie, D., Onishi, K., Hogan, M.V., and Wang, J.H.-C. (2020). Effect of
Metformin on Development of Tendinopathy Due to Mechanical Overloading in an Animal
Model. Foot Ankle Int. *41*, 1455–1465. https://doi.org/10.1177/1071100720966318.

49. Rapisarda, V., Borghesan, M., Miguela, V., Encheva, V., Snijders, A.P., Lujambio, A., and
O'Loghlen, A. (2017). Integrin Beta 3 Regulates Cellular Senescence by Activating the TGF-
β Pathway. Cell Reports *18*, 2480–2493. https://doi.org/10.1016/j.celrep.2017.02.012.

50. Georgiadou, M., Lilja, J., Jacquemet, G., Guzmán, C., Rafaeva, M., Alibert, C., Yan, Y.,
Sahgal, P., Lerche, M., Manneville, J.-B., et al. (2017). AMPK negatively regulates tensin-
dependent integrin activity. Journal of Cell Biology *216*, 1107–1121.
https://doi.org/10.1083/jcb.201609066.

51. Schafer, M.J., White, T.A., Iijima, K., Haak, A.J., Ligresti, G., Atkinson, E.J., Oberg, A.L.,
Birch, J., Salmonowicz, H., Zhu, Y., et al. (2017). Cellular senescence mediates fibrotic
pulmonary disease. Nat Commun *8*, 14532. https://doi.org/10.1038/ncomms14532.

52. Werner, C., Hanhoun, M., Widmann, T., Kazakov, A., Semenov, A., Pöss, J., Bauersachs,
J., Thum, T., Pfreundschuh, M., Müller, P., et al. (2008). Effects of Physical Exercise on
Myocardial Telomere-Regulating Proteins, Survival Pathways, and Apoptosis. Journal of the
American College of Cardiology *52*, 470–482. https://doi.org/10.1016/j.jacc.2008.04.034.

53. Luzina, I.G., Rus, V., Lockatell, V., Courneya, J.-P., Hampton, B.S., Fishelevich, R.,
Misharin, A.V., Todd, N.W., Badea, T.C., Rus, H., et al. (2022). Regulator of Cell Cycle
Protein (RGCC/RGC-32) Protects against Pulmonary Fibrosis. Am J Respir Cell Mol Biol *66*,
146–157. https://doi.org/10.1165/rcmb.2021-0022OC.

54. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.
https://doi.org/10.1186/s13059-014-0550-8.

55. Sherman, B.T., Hao, M., Qiu, J., Jiao, X., Baseler, M.W., Lane, H.C., Imamichi, T., and
Chang, W. (2022). DAVID: a web server for functional enrichment analysis and functional
annotation of gene lists (2021 update). Nucleic Acids Research *50*, W216–W221.
https://doi.org/10.1093/nar/gkac194.

56. Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative
analysis of large gene lists using DAVID bioinformatics resources. Nat Protoc *4*, 44–57.
https://doi.org/10.1038/nprot.2008.211.

57. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet j. *17*, 10. https://doi.org/10.14806/ej.17.1.200.

58. Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.
https://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

59. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15–21. https://doi.org/10.1093/bioinformatics/bts635.

60. Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome. BMC Bioinformatics *12*, 323. https://doi.org/10.1186/1471-2105-12-323.

61. Ewels, P., Magnusson, M., Lundin, S., and Käller, M. (2016). MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics *32*, 3047–3048. https://doi.org/10.1093/bioinformatics/btw354.

62. Bankhead, P., Loughrey, M.B., Fernández, J.A., Dombrowski, Y., McArt, D.G., Dunne, P.D., McQuaid, S., Gray, R.T., Murray, L.J., Coleman, H.G., et al. (2017). QuPath: Open source software for digital pathology image analysis. Sci Rep *7*, 16878. https://doi.org/10.1038/s41598-017-17204-5.

63. Weigert, M., and Schmidt, U. (2022). Nuclei instance segmentation and classification in histopathology images with StarDist. https://doi.org/10.48550/ARXIV.2203.02284.

STAR★METHODS

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
P16-INK4A Polyclonal antibody	Proteintech	Cat# CL555-10883, RRID:AB_2923890
P21 Polyclonal antibody	Proteintech	Cat# CL594-10355, RRID:AB_2934700
ImmPRESS (Peroxidase) Polymer Anti-Rabbit IgG Reagent	Vector Labs	Cat# MP-7401-15
Chemicals, peptides, and recombinant proteins		
Harris Hematoxylin	StatLabs	Cat# HXHHEPT
Eosin Y	StatLabs	Cat# STE0157
Picro-Sirius Red Pint Stain Kit, Pint	StatLabs	Cat# KTPSRPT
Hematoxylin QS Counterstain	VectorLabs	Cat# H-304-100
Silver Nitrate	FisherScientific	Cat# BP2546100
Sodium Thiosulfate	FisherScientific	Cat# A118P-500
Formic Acid	FisherScientific	Cat# S446-500

Antigen Unmasking Solution, Citrate-Based	Vector Labs	Cat# H-3300-250
Epredia™ Shandon-Mount™	FisherScientific	Cat# 1900331
ImmPACT DAB Substrate Kit, Peroxidase (HRP)	Vector Labs	Cat# SK-415
DMEM/F12 Gibco	ThermoFisher Scientific	Cat# 11320033
RNAlater	ThermoFisher Scientific	Cat # AM7020
Critical commercial assays		
PureLink RNA mini kit	ThermoFisher	Cat# 12183018A
RNase-free DNase	Qiagen	Cat# 79254
SuperScript IV VILO Master Mix	ThermoFisher	Cat# 11756050
Power SYBR Green PCR Master Mix	ThermoFisher	Cat# 4367659
Senescence Beta-Galactosidase Staining Kit	Cell Signaling	Cat# 9860S
RNeasy Mini Fibrous Tissue Kit	Qiagen	Cat # 74704
NEBNext Ultra II RNA Library Preparation for Illumina Kit	New England Bio Labs	Cat # E7770S
Dual Index Primers	New England Bio Labs	Cat # E3392S
Deposited data		
Human Tendinopathy Bulk RNA Sequencing	This Paper; Gene Expression Omnibus	
Prkaa1$^{Scx-Cre}$ cKO Bulk RNA dataset	This Paper; Gene Expression Omnibus	
Experimental models: Organisms/strains		
C57BL/6-*AMPKa1$^{flox/flox}$*	The Jackson Laboratory	JAX:014141
C57BL/6-Tg(Scx-GFP/cre)1Stzr	Gifted from Ronen Schweitzer	MGI ID: 5317398
Software and algorithms		
SciPy Stats library	Python	
QuPath 0.5.1	QuPath	https://qupath.github.io
ImageJ	NIH	https://imagej.nih.gov/ij/

Prism v10	GraphPad	
MetaShape	Agisoft	
Adobe Illustrator	Adobe	https://adobe.com
RStudio	The R Foundation	https://cran.r-project.org/
R	The R Foundation	https://cran.r-project.org/
BioTek Gen5 Software for Imaging & Microscopy	Agilent	
MATLAB	Mathworks	
Database for Annotation, Visualization, and Integrated Discovery (DAVID)	NIH	https://davidbioinformatics.nih.gov/summary.jsp; doi:10.1093/nar/gkac194; DOI: 10.1038/nprot.2008.211
BCL Convert Conversion Software v4.0	Illumina	
Cutadapt v4.8		DOI:10.14806/ej.17.1.200 .
FastQC v0.11.8	Babraham Bioinformatics	https://bioinformatics.babraham.ac.uk/projects/fastqc/
Fastq Screen v0.15.3	Babraham Bioinformatics	https://www.bioinformatics.babraham.ac.uk/projects/fastq_screen/
GRCm38	National Library of Medicine	https://api.ncbi.nlm.nih.gov/datasets/v2/genome/accession/GCF_000001635.20/download?include_annotation_type=GENOME_FASTA&include_annotation_type=GENOME_GFF&include_annotation_type=RNA_FASTA&include_annotation_type=CDS_FASTA&include_annotation_type=PROT_FASTA&include_annotation_type=SEQUENCE_REPORT&hydrated=FULLY_HYDRATED
STAR v2.7.8a		https://doi.org/10.1093/bioinformatics/bts635
RSEM v1.3.3		https://doi.org/10.1186/1471-2105-12-323
Multiqc v1.20	Seqera	https://docs.seqera.io/multiqc/getting_started/installation
Other		
Kiralux Polarization Camera, 5 MP Monochrome CMOS Sensor, USB 3.0 Interface	ThorLabs	Cat# CS505MUP1
V500C	Biomomentum	

NovaSeqXPlus	Illumina	
ECM Select® Array Kit Ultra-36	Advanced Biomatrix	Cat# 5170
Eclipse Ni-U	Nikon	
THUNDER	Leica	
Cytation 10	Agilent	
50mm Dia. Left-Handed Plastic Circular Polarizer	Edmund Optics	Cat# 88-099
Thin Film Grip	Imada	
MA160	Biomomentum	
Ball Mill Homogenizer	Qiagen	
Qubit fluorometer	Life Technologies	
Bioanalyzer	Agilent	
Illumina NextSeq500 Instrument	Illumina	Cat # SY-415-1002

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Human

Late-stage RC (supraspinatus) tendon samples and control hamstring (semitendinosus) tendon samples were provided by the School of Infection and Immunity at the University of Glasgow. Informed consent was obtained from participants, and the use of tissue for research was approved through the West of Scotland Research Ethics Service (REC 16/WS/0207).

Animal Models

All procedures were approved by the Institutional Animal Care and Use Committees at the University of Michigan. ScxCre; Prkaa1$^{fl/fl}$ male mice were bred with Prkaa1$^{fl/fl}$ female mice to generate Prkaa1$^{Scx-Cre}$ (cKO). Prkaa1$^{fl/fl}$ (WT) and Prkaa1$^{Scx-Cre}$ cKO (cKO) mice were generated for evaluation of Achilles tendon structure and function at 1-, 3-, and 9-months (1M, 3M, 9M). All mice were kept in standard housing conditions with ad libitum access to water and chow throughout the experiment.

METHOD DETAILS

RNA isolation and sequencing of human tendon

Tissues were collected and stored in RNALater, these were subsequently physically disrupted using a ball mill homogenizer. RNA extraction was performed using RNeasy Mini Fibrous Tissue Kit with on-column deoxyribonuclease I treatment according to manufacturer's instructions. RNA quality and quantity was obtained using Qubit fluorometer and the RNA Pico kit and Bioanalyzer 2100 system.

The NEBNext Ultra II RNA Library Preparation for Illumina Kit was used to prepare the samples for sequencing following the manufacturer protocol. Dual Index Primers were used in the NEXNext Ultra II

RNA Library Preparation for Illumina protocols. Library qualities were confirmed using Qubit fluorometer
and high sensitivity DNA kit and Bioanalyzer. Samples were then sequenced using an Illumina
NextSeq500 Instrument (at University of Glasgow Polyomics Facility (Glasgow Polyomics, University of
Glasgow).

Differential gene expression was determined from count matrices with a paired design (e.g. Healthy vs.
Diseased) in DESeq2 in R/Bioconductor[54].We used the Database for Annotation, Visualization, and
Integrated Discovery (DAVID) to analyze biological processes and Kyoto Encyclopedia of Genes and
Genomes (KEGG) pathways[55,56].

Wheel Running

At 1M of age WT and Prkaa1[Scx-Cre] cKO mice were randomly assigned to either remain at cage activity
(CA) or were singly housed with voluntary access to running wheels (WR) for 8 weeks in standard mouse
cages. Daily wheel rotations were collected and recorded. One week prior to euthanasia mice were fasted
for 6hr to collect fasting blood glucose (FBG) measurements.

Biomechanical testing of Achilles tendon

At time of euthanasia, one hindlimb from each mouse was immediately frozen at -20°C. Simple
randomization of samples prior to Achille's tendon dissection was performed and tester was blinded to all
experimental groups. 24 hours before the start of mechanical testing, limbs were thawed at 4°C and
prepared for photogrammetry and biomechanical testing. Both the plantaris tendon and muscle were
carefully removed and the Achilles tendons and calcaneus were left intact. Cross-sectional areas (CSA)
of the Achilles tendon were measured using photogrammetry with a custom pinch clamp holder attached
to a motor controller (Arduino, Ivrea, Italy). With the muscle and Achilles hanging free of the pinch clamp,
at least 50 consecutive images of the pinch clamped tendon were acquired using a 12mm focal length
lens (Basler Fujinon Lens, Ahrensburg, Germany). Using Metashape software (Agisoft, St. Petersburg,
Russia), images were aligned and converted first to a sparse point then to a dense point cloud, to create
an STL surface mesh of the tendon. CSA was defined as the smallest area from the STL surface mesh
generated from the Achilles tendon and was measured using a slice analysis tool in Dragonfly (Comet
Technologies Canada Inc., Montreal, Canada).

A custom 3D printed fixture secured the calcaneus into place (FormLabs 3B, Somerville, MA, USA), and
the proximal Achilles tendon was clamped in a textured grip and screwed into place (Imada, Northbrook,
IL, USA). The assembled grip with the secured Achilles tendon was placed into a phosphate buffered
solution (PBS) bath at 37°C using a temperature controller (MA160, Biomomentum Inc., Laval, Quebec,
Canada) and secured with a pin to a tensile testing frame with a multi-axis load cell (±70 N; Mach-1
VS500CST, Biomomentum, Laval, Quebec, Canada). Samples were preloaded to 0.1 N, and gauge
length was measured as the distance between the secured calcaneus and the textured grip.

The Achilles tendons were then preconditioned for 10 cycles (±0.1 N at 0.1mm/sec) followed by the load
to failure test at 0.1mm/sec. Off-axis load (forces in X and Y, torques in X, Y, Z) were collected to assess
off-axis loading for the duration of the experiment. Instantaneous grip-to-grip strain was calculated as the
displacement divided by the original gauge length, L_0. For the 1M, 3M and 9M WT and cKO age-
dependent tests maximum load, maximum stress, and maximum strain were calculated using a custom
MATLAB script (R2017 or later, MathWorks, Natick, USA). Data between cKO and WT Achilles tendons
at 1M, 3M and 9M were compared using a two-way ANOVA with post-hoc Sidak's multiple comparisons
to identify specific differences (Genotype, Time).

RNA isolation and sequencing of murine tendon

1M and 3M old as well as WR and CA WT and Prkaa1[Scx-Cre] cKO mice were euthanized and Achilles
tendons were immediately dissected under ribonuclease (RNase)–free conditions and flash frozen ($n =$
3/group). Tissues were mechanically pulverized in TRIzol, and total RNA was isolated using spin-columns

(with on column genomic DNA digestion. RNA Integrity Number (RIN) was measured using Bioanalyzer
(Agilent). This pool was subjected to 151bp paired-end sequencing according to the manufacturer's
protocol (Illumina NovaSeqXPlus). BCL Convert Conversion Software v4.0 (Illumina) was used to
generate de-multiplexed Fastq files. Reads were trimmed using Cutadapt v4.8[57] and FastQC v0.11.8 was
used to ensure the quality of data[58]. Next, Fastq Screen v0.15.3 was used to screen for various types of
contamination (Wingett and Andrews, 2018). Reads were then mapped to the reference genome
GRCm38 (ENSEMBL), using STAR v2.7.8a [59] and assigned count estimates to genes with RSEM v1.3.3
[60]. Alignment options followed ENCODE standards for RNA-seq. Multiqc v1.20 compiled the results from
several of these tools and provided a detailed and comprehensive quality control report [61]. Differential
gene expression was determined from count matrices with a paired design (e.g., WT v cKO, WT WR v.
CA or cKO WR v. CA) in DESeq2 in R/Bioconductor[54]. We used the Database for Annotation,
Visualization, and Integrated Discovery (DAVID) to analyze biological processes and Kyoto Encyclopedia
of Genes and Genomes (KEGG) pathways[55,56].

Histology Preparation and Imaging

Distal hindlimbs from WT and cKO tendons (n = 3 per sex/group) were dissected, fixed in 4%
paraformaldehyde (PFA), and decalcified in 14% ethylenediaminetetraacetic acid (EDTA) and processed
for paraffin embedding. Tissues were sectioned at 5 μm in the sagittal plane and stained with either
Hematoxylin and Eosin (H&E), Silver Nitrate, Picrosirius Red (PSR) or processed for
immunohistochemistry (IHC), and cover-slipped with an acrylic mounting media. H&E, Silver Nitrate and
(IHC) slides were imaged on a bright-field microscope (ECLIPSE Ni-U, Nikon) at 10x and analyzed using
QuPath. PSR slides were imaged with a 10x objective on an epi-fluorescent microscope (dmi600b, Leica)
and analyzed using quantitative polarized light imaging (qPLI) analysis.

Immunohistochemistry

For immunohistochemistry, paraffin sections were deparaffinized followed by heat-mediated antigen
retrieval with an Antigen Unmasking Solution, Citrate-Based or Tris-EDTA (pH 9.0) for 2hrs at 65°C and
blocking (2.5% horse serum). Slides were then incubated in primary P16-INK4A and p21 antibodies
overnight at 4°C, followed by washing and incubation with appropriate secondary antibodies. ImmPACT
DAB Substrate Kit, Peroxidase (HRP) was used for detection, and slides were counterstained with
hematoxylin and cover-slipped with Shandon Mount. Antibody Information Table 2.

qPLI Analysis

The degree of linear polarization (DoLP) and angle of linear polarization (AOP) images were acquired
using a polarization camera and a circular polarizing lens. The mean DoLP and standard deviation of the
AOP were analyzed using Math and SciPy Stats libraries in. Data from cKO mice were compared to WT
controls at 1M, 3M and 9M using a two-way ANOVA with post-hoc Sidak's multiple comparisons to
identify specific differences (Genotype, Time).

FIJI Analysis

Silver Nitrate slides were imaged and collected for percent lesion area analysis by FIJI (NIH, Bethesda,
MD). First, the Achilles tendon region of interest area was selected using morphological features to
determine total area. Next, using the polygon shape tool, the positive lesion was outlined and the area
was measured. All measurements were taken three times to ensure reproducibility. Percent lesion area
was calculated by dividing lesion area(s) by the whole tendon area. Data from Prkaa1[Scx-Cre] cKO mice
were compared to WT controls at 9M using an unpaired two-tailed t-test.

QuPath Analysis

IHC slides were imaged and collected for percent positive analysis by QuPath[62], an open-source software
for digital pathology. First, the regions of interest in Achilles tendons were selected using morphological

features and areas were measured. Next, color deconvolution was performed by QuPath to digitally
separate stains (Hematoxylin and DAB) within each tissue sample. Positive cell detection, highlighting
DAB positive cells, was then performed to differentiate cells. Total cell detection was collected from
counting hematoxylin positive nuclei. Data from Prkaa1[Scx-Cre] cKO mice were compared to WT controls at
3M and 9M using an Ordinary two-way ANOVA with uncorrected Fisher's LSD, with a single pooled
variance to identify specific differences (Genotype, Time).

Nuclear Circularity

752 Slides were deparaffinized and stained with antifade mountant with NucBlue. Slides were imaged with
753 widefield fluorescence on a Cytation 10 at 20x. Nuclei counts and circularity were calculated with the
754 Gen5 cellular analysis tool. Data from Prkaa1[Scx-Cre] cKO mice were compared to WT controls using a one-
755 way ANOVA with Tukey's multiple comparisons test with a single pool variance.

Cell Isolation and Culture

757 Prkaa1[Scx-Cre] cKO mice were euthanized and tails were removed. Skin was removed from tail and tendon
758 fascicles were carefully dissected and placed in a 60mm petri dish with Dulbecco's Modified Eagle's
759 Medium/Nutrient Mixture F-12 (DMEM: F12), 10 % Fetal Bobine Serum (FBS) and 1%
760 Penicillin/Streptomycin (P/S), media was changed every other day. Tail TFs were allowed to "crawl out"
761 for two weeks before passaging for β-Galactosidase staining and ECM arrays.

β-Galactosidase (Beta-Galactosidase) Staining

763 Prkaa1[Scx-Cre] cKO Tail TFs were passaged into a 12w plate in DMEM: F12, 10% FBS and 1% P/S for 24h.
764 Media was changed to DMEM: F12, 1% FBS and 1% P/S for 96h, media was changed every other day.
765 Cells were fixed and stained per manufacturer's instructions with the Senescence β-Galactosidase
766 Staining Kit and imaged on an Agilent Lionheart RX.

ECM Arrays

768 Prkaa1[Scx-Cre] cKO Tail TFs were passaged onto a per manufacturer's instructions in DMEM: F12, 10%
769 FBS and 1% P/S for 24h. After 24h, cells were fixed with 4% PFA and, rinsed with PBS stained with
770 Hoechst. Slides were imaged using a Leica THUNDER and nuclei were counted using FIJI with the
771 Stardist plugin[63]. Counts were collected and analyzed with Bayesian statistical modeling.

Bayesian Statistical Modeling

773 Nuclear Counts for the WT and Prkaa1[Scx-Cre] cKO Tail TFs on different substrates were
774 compared with the fit of a range of two Bayesian regression models, baseline and interaction. In the
775 baseline model the cell counts are assumed to occur assumed to occur independently at random only.
776 With the interaction model, the cell counts are dependent on the genotype and substrate.
777
QUANTIFICATION AND STATISTICAL ANALYSIS

779 For graphs in figures 2,4 and 5 data are shown as mean ± standard deviation (SD). GraphPad (GP) p-
780 value style, * p<0.05, ** p<0.01, ***p<0.001, ****p<0.0001. Statistical differences were evaluated using
781 two-way ANOVA with Sidak correction or otherwise stated in figure legends. Graphs were produced and
782 statistical analyses were performed using GraphPad Prism. For figure 3 data were compared with the fit
783 of a range of two Bayesian regression models, baseline and interaction.
784

A. Biological processes

Mitochondrial electron transport

FBP metabolic process

Energy homeostasis

Cytokine-mediated signaling pathway

Cellular response to glucose stimulus

Enrichment
0   5   10   15   20

B. Molecular functions

Phosphofructokinase activity

Fructose-6-phosphate binding

AMP-activated protein kinase activity

Enrichment
0   10   20   30   40   50

 # DEGs
10   20
30

C. KEGG Pathways

AMPK signaling pathway

Metabolic pathways

Glycolysis / Gluconeogenesis

Cytokine-cytokine receptor interaction

Oxidative phosphorylation

Enrichment
0   1   2   3   4   5

D. Human Tendinopathy Signaling Diagram

MT-ND1

NAD+/NADH ratio

PFKP

PFKFB1

PRKAA2

PPARGC1A

IL36G

IL13RA2

IL1RN

TNFSF9

TNFSF11

From KEGG Pathways:
Metabolic Pathways ○
AMPK Signaling Pathway ○
Cytokine-cytokine Receptor interaction ●

E. Relative expression of DEGs from signaling diagram

log2FC
0
-1
-2

MT-ND1

PRKAA2
PFKP
PPARGC1A
PFKFB1

IL36G
IL13RA2
IL1RN
TNFSF9
TNFSF11

A. Biological Processes

Response to stress
Response to glucose
Glucose metabolic process
Glucose homeostasis
ECM organization
Collagen catabolic process
Cellular response to oxidative stress
Cell adhesion mediated by integrin

Enrichment
0 1 2 3 4 5 6

#DEGs
10 20 30

B. Cell Cycle Diagram

Cdkn1a → Senescence

Ccnd3 → Cell Cycle Progression

Bop1

C. Tendon Maturation Diagram

Mesenchymal Stem Cell

Osteoblast Progenitor — *Runx2*
Tendon Progenitor — *Scx Sox9*

Osteoblast — *Sparc*
Immature tendon cells — *Mkx*
Mature tendon cells — *Col1a1 Lox Dcn Col1a2*

D.

Max load (N) — 1M, 3M, 9M — ****

E.

Force (N) vs Displacement (mm)
1M, 3M, WT, cKO

F.

CSA (mm²) — 1M, 3M, 9M — * — WT, cKO

G.

Max Stress (MPa) — 1M, 3M, 9M — ** ** — WT, cKO

H.

WT   cKO

Male

Female

I.

WT   cKO

J.

Lesion Area at 9M (%) — WT, cKO — *

K.

Nuclear Circularity — **** **** — WT, cKO — H H L

A.

WT   cKO

COL I

COL III

Hoescht

B. Baseline Model

cKO
Col1
Col3
Col4
Col5
Col6
Fibronectin
Laminin
Vitronectin
Tropoelastin

30%   1x   3x
Effect Size

C. Interaction Model

WT

cKO
Col1
Col3
Col4
Col5
Col6
Fibronectin
Laminin
Vitronectin
Tropoelastin

30%   1x   3x
Effect Size

cKO

cKO
Col1
Col3
Col4
Col5
Col6
Fibronectin
Laminin
Vitronectin
Tropoelastin

30%   1x   3x
Effect Size

A. *in vitro* Beta-Gal

WT   cKO

B. p21

WT   cKO

C. p16

WT   cKO

9M

D.

***

p21 Positive Cells (%)

WT   cKO

E.

****

p16 Positive Cells (%)

WT   cKO