Using artificial intelligence to optimize ecological restoration for climate and biodiversity

DANIELE SILVESTRO[1,2,3], STEFANO GORIA[3,4], BEN GROOM[5,6], THOMAS STERNER[3,7], ALEXANDRE ANTONELLI[3,8,9,10]

[1] *Department of Biosystems Science and Engineering, ETH Zurich, Basel, Switzerland*

[2] *Gothenburg Global Biodiversity Centre, Department of Biological and Environmental Sciences, University of Gothenburg, Sweden*

[3] *Captain Technologies Ltd, London, UK*

[4] *Thymia Ltd, London, UK*

[5] *LEEP Institute, Department of Economics, University of Exeter Business School, Exeter University, UK*

[6] *Grantham Research Institute on Climate Change and the Environment, London School of Economics, London, UK*

[7] *Department of Economics, University of Gothenburg, Sweden*

[8] *Royal Botanic Gardens, Kew, Richmond, UK*

[9] *Antonelli Foundation for Biodiversity Research and Conservation Rio de Janeiro, Brazil*

[10] *Department of Biology, University of Oxford, UK*

Correspondence: daniele[at]captaintech.ai, a.antonelli[at]kew.org

Abstract

The restoration of degraded ecosystems can provide important contributions to help mitigate climate change and bend the curve of biodiversity loss. Depending on the primary objective – such as maximizing carbon storage, protecting threatened species, or reducing overall costs – different spatial priorities have been identified at global and regional levels. Funding mechanisms to support such work comprise public sources, philanthropy, and the private sector, including the sales of carbon and biodiversity credits. However, effectively exploring tradeoffs between restoration objectives and estimating the price of biodiversity and carbon credits to design financially viable projects remain challenging. Here we harness the power of artificial intelligence in our software CAPTAIN, which we further develop to identify spatial priorities for ecological restoration that maximize multiple objectives at once and which allows a robust evaluation of biodiversity and climate outcomes. We find through a series of realistic simulations that even low to moderate consideration of biodiversity in restoration projects leads to the selection of restored areas that substantially improve the conservation of threatened species, while resulting in a relatively small decrease in total carbon captured. We propose a data-driven valuation of biodiversity credits in relation to carbon credits enabling the design of a blended financial model that could support restoration efforts even in areas previously excluded for economic reasons. This study shows how the use of a robust methodological framework can lead to significant improvements in outcomes for climate and nature, while minimizing costs.

Keywords: *multi-objective reinforcement learning, Global Biodiversity Framework, Paris Agreement, CAPTAIN*

Introduction

Ecological restoration is one of the most powerful nature-based solutions to tackle the twin planetary crises of biodiversity loss and climate change (Seddon et al., 2020). This is why the United Nations has declared 2021-2030 the Decade on Ecosystem Restoration and motivates the commitment under Target 2 of the Kunming-Montreal Global Biodiversity Framework – ratified by 196 countries – that "*by 2030 at least 30 per cent of areas of degraded terrestrial, inland water, and marine and coastal ecosystems are under effective restoration*". Alongside the conservation of natural ecosystems – the preferred choice whenever there is an option – ecological restoration that is well designed and implemented can capture and store carbon, increase the populations of threatened species, enhance ecosystem services and multiple other contributions to people and the natural environment (e.g., Di Sacco et al., 2021).

Despite the manifold benefits of restoration and its urgency in the face of global environmental change, two major challenges hamper its implementation. Firstly, much like prioritizing conservation areas (Jung et al., 2021), there are clear trade-offs when designing restoration projects aiming to maximize carbon capture, biodiversity recovery, or reducing costs (Strassburg et al., 2020). For example, the plantation of exotic trees across Africa, while growing fast and capturing large amounts of carbon (in principle helping to mitigate climate change, if the biomass produced were retained long-term), has inadvertently led to detrimental effects on the native biodiversity (Parr et al., 2024). While such trade-off effects have been widely discussed in the literature (e.g. Holl and Brancalion, 2020; Hua et al., 2022), to our knowledge no tools have been yet developed to allow their quantification that take into account the complexity of the real world – including the spatial heterogeneity in biodiversity, carbon storage potential, costs – and their changes over time.

Secondly, there is a major funding gap – estimated between USD 600 billion and over 800 billion (Deutz et al., 2020) – to implement the Global Biodiversity Framework, including its restoration target. Given that public funding from institutions such as the Global Environment Facility falls far short of reaching the required levels, many restoration projects today are supported instead by the sales of carbon credits, which can be voluntary or mandatory depending on the country. However, the simplicity of carbon credits – where one credit equates to one tonne of atmospheric carbon captured, regardless of location – is also its pitfall. As market forces push trade towards low prices, this incentivizes restoration projects that minimize costs (such as reforestation on marginal and highly degraded land) and focus exclusively on maximum carbon recovery, even at the cost of biodiversity benefits. To improve this situation, carbon credits are being designed to include additional benefits such as biodiversity, and, although this is much more complicated, biodiversity credits are emerging following strict principles for high-integrity and governance (Antonelli et al., 2024). However, we are not aware of attempts to quantify the cost relationship between carbon and biodiversity credits to make them financially interchangeable

(allowing, for instance, a restoration project to become viable by selling biodiversity credits rather than carbon credits), nor the economic opportunities for bundling carbon and biodiversity credits for the same area making it viable to restore ecosystems in expensive areas, which have so far been excluded due to insufficient profit margins.

Here, we propose an approach based on multi-objective reinforcement learning to address these shortcomings. Specifically, we i) demonstrate through simulations that this approach can help spatially guide restoration programs while monitoring their progress over time; ii) explore trade-offs and synergies that exist between restoration efforts focused on carbon, biodiversity and costs, through multi-objective optimization to estimate socially optimal solutions; and iii) link our predictions to the emerging concept of biodiversity crediting alongside carbon credits to design financially viable restoration programs. Our results show that identifying trade-offs between carbon and biodiversity is crucial for appropriately valuing biodiversity credits to incentivize restoration programs that maximize benefits for both climate and biodiversity.

Methods

A reinforcement learning framework for restoration

We optimize spatial restoration planning within a reinforcement learning framework developing new functionalities within the CAPTAIN (Conservation Area Prioritization Through Artificial Intelligence) software (Silvestro et al., 2022). The implementation includes a spatially explicit dynamic *environment* characterized by spatial and climatic features, species and their geographic ranges, and anthropogenic disturbances. The disturbance could be any kind of anthropogenic activity including settlements, roads, farming or logging on land, fishing or mining in the sea. The framework also includes an *agent* (i.e. a manager or social planner) who monitors the environment and implements a restoration program pursuing one or multiple pre-defined restoration targets. The system (environment and agent) evolves over time through a number of steps (e.g. years) at which multiple processes take place, including dispersal and mortality among species, changes to climate or disturbances, and the establishment of restoration areas, where the disturbance is reversed.

In this study we use environments and species generated through simulations to assess the performance and results of different restorations models under fully controlled settings. However, the framework can be used with empirical data, for instance by using species distribution models, climate data, and land use and other indexes of human footprint, and real costs of restoration. Similarly empirical data, when available, can be used to inform the model about species dispersal ability, sensitivity to disturbances, and -in the case of plants- net sequestrated carbon.

Environment – The environment is discretized into spatial units of arbitrary size that represent the cells of a grid and are used as units for the description of environmental variables,

species ranges, and restoration actions. Each cell features environmental variables distinguishing land from sea and providing elevation and climate data (such as temperature and precipitation – e.g. their mean, monthly and seasonal variations; for ocean environments, specific variables such as water depth, oxygen levels, and salinity could be used). Cells are also characterized by a value of anthropogenic disturbance, indicated with $\delta \in [0,1]$, that alters the ability of species to live in the cell and their abundance (see below) and by a restoration cost (ξ), quantifying the relative cost of restoring the cell to a low level of disturbance. This could include, for instance, the cost of buying the cell, displacing or removing the cause of disturbance, and potentially taking action to reintroduce native species. Note that in real conservation work, restoration costs may be complex depending on the legal and socio-economic setting. For example, buying land is not always sufficient to stop all logging, but we assume that we can model appropriate total costs here. All these features can be set to change through time, e.g., capturing the dynamics of predicted climate change, variation in disturbance patterns, and evolving costs per cell.

The environment is also characterized by the species that live (or can live) in it. Specifically, for each cell and each species in the system, there is a value describing the number of individuals that can occupy the cell in the absence of disturbance (i.e. the natural state). This is a cell- and species-specific carrying capacity $K_{c,s}$ which depends on both the habitat suitability of the cell for that species (H_s) and a species-specific maximum abundance per cell (K_s):

$$K_{c,s} = H_{c,s} \times K_s, \text{ where } H \in [0,1] \text{ and } K_s \in \mathbb{R}^+ \tag{1}$$

The environment also includes the actual number of individuals of a species at any given time t, $N_{c,s,t}$, which results from the interaction between the species and the environment, including the effects of disturbance, growth rates, and dispersal.

For each species, we define a specific value of sensitivity $S_s \in [0,1]$ that determines individual mortality fractions in response to a disturbance in a cell, $\delta_{c,t}$. Species are also assigned a growth rate $G_s > 1$, which describes the demographic growth in one time step, and a dispersal rate $D_s \geq 0$, which determines the number of individuals that disperse from one cell to another. In the current implementation, we assume individuals to be static (e.g., plants). Therefore, dispersal does not move individuals from one cell to another. Instead, dispersal only adds individuals to the receiving cell (replicating for instance the dispersal of seeds). The number of individuals dispersing from cell i to j follows a truncated exponential distribution:

$$I_{i \rightarrow j,s,t} = \begin{cases} N_{i,s,t} \times \exp(-d_{ij}D_s^{-1}) \text{ if } d_{ij} \leq T \\ 0 \text{ if } d_{ij} > T \end{cases} \tag{2}$$

where d_{ij} is the distance between the two cells and T is the maximum distance that can be reached in a single time step. Species distributions and abundances change over time based on

growth, dispersal, and mortality. Specifically, at each time step, the distribution and abundance of species is updated according to the following equation:

$$N_{c,s,t} = \min\left(N_{c,s,t-1} \times (1 - S_s\delta_{c,t}) \times G_s + I_{c,s,t}, K_{c,s}\right) \tag{3}$$

where $N_{c,s,t-1}$ is the number of individuals of species s in cell c at the previous time step, $S_s\delta_{c,t}$ is the product between the disturbance in the cell at time t and the species sensitivity, with $(1 - S_s\delta_{c,t})$ representing the mortality driven by disturbance. The term G_s represents the contribution of demographic growth, while

$$I_{c,s,t} = \sum_{\forall j \neq i} I_{j \to i, s, t-1} \tag{4}$$

is the sum of the individuals dispersing from all other cells and represents the contribution of dispersal to the cell's population.

Each species in the environment is assigned a value representing its expected carbon value $C_s \in \mathbb{R}^+$. This is a single value and can be interpreted as the net amount of carbon sequestered by one individual of species s. As our simulated system in its current implementation does not track the age of each individual, we define the total carbon in a cell as the sum of carbon storage from all individuals and all species in the cell:

$$C_{c,t} = \sum_s C_s N_{c,s,t} \tag{5}$$

This quantity could in principle extended to include carbon in the soil. We note that, as the age of individuals is not modeled here, the simulated accumulation of carbon in the current implementation might occur more rapidly than in empirical systems.

Finally, we also include a 5-tiered classification of species extinction risk, inspired by the International Union for Conservation of Nature (IUCN) Red List of Threatened Species (2023). When working on empirical data, this can be effectively replaced by actual IUCN Red List assessments or model-based predictions (Zizka et al., 2022; Antonelli et al., 2023). Hereafter, we will use the Red List-inspired labels to describe increasing extinction risks: Least Concern (LC), Near Threatened (NT), Vulnerable (VU), Endangered (EN), and Critically Endangered or (locally) extinct (CR). As part of the evaluation of the implemented restoration action, the initial extinction risk of a species can be updated based on the simulated dynamics of the species demography or geographic range, for instance, increasing or decreasing the extinction risk depending on whether the census population size of a species is decreasing or increasing, respectively. We indicate the extinction risk of each species s at time t as:

$$X_{s,t} \in \{1, ..., 5\}$$

Agent – The agent in the system observes the environment at each time step extracting features for each grid cell about its disturbance level, its cost for restoration, the species that occur there, those that could occur there naturally in the absence of disturbance, their current extinction risk and carbon contribution (a detailed list of the features used in our experiments is provided in the next section). Following the modeling framework developed in the initial implementation of CAPTAIN (Silvestro et al., 2022), the features of each cell feed into a fully connected deep neural network with `tanh` activation function (Apicella et al., 2021) in the hidden layers and a single output node. The parameters of the neural networks are shared across all cells and their output values are pooled into a single `softmax` function providing a restoration rank of each cell. The agent will then select deterministically or probabilistically a pre-defined number of cells based on their ranking, within which restoration occurs. Here we assume restoration to result in the instantaneous removal of disturbance from the cells, although different dynamics such as a gradual decrease or partial reduction in disturbance can be implemented. The restoration action is associated with the cumulative cost of the selected cells. The agent is given an initial budget that can be spent for restoration, and only cells with a cost lower than the current budget can be selected for restoration. Once the disturbance is removed from a cell, its populations will continue to evolve according to Equation 3 with reduced (or removed) disturbance.

Rewards and optimization – The actions of the agent result in changes in the environment that are quantified through a *total reward*, defined based on the objective of the restoration policy. The parameters of the neural networks determining the restoration action are chosen to maximize the cumulative net reward at the end of a simulation of a fixed number of steps. The net reward or payoff is here defined as the weighted sum of three components quantifying net carbon gain (carbon reward), net biodiversity outcomes (biodiversity reward), and costs (budget reward). These quantities allow us to optimize policies with one or multiple objectives, generating socially optimal solutions when multiple objectives are used. At each time step, the carbon reward of a restoration action is computed as the fraction between the net change in total carbon stored in the system compared with its initial state (at time $t = 0$) and the amount of carbon lost between the natural state (in the absence of any human disturbance) and the initial state:

$$\mathcal{R}_{C,t} = \frac{\sum_c C_{c,t} - \sum_c C_{c,t=0}}{\sum_c C_{c,\text{natural}} - \sum_c C_{c,t=0}} - \mathcal{R}_{C,t-1} \tag{6}$$

We note that with simulated datasets, like the ones used in this study, the natural carbon amount C_c, natural is known *a priori*. When working with empirical data, this quantity might

have to be approximated based on measurements, such as through survey data (for trees, this can be derived indirectly from measurements of height and stem diameter at breast height (dbh), or more directly from biovolumes estimated from lidar scans).

The total reward of a restoration action also includes a measurement of the resulting biodiversity outcome, which can be defined in many different ways, as shown by a recent compendium (Burgess et al., 2024) currently listing over 570 biodiversity metrics. In the current implementation, we exemplify the calculation of biodiversity gains in a similar way as recently calculated in Silvestro et al. (2023), who used the number of extinct or threatened species as a negative reward. The focus on extinction risk is also aligned with other recently proposed metrics for conservation priorities and threat abatement (Durán et al., 2020; Mair et al., 2021; Eyres et al., 2024). Here we use an average of the restored species population weighted by their current conservation status. Specifically, for any given extinction risk status $X = i$, we identify the subset of species S_i such that:

$$X_{s,t} = i, \forall s \in S_i \tag{7}$$

and compute the respective reward as

$$\mathcal{R}_{X=i,t} = \begin{cases} \dfrac{w_i}{\sum_j w_j} & \text{if } S_i = \emptyset \\ \displaystyle\sum_{s \in S_i} \dfrac{w_i R_{s,t}}{\sum_j w_j} & \text{otherwise} \end{cases} \tag{8}$$

where w_i is a weight associated with extinction risk status i, and $R_{s,t}$ is the ratio between the number of individuals of species s at time t found within restored cells (indicated with M) and the total population of the species in the environment:

$$R_{s,t} = \frac{\sum_{\varkappa \in M} N_{\varkappa,s,t}}{\sum_c N_{c,s,t}} \tag{9}$$

Thus, $R_{s,t} = 0$ when the species is not found in any restored areas and $R_{s,t} = 1$ when all its population falls inside restored areas. The weights are set to favor scenarios where the highest average fraction of population is restored in species classified in a high-risk category:

$$\begin{cases} w_1 = 1 \\ w_2 = 8 \\ w_3 = 16 \\ w_4 = 32 \\ w_5 = 64 \end{cases} \tag{10}$$

so that, e.g., protecting 30% of the individuals in a least concern species ($X = 1$) contributes 0.0025 to the reward, while the same fraction in a critically endangered species ($X = 5$) contributes 0.1587. The overall biodiversity reward is set to the sum of $R_{X,t}$ over all extinction risk categories. The use of increasing weights associated with extinction risk is consistent with several metrics developed to quantify conservation priorities (Pimiento et al., 2020; Mair et al., 2021; Gumbs et al., 2023).

Finally, we include restoration costs ε_c as part of the total reward. Based on an initial budget $B_{t=0}$ we compute the budget reward $R_{\varepsilon,t}$ as the ratio between the current budget and the initial one. Here we set the initial budget as a function of the restoration target (30%) and the total cost of restoration. Specifically, given a restoration cost ε_c (defined in the next section) for cell c and a restoration target $r \in [0,1]$ specifying the fraction of cells to be restored, we set the initial budget to a value arbitrarily larger than the total average cost a number of cells required to meet the target:

$$B_{t=0} > r \sum_c \varepsilon_c \tag{11}$$

The rewards can be used alone or in combination to optimize policies that pursue one or more targets. Thus, we defined the total reward as:

$$\mathcal{R}_t = w_C \mathcal{R}_{C,t} + w_X \mathcal{R}_{X,t} + w_\varepsilon \mathcal{R}_{\varepsilon,t} \tag{12}$$

were the pre-defined weights (w) quantify the proportional importance assigned to carbon, biodiversity, and budget. Here we set the weight associated with the budget (w_ε) to a constant and set the others to express their relative importance. Under this formulation, a policy targeting a single objective (e.g., focusing exclusively on budget) can be optimized by setting the respective weight to 1 and the others to 0 (e.g., $w_C, w_X = 0$, and $w_\varepsilon = 1$). The weights can also be used to estimate different solutions where the total reward includes two or more potentially competing optimality criteria.

We use the parallel genetic strategies algorithm used in Silvestro et al. (2022) to estimate the parameters of the neural networks in the model that maximize the reward. Here, multiple environments are run in parallel with an optimization step occurring at the end of each training epoch.

Simulations

We use simulations of biodiversity to evaluate the ability of restoration policies optimized through reinforcement learning to achieve different targets, focusing on carbon and biodiversity outcomes. Although the same policies could be applied to empirical data, simulations allow

us to benchmark and compare different restoration actions across many simulated scenarios in which we can accurately measure all outcomes.

Parameters of the simulated environment – We simulate environments based on a raster of 128×128 cells inspired from physical properties from an arbitrary region spanning a portion of Central and South America (Fig. 1a). The properties include its coastline, elevation and cell-specific climate average values obtained from CHELSA (Karger et al., 2023), here including the mean annual temperature, mean annual precipitation and the minimum temperature of the coldest month. The selection of variables is not intended to provide an exhaustive account of the climate variables determining a species niche but were used to generate artificial species distributions with realistically constrained geographic ranges. After z-transforming these topographic and climate variables we indicate them as $V = V_1, ..., V_v$, where $V_{i,x,y}$ represents the value of the i^{th} variable corresponding to the longitude and latitude coordinates x, y.

We simulated species geographic range by first drawing a random pair of coordinates (m, n), based on which we extracted the corresponding elevation and climate data. We use a normal distribution to describe the environmental and climatic niche of the species $\text{MVN}(V_{m,n}, \Sigma)$, where the diagonal of the covariance matrix is a vector $\sigma = \{\sigma_1, ..., \sigma_V\}$ is randomly drawn for each species from a uniform distribution $\mathcal{U}(0.05, 1)$ and off-diagonal values (covariances) are set to 0. The σ vector will therefore generate species with a range of narrow to wide climatic tolerances. We also introduce a distance factor, to ensure that species ranges are also spatially constrained e.g. through limited dispersal ability and not only determined by habitat suitability. The effect of distance was simulated through an exponential distribution with rate parameter d randomly drawn for each species from a uniform distribution $\mathcal{U}(0.05, 1)$. Thus, for any set of coordinates (x, y) the relative population density of a species is $\text{MVN}(V_{x,y}, V_{m,n}, \Sigma) \exp(-d\Delta(x, y, m, n))$, where $\Delta(\cdot)$ is the cartesian distance between (m, n) and (x, y). We use this formula to define the habitat suitability $H_{c,s}$ of a species s in each cell c, as used in Equation 1.

We used the empirical mean annual precipitation layer (rescaled between 0 and 1) to split cells into three categories with high (> 0.4), intermediate (< 0.4) and low precipitation (< 0.25). We use these categories to simulate different vegetation types, e.g. representing one dominated by trees, a mixed open habitat, and a fully open habitat (Fig. 1a). These vegetation types are then used to characterize properties of the species that inhabit them. Specifically, we simulated three categories of species to represent different growth forms, e.g. herbaceous plants, shrubs, trees. We assigned each species to one of the categories randomly with probability set equal to the proportion of the overlap of their simulated range with low, intermediate and high precipitation, respectively.

We drew the species demographic growth rates from a Weibull distribution:

$$G_s \sim 1 + W(0.75, s)$$

where $s = 10$, 2, and 0.1 for the first, second, and third growth forms, respectively. This results in faster average demographic growth for simulated 'herbaceous' species. Similarly, we drew the species-specific carbon value from a gamma distribution:

$$C_s \sim \Gamma\left(2, \frac{m}{2}\right)$$

where $m = 0.5$, 1, or 2 is the mean of the distribution applied to the first, second, and third growth forms, respectively. We draw the species-specific carrying capacity K_s (i.e. the maximum number of individuals that can live in a cell; Equation 1) from a beta distribution that generates many species with low abundance and few species with high abundance:

$$K_s \sim 2 + 100 \times B(0.6, 1.2)$$

This leads to a mean carrying capacity a species per cell of ≈ 35 ranging from ≈ 2 to ≈ 102. We assigned these values by setting the largest ones to species of the first growth forms, the intermediate values to those of the second, and the smallest values to species of the third. Thus, species with fast growth and small carbon tended to have higher carrying capacity than slow growing species with high carbon value (reflecting the realistic differences in wood densities and carbon storage among woody species). Examples of the simulated species' natural abundances resulting from both habitat suitability and species-specific carrying capacity are shown in Fig. 1d.

Disturbance and restoration costs – We simulated spatially heterogeneous disturbance through random multivariate normal distributions rescaled to range between 0 and 1, which we multiplied by exp(-rescaled elevation) to simulate a scenario in which anthropogenic disturbance decreases on average with elevation (given the increased difficulties in accessing natural resources or performing agriculture; see example of simulated disturbance in Fig. 1c). We assigned a sensitivity to disturbance for each species based on a symmetric U-shaped beta distribution:

$$S_s \sim B(0.5, 0.5)$$

thus generating several species with either very low or very high sensitivity. Prior to applying any restoration policy, the environment was set to evolve for a number of steps until reaching an equilibrium under the simulated disturbance.

The restoration costs were linked to disturbance, so that on average the cost of restoration was higher for highly degraded cells (reflecting lesser opportunities for natural regeneration and higher distance to seed sources). We added some stochasticity in this pattern by computing costs as the product between disturbance and another rescaled random multivariate normal distribution (see example of simulated disturbance in Fig. 1c). We set the top 5% cells with the highest cost as unrestorable, further raising their cost to exceed the budget. These cells are meant to represent

areas where restoration is not possible e.g. because that would imply displacing inhabited areas or activities and infrastructure that cannot be moved.

Evolution of the species extinction risk – We initialized the species extinction risk based on their range size, on their sensitivity, and on the effect of disturbance:

$$r = \left(\frac{A_s}{\max(A_s)}\right)^l \times \frac{A_s(1 - S_s \delta)}{A_s} \tag{13}$$

where A_s is the natural range of the species (number of occupied cells) in the absence of disturbance. Thus, the value r decreases with small natural range sizes and with increasing sensitivity and/or disturbance (Strassburg et al., 2020). The exponent l (here set to $l = 0.3$) has the effect of flattening the difference in initial extinction risk based on natural range size. We then mapped the value to an initial extinction risk $X_t = 0$ (Table 1). We note that this approach to simulate species extinction risk, while based on common assumptions about species range size and decline due to anthropogenic disturbance, is arbitrary and should be replaced with actual risk assessments (e.g. from the IUCN Red List) when running the optimization with real world data.

Throughout the simulations, the extinction risks of all species were re-evaluated at each step, and we used two criteria to re-assess the conservation status of a species. First, we calculated the number of restored cells within which a species s occurred at time t, indicated as $O_{s,t} \cap M_t$ relative to the species natural range size A_s. Thus, increasing values of $(O_{s,t} \cap M_t)/A_s$ indicate that a proportionally higher number of restored cells is occupied (potentially after re-colonization) by species s, leading to a decreased extinction risk (Table 1, Criterion 1). Second, we calculate the ratio between the occupied geographic range at time t and the occupied range at the beginning of the simulation $O_{s,t} = 0$, i.e. prior to enforcing restoration programs. This ratio is used to verify if the species is in decline, e.g., if the restoration is ineffective or if other factors, such as climate change, negatively affect a species' range. We used the thresholds reported in Table 1 (Criterion 2) to determine if the species was in decline and to what extent that affected the species extinction risk. Finally, any species among the initial simulated pool with occupied range size of 0 was moved to the highest risk category and considered as possibly extinct in the study system. These criteria are inspired by some of the Red List guidelines (IUCN Standards and Petitions Committee, 2025), specifically looking at species range sizes, and recovery or declining trends through time. While we used the occupied range as the main metric, the criteria can be defined based on alternative metrics, e.g., based on census population size and its temporal variation or fragmentation of the species range.

Observed features – The agent was set to observe the entire environment at each step, extracting several features for each cell, which form the input of the neural network and determine the next action. We used the same 16 features for all policies, independently of the

Table 1: Thresholds used to set species extinction risks and their reassessment.

$X_{s,t=0}$	r	Criterion 1	$(O_{s,t} \cap M_t)/A_s$	Criterion 1	$O_{s,t}/O_{s,t=0}$
LC	>0.95	$X_{s,t} = X_{s,t=0}$	< 0.10	$X_{s,t} = X_{s,t=0}$	< 0.10
NT	0.9-0.95	$X_{s,t} = X_{s,t=0} - 1$	0.10 - 0.25	$X_{s,t} = X_{s,t=0} + 1$	0.10 - 0.25
VU	0.8-0.9	$X_{s,t} = X_{s,t=0} - 2$	0.25 - 0.50	$X_{s,t} = X_{s,t=0} + 2$	0.25 - 0.50
EN	0.6-0.8	$X_{s,t} = X_{s,t=0} - 3$	0.50 - 0.80	$X_{s,t} = X_{s,t=0} + 3$	0.50 - 0.80
CR	<0.6	$X_{s,t} = X_{s,t=0} - 4$	> 0.80	$X_{s,t} = X_{s,t=0} + 4$	> 0.80

reward. The first features were the disturbance in the cell and the mean disturbance in the 5×5 cells around it (features 1-2). The latter feature provides a measure of the spatial distribution of disturbance, for instance, differentiating between a disturbed cell neighboring a pristine area and one placed within a broader degraded area. We then included the potential carbon net change in the cell calculated as the difference between the total carbon in the cell under natural conditions and the current amount of carbon, i.e. $C_{c,\text{natural}} - C_{c,t}$. The same measure is also provided as a mean value in the 5×5 cells around the cell (features 3-4). The features additionally include the number of CR, EN, VU and NT species in the cell based on the current populations occupying the cells and the mean number of CR and EN in the surrounding 5×5 cells (features 5-10). We also included the number of species in each of the five extinction risk categories, but counting only species that do not occur in any of the previously restored cells, if any (features 11-15). These features serve as a measure of complementarity among cells so that the model can choose to restore cells with a different species composition compared with areas already restored. Finally, we included the cell's cost (feature 16). All features were rescaled between 0 and 1 before being fed to the neural network.

Training – We trained models with 250 species running simulations for 30 time steps (such as years). The planner was set to restore 100 cells at each step, i.e., reaching 3,000 restored cells at the end of the simulation, i.e., corresponding to about 30% of the land cells in our grid to mimic the 30% restoration target of the Kunming-Montreal Global Biodiversity Framework. We used a parallel genetic strategies algorithm with 8 environments evolving in parallel and ran 5,000 epochs to train the models. We used small neural networks for each cell, connecting the 16 features to a first hidden layer of 3 nodes, followed by a second hidden layer of 2 nodes. The parameters of the network were shared among all cells. We trained different models exploring a gradient of optimality (Lin et al., 2019) going from favoring the carbon outcome to favoring the biodiversity outcome. We achieved this by assigning different weights to the reward (see Equation 12). Specifically, we trained five models, keeping the weight of the budget reward constant at 0.5 and assigning different weights to carbon and biodiversity such that the sum of the weights was kept constant and equal to 1 (Table 2).

Experiments – After training the models, we used them on 100 simulations each including 500 species and initialized following the same approach utilized to generate the training data.

Table 2: Weights used in the different trained models and assigned to the carbon reward (w_C), biodiversity reward (w_X), and budget reward (w_ε). We considered a range of models maintaining constant the weight on the budget reward, while progressively moving from a carbon-focused optimization (Policy 1) to a biodiversity-focused one (Policy 5). The intermediate policies allowed us to explore the trade-offs between these potentially competing objectives.

	w_C	w_X	w_ε
Policy 1	0.5	0	0.5
Policy 2	0.4	0.1	0.5
Policy 3	0.25	0.25	0.5
Policy 4	0.1	0.4	0.5
Policy 5	0	0.5	0.5

Although the 100 simulations were based on the same simulation settings, they differed among each other being based on different random draws for all parameters. Thus, the distribution of species, disturbance, and costs and the traits associated with each species (carbon, sensitivity, growth rates) varied randomly across simulations. This allowed us to evaluate the results of different models across a range of different The simulated environments in their initial state, i.e. prior to restoration, included on average 30 CR species (standard deviation, sd: 18), 87 EN (sd: 39), 156 VU (sd: 31), 139 NT (sd: 30), and 87 LC (sd: 56) species. As during training, the simulated datasets were then let evolve for 30 time-steps with the planner interacting with the environment based on each of the five trained policies and restoring 3,000 cells. We evaluated the outcomes of the different policies by calculating the amount of carbon gained through restoration, the changes achieved in terms of number of species falling in each of the five extinction risk categories, the number of species occurring within restored areas at the end of the simulation, and the total cost of restoration. This allowed us to quantify the trade-off between carbon and biodiversity outcomes along a gradient of optimal solutions.

Carbon and biodiversity credits

We implemented a simplified concept of carbon credits by assigning a monetary value to the amount of net carbon gain (or potentially loss) incurred during the simulations. As carbon (like restoration costs and budget) is set in arbitrary units in our simulations, we defined the value of a carbon credit unit (χ) as the value of the net carbon gain obtained through Policy 1, namely the policy placing the highest weight on carbon, leading to the full compensation of the restoration costs:

$$\chi \frac{\sum_c C_{c,t} - \sum_c C_{c,t=0}}{\sum_c \sum_t \varepsilon_{c,t}} := 1 \tag{14}$$

where χ is the value of one carbon credit unit based on the simulated systems used for carbon and budget, the fractions numerator is the net carbon gain at time t compared to the initial

Figure 1: Maps of the physical properties of the arbitrary area used for simulations, based on the topography and climate of a portion of Central and South America (a). The area was gridded into 128×128 cells and included empirical climate and elevation layers, while the 'natural' state of biodiversity and carbon (b) and the disturbance patterns and costs (c) were simulated and changed randomly across simulations. Species richness and carbon maps are obtained after stacking 500 simulated species (see examples in d). Note that only the map in panel a) is based on real data, while all other the panels show simulated data.

state ($t = 0$) and the denominator is the total cost of restoration. Thus, under this definition, the carbon credits accumulated through the implementation of Policy 1 results, by construction, in full compensation of the cumulative restoration costs, with no economic loss at the end of the simulation. We note that this definition is arbitrary and could be replaced by other definitions, including empirical valuations of carbon in the cap-and-trade system of the European Union, or others.

The same carbon credit value χ was then applied to the other four policies where carbon was given lower weights (Policies 2–5) in the training phase with the inclusion of a positive weight assigned to the biodiversity reward. This allowed us to evaluate to what extent the carbon credits accumulated through restoration policies that optimized for both carbon and biodiversity (Policies 2–4) or biodiversity only (Policy 5) were able to compensate for the restoration costs.

Next, we selected a simple metric as a reference to define a biodiversity credit and developed an approach to valuate the credit relative to the value of a carbon credit. Specifically, we quantified the biodiversity credits as the number of species that moved from a threatened class (CR, EN, VU) at the beginning of the simulation to a non-threatened class (NT, LC) after restoration. This differs to current practice in some of the emerging biodiversity credit markets, such as in Colombia where one biodiversity credit unit equals one hectare regardless of how many species were benefited by the restoration and in which ways. However, the same approach utilized here can be used for alternative definitions of biodiversity credits.

We then evaluated the net financial loss or gain of each policy after stacking the value of carbon and biodiversity credits under different valuations of biodiversity credits. We indicate with β the value of one biodiversity credit obtained from moving a species from a threatened to a non-threatened status after restoration. Given that we are using arbitrary units for costs and carbon credits, we defined the value of a biodiversity credit relative to that of a carbon credit. This allowed us to assess which policy would become financially viable or preferable if e.g. the value of one biodiversity credit equals that of a carbon credit (i.e. $\beta = \chi$). We tested valuations of $\beta = 0, \chi, 2\chi,$ and 4χ and calculated the net financial loss or gain of a policy (Φ) as a percentage of the restoration cost:

$$\Phi_{\text{Pol. } i} = \left(\frac{\chi C_{\text{Pol. } i} + \beta S_{\text{Pol. } i}}{\varepsilon_{\text{Pol. } i}} - 1 \right) \times 100 \tag{15}$$

where $C_{\text{Pol}.i}$ is the net carbon gain of the i^{th} policy, $S_{\text{Pol}.i}$ is the policy's net biodiversity outcome (as defined above), and $\varepsilon_{\text{Pol}i}$ is the cost of restoration.

Finally calculated, for any given policy i, the value that should be attributed to a biodiversity credit so that the policy results in the same net financial outcome as a restoration program focused on carbon (i.e. our Policy 1):

$$\frac{\chi C_{\text{Pol. } 1} + \beta S_{\text{Pol. } 1}}{\varepsilon_{\text{Pol. } 1}} := \frac{\chi C_{\text{Pol. } i} + \beta S_{\text{Pol. } i}}{\varepsilon_{\text{Pol. } i}} \tag{16}$$

The value of one biodiversity credit is therefore, after solving Equation 16 for β:

$$\beta = \chi \frac{C_s \varepsilon_c - C_c \varepsilon_s}{S_c \varepsilon_s - S_s \varepsilon_c} \tag{17}$$

Using this formula, we derive the value of a biodiversity credit (relative to the value of a carbon credit) required to make the financial outcome (defined in Eq. 15) of any optimal restoration policy equal to the profit of a carbon-oriented policy (our Policy 1), that is such that $\Phi_{\text{Pol. } i} = \Phi_{\text{Pol. } 1}$. We note that under this definition, the value of biodiversity credits is context specific, as it depends on the overall area considered and its restorable carbon and biodiversity. Within the area, the restoration of cells that contribute the most to reducing the extinction risk of species (e.g., cells that protect rare and threatened species) will lead to the highest biodiversity credits (Drupp et al., 2024).

Results

Optimization of restoration policies and trade-offs

Models trained through multi-objective reinforcement learning using differently weighted rewards ranged from one focusing on carbon (Policy 1) to one targeting biodiversity outcomes

(Policy 5), with intermediate models exploring the gradient of solutions between the two extremes. All models also included a budget reward to attempt to minimize the restoration costs (Table 2), thus all policies targeted two to three objectives.

The five trained policies applied on a set of 100 simulations resulted in substantial spatial differences in the prioritization of restoration areas and led to widely different carbon and biodiversity outcomes. Since all simulations differed in the distribution of species, threats, and costs the placement of the selected restored areas varied among them. However, if we look at an individual simulation, it is apparent that the policies prioritize different areas (Fig. 2). Policy 1 favored the restoration of areas that, in the specific settings of this simulated dataset, resulted in the highest net carbon gain, as shown by the resulting steady increase of carbon in the system (Fig. 3a) that exceeds a 20% net gain over the initial amount at the end of the 30 times steps. In contrast, Policy 5 prioritized areas that contributed the most to reducing species extinction risk, therefore considering threats, species richness, and complementarity of species composition (Fig. 2e). The result was a more modest carbon gain (just above 10% in the simulation shown in Fig. 3c) but a substantially stronger reduction of the extinction risk compared with Policy 1. For instance, the number of critically endangered species at the end of the simulation was 50% lower under Policy 5 compared to Policy 1 (Fig. 3). Similarly, the number of endangered species decreased by 70% when giving priority to biodiversity over carbon. These experiments also showed that while carbon accumulated following simple almost linear trajectories, changes in species extinction risks followed more complex pathways, reflecting the fact that species moved across rick categories and that their conservation status was determined by their ability to re-colonize restored areas based on their dispersal ability and connectivity and disturbance among areas.

When looking at the outcome at the end of the simulated 30 time-step restoration, we can appreciate the variability among simulations and the consistency in the patterns observed at the level of an individual simulation. The net carbon gain in Policy 1, which was trained without considering a biodiversity reward, resulted in an average 2.2-fold increase (standard deviation among simulations: ±0.26) in carbon storage compared to Policy 5, which focuses on biodiversity but does not include carbon in the reward (Fig. 4a). Optimized policies with positive weights associated with both carbon and biodiversity rewards revealed a gradient of outcomes with Policy 3, which assigned equal weights to carbon and biodiversity, resulting in intermediate levels of carbon net gain (Figs. 3b, 4a).

Conversely, the biodiversity outcomes revealed that a carbon-focused policy resulted in significantly worse impacts on extinction risks compared to most other policies (Fig. 4b, d–i). For instance, whereas on average 30% of the species decreased their extinction risk under Policy 1 (150 out of the 500 simulated species), the percentage raised to 71–76% (354–378 species) under all other policies (Fig. 4b). The fraction of critically endangered species remained virtually

unchanged under Policy 1, while it decreased on average by 29% under Policy 5, and by 27% under all other policies (Fig. 4d). When combining all threatened species (CR, EN, VU), these decreased by about 25% under Policy 1 (i.e. 68 species) but dropped by 62–69% (i.e. 170–190 species) units under the others (Fig. 4i), thus showing a 2.6-fold improvement. These results show that even when biodiversity outcomes are given a smaller weight compared to carbon in the optimization of the restoration policy (e.g., Policy 2) the impact on extinction risk reduction is large.

Finally, the costs of implementation of the different policies, i.e. the cumulative cost of all restored cells, were relatively similar, with changes between 4 and 17% among policies (Fig. 4c). Thus, at least under our simulated scenarios where the same weight was assigned to the budget reward in all policies, the different restoration programs had comparable costs of implementation.

Figure 2: Example of the restored areas in one of the 100 simulations as selected under five policies (Table 2) trained with different weights on carbon (highest in a) and biodiversity outcomes (highest in e). Note that the placement of the restored areas depends on the specific conditions of the simulated dataset. Restored areas will therefore vary among simulations, reflecting the randomly initialized distribution of species, disturbance, and costs. The outcomes of this specific simulation in terms of resulting net carbon gain and changes in species extinction risks are shown in Fig. 3

Carbon credits and valuation of biodiversity credits

Our simulations using multi-objective policies optimized under different targets, they led to vastly different carbon and biodiversity outcomes, but similar implementation costs, prompting the question of whether they are equally feasible in financial terms. We therefore first explored their feasibility in a context in which carbon credits were used to compensate for the restoration costs, under a valuation of carbon credits such that the net carbon obtained from Policy 1 results in full compensation of the costs.

Given our definition of carbon credit (Eq. 14), Policy 1 resulted in the cumulative value of carbon credits compensating exactly the costs of restoration (Fig. 5a). In contrasts, all other policies resulted in a net loss that ranged from 10% on average in Policy 2 (optimized with a weight on carbon four times higher than on biodiversity) to about 50% in Policy 5 (Fig. 5a).

Figure 3: Accumulation of carbon and evolution of extinction risks across 30 time-steps in one of the 100 simulations. The results are shown for Policy 1 where the reward focused on the carbon outcomes (a); Policy 3 where equal weights were assigned to carbon and biodiversity rewards (b); and Policy 5 in which the reward focused on biodiversity with no input from carbon. The plots show that when biodiversity is factored in, the carbon accumulation curve reaches lower levels decreasing from ca. 22% (a) to 14% (b) and 10% (c). This trend is counterbalanced by a substantial improvement of species conservation status, with a ~30–50% drop in critically endangered species.

Figure 4: Summary of 100 simulations run under five trained restoration models, Policies 1–5. The net carbon (a) represents the net gain in amount of carbon (here expressed in arbitrary units – see Methods) in the system after 30 time-steps during which 3,000 cells were restored. As the weight assigned to the carbon reward decreases from 0.5 to 0 in favor of the biodiversity reward, the net carbon gain decreases too, although we note that all restoration policies resulted in a net gain. In contrast, the number of species with a reduced extinction risk at the end of the simulation compared with their initial status quickly increased as soon as a non-zero weight was assigned to the biodiversity reward (Policies 2–5; b). Yet, the implementation cost was similar across all policies (c), reflecting the fact that all models were trained with the same weight assigned to the budget reward. Plots d–h) show the difference between the number of species in each extinction risk category, going from highest risk (CR) to lowest (LC), at the beginning of the simulation and the corresponding number at the end of the simulation. Thus, negative numbers, here observed in d–g) mean that the number of species in the corresponding risk categories decreased. Positive numbers, here mostly observed in h), mean that the number of species in these categories increased. These plots show that all policies that considered biodiversity rewards in the optimization (Policies 2–5) resulted in substantial reductions of threatened species compared to a policy targeting carbon only. This is also evident when lumping together all threatened species (i).

The results changed substantially when assigning a non-zero value to biodiversity credits (while leaving unchanged the value of carbon credits). When setting the value of one biodiversity credit equal to that of a carbon credit (i.e. $\beta = \chi$), both Policy 1 and 2 became profitable with financial gains above 10% for the latter. Under this valuation, the implementation of Policies 3–5 continued to lead to significant financial loss (Fig. 5b). Increasing the relative value of biodiversity credits ($\beta = 2\chi$ and $\beta = 4\chi$) led to improved financial outcomes of all policies (Fig. 5c,d). Yet, we found that Policy 2 consistently led to the best outcomes, reflecting the non-linear improvement in biodiversity outcomes observed in Fig. 4b.

We quantified the value of a biodiversity credit such that a biodiversity-oriented policy reached a net monetary profit relative to restoration costs matching that of a policy designed to maximize carbon only (Policy 1; Eq. 17). We found that policies 3–5 could all reach profits equal to Policy 1 with a similar median biodiversity credit value of about $\beta = 3.2 - -4.4\chi$, while for Policy 2, a value of $\beta = 1.2\chi$ was sufficient. This means that if a biodiversity credit is valued at 1.2 times the value of a carbon credit, then Policies 1 and 2 result in the same net relative profit based on the cumulative carbon and biodiversity credits. These results indicate that the value of biodiversity credits converge to a relatively constrained range across different optimal policies in which biodiversity outcomes are assigned different weights. Consequently, for a given valuation of biodiversity credits there is some room for favoring carbon or biodiversity in restoration programs with similar economic incentives. In contrast assigning a null value to biodiversity credits ($\beta = 0$) leaves limited options to target biodiversity in restoration programs without drastically the financial viability of the policy (Fig 5a).

Figure 5: Financial outcomes (Φ) of restoration policies (net percentage loss or gain) under different valuations of biodiversity credits. Negative values indicate a net economic loss where restoration costs exceed the cumulative value of carbon and biodiversity credits, while positive values indicate a net gain. The results are shown when only carbon credits are included (i.e. $\beta = 0$; a), and when biodiversity credits are valuated at $\beta = \chi, 2\chi$ and 4χ, respectively (b–d).

Discussion

Reinforcement learning as tool for guiding restoration programs

We developed a multi-objective reinforcement learning approach, building upon the CAP-TAIN software (Silvestro et al., 2022), to optimize ecological restoration strategies targeting climate and biodiversity positive outcomes while accounting for costs of implementation. The simulation and optimization framework allows us to evaluate the trade-offs among carbon, costs, and species protection and conservation status. Using weighted rewards in the training phase allowed us to calibrate models that explore the boundary of optimality (Lin et al., 2019) across a set of weights assigned to the net carbon gain and to the reduction in extinction risks obtained through restoration.

As our models are trained within a dynamic environment, the solutions incorporate the effects of connectivity among restored areas, that favor migration processes and range expansion underlying the simulation of species recovery. They also account for species-specific features such as climatic niche, growth rates, population density, and sensitivity to anthropogenic disturbance. The simulations show that the spatial distribution of biodiversity and carbon priorities differ, as observed in real-world scenarios (Jung et al., 2021; Alejo et al., 2025). This mismatch means that a restoration program focusing on the maximization of carbon will likely not result in the best biodiversity outcome.

Toward a data-driven valuation of biodiversity credits

Our results show that simulation experiments coupled with multi-objective reinforcement learning optimization allow us to explore the trade-offs between carbon and biodiversity outcomes within the constraints of a limited restoration budget. We propose that this systematic evaluation of the predicted restoration outcomes and associated costs can be used as a data-driven approach to valuate biodiversity credits relative to carbon credits, such that restoration policies targeting both carbon fixation and extinction risk reduction become financially sustainable. We showed that even a relatively small value assigned to biodiversity outcomes can create incentives favoring a multi-objective restoration policy.

While we demonstrated our approach with a fixed definition of biodiversity credits (based on species extinction risks), our approach could also be used to evaluate the effect of different metrics as the basis for biodiversity crediting. For instance, other approaches could be based on population recovery or geographic range expansion (Farnsworth et al., 2015; Schipper et al., 2020). In a simulated setting, virtually any metric can be tested as we can afford to quantify the evolution of an environment through restoration at any level of demographic and spatial detail. In practice, however, the reference metric will have to be chosen so that it can be

monitored accurately and continuously in the real world, for instance, through the re-assessment of extinction risks of selected species (Durán et al., 2020; Antonelli et al., 2023).

Biodiversity credits can be defined such that an optimal solution for both carbon and biodiversity is preferable compared to a policy targeting carbon alone. Unlike carbon, a biodiversity outcome (and valuation) is inherently context dependent (Farnsworth et al., 2015). That is because a ton of fixated carbon provide the same contribution to climate wherever it is sequestrated (except for potentially losing value through time if the sequestration is not permanent (Groom and Venmans, 2023)), while the value of the restoration and recovery of one species is necessarily linked to where the species occurs and its environmental context. Estimating trade-offs can help us defining actionable and context-dependent valuation for biodiversity credits. In our experiments, we found that setting the value of one species recovering from threatened to a non-threatened status to between 1 and 4 carbon credits was required to make a biodiversity-oriented policy financially competitive against a carbon-focused one. The value will likely differ in different geographic contexts and depending on the spatial extent considered and the selection of reference species. Yet, this approach to value biodiversity credits relative to carbon credits can be in principle applied to different contexts and different quantities used to measure a biodiversity credit.

Applicability to real-world scenarios

The current framework could be applied to real world scenarios by replacing some variables with actual data. For instance, species distribution models based on species occurrence data and biodiversity monitoring can be used to replace the simulated biodiversity data and habitat suitability in our model. Additionally, carbon estimates in both degraded and restored areas can be used to quantify the amount of fixated carbon in disturbed habitats and its potential amount after restoration of natural vegetation. Further, real land use and predicted restoration costs can replace the respective simulated variables in our reinforcement learning framework. Extinction risks can be initialized based on existent or predicted species extinction risks evaluated at a global or regional level (Red List of Threatened Species, 2023; Zizka et al., 2022). In our simulations, we used simple rules based on simulated population recovery to model the changes in extinction risks after implementing restoration policies. These rules can be adapted to reflect the actual criteria used to evaluate the conservation status of species, for instance, based on IUCN guidelines for global assessments or other criteria implemented in regional Red Lists (IUCN Standards and Petitions Committee, 2025). Our approach can therefore, be informed by actual data and provide restoration priorities while leveraging potentially conflicting targets for carbon sequestration and biodiversity outcomes. Its application to real-world scenarios will likely result in different valuations of biodiversity credits, reflecting their context-dependent nature. Yet, these valuations will be obtained through a data-driven approach providing an robust way to assign a monetary value to something as complex and multifaceted as biodiversity.

Conclusion

The controlled simulations performed here indicate that it is possible to select areas for ecological restoration that substantially benefit both climate and biodiversity. Disregarding one or the other would result in missed opportunities, given the relatively small reduction of carbon capture or biodiversity protection associated with a multi-objective prioritization. Future work could examine the performance of this approach applied to empirical data, in comparison with other data-driven or expert-led prioritization approaches. As ecosystem restoration is increasingly recognized as a crucial step in the conservation of nature and stabilization of Earth's climate (Strassburg et al., 2020; Seddon et al., 2020), the use of robust models and AI-based optimization able to capture the complexity of biodiversity and its spatial and temporal dynamics can play an important role in guiding our efforts to prevent further extinctions and tackle climate change.

Funding

A.A., B.G. T.S. and D.S. are members of the BIOPATH research programme funded by the Swedish Foundation for Strategic Environmental Research MISTRA (F 2022/1448). A.A. is also funded by the Swedish Research Council (VR: 2019-05191, 2024-04303) and the Royal Botanic Gardens, Kew; B.G. is funded by Dragon Capital and the NERC/UKRI projects BIOADD (ref: NE/X002292/1), BIOESG (NE/X016560/1) and RENEW (ref: NE/W004941/1); D.S. by ETH Zurich and the Swiss National Science Foundation (PCEFP3_187012); and T.S. by the Kamprad Foundation. The funders had no role in the contents of this study.

*

References

Alejo, C., A. Luers, A. Ventimiglia, and H. D. Matthews. 2025. Maximizing nature-based solutions using artificial intelligence to align global biodiversity, climate, and water targets. bioRxiv Pages 2025–01.

Antonelli, A., K. L. Dhanjal-Adams, and D. Silvestro. 2023. Integrating machine learning, remote sensing and citizen science to create an early warning system for biodiversity. Plants, people, planet 5:307–316.

Antonelli, A., X. Rueda, R. Calcagno, and P. Nantongo Kalunda. 2024. How biodiversity credits could help to conserve and restore nature. Nature 634:1045–1049.

Apicella, A., F. Donnarumma, F. Isgrò, and R. Prevete. 2021. A survey on modern trainable activation functions. Neural Networks 138:14–32.

Burgess, N. D., N. Ali, J. Bedford, N. Bhola, S. Brooks, A. Cierna, R. Correa, M. Harris, A. Hargey, J. Hughes, et al. 2024. Global metrics for terrestrial biodiversity. Annual Review of Environment and Resources 49:673–709.

Deutz, A., G. M. Heal, R. Niu, E. Swanson, T. Townshend, L. Zhu, A. Delmar, A. Meghji, S. A. Sethi, and J. Tobin-de la Puente. 2020. Financing nature: Closing the global biodiversity financing gap. The Paulson Institute, The Nature Conservancy, and the Cornell Atkinson Center for Sustainability 256.

Di Sacco, A., K. A. Hardwick, D. Blakesley, P. H. Brancalion, E. Breman, L. Cecilio Rebola, S. Chomba, K. Dixon, S. Elliott, G. Ruyonga, et al. 2021. Ten golden rules for reforestation to optimize carbon sequestration, biodiversity recovery and livelihood benefits. Global Change Biology 27:1328–1348.

Drupp, M., M. Hänsel, E. Fenichel, M. Freeman, C. Gollier, B. Groom, G. Heal, P. Howard, A. Millner, F. Moore, et al. 2024. Accounting for the increasing benefits from scarce ecosystems. Science 383:1062–1064.

Durán, A. P., J. M. Green, C. D. West, P. Visconti, N. D. Burgess, M. Virah-Sawmy, and A. Balmford. 2020. A practical approach to measuring the biodiversity impacts of land conversion. Methods in Ecology and Evolution 11:910–921.

Eyres, A., T. Ball, M. Dales, T. Swinfield, A. Arnell, D. Baisero, A. P. Durán, J. Green, R. E. Green, A. Madhavapeddy, et al. 2024. Life: A metric for quantitively mapping the impact of land-cover change on global extinctions. Earth and Environmental Sciences .

Farnsworth, K. D., A. H. Adenuga, and R. De Groot. 2015. The complexity of biodiversity: A biological perspective on economic valuation. Ecological Economics 120:350–354.

Groom, B. and F. Venmans. 2023. The social value of offsets. Nature 619:768–773.

Gumbs, R., C. L. Gray, M. Böhm, I. J. Burfield, O. R. Couchman, D. P. Faith, F. Forest, M. Hoffmann, N. J. Isaac, W. Jetz, et al. 2023. The edge2 protocol: Advancing the prioritisation of evolutionarily distinct and globally endangered species for practical conservation action. PLoS Biology 21:e3001991.

Holl, K. D. and P. H. Brancalion. 2020. Tree planting is not a simple solution. Science 368:580–581.

Hua, F., L. A. Bruijnzeel, P. Meli, P. A. Martin, J. Zhang, S. Nakagawa, X. Miao, W. Wang, C. McEvoy, J. L. Peña-Arancibia, et al. 2022. The biodiversity and ecosystem service contributions and trade-offs of forest restoration approaches. Science 376:839–844.

IUCN Standards and Petitions Committee. 2025. Guidelines for Using the IUCN Red List Categories and Criteria. Version 16 (http://www.iucnredlist.org/documents/RedListGuidelines.pdf).

Jung, M., A. Arnell, X. De Lamo, S. García-Rangel, M. Lewis, J. Mark, C. Merow, L. Miles, I. Ondo, S. Pironon, et al. 2021. Areas of global importance for conserving terrestrial biodiversity, carbon and water. Nature Ecology & Evolution 5:1499–1509.

Karger, D. N., M. P. Nobis, S. Normand, C. H. Graham, and N. E. Zimmermann. 2023. Chelsa-trace21k–high-resolution (1 km) downscaled transient temperature and precipitation data since the last glacial maximum. Climate of the Past 19:439–456.

Lin, X., H.-L. Zhen, Z. Li, Q.-F. Zhang, and S. Kwong. 2019. Pareto multi-task learning. Advances in neural information processing systems 32.

Mair, L., L. A. Bennun, T. M. Brooks, S. H. Butchart, F. C. Bolam, N. D. Burgess, J. M. Ekstrom, E. Milner-Gulland, M. Hoffmann, K. Ma, et al. 2021. A metric for spatially explicit contributions to science-based species targets. Nature Ecology & Evolution 5:836–844.

Parr, C. L., M. Te Beest, and N. Stevens. 2024. Conflation of reforestation with restoration is widespread. Science 383:698–701.

Pimiento, C., F. Leprieur, D. Silvestro, J. Lefcheck, C. Albouy, D. Rasher, M. Davis, J. Svenning, and J. Griffin. 2020. Functional diversity of marine megafauna in the anthropocene. sci adv 6: eaay7650.

Red List of Threatened Species. 2023. Red List of the International Union for the Conservation of Nature (http://www.iucnredlist.org).

Schipper, A. M., J. P. Hilbers, J. R. Meijer, L. H. Antão, A. Benítez-López, M. M. de Jonge, L. H. Leemans, E. Scheper, R. Alkemade, J. C. Doelman, et al. 2020. Projecting terrestrial biodiversity intactness with globio 4. Global Change Biology 26:760–771.

Seddon, N., A. Chausson, P. Berry, C. A. Girardin, A. Smith, and B. Turner. 2020. Understanding the value and limits of nature-based solutions to climate change and other global challenges. Philosophical Transactions of the Royal Society B 375:20190120.

Silvestro, D., S. Goria, B. Groom, T. Sterner, and A. Antonelli. 2023. The 30 by 30 biodiversity commitment and financial disclosure: Metrics matter. Available at SSRN 4594937 Page 10.2139/ssrn.4594937.

Silvestro, D., S. Goria, T. Sterner, and A. Antonelli. 2022. Improving biodiversity protection through artificial intelligence. Nature sustainability 5:415–424.

Strassburg, B. B., A. Iribarrem, H. L. Beyer, C. L. Cordeiro, R. Crouzeilles, C. C. Jakovac, A. Braga Junqueira, E. Lacerda, A. E. Latawiec, A. Balmford, et al. 2020. Global priority areas for ecosystem restoration. Nature 586:724–729.

Zizka, A., T. Andermann, and D. Silvestro. 2022. Iucnn–deep learning approaches to approximate species' extinction risk. Diversity and Distributions 28:227–241.