BRIEF REPORT

Complete mitochondrial genome assembly of two searobin species

(genus *Prionotus*) and their phylogenetic relationships (Triglidae:

Prionotinae)

Alan Marín[1], Ruben Alfaro[2], Eliana Zelada-Mázmela[1]

[1] *Laboratorio de Genética, Fisiología y Reproducción, Facultad de Ciencias, Universidad*
Nacional del Santa, Chimbote, Perú

[2] *Ecobiotech Lab S.A.C. Trujillo, Perú*

25 Corresponding author: Alan Marín amarin@uns.edu.pe

Abstract

Background The genus *Prionotus* "searobins" are marine ray-finned fish of the subfamily Prionotinae that have evolved distinctive morphological features and specialized behavior for walking over the seafloor detecting buried preys, making them excellent candidates for biological models in evolutionary genetics studies. Despite that, their phylogenetic relationships are still not well understood. Here, we assembled the first complete mitochondrial genomes of two searobin species namely *Prionotus alatus* and *P. stephanophrys* and performed a comparative mitogenome examination including the first phylogenetic analysis of Prionotinae based on complete mitogenomic information.

Methods and results Both mitogenomes were assembled using Illumina genomic reads mined from the GenBank database. Single circular contigs of 16,602 and 16,896 bp were obtained for *P. alatus* and *P. stephanophrys* respectively. Both mitogenomes displayed a standard vertebrate organization (13 PCGs, 2 rRNAs, 22 tRNAs, and a putative control region) except for a third tRNA-Leu and an extra non-coding region detected in *P. stephanophrys*. A Bayesian phylogenetic analysis recovered all *Prionotus* species into a monophyletic clade within Prionotinae, including the sister taxon *Bellator* grouped in a discrete subclade. Species of the subfamily Triglinae were resolved in a basal monophyletic group within Triglidae.

Conclusion The results presented herein represent an important step into a deeper understanding of the phylogenetic relationships and evolution of *Prionotus*. Taxonomic clarification may also support future management and conservation efforts of these economically valuable species.

Keywords: Mitogenomics, Phylogeny, Scorpaenoidei, GenBank data mining

Introduction

Species of the genus *Prionotus* Lacepède, 1801 (Triglidae: Prionotinae), commonly known as

"searobins", not only support important fisheries throughout their distribution ranges but also

possess specialized morphological and biological characteristics such as wing-like pectoral fins

with three modified lower rays used to "walk" along the sea bottom and certain species use

those rays to detect buried preys due to their sensitiveness to chemical and mechanical stimuli

(Portnoy et al., 2017; Allard et al., 2024). Such unique features make them ideal biological

models for evolutionary genetics and adaptation studies (Herbert et al., 2024).

According to the World Register of Marine Species (WORMS) database, there are currently

23 extant *Prionotus* species (Froese and Pauly, 2025), whose natural distribution is restricted

to the Americas (Portnoy et al., 2017), including 16 western Atlantic and 7 eastern Pacific

species. Nevertheless, taxonomic research on the classification of *Prionotus* remains scarce,

and only a few studies have addressed their taxonomy using morphological characters

(Imamura, 1996; Richards and Jones, 2002), while only two studies performed molecular

phylogenetic analysis employing information from partial mitochondrial and nuclear gene

fragments (Portnoy et al., 2017; Smith et al., 2018). Although there has been some degree of

consistency between the results of prior morphological and molecular phylogeny studies, the

molecular phylogenetic results of Portnoy et al. (2017) rendered *Prionotus* as paraphyletic

within Triglidae, suggesting that the currently accepted genus *Bellator*, its only sister genus

within the subfamily Prionotinae, should be considered as a junior synonym of *Prionotus* until

further studies using robust phylogenetic analysis and broader species coverage shed much

additional light on the taxonomic status of Prionotinae species.

Since phylogeny relationships inferred from morphological characters and/or partial gene

fragments may lead to limited taxonomic resolution, the use of more robust genome-wide data

has become necessary to resolve complex phylogenetic problems in fish lineages (Miya and

Nishida, 2015). In this context, nucleotide information obtained from the complete

mitogenome has proven effective in resolving the phylogeny of several fish taxa on many

taxonomic levels (Miya and Nishida, 2015). Moreover, the unique properties of the

mitogenome (e.g. fast evolutionary rates, lack of recombination, high copy number) make this

organelle a great source of powerful molecular markers suitable for diverse applications

including accurate species identification, wildlife forensics, environmental DNA, population

genetics and evolutionary studies (Maduna et al., 2022).

Despite the existence of 6 mitogenomes from Atlantic *Prionotus* species available in the

GenBank database and the unresolved taxonomic status of Prionotinae species, no study has

yet analyzed those mitogenomes to investigate their phylogenetic relationships. Furthermore,

the vast amount of genomic next-generation sequencing (NGS) data publicly available within

the GenBank database provides a rich source of organelle genomes that can be screened to

identify fish species of taxonomic interest (Eldem et al., 2024; Marín et al., 2025). The

objective of the present work was: a) to assemble the mitochondrial genomes of two

commercially important searobin species namely *P. alatus* and *P. stephanophrys* using SRA

reads mined from GenBank, b) to describe the mitogenome characteristics and perform

comparative analyses, and c) to investigate the phylogenetic relationships of Triglidae species

using the two *Prionotus* mitogenomes assembled herein and the mitogenomes of all Triglidae

species available in GenBank.

Materials and methods

Genomic NGS reads mining, mitogenome assembly and annotation

The NGS reads used to assemble the two mitochondrial genomes came from two BioSamples

deposited in the GenBank database. BioSample metadata records indicate that the *P. alatus*

(BioSample SAMN36762230) and the *P. stephanophrys* (BioSample SAMN31811746)

individuals were collected from the Atlantic Ocean in USA (34.6094 N, 75.7052 W) on

September 17, 2022, and from the Pacific Ocean in El Salvador (12.7459 N, 89.0088 W) on

December 10, 2010, respectively, by the Smithsonian National Museum of Natural History in

their "NOAA Genome Skimming of Marine animals inhabiting the US Exclusive Economic

Zone" project (BioProject accession PRJNA720393). Paired-end FASTQ genomic reads

(Platform: Illumina NovaSeq 6000, *P. alatus*: 1.9 Gbp within 12.50 million reads; *P.*

stephanophrys: 1.8 Gbp within 12.13 million reads) were downloaded from the public

GenBank SRA repository (*P. alatus* SRA accession SRR25465369; *P. stephanophrys* SRA

accession SRR22396965). Raw reads were subjected to quality control using FastQC and

trimmed with BBDuk as implemented in Geneious Prime v. 2025.0 (Biomatters Ltd., Auckland,

New Zealand) resulting in 12.49 and 12.07 million reads in *P. alatus* and *P. stephanophrys*

respectively. To validate the mitogenomes assembled herein, we obtained consensus

mitogenomes from two different strategies namely "map to reference" approach and a hybrid

"map to reference/de novo" approach. For the first approach, trimmed reads from *P. alatus* and

P. stephanophrys were assembled against the complete mitogenome reference sequences of *P.*

nudigula (GenBank accession NC_088019) and *P. evolans* (GenBank accession NC_083024)

respectively, using the "Map to Reference" tool in Geneious Prime with default parameter

settings recommended by the software developers (Sensitivity: Medium-Low Sensitivity/Fast,

five iterations). For the second approach, the contigs obtained from the first assembly approach

were dissolved and reassembled "de novo" following the parameter settings recommended by

the software developers (Sensitivity: Medium Sensitivity/Fast) and selecting the "circularize

contigs" option. The consensus sequences obtained from both approaches were extracted and

checked manually using the Genious Prime alignment tool.

Gene annotation was performed using the "Annotate and Predict" feature of Geneious Prime

by comparing the novel *P. alatus* and *P. stephanophrys* mitogenomes to other complete

mitogenomes of closely related species. The start and stop codon positions of the protein-

coding genes (PCGs) were verified using the Open Reading Frame Finder tool (available at

https://www.ncbi.nlm.nih.gov/orffinder/) using the vertebrate mitochondrial genetic code. The

positions of the transfer RNA genes (tRNAs) were verified using tRNAscan-SE (Chan and

Lowe 2019). The circular mitogenome figures were generated with Organellar Genome Draw

"OGDRAW" version 1.3.1 (Greiner et al., 2019) and edited with Inkscape macOS version 1.4

(Inkscape Project, 2020).

Bayesian phylogenetic analysis

For the Bayesian phylogenetic analysis, we used the 13 PCGs of the two mitogenomes

assembled herein plus those of all Triglidae species available in GenBank (17 species: 8 from

Prionotinae and 9 from Triglinae). A Hoplichthyidae species, *Hoplichthys gilberti*, was used

as outgroup. Nucleotide sequences from each PCG were multialigned separately using MEGA

7 (Kumar et al., 2016) and concatenated with SeaView version 5.0.4 (Guoy et al., 2021),

resulting in a final matrix of 11,444 nucleotides. The program jModelTest 2 (Darriba et al.,

2012) was used to find the most appropriate model of nucleotide evolution for each one of the

13 PCGs under the corrected Bayesian Information Criterion (BICc). Levels of substitution

saturation at single codon positions from each PCG were assessed with Xia´s method

implemented in DAMBE7 (Xia, 2018). The third codon positions of three PCGs (ATP8,

NADH3, and NADH6) showed clear signs of saturations and thereby the Bayesian analysis

was run with the data partitioned by gene and by codon position excluding the third codon

position of those genes. Bayesian phylogenetic trees were constructed with MrBayes 3.2.7

(Ronquist et al., 2012) as implemented on the CIPRES science gateway 3.3 server (Miller et

al. 2010), using two independent runs of four Markov chains each, conducted for 10,000,000

generations, with samplings frequency of 1000. The first 25% of the sampled trees were

discarded as burn-in. The final consensus phylogenetic tree was drawn using Figtree version

1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

Results and discussion

In this study, we successfully assembled and annotated the first mitochondrial genomes of the

Atlantic spiny searobin *P. alatus* (occurring from the western Atlantic from Virginia in the

USA to the Campeche Shelf in Mexico) and the Pacific lumptail searobin *P. stephanophrys*

(occurring from the Columbia River in Washington in the USA to Ilo in Peru) (Russell et al.,

1992; Chirichigno and Cornejo, 2001). There were no differences between the consensus

mitogenome contigs obtained with the two different assembly approaches applied herein. A

total of 30,235 trimmed reads from *P. alatus* and 24,089 from *P. stephanophrys* were

assembled against the selected reference sequences, allowing the generation of a single circular

contig of 16,602 bp in length (mean per base coverage of 121.2 reads) in *P. alatus* and 16,896

bp (mean per base coverage of 133.1 reads) in *P. stephanophrys* (GenBank accessions pending)

(Fig. 1), which were within the size range of other reported *Prionotus* mitogenomes ranging

from 16,599 bp in *P. nudigula* (NC_088019.1) to 16,907 in *P. evolans* (NC_083024.1). Overall,

the novel mitogenome sequences showed highly similar base frequencies of A=26.3%,

C=29.6%, G=17.3%, and T=26.8% in *P. alatus* and of A=26.6%, C= 29.6%, G=17.6%, and

T=26.2% in *P. stephanophrys*.

Fig. 1. Circular mitochondrial genome organization of A) *P. alatus* (photo kindly provided by

R. Robertson, Robertson and Van Tassell, 2015) and B) *P. stephanophrys* (Photo by R. Alfaro).

Genes inside the circle (negative strand) are transcribed clockwise and genes outside the circle

counter clockwise. The circle inside the GC content graph (innermost circle) indicates the 50%

threshold where darker and lighter grey corresponds to the GC and AT content respectively.

NADH dehydrogenase, cytochrome c oxidase, ATP synthase, transfer RNA, and ribosomal

RNA genes are depicted in yellow, pink, green, purple, blue, and red bars respectively. The bar

representing the control region is shown in coral red color.

Both mitogenomes showed a standard vertebrate organization (i.e. 13 PCGs, 2 rRNAs, 22

tRNAs, and a putative control region) (Winn et al., 2024), except for a third tRNA-Leu and an

extra non-coding region detected in the mitogenome of *P. stephanophrys*. The third tRNA-Leu

was located in the positive strand between a 220 bp non-coding region downstream of the

conserved block NADH4/His/Ser/Leu and the NADH5 gene (Table 1 and Fig. 1). This unusual

193 **Table 1.** Complete mitochondrial annotation and number of amino acids in each of the genes

194 detected in the mitochondrial genomes of *Prionotus alatus* (PA) and *P. stephanophrys* (PS).

195 The plus "+" and minus "-" signs stand for the heavy and light strands respectively. NCR: non-

196 coding region.

Gene	Strand	Location				Size (bp)		Anticodon		START codon		STOP codon		No. of amino acids	
		Start		End											
		PA	PS	PA	PS	PA	PS	PA	PS	PA	PS	PA	PS	PA	PS
Phe (F)	+	1	1	68	68	68	68	GAA	GAA	—	—	—	—	—	—
12S rRNA	+	69	69	1016	1017	948	949	—	—	—	—	—	—	—	—
Val (V)	+	1017	1018	1090	1091	74	74	UAC	UAC	—	—	—	—	—	—
16S rRNA	+	1091	1092	2782	2789	1692	1698	—	—	—	—	—	—	—	—
Leu (L)	+	2783	2790	2856	2863	74	74	UAA	UAA	—	—	—	—	—	—
NADH1	+	2857	2864	3831	3833	975	975	—	—	ATG	ATG	TAA	TAG	324	324
Ile (I)	+	3836	3843	3905	3912	70	70	GAU	GAU	—	—	—	—	—	—
Gln (Q)	-	3905	3912	3975	3982	71	71	UUG	UUG	—	—	—	—	—	—
Met (M)	+	3975	3982	4043	4051	69	70	CAU	CAU	—	—	—	—	—	—
NADH2	+	4044	4052	5089	5097	1046	1046	—	—	ATG	ATG	TA-	TA-	348	348
Trp (W)	+	5090	5098	5161	5168	72	71	UCA	UCA	—	—	—	—	—	—
Ala (A)	-	5163	5170	5231	5238	69	69	UGC	UGC	—	—	—	—	—	—
Asn (N)	-	5233	5440	5305	5312	73	73	GUU	GUU	—	—	—	—	—	—
Cys (C)	-	5344	5351	5408	5415	65	65	GCA	GCA	—	—	—	—	—	—
Tyr (Y)	-	5409	5416	5478	5485	70	70	GUA	GUA	—	—	—	—	—	—
COI	+	5480	5487	7030	7037	1551	1551	—	—	GTG	GTG	TAA	TAA	516	516
Ser (S)	-	7031	7038	7101	7108	71	71	UGA	UGA	—	—	—	—	—	—
Asp (D)	+	7105	7112	7177	7184	73	73	GUC	GUC	—	—	—	—	—	—
COII	+	7188	7193	7878	7883	691	691	—	—	ATG	ATG	T--	T--	230	230
Lys (K)	+	7879	7884	7952	7957	74	74	UUU	UUU	—	—	—	—	—	—
ATP8	+	7956	7959	8123	8126	168	168	—	—	ATG	ATG	TAA	TAA	55	55
ATP6	+	8114	8117	8796	8799	683	683	—	—	ATG	ATG	TA-	TA-	227	227
COIII	+	8797	800	9581	9584	785	785	—	—	ATG	ATG	TA-	TA-	261	261
Gly (G)	+	9582	9585	9653	9655	72	71	UCC	UCC	—	—	—	—	—	—
NADH3	+	9654	9656	10002	10004	349	349	—	—	ATG	ATG	T--	T--	116	116
Arg (R)	+	10003	10005	10071	10073	69	69	UCG	UCG	—	—	—	—	—	—
NADH4L	+	10072	10074	10368	10370	297	297	—	—	ATG	ATG	TAA	TAA	98	98
NADH4	+	10362	10364	11742	11744	1381	1381	—	—	ATG	ATG	T--	T--	460	460
His (H)	+	11743	11745	11812	11813	70	69	GUG	GUG	—	—	—	—	—	—
Ser (S)	+	11813	11814	11879	11880	67	67	GCU	GCU	—	—	—	—	—	—
Leu (L)	+	11887	11887	11959	11959	73	73	UAG	UAG	—	—	—	—	—	—
NCR	+/-	—	11960	—	12179	—	220	—	—	—	—	—	—	—	—
Leu (L)	+	—	12180	—	12252	—	73	—	UAG	—	—	—	—	—	—
NADH5	+	11960	12253	13798	14091	1839	1839	—	—	ATG	ATG	AGA	TAA	612	612
NADH6	-	13794	14088	14315	14609	522	522	—	—	ATG	ATG	TAG	TAA	173	173
Glu (E)	-	14316	14610	14384	14677	69	68	UUC	UUC	—	—	—	—	—	—
CYTb	+	14388	14683	15528	15823	1141	1141	—	—	ATG	ATG	T--	T--	380	380
Thr (T)	+	15529	15824	15600	15895	72	72	UGU	UGU	—	—	—	—	—	—
Pro (P)	-	15601	15896	15670	15965	70	70	UGG	UGG	—	—	—	—	—	—
Control region	+	15671	15696	16602	16896	932	1200	—	—	—	—	—	—	—	—

non-coding region was also observed in the mitogenomes of *P. rubio* (219 bp), *P. tribulus* (219

bp), and *P. evolans* (235 bp). Similarly, a non-coding region of 533 bp and 165 bp were

reported within the cluster NADH4/His/Ser/Leu/NADH5 in the freshwater gobioid

Odontobutis platycephala (GenBank accession NC_010199; Ki et al., 2008) and in the

threadfin *Eleutheronema rhadinum* (Genbank accession MW845829; Zhong et al., 2021)

respectively. Although the unusual non-coding region observed in the four previously

mentioned *Prionotus* species may have been formed as a result of an unknown evolutionary

process due to their close phylogenetic relationship (Fig. 2), we cannot rule out the possibility

of an artefactual origin due to the presence of a C homopolymer at the 5' end of this non-coding

region, which is known to bias short read sequencing (Laehnemann et al., 2016).

The size of the PCGs ranged from 168 bp (55 amino acids) in the ATP8 to 1,839 (612 amino

acids) in the NADH5 of both mitogenomes. All PCGs were encoded in the heavy strand, except

for the NADH6 which was encoded in the light strand. Most PCGs of both mitogenomes used

the start codon ATG, except for the COI gene which was initiated with the alternative GTG

codon. The use of alternative stop codons was observed only in two genes of *P. alatus*

mitogenome where the NADH5 and NADH6 genes ended with the AGA and TAG codons

respectively, and the NADH1 gene in *P. stephanophrys* that used the stop codon TAG. In both

mitogenomes, the heavy strand contained the two rRNA and 14 tRNA genes, including the

additional tRNA-Leu detected in *P. stephanophrys*, while the light strand contained the

remaining 8 tRNA genes.

The Bayesian phylogenetic analysis recovered high to maximum statistical support in all clades

(Fig. 2), grouping all *Prionotus* and *Bellator* species within a major Prionotinae clade. The 8

analyzed *Prionotus* species were recovered within three discrete subclades indicated from

apical towards more basal positions within the phylogenetic tree: subclade I) *P. rubio* + *P.*

tribulus + *P. evolans* + *P. stephanophrys*, subclade II) *P. carolinus* + *P. scitulus*, and subclade

III) *P. alatus + P. nudigula*. The *Bellator* subclade (*B. egretta + B. brachychir*) was positioned

at the basal node of Prionotinae. The well supported sister relationship between *Prionotus* and

Bellator supports the current accepted genus status of *Bellator*, which is also in agreement with

the results of the morphological taxonomic analysis reported by Miller and Richards (1991)

and Imamura (1996). In our phylogenetic results, the subfamily Triglinae, which consisted of

9 species within 3 genera (*Chelidonichthys*, *Eutrigla*, and *Lepidotrigla*), was recovered in the

basal position of the phylogenetic tree. These results support the molecular phylogeny of

Portnoy et al. (2017), where the Triglinae group was regarded as more basal than Prionotinae.

However, our results are in disagreement with those of Richards an Jones (2002), whose

morphological phylogeny analysis recovered *Prionotus* and *Bellator* in adjacent but different

clades, identifying the Prionotinae as the most ancestral subfamily within Triglidae. In contrast

to our findings, the results of Portnoy et al. (2017) rendered *Prionotus* as paraphyletic within

Prionotinae with *Bellator* recovered within the genus *Prionotus*. To address this issue, the

authors suggested two alternative solutions: to retain *Bellator* and split *Prionotus* into different

genera or to lump *Bellator* into the *Prionotus* genus.

Our results recovered *P. alatus* and *P. stephanophrys* within different subclades of Prionotinae,

showing the closest relationships with *P. nudigula* and *P. evolans* respectively. The estimated

levels of divergence (Kimura two-parameter model) between *P. alatus-P. nudigula* and *P.*

stephanophrys-P. evolans were 0.05 and 0.12 respectively. Taking into account that *P.*

stephanophrys was the only Pacific species included in our study and that there are still several

Atlantic species without sequenced mitogenomes, our results cannot determine which Atlantic

species represent the transisthmian sister taxon of *P. stephanophrys*. However, based on the

results reported by Portnoy et al. (2017), *P. longispinosus* represents so far, the closest Atlantic

relative of *P. stephanophrys*, and likely the transisthmian sister species of *P. stephanophrys*.

Fig. 2. Bayesian phylogenetic tree constructed based on the concatenation of 13 mitochondrial

protein-coding genes of 19 Triglidae species (GenBank accessions are shown between

parenthesis). The two species with newly determined mitogenomes assembled in this study

(*Prionotus alatus* and *P. stephanophrys*, subfamily Prionotinae) are highlighted in bold. The

three subclades formed by *Prionotus* species are shaded in sky blue, violet, and blue

respectively. Posterior probabilities are shown at correspondent nodes. The Gilbert's spiny

flathead *Hoplichthys gilberti* was used as outgroup.

The shortest interspecific genetic distance (K2P=0.0014; 23 variable sites) was observed

between the mitogenomes of *P. rubio* (Genbank accession OP056884, Biosample

SAMN29555344)　and　*P. tribulus* (Genbank accession OP056925, BioSample

SAMN29555068), displaying the same length of 16,891 bp. Further analysis of the COI gene

sequence of both mitogenomes in the BOLD database identified both sequences as *P. tribulus*.

Taken all these results together, it is likely that Biosample SAMN29555344, identified as *P.*

rubio, may be a hybrid between a male *P. rubio* and a female *P. tribulus*. Hybridization

between different *Prionotus* species has been previously reported in the wild and under

laboratory conditions (McClure and McEachran, 1992; Herbert et al., 2024), which suggests

that the possible hybridization between the sympatric *P. rubio* and *P. tribulus* is likely.

Conclusion

The phylogenetic analysis performed herein recovered *Prionotus* and *Bellator* species as sister

taxon within Prionotinae, which is in agreement with some previous related studies. The

monophyletic condition of *Prionotus* is in disagreement with the findings reported by Portnoy

et al. (2017), where *Prionotus* was rendered as paraphyletic, suggesting that *Bellator* should be

considered as a junior synonym of *Prionotus*. However, the only two *Bellator* species with

complete mitogenomes included in our analysis (*B. egretta* and *B. brachychir*) were different

from those analyzed by Portnoy et al. (2017). Thereby, the putative paraphyletic status of

Prionotus is still uncertain, highlighting the need for the complete mitogenome sequencing of

more *Prionotus* and *Bellator* species that may allow robust molecular phylogenetic analysis

with broader taxon coverage.

References

Allard CA, Herbert AL, Krueger SP, Liang Q, Walsh BL, Rhyne AL, ... Bellono NW (2024)

Evolution of novel sensory organs in fish with legs. Curr Biol 34(19): 4349-4356.

Chirichigno N, Cornejo RM (2001) Catálogo comentado de los peces marinos del Perú. Lima:

Instituto del Mar del Perú. 2001: 314 pp.

David S, Stuart C, Dominic G, John R, Kevin W (2017) Molecular phylogenetics of New

World searobins (Triglidae; Prionotinae). Mol Phylogenet Evol 107:382-387.

Eldem V, Balcı MA (2024) Mining NCBI Sequence Read Archive Database: An Untapped

Source of Organelle Genomes for Taxonomic and Comparative Genomics

Research. Divers 16(2): 104.

Froese R, Pauly D Editors (2025) FishBase. Prionotus Lacepède, 1801. Accessed through:

World Register of Marine Species at:

https://www.marinespecies.org/aphia.php?p=taxdetails&id=159569 on 2025-01-30

Gouy M, Tannier E, Comte N, Parsons DP (2021) Seaview version 5: a multiplatform software

for multiple sequence alignment, molecular phylogenetic analyses, and tree

reconciliation. Multiple sequence alignment: Methods Protoc 241-260.

Greiner S, Lehwark P, Bock R (2019) OrganellarGenomeDRAW (OGDRAW) version 1.3. 1:

expanded toolkit for the graphical visualization of organellar genomes. Nucleic Acids Res

47(W1): W59-W64.

Herbert AL, Allard CA, McCoy MJ, Wucherpfennig JI, Krueger SP, Chen HI, ...Kingsley DM

(2024) Ancient developmental genes underlie evolutionary novelties in walking fish. Curr

Biol 34(19): 4339-4348.

Imamura H (1996) Phylogeny of the family Platycephalidae and related taxa (Pisces:

Scorpaeniformes). Species Divers 1(2): 123-233.

Inkscape Project (2020) Inkscape. Retrieved from https://inkscape.org. Accessed 30 January

2025

Ki JS, Jung SO, Hwang DS, Lee YM, Lee JS (2008) Unusual mitochondrial genome structure

of the freshwater goby *Odontobutis platycephala*: rearrangement of tRNAs and an

additional non-coding region. J Fish Biol 73(2): 414-428.

Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis

version 7.0 for bigger datasets. Mol Biol Evol 33(7): 1870-1874.

Laehnemann D, Borkhardt A, McHardy AC (2016) Denoising DNA deep sequencing data—

high-throughput sequencing errors and their correction. Brief Bioinform 17(1): 154-179.

Maduna SN, Vivian-Smith A, Jónsdóttir ÓDB, Imsland AK, Klütsch CF, Nyman T, ...Hagen

SB (2022) Mitogenomics of the suborder Cottoidei (Teleostei: Perciformes): improved

assemblies,   mitogenome   features,   phylogeny,   and   ecological

implications. Genomics 114(2): 110297.

Marín A (2025) The first complete mitochondrial genome of the endangered spotted eagle ray

Aetobatus ocellatus. bioRxiv 2025.01.13.632509.

McClure MR, McEachran JD (1992) Hybridization between *Prionotus alatus* and *P. paralatus*

in the Northern Gulf of Mexico (Pisces: Triglidae). Copeia 1992(4): 1039-1046.

Miya M, Nishida M (2015) The mitogenomic contributions to molecular phylogenetics and

evolution of fishes: a 15-year retrospect. Ichthyol Res 62: 29-71.

Richards WJ, Jones DL (2002) Preliminary classification of the gurnards (Triglidae:

Scorpaeniformes). Mar Freshw Res 53(2): 274-282.

Robertson DR, Van Tassell J (2015) Shorefishes of the Greater Caribbean: online information

system. Version 3.0 Smithsonian Tropical Research Institute, Balboa, Panamá.

Ronquist F, Teslenko M, Van Der Mark P, Ayres DL, Darling A, Höhna S, ...Huelsenbeck JP

(2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a

large model space. Syst Biol 61(3): 539-542.

Russell M, Grace M, Gutherz EJ (1992) Field Guide to the searobins (*Prionotus* and *Bellator*)

in the western North Atlantic. NOAA Technical Report NMFS 107. U.S. Department of

Commerce. March 1992.

Smith WL, Everman E, Richardson C (2018) Phylogeny and taxonomy of flatheads,

scorpionfishes, sea robins, and stonefishes (Percomorpha: Scorpaeniformes) and the

evolution of the lachrymal saber. Copeia 106(1): 94-119.

Winn JC, Maduna SN, Bester-van der Merwe AE (2024) A comprehensive phylogenomic

study unveils evolutionary patterns and challenges in the mitochondrial genomes of

Carcharhiniformes: A focus on Triakidae. Genomics 116(1): 110771.

Xia X (2018) DAMBE7: New and improved tools for data analysis in molecular biology and

evolution. Mol Biol Evol 35(6): 1550-1552.

Zhong L, Wang M, Li D, Tang S, Chen X (2021) Mitochondrial genome of *Eleutheronema*

rhadinum with an additional non-coding region and novel insights into the

phylogenetics. Front Mar Sci 8: 746598.