1   ROR2 drives right ventricular heart failure via disruption of proteostasis
2

Hali Hartman B.S.[1], Genevieve Uy B.S.[1], Keita Uchida Ph.D.[2], Emily A. Scarborough Ph.D.[2],
Yifan Yang M.S.[3], Eric Barr B.S.[1,3], Spencer Williams B.S.[1], Ryan T. Santilli B.S.[1], Sanam L.
Kavari B.S.i[3], Jeff Brandimarto M.S.[3], Li Li B.S.[2], Ling Lai M.D.[3], Joanna Griffin[3], Nora Yucel
Ph.D.[3], Swapnil Shewale Ph.D.[3], Hari Rajagopal[1], Deborah M. Eaton Ph.D.[3], Tanis Dorwart
B.S.[4], Kenneth C. Bedi Jr B.S.[3], Crystal S. Conn Ph.D.[4], Kenneth B. Margulies M.D.[3], Benjamin
Prosser Ph.D.[2], *Zoltan Arany M.D., Ph.D.[2,3], *Jonathan J. Edwards M.D.[1,3]
9
[1]Cardiovascular Institute, Children's Hospital of Philadelphia, Philadelphia, PA, USA

[2]Department of Physiology, Pennsylvania Muscle Institute, Perelman School of Medicine,

University of Pennsylvania, Philadelphia, PA, USA

[3]Cardiovascular Institute, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA, USA

[4]Department of Radiation Oncology, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA, USA

17
18   *Contributed equally
19
20   Short Title: ROR2 drives right ventricular heart failure
21
Correspondence to:
Jonathan Edwards, MD
University of Pennsylvania
Cardiovascular Institute
3401 Civic Center Blvd
Smilow Center for Translational Research, Room 11-114
Philadelphia, PA 19104
EdwardsJ6@chop.edu

30
Zoltan Arany, MD, PhD
University of Pennsylvania
Cardiovascular Institute
3401 Civic Center Blvd
Smilow Center for Translational Research, Room 11-106
Philadelphia, PA 19104
zarany@pennmedicine.upenn.edu

Word Count: 4993 (limit 5000)

Abstract

Background: No therapies exist to reverse right ventricular failure (RVF), and the molecular

mechanisms that drive RVF remain under studied. We recently reported that the

developmentally restricted noncanonical WNT receptor ROR2 is upregulated in human RVF in

proportion to severity of disease. Here we test the mechanistic role of ROR2 in RVF

pathogenesis.

Methods: ROR2 was overexpressed or knocked down in neonatal rat ventricular myocytes

(NRVMs) and then characterized using confocal microscopy, RNAseq, proteomics, proteostatic

functional assays, and pacing to assess contractile properties. The impact of cardiac ROR2

expression was evaluated in mice by AAV9-mediated overexpression and by AAV9-mediated

delivery of shRNA to knockdown ROR2 in a pulmonary artery banded pressure overload model

of RVF. ROR2-modified mice were evaluated by echocardiography, histology, and RV protein

synthesis and proteasome capacity.

Results: In NRVMs, we find that ROR2 profoundly dysregulates the coordination between

protein translation and folding. This imbalance leads to excess protein clearance by the ubiquitin

proteasome system (UPS) with dramatic impacts on sarcomere and cytoskeletal structure and

function. Inhibiting the UPS or restoring chaperone expression is sufficient to partially rescue

ROR2-induced structural and contractile deficits in cardiomyocytes. In mice, forced cardiac

ROR2 expression is sufficient to disrupt proteostasis and drive RVF, while conversely ROR2

knockdown partially rescues proteostasis and RV structure and function in a pressure overload

model of RVF.

Conclusions: In sum, ROR2 is a key driver of RVF pathogenesis through proteostatic

disruption and, thus, provides a promising target to treat RVF.

Introduction

Right ventricular failure (RVF) is a widespread problem for which there are no proven therapies. Nearly half of patients with left ventricular failure (LVF) develop RVF, and RVF doubles their risk of death.[1,2] RVF is also the leading cause of hospital admission and death in pulmonary hypertension.[3-5] For LVF, there are several therapies that can reduce patient morbidity and mortality.[6] In contrast, no therapies have proven efficacious for patients with RVF, and in fact, some medications that benefit LVF led to worse outcomes.[7-13] RVF thus fundamentally differs from LVF, and finding new therapies for RVF will require a deeper understanding of targetable molecular drivers of RVF.[14,15]

We recently analyzed a large cohort of human hearts from patients transplanted for dilated or ischemic cardiomyopathy, and discovered that the noncanonical WNT receptor ROR2—usually developmentally restricted—is robustly upregulated in failing RVs, and its expression correlated with RVF severity.[16] ROR2 is essential for the development of multiple organ systems, including the heart.[17] *Ror2* knockout caused ventricular septal defects in mice, while loss of *Ror1* did not impact heart development.[17-19] However, combined *Ror1/Ror2* double knockout caused more severe developmental pathologies including complex conotruncal defects and ventricular noncompaction. The mechanistic role of ROR2 in RVF, however, has not been explored.

We investigated whether ROR2 induction contributes to RVF and identified a previously unrecognized role for ROR2 in regulating cardiomyocyte proteostasis. ROR2 induction disrupts the balance between protein synthesis and folding capacity and increases ubiquitin proteasome system (UPS)-mediated protein degradation. This UPS response is either inadequate or excessive to the point being maladaptive, further contributing to RVF pathogenesis. In a mouse

85   model of RVF, preventing ROR2 induction ameliorated both RV proteostatic disruption and

86   RVF. These findings indicate that targeting ROR2-driven proteostatic disruption may offer a new

87   therapeutic approach for RVF.

Methods

Detailed methods are provided in the Supplemental Material. Source data supporting the

findings and displayed in main figures and supplemental figures are also included in the

Supplemental Material. RNAsequencing data generated for this study was deposited in the

Gene Expression Omnibus under the accession number GSE278627. The mass spectrometry

proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE

partner repository with the dataset identifier PXD05655.[20]

Animal Studies

All animal work was performed under University of Pennsylvania Institutional Animal Care and

Use Committee approved protocols (#805255, #805309, and #807751). Neonatal rat ventricular

myocytes (NRVMs) were isolated from 1-2 day old Sprague Dawley rats after obtaining

pregnant rats from Charles River Laboratory and maintained in culture as previously

described.[21] C57BL/6J mice obtained from Jackson Laboratory were used for all mouse

experiments studies.

Human Studies

Procurement of all human RV myocardial tissue was performed in accordance with protocols

approved by the University of Pennsylvania Institutional Review Board (approval 802781) and

the Gift-of-Life Donor Program, Philadelphia, PA. Prospective informed consent for research

use of surgically explanted tissue was obtained from the transplant recipients or from the next-

of-kin in the case of deceased organ donors, as previously described.[22]

Statistical Analyses

RNAseq and proteomics data were analyzed in RStudio using the voom function from the

Limma package (version 3.56.1) to transform and normalize the data, and the toptable function

to output the differential analysis result. Two group and three group statistical comparisons were

performed in GraphPad Prism v10 using T-test/Mann Whitney U or One-Way ANOVA/Kruskal-

Wallis/Two-Way ANOVA based on normality and number of independent variables.

<u>Results</u>

<u>Cardiomyocyte ROR2 expression is developmentally restricted and reactivated in RV failure</u>

ROR2 is necessary for heart development and its expression is broadly absent in healthy adult

tissues, but the temporal nature for its loss and its histologic pattern in the early postnatal heart

has not been well characterized.[23] By qPCR and western blot, we observe maturity dependent

loss of ROR2 with about 70% reduction in ventricular ROR2 from P2 to P7 and near loss of

ROR2 by P21 (**Fig. 1a, Supplementary Fig. 1a/b**). Myocardial ROR2 at birth is observed

predominantly in cardiomyocytes with localization to the cell membrane and nucleus (**Fig 1b**

and **Supplementary Fig. 1c**). With postnatal maturation, ROR2 is minimally retained and

mostly observed in the atria and subepicardium (**Fig 1a** and **Supplementary Fig. 1c/d**),

consistent with prior single nuclear RNAseq datasets.[24,25] Nuclear localizing ROR2 has been

observed in some cancers, which has been linked to worse outcomes. Therefore, we further

evaluated ROR2 localization using a protein fractionation method that separates nuclear

depleted cytoplasmic proteins ("NCD") from a nuclear enriched fraction that also contains

cytoskeleton proteins ("NCE") (**Supplementary Fig. 1e/f**).[26-28] We found most ROR2 localized

to the NCD fraction, but the temporal loss of ROR2 is comparable across total, NCD, and NCE

fractions.

We previously found that ROR2 is re-expressed in human RVF in a severity dependent manner,

localizing to both cardiomyocytes and fibroblasts.[16] This pattern differs from a prior study in

mouse pressure overload LVF where ROR2 induction was restricted to noncardiomyocytes, so

we next tested if and where ROR2 is reactivated in the failing mouse RV.[29] To that end, we

used a pulmonary artery band (PAB) pressure overload model to generate RVF. With an

average pressure gradient of ~60-70mmHg, PAB mice developed clear markers of pathologic

RV remodeling and RVF with increased RV mass, and echocardiographic RV hypertrophy,

dilation, systolic dysfunction, and tricuspid insufficiency (**Fig. 1c-f** and **Supplementary Table 1**).

PAB also caused RV wall thickening and diffuse RV fibrosis by histology (**Fig. 1g**). PAB RVs

also demonstrate increased expression of the classic heart failure markers fetal myosin heavy

chain protein MYH7 and the natriuretic peptide transcript *Nppb* (**Fig. 1h/i**). In parallel, ROR2

was induced ~5-fold in the NCD fraction and more than 30-fold in the NCE fraction. (**Fig. 1j/k**).

Thus, the normally developmentally restricted ROR2 is robustly reexpressed in mouse RVF (**Fig**

1l). Interestingly, NCE ROR2 strongly correlated to RV functional and structural changes in PAB

including the severity of RV dilation (r=0.66), TAPSE (r= -0.69), and fractional area change (r= -

0.60) (all p<0.05, **Supplementary Table 1**). In contrast, NCD ROR2 showed no significant

correlation to any of these RV functional or structural parameters. Induced ROR2 localized to

both cardiomyocytes and fibroblasts in the PAB RV by immunohistochemistry and RNAscope.

ROR2 protein in cardiomyocytes localized to the cytoplasm, membrane, and nucleus (**Fig. 1m**)

similar to ~P2-P7 mouse RV and consistent with the NCD/NCE fractionation. This was also

consistent with our previous findings in human RVF immunohistochemistry and further

supported by complementary immunohistochemistry and RNAscope of human RVs in this study

(**Supplementary Fig. 2a/b).** We conclude that ROR2 expression, particularly the NCE fraction,

is strongly induced in mouse RVF and increases in proportion to RVF severity.

<u>ROR2 regulates cardiomyocyte morphology</u>

Since little is known about the role of ROR2 in cardiomyocytes, we directly tested the impact of

ROR2 induction and loss using NRVMs. We cultured cells on patterned substrates to promote

sarcomere alignment, allowing more robust structural characterization. We overexpressed

human ROR2 (ROR2^{OE}) or knocked down native *Ror2* by shRNA (ROR2^{KD}) using adenovirus

with a GFP reporter (**Fig. 2a/b**). Adeno-GFP alone was used as a control. ROR2^{OE} and ROR2^{KD}

caused dichotomous morphologic changes. ROR2^{OE} NRVMs had fragmented sarcomeres and

were significantly smaller, driven by reduction in the longitudinal major axis, with a rounded

appearance and smaller aspect ratio (major/minor axis) (**Fig. 2c-e**). In contrast, ROR2^{KD}

NRVMs were slightly larger without affecting the aspect ratio. We also observed opposing

effects on cell-cell junctions with ROR2^{KD} increasing and ROR2^{OE} decreasing peripheral

enrichment of the intercalated disc (ICD) localizing proteins α-actinin, β-catenin, and Cx43 (**Fig.**

2f-h and **Supplementary Fig. 3a**). Total Cx43 and α-actinin staining intensity and protein

expression by western blot increased in ROR2^{OE} NRVMs, while β-catenin levels were stable,

indicating that reduced peripheral staining represents redistribution of the proteins rather than

loss (**Fig 2i-l**). Thus, ROR2 regulates gross cardiomyocyte morphology and both abundance

and peripheral localization of ICD/junction proteins.

ROR2 affects cardiomyocyte ubiquitin-proteasome system, protein folding, and synthesis

We used RNAseq of ROR2^{OE} and ROR2^{KD} NRVMs to investigate potential mechanisms for

these NRVM structural phenotypes (**Fig. 3a** and Supplementary Data 4). We observed that β-

catenin responsive transcriptional targets were not significantly impacted, indicating no

significant role for canonical WNT signaling (**Supplementary Table 2** and **Data 5**). Instead,

gene-set enrichment analysis revealed changes in protein turnover gene programs, which

included upregulation ubiquitin ligase activity genes and downregulation for ubiquitin-dependent

protein catabolism genes in ROR2^{OE} (**Fig. 3b**). Interestingly, at the protein level, total ubiquitin

was significantly decreased in ROR2^{OE} but unchanged in ROR2^{KD} (**Fig 3c/d**). We tested

whether the reduction of ubiquitinated proteins in ROR2^{OE} NRVMs was driven by increased

turnover or decreased ubiquitination by blocking clearance of ubiquitinated proteins using the

competitive proteasome inhibitor MG132. As expected, MG132 increased accumulation of total

ubiquitin across all groups, with ROR2^{OE} NRVMs demonstrating significantly greater

accumulation of proteasome directing K48 ubiquitin (**Fig 3e/f** and **Supplementary Fig 3b**).

Interestingly, ROR2^{KD} demonstrated lower basal K48 ubiquitin with intact accumulation of both

total and K48-linked ubiquitin. We next measured maximal proteasome capacity by incubating

cell lysates with Suc-LLVY-AMC—a fluorescent reporter containing a peptide substrate of the

chymotrypsin-like proteasome subunit (PSMB5). ROR2^{OE} increased PSMB5 maximal capacity

at 0 and 3.5µM ATP, and trended towards an increase at 7µM ATP, without affecting total

PSMB5 expression (**Fig 3g** and **Supplementary Fig 3c)**. Whereas, ROR2^{KD} did not impact

PSMB5 expression or maximal capacity, mirroring the intact accumulation of ubiquitin. Thus,

these data indicate that ROR2 expression significantly alters the UPS, with high ROR2

promoting K48 ubiquitination and proteasome-mediated turnover while low ROR2 decreases

K48 ubiquitination without affecting proteasome capacity, indicating additional levels of

proteostatic regulation.

In addition to UPS related transcriptional changes, the two HSP70 paralogs *Hspa1a* and

Hspa1b exhibited the highest magnitude reduction in ROR2^{OE} (log$_2$FC <-4.7, adj p =0.059,

both). Multiple other chaperones were also trending as significantly downregulated (adj p <0.1)

including *Hsph1/Hspa4* (HSP105/110), *Hsp90aa1/ab1*, *Dnajb1/a1 (*HSP40), and *Cryab* (**Fig 3a)***.

At the protein level, most tested chaperones including HSP70 were significantly increased in

ROR2^{KD}. Whereas, in ROR2^{OE}, HSP70 was significantly decreased, and other tested

chaperones were unchanged (**Fig. 3c/h** and **Supplementary Fig 3d**). To directly measure

protein folding capacity, we used a luciferase refolding assay in which heat-shocked

recombinant luciferase is allowed to refold in cell lysates supplemented with protease inhibitor

and then interrogated by its ability to cleave luciferin thereby generating light (**Fig. 3i**).

Consistent with reduced HSP70 expression, refolding of heat-shocked recombinant luciferase

was significantly impaired in ROR2^{OE} lysates. Despite globally increased chaperone expression,

luciferase refolding was unaffected in ROR2^{KD}. This result may be explained by GFP control

lysates exceeding a minimum HSP70 threshold to sufficiently refold heat shocked luciferase or

by ROR2^{KD} also increasing HSP105/110, which can lead to premature release of protein

substrates that diminishes refolding efficiency (**Supplementary Fig 3c**).[30]

Given the ROR2-responsive effects on protein folding and UPS, we next tested whether other

aspects of protein production or stability may also be disrupted. We quantified global protein

synthesis rates by using the tyrosyl tRNA analogue puromycin and radiolabeled ^{35}S-Met to label

newly synthesized proteins. By puromycin labeling, ROR2^{OE} resulted in ~2-3-fold increase in

protein synthesis and ROR2^{KD} modestly decreased translation (**Fig. 3j/k**). ^{35}S-Met labeling

revealed similar results with ~1.5-fold increase in ROR2^{OE} and ~0.6-fold reduction in ROR2^{KD}

(**Supplementary Fig. 3e**). ROR2-driven effects on protein synthesis were unlikely a secondary

effect of altered chaperone expression, as HSP70 overexpression (HSP70^{OE}) did not reverse

the differential translation rates or ubiquitin levels (**Supplementary Fig. 3f/g**). However, given

the observed differences in protein synthesis, folding, and proteasome capacity with ROR2

modulation, we considered whether the stability of newly synthesized proteins differed across

groups. To that end, we performed a pulse chase experiment consisting of a 1 hour of

puromycin labeling followed by a 1hr incubation with cycloheximide to block further protein

synthesis with or without MG132 or HSP70^{OE}. We found that turnover of newly synthesized

proteins was ~2-fold higher in ROR2^{OE} (~50% vs. 25%) compared to controls, and ROR2^{KD}

demonstrated decreased turnover (<10%) indicating increased protein stability. Strikingly,

MG132 and HSP70^{OE} equivalently increased the stability of newly synthesized proteins ROR2^{OE}

and control NRVMs, but these only reached statistical significance for ROR2^{OE}. We also tested

protein stability using the constitutive UPS substrate GFPµ, which required generating ROR2

modified NRVMs lacking a GFP reporter (**Supplementary Fig. 3h-k**). By microscopy, flow

cytometry, and western blot, GFPµ was consistently lowest in ROR2^{OE} and highest in ROR2^{KD},

even in the presence of MG132. Using bafilomycin, we excluded the possibility that autophagy

contributed to GFPµ clearance independently or in combination with proteasome inhibition. Of

note, although GFPµ remained lowest in ROR2^{OE}, the relative increase of GFPµ by MG132 was

most pronounced in ROR2^{OE}. Comparing these results with the puromycin pulse chase

experiments highlights the importance of considering total protein synthesis rates and baseline

GFPµ levels when interpreting GFPµ flux. These results also suggest that ROR2 increases

protein synthesis in a biologically selective manner. Taken together, these data show that ROR2

profoundly disrupts cardiomyocyte proteostasis by promoting protein synthesis while

simultaneously suppressing folding capacity, priming NRVMs to clear proteins through the UPS.

<u>ROR2 impairs cardiomyocyte structure and function via Proteostatic Disruption</u>

Next, to determine if changes in the proteostatic balance were responsible for the NRVM

morphologic changes observed with ROR2^{OE}, we repeated the morphology evaluations with a

24-hour incubation of 1 µM MG132. Proteasome inhibition significantly abrogated the impact of

ROR2^{OE} on NRVM morphology with improvement in the sarcomere pattern, aspect ratio, and

peripheral localization of α-actinin localization (**Fig. 4a-c** and **Supplementary Fig. 3m**). We also

tested the impact of increased folding capacity by HSP70^{OE}, which similarly improved ROR2^{OE}

NRVM morphology by increasing cell area and aspect ratio, but not to the extent of proteasome

inhibition (**Fig. 4d-f**).

We next tested the impact of ROR2 modulation on cardiomyocyte contractile properties. Using

an IonOptix Cytocypher pacing apparatus, we found that ROR2^{OE} significantly slowed

contractile and relaxation kinetics, and that ROR2^{KD} accelerated relaxation (**Fig. 4g-k**).

Strikingly, HSP70^{OE} accelerated contraction and relaxation kinetics in both ROR2^{OE} and control

NRVMs to match ROR2^{KD} (**Fig. 4g-k**). Taken together, these data demonstrate that disruptions

in proteostatic balance—including UPS-mediated turnover and folding capacity—are primary

mechanistic mediators of ROR2-responsive effects on NRVM morphology and function.

ROR2 regulates the cardiomyocyte proteome

In light of the profound effects of ROR2 on proteostatic balance and evidence that ROR2

impacts protein synthesis and turnover in a selective manner, we next sought to characterize

the interaction between ROR2 expression and proteasome inhibition on the global proteome

(**Fig. 5a-c**). By principal component analysis plot, we observed tight clustering by groups with

PC1 correlating to MG132 and PC2 correlating to ROR2^{OE} effects (**Fig 5c**). Accordingly,

ROR2^{OE} dramatically altered the NRVM proteome, which was broadly reversed by proteasome

inhibition (**Fig 5a/b**). We considered whether intrinsic protein characteristics contributed to

ROR2^{OE}-mediated effects on the global NRVM proteome. Using a comprehensive dataset of

protein half-lives in mouse fibroblasts and known molecular masses as indicators of stability and

complexity, we found that upregulated proteins (adj P<0.05) were smaller and exhibited longer

predicted half-lives compared to downregulated proteins (65.8 kDa vs. 87.5 kDa and 134 vs. 55

hours, p<0.0001, both, **Supplementary Fig a/b**).[31] However, the magnitude of up- or down-

regulation did not correlate directly to protein mass or predicted half-lives, suggesting additional

levels of regulation **(Supplementary Fig 4c-f)**. ROR2^{OE} did not significantly affect the

expression of proteasome subunits, consistent with similar PSMB5 levels by western blot as

noted above. In contrast, we found ROR2^{OE} significantly altered E3 ubiquitin ligase expression

(20 upregulated and 26 downregulated), some of which was reversed by proteasome inhibition,

indicating substantial protein substrate-specificity for ROR2-driven UPS-mediated clearance

(**Fig**. **5d/e**, Supplementary Data 20 and 21). These included upregulation of OZZ-E3 (*Neurl2*)—

a ubiquitin ligase that targets β-catenin specifically at the intercalated disc—and ITCH—which

targets the non-canonical WNT signaling protein DVL (**Fig 5d**). PDZRN3, an E3 ligase that was

recently identified as a likely indirect ROR2 phosphorylation target and which plays a role in

postnatal sarcomerogenesis and ICD formation, was also significantly reduced.[32,33] In contrast,

we observed no changes in the components of the E3 ligase complex SCF$^{β-TRCP}$ that targets the

transcriptional pool of β-catenin, which is consistent with no substantial changes in canonical

WNT signaling by RNAseq (**Supplementary Fig. 4a**).

Pathway enrichment analysis revealed that proteins which were upregulated by ROR2[OE] were

enriched for multiple biological processes including triglyceride biosynthesis, extracellular matrix

proteins, integrin-mediated signaling, heart contraction, and response to insulin. Notable

upregulated proteins included myogenic differentiation factors VGLL3 and MEF2A—consistent

with an immature state—and regulators of calcium homeostasis PLN, CAMK2D, and

CAMK2A—which may contribute to altered contractile kinetics. Moreover, in the setting of

junctional proteins being mislocalized in ROR2[OE] by microscopy, the increase in integrin

associated proteins may reflect a compensatory attempt to establish cell-cell junctions.

Interestingly, of 426 proteins with at least logFC >0.5 upregulation (adj p<0.05), 60% were

significantly decreased by proteasome inhibition compared to 2% that were further increased.

Downregulated proteins were enriched for regulators of proline hydroxylation, sarcomere and

actin filament organization, supramolecular organization, and inhibitors of apoptosis (**Fig. 5f**).

Mirroring the behavior of upregulated proteins, 60% of the 559 downregulated proteins (logFC <

-0.5, adj p<0.05) were significantly increased by MG132 with 6% demonstrating further

reduction. Importantly, reversal of downregulated proteins included many actin cytoskeleton and

sarcomere associated proteins including: TPM1/2/3, XIRP1/2, MYOZ2, LMOD2, FLNC, NRAP,

and TNNT2 (**Fig. 5g** and **Supplementary Fig. h/i**). Thus, we conclude that ROR2-mediated

disruption of the cardiomyocyte proteome impinges on cardiomyocyte differentiation, structural,

functional, and adhesion proteins, that are largely dependent on altered UPS activity.

<u>Cardiac ROR2 expression causes RV failure and proteostatic disruption</u>

Given the marked effects of ROR2^{OE} in cardiomyocytes in cell culture, we next sought to test

the effect of ROR2 expression in the RV *in vivo*. To this end, we generated cardiac ROR2^{OE}

mice using AAV9 to deliver HA-tagged ROR2 versus GFP in 4-week-old male and female mice

(**Fig. 5a**). At 8 weeks after injection, we observed robust RV and LV ROR2 expression, which

localized to the cardiomyocyte membrane and cytoplasm (**Fig. 6b, Supplementary Figs. 1a**

and 5a). ROR2^{OE} led to significant RV hypertrophy in male mice, marked by an increase in RV

mass, RV wall area, and RV cardiomyocyte cross-sectional area by minimum Feret Diameter

(**Fig. 6b-h**). Echocardiography (n=8/group, 50% male) revealed biventricular dilation and

systolic dysfunction in male ROR2^{OE} mice (**Fig. 6i-l** and **Supplementary Table 3**). Cardiac

catheterization revealed a male-specific increase in RV end diastolic pressure (**Fig. 6m**).

ROR2^{OE} male mice also exhibited induction of *Nppb* mRNA and fetal MYH7 **(Fig. 6n/o)** without

causing an increase in fibrosis (**Fig. 6p**). The cardiac dysfunction seen in male ROR2^{OE} mice

was accompanied by proteostatic disruptions similar to those seen in NRVMs including lower

Hspa1b mRNA, decreased cardiomyocyte ubiquitin staining, increased protein synthesis, and

increased proteasome capacity **(Fig. 6q-u** and **Supplementary Table 4**). We also observed

reduced peripheral β-catenin in RV cardiomyocytes consistent with disrupted cell-cell junctions

that mirror effects seen in NRVMs (**Supplementary Fig. 5b/c**). We conclude that ROR2

expression is sufficient to drive RV failure *in vivo* in male mice, concomitant with disruption of

proteostasis, whereas female ROR2^{OE} mice were generally much less affected.

<u>Suppression of ROR2 induction in RV pressure overload abrogates RV failure and proteostatic</u>

<u>disruption</u>

The results above demonstrated that ROR2 expression was sufficient to cause RV dysfunction

and disrupt proteostasis. We next sought to test if ROR2 expression was necessary for RVF in

the PAB mouse model and its role in proteostasis. We first evaluated the proteostatic signature

of PAB-induced RVF, noting that PAB increased RV protein synthesis, HSP70, and total

ubiquitin—with K48 ubiquitin and PSMB5 increased in the NCE fraction—and proteasome

capacity (**Fig. 7a-g** and **Supplementary Table 5**). Thus, the PAB RVF mouse model

demonstrates clear proteostatic disruption with an increase in both protein synthesis and UPS-

mediated turnover, to some extent analogous to that found with ROR2^{OE}. Next, to determine if

ROR2 induction in PAB contributes to RVF and proteostatic disruption, we injected male mice

with AAV9 to deliver *shRor2* (PAB^{shRor2}, n = 15) or scrambled shRNA (PAB^{shScr}, n=15) prior to

PAB. Since ROR2 localization in the NCE fraction correlated to PAB severity, we assessed the

relative ROR2 knockdown in NCD and NCE fractions and found PAB^{shRor2} had ~50% lower NCE

and ~25% lower NCD ROR2 in the RV (**Fig. 7i/j**). Strikingly, PAB^{shRor2} mice demonstrated less

severe RV dilation, decreased incidence of tricuspid insufficiency, better RV systolic function

measured by TAPSE (**Fig. 7j-l**), improved diastolic function by e' (**Fig. 7m**), less RV fibrosis,

and reduction in RV wall area by histology (**Fig. 7k-p** and **Supplementary Table 6**). ROR2

knockdown also ameliorated the proteostatic dysfunction in PAB with decreases in protein

synthesis, HSP70, and K48-linked ubiquitin without affecting proteasome capacity (**Fig. 7q-t**),

partially reversing the PAB-induced defects. Taken together, these data demonstrate that ROR2

knockdown protects from pathologic RV remodeling and improves RV function in response to

pressure overload, concomitant with partial reversion of altered proteostasis.

<u>ROR2 expression in human RVF correlates to proteasome capacity</u>

Finally, we sought to determine in human RVF the relationship between ROR2 and

proteostasis. To do so, we leveraged a collection of RV tissue obtained from non-failing organ

donors compared to DCM RVs collected at transplant. DCM RVs were stratified by having

preserved RV function versus RVF, and RVF hearts were further stratified by the level of NCD

and NCE ROR2. We evaluated for demographic, clinical, and anthropometric balance across all

relevant comparisons (**Fig 8a** and **Supplementary Tables 7-10**). We observed that high ROR2

expressing RVF, stratified by NCE or NCD fractions, demonstrated significantly increased

proteasome capacity at 0, 3.5, and 7 µM ATP (**Fig 8, Supplementary Fig 6a**, and

Supplementary Table 11; p<0.05). Interestingly, NCE ROR2 showed significant Pearson

correlation to proteasome capacity at each of the tested ATP levels (r=0.68 – 0.82, p≤ 0.0008;

Fig 8c and **Supplementary Tables 12**), whereas NCD ROR2 showed no clear relationship. We

also assessed the abundance of other proteostatic markers and found that high NCE ROR2

associated with a trend for decreased NCE HSP70 and that high NCD ROR2 associated with

increased NCD K48 ubiquitin (**Supplemental Fig 6b-e**). However, we found no clear patterns

for PSMB5 and total ubiquitin (**Supplemental Figure 6f-i**). We conclude that in human

cardiomyopathy, as in mice, high ROR2 expression—particularly NCE ROR2—is associated

with an increase in proteasome capacity.

Discussion

RVF remains a widespread, challenging problem for which there are no proven therapies.[14] The

breadth of LVF medications have not shown clear benefit for RVF, indicating the presence of

distinct RVF mechanisms.[7-13,34] In severe human RVF, we found that the developmentally

restricted noncanonical WNT receptor ROR2 is pathologically reactivated by both

cardiomyocytes and fibroblasts.[16] We now find this expression pattern is recapitulated in mouse

pressure overload RVF, which contrasts with a prior report in mouse LVF for which induced

ROR2 is restricted to noncardiomyocytes.[29] Using NRVMs, in vivo murine models, and human

heart biospecimens, we now reveal a previously unrecognized role for ROR2 regulating

cardiomyocyte proteostasis with implications on contractile function and structure. Interestingly,

in mouse and human RVF, NCE localizing ROR2 demonstrates a stronger association than

NCD ROR2 with pathologic RV remodeling and proteostatic disruption. Our data point to a

model wherein ROR2 promotes protein synthesis without sufficient folding capacity (**Fig. 8d**).

Although UPS-mediated clearance is upregulated in this setting, this is either not sufficient to

meet the demands or, potentially, chronically maladaptive and ultimately converges on

cardiomyocyte functional compromise. Thus, we propose that addressing ROR2-mediated

proteostatic disruption represents a potential novel therapeutic avenue for RVF, for which no

proven therapies currently exist.

ROR2 is a noncanonical cell surface WNT receptor pseudokinase that signals through at least

three broad pathways: RAC/RHO regulation of cell polarity, activation of a JNK/cJUN

phosphorylation cascade, and activation of Ca^{2+}-responsive targets (e.g. PKC, CAMKII).[35] The

molecular and functional impact of pathogenic ROR2 has been best studied in cancer, for which

ROR2 causes context-specific pleiotropic phenotypes with mixed impact on cancer cell death,

invasion, and metastasis resulting in both positive and negative associations with patient

mortality.[36,37] Little is known about the role of ROR2 in cardiac pathology, but upregulation of

ROR2 leading to proteostatic disruption in RVF fits within the well-recognized paradigm of fetal

gene program reactivation during heart failure.[38,39] As the first functional organ, proliferating fetal

cardiomyocytes must maintain a high quality pool of contractile proteins, including some of the

largest in the proteome (e.g. Titin, Obscurin, Dystrophin, etc.), which is energetically demanding

and requires high protein synthesis matched to a robust folding capacity.[40-42] During human

heart development, *ROR2* and *HSPA1A/B* (HSP70) mRNA levels are positively correlated and

taper during later prenatal stages, consistent with amplified but balanced proteostasis of the

immature heart.[43] Postnatally, in mice we found age-dependent reductions in ventricular HSP70

and ROR2, with reductions in ROR2 preceding HSP70. In mouse PAB RV, translation, folding,

and proteasome capacity are all increased, but the presence of ubiquitin accumulation suggests

there is imbalance in these three components that favors excess translation.[42] Moreover, given

that ROR2 knockdown in PAB improves RV remodeling and function while most substantially

affecting protein synthesis, attenuating the translational hyperactivation of RVF may be the

critical factor to restore proteostatic balance. Prior work has shown that slowing translation with

rapamycin can improve translational fidelity in mTOR activated embryonic fibroblasts and can

rescue multiple preclinical LVF models.[44-46] In kind, rapamycin was shown to abrogate PAB-

induced RV hypertrophy and contractile dysfunction in rats.[44] Although protein translation and

UPS capacity were not measured, rapamycin did further increase PAB-induced ubiquitin

accumulation suggesting a decrease in UPS-mediated clearance, which may be the result of

slowing translation leading to improved overall proteome quality.

Proteostatic disruption is strongly implicated in heart failure, and its role in RVF is increasingly

recognized, yet developing targeted therapies has remained elusive.[47-49] Restoring proteostatic

balance would require precise control of synthesis, folding, and turnover, or a more practical

strategy of eliminating a pathogenic upstream regulator. Before this study, few links connected

ROR2—or noncanonical WNT signaling more broadly—to proteostasis. In cultured ovarian

carcinoma cells that typically express low ROR2, adenoviral ROR2 delivery activated the

unfolded protein response and triggered apoptosis.[50] These ROR2 effects in ovarian carcinoma

were rescued by blocking induction of the endoplasmic reticulum stress protein IRE1a. In

NRVMs, treatment with WNT5A—the only known ligand for ROR2—was shown to promote

protein synthesis and drive cardiomyocyte hypertrophy.[51,52] These WNT5A effects could be

blocked by preventing the membrane localization of the ROR2 binding partner VANGL2 or by

inhibiting JNK. Although JNK/cJUN are not classic regulators of protein synthesis, extensive

evidence shows crosstalk with mTOR and ERK. For example, both canonical and noncanonical

WNT activation can inhibit the signaling hub protein GSK3 through phosphorylation.[53,54]

Inhibiting GSK3 prevents it from phosphorylating several key substrates including β-catenin—

(leading to canonical WNT activation), TSC2 (activating mTORC1), and Raptor (suppressing

mTORC1).[55-57] JNK can also activate mTORC1 through Raptor phosphorylation, as observed in

HEK293 cells under osmotic stress.[58] ERK signaling provides an additional link. In A375

melanoma cells which have low endogenous ROR2, induction of ROR2 caused ERK

hyperactivation and cell death.[36] In isolated mouse cardiomyocytes, ERK1/2 activation and

HSP70 expression show reciprocal antagonism, and HSP70 knockdown was sufficient to cause

contractile and relaxation deficits similar to those we observed with ROR2^{OE} NRVMs.[59] Thus,

ROR2 could lead to a pathologic increase in protein translation, thereby leading to proteostatic

disruption in cardiomyocytes, by activating mTOR or ERK signaling through modulating activity

of other true kinases (e.g. JNK/cJUN or GSK3)

Using orthogonal methods, we found that ROR2 can localize to the nucleus in the immature

heart and RVF. Nuclear enriched ROR2 in the NCE fraction strongly correlated to pathogenic

RV remodeling and proteostatic disruption. The functional role of nuclear ROR2 remains poorly

characterized, but in cancer nuclear ROR2 correlates to worse disease with increased

metastasis in melanoma and invasiveness in pancreatic cancer.[26,28] These pathophysiologic

associations with nuclear ROR2 underscore the importance of understanding the regulatory

mechanisms and functional roles (e.g. transcriptional modulation, scaffolding, etc.) of nuclear

localizing ROR2, which may further enable selective therapeutic targeting. For example,

ERK1/2 homeostatic and pathogenic roles are compartmentalized into the cytoplasm and

nucleus, respectively, and blocking its nuclear localization was shown to prevent pressure

induced cardiomyocyte hypertrophy and heart failure.[60,61]

Limitations

More work will be necessary to establish the therapeutic potential for targeting ROR2 in RVF.

While we have not observed sexual dimorphism for the impact of ROR2 in humans, we found

that overexpressing ROR2 in female mice led to a milder phenotype compared to males. This

sex difference in mice may fit with prior reports showing female mice are protected from

proteostatic disruption through higher basal chaperone expression and UPS capacity.[62-64] We

focused our in vitro mechanistic studies in cardiomyocytes, but ROR2 upregulation was also

observed in fibroblasts of human and mouse RVF. Importantly, in a prior mouse model of left

ventricular pressure overload where ROR2 induction was restricted to fibroblasts and

leukocytes, inducible total body combined ROR1/ROR2 knockout was lethal, diminished early

myofibroblast activation, and increased the abundance of inflammatory cells in the LV.[29] The

differences between that study and ours may be due to multiple factors including ubiquitous

knockout of both genes compared to knockdown of only ROR2 and by differences in the models

(RV vs LV pressure overload and the implicated cell populations) that may alter the impact of

loss of ROR2. Since ROR2 is known to be pleiotropic, understanding the role of ROR2 across

the spectrum of fibroblast activation states could have significant implications for its therapeutic

targetability.[29, 65] Finally, our analysis on the relationship between ROR2, RVF, and

proteostasis in humans reflects a single, uncontrolled timepoint which prevents us from

measuring dynamic processes (e.g. UPS flux, protein synthesis rates) that would more fully

capture this relationship.

Conclusions

In summary, we find compelling evidence to implicate ROR2 in RVF pathogenesis. Pathologic

reactivation of ROR2 causes proteostatic disruption in the RV, leading to RVF. In humans,

ROR2 is otherwise broadly absent from healthy mature tissues with its only other well

recognized induction occurring in cancer.[23] Thus, we conclude that targeting ROR2 offers a

potential therapeutic strategy to restore proteostatic balance and treat RVF, for which no

specific therapies currently exist.

Acknowledgements

We thank Julian Mintseris, Rachel Rodrigues, and Jonathan Van Vraken at Thermo Fisher
Scientific Center for Multiplexed Proteomics at Harvard Medical School for support in performing
proteomics. We thank the Day Lab at University of Pennsylvania for kindly providing adenovirus
to express GFPµ. The mouse PAB and echocardiography were performed by the Rodent
Cardiovascular Phenotyping Core (RRID: SCR_022419) at the University of Pennsylvania
supported by the Penn Cardiovascular Institute and NIH S10OD016393. Figures were
generated with assistance of BioRender.

Sources of Funding

J.J.E and Z.A. was supported by the Children's Hospital of Philadelphia Frontier Program
"Advanced Cardiac Therapies for Heart Failure". J.J.E. was also supported by National Institute
of Health (NIH) K08HL159311 and Z.A. was also supported by NIH HL152446 and HL152446.
K.U. was supported by the NIH (T32 HL007843). B.P. was supported by the NIH R01 HL149891
and the Foundation Leducq Research Grant no. 20CVD01. C.S.C. was supported by
Department of the Army (HT94252310080) and the NIH R35 GM154896. Human heart tissues
were procured with funding support from multiple NIH grants to K.M.: R01 AG017022, R01
HL089847, and R01 HL105993, and R01 HL133080.

Author Contributions

H.H., J.J.E., and Z.A. developed the concept and designed the experiments. J.J.E, Z.A., and
B.L.P. supervised and funded the project. J.J.E., H.H., K.U., E.A.S., G.U., S.K., and S.W.

conducted NRVM experiments. J.J.E., H.H., D.E., and R.T.S. conducted NRVM

contractility/relaxation experiments. J.J.E and H.H. performed mouse experiments. J.J.E, T.D.,

and C.S.C. conducted luciferase refolding and ^{35}S-Met labeling assays. J.B. and Y.Y. performed

RNAseq analysis. L. Lai, J.G., and S.S. performed mouse surgeries and physiologic

characterization. K.C.B. and K.M. supported human heart tissue procurement and clinical

characterization. H.R. contributed to microscopy image analysis. L.Li performed tissue

histology. All authors contributed to the manuscript preparation and approved the final version of

the manuscript.

Disclosures

There are no relevant disclosures.

Corresponding author

Correspondence to Jonathan J. Edwards and Zoltan Arany.

References

References

1 Lampert, B. C. & Teuteberg, J. J. Right ventricular failure after left ventricular assist

devices. *J Heart Lung Transplant* **34**, 1123-1130 (2015).

https://doi.org/10.1016/j.healun.2015.06.015

2 Zakeri, R. & Mohammed, S. F. Epidemiology of Right Ventricular Dysfunction in Heart

Failure with Preserved Ejection Fraction. *Curr Heart Fail Rep* **12**, 295-301 (2015).

https://doi.org/10.1007/s11897-015-0267-3

3 Khairy, P. *et al.* Long-term survival, modes of death, and predictors of mortality in

patients with Fontan surgery. *Circulation* **117**, 85-92 (2008).

https://doi.org/10.1161/CIRCULATIONAHA.107.738559

4 Piran, S., Veldtman, G., Siu, S., Webb, G. D. & Liu, P. P. Heart failure and ventricular

dysfunction in patients with single or systemic right ventricles. *Circulation* **105**, 1189-

1194 (2002).

5 Campo, A. *et al.* Outcomes of hospitalisation for right heart failure in pulmonary arterial

hypertension. *Eur Respir J* **38**, 359-367 (2011).

https://doi.org/10.1183/09031936.00148310

6 D'Amario, D. *et al.* Eligibility for the 4 Pharmacological Pillars in Heart Failure With

Reduced Ejection Fraction at Discharge. *J Am Heart Assoc* **12**, e029071 (2023).

https://doi.org/10.1161/JAHA.122.029071

7 Babu-Narayan, S. V. *et al.* Randomised trial of ramipril in repaired tetralogy of Fallot and

pulmonary regurgitation: the APPROPRIATE study (Ace inhibitors for Potential

PRevention Of the deleterious effects of Pulmonary Regurgitation In Adults with repaired

TEtralogy of Fallot). *Int J Cardiol* **154**, 299-305 (2012).

https://doi.org/10.1016/j.ijcard.2010.09.057

8   Dos, L. *et al.* Eplerenone in systemic right ventricle: double blind randomized clinical

trial. The evedes study. *Int J Cardiol* **168**, 5167-5173 (2013).

https://doi.org/10.1016/j.ijcard.2013.07.163

9   Doughan, A. R., McConnell, M. E. & Book, W. M. Effect of beta blockers (carvedilol or

metoprolol XL) in patients with transposition of great arteries and dysfunction of the

systemic right ventricle. *Am J Cardiol* **99**, 704-706 (2007).

https://doi.org/10.1016/j.amjcard.2006.10.025

10   Shaddy, R. E. *et al.* Carvedilol for children and adolescents with heart failure: a

randomized controlled trial. *JAMA* **298**, 1171-1179 (2007).

https://doi.org/10.1001/jama.298.10.1171

11   Bokma, J. P. *et al.* Effect of Losartan on Right Ventricular Dysfunction: Results From the

Double-Blind, Randomized REDEFINE Trial (Right Ventricular Dysfunction in Tetralogy

of Fallot: Inhibition of the Renin-Angiotensin-Aldosterone System) in Adults With

Repaired Tetralogy of Fallot. *Circulation* **137**, 1463-1471 (2018).

https://doi.org/10.1161/CIRCULATIONAHA.117.031438

12   Roche, S. L. & Redington, A. N. Right ventricle: wrong targets? Another blow for

pharmacotherapy in congenital heart diseases. *Circulation* **127**, 314-316 (2013).

https://doi.org/10.1161/CIRCULATIONAHA.112.155887

13   Hartupee, J. & Mann, D. L. Neurohormonal activation in heart failure with reduced

ejection fraction. *Nat Rev Cardiol* **14**, 30-38 (2017).

https://doi.org/10.1038/nrcardio.2016.163

14   Houston, B. A., Brittain, E. L. & Tedford, R. J. Right Ventricular Failure. *N Engl J Med*

388, 1111-1125 (2023). https://doi.org/10.1056/NEJMra2207410

15   Jurida, L. *et al.* A common gene signature of the right ventricle in failing rat and human

hearts. *Nat Cardiovasc Res* **3**, 819-840 (2024). https://doi.org/10.1038/s44161-024-

00485-1

16   Edwards, J. J. *et al.* Noncanonical WNT Activation in Human Right Ventricular Heart

Failure. *Front Cardiovasc Med* **7**, 582407 (2020).

https://doi.org/10.3389/fcvm.2020.582407

17   Takeuchi, S. *et al.* Mouse Ror2 receptor tyrosine kinase is required for the heart

development and limb formation. *Genes Cells* **5**, 71-78 (2000).

18   Guo, R. & Xing, Q. S. Roles of Wnt Signaling Pathway and ROR2 Receptor in

Embryonic Development: An Update Review Article. *Epigenet Insights* **15**,

25168657211064232 (2022). https://doi.org/10.1177/25168657211064232

19   Nomi, M. *et al.* Loss of mRor1 enhances the heart and skeletal abnormalities in mRor2-

deficient mice: redundant and pleiotropic functions of mRor1 and mRor2 receptor

tyrosine kinases. *Mol Cell Biol* **21**, 8329-8335 (2001).

https://doi.org/10.1128/MCB.21.24.8329-8335.2001

20   Perez-Riverol, Y. *et al.* The PRIDE database at 20 years: 2025 update. *Nucleic Acids*

Res **53**, D543-D553 (2025). https://doi.org/10.1093/nar/gkae1011

21   Scarborough, E. A. *et al.* Microtubules orchestrate local translation to enable cardiac

growth. *Nat Commun* **12**, 1547 (2021). https://doi.org/10.1038/s41467-021-21685-4

22   Chen, C. Y. *et al.* Suppression of detyrosinated microtubules improves cardiomyocyte

function in human heart failure. *Nat Med* **24**, 1225-1233 (2018).

https://doi.org/10.1038/s41591-018-0046-2

23   Morioka, K. *et al.* Orphan receptor tyrosine kinase ROR2 as a potential therapeutic

target for osteosarcoma. *Cancer Sci* **100**, 1227-1233 (2009).

https://doi.org/10.1111/j.1349-7006.2009.01165.x

24   Chaffin, M. *et al.* Single-nucleus profiling of human dilated and hypertrophic

cardiomyopathy. *Nature* **608**, 174-180 (2022). https://doi.org/10.1038/s41586-022-

04817-8

25   Litvinukova, M. *et al.* Cells of the adult human heart. *Nature* **588**, 466-472 (2020).

https://doi.org/10.1038/s41586-020-2797-4

26   Carbone, C. *et al.* Adipocytes sustain pancreatic cancer progression through a non-

canonical WNT paracrine network inducing ROR2 nuclear shuttling. *Int J Obes (Lond)*

42, 334-343 (2018). https://doi.org/10.1038/ijo.2017.285

27   Lopez-Bergami, P. ROR2, a driver of "phenotype switching" in melanoma? *Cancer Cell*

Int **22**, 288 (2022). https://doi.org/10.1186/s12935-022-02711-x

28   O'Connell, M. P. *et al.* The orphan tyrosine kinase receptor, ROR2, mediates Wnt5A

signaling in metastatic melanoma. *Oncogene* **29**, 34-44 (2010).

https://doi.org/10.1038/onc.2009.305

29   Chavkin, N. W. *et al.* The Cell Surface Receptors Ror1/2 Control Cardiac Myofibroblast

Differentiation. *J Am Heart Assoc* **10**, e019904 (2021).

https://doi.org/10.1161/JAHA.120.019904

30   Mandal, A. K. *et al.* Hsp110 chaperones control client fate determination in the hsp70-

Hsp90 chaperone system. *Mol Biol Cell* **21**, 1439-1448 (2010).

https://doi.org/10.1091/mbc.e09-09-0779

31   Schwanhausser, B. *et al.* Global quantification of mammalian gene expression control.

Nature **473**, 337-342 (2011). https://doi.org/10.1038/nature10098

32   Konopelski Snavely, S. E. *et al.* Proteomic analysis identifies the E3 ubiquitin ligase

Pdzrn3 as a regulatory target of Wnt5a-Ror signaling. *Proc Natl Acad Sci U S A* **118**

(2021). https://doi.org/10.1073/pnas.2104944118

33   Pernot, M. *et al.* Decrease of Pdzrn3 is required for heart maturation and protects

against heart failure. *Sci Rep* **12**, 8 (2022). https://doi.org/10.1038/s41598-021-03795-7

34   Jurida, L. *et al.* A common gene signature of the right ventricle in failing rat and human

hearts. *Nature Cardiovascular Research* **3**, 819-840 (2024).

https://doi.org/10.1038/s44161-024-00485-1

35   Debebe, Z. & Rathmell, W. K. Ror2 as a therapeutic target in cancer. *Pharmacol Ther*

150, 143-148 (2015). https://doi.org/10.1016/j.pharmthera.2015.01.010

36   Castro, M. V. & Lopez-Bergami, P. Cellular and molecular mechanisms implicated in the

dual role of ROR2 in cancer. *Crit Rev Oncol Hematol* **170**, 103595 (2022).

https://doi.org/10.1016/j.critrevonc.2022.103595

37   Ford, C. E., Qian Ma, S. S., Quadir, A. & Ward, R. L. The dual role of the novel Wnt

receptor tyrosine kinase, ROR2, in human carcinogenesis. *Int J Cancer* **133**, 779-787

(2013). https://doi.org/10.1002/ijc.27984

38   van der Pol, A., Hoes, M. F., de Boer, R. A. & van der Meer, P. Cardiac foetal

reprogramming: a tool to exploit novel treatment targets for the failing heart. *J Intern Med*

288, 491-506 (2020). https://doi.org/10.1111/joim.13094

39   Poon, E. *et al.* Proteomic Analysis of Human Pluripotent Stem Cell-Derived, Fetal, and

Adult Ventricular Cardiomyocytes Reveals Pathways Crucial for Cardiac Metabolism and

Maturation. *Circ Cardiovasc Genet* **8**, 427-436 (2015).

https://doi.org/10.1161/CIRCGENETICS.114.000918

40   Kotter, S. & Kruger, M. Protein Quality Control at the Sarcomere: Titin Protection and

Turnover and Implications for Disease Development. *Front Physiol* **13**, 914296 (2022).

https://doi.org/10.3389/fphys.2022.914296

41   Nagai, R., Low, R. B., Stirewalt, W. S., Alpert, N. R. & Litten, R. Z. Efficiency and

capacity of protein synthesis are increased in pressure overload cardiac hypertrophy.

Am J Physiol **255**, H325-328 (1988). https://doi.org/10.1152/ajpheart.1988.255.2.H325

42   Lewis, Y. E. *et al.* Localization of transcripts, translation, and degradation for

spatiotemporal sarcomere maintenance. *J Mol Cell Cardiol* **116**, 16-28 (2018).

https://doi.org/10.1016/j.yjmcc.2018.01.012

43 Pervolaraki, E., Dachtler, J., Anderson, R. A. & Holden, A. V. The developmental

transcriptome of the human heart. *Sci Rep* **8**, 15362 (2018).

https://doi.org/10.1038/s41598-018-33837-6

44 Harston, R. K. *et al.* Rapamycin treatment augments both protein ubiquitination and Akt

activation in pressure-overloaded rat myocardium. *Am J Physiol Heart Circ Physiol* **300**,

H1696-1706 (2011). https://doi.org/10.1152/ajpheart.00545.2010

45 Fleming, N. D., Samsa, L. A., Hassel, D., Qian, L. & Liu, J. Rapamycin attenuates

pathological hypertrophy caused by an absence of trabecular formation. *Sci Rep* **8**, 8584

(2018). https://doi.org/10.1038/s41598-018-26843-1

46 Li, S. *et al.* Mechanism of Eccentric Cardiomyocyte Hypertrophy Secondary to Severe

Mitral Regurgitation. *Circulation* **141**, 1787-1799 (2020).

https://doi.org/10.1161/CIRCULATIONAHA.119.043939

47 Buhringer, H., Doroudgar, S., Wang, X., Frey, N. & Rangrez, A. Y. Editorial: Proteostasis

in cardiac health and disease. *Front Mol Biosci* **11**, 1433721 (2024).

https://doi.org/10.3389/fmolb.2024.1433721

48 Heitmeier, T. *et al.* Altered proteasome function in right ventricular hypertrophy.

Cardiovasc Res **116**, 406-415 (2020). https://doi.org/10.1093/cvr/cvz103

49 Rajagopalan, V. *et al.* Altered ubiquitin-proteasome signaling in right ventricular

hypertrophy and failure. *Am J Physiol Heart Circ Physiol* **305**, H551-562 (2013).

https://doi.org/10.1152/ajpheart.00771.2012

50 Li, R. *et al.* ROR2 induces cell apoptosis via activating IRE1alpha/JNK/CHOP pathway

in high-grade serous ovarian carcinoma in vitro and in vivo. *J Transl Med* **17**, 428 (2019).

https://doi.org/10.1186/s12967-019-02178-x

51 Hagenmueller, M. *et al.* Dapper-1 is essential for Wnt5a induced cardiomyocyte

hypertrophy by regulating the Wnt/PCP pathway. *FEBS Lett* **588**, 2230-2237 (2014).

https://doi.org/10.1016/j.febslet.2014.05.039

52   Zhang, P. *et al.* Vangl2 is essential for myocardial remodeling activated by Wnt/JNK

signaling. *Exp Cell Res* **365**, 33-45 (2018). https://doi.org/10.1016/j.yexcr.2018.02.012

53   Landin Malt, A. *et al.* Non-Canonical Wnt Signaling Regulates Cochlear Outgrowth and

Planar Cell Polarity via Gsk3beta Inhibition. *Front Cell Dev Biol* **9**, 649830 (2021).

https://doi.org/10.3389/fcell.2021.649830

54   Wu, D. & Pan, W. GSK3: a multifaceted kinase in Wnt signaling. *Trends Biochem Sci* **35**,

161-168 (2010). https://doi.org/10.1016/j.tibs.2009.10.002

55   Zhai, P., Sciarretta, S., Galeotti, J., Volpe, M. & Sadoshima, J. Differential roles of GSK-

3beta during myocardial ischemia and ischemia/reperfusion. *Circ Res* **109**, 502-511

(2011). https://doi.org/10.1161/CIRCRESAHA.111.249532

56   Stretton, C. *et al.* GSK3-mediated raptor phosphorylation supports amino-acid-

dependent mTORC1-directed signalling. *Biochem J* **470**, 207-221 (2015).

https://doi.org/10.1042/BJ20150404

57   Robertson, H., Hayes, J. D. & Sutherland, C. A partnership with the proteasome; the

destructive nature of GSK3. *Biochem Pharmacol* **147**, 77-92 (2018).

https://doi.org/10.1016/j.bcp.2017.10.016

58   Kwak, D. *et al.* Osmotic stress regulates mammalian target of rapamycin (mTOR)

complex 1 via c-Jun N-terminal Kinase (JNK)-mediated Raptor protein phosphorylation.

J Biol Chem **287**, 18398-18407 (2012). https://doi.org/10.1074/jbc.M111.326538

59   Kim, Y. K. *et al.* Deletion of the inducible 70-kDa heat shock protein genes in mice

impairs cardiac contractile function and calcium handling associated with hypertrophy.

Circulation **113**, 2589-2597 (2006).

https://doi.org/10.1161/CIRCULATIONAHA.105.598409

60   Lorenz, K., Schmitt, J. P., Schmitteckert, E. M. & Lohse, M. J. A new type of ERK1/2

autophosphorylation causes cardiac hypertrophy. *Nat Med* **15**, 75-83 (2009).

https://doi.org/10.1038/nm.1893

61   Tomasovic, A. *et al.* Interference with ERK-dimerization at the nucleocytosolic interface

targets pathological ERK1/2 signaling without cardiotoxic side-effects. *Nat Commun* **11**,

1733 (2020). https://doi.org/10.1038/s41467-020-15505-4

62   Kararigas, G. *et al.* Transcriptome characterization of estrogen-treated human

myocardium identifies myosin regulatory light chain interacting protein as a sex-specific

element influencing contractile function. *J Am Coll Cardiol* **59**, 410-417 (2012).

https://doi.org/10.1016/j.jacc.2011.09.054

63   Olazabal, U. E., Pfaff, D. W. & Mobbs, C. V. Sex differences in the regulation of heat

shock protein 70 kDa and 90 kDa in the rat ventromedial hypothalamus by estrogen.

Brain Res **596**, 311-314 (1992). https://doi.org/10.1016/0006-8993(92)91563-t

64   Voss, M. R. *et al.* Gender differences in the expression of heat shock proteins: the effect

of estrogen. *Am J Physiol Heart Circ Physiol* **285**, H687-692 (2003).

https://doi.org/10.1152/ajpheart.01000.2002

65   Mitra, A. *et al.* LOSS OF ROR2 TYROSINE KINASE RECEPTOR IS ASSOCIATED

WITH ENDOTHELIAL DYSFUNCTION IN PAH VIA INAPPROPRIATE INTEGRIN β1

ACTIVATION. *bioRxiv*, 2025.2004.2028.651135 (2025).

https://doi.org/10.1101/2025.04.28.651135

66   Medvar, B., Raghuram, V., Pisitkun, T., Sarkar, A. & Knepper, M. A. Comprehensive

database of human E3 ubiquitin ligases: application to aquaporin-2 regulation. *Physiol*

Genomics **48**, 502-512 (2016). https://doi.org/10.1152/physiolgenomics.00031.2016

67   Droste, P. *et al.* Semiautomated pipeline for quantitative analysis of heart

histopathology. *J Transl Med* **21**, 666 (2023). https://doi.org/10.1186/s12967-023-04544-

2

68   McAfee, Q. *et al.* Truncated titin proteins in dilated cardiomyopathy. *Sci Transl Med* **13**,

eabd7287 (2021). https://doi.org/10.1126/scitranslmed.abd7287

69   Rabkin, S. W., Lodha, P. & Kong, J. Y. Reduction of protein synthesis and statin-induced

cardiomyocyte cell death. *Cardiovasc Toxicol* **7**, 1-9 (2007).

https://doi.org/10.1007/s12012-007-0003-7

70   Powell, S. R., Davies, K. J. & Divald, A. Optimal determination of heart tissue 26S-

proteasome activity requires maximal stimulating ATP concentrations. *J Mol Cell Cardiol*

42, 265-269 (2007). https://doi.org/10.1016/j.yjmcc.2006.10.010

71   Conn, C. S. & Qian, S. B. Nutrient signaling in protein homeostasis: an increase in

quantity at the expense of quality. *Sci Signal* **6**, ra24 (2013).

https://doi.org/10.1126/scisignal.2003520

72   Navarrete-Perea, J., Yu, Q., Gygi, S. P. & Paulo, J. A. Streamlined Tandem Mass Tag

(SL-TMT) Protocol: An Efficient Strategy for Quantitative (Phospho)proteome Profiling

Using Tandem Mass Tag-Synchronous Precursor Selection-MS3. *J Proteome Res* **17**,

2226-2236 (2018). https://doi.org/10.1021/acs.jproteome.8b00217

73   Li, J. *et al.* TMTpro-18plex: The Expanded and Complete Set of TMTpro Reagents for

Sample Multiplexing. *J Proteome Res* **20**, 2964-2972 (2021).

https://doi.org/10.1021/acs.jproteome.1c00168

74   Schweppe, D. K. *et al.* Full-Featured, Real-Time Database Searching Platform Enables

Fast and Accurate Multiplexed Quantitative Proteomics. *J Proteome Res* **19**, 2026-2034

(2020). https://doi.org/10.1021/acs.jproteome.9b00860

75   Rad, R. *et al.* Improved Monoisotopic Mass Estimation for Deeper Proteome Coverage.

J Proteome Res **20**, 591-598 (2021). https://doi.org/10.1021/acs.jproteome.0c00563

Fig. 1 ROR2 is a developmentally restricted protein that is robustly upregulated in RVF.

(**a**) ROR2 protein expression was measured in the ventricular tissue using thiourea extraction by

western from P2 to P21 and in P21 atria (N=3-7/age group; statistics by One-Way ANOVA with

Two step FDR correction and T-test for P21 ventricle vs atria). (**b**) ROR2 immunohistochemistry

was performed to evaluate localization, which is observed at the cell membrane and nucleus.

(**c-f**) PAB was performed in 9-week-old C57BL/6J male mice with echocardiogram performed

after 2 weeks to assess RV remodeling and systolic function, Sirius red staining to assess for

fibrosis quantified using ImageJ (**g,** 10X magnification), western blot to assess induction of fetal

MYH7 (**h**) and qPCR to assess expression *Nppb* in the RV free wall. Protein fractionation was

performed to assess expression of ROR2 in the NCD (cytoplasmic, triton soluble) and the NCE

768   fractions (triton-insoluble) (**j-l**). ROR2 immunohistochemistry (**m**, 20X magnification) and

769   RNAscope (**n**, 63X magnification), was performed to assess cell type pattern of expression for

770   ROR2 induction in PAB using α-actinin and *Ryr2* to identify cardiomyocytes and *Dcn* to identify

771   fibroblasts in PAB RV. Statistics for PAB analyses by unpaired t-test or Mann-Whitney,

772   depending on normality (n= 3-6 for sham, n=5-12 for PAB).

Fig. 2 ROR2 regulates in vitro cardiomyocyte morphology. (**a/b**) NRVMs were cultured on

nanopatterned surface and received adenovirus to overexpress human ROR2, knockdown

ROR2, or express GFP alone. NRVM structure was evaluated by Airyscan confocal microscopy

(**c-e**, 63X magnification) using FIJI to generate maximum intensity projections and quantify cell

size and aspect ratio (major/minor axis using the fit ellipse). (**f-k**) FIJI was used to quantify mean

staining intensity; peripheral intensity enrichment was further measured as described in the

methods by subtracting the inner cell integrated intensity (total area shrunken by 2µm) from the

full cell integrated for α-actinin, β-catenin, and Cx43 (N=8-9 NRVM isolations, n= 341-411

782   cardiomyocytes/group). Total protein expression for these were also assessed by western blot

783   (n=3/group). One-Way ANOVA with two-stage step up Benjamini correction).

Fig. 3 ROR2 regulates in vitro cardiomyocyte proteostasis.

RNAseq with Voom/Limma was performed to assess differential gene expression between

ROR2[OE] and ROR2[KD] NRVMs, with volcano plot (**a**) highlighting notable genes including

transcriptional changes in the proteostatic pathways including folding (e.g. *Hspa1a/b, Hsph1,*

Cryab) and pathway enrichment analysis (**b**) performed by Gene Ontology Biological Processes

(n=4/group). Western blot was used to evaluate total ubiquitin (**c/d**, n=16/group) and HSP70

(**c/h**, n=10/group) between ROR2[OE], ROR2[KD], and controls. NRVMs were treated with 1μM

MG132 or DMSO for 24 hours prior to collection to assess the flux of K48-linked ubiquitin

clearance by the UPS (**e/f**, n=7/group). Proteasome capacity was measured using the

chymotrypsin proteasome substrate Suc-LLVY-AMC with cell lysates in the absence or

presence of indicated supplemented ATP (**g**, n=7-11/group and ATP condition). Protein

refolding was measured using heat shocked luciferase (**i**, n=11/group), measured by

luminescence after addition of luciferin. Protein synthesis rates were measured using 0.55

µg/mL tyrosyl-tRNA analogue puromycin for 1 hour pulse with immediate collection (**j**) or with a

subsequent 1 hour chase (**k**) including cycloheximide (20 µg/mL) to block further translation with

or without MG132 (20 µM) or HSP70^{OE} (n=6-8/group). Statistical comparisons were performed

with One-Way ANOVAs or Two-Way ANOVAs as indicated on graphs, with two-stage step up

Benjamini correction.

Fig 4. **Proteostatic disruption is responsible for ROR2-mediated cardiomyocyte structural and functional defects**

The impact of proteasome inhibition (**a-c**; MG132 1µM for 24 hours, n=129-183/group) and increased HSP70 expression (**d-f**; 24 hours of adenoviral expression of *Hspa1b*, n=65-111/group) on ROR2^{OE}-mediated change in cell area and aspect ratio was evaluated as described above. Using a Cytocypher pacing apparatus, the impact of ROR2^{OE} and ROR2^{KD} on contractile and relaxation kinetics (**g-k**) were measured using CytoMotion Pixel Correlation with IonOptix software (n=48-75/group from three separate NRVM isolations). Statistics by Two Way ANOVA without correction (**a-f**) or with two-stage step up Benjamini correction (**i-l**).

Fig. 5 ROR2 mediated proteostatic disruption impacts cardiomyocyte structural proteins

The interaction of ROR2[OE] and proteasome inhibition on the impact of the NRVM proteome was

investigated by TMT-labeled mass spectrometry-based proteomics using Voom/Limma to

evaluate differential protein abundance, with visualization by volcano (**a/b**) and PCA plots (**c**).

Normalized abundance of representative (**d**) E3 ligases is displayed by dotplot and heatmap (**e**)

using a previously published comprehensive list of human E3 ligases.[66] (**d/e**) Differential protein

abundance enrichment analysis was performed for GO Biological Processes (**f**). Representative

change in actin and sarcomere associated proteins with ROR2[OE] and MG132 are displayed by

dotplot (**g**).

Fig. 6 Cardiac ROR2 induction is sufficient to cause RVF and proteostatic disruption in

mice. (**a**) The impact of cardiac ROR2^{OE} was tested using AAV9 delivery of HA-tagged *Ror2* vs

GFP in male and female mice at age 4 weeks with phenotyping performed at age 12 weeks.

Cardiac ROR2^{OE} caused a male predominant phenotype with increase in RV wall area **(b**

representative 10X magnification and **c**) quantified using FIJI with color deconvolution to identify

epi and endocardial borders. Total heart mass after trimming great vessels and RV free wall and

LV mass were measure after removal of atria and separation of RV free wall (**d-f**). RV

cardiomyocyte cross sectional area was measured using HeartJ[67] FIJI package on WGA

stained section and quantifying the minimum Feret Diameter (**g/h**, 40X magnification).

Echocardiography was performed to assess biventricular size and systolic function (**i-l**) and

jugular catheterization was performed to assess RV end diastolic pressure (**m**). Expression of

Nppb was measured by qPCR (**n**) and fetal MYH7 was measured by western blot (**o**). Fibrosis

was measured using trichrome staining of RV free wall tissue (**p**). Proteostatic characterization

including assessment of *Hspa1b* by qPCR (**q**), ubiquitin immunohistochemistry of HA-tag (i.e.

AAV infected cardiomyocytes) (**r/s**, 60X magnification), puromycin labeling to measure protein

synthesis rates using a 1-hour puromycin pulses by intraperitoneal injection (**t**) and proteasome

capacity using Suc-LLVY-AMC (**u**). Statistical comparisons performed using t-test or Mann

Whitney depending on normality, (n=3-8 mice/group and 32-72 RV cardiomyocytes/group for

panel **s**).

Fig. 7 ROR2 knockdown improves RVF and proteostatic burden in RV pressure overload.

Proteostatic characterizations were performed for PAB and sham mice (**a**) by puromycin

labeling to measure protein synthesis (**b**), expression of HSP70 (**c**), total ubiquitin (**d**), K48-

linked ubiquitin (**e**), and PSMB5 expression (**f**) by western blot in NCD and NCE fractions.

Proteasome capacity was measured using Suc LLVY-AMC (**g**). ROR2^{KD} was achieved by

AAV9-delivery of *Ror2* targeting shRNA (PAB^{shRor2}) delivered by retroorbital injection 4-weeks

prior to PAB/sham, which reduced NCD (~25%) and NCE (~50%) ROR2 expression compared

to scrambled shRNA (PAB^{Scr}; **h-j**). Echocardiography was performed 2 weeks after PAB to

assess RV remodeling, systolic and diastolic function (**k-m**). Histology using Sirius red of RV

free wall and FIJI to assess RV area and fibrosis percent using colour deconvolution (**n-p**).

Proteostatic state in PAB^{shScr} compared to PAB^{shRor2} was characterized using puromycin

labeling, K48 ubiquitin and HSP70 expression, and proteasome capacity by Suc-LLVY-AMC as

above (**q-t**). Statistics by t-test or Mann Whitney depending on normality (n=6-7 for Sham vs

PAB n=9-12 and n=5-15 PAB^{shScr} vs n=7-14 for PAB^{shRor2}), or Two-Way ANOVA for proteasome

capacity.

Fig. 8 ROR2 expression is associated with proteostatic disruption in human RVF. (**a**)

ROR2 expression in NCE (filled in symbols) and NCD (open symbols) fractions was measured

in the RVs from humans with RVF from dilated cardiomyopathy with (purple), preserved RV

function (orange triangles, pRV), or nonfailing (NF, black). RVF samples were further stratified

by high (red symbols) and low (blue symbols) NCE/NCD ROR2. (**b**) Proteasome capacity was

compared across human RV functional groups. (**c**) Within RVF, NCD/NCE ROR2 levels were

assessed for Pearson correlation against proteasome capacity (noted below graph) and for

differences in linear slopes. Statistics by Kruskal-Wallis (**a**) or Two Way ANOVA (**b**) with two-

stage step up Benjamini correction (n=9-11/group). (**d**) Taken together, our work reveals a

model wherein high ROR2 expression in cardiomyocytes disrupts proteostasis with increased

protein synthesis (1) and reduced folding (2) that results in a poor-quality proteome, leading

876   cardiomyocytes to clear proteins by the UPS (3), which disrupts cardiomyocyte structure and

877   function (4).