Title: Sialidases derived from *Gardnerella vaginalis* and *Prevotella timonensis* remodel the

sperm glycocalyx and impair sperm function

Authors: Sarah Dohadwala[1], Purna Shah[2], Maura K. Farrell[2], Joseph A. Politch[1], Jai Marathe[1],

Catherine E. Costello[1,2], Deborah J. Anderson[1]

*correspondence to: Deborah.Anderson@bmc.org

670 Albany Street, Rm 516

Boston, MA 02118

Affiliations

[1]Boston University Chobanian and Avedisian School of Medicine, Boston MA

[2]Boston University, Boston MA

Running head: Sialidases from Dysbiotic Vaginal Bacteria Impair Sperm Function

Keywords: bacterial vaginosis, sialidase, reproductive immunology, sperm, infertility

Supplementary data: Figures S1 – S10 attached.

Abstract

Bacterial vaginosis (BV), a dysbiosis of the vaginal microbiome, affects approximately 30 percent of women worldwide (up to 50% in some regions) and is associated with several adverse health outcomes including preterm birth and increased incidence of sexually transmitted infections (STIs). BV-associated bacteria such as *Gardnerella vaginalis* and *Prevotella timonensis* damage the vaginal mucosa through the activity of sialidase enzymes that remodel the epithelial glycocalyx and degrade mucin glycoproteins. This damage may contribute to adverse health outcomes. However, whether BV-associated glycolytic enzymes also damage sperm has not yet been determined. Here, we show that sialidase-mediated glycocalyx remodeling of human sperm increases sperm susceptibility to damage and adversely affects their function in vitro. Specifically, we report that sperm motility was not adversely affected by sialidase treatment, but desialylated human sperm demonstrate increased susceptibility to agglutination and complement-mediated cytotoxicity as well as impaired transit through cervical mucus. Our results demonstrate mechanisms by which BV-associated sialidases affect sperm survival and function and potentially contribute to adverse reproductive outcomes such as infertility.

            Introduction

33   Bacterial vaginosis (BV) is a condition characterized by vaginal dysbiosis that affects

approximately one third of women globally (Peebles et al. 2019). In some regions in the global

South, up to 50% of women have BV (Kenyon et al. 2013). BV is an extremely variable

condition, characterized by high diversity in the vaginal microbiome, and clinical symptoms such

as irritation, discharge, itching, and fishy odor (Abou Chacra et al. 2022). To more precisely

define the composition of vaginal microbiota found in humans, the field has adopted community

state types (CSTs) numbered I to V to describe various proportions of different bacterial taxa

(Ravel et al. 2011). A healthy vaginal microbiome, or CST I, consists primarily of *lactobacillus*

crispatus, which maintains low vaginal pH and inhibits inflammation. A microbiome dominated

by *lactobacillus iners*, termed CST III is thought to be an intermediary microbiome between a

healthy and disease state. CST IV (i.e. BV) is characterized by a relative lack of lactobacilli and

the presence of dysbiotic anaerobic vaginal bacterial species such as *Gardnerella vaginalis* and

Prevotella timonensis. BV is associated with spontaneous preterm birth, increased risk of

preclinical pregnancy loss, and the acquisition of sexually transmitted infections (STIs) (Howe et

al. 1999; Ralph et al. 1999; Cauci and Culhane 2011; van Oostrum et al. 2013; Armstrong and

Kaul 2021; Abou Chacra et al. 2023; Mohanty et al. 2023, Hadhoum et al 2025.). In addition,

emerging evidence suggests a link to infertility and pre-eclampsia, as well as adverse neonatal

outcomes even for full term infants (Denney and Culhane 2009; van Oostrum et al. 2013;

Dingens et al. 2016; Kindschuh et al. 2024; Ravel et al. 2021).

52   To date, the mechanisms by which BV increases the risk of adverse reproductive

outcomes are poorly understood (Ugwumadu 2002; Mania-Pramanik et al. 2009; Elovitz et al.

2019; Gudnadottir et al. 2022); several have been proposed, including higher vaginal pH, an

inflammatory vaginal milieu, epithelial damage and altered innate immunity (Cauci et al. 2008).

Recent findings have implicated sialidases, enzymes that cleave sialic acid residues from

glycoproteins and glycolipids, in BV-associated adverse reproductive outcomes. Several BV-

associated bacteria such as *Gardnerella vaginalis and Prevotella timonensis* can produce high

levels of sialidases, and sialylation of the vaginal epithelium and mucus is markedly lower in

women with BV than those with a healthy microbiome (Agarwal et al. 2023; Segui-Perez et al.

2024). In particular, elevated sialidase activity at 12-weeks gestation has been correlated with a

higher risk of early spontaneous preterm birth and miscarriage, with a stronger correlation than

for BV generally (Cauci et al 2008). BV-associated sialidases have been shown to degrade the

vaginal glycocalyx and mucus (Pelayo et al 2024; Agarwal et al 2023). While BV-associated

degradation of the vaginal mucosa certainly contributes to clinical symptoms of BV (Roselletti et

al. 2020; Cheu et al. 2022), the mechanistic connection between BV-associated vaginal

changes and adverse reproductive outcomes, which primarily occur in the upper female

reproductive tract (FRT), remain unclear. Sperm migrate from the vagina to the upper FRT and

it has been postulated that abnormal sperm play a role in female reproductive disorders

(Mahdavinezhad et al. 2022). In addition, recent evidence suggests that sperm can modulate

the uterine immune milieu which may have an impact on the timing of gestation (Green et al.

2020; Katila et al. 2001; Schjenken et al 2021). Accordingly, the purpose of this study was to

determine whether BV-associated sialidases derived from *Gardnerella vaginalis and Prevotella*

timonensis affect the human sperm glycocalyx and impact sperm function. The sialylated sperm

glycocalyx protects them from the female immune system, facilitates their passage through

cervical mucus, and prevents premature capacitation, a process necessary for fertilization

(Tecle and Gagneux 2015; Fliniaux et al. 2022). It is estimated that mature sperm carry

approximately 30 million sialic acid molecules per cell (Ghaderi et al. 2011, PNAS). Thus, BV-

associated sialidases may adversely affect sperm migration, immune signaling, and fertilization

capacity.

We used three recombinant sialidases, GvNanH2, PtNanH2 and PtNanH1, derived from

clinically relevant strains of *Gardnerella vaginalis* (*Gv*) and *Prevotella timonensis (Pt)*, to study

the effects of BV-associated sialidases on human sperm. These enzymes reportedly have

different activities: the NanH2 enzymes cleave both *N-* and *O-*linked glycans and play an

important role in degrading the protective sialoglycan mucus barrier lining the lower female

reproductive tract, whereas the NanH1 enzyme demonstrates reduced enzymatic activity

including towards sialic acids linked to human MUC5B and other mucin substrates (Agarwal et

al. 2023, Pelayo et al 2024). We employed lectin flow cytometry and zeta potential to

characterize the changes to surface glycan moieties and charge on the remodeled sperm

glycocalyx after sialidase treatment. We next investigated the effect of desialylation on sperm

motility, the ability of sperm to traverse through cervical mucus, and on spontaneous and

antibody/lectin-induced sperm agglutination. Lastly, we investigated the susceptibility of

desialylated sperm to complement-mediated cytotoxicity. Our data provide strong evidence that

BV-associated sialidases adversely affect several sperm functions, potentially contributing to the

adverse reproductive outcomes such as subfertility and preterm birth that are associated with

bacterial vaginosis.

Results

Sialic acids are depleted from the sperm glycocalyx after exposure to recombinant BV sialidase, exposing cryptic epitopes and reducing negative surface charge.

Sperm are rich in sialylated cell surface proteins that may be susceptible to hydrolysis by BV-associated sialidases. Accordingly, to investigate BV-associated damage to sperm, we produced recombinant, low endotoxin GvNanH2 as described previously (Robinson et al. 2019, **Fig S1 – S2**). GvNanH2 is a truncated variant of a sialidase derived from *Gardnerella vaginalis.* Of note, the *Prevotella* species do produce endotoxin in the vaginal tract; in our study, *Prevotella* enzymes were produced in BL21 E.coli, and as a result likely contain endotoxin (Aroutcheva et al. 2009). Controls to determine the effects of endotoxins were incorporated into the experiments as described below. To ensure that our experiments employed physiologically relevant amounts of sialidase, we used a range of activity levels of enzyme derived from cervicovaginal swabs obtained from women with BV (0 to 2 uM 4MU/min sialidase activity, mean = 0.5) from Agarwal et al. 2023, Figure 1B. Given those 4-MU activity levels from vaginal swabs, we used 0.88 and 0.11 uM 4-MU/min/uL of enzyme activity to represent BV-like conditions and standardize our assays. To confirm that BV-associated sialidases desialylated sperm-surface glycans, sperm were collected from healthy, reproductive aged men, isolated by density gradient, and treated with either recombinant GvNanH2, AUS (a commercially available sialidase derived from *A. ureafaciens*, positive control), or no treatment (negative control). Sperm were incubated in non-capacitating conditions. Then, we assessed removal of terminal sialic acids with lectin flow cytometry using SNA-Cy5 and MAL II-biotin to measure a decrease in 2,6 and 2,3-linked terminal sialic acids, respectively (**Fig 1A**). The untreated controls demonstrated the presence of 2,3 and 2, 6 linked sialic acids on the sperm surface (**Fig 1B, 1C).** This corresponds to results from previous studies using lectin microarrays to profile the sperm surface glycome (Xin et al. 2014). For SNA-Cy5 staining, the direct conjugate enabled a direct comparison of the activity of AUS and GvNanH2. AUS reduced the signal by roughly 50%

relative to the control, while GvNanH2 reduced it by approximately 15% (**Fig 1C**). Similarly, a

significant reduction was observed in the staining for MAL-II, though due to a secondary

amplification step, a linear relationship to compare AUS and GvNanH2 was not possible.

Although not statistically significant, a three-fold decrease in signal was observed with NanH2

compared to AUS (**Fig 1B**).

Sperm, being foreign to the female reproductive tract, have immunogenic epitopes

capped with terminal sialic acids that function in processes such as the acrosome reaction and

sperm-egg interaction (Ma et al 2016). To determine if the glycan moieties underlying terminal

sialic acids were exposed or degraded after sialidase treatment, we used lectin flow cytometry

with pea savitum agglutinin (PSA), a lectin which binds *N*-acetyllactosamine residues and is a

marker for the acrosome reaction (**Fig 1A**). In untreated (control) sperm, robust PSA staining

was observed (**Fig 1D**). In both sialidase treatment conditions (AUS and NanH2), sperm PSA

staining was unchanged from that of the control (**Fig 1D**). This indicates that sialidase treatment

does not result in a premature acrosome reaction under these conditions.

Sperm surface sialic acids contribute to a negative surface charge. To determine if the

surface charge of sperm is modified in response to sialidase treatment, the zeta potential of

sperm cultures was measured after treatment with different doses of enzyme. First, we

optimized the diluent for the assay and confirmed the effect of different culture conditions on the

measurement (**Fig S4A**)**.** A decrease in sperm surface charge was observed when whole

semen was treated, though the magnitude of the change observed was lower than when

washed sperm were treated (**Fig S4B**). It is possible that sialylated glycoproteins in seminal

plasma protect the sperm surface from the full force of the sialidase enzymes through a

competition mechanism. Next, we measured the change in zeta potential to compare the same

doses of recombinant NanH2 and AUS and found that effects were similar (**Fig S4C**). Finally,

we used a series of doses of GvNanH2, PtNanH2, PtNanH1, and AUS sialidase and measured

the zeta potential of the sperm. We observed a dose dependent decrease in the magnitude of

the negative surface charge after a 1-hour incubation for AUS, PtNanH2, and GvNanH2, though

we did not observe a significant decrease with PtNanH1 (**Fig 1F**). This indicates that glycans

that cannot be cleaved by PtNanH1 are a major contributor to the negative surface charge of

sperm. Based on the limited data on the substrate specificity of PtNanH1 and PtNanH2, we

surmise that these sialylated proteins may be similar to mucin glycan substrates such as

MUC5B in structure or accessibility. We also measured the donor-to-donor variance in the

biological replicates and found that there was some variance in how the sperm from different

donors were affected by the sialidases, though overall trends in the data were retained (**Fig**

S4D).

159   **Desialylated sperm are more susceptible to agglutination**

Agglutination is a phenomenon driven by cell motility and surface charge wherein motile

sperm collide and stick to one another, forming large, aggregated networks. We investigated the

propensity of highly motile sperm to agglutinate after sialidase treatment. We hypothesized that

by reducing the negative charge, sperm desialylation would make the sperm stickier and prone

to agglutination.

To measure this, washed sperm were treated with different doses of PtNanH2 sialidase

and observed over time. We observed variable degrees of spontaneous sperm agglutination

under these conditions, largely because agglutination is a concentration dependent

phenomenon that relies on cell-cell collisions. The variable characteristics in these spontaneous

agglutinates included size and shape of agglutinates (**Fig S9, S10**). Next, we tested whether

sperm desialylation affects antibody- or lectin-induced sperm agglutination. To quantify this

effect, we performed a kinetic agglutination assay on sperm treated with sialidase or heat

inactivated enzyme (negative control), followed by Galectin-1, the Human Contraception

Antibody (HCA), or Pea savitum agglutinum (PSA) **(Fig 3A).** These entities were selected due

to their importance in reproductive biology. Galectin-1, an endogenous lectin, is involved in

binding STI pathogens and is expressed in FRT tissues. HCA, a sperm-binding monoclonal

antibody derived from the blood of an infertility patient, is part of a contraceptive product that is

poised to enter Phase II clinical trials (Anderson et al. 2020). Finally, PSA is a marker for the

acrosome reaction, a process crucial to sperm-egg interaction. These three entities bind distinct

glycans-- HCA binds a male reproductive tract specific, but largely uncharacterized sperm *N*-

glycoprotein; Galectin-1 binds repeat *N*-acetyllactosamine units, and PSA binds terminal

mannose residues. None of these three entities bind sialic acid moieties specifically, though

they may all be tolerant of sialylated glycans.

We optimized the concentrations of the agglutinating antibodies and lectins, selecting

low concentrations that were weakly agglutinating to distinguish the effect of sialidases

treatment. We found that sperm agglutinated significantly faster in response to each of the three

agglutinating entities after treatment with either AUS or GvNanH2 sialidase (**Fig 3B, 3E**). We

also tested the *Prevotella* sialidases using HCA to assess whether the specificity differences

between PtNanH1 and PtNanH2 would have different functional implications related to cell

adhesion and agglutination. We found that PtNanH1 did not increase agglutination speed, while

PtNanH2 did, similar to GvNanH2 in a dose dependent manner (**Fig 3C, 3F**). This indicates that

sialylated glycans that are cleaved by PtNanH2 but not by PtNanH1 have important roles in

preventing agglutination and cell-cell sticking. Given reports in the literature that demonstrate an

impact of endotoxin (lipopolysaccharides, LPS) on sperm capacitation, we used polymyxin-B, an

antibiotic that neutralizes the effects of LPS, to isolate the role of sialidases and exclude

contributions from endotoxin in our assays (Bhagwat et al. 2025). Sperm agglutination assays

were repeated in the presence of sialidase pre-treated with polymyxin-B, and there was no

difference in activity, indicating that the effect on sperm agglutination can indeed be attributed to

sialidase activity (**Fig 3D**). In addition, for PSA, flow cytometry data indicated that surface

staining is unchanged in response to sialidase treatment, suggesting charge-mediated repulsion

in preventing agglutination, rather than enhanced binding of underlying glycan epitopes in

treated sperm (**Fig 1D).** In sum, removal of surface sialic acids from sperm increased their

propensity to agglutinate spontaneously and in the presence of several distinct lectins and

antibodies.

Desialylated sperm are more susceptible to complement-mediated immobilization and

cytolysis

Sialic acids prevent aberrant complement activation by binding several complement

inhibitors including Factor H (Blaum et al. 2015). Both complement and complement inhibitors

can be found in human cervical mucus and seminal plasma (Price and Boettcher 1979;

Petersen et al. 1980) suggesting that complement biology may be important to fertility and

reproduction, though specific mechanisms are not yet fully understood. For instance,

complement proteins may serve as a bridge between sperm and egg (Anderson et al. 1993),

and some antisperm antibodies demonstrate potent complement-dependent immobilization and

cytolysis of sperm (Baldeon-Vaca et al. 2021).

We used the complement-mediated sperm immobilization assay to determine the

potential role of sialidase enzymes in spontaneous- and lectin- mediated complement activation.

We first investigated whether sialidase treatment alone affected sperm motility. To isolate the

measurement of sperm motility from the spontaneous agglutination we sometimes observed,

these motility assays were done at low concentrations of sperm at which agglutination was not

observed. We did not observe a significant effect of any of the tested sialidases on sperm

motility through 60 minutes, after which time we observed a slight decrease in sperm motility in

some donors (**Fig 2A, 2C).** In addition, we plotted average sperm motility parameters that relate

to capacitation, including curvilinear velocity, amplitude of lateral head displacement, and

linearity over time to determine if sialidase treatment impacted these parameters rather than

total motility. We found that, under non-capacitating experimental conditions for 90 minutes,

sialidases do not impact hyperactivation motility parameters associated with sperm capacitation

(**Fig 2B-D, 2G-H**). Next, we examined the effect of sialidases on sperm motility in the presence

of complement. Addition of GvNanH2 and PtNanH2 with complement significantly reduced

sperm motility compared to all three negative controls (no enzyme treatment, heat inactivated

complement, addition of compstatin, a C3 complement inhibitor) (**Fig 4A, 4B**). We found that

sperm exposure to PtNanH1 did not mediate complement immobilization, which indicates that

the sialylated glycans that are cleaved by the NanH2s, but not cleaved by PtNanH1 play

another critical role on the surface of sperm in protection from complement (**Fig 4B, 4C**). Finally

we confirmed that complement-mediated sperm immobilization is a surrogate for cell lysis with

trypan blue staining (**Fig S10**). After complement treatment of sialidase-treated sperm the

majority of immotile sperm stained positive for trypan blue, indicating that they were lysed (**Fig**

S10). The majority of lysed sperm were present in agglutinates, which could indicate that sperm

become sticky after lysis, or that agglutination facilitates complement activation on the surface

of sperm due to increased surface area of the agglutinate (Dalia et al 2011). We confirmed that

complement activation could be abrogated using sialidase inhibitors, including DANA, a

competitive inhibitor, and Zanamivir, a sialidase inhibitor used for treatment of flu. PtNanH2 was

susceptible to inhibition by Zanamivir, while GvNanH2 was susceptible to DANA (**Fig 4D**). In

addition, we explored antibody-mediated complement lysis, and found that sperm were more

prone to complement lysis through the classical complement pathway in the presence of

sialidase (**Fig S5**). In sum, these data provide evidence that surface sialic acid moieties protect

sperm from complement-mediated lysis. Finally, given prior reports that endotoxin (which can be

found in recombinant bacterial enzymes) may itself impact sperm health, we repeated

complement assays in the presence of 100 ug/mL polymyxin B to neutralize endotoxin

(Bhagwat et al. 2025). We observed a small (10.5%) but statistically significant motility

difference between the polymyxin pre-treated GvNanH2 and GvNanH2 alone (**Fig 4E**). Thus,

endotoxin may account for some of the observed sperm immobilization in our assays. However,

in the presence of polymyxin B, sialidase-treatment resulted in an average motility reduction of

approximately 40% (GvNanH2) and 70% (PtNanH2), compared to polymyxin only controls (**Fig**

4E). The differences in motility between the sialidase-treated and no treatment group in the

presence of polymyxin remain highly significant indicating that sialidase treatment does account

for the increased sensitivity to complement-mediated immobilization.

Desialylated sperm are less able to transit through human cervical mucus

Human cervical mucus varies in physical properties and composition throughout the

menstrual cycle and is rich in negatively charged mucins. Sperm must transit across this

mucosal barrier to access the upper reproductive tract and fertilize an egg. To determine how

sperm transit through mucus may be affected by sialidase treatment, we assessed numbers of

sperm observed at progressive distances through a capillary tube filled with ovulatory human

cervical mucus. We started with a model that simulates sperm transit through the vagina and

cervix after intercourse (**Fig** 5A). Motile sperm were isolated, resuspended at a concentration of

80 million/mL in 50% seminal plasma, and placed into a capillary tube to measure migration

through a small region (0.5 cm) of GvNanH2 sialidase or media (control), followed by midcycle

cervical mucus (**Fig** 5A). We found a significant reduction in the numbers of total sperm that

were able to transit through the mucus after exposure to sialidase (**Fig** 5B).

Next, we measured different methods of sialidase exposure since the localization of the

bacterial enzyme in the cervix is unknown. We mixed human midcycle cervical mucus with

recombinant sialidase (amount equivalent to 0.5 µM/min 4-MU hydrolysis activity) or media

(control) and counted the number of sperm that penetrated to various depths of the capillary

tube after 90 minutes. Pre-treatment of mucus with sialidase initially facilitated sperm transit, but

the sperm became immotile as they traversed the sialidase-containing mucus (**Fig S7**). While

the numbers of sperm found in mucus increased in the sialidase-containing mucus, the percent

motile sperm decreased, which was the opposite of the trend observed in untreated mucus (**Fig**

S7). This may be due to more efficient complement lysis of the sperm exposed to sialidase

during transit through the cervical mucus. Next, we tested treatment of sperm with sialidase

prior to mucus exposure and found that this also reduced their transit through cervical mucus

(**Fig S7**). We also tested these phenomena with and without seminal plasma, which contains

several proteins known to coat the sperm surface (**Fig S6**). We found that seminal plasma

protects sperm from the immobilization observed in sialidase-containing mucin, perhaps due to

abundant seminal sialoglycoproteins in seminal plasma (**Fig S6**). Finally, we also compared

sperm transit through human cervical mucus to a commonly used commercially available mucin

model, bovine submaxillary mucin. In the absence of seminal plasma, a similar pattern was

observed; however, in the presence of seminal plasma, human cervical mucus was much more

permissive to sperm transit (**Fig S8**). This indicates some specific interaction between

components of human cervical mucin and seminal plasma that is not recapitulated with bovine

mucin. In sum, our data indicate that BV-associated sialidases affect sperm transit through

human midcycle cervical mucus, both by altering the sperm's forward progression in mucus,

and by altering the properties of the mucus itself.

Discussion

Our study demonstrates that sialidases derived from BV-associated bacteria alter the sperm glycocalyx by removing terminal sialic acid residues. Furthermore, it provides evidence that the unperturbed sperm glycocalyx facilitates the transit of sperm through cervical mucus and likely protects sperm from immune effectors such as complement in the female reproductive tract. Desialylated sperm were highly susceptible to complement-mediated damage and agglutination, and demonstrated impaired penetration through midcycle cervical mucus. Prior to our work, most studies on the effects of BV-associated sialidases were limited to the vaginal mucosa (Agarwal et al. 2023; Segui-Perez et al. 2024), and the importance of sperm surface sialic acids in bacterial infection and capacitation (Khosravi et al. 2016). Studies related to the effects of sialidases on sperm were mostly conducted in animal models which supported the idea that exposure to sialidases affects sperm function in the reproductive tract (Ma et al. 2016; Fernandez-Fuertes et al. 2018).

Several orthogonal lines of evidence support the idea that BV-associated sialidases may damage sperm and affect fertility: 1) *G.vaginalis* is found at a higher frequency in semen from infertile men (Andrade-Rocha 2009) and oligozoospermic seminal plasma contains higher levels of free sialic acid relative to normozoospermic samples (Menevse et al. 2022), 2) other bacterial conditions that involve glycolytic enzymes are mechanistically linked to male infertility (Khosravi et al. 2016), 3) multiple studies with sperm from various other mammalian species have pointed to the importance of the glycocalyx in protecting from innate immunity in the FRT (Fliniaux et al. 2022), 4) the semen microbiome is associated with sperm health (He et al. 2024; Osadchiy et al. 2024) and 5) natural reproductive processes, such as capacitation, the acrosome reaction and sperm-egg fusion, are sialidase-dependent and rely on the timely exposure of relevant surface glycans (Feng et al. 2016). Finally, several animal models demonstrate that sialic acid moieties on sperm can impact immune cells and endometrial reproductive processes (Toshimori et al. 1991; Froman and Thursam 1994; Machado et al. 2014; Álvarez-Rodríguez et al. 2020;

Batra et al. 2020; Schjenken et al. 2021; Donnellan et al. 2023; Li et al. 2024). In summary,

human sperm are an underappreciated target for bacterial sialidase enzymes associated with

BV; premature desialylation likely impacts sperm function and the immunology of the female

reproductive tract.

We found that desialylated sperm provoke complement activation resulting in sperm

immobilization and death. In addition, sperm exposure to endotoxin may synergize with

sialidase treatment to further sensitize sperm to complement immobilization (Bhagwat et al.

2025). The premature death of sperm in the vaginal and cervical environment may explain the

association of BV with infertility (Ravel et al. 2021). The appropriate activation of the

complement cascade is important to several reproductive processes, including fertilization and

early pregnancy, and disturbances resulting in aberrant complement activation are associated

with adverse pregnancy outcomes including preterm birth (Anderson et al. 1993; Riley-Vargas

et al 2005; Lynch et al. 2011; Lynch et al 2008; Regal et al. 2015). In particular, higher levels of

complement components C3b and C5, and mediators such as mannose-binding lectin have

been found in vaginal swabs from pregnant women who deliver preterm compared to women

who delivered term (Chan et al. 2022). Given that sperm transit through the lower reproductive

tract to the upper tract, we hypothesize that desialylated sperm may initiate and carry some of

these aberrant complement cascades into the upper tract prior to fertilization (Schjenken et al.

2021; Denny et al 2012). Early processes in pregnancy, including implantation and placentation,

are inflammatory in nature, but overactivation of these pathways results in pathology; perhaps

desialylated sperm are partially responsible for the imbalance. (Tantengco and Menon 2022;

Norwitz et al. 2001; Norwitz 2006).

We found that sialidase did not affect sperm motility including hyperactivated motility.

Given that this study focuses on the impacts to sperm that may occur in the vaginal tract or the

cervix, we did not study impacts on sperm under capacitating conditions, which typically are

considered to represent the uterine environment. Given that sperm do alter their surface

sialylation during capacitation, further studies to clarify whether endogenous desialylation is

required for capacitation or whether bacteria mediated desialylation impacts capacitation could

be informative.

Next, we found that desialylated sperm are more prone to agglutination, probably due to

a reduction in the surface charge that removes inherent cell-cell repulsion. Our data indicate

that the surface charge of sperm is largely due to the presence of specific sialylated glycans

(those cleaved by NanH2s but not PtNanH1), which are crucial for cell-cell repulsion. Enhanced

agglutination may mean that contraceptive antibodies and lectins will be more efficacious in

women with sialidase-positive BV (Tsuji et al. 1988; Baldeon-Vaca et al. 2021). The process of

agglutination is entirely dependent on sperm concentration, due to its dependence on cell-cell

collisions. Accordingly, the extent and nature of spontaneous agglutinates varies and it remains

unclear how best to accurately measure spontaneous agglutination in a context that would

reflect in vivo conditions. Given that several leading clinical contraceptive candidate antibodies

as well as a few anti-HIV antibodies rely on glycan binding, it will be important to investigate

how women with sialidase-positive BV respond to these topical glycan-binding drug products

(Anderson et al. 2017). We believe the enhanced agglutination is due to reduced surface charge

of the sperm, rendering them "stickier. Several groups have proposed that sperm may carry STI

pathogens into the upper tract via cell-cell sticking, and the changes in surface charge due to

vaginal sialidases may exacerbate this phenomenon (Young et al. 2019).

We found that desialylated sperm traverse less successfully through ovulatory cervical

mucus. This supports data from cervical mucus in the sheep model, where differences in sialic

acid content in cervicovaginal tissue and mucus were associated with impaired fertility (Abril-

Parreño et al. 2022). The human cervical mucus penetration assay was originally designed as a

fertility test, so the impaired ability of desialylated sperm in this assay suggests that sialidases

may impact fertility (Hull et al. 1984). Studies on the nature of ovulatory human cervical mucus

are hindered by the difficulty of sample collection, particularly from women with untreated

bacterial vaginosis and in the absence of hormonal birth control, which can modify mucus

characteristics (Moncla et al. 2016). It is known that women with BV produce more mucus,

perhaps to counteract the bacterial metabolism of mucin glycoproteins observed in BV (Moncla

et al. 2016). However, it is unknown whether sialidases remain in the vagina or ascend into the

cervical environment and upper reproductive tract in women with sialidase-positive BV.

Given that mucin-like glycans contribute significantly to the surface charge of sperm and

mediate protection from complement lysis and agglutination, it is important to understand which

sialylated proteins mediate these functions, and to more granularly define the specificity

differences that exist between the NanH2s and PtNanH1. Much work remains in understanding

the roles of sialylated glycoforms of proteins on the sperm surface, and even how some

sialylated proteins, such as the semenogelins and beta defensin 126, which are abundant

mucin-like proteins critical for semen liquefaction and mucus penetration, respectively, adhere

to the sperm surface (Luo et al 2023, Diao et al 2014; Tollner at al 2011, Sci Trans, Med, Dorin

and Barratt 2014 Hum Reprod.)

This work has some limitations. The fertility status of our sperm donors was unknown;

we did not collect samples from infertile or subfertile men, so we cannot make assertions about

male infertility outside the FRT, or how bacterial sialidases may exacerbate pre-existing male

infertility. In addition, we did not screen sperm donors for BV-associated bacteria, though sperm

samples with poor motility were excluded. In addition to sialidases, several other glycolytic

enzymes including amylases and fucosidases are produced by BV-associated bacteria that we

have not yet explored. Due to limitations in human cervical mucin availability, we did not assay

the *Prevotella* sialidases' impact on sperm movement through mucus. Finally, the diversity in

bacterial vaginosis cases should also be further considered, given that molecular profiling has

revealed varying presentations of vaginal flora (Savicheva et al. 2023). *Gardnerella vaginalis*

and Prevotella timonensis, which produce the sialidases of focal interest in this study, are a

primary cause of BV (Morrill et al. 2020). However, it should be noted that not all genotypes of

G. *vaginalis* produce sialidase, and this may cause variation in clinical presentation of BV cases.

Furthermore, other species such as mycoplasma, express additional sialidases (Hardy et al.

2017). Clearly, in adverse reproductive outcomes associated with BV, there is also a role of the

damage to the FRT. This work did not study the interactions between damaged FRT and sperm.

The *in vitro* environment may not recapitulate certain aspects of physiology, such as fluid flow,

muscle contractions, as well as the menstrual cycle and hormone signaling on sperm. In

addition, we did not assess interactions between sperm and uterine immune cells which may be

impacted by desialylation, such as targeting of sperm by neutrophils via NETosis, sperm

phagocytosis mediated by uterine macrophages, or sperm immunomodulation of uterine NK

cells.

In summary, we demonstrate that BV-associated sialidases increase sperm

susceptibility to damage in the FRT and impede their penetration through cervical mucus,

providing evidence that these mechanisms may play an underappreciated role in adverse

reproductive outcomes. Understanding aberrant immune damage to sperm in bacterial

vaginosis raises new questions about how sperm may modify the immune milieu of the female

reproductive tract in both a eubiotic and dysbiotic condition. These findings highlight the need

for further investigation of the role of sperm in decidualization and ascending inflammation in the

upper reproductive tract, as well as a better foundational understanding of important processes

of early pregnancy in humans.

 Materials and Methods:

Materials:

Enzymes and substrates: AUS refers to Neuraminidase A (New England Biolabs P0722L). 4-

MU Sialic acid refers to 2'-(4-methylumbelliferyl)-α-D-*N*-acetylneuraminic acid sodium salt

hydrate (Sigma Aldrich 69587). GvNanH2 refers to truncated recombinant *Gardnerella vaginalis*

derived NanH2, as described below. PtNanH1 and PtNanH2 refer to enzymes derived from

Prevotella timonensis, isolated and produced as described in Pelayo et al 2024.

Bacterial strains, plasmids and DNA: ClearColi bacteria strains that produce low endotoxin

protein were a gift of the Dempsey lab at BUMC. The GvNanH2 plasmid was a gift of the Lewis

lab at UCSD.

Production of recombinant NanH2 (GvNanH2): The GvNanH2 plasmid contains a functional,

soluble, His-tagged, truncated form of NanH2, a sialidase expressed by *Gardnerella vaginalis*,

and a gene encoding kanamycin resistance on a pET28a backbone. The plasmid was

transfected into ClearColi by heat shock, then streaked for single colonies on kanamycin

selective Luria Bertani (LB) agar plates. The identity of the plasmid was confirmed by whole

plasmid sequencing (Plasmidsaurus). Colonies were picked after 2 days (ClearColi grows at

half the rate of BL21 E.coli), and inoculated overnight at 37° C with shaking in a 10 mL LB-

Kanamycin culture. Larger 250 mL cultures were inoculated and grown until they reached

optical density 0.4. Cultures were induced with 250 mM Isopropyl ß-D-1-thiogalactopyranoside

(IPTG) and grown overnight at 37° C after which time the bacteria were pelleted, and stored in a

lysis buffer (50 mM NaH2PO4 at pH 7.4, and 300 mM NaCl) at -80°C until use. To purify

protein, pellets were thawed, mixed thoroughly with a serological pipette, and then lysed in a

French pressure cell press thrice. The lysate was clarified by centrifugation thrice, then

incubated with Nickel NTA resin while shaking at room temperature for one hour. The resin was

pelleted and washed thrice in wash buffer (20 mM sodium phosphate, 300 mM sodium chloride,

25 mM imidazole), and then bound enzyme was eluted thrice in the elution buffer (20 mM

sodium phosphate, 300 mM sodium chloride, 250 mM imidazole). The eluants were pooled,

buffer exchanged (to remove imidazole) into storage buffer (50 mM NaCl, 20 mM Tris-HCl, 1

mM EDTA, pH 7.5), assessed for protein concentration via BCA, run on an SDS page gel (**Fig**

S1), then aliquoted into single use aliquots stored at -80° C until use.

<u>Lectins and antibodies:</u> Biotinylated Maackia Amurensis Lectin II (MAL II, Cat #:B-1265-1) and

Cy5-conjugated Sambucus Nigra Lectin (SNA, Cat #: CL-1305-1) were purchased from Vector

Labs. Human Contraception Antibody (HCA), was a gift from ZabBio. Recombinant human

Galectin-1 was purchased from ACROBiosystems (Cat# 50-210-6461) and Pea savitum

agglutinin (PSA-FITC) was purchased from Sigma (cat #L0770).

Methods:

<u>LC/MS-MS to confirm sequence of GvNanH2:</u> In-gel digestion using trypsin/Lys-C and LC-

MS/MS were used to confirm the protein sequence of the recombinant enzyme. LC-MS/MS

analysis was carried out using a nanoAcquity UPLC (Waters Technology) interfaced to a Q-

Exactive HF mass spectrometer (ThermoFisher Scientific.) Reversed-phase C-18 analytical and

trapping columns were used with a 75 minute method with a gradient of 2% B to 40% B in 40

minutes, using 99% water/1% acetonitrile/0.1% formic acid as mobile phase A and 99%

acetonitrile/1% water/0.1% formic acid as mobile phase B at a flow rate of 0.5 μl/min. Data-

dependent tandem MS spectra were acquired in positive mode for the top 20 most abundant

precursors. Full MS scans were acquired from *m/z* 350 to 2000 with a resolution of 60,000 using

an automatic gain control target of 3e6 and maximum injection time (IT) of 100 ms. Dynamic

exclusion (12 s) was enabled. Precursor ions were fragmented using a resolution of 15,000 with

a maximum injection time of 50 ms and an automatic gain control value of 2e5 using higher

energy collision-induced dissociation with a stepped normalized collision energy of 27 and 35 V.

The MS/MS spectra were searched using Byonic against a fasta file containing the truncated

NanH2 sequence and provided assignments for 81 peptides; these represented 83.2% of the

predicted sequence (**Fig S2**).

<u>Sperm isolation:</u> Fresh semen was acquired from healthy donors after obtaining written

informed consent under the approved BUMC Institutional Review Board (IRB) protocol

#H36843, and processed as described previously in Mausser et al. 2023. In brief, samples were

processed within one hour of acquisition; motile sperm were separated by layering on a 90%

ISolate density gradient (FUJIFILM Irvine Scientific; Santa Ana, CA, USA) and centrifugation at

300g for 20 minutes. The motile sperm pellet was resuspended in Multipurpose Handling

Medium (MHM; FUJIFILM Irvine Scientific; Santa Ana, CA, USA). The concentration and motility

of sperm in whole semen and isolated motile sperm were assessed using a Computer-Assisted

Sperm Analysis system (CASA; Human Motility II software, CEROS II, Hamilton Thorne,

Beverly, MA, USA). Samples with low concentrations (<10 million/mL) or motility (<40%) post-

processing were not used for agglutination and complement experiments.

<u>Sperm sialidase treatment:</u> Motile sperm were resuspended to a concentration of 30 million per

mL in MHM. 40 µL of sperm suspension were incubated with 1 µL of enzyme at 37°C for the

chosen time-points and enzyme concentration. All enzymes were thawed on ice, used fresh,

and diluted into identical storage buffers (50 mM NaCl, 20 mM Tris-HCl, 1 mM EDTA, pH 7.5) to

maintain identical media conditions and concentrations, even as enzyme concentration varied.

<u>Sialidase activity assay:</u> Activity was confirmed by 4MU hydrolysis assay as previously

described (Robinson et al. 2019). In brief, to measure sialidase activity and match levels of

AUS, GvNanH2, PtNanH1 and PtNanH2 activity, a 4-methylumbelliferone-sialic acid assay was

used. 20 µL of enzyme solution was diluted into 100 µL of 600 µM 4MU-sialic acid (Sigma

Aldrich 69587), and immediately measured for fluorescence in a BioTek plate reader. The

measurement was performed in a 96 well plate every 1 minute for 2 hours at 37°C (Ex/Em:

365nm/448nm, Gain=43). A standard curve of 2-fold dilutions in triplicate of 4MU starting at 150

µM and ending with 2.34 µM was used to convert arbitrary fluorescence units into µM 4MU

hydrolysis per minute. The first 12 minutes of the reaction (<10% substrate consumption) was

used as a linear portion of the curve to calculate activity. rNanH2 (produced recombinantly as

described above) was used at a final concentration of at 0.27 ng/µL. AUS enzyme (NE BioLabs

P0722L) at 20000 U/mL was diluted into 100 mM sodium acetate buffer (pH 5.5) for a final

dilution of 1.8U/µL. Enzymes were thawed and kept on ice until use. *Prevotella* enzymes were

tested for activity to match activity levels to those used for GvNanH2 in various assays (**Fig** S1).

<u>Normalization of sialidase activity between different enzymes and calculations of enzyme doses:</u>

To normalize activity of enzymes, matched 4-MU activity levels were used on the same number

of sperm in our assays. In our studies, sperm were used at a concentration of 35 million per mL,

with volumes of 40 uL per treatment condition. Our activity levels in the data are described as

concentrations per total volume, as 0.88 uM 4-MU hydrolysis per minute per uL in the highest

treatment condition. An additional way is to describe enzyme activity per million sperm, which is

25.1 4-MU hydrolysis per minute per million sperm.

<u>Inhibition of Sialidase Activity Assay:</u> To measure activity of inhibited sialidase enzyme, activity

assays were performed using purified NanH2 enzyme and N-Acetyl-2,3-dehydro-2-

deoxyneuraminic acid inhibitor (DANA, Sigma Aldrich D9050). rNanH2 was used at a final

concentration of 0.27 ng/µL. DANA was used at final concentrations ranging from 1 mM to 10

pM. Enzyme and inhibitor were added in triplicate in a black-bottom 96 well plate. The substrate,

4-methylumbelliferone-sialic acid (Sigma), was added immediately prior to reading the plate.

Fluorescence activity was measured in a BioTek plate reader every 1 minute for 2 hours at 37°C

(Ex/Em: 365nm/448nm, Gain=43). A standard curve of 2-fold dilutions in triplicate of 4MU

starting at 150 µM and ending with 2.34 µM was used to convert arbitrary fluorescence units

into µM 4MU.

Endotoxin neutralization: To isolate the impact of endotoxin contamination in our experiments,

enzymes were pretreated with 100 ug/mL of Polymyxin B (Sigma Aldrich, Cat #: 81271) for 10

minutes prior to testing in agglutination and complement assays as described. Polymyxin binds

to the lipid A region on endotoxins and neutralizes their activity (Domingues et al 2012).

Lectin flow cytometry: To measure the effect of sialidases on sperm surface glycopeptides,

sperm were incubated with 160 U NanH2, 20 U AUS, or no treatment/heat-inactivated enzyme

for 1 hour and were fixed in formaldehyde. After fixation, $5x10^4$ sperm were stained per

condition at a final lectin concentration of 10 µg/mL for primary conjugates and 5 ug/mL for

biotin conjugates for 30 minutes. Biotin conjugates were reacted with Neutravidin-Alexa Fluor

488 at 2 µg/mL for 45 minutes. Fluorescence on sperm was analyzed using a Cytek Aurora full

spectral flow cytometer. Cells were gated by forward and side scatter and doublets were

removed by gating the linear FSC-A vs FSC-H. Populations were compared by determining

median fluorescence intensity for each condition in FlowJo software.

Zeta potential measurements: Zeta potential was measured by diluting 20 µL of sperm at a

concentration of 30 million per mL into 1 mL of distilled water immediately before measurement.

A Malvern zetasizer instrument was used at the following settings: the diluent setting was 10mM

NaCl, and seven measurements were taken per sample with no cooldown time between

measurements.

Sperm agglutination assay: Sperm were treated with sialidase as described above. Sperm with

and without sialidase pretreatment were exposed to three sperm agglutinating agents: Human

Contraception Antibody (HCA), and Galectin-1 and Pisum sativum agglutinin (PSA) lectins. A

kinetic agglutination assay was performed as previously described (Mausser et al 2023.) to

measure time to 95% agglutination.

<u>Complement-mediated sperm immobilization:</u> Sperm were treated with sialidase as described,

then exposed to human serum complement for 1 hour. The assay was performed as previously

described (Mausser et al. 2023), except antibody treatment was excluded. In brief, motile sperm

were quantified with CASA software after complement exposure. Heat-inactivated (Hi) sialidase

enzyme and Hi complement were used as negative controls. A human complement inhibitor,

compstatin (MedChemExpress, USA), was used as an additional negative control. Complement

and compstatin were incubated together for 15 mins at room temperature and then sialidase-

treated sperm were added as previously described. Classical complement-mediated sperm

immobilization was measured with an identical protocol, but including human contraception

antibody (HCA). As additional controls, sialidase activity was inhibited by addition of DANA(N-

Acetyl-2,3-dehydro-2-deoxyneuraminic acid) at 1mM or Zanamivir at 1mM (MedChem Express,

Cat. No.: HY-13210) to the maximum dose of enzyme (0.88 micromolar 4MU

hydrolysis/min/uL).

<u>Human cervical mucus penetration assay:</u> Cervical mucus samples were acquired after

obtaining written informed consent with the BUMC Institutional Review Board (IRB) approved

protocol #H41454. Midcycle ovulatory cervical mucus was collected from reproductive aged

women using an endocervical pipelle (Aspirette Endocervical Pipelle, Cooper Surgical,

Trumbull, CT, USA) within 48 hours of a positive ovulation test (Digital Ovulation Predictor Kit,

Clearblue). Participants had regular menstrual cycles, were not taking any hormonal birth

control, and refrained from intercourse for at least 48hrs prior to collection. The samples were

stored at 4° C until use (within 5 days). Cervical mucus was diluted at a 1:3 ratio of mucus: PBS

and aspirated into capillary tubes (Borosilicate Capillary Glass Slide, 0.30x3.0 mm, 50 mm,

Electron Microscopy Sciences, Hatfield, PA, USA) using a squirrel feeder and syringe. One end

of the tube was sealed using nail enamel, another end was filled with 10 µl of NanH2 sialidase

or media alone as a control. The capillary tube was inserted horizontally in a 1.5ml

microcentrifuge tube containing 80 µL of 20 x10^6/mL motile sperm. The assay was performed

as previously reported (Mausser et al, 2023). CASA images were taken at different points along

the capillary tube to evaluate the number of motile sperm reaching distances of 1, 2, 3 and 4

cm.

Sperm motility assay: Sperm motility was quantified over time using CASA software and

specialized microscope slides with a shallow chamber depth. Sperm at 30 million per mL were

treated with sialidase, incubated at 37C in a humidified chamber, and measurements were

taken at specific time points in the CASA. Fields of view lacking spontaneous agglutinates were

captured in this assay to enable accurate measurement of motility in the CASA software. At

least 300 sperm were measured per condition to determine a percent motility corresponding to 3

to 7 fields of view per time point per condition. To measure hyperactivation parameters including

curvilinear velocity, amplitude of lateral head displacement, and linearity for motile sperm, the

average values calculated by the CASA software were plotted. In particular, these average

values were calculated for motile sperm only, to exclude dead sperm with may skew the

averages towards zero motility and hide the signal from hyperactivated motility.

Sperm Live/Dead staining after complement treatment: Sperm were incubated with sialidase

and serum complement, as described for the complement-mediated immobilization assay.

Motility was measured with the Computer Assisted Sperm Analysis (CASA) software, then cells

were stained with trypan blue (1:1) to confirm whether percent motility inversely correlated with

the percentage of dead cells (trypan positive).

<u>Statistical analysis:</u> Continuous variables with >2 conditions were analyzed by analysis of

variance (ANOVA) or mixed-model analysis. A significant effect was followed by Holms-Sidak

multiple-comparison tests, or a Tukey multiple-comparison test when Holms-Sidak could not be

calculated. Variables were tested for normality by the Shapiro-Wilk test. If continuous variables

were not normally distributed, they were log (natural)- transformed prior to analysis. For

repeated measures ANOVA, the equality of variances of the pairwise differences between

within-subject conditions was assessed through tests of sphericity (Greenhouse-Geisser epsilon

and Huynh-Feldt epsilon). For 2-group/condition comparisons of continuous variables, unpaired

or paired t-tests or Mann-Whitney U tests or Wilcoxon matched-pairs signed-rank tests

comparing the slopes of linear regression lines were performed. Statistical significance was

assumed when $P<.05$. Data analysis and graphing were performed with Prism, Version 9.4.1

(GraphPad Software, Inc, La Jolla, CA) software.

Funding: This work was supported by the National Institutes of Health [P50 HD096957 to D.J.A

and T32-A1007309 to S.D] and a Boston University Sexual Medicine Pilot Grant to J.M.

Acknowledgements: We thank Warren and Amanda Lewis for generously providing the

plasmids encoding the *Gardnerella* sialidases, and Emily Balskus, and her students Claire

Xiang and Paula Pelayo for generously providing the *Prevotella* sialidase enzymes. We also

thank Daniel Dempsey, Nader Rahimi, Bjoern Reinhard, and Joseph Zaia, and members of their

lab(s), especially Laura Carretero Santos, Koustav Kundu and Margaret Downs, whose

assistance and willingness to share equipment greatly aided this study. We thank Kevin Whaley

for provision of HCA for these studies. We thank Shari Brezinsky from the BUMC Flow

Cytometry Core Facility for outstanding technical help, and Jolinada Zhang and James Doud for

contributing to motility and complement assays. Additionally, many thanks to Ashley Rebello,

Seth Bloom, Cristina Zamora, Allysa Allen, and Matt Geib for providing advice regarding the

study.

Abbreviations: Bacterial vaginosis (BV); sexually transmitted infection (STI); A. ureafaciens

sialidase (AUS); Female Reproductive Tract (FRT); Human Contraception Antibody (HCA);

Pisum sativum agglutinin (PSA)

Data availability statement: The data that support the findings of this study are available from

the corresponding author, [D.J.A], upon reasonable request.

Author contributions: SD designed the research. SD, PS, and MF generated data. JP, SD and

PS completed statistical analysis of the data. JP and JM obtained IRB approval and provided

human samples for the study. CEC and DJA supervised the research and contributed to

interpretation of the results. DJA and JM acquired funding for the study. SD took in the lead in

writing the manuscript. All authors discussed the results and contributed to the final manuscript.

Conflicting interests: The authors declare no conflicts of interest. The funders had no role in

the design of the study; in the collection, analyses, or interpretation of data; in the writing of the

manuscript; or in the decision to publish the results.

Figure 1: **BV-associated sialidases remove surface sialic acid from sperm**. A) Schematic describing lectin flow cytometry and representative glycan epitopes. B, C, D) Sperm were exposed to sialidases (AUS 0.11 4-MU hydrolysis/min/uL or GvNanH2 0.88 4-MU hydrolysis/min/uL) for 1 hour, fixed in formalin, and stained with MAL-II-biotin/Neutravidin-Alexa Fluor 488, SNA-Cy5, or PSA-FITC respectively. Each point represents the median fluorescence intensity (MFI) from one donor. Statistics represent a mixed model comparison with post-hoc Holm- Šidák tests. E) Schematic representing change in zeta potential after removal of negatively charged sialic acid on sperm surface. F) Sperm were treated with sialidase at various doses for 1 hour, and zeta potential was measured via dynamic light scattering. Each measurement represents the mean of seven technical replicates, and statistics represent a mixed model comparison with post hoc Holm- Šidák tests. * indicates $p \leq 0.05$, ** $p \leq 0.01$, ***$p \leq 0.001$, and **** $p \leq 0.0001$. Hi indicates heat-inactivated at the highest dose of enzyme (0.88 for PtNanH1, PtNanH2, and GvNanH2, and 0.11 for AUS).

Figure 2: Effect of sialidase treatment on sperm motility A) Motile sperm were isolated, and motility
was measured via computer assisted sperm analysis (CASA) after sialidase treatment (0.88 4-MU
hydrolysis/min/uL GvNanH2 and 0.11 4-MU hydrolysis/min/uL AUS). Statistics represent a repeated
measures comparison, and neither time nor enzyme condition were statistically significant. E) same as
A, except using PtNanH1 and PtNanH2 at 0.88 4-MU hydrolysis/min/uL. B, C, D) Average curvilinear
velocity (VCL), Amplitude of lateral head displacement (ALH), and Linearity (LIN), respectively, as
measured by CASA on the motile sperm population after treatment with enzyme (0.88 4-MU
hydrolysis/min/uL GvNanH2 and 0.11 4-MU hydrolysis/min/uL AUS). F, G, H) Same as B, C, D, except
using PtNanH1 and PtNanH2 at 0.88 4-MU hydrolysis/min/uL. Each measurement represents the mean
of four technical replicates. Statistics represent mixed model comparisons with post-hoc holms Sidak
tests, and neither time nor enzyme condition were a significant source of variation for LIN values. For
ALH and VCL values, only time was statistically significant (p<0.01).

674
Figure 3: Desialylation results in faster sperm agglutination. A) Schematic and representative
images representing sperm agglutination after sialidase treatment. B C, E, F) Time to agglutination in
the presence of different lectins and glycan-binding antibodies after treatment with GvNanH2,
PtNanH2, AUS and PtNanH1, respectively. Galectin-1 was used at a concentration of 10 µg/mL, PSA
was used at a concentration of 0.05 µg/mL, and HCA was used at 5 µg/mL. D) Time to agglutination in
the presence of 0.88 4-MU hydrolysis/min/uL of enzyme in the presence and absence of polymyxin B
(PMB). Each measurement represents the mean of three technical replicates. Statistics represent
repeated measures ANOVA with post hoc Holm-Šidák tests for GvNanH2, PtNanH2 ,and PtNanH1 (1B,
1C, 1D,1F), and unpaired t-tests for AUS (1E). * indicates $p \leq 0.05$, ** $p \leq 0.01$, ***$p \leq 0.001$, and **** p
≤ 0.0001. Hi indicates heat-inactivated enzyme at the highest concentration.

Figure 4: Effect of sialidase treatment on susceptibility to complement-mediated immobilization.
A, B, C) Sperm motility was measured after treatment with sialidase in the presence of complement. Heat-inactivated complement, or 125 uM compstatin, a human C3 inhibitor, were used as negative controls. Statistics represent mixed model comparisons with post-hoc Holms-Sidak tests. Hi indicates heat inactivated. D) The effect of sialidase inhibitors on complement immobilization of sperm with GvNanH2 and PtNanH2. DANA indicates N-acetyl-2,3-dehydro-2-Deoxyneuraminic Acid, and ZANA indicates Zanamivir. E) The effect of LPS inhibitor polymyxin B (PMB) on complement immobilization of sperm with GvNanH2 and PtNanH2 at 0.88 4-MU hydrolysis/min/uL of activity. Each measurement represents the mean of three technical replicates. Statistics represent a mixed model with post-hoc Holms-Sidak tests. * indicates $p \leq 0.05$, ** $p \leq 0.01$, ***$p \leq 0.001$, and **** $p \leq 0.0001$.

697
Figure 5: Sperm transit through cervical mucus is reduced after desialylation. A) Schematic
depicting experimental setup and mucus-sperm interaction model. Sperm were incubated with seminal
plasma, then inserted into a capillary tube containing sialidase at the entrance and ovulatory cervical
mucin. B) Total sperm were quantified at each distance in the capillary tube containing ovulatory
cervical mucus, and either 0.5 4-MU hydrolysis/min/uL GvNanH2 sialidase or media at the entrance.
Statistics represent an unpaired test comparison of the slopes of the lines derived from the four points.
Data represent three biological replicates with either two or three technical replicates depending on
donor volume.
706

707
Supplementary Figure 1: Production of GvNanH2 and characterization of Prevotella enzymes. A)
SDS PAGE of purified GvNanH2. B) Activity determination of different concentrations of GvNanH2 with
4-MU hydrolysis assay. C) Activity determination of Prevotella enzymes stocks of with 4-MU hydrolysis
assay. N= 3. D) Activity comparison to 5 units of commercially available AUS sialidase. N= 5 E)
Inhibition of GvNanH2 by DANA, a sialidase inhibitor. 95% confidence intervals are represented by the
pink dashed line. N = 3. Error bars represent standard error of the mean.
714

Supplementary Figure 2: LC-MS/MS- based sequence coverage of in-gel digest confirms sequence

of GvNanH2. The 75 Kda band from the SDS PAGE gel in Fig S1A was cut and digested with trypsin,

and peptides were subjected to LC-MS/MS analysis to confirm the identity of the protein.

Supplementary Figure 3: Flow cytometry gates for sperm. Sperm were fixed in formalin, stained with

MAL-II, and analyzed by flow cytometry. Gates were selected to select sperm and remove doublets

based on forward and side scatter.

Supplementary Figure 4: Zeta potential method development. A) Diluent optimization of reducing variability within technical replicates. B) Zeta potential and sperm culture conditions. C) Comparison of AUS (green) and NanH2 (purple). D) Biological replicates shown individually to show technical replicates within each sample, and highlight donor to donor variability. 2000ng/mL corresponds with 0.88 4-MU hydrolysis/min/uL.

734
Supplementary Figure 5: Human Contraception Antibody (HCA)-mediated complement
immobilization of sperm is enhanced in the presence of sialidase. AUS sialidase at 20 units was
incubated with sperm for 1 hour, then complement was added and incubated for 1 hour. Motility of
these samples was quantified by CASA software. Each point represents the average of biological
replicates and error bars represent standard deviations. A mixed effects analysis revealed that there
was a statistically significant effect on motility due to the sialidase treatment and antibody dose
(p<0.001), and post-hoc Holms-Sidak testing revealed a difference in the sialidase treatment conditions
in the absence of antibody. N = 3 per biological replicate. Hi=heat inactivated.

743
744 **Supplementary Figure 6: Effect of seminal plasma on cervical mucus transit of sperm.** A cervical
745 mucus transit assay was performed with one cervical mucus sample containing sialidase enzyme in the
746 presence and absence of seminal plasma. The bubble size represents total sperm count at each point in
747 the capillary tube, and the coloring represents the percent motility. Each point represents a technical
748 replicate.
749

750
Supplementary Figure 7: Method development for cervical mucus penetration To optimize the
cervical mucus transit assay, several variants of the assay were performed with a single cervical mucus
sample. In A) no sialidase was added as a control condition. In B), sialidase was added to sperm samples
only. In C), sialidase was per-mixed with cervical mucus. Each point represents a technical replicate.
Data in the top row are stacked graphs representing both sperm counts and contribution of each
motility category divided by experimental condition, and the bottom row is split by motility category to
compare different experimental conditions.
758

Supplementary Figure 8: Comparison of human cervical mucin (HCM) and bovine submaxillary

mucin (BSM) in sperm transit assay. Cervical mucus transit assay was performed with 1.5% bovine

submaxillary mucin in PBS, to compare sperm transit with and without seminal plasma as part of assay

development. Each point represents the average of three technical replicates, and error bars represent

standard deviations. In the absence of seminal plasma, the overall sperm transit pattern was retained,

but in the presence of seminal plasma, the pattern of sperm transit deviated between the bovine and

human mucus.

Sperm in MHM media

Sialidase concentration
(micromolar 4-MU hydroylysis/min/million sperm)

**In Human Cervical Mucus
20 million sperm/mL**

Sialidase concentration
(micromolar 4-MU hydroylysis/min/uL)

Supplementary figure 9: Spontaneous agglutination in the presence of sialidases. Motile sperm were exposed to PtNanH2 sialidase (0.88 4MU uM hydrolysis per min) for 1 hour and imaged for the presence of agglutinates. Some small clusters of sperm sticking together were observed. In cervical mucus, large agglutinates were sometimes observed in the presence of sialidase.

In 10% human serum
35 million sperm/mL

0.88 micromolar 4-MU hydroylsis/min/uL

776
777 **Supplementary Figure 10:** Representative images of sperm agglutination after treatment with NanH2
778 sialidase and complement for 1 hour. Sperm were stained with trypan blue to confirm cell death. Dead
779 sperm were observed, along with motile live sperm (sometimes seen as tracks in the images due to
780 movement).
781

References

Abou Chacra L, Fenollar F, Diop K. 2022. Bacterial Vaginosis: What Do We Currently Know?

Front Cell Infect Microbiol. 11. doi:10.3389/fcimb.2021.672429. [accessed 2025 Jan 21].

https://www.frontiersin.org/journals/cellular-and-infection-

microbiology/articles/10.3389/fcimb.2021.672429/full.

Abou Chacra L, Ly C, Hammoud A, Iwaza R, Mediannikov O, Bretelle F, Fenollar F. 2023.

Relationship between Bacterial Vaginosis and Sexually Transmitted Infections: Coincidence,

Consequence or Co-Transmission? Microorganisms. 11(10):2470.

doi:10.3390/microorganisms11102470.

Abril-Parreño L, Krogenæs A, Druart X, Saldova R, Cormican P, Meade K, Fair S. 2022. Ewe

breed differences in cervical mucus biochemical and molecular composition: the role of

sialylated cervical mucins. Animal Reproduction Science. 247:107103.

doi:10.1016/j.anireprosci.2022.107103.

Abril-Parreño L, Wilkinson H, Krogenæs A, Morgan J, Gallagher ME, Reid C, Druart X, Fair S,

Saldova R. 2022. Identification and characterization of *O* -linked glycans in cervical mucus as

biomarkers of sperm transport: A novel sheep model. Glycobiology. 32(1):23–35.

doi:10.1093/glycob/cwab085.

Agarwal K, Choudhury B, Robinson LS, Morrill SR, Bouchibiti Y, Chilin-Fuentes D, Rosenthal

SB, Fisch KM, Peipert JF, Lebrilla CB, et al. 2023. Resident microbes shape the vaginal

epithelial glycan landscape. Science Translational Medicine. 15(724):eabp9599.

doi:10.1126/scitranslmed.abp9599.

Agarwal K, Lewis AL. 2021. Vaginal sialoglycan foraging by *Gardnerella vaginalis*: mucus

barriers as a meal for unwelcome guests? Glycobiology. 31(6):667–680.

doi:10.1093/glycob/cwab024.

Álvarez-Rodríguez M, Martinez CA, Wright D, Rodríguez-Martinez H. 2020. The role of semen

and seminal plasma in inducing large-scale genomic changes in the female porcine peri-

ovulatory tract. Sci Rep. 10(1):5061. doi:10.1038/s41598-020-60810-z.

Anderson DJ, Abbott AF, Jack RM. 1993. The role of complement component C3b and its

receptors in sperm-oocyte interaction. Proceedings of the National Academy of Sciences.

90(21):10051–10055. doi:10.1073/pnas.90.21.10051.

Anderson DJ, Politch JA, Cone RA, Zeitlin L, Lai SK, Santangelo PJ, Moench TR, Whaley KJ.

2020. Engineering monoclonal antibody-based contraception and multipurpose prevention

technologies†. Biol Reprod. 103(2):275–285. doi:10.1093/biolre/ioaa096.

Anderson DJ, Politch JA, Zeitlin L, Hiatt A, Kadasia K, Mayer KH, Ruprecht RM, Villinger F,

Whaley KJ. 2017. Systemic and topical use of monoclonal antibodies to prevent the sexual

transmission of HIV. AIDS. 31(11):1505–1517. doi:10.1097/QAD.0000000000001521.

Andrade-Rocha FT. 2009. Colonization of *Gardnerella vaginalis* in Semen of Infertile Men:

Prevalence, Influence on Sperm Characteristics, Relationship with Leukocyte Concentration and

Clinical Significance. Gynecologic and Obstetric Investigation. 68(2):134–136.

doi:10.1159/000228583.

Armstrong E, Kaul R. 2021. Beyond bacterial vaginosis: vaginal lactobacilli and HIV risk.

Microbiome. 9(1):239. doi:10.1186/s40168-021-01183-x.

Aroutcheva A, Ling Z, Faro S. Prevotella bivia as a source of lipopolysaccharide in the vagina.

Anaerobe. 2008 Nov;14(5):256-60. doi: 10.1016/j.anaerobe.2008.08.002. Epub 2008 Sep 20.

PMID: 18849004; PMCID: PMC2651005.

Baldeon-Vaca G, Marathe JG, Politch JA, Mausser E, Pudney J, Doud J, Nador E, Zeitlin L,

Pauly M, Moench TR, et al. 2021. Production and characterization of a human antisperm

monoclonal antibody against CD52g for topical contraception in women. EBioMedicine.

69:103478. doi:10.1016/j.ebiom.2021.103478.

Batra V, Dagar K, Nayak S, Kumaresan A, Kumar R, Datta TK. 2020. A Higher Abundance of

O-Linked Glycans Confers a Selective Advantage to High Fertile Buffalo Spermatozoa for

Immune-Evasion From Neutrophils. Front Immunol. 11:1928. doi:10.3389/fimmu.2020.01928.

Bhagwat S., Asadi, L. McCarthy R., Ferreira J., Li P., Li E., Spivak S., Gaydon A., Reddy V.,

Armstrong C., Morrill S., Zhou H., Lewis AL., Lewis WG., Santi CM. 2025. Bacterial vaginosis

toxins impair sperm capacitation and fertilization. *Human Reproduction*.

deaf132, https://doi.org/10.1093/humrep/deaf132

Blaum BS, Hannan JP, Herbert AP, Kavanagh D, Uhrín D, Stehle T. 2015. Structural basis for

sialic acid–mediated self-recognition by complement factor H. Nat Chem Biol. 11(1):77–82.

doi:10.1038/nchembio.1696.

Cauci S, Culhane JF. 2011. High sialidase levels increase preterm birth risk among women who

are bacterial vaginosis-positive in early gestation. Am J Obstet Gynecol. 204(2):142.e1–9.

doi:10.1016/j.ajog.2010.08.061.

Cauci S, Culhane JF, Di Santolo M, McCollum K. 2008. Among pregnant women with bacterial

vaginosis, the hydrolytic enzymes sialidase and prolidase are positively associated with

interleukin-1beta. Am J Obstet Gynecol. 198(1):132.e1–7. doi:10.1016/j.ajog.2007.05.035.

Chan D, Bennett PR, Lee YS, Kundu S, Teoh TG, Adan M, Ahmed S, Brown RG, David AL,

Lewis HV, et al. 2022. Microbial-driven preterm labour involves crosstalk between the innate

and adaptive immune response. Nat Commun. 13(1):975. doi:10.1038/s41467-022-28620-1.

Cheu RK, Mohammadi A, Schifanella L, Broedlow C, Driscoll CB, Miller CJ, Reeves RK, Yudin

MH, Hensley-McBain T, Kaul R, et al. 2022. Altered Innate Immunity and Damaged Epithelial

Integrity in Vaginal Microbial Dysbiosis. Front Reprod Health. 4. doi:10.3389/frph.2022.876729.

Dalia AB, Weiser JN. 2011. Minimization of Bacterial Size Allows for Complement Evasion and

Is Overcome by the Agglutinating Effect of Antibody. Cell Host & Microbe. 10(5):486–496.

doi:10.1016/j.chom.2011.09.009.

Denney JM, Culhane JF. 2009. Bacterial vaginosis: a problematic infection from both a perinatal

and neonatal perspective. Semin Fetal Neonatal Med. 14(4):200–203.

doi:10.1016/j.siny.2009.01.008.

Denny KJ, Woodruff TM, Taylor SM, Callaway LK. 2012. Complement in Pregnancy: A Delicate

Balance. American Journal of Reproductive Immunology. 69(1):3–11.

doi:https://doi.org/10.1111/aji.12000.

Domingues MM, Inácio RG, Raimundo JM, Martins M, Castanho MARB, Santos NC. 2012.

Biophysical characterization of polymyxin b interaction with LPS aggregates and membrane

model systems. Biopolymers. 98(4):338–344. doi:https://doi.org/10.1002/bip.22095.

Dingens AS, Fairfortune TS, Reed S, Mitchell C. 2016. Bacterial vaginosis and adverse

outcomes among full-term infants: a cohort study. BMC Pregnancy Childbirth. 16(1):278.

doi:10.1186/s12884-016-1073-y.

Donnellan EM, Cormican P, Reid C, Duggan G, Stiavnicka M, Meade KG, Fair S. 2023. The

transcriptomic response of bovine uterine tissue is altered in response to sperm from high- and

low-fertility bulls†. Biology of Reproduction. 108(6):912–921. doi:10.1093/biolre/ioad031.

Dorin JR, Barratt CLR. 2014. Importance of β-defensins in sperm function. Molecular Human

Reproduction. 20(9):821–826. doi:https://doi.org/10.1093/molehr/gau050.

Elovitz MA, Gajer P, Riis V, Brown AG, Humphrys MS, Holm JB, Ravel J. 2019. Cervicovaginal

microbiota and local immune response modulate the risk of spontaneous preterm delivery. Nat

Commun. 10(1):1305. doi:10.1038/s41467-019-09285-9.

Feng Y, Wang L, Wu Y-L, Liu H-H, Ma F. 2016. [Roles of sialic acids in sperm maturation and

capacitation and sperm-egg recognition]. Zhonghua Nan Ke Xue. 22(10):944–948.

Fernandez-Fuertes B, Blanco-Fernandez A, Reid CJ, Meade KG, Fair S, Lonergan P. 2018 Jun

1. Removal of sialic acid from bull sperm decreases motility and mucus penetration ability but

increases zona pellucida binding and polyspermic penetration in vitro. doi:10.1530/REP-17-

0429. [accessed 2025 Jan 21]. https://rep.bioscientifica.com/view/journals/rep/155/6/REP-17-

0429.xml.

Fliniaux I, Marchand G, Molinaro C, Decloquement M, Martoriati A, Marin M, Bodart J-F,

Harduin-Lepers A, Cailliau K. 2022. Diversity of sialic acids and sialoglycoproteins in gametes

and at fertilization. Front Cell Dev Biol. 10. doi:10.3389/fcell.2022.982931. [accessed 2025 Jan

21]. https://www.frontiersin.org/journals/cell-and-developmental-

biology/articles/10.3389/fcell.2022.982931/full.

Froman DP, Thursam KA. 1994. Desialylation of the rooster sperm's glycocalyx decreases

sperm sequestration following intravaginal insemination of the hen. Biol Reprod. 50(5):1094–

1099. doi:10.1095/biolreprod50.5.1094.

Green, E.S., Arck, P.C. 2020. Pathogenesis of preterm birth: bidirectional inflammation in

mother and fetus. Semin Immunopathol 42, 413–429. https://doi.org/10.1007/s00281-020-

00807-y

D. Ghaderi, S.A. Springer, F. Ma, M. Cohen, P. Secrest, R.E. Taylor, A. Varki,

& P. Gagneux. 2011. Sexual selection by female immunity against paternal antigens can fix loss

of function alleles, Proc. Natl. Acad. Sci. U.S.A. 108 (43) 17743-17748,

https://doi.org/10.1073/pnas.1102302108.

Gudnadottir U, Debelius JW, Du J, Hugerth LW, Danielsson H, Schuppe-Koistinen I, Fransson

E, Brusselaers N. 2022. The vaginal microbiome and the risk of preterm birth: a systematic

review and network meta-analysis. Sci Rep. 12(1):7926. doi:10.1038/s41598-022-12007-9.

Hadhoum S, Subtil D, Labreuche J, Couvreur E, Brabant G, Dessein R, Le Guern R.

Reassessing the association between bacterial vaginosis and preterm birth: A systematic review

and meta-analysis. J Gynecol Obstet Hum Reprod. 2025 Jan;54(1):102871. doi:

10.1016/j.jogoh.2024.102871. Epub 2024 Oct 21. PMID: 39442804.

Hardy L, Jespers V, Van den Bulck M, Buyze J, Mwambarangwe L, Musengamana V,

Vaneechoutte M, Crucitti T. 2017. The presence of the putative *Gardnerella vaginalis* sialidase

A gene in vaginal specimens is associated with bacterial vaginosis biofilm. PLoS One.

12(2):e0172522. doi:10.1371/journal.pone.0172522.

He J, Ma M, Xu Z, Guo J, Chen H, Yang X, Chen P, Liu G. 2024. Association between semen

microbiome disorder and sperm DNA damage. Microbiology Spectrum. 12(8):e00759-24.

doi:10.1128/spectrum.00759-24.

Howe L, Wiggins R, Soothill PW, Millar MR, Horner PJ, Corfield AP. 1999. Mucinase and

sialidase activity of the vaginal microflora: implications for the pathogenesis of preterm labour.

Int J STD AIDS. 10(7):442–447. doi:10.1258/0956462991914438.

Hull MGR, Mcleod FN, Joyce DN, Ray BD, Mcdermott A. 1984. HUMAN IN-VITRO

FERTILISATION, IN-VIVO SPERM PENETRATION OF CERVICAL MUCUS, AND

UNEXPLAINED INFERTILITY. The Lancet. 324(8397):245–246.

doi:https://doi.org/10.1016/s0140-6736(84)90296-4.

Katila T. 2001. Sperm–uterine interactions: a review,. Animal Reproduction Science,. 68:267–

272

Kenyon C, Colebunders R, Crucitti T. 2013. The global epidemiology of bacterial vaginosis: a

systematic review. American Journal of Obstetrics and Gynecology. 209(6):505–523.

doi:10.1016/j.ajog.2013.05.006.

Khosravi F, Michel V, Galuska CE, Bhushan S, Christian P, Schuppe H-C, Pilatz A, Galuska

SP, Meinhardt A. 2016. Desialylation of Spermatozoa and Epithelial Cell Glycocalyx Is a

Consequence of Bacterial Infection of the Epididymis. J Biol Chem. 291(34):17717–17726.

doi:10.1074/jbc.M116.718072.

Kindschuh WF, Austin GI, Meydan Y, Park H, Urban JA, Watters E, Pollak S, Saade GR, Chung

J, Mercer BM, et al. 2024. Early prediction of preeclampsia using the first trimester vaginal

microbiome. :2024.12.01.626267. doi:10.1101/2024.12.01.626267. [accessed 2025 Jan 21].

https://www.biorxiv.org/content/10.1101/2024.12.01.626267v1.

Li Y, Zhai Y, Fu B, He Y, Feng Y, Ma F, Lu H. 2024. A comprehensive N-glycome map of

porcine sperm membrane before and after capacitation. Carbohydrate Polymers. 335:122084.

doi:10.1016/j.carbpol.2024.122084.

Luo M, Su T, Cheng Q, Zhang X, Cai F, Yin Z, Li F, Yang H, Liu F, Zhang Y. 2023. GlycoTCFM:

Glycoproteomics Based on Two Complementary Fragmentation Methods Reveals Distinctive O-

Glycosylation in Human Sperm and Seminal Plasma. J Proteome Res. 22(12):3833–3842.

doi:10.1021/acs.jproteome.3c00489.

Lynch, Anne M. MD, MSPH; Gibbs, Ronald S. MD; Murphy, James R. PhD; Giclas, Patricia C.

PhD; Salmon, Jane E. MD; Holers, V. Michael MD. Early Elevations of the Complement

Activation Fragment C3a and Adverse Pregnancy Outcomes. Obstetrics & Gynecology 117(1):p

75-83, January 2011. | DOI: 10.1097/AOG.0b013e3181fc3afa

Lynch AM, Murphy JR, Byers T, Gibbs RS, Neville MC, Giclas PC, Salmon JE, Holers VM.

2008. Alternative complement pathway activation fragment Bb in early pregnancy as a predictor

of preeclampsia. American Journal of Obstetrics and Gynecology. 198(4):385.e1–385.e9.

doi:https://doi.org/10.1016/j.ajog.2007.10.793.

Ma X, Pan Q, Feng Y, Choudhury BP, Ma Q, Gagneux P, Ma F. 2016. Sialylation Facilitates the

Maturation of Mammalian Sperm and Affects Its Survival in Female Uterus. Biol Reprod.

94(6):123. doi:10.1095/biolreprod.115.137810.

Machado SA, Kadirvel G, Daigneault BW, Korneli C, Miller P, Bovin N, Miller DJ. 2014. LewisX-

Containing Glycans on the Porcine Oviductal Epithelium Contribute to Formation of the Sperm

Reservoir1. Biology of Reproduction. 91(6):140, 1–9. doi:10.1095/biolreprod.114.119503.

Mahdavinezhad F, Gharaei R, Farmani AR, Hashemi F, Kouhestani M, Amidi F. 2022. The

Potential Relationship Between Different Human Female Reproductive Disorders and Sperm

Quality in Female Genital Tract. Reprod Sci. 29(3):695–710. doi:10.1007/s43032-021-00520-7.

Mania-Pramanik J, Kerkar SC, Salvi VS. 2009. Bacterial vaginosis: a cause of infertility? Int J

STD AIDS. 20(11):778–781. doi:10.1258/ijsa.2009.009193.

Mausser E, Nador E, Politch JA, Pauly MR, Marathe JG, Moench TR, Zeitlin L, Whaley KJ,

Anderson DJ. 2023. LALAPG variant of the Human Contraception Antibody (HCA) reduces Fc-

mediated effector functions while maintaining sperm agglutination activity. PLoS One.

18(3):e0282147. doi:10.1371/journal.pone.0282147.

Menevse E, Dursunoglu D, Cetin N, Korucu EN, Erbayram FZ. 2022. Evaluation of sialic acid,

malondialdehyde and glutathione levels in infertile male. Rev Int Androl. 20(4):266–273.

doi:10.1016/j.androl.2021.05.002.

Mohanty T, Doke PP, Khuroo SR. 2023. Effect of bacterial vaginosis on preterm birth: a meta-

analysis. Arch Gynecol Obstet. 308(4):1247–1255. doi:10.1007/s00404-022-06817-5.

Moncla BJ, Chappell CA, Debo BM, Meyn LA. 2016. The Effects of Hormones and Vaginal

Microflora on the Glycome of the Female Genital Tract: Cervical-Vaginal Fluid. PLOS ONE.

11(7):e0158687. doi:10.1371/journal.pone.0158687.

Morrill S, Gilbert NM, Lewis AL. 2020. *Gardnerella vaginalis* as a Cause of Bacterial Vaginosis:

Appraisal of the Evidence From in vivo Models. Front Cell Infect Microbiol. 10:168.

doi:10.3389/fcimb.2020.00168.

Norwitz ER. 2006. Defective implantation and placentation: laying the blueprint for pregnancy

complications. Reproductive BioMedicine Online. 13(4):591–599. doi:10.1016/S1472-

6483(10)60649-9.

Norwitz ER, Schust DJ, Fisher SJ. 2001. Implantation and the Survival of Early Pregnancy. New

England Journal of Medicine. 345(19):1400–1408. doi:10.1056/NEJMra000763.

van Oostrum N, De Sutter P, Meys J, Verstraelen H. 2013. Risks associated with bacterial

vaginosis in infertility patients: a systematic review and meta-analysis. Hum Reprod.

28(7):1809–1815. doi:10.1093/humrep/det096.

Osadchiy V, Belarmino A, Kianian R, Sigalos JT, Ancira JS, Kanie T, Mangum SF, Tipton CD,

Hsieh T-CM, Mills JN, et al. 2024. Semen microbiota are dramatically altered in men with

abnormal sperm parameters. Sci Rep. 14(1):1068. doi:10.1038/s41598-024-51686-4.

Peebles K, Velloza J, Balkus JE, McClelland RS, Barnabas RV. 2019. High Global Burden and

Costs of Bacterial Vaginosis: A Systematic Review and Meta-Analysis. Sexually Transmitted

Diseases. 46(5):304. doi:10.1097/OLQ.0000000000000972.

Pelayo P, Hussain FA, Werlang CA, Wu CM, Woolston BM, Xiang CM, Rutt L, France MT,

Ravel J, Ribbeck K, et al. 2024. Prevotella are major contributors of sialidases in the human

vaginal microbiome. Proc Natl Acad Sci U S A. 121(36):e2400341121.

doi:10.1073/pnas.2400341121.

Petersen BH, Lammel CJ, Stites DP, Brooks GF. 1980. Human seminal plasma inhibition of

complement. The Journal of Laboratory and Clinical Medicine. 96(4):582–591.

doi:10.5555/uri:pii:0022214380900621.

Price RJ, Boettcher B. 1979. The Presence of Complement in Human Cervical Mucus and its

Possible Relevance to Infertility in Women with Complement-Dependent Sperm-Immobilizing

Antibodies. Fertility and Sterility. 32(1):61–66. doi:10.1016/S0015-0282(16)44117-8.

Ralph SG, Rutherford AJ, Wilson JD. 1999. Influence of bacterial vaginosis on conception and

miscarriage in the first trimester: cohort study. BMJ. 319(7204):220–223.

Ravel J, Gajer P, Abdo Z, Schneider GM, Koenig SSK, McCulle SL, Karlebach S, Gorle R,

Russell J, Tacket CO, et al. 2011. Vaginal microbiome of reproductive-age women. Proc Natl

Acad Sci U S A. 108 Suppl 1(Suppl 1):4680–4687. doi:10.1073/pnas.1002611107.

Ravel J, Moreno I, Simón C. 2021. Bacterial vaginosis and its association with infertility,

endometritis, and pelvic inflammatory disease. American Journal of Obstetrics and Gynecology.

224(3):251–257. doi:https://doi.org/10.1016/j.ajog.2020.10.019.

1019

1020   Regal JF, Gilbert JS, Burwick RM. 2015. The complement system and adverse pregnancy

1021   outcomes. Molecular Immunology. 67(1):56–70.

1022   doi:https://doi.org/10.1016/j.molimm.2015.02.030.

1023

Robinson LS, Schwebke J, Lewis WG, Lewis AL. 2019. Identification and characterization of

NanH2 and NanH3, enzymes responsible for sialidase activity in the vaginal bacterium

Gardnerella vaginalis. J Biol Chem. 294(14):5230–5245. doi:10.1074/jbc.RA118.006221.

Roselletti E, Sabbatini S, Perito S, Mencacci A, Vecchiarelli A, Monari C. 2020. Apoptosis of

vaginal epithelial cells in clinical samples from women with diagnosed bacterial vaginosis. Sci

Rep. 10(1):1978. doi:10.1038/s41598-020-58862-2.

Riley-Vargas RC, Lanzendorf S, Atkinson JP. 2005. Targeted and restricted complement

activation on acrosome-reacted spermatozoa. Journal of Clinical Investigation. 115(5):1241–

1249. doi:https://doi.org/10.1172/jci23213.

1033
Savicheva AM, Krysanova AA, Budilovskaya OV, Spasibova EV, Khusnutdinova TA, Shalepo

KV, Beliaeva NR, Safarian GK, Sapozhnikov KV, Tapilskaya NI, et al. 2023. Vaginal Microbiota

Molecular Profiling in Women with Bacterial Vaginosis: A Novel Diagnostic Tool. International

Journal of Molecular Sciences. 24(21):15880. doi:10.3390/ijms242115880.

Schjenken JE, Sharkey DJ, Green ES, Chan HY, Matias RA, Moldenhauer LM, Robertson SA.

2021. Sperm modulate uterine immune parameters relevant to embryo implantation and

reproductive success in mice. Commun Biol. 4(1):1–14. doi:10.1038/s42003-021-02038-9.

Segui-Perez C, de Jongh R, Jonkergouw RLW, Pelayo P, Balskus EP, Zomer A, Strijbis K.

2024. *Prevotella timonensis* degrades the vaginal epithelial glycocalyx through high fucosidase

and sialidase activities. mBio. 15(9):e00691-24. doi:10.1128/mbio.00691-24.

Tantengco OAG, Menon R. 2022. Breaking Down the Barrier: The Role of Cervical Infection and

Inflammation in Preterm Birth. Front Glob Womens Health. 2. doi:10.3389/fgwh.2021.777643.

[accessed 2025 Jan 21]. https://www.frontiersin.org/journals/global-womens-

health/articles/10.3389/fgwh.2021.777643/full.

Tecle E, Gagneux P. 2015. Sugar-coated sperm: Unraveling the functions of the mammalian

sperm glycocalyx. Mol Reprod Dev. 82(9):635–650. doi:10.1002/mrd.22500.

Diao R, Fok KL, Chen H, Yu MK, Duan Y, Chung CM, Li Z, Wu H, Li Z, Zhang H, et al. 2014.

Deficient human -defensin 1 underlies male infertility associated with poor sperm motility and

genital tract infection. Science Translational Medicine. 6(249):249ra108–249ra108.

doi:https://doi.org/10.1126/scitranslmed.3009071.

1054
Toshimori K, Araki S, Öra C, Eddy EM. 1991. Loss of Sperm Surface Sialic Acid Induces

Phagocytosis: An Assay with a Monoclonal Antibody T21, Which Recognizes a 54K

Sialoglycoprotein. Archives of Andrology. 27(2):79–86. doi:10.3109/01485019108987656.

Tsuji Y, Clausen H, Nudelman E, Kaizu T, Hakomori S, Isojima S. 1988. Human sperm

carbohydrate antigens defined by an antisperm human monoclonal antibody derived from an

infertile woman bearing antisperm antibodies in her serum. Journal of Experimental Medicine.

168(1):343–356. doi:10.1084/jem.168.1.343.

Ugwumadu AHN. 2002. Bacterial vaginosis in pregnancy. Current Opinion in Obstetrics and

Gynecology. 14(2):115.

Xin A-J, Cheng L, Diao H, Wang P, Gu Y-H, Wu B, Wu Y-C, Chen G-W, Zhou S-M, Guo S-J, et

al. 2014. Comprehensive profiling of accessible surface glycans of mammalian sperm using a

lectin microarray. Clinical Proteomics. 11(1):10. doi:10.1186/1559-0275-11-10.

Young CD, Tatieng S, Kongmanas K, Fongmoon D, Lomenick B, Yoon AJ, Kiattiburut W,

Compostella F, Faull KF, Suree N, et al. 2019. Sperm can act as vectors for HIV-1 transmission

into vaginal and cervical epithelial cells. Am J Reprod Immunol. 82(1). doi:10.1111/aji.13129.

https://onlinelibrary.wiley.com/doi/10.1111/aji.13129

A

MAL-II lectin sialidase

SNA lectin PSA lectin

sialidase

B
MAL-II biotin
Alexa Fluor 488

MFI

**

**

None   AUS   GvNanH2

C
SNA-Cy5

MFI

*

*

None   AUS   GvNanH2

D
PSA-FITC

MFI

ns

ns

None   AUS   GvNanH2

E

sialidase

F

zeta potential (mV)

***

ns

ns

ns

****

ns

ns

***

Hi 0.002 0.02 0.11   Hi 0.002 0.02 0.11 0.88

AUS   GvNanH2

zeta potential (mV)

ns

ns

ns

ns

****

ns

****

Hi 0.002 0.02 0.11 0.88   Hi 0.002 0.02 0.11 0.88

PtNanH1   PtNanH2

Enzyme (4-MU hydrolysis/min/uL)

A) A. Ureafaciens Sialidase (AUS)

B) Gardnerella sialidase (GvNanH2)

C) Prevotella sialidases (PtNanH1, PtNanH2)

D) sialidase inhibitors

E) LPS inhibitors

A

sperm in 50% seminal plasma | sialidase or media control

human ovulatory cervical mucus

1 cm 2 cm 3 cm 4 cm

sialidase

B

**

● GvNanH2
■ Control