Optimal membrane fluidity modulates biofilm formation in *Escherichia coli*

Yaoqin Hong[1,2,3,*], Jilong Qin[1,2,4], Makrina Totsika[1,2,*]

[1] Centre for Immunology and Infection Control, School of Biomedical Sciences, Queensland
University of Technology, Herston, Queensland 4006, Australia

[2] Max Planck Queensland Centre, Queensland University of Technology, Herston, Queensland 4006,
Australia

[3] Biomedical Sciences and Molecular Biology, College of Medicine and Dentistry, James Cook
University, Douglas, Queensland 4814, Australia

[4] Institute for Molecular Bioscience, University of Queensland, St Lucia, Queensland 4067, Australia

* For correspondence, contact Yaoqin Hong (yaoqin.hong@jcu.edu.au) and Makrina Totsika
(makrina.totsika@qut.edu.au)

ABSTRACT

Changes in membrane composition govern the fitness and survival of bacteria in response to environmental stresses. One critical parameter is membrane fluidity, which dictates overall membrane integrity, adaptability to environmental fluctuations (e.g., temperature changes), and the functionality and interactions between integral and peripheral membrane proteins. In recent years, substantial evidence has shown that membrane fluidity is finely tuned for biofilm formation. While high membrane fluidity promotes robust biofilm formation in Klebsiella pneumoniae, this is challenged by reports from multiple species indicating that biofilms are enriched in saturated fatty acids (i.e., reduced fluidity). We hypothesize that this paradox arises from the complex relationship between lipid composition and biofilm formation. Our findings demonstrate that an optimal balance of lipid fluidity is required for biofilm formation. We propose that the precise fluidity required for biofilm growth, relative to that required for planktonic growth, may differ between species, and is readily adjusted during lifestyle transitions.

RESULTS AND DISCUSSIONS

Microbes in natural environments often exist in communities, called biofilms, where cells are encased in an extracellular polymeric matrix (EPS) that provides protection from external stresses such as detergents and antibiotics. This EPS, is primarily composed of polysaccharides, proteins, lipids, and extracellular DNA, with its composition and spatial organization tightly regulated by molecular signals [1]. Identifying the molecular cues that bacteria use to regulate biofilm formation and dispersal is critical for both combating infections and harnessing beneficial biofilms. The contribution of bacterial cell membrane fluidity, as a modulator of these processes, remains an underexplored area.

The bacterial membrane plays a vital role in supporting essential processes, including energy generation, fitness in different environments and changes in environmental conditions necessitates membrane lipid adjustments for survival and fitness [2]. Changes in membrane lipids serve as a molecular cue in the Envelope Stress Response (ESR) that can influence various pathways involved in biofilm development [3]. A typical bacterial phospholipid contains two fatty acid (FA) chains, and the chemical nature of these FAs dictates the membrane fluidity [2]. Previous studies have demonstrated that biofilms of *Salmonella enterica, Staphylococcus aureus, Listeria monocytogenes,* and *Pseudomonas aeruginosa* contain elevated levels of saturated fatty acids (SFAs) compared to their planktonic counterparts [4]. This suggests that bacterial membrane composition and fluidity may play roles in biofilm formation, although the underlying mechanisms remain unclear [4,5].

Most bacterial species *de novo* synthesise the acyl chains to be integrate into membrane
lipids by the highly conserved type II fatty acid synthesis pathway distinctive from that used by
eukaryotes [2]. As such, the pathway had served as an important target for antimicrobial discovery [6,7],
with the recent progress shown the potential of potentiating susceptibility to otherwise ineffective
antibiotics [8,9]. In *Escherichia coli*, the major unsaturated fatty acids (UFAs) are palmitoleate (C16:1)
and *cis*-vaccenate (C18:1). Bacterial phospholipids exhibit chemical asymmetry, with SFAs typically
occupying the *sn*1 position and unsaturated fatty acids (UFAs) occupying the *sn*2 position of the
glycerol backbone owed to the substrate preference of acyltransferases involved in phosphatidic
acid synthesis [2,10]. The UFA/SFA ratio that affects membrane fluidity is predominantly determined by
the diversity of FAs at the *sn*2 position [2,10]. The only exception to this substrate selectivity at position
*sn*1 is the integration of *cis*-vaccenate, which leads to higher membrane fluidity that is imperative for
bacterial fitness under certain conditions, such as low temperature growth [11]. As such, this
phospholipid asymmetry plays a pivotal role in modulating membrane fluidity, which, in turn, may
influences biofilm formation. In *E. coli,* UFA synthesis is mediated by ketoacyl-ACP synthase I
(FabB), while the conversion of palmitoleate to *cis*-vaccenate requires ketoacyl-ACP synthase II
(FabF) [12,13]. Thus, FabB and, to a lesser extent, FabF, are key players governing membrane fluidity.

The *E. coli* K-12 strain AR3110 exhibits a distinct macrocolony architecture when grown on
agar surfaces, characterized by a radial, rugose, three-dimensional structure with delaminated
buckling and skin-like elasticity [14]. This strain also forms a robust pellicle highly resistant to
mechanical disturbance at the liquid-air interface (Fig. 1B). Remarkably, both the rugose colony
architecture and pellicle formation disappeared in the Δ*fabF* mutant, suggesting that reduced
membrane fluidity underlies these phenotypic changes (Fig. 1A and 1B).

Aside from the SFA/UFA ratio, membrane fluidity is also influenced by the profile of
phospholipid headgroups. In this work, rather than using standard FA profiling, we utilised DPH
fluorescence anisotropy, a common method for measuring membrane fluidity. A reduced anisotropy
correlates with higher membrane fluidity, while higher anisotropy indicates a more ordered and less
fluid membrane. The measured anisotropy in the Δ*fabF* mutant was significantly lower compared to
the wild-type AR3110 (Fig. 1C), which supported the hypothesis that membrane fluidity is crucial for
biofilm formation in *E. coli* (Fig. 1C).

Cerulenin inhibits both FabF and FabB, enzymes crucial for the synthesis of UFAs, thereby
reducing membrane fluidity [15]. To investigate the role of membrane fluidity in biofilm formation,
AR3110 were treated with subinhibitory concentrations of cerulenin in both rugose and pellicle
biofilm models (Supplemental material, Fig. S1). This treatment reduced membrane fluidity to levels
similar those observed in the Δ*fabF* mutant (Fig. 1C). The macrocolony of AR3110 typically exhibits
a skin-like consistency, which prevents full resuspension of the biomass in solution (Supplemental

material, Fig. S2). However, both the Δ*fabF* mutant and cerulenin-treated wild-type macrocolonies

acquired a buttery consistency, allowing for easy resuspension (Supplemental material, Fig. S2).

The data shown in Fig. 1A and 1B demonstrated the absence of rugose macrocolony architecture

and pellicle formation in both the Δ*fabF* mutant and cerulenin-treated wild-type AR3110, suggesting

that reduced membrane fluidity hinders the development of stable biofilm structures.

The ectopic expression of FabB counteracted the fluidity reduction of AR3110 caused by

cerulenin treatment (Fig. 1F), thereby restoring pellicle formation in cerulenin-treated AR3110 (Fig.

1D). This observation suggests that a reduction in membrane fluidity antagonises biofilm

development. While this manuscript was being prepared, a similar observation was reported in

Klebsiella pneumoniae [16]. FabR negatively regulates the expression of FabA and FabB in known

Enterobacteriaceae species, including *K. pneumoniae*. Consequently, the activity of UFA synthesis

in the *K. pneumoniae* Δ*fabR* mutant is elevated [16], leading to higher membrane fluidity compared to

the WT during biofilm growth. In agreement to our observation, *K. pneumoniae* Δ*fabR* mutant was

found to produce robust biofilm capable of withstanding mechanical disruptions [16]. Dramé *et al.* [16]

and this work collectively suggest that high membrane fluidity is beneficial for the robust

development of microbial biofilms.

An intriguing aspect of biofilm lipid composition is the observed higher SFA content in

biofilms compared to planktonic cells in several species, including *S. enterica* [4]. This observation

seems to contradict the idea that increased membrane fluidity is essential for biofilm formation.

However, we hypothesize that this paradox arises from the complex relationship between lipid

composition and biofilm formation. While membrane fluidity is crucial, an overabundance of UFAs

may lead to excessive fluidity, compromising biofilm robustness.

In this work, we ectopically overexpressed FabB to suppress the biofilm-inhibitory activity of

cerulenin (Fig. 1D). Strikingly, in the absence of cerulenin treatment, AR3110 cells ectopically

expressing FabB exhibited a significantly more fluid membrane (Fig. 1F), and a pellicle was no

longer formed in liquid culture (Fig. 1E), supporting our hypothesis that an optimal balance of lipid

fluidity, instead of simply high fluidity, is critical for biofilm formation.

To this end, we argue that the precise UFA/SFA ratio optimal for biofilm growth relative to

that required for planktonic growth may vary across species if not at strain levels, and is readily

adjusted to facilitate the transition into biofilm lifestyle [4,5,16]. Our findings highlight the critical role of

optimised membrane fluidity for robust biofilm development, and these insights could pave the way

for future studies exploring lipid composition as a potential target for biofilm control in clinical and

industrial applications.

Acknowledgement

This work is funded by an Early Career Researcher Scheme Grant from the Institute of Health and

Biomedical Innovations, and an Early Career Researcher Scheme Grant from the Faculty of Health,

Queensland University of Technology (Australia) to YH. YH and MT were supported by the Max

Planck Queensland Centre on the Materials Science of Extracellular Matrices. The Ian Potter

Foundation sponsored the CLARIOStar high-performance microplate reader (BMG, Australia). The

funders had no role in study design, data collection and analysis, decision to publish, or preparation

of the manuscript. The AR3110 strain used in this work was generously shared by Dr Cécile Bidan

(Max Planck Institute of Colloids and Interfaces, Germany).

Author contributions

YH conceptualised the project, designed and conducted the experiments, and contributed to data

collection and analysis; JQ contributed to experimental resources; YH and MT supervised the study

and obtained the funding. YH wrote the manuscript, and all authors edited the manuscript.

Competing interests

MT is an employee of the GSK group of companies. All remaining authors declare no competing

interests. This research was conducted in the absence of any commercial or financial relationships

that could be constructed as a potential conflict of interest.

Data availability

All data generated or analyzed in this study were included in this article.

References

1.   Sauer, K., Stoodley, P., Goeres, D.M., Hall-Stoodley, L., Burmølle, M., Stewart, P.S.,
and Bjarnsholt, T. (2022). The biofilm life cycle: expanding the conceptual model of
biofilm formation. Nature Reviews Microbiology *20*, 608-620. 10.1038/s41579-022-
00767-0.
2.   Cronan, J.E., Jr., and Rock, C.O. (2008). Biosynthesis of Membrane Lipids. EcoSal
Plus *3*. 10.1128/ecosalplus.3.6.4.
3.   Mitchell, A.M., and Silhavy, T.J. (2019). Envelope stress responses: balancing
damage repair and toxicity. Nature Reviews Microbiology *17*, 417-428.
10.1038/s41579-019-0199-0.

4. Dubois-Brissonnet, F., Trotier, E., and Briandet, R. (2016). The Biofilm Lifestyle
Involves an Increase in Bacterial Membrane Saturated Fatty Acids. Frontiers in
Microbiology *7*, 1673. 10.3389/fmicb.2016.01673.
5. Nepper Julia, F., Lin Yin, C., and Weibel Douglas, B. (2019). Rcs Phosphorelay
Activation in Cardiolipin-Deficient Escherichia coli Reduces Biofilm Formation.
Journal of Bacteriology *201*, 10.1128/jb.00804-00818. 10.1128/jb.00804-18.
6. Campbell, J.W., and Cronan Jr, J.E. (2001). Bacterial fatty acid biosynthesis: targets
for antibacterial drug discovery. Annual Review of Microbiology *55*, 305-332.
7. Yao, J., and Rock, C.O. (2017). Bacterial fatty acid metabolism in modern antibiotic
discovery. Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of
Lipids *1862*, 1300-1309. https://doi.org/10.1016/j.bbalip.2016.09.014.
8. Hong, Y., Qin, J., Verderosa, A.D., Hawas, S., Zhang, B., Blaskovich, M.A.T., Cronan,
J.E., and Totsika, M. (2022). Loss of β-ketoacyl acyl carrier protein synthase III
activity restores multidrug-resistant Escherichia coli sensitivity to previously
ineffective antibiotics. mSphere. 10.1101/2022.02.18.481121.
9. Guo, Q., Su, J., Liao, Y., Yu, Y., Luo, L., Weng, X., Zhang, W., Hu, Z., Wang, H.,
Beattie Gwyn, A., and Ma, J. (2024). An atypical 3-ketoacyl ACP synthase III
required for acyl homoserine lactone synthesis in Pseudomonas syringae pv.
syringae B728a. Applied and Environmental Microbiology *90*, e02256-02223.
10.1128/aem.02256-23.
10. Marr, A.G., and Ingraham, J.L. (1962). Effect of temperature on the composition of
fatty acids in Escherichia coli. Journal of Bacteriology *84*, 1260-1267.
10.1128/jb.84.6.1260-1267.1962.
11. Goelz, S.E., and Cronan, J.E. (1980). The positional distribution of fatty acids in
Escherichia coli phospholipids is not regulated by sn-glycerol 3-phosphate levels.
Journal of Bacteriology *144*, 462-464. 10.1128/jb.144.1.462-464.1980.
12. Gelmann, E.P., and Cronan, J.E., Jr. (1972). Mutant of Escherichia coli deficient in
the synthesis of cis-vaccenic acid. Journal of Bacteriology *112*, 381-387.
10.1128/jb.112.1.381-387.1972.
13. Garwin, J.L., Klages, A.L., and Cronan, J.E., Jr. (1980). Beta-ketoacyl-acyl carrier
protein synthase II of Escherichia coli. Evidence for function in the thermal regulation
of fatty acid synthesis. J Biol Chem *255*, 3263-3265.
14. Serra, D.O., Richter, A.M., and Hengge, R. (2013). Cellulose as an architectural
element in spatially structured Escherichia coli biofilms. Journal of Bacteriology *195*,
5540-5554. 10.1128/jb.00946-13.
15. Price, A.C., Choi, K.H., Heath, R.J., Li, Z., White, S.W., and Rock, C.O. (2001).
Inhibition of beta-ketoacyl-acyl carrier protein synthases by thiolactomycin and
cerulenin. Journal of Biological Chemistry *276*, 6551-6559. 10.1074/jbc.M007101200.
16. Dramé, I., Rossez, Y., Krzewinski, F., Charbonnel, N., Ollivier-Nakusi, L., Briandet, R.,
Dague, E., Forestier, C., and Balestrino, D. (2024). FabR, a regulator of membrane
lipid homeostasis, is involved in Klebsiella pneumoniae biofilm robustness. mBio *15*,
e01317-01324. 10.1128/mbio.01317-24.

Fig 1. Optimal membrane fluidity governs *E. coli* biofilm rugosity and pellicle formation. (A) The *fabF* null mutation or cerulenin treatment eliminated rugosity in AR3110 macrocolony biofilms grown on YESCA agar at 26°C; (B) The *fabF* null mutation or cerulenin treatment abolished pellicle formation in AR3110 on Muller Hinton broth. Note that identical results were observed on YESCA, LB Lennox, and LBON; (C) DPH fluorescence anisotropy was used to measure membrane fluidity (fluidity is inversely proportional to fluorescence); (D) Ectopic expression of FabB (pSU2718 vector, induced with 250 µM IPTG) restored pellicle

219   formation inhibited by cerulenin treatment (E) Overexpression of FabB blocked pellicle formation in AR3110,
220   but this defect was alleviated by cerulenin treatment. (F) Ectopic expression of FabB significantly increased
221   membrane fluidity. Note that the increased fluidity can be ameliorated with cerulenin treatment. Panels A, B, D
222   and F were representative of at least three independent observations. Data in Panels C and F were obtained
223   from three independent repeats. Error bar is indicative of standard deviation values.