Internet images reveal bumblebee-mimicking hoverflies have

evolved to follow flower colour preferences of their models

James D. J. Gilbert[1][✉], Joseph Craig-Jackson[1], Lesley Morrell[1], Francis Gilbert[2]

[✉]To whom correspondence should be addressed: james.gilbert@hull.ac.uk

[1] University of Hull, Cottingham Road, Hull, UK HU6 7RX

[2] University of Nottingham, UK NG7 2RD

Abstract

Batesian mimics, such as hoverflies, can gain protection by resembling noxious species that

predators avoid, such as bees. Such resemblance can be behavioural as well as morphological.

Bees commonly have a preference for blue flowers, whilst flies generally prefer yellow and

white. We predicted that, driven by selection for enhanced mimicry, bee-mimicking hoverflies

would have switched flower preference from yellow to blue, to follow the flower choices of their

models. We gathered internet photographs of flower-visiting bees and wasps, their hoverfly

mimics, and non-mimicking hoverflies, and compared the colours of the visited flowers. Flowers

visited by bumblebees were "bluer" than those visited by other hymenopterans (honeybees,

wasps and solitary bees). Correspondingly, flowers visited by bumblebee-mimicking hoverflies

were bluer than those visited by non-mimics, and were indistinguishable in blueness from those

visited by bumblebees. There was no such pattern in flower "yellowness", where all flies and

most hymenopterans preferred similar yellowness, and only bumblebees chose less yellow

flowers than the others. Our study demonstrates widespread changes in microhabitat choice

(specifically flower preference) among a diverse group of Batesian mimics, consistent with the

idea of behavioural mimicry. Mildly noxious models such as bumblebees are thought to exert

particularly strong selection for mimetic accuracy, suggesting that habitat-choice mimicry may

be selected as a way of enhancing morphological mimicry.

Introduction

Batesian mimics such as hoverflies (Diptera:Syrphidae) are palatable to predators, but gain protection by resembling noxious model species that predators avoid (Rettenmeyer, 1970), such as bees (Hymenoptera:Apoidea). Mimics often experience strong directional selection to maintain or increase similarity to models (Mappes & Alatalo, 1997; Sherratt, 2002). To do this they mimic cues that predators normally use to recognise noxious prey, which can be subtle (Ruxton, 2009). For example, many species employ behavioural as well as morphological mimicry (Ceccarelli, 2008; Kitamura & Imafuku, 2015).

Bees and flies are both pollinators, and are typically associated with distinctive suites of floral characteristics ("pollination syndromes"), including distinct floral colours (Faegri & van der Pijl, 1979). Bees have long been known to prefer blue flowers (Giurfa et al., 1995; Gumbert, 2000; Lubbock, 1882; Raine et al., 2006; Raine & Chittka, 2007; Robertson, 1928), while flies (Diptera) prefer visiting yellow and/or white flowers (de Buck, 1990; Lunau, 2014).

We suggest that selection on mimetic accuracy may include selection for mimetic flies to mimic the flower choices of their models. A large part of the predation of dipteran species takes place on and around flowers (Dlusski, 1984), and closer mimics are attacked by predators at a higher frequency (F. Gilbert, 2005; Morse, 1979). Association with "fly flowers" could potentially be used by predators as a cue to a mimic's identity. As one potential example, bee-flies (Bombyliidae) are known for their exceptionally close mimicry of bumblebees, and unusually for flies, they also prefer visiting blue and violet flowers (Kastinger & Weber, 2001); so much so, in fact, that they have evolved long tongues with which to exploit them (Szucsich & Krenn, 2000).

Hoverflies are an ideal clade in which to test this hypothesis. The Syrphidae contain several origins of mimicry of various hymenopterans (Leavey et al., 2021), varying from imperfect to astonishingly accurate (Gilbert 2005). While hoverflies commonly resemble their models morphologically, in some species elaborate behavioural mimicry has also been documented (Golding & Ennos, 2005) such as mock stinging, leg waving and wing wagging (Penney et al., 2014). In many cases floral visitation patterns of syrphid mimics follows that of their models, whether seasonally (Howarth & Edmunds, 2000), daily (Howarth et al., 2004) or in second-to-second foraging patterns (Golding & Edmunds, 2000).

Like other flies, syrphids are well known to have a general and presumably ancestral preference
for yellow and white flowers (Hannah et al., 2019; Ilse, 1949; Kugler, 1950; Lunau et al., 2018;
Robertson, 1928; Rodríguez-Gasol et al., 2019), and only rarely visit blue flowers (Ssymank,
1989, 2001). We predicted that hoverflies mimicking hymenopteran species that prefer blue
flowers, such as bees, should have switched their own preferences towards bluer flowers than
non-mimics, or related species that mimic other models. In addition, behavioural mimicry has
mostly evolved among larger and morphologically closer mimics (Penney et al., 2014), lending
credence to the established idea that larger body size selects for more faithful mimicry (Penney
et al., 2012). Thus, we also predicted that flower preferences are most likely to have switched
in the largest hoverflies.

For assessing visual traits of organisms, images obtained from the internet have been shown to
provide a good substitute for field data which can be difficult to obtain (Leighton et al., 2016).
This can include studies assessing colour-based traits (Atsumi & Koizumi, 2017; Austen et al.,
2018; M. P. Moore et al., 2019) despite variation due to camera- and environment-related
differences (Laitly et al., 2021; Stevens et al., 2007). In this study, we obtained internet images
of flower-visiting syrphid mimics and their hymenopteran models, as well as non-mimetic
syrphids. We extracted data from those images on the RGB colour of the flowers being visited.
We then used these data to test the hypotheses that (a) mimetic syrphids should follow the
flower colour choices of their hymenopteran models, rather than the yellow/white preferences of
their dipteran (and non-mimetic syrphid) ancestors, and (b) larger bodied syrphids should show
greater fidelity to the colour choices of their models.

Methods

Broadly speaking, we collected images depicting insects on flowers, and processed them to
remove digital colour profiles (see below). Then, we digitally cut out the areas within the images
showing the flowers, calculated their mean RGB colours and compared these values
statistically among groups. By using internet-obtained images, we accepted that we had to
ignore colours outside of the human-visible spectrum that might be relevant to predators, e.g. in
the UV range; and that unknown camera types and settings result in a degree of uncontrolled
colour bias in all photographs (Laitly et al., 2021; Stevens et al., 2007). We made the
simplifying assumption that photographs would not be systematically biased when depicting
different groups (such as models and mimics) towards particular camera settings or lighting

conditions. For example, we assumed that models and mimics would not systematically be

photographed at different times of day in a way that would affect the imaged flower colour.

Image collection

Mimetic syrphid species were matched to putative Hymenopteran models (Table S1) using a

near-exhaustive list of descriptions of their mimicry in syrphid-related literature, collated by FG.

Each mimic species and its set of one or more putative model species were grouped as a

"model-mimic pair" (e.g. *Eristalis pertinax* with *Apis mellifera*; *Volucella bombylans* var. *plumata*

with a pooled set comprising *Bombus terrestris*, *B. lucorum*, *B. hortorum*, *B. soroensis* and *B.*

jonellus; see Table S1).

Model-mimic pairs were then grouped into four distinct morphotypes, determined by the identity

of the model identified in Table S1. These were: Bumblebee [Apoidea:*Bombus*], Honeybee

[Apoidea:*Apis*], Solitary bee [all non-eusocial Apoidea i.e. not *Apis*, *Bombus* or other social

bee], and Social wasp [Aculeata:Vespidae]). We also included a final morphotype "Non-mimic

hoverflies" which were identified as such in the literature (e.g. *Ferdinandea cuprea*, *Baccha*

elongata; Table S1).

Internet images showing flowers being visited by mimics, their corresponding models, and non-

mimic hoverflies were obtained from web searches in March-April 2020, using common (e.g.

"Tapered drone fly") and binomial names (e.g. "Eristalis pertinax") as search terms. For each of

the four model morphotypes, we gathered photographs of 10 mimic and 10 model species with

10 photographs per species, as well as 5 non-mimic syrphid species with 20 photographs each.

Thus, in total we gathered 100 photographs for each morphotype. For each morphotype we

made sure to gather images of syrphids from across the phylogeny, and not from a single

evolutionary origin (Leavey et al., 2021; see Figure 1).

All images were collected from nationally recognised nature ID and photography sites, such as

bugguide (bugguide.net) and iNaturalist (inaturalist.org); specialised forums for ID of images of

Diptera, such as Diptera.info (diptera.info); or sites with clear author attribution of images, e.g.

Flickr (www.flickr.com) and where the author was a known authority, such as the Flickr gallery

maintained by S. Falk (https://www.flickr.com/people/63075200@N07/). Images were discarded

if the putative species ID could not be verified.

We chose only images showing the insect on a clearly visible flower and with dimensions above
approximately 275x183 pixels to ensure sufficient number of pixels per image for effective
flower colour comparison and consistency between monitors of varying resolution (Tkalcic and
Tasic, 2003). While this limited the sample size in the case of some species with few available
images of confirmed ID and sufficient quality, such as *Brachypalpus laphriformis,* this did not
significantly affect the sample sizes per group (N=100 images) or, we assumed, average RGB
values for each image.

Image processing

We modified all images to reduce the effect of camera setting adjustments applied by camera
profiles (exposure, brightness etc.) using the program RawTherapee (RawTherapee Team,
2020). To do this, we modified the RGB values obtained from each image directly, using
RawTherapee's "processing profile" function to remove any profile. Typically this brightened
each image, bringing R, G and B values of consistent blocks of colour closer to 'true' R, G and
B values (Stevens et al., 2007; Tkalcic & Tasic, 2003).

We then digitally extracted the part of each image containing the flower(s) using the GNU
Image Manipulation Program (GIMP) 2.10.14 (The GIMP Team, 2020), using GIMP's "free
select" tool. We used the histogram tool with a linear axis to extract RGB profiles for each
cutout, and recorded the median RGB values. A linear rather than logarithmic axis was used for
this process (Stevens et al., 2007), since it is best suited for use with photographs, particularly
those taken in the field, whereas a logarithmic axis is more commonly used for images
containing substantial areas of constant colour (Smith & Joost, 2012).

Body size

We used head width as a proxy for body size using a dataset, compiled by FSG, that combined
our previously published data (F. S. Gilbert, 1985) with subsequent unpublished measurements.
Twenty species were directly measured in this dataset (Table S2). For the 20 remaining species
not directly measured, we interpolated species-typical head width from intra-generic regressions
(typically very tight; F. S. Gilbert, 1985), using congeners in FSG's dataset, based on other
structural measurements such as wing length and body length taken from the literature (Bot &
Van de Meutter, 2023; Skevington et al., 2019; Thompson, 1991).

Data analysis

All analyses were conducted in R 4.4.1 (R Core Team, 2023). We log-transformed the original

three RGB values of the flower cutouts to conform with linear assumptions. We then reduced

their dimensions to scores along two principal axes of variation using Principal Component

Analysis (Figure 2). Broadly speaking, PCA1 equated to yellow (a combination of red and

green) and PCA2 to blue, and we refer to them henceforth as "yellow" and "blue" respectively.

To compare colour preferences, we first compared all syrphid morphotypes including non-

mimics (bumblebee, honeybee, solitary bee, social wasp, and non-mimic) with model

morphotypes, in a one-way analysis. We did not include phylogenetic information here

because, when hymenopteran models were included, the evolutionary distances between

syrphids and their models dwarfed the inter-species distances within these clades. We used

mixed-effects models, implemented in the lme4 package in R (Bates, 2010), with either PCA1

or PCA2 as a response, "morphotype" as a fixed effect and "species" as a random effect. We

used the emmeans package (Lenth, 2023) to conduct specific comparisons among

morphotypes.

Then we looked specifically at the effect of "being a model versus a mimic" within the different

morphotypes. For this, we trimmed the dataset to exclude non-mimic hoverflies, and fitted a

mixed effect model with "morphotype" and "model or mimic" plus their interaction as fixed

effects and "model-mimic species pair" as a random effect.

Finally, we investigated whether mimics had departed from non-mimics in their flower colour

preference, and whether body size could explain any such departure. For this, we compared

just the hoverfly species across the syrphid phylogeny, excluding hymenopteran models. We

used the MCMCglmm package in R to fit the model while also accounting for nonindependence

due to shared evolutionary history (the syrphid phylogeny), and also multiple observations for

each tip of the phylogeny (by including species as a random effect in the model). We used

"morphotype", "body size" and their interaction as fixed predictors, with 100000 MCMC

iterations, a thinning interval of 50 and a burnin of 2000, checking for convergence using trace

plots and Gelman-Rubin diagnostics over 3 replicate chains. We used weakly informative priors

with V=1 and nu=0.2 for both fixed and random effects.

The phylogeny (Figure 1) was developed following recommendations of Beaulieu et al (2012).

The base phylogeny was based on Leavey et al (2021) at the genus level, with species within

genera added as star phylogenies where necessary. Branch lengths for added species were

unknown, so branch lengths across the tree were scaled arbitrarily according to node depth,

following Grafen (1989), using a power of 0.8.

Results

181   The 45 syrphids in the dataset, their mean colour preferences, and those of their models, can be

182   seen in Figure 1. Flower colour data can be found in Table S3.

Figure 1. Syrphid phylogeny used in this study (with arbitrarily scaled branch lengths) showing

morphotype, body size, and the flower colour preferences of mimics and models, as assessed

by internet images depicting the species visiting a flower. Photos of representative species are

given, but are not necessarily part of the image dataset.

Two axes explained 83% of the variance in the colour of flowers photographed being visited by

models or mimics. Axis 1 (49% of the variance) was aligned strongly and identically with red (71%)

and green (70%) and will be known henceforth as yellowness. Axis 2 (34% of the variance) was

exclusively aligned with blue (98%) and we refer to this axis as blueness.

Scores along the yellowness and blueness axes for different morphotypes (models and mimics) are
shown in Figure 2.

Figure 2. Blueness (PCA2) plotted against yellowness (PCA1) for models and mimics of different
morphotypes, with means and 95% confidence ellipses shown in solid colours. Inset shows RGB
colours for the whole dataset, with representative extremes marked for comparison.

Hymenopteran models of different morphotypes differed in the yellowness of the flowers they chose
to visit, while syrphids (mimics and non-mimics) all chose flowers of similar yellowness (Linear
mixed model, X2=32.5, df=8, p<0.0001, Figure 3). When we examined effects of mimicry and
morphotype independently, excluding non-mimic hoverflies, there was a significant "mimic/model" x
morphotype interaction (X2=8.45, df=3, p=0.037, Table 2). Bumblebees chose less yellow flowers

than other models, which did not differ from each other (honeybee, solitary bee and social wasp). In
contrast, all mimic syrphids chose a similar yellowness, which was also similar to that chosen by
honeybees, solitary bees and social wasps, but was higher than the yellowness chosen by
bumblebees.

We then focused only on syrphids so that we could use phylogenetic information to ask whether
mimics and non-mimics had diverged in preference, and whether body size as well as morphotype
could explain flower colour preference.

Phylogenetic models of yellowness preference were practically indistinguishable from each other
according to their DIC, including the model with no explanatory factors at all (delta DIC < 2.5).
Across the phylogeny, syrphids of all morphotypes, mimics and non-mimics, preferred similar flower
yellowness (see Table 3). In the model with technically the lowest DIC (delta ~ 0.5), larger syrphids
had a very slight tendency to prefer less yellow flowers which we regard as negligible.

Table 1. Linear mixed-effect model of flower yellowness against morphotype, with species as random effect.

	Estimate	95% CI[1]	Chi-sq	DF	p-value
Intercept	0.07	-0.24, 0.38			
group			32.5	8	<0.001
Bumblebee mimic	—	—			
Bumblebee model	-0.83	-1.3, -0.39			
Honeybee mimic	-0.06	-0.50, 0.37			
Honeybee model	-0.04	-0.55, 0.46			
Solitary bee mimic	0.11	-0.32, 0.55			
Solitary bee model	-0.29	-0.72, 0.14			

Social wasp mimic	0.16	-0.27, 0.59
Social wasp model	0.00	-0.43, 0.43
Non-mimic	0.40	-0.08, 0.88

[1] CI = Confidence Interval

220 *Table 2. Linear mixed-effect model of flower yellowness against morphotype and "model or mimic",*
221 *with model-mimic pair as random effect.*

	Estimate	95% CI[1]	Chi-sq	DF	p-value
(Intercept)	0.02	-0.29, 0.34			
Morphotype			-	-	-
Bumblebee	—	—			
Honeybee	-0.12	-0.68, 0.43			
Solitary bee	0.11	-0.32, 0.55			
Social wasp	0.22	-0.22, 0.65			
Model or mimic			—	—	—
Mimic	—	—			
Model	-0.82	-1.3, -0.33			
Morphotype × Mimic			8.34	3	0.038

Honeybee model	1.1	0.28, 2.0	
Solitary bee model	0.46	-0.19, 1.1	
Social wasp model	0.64	0.00, 1.3	

[1] CI = Confidence Interval

222 *Table 3. Table of fitted phylogenetic models of yellowness preference among syrphids, showing*
223 *coefficients and ranked DIC values.*

(Intercept)	Morphotype	Head width	Morphotype × head width	df	logLik	DIC	delta
0.55		+		4	-795.83	1607.17	0.00
2.37	+	+	+	12	-793.67	1607.86	0.70
0.16				3	-796.72	1608.04	0.88
0.52	+	+		8	-795.41	1608.87	1.71
0.02	+			7	-796.42	1609.60	2.44

Figure 3: Yellowness preferences by morphotype and mimicry status. Syrphids (mimetic and non-mimetic) showed similar yellowness preference, while hymenopteran models differed (see text for details).

In contrast, there were significant differences among groups in blueness preference (Linear Mixed model, $X2=30.7$, $df=8$, $p<0.001$). Solitary bee morphotypes and non-mimic hoverflies preferred flowers with the least blue. Honeybee and social wasp morphotypes were intermediate. Both bumblebee models and mimics preferred flowers with more blue than others.

When we excluded non-mimics and looked specifically at effects of mimicry and morphotype, there was no interaction between "model/mimic" and "morphotype", and also no effect of "model/mimic". This indicated that models tended to choose flowers with similar B values to those chosen by mimics. In contrast, there was a strong effect of morphotype ($X2=26.0$, $df=3$, $p<0.0001$), showing that different hymenopteran groups (and their syrphid mimics which follow their preferences) tend to choose flowers with distinctive blueness values.

The three phylogenetic models of blueness preference containing "morphotype" were
indistinguishable from each other by their DIC ("morphotype only", "morphotype + body size",
"morphotype X body size"; delta < 0.57). However, all were better than models without "morphotype"
by > 4 DIC points (Table 6). This indicated that (as with the non-phylogenetic models above)
morphotype, but not body size, was associated with blueness preference. Bumblebee mimics and,
marginally, honeybee mimics had different blueness preferences from those of non-mimics, and
from those of mimics of solitary bees and social wasps (see table 7).

Table 4. Linear mixed-effect model of flower blueness against morphotype, with species as random
effect.

Characteristic	Beta	95% CI[1]	Chi-sq	DF	p-value
(Intercept)	0.39	0.13, 0.66			
group			30.721	8	<0.001
Bumblebee mimic	—	—			
Bumblebee model	0.03	-0.35, 0.40			
Honeybee mimic	-0.21	-0.58, 0.17			
Honeybee model	-0.51	-0.95, -0.07			
Non-mimic	-0.56	-0.98, -0.14			
Solitary bee mimic	-0.60	-0.98, -0.23			
Solitary bee model	-0.81	-1.2, -0.43			
Social wasp mimic	-0.45	-0.82, -0.07			

Social wasp model	-0.34	-0.72, 0.03

[1] CI = Confidence Interval

252 *Table 5. Linear mixed-effect model of flower blueness against morphotype and "model or mimic",*
253 *with model-mimic pair as random effect.*

	Estimate	95% CI[1]	Chisq	DF	p-value
Intercept)	0.42	0.19, 0.66			
Morphotype			21.31	3	<0.0001
Bumblebee	—	—			
Honeybee	-0.34	-0.72, 0.03			
Solitary bee	-0.69	-0.98, -0.40			
Social wasp	-0.36	-0.65, -0.08			
Model/mimic			0.66	1	0.42
Mimic	—	—			
Model	-0.09	-0.30, 0.13			

[1] CI = Confidence Interval

255 *Table 6. Table of fitted phylogenetic models of blueness preference among syrphids, showing*
256 *coefficients and ranked DIC values.*

	Model terms						
Intercept	Morphotype	Head width	Morphotype × head width	df	logLik	DIC	delta
+	+			7	-693.95	1406.80	0.00
+	+	+	+	12	-692.03	1407.27	0.47
+	+	+		8	-694.00	1407.38	0.58
+				3	-695.44	1411.51	4.70
+		+		4	-696.67	1413.37	6.57

260 *Table 7. Table of fixed coefficients for top phylogenetic model of blueness preference among*
261 *syrphids*

	Posterior mean	95% CI[1]	p(MCMC)
Intercept	-0.16	-0.59, 0.30	0.45
Non-mimic	—	—	—
Bumblebee mimic	0.50	0.13, 0.87	0.01
Honeybee mimic	0.34	-0.06, 0.69	0.09
Solitary bee mimic	-0.11	-0.50, 0.27	0.62
Solitary wasp mimic	0.05	-0.34, 0.41	0.78

[1] CI = Confidence Interval

269 *Figure 4. Blueness preferences by morphotype and mimicry status. Syrphid mimics were different*

270 *from non-mimics, and tended to follow the preferences of their models such that morphotypes*

271 *differed but models and mimics were statistically similar (see text for details).*

Discussion

The "blueness" of flowers preferred by mimetic syrphids, and in particular bumblebee mimics,

was statistically similar to that of their hymenopteran models - and in the case of bumblebee

and honeybee mimics, was bluer than those preferred by non-mimetic syrphids. Bumblebee

mimics such as *Volucella bombylans* have been documented as preferring blue flowers (Lunau,

2014), but to our knowledge this is the first systematic study to demonstrate a shift in flower

colour preference in bumblebee mimics across the syrphid phylogeny. For their part,

hymenopteran models showed differences in the blueness of flowers on which they tended to

be photographed. Specifically, bumblebees, and to a lesser extent honeybees, were

photographed on bluer flowers than other bees and wasps. This reflects a known preference

towards blue and violet flowers as food sources (Giurfa et al., 1995; Gumbert, 2000; Lubbock,

1882; Raine et al., 2006) that probably indicates availability of floral nectar in suitably accessible

locations for their generally long tongues (Raine & Chittka, 2007).

In yellowness preference, syrphid mimics also mostly tended to resemble their models, but their

preferences were also similar to those of their non-mimic relatives. So, while this finding could

be interpreted as behavioural mimicry of the models' preferences by the syrphids, it could also

simply reflect syrphid evolutionary ancestry, and/or dipteran ancestry generally. Dipterans,

including hoverflies, have strong preferences towards yellow and white flowers in many species

(Arnold et al., 2009) which again is probably related to accessibility of floral nectar for their

relatively short tongues (see Lunau, 2014). Bumblebees were associated with less-yellow

flowers than the other Hymenopterans, but bumblebee mimics did not follow this preference.

Instead, they retained the preference of their non-mimic relatives and of all other syrphids for

more-yellow flowers.

Why would syrphids mimicking social bees, in particular bumblebees, evolve away from their

ancestral flower colour preferences to follow the flower blueness preferences of their models? It

is difficult to escape the conclusion that their flower colour preferences have shifted to enhance

their mimicry, either by providing a multicomponent stimulus (Rowe, 1999) or by bringing

models and mimics into closer proximity to enhance protection (F. Gilbert, 2005). Bumblebee-

mimicking syrphids include some of the most faithful of all syrphid mimics (F. Gilbert, 2005).

Selection for high mimetic accuracy may be driven by both (1) large body size (Penney et al.,

2012) and (2) mimicry of mildly noxious models such as bumblebees (Sherratt, 2002). On the

one hand, Penney et al (2012) found that the largest syrphid mimics were the most faithful, and

that small body size (and hence meagre reward for predators) was the best explanation for the

occurrence of imperfect mimicry across species. Syrphids that show behavioural mimicry (mock

stinging, wing-wagging etc.) tend to be those with large bodies and high-quality mimicry

(although of wasps; Penney et al., 2014). While bumblebee mimics fit this description, in this

study we found that body size did not affect preference for either flower blueness or yellowness

in syrphids mimicking any kind of model.

On the other hand, Sherratt's (2002) hypothesis holds that models such as wasps are so

noxious to predators that any risk of misidentification confers protection even for poor quality

mimics. For models such as bumblebees that are less dangerous to predators than wasps (F.

Gilbert, 2005), the morphological "zone of protection" is more restricted, such that only near-
perfect mimics are protected. Strong enough selection on mimetic accuracy may extend to
selection for behavioural mimicry of flower colour preference. One interesting further question
would be whether any such selection has been strong enough to distinguish finer-scale
differences in colour preference among mimics of different bumblebee species. Fussell &
Corbet (1992) report that bumblebee species "colour groups" (e.g. "brown", "black with red
abdominal band") show distinct differences in flower usage, perhaps reflecting flowers with
suitable corolla depths for their tongues (Brian, 1957; Faegri & van der Pijl, 1979). Flower colour
loosely predicts corolla depth (Binkenstein et al., 2016) which may translate into colour
preferences that mimics could exploit. Looking at the flower preferences (and tongue lengths) of
the distinct morphs of polymorphic mimics such as *V. bombylans* and *Merodon equestris* might
provide a good test of this hypothesis. Another natural further question is whether bumblebee-
mimicking syrphids have evolved longer tongues to facilitate exploitation of "bee flowers" in the
same way as bee-flies. Tongue length is particularly labile among hoverfly species
independently of body size (F. S. Gilbert, 1985). *Rhingia*, a long-tongued genus of syrphids,
although non-mimetic, are unusual amongst syrphids in preferring to visit red and blue flowers
rather than yellow ones (Haslett, 1989; Robertson, 1928); more generally, flies that visit blue
flowers often have elongate mouthparts (reviewed in Lunau, 2014).

Why mimic specifically the blueness and not the yellowness of bumblebee flower preferences?
Some syrphids show an innate preference for yellow flowers that cannot be extinguished by
training (Lunau et al., 2018), so it may be that the strong preference for yellow is stabilised by
other evolutionary mechanisms and does not respond to selection for copying the preferences
of models.

Ultimately, selection favouring mimicry arises from the visual systems of predators (Rowe,
1999; Speed & Turner, 1999). Internet photographs cannot possibly capture the range of colour
involved in flower-pollinator-predator interactions (Stevens et al., 2007). Bees mostly have three
kinds of photoreceptor: broadly, UV, blue and green (Briscoe & Chittka, 2001; Goldsmith, 1990),
while syrphids and other flies have an effectively tetrachromatic system with an additional
receptor in the UV range (An et al., 2018; Lunau, 2014). Birds also have potentially four
receptors, but their additional receptor is in the red range (Hart & Hunt, 2007) which appears
achromatic to insects - despite this, bird pollinators do not appear to show particular flower
colour preferences that might exploit this private colour channel (Lunau et al., 2011). Whether

avian predators pay more attention to the blueness of flowers visited by potential prey than to

their yellowness remains a question to be investigated.

Of course, aspects of flowers other than colour are important in determining visitation by

hoverflies, e.g. flower size (Sutherland et al., 1999) or height (Ssymank, 2001) and many

others. Nevertheless, despite the obvious limitations of using internet photographs, we have

managed to discern broad patterns consistent with behavioural mimicry of the flower-colour

choices of hymenopteran models by syrphid mimics across the phylogeny, specifically in flower

"blueness". These patterns were most evident in bumblebee mimics, but were sufficiently strong

that models and mimics generally were not statistically separable in their blueness preferences.

How diverse and multimodal traits interact to determine the effectiveness of antipredator

defence is largely unknown (Kikuchi et al., 2023) - some traits combine with additive effects,

others synergistic, and others antagonistic. Mimicry often involves multiple modalities (Rowe,

1999), such as morphology, colouration, behaviour, phenology and habitat use. Habitat-choice

mimicry, such as we identify here, likely contributes to a "portfolio" of mimicry-enhancing traits in

any given mimic species. Its costs and benefits likely depend upon factors such as predator-,

model- and habitat-specific selection pressures (see e.g. Valkonen et al., 2012), competing

allocation demands within the organism (Kikuchi et al., 2023), and how strong selection there is

for mimetic accuracy in the first place (F. Gilbert, 2005; Penney et al., 2012). The hoverfly clade

contains large numbers of origins of mimicry (Leavey et al., 2021) with members resembling

diverse different models and employing multiple different modalities (Howarth & Edmunds,

2000; C. D. Moore & Hassall, 2016; Penney et al., 2014). We suggest that hoverflies are

potentially an exceptional system in which to explore the evolution of multimodal mimicry,

including the habitat-choice mimicry we have identified here.

Acknowledgements

370   We thank M. Stevens for helpful advice on image processing; G. Ruxton for helpful comments

371   on the manuscript; J. Boynton, K. Lester, E. Howard and other undergraduate students involved

372   in this and similar projects over the years.

Ethical Statement

All modification and usage of internet images in this study was for non-commercial research purposes (Levering, 1999, Jaszi, 2013). The research received ethical approval (in the form of a waiver) from the University of Hull Faculty Ethics Committee.

Author contributions

JDJG conceived and co-designed the study, co-supervised data collection, conducted all data analysis and wrote the manuscript. JCJ collected all data and wrote an initial draft of the project as an undergraduate dissertation. LM co-designed the study, co-supervised data collection and edited the manuscript. FG provided data on models for different mimics, morphological data and phylogenetic information, and edited the manuscript.

References

An, L., Neimann, A., Eberling, E., Algora, H., Brings, S., & Lunau, K. (2018). The yellow specialist: dronefly Eristalis tenax prefers different yellow colours for landing and proboscis extension. *The Journal of Experimental Biology*, *221*(Pt 22), jeb184788.

Arnold, S. E. J., Savolainen, V., & Chittka, L. (2009). Flower colours along an alpine altitude gradient, seen through the eyes of fly and bee pollinators. *Arthropod-Plant Interactions*, *3*(1), 27–43.

Atsumi, K., & Koizumi, I. (2017). Web image search revealed large-scale variations in breeding season and nuptial coloration in a mutually ornamented fish, *Tribolodon hakonensis*. *Ecological Research*, *32*(4), 567–578.

Austen, E. J., Lin, S.-Y., & Forrest, J. R. K. (2018). On the ecological significance of pollen color: a case study in American trout lily (Erythronium americanum). *Ecology*, *99*(4), 926–937.

Bates, D. (2010). lme4 : linear mixed-effects models using S4 classes. *Http://cran.r-project.org/package=lme4*. https://cir.nii.ac.jp/crid/1571698600750746240

Binkenstein, J., Stang, M., Renoult, J. P., & Schaefer, H. M. (2016). Weak correlation of flower color and nectar-tube depth in temperate grasslands. *Journal of Plant Ecology*, *10*(2), rtw029.

Bot, S., & Van de Meutter, F. (2023). *Hoverflies of Britain and North-west Europe: A*

photographic guide. Bloomsbury Wildlife.

Brian, A. D. (1957). Differences in the flowers visited by four species of bumble-bees and their

causes. *The Journal of Animal Ecology*, *26*(1), 71.

Briscoe, A. D., & Chittka, L. (2001). The evolution of color vision in insects. *Annual Review of*

Entomology, *46*(1), 471–510.

Ceccarelli, F. S. (2008). Behavioral mimicry in Myrmarachne species (Araneae, Salticidae) from

North Queensland, Australia. *The Journal of Arachnology*, *36*(2), 344–351.

de Buck, N. (1990). *Bloembezoek en bestuivingsecologie van zweefvliegen (Diptera,*

Syrphidae) in het bijzonder voor België. Royal Belgian Institute of Natural Sciences.

Dlusski, G. M. (1984). Are dipteran insects protected by their similarity to stinging Hymenoptera.

Byulleten'Moskovskogo Obshchestva Ispytatelei Prirody, Otdel Biologicheskii, *89*, 25–40.

Faegri, K., & van der Pijl, L. (1979). *The principles of pollination ecology - 3rd edition*.

https://agris.fao.org/search/en/providers/122621/records/647396dae0110688009805d2

Fussell, M., & Corbet, S. A. (1992). Flower usage by bumble-bees: A basis for forage plant

management. *The Journal of Applied Ecology*, *29*(2), 451.

Gilbert, F. (2005). The evolution of imperfect mimicry. *Symposium-Royal Entomological Society*

of London.

https://www.cabidigitallibrary.org/doi/pdf/10.1079/9780851998121.0000#page=244

Gilbert, F. S. (1985). Morphometric patterns in hoverflies (Diptera, Syrphidae). *Proceedings of*

the Royal Society of London, *224*(1234), 79–90.

Giurfa, M., Núñez, J., Chittka, L., & Menzel, R. (1995). Colour preferences of flower-naive

honeybees. *Journal of Comparative Physiology. A, Neuroethology, Sensory, Neural, and*

Behavioral Physiology, *177*(3), 247–259.

Golding, Y. C., & Edmunds, M. (2000). Behavioural mimicry of honeybees (Apis mellifera) by

droneflies (Diptera: Syrphidae: Eristalis spp.). *Proceedings of the Royal Society of London.*

Series B: Biological Sciences, *267*(1446), 903–909.

Golding, Y. C., & Ennos, A. R. (2005). Biomechanics and behavioural mimicry in insects. In A.

Herrel, T. Speck, & N. P. Rowe (Eds.), *Ecology and biomechanics: a mechanical approach*

to the ecology of animals and plants. Taylor & Francis.

Goldsmith, T. H. (1990). Optimization, constraint, and history in the evolution of eyes. *The*

Quarterly Review of Biology, *65*(3), 281–322.

Gumbert, A. (2000). Color choices by bumble bees (Bombus terrestris): innate preferences

and generalization after learning. *Behavioral Ecology and Sociobiology*, *48*(1), 36–43.

Hannah, L., Dyer, A. G., Garcia, J. E., Dorin, A., & Burd, M. (2019). Psychophysics of the

hoverfly: categorical or continuous color discrimination? *Current Zoology*, *65*(4), 483–492.

Hart, N. S., & Hunt, D. M. (2007). Avian visual pigments: characteristics, spectral tuning, and

evolution. *The American Naturalist*, *169 Suppl 1*(S1), S7–S26.

Haslett, J. R. (1989). Interpreting patterns of resource utilization: randomness and selectivity in

pollen feeding by adult hoverflies. *Oecologia*, *78*(4), 433–442.

Howarth, B., & Edmunds, M. (2000). The phenology of Syrphidae (Diptera): are they Batesian

mimics of Hymenoptera? *Biological Journal of the Linnean Society. Linnean Society of*

London, *71*(3), 437–457.

Howarth, B., Edmunds, M., & Gilbert, F. (2004). Does the abundance of hoverfly (Syrphidae)

mimics depend on the numbers of their hymenopteran models? *Evolution; International*

Journal of Organic Evolution, *58*(2), 367–375.

Ilse, D. (1949). Colour discrimination in the dronefly, Eristalis tenax. *Nature*, *163*(4137), 255.

Kastinger, C., & Weber, A. (2001). Bee-flies (Bombylius spp., Bombyliidae, Diptera) and the

pollination of flowers. *Flora*, *196*(1), 3–25.

Kikuchi, D. W., Allen, W. L., Arbuckle, K., Aubier, T. G., Briolat, E. S., Burdfield-Steel, E. R.,

Cheney, K. L., Daňková, K., Elias, M., Hämäläinen, L., Herberstein, M. E., Hossie, T. J.,

Joron, M., Kunte, K., Leavell, B. C., Lindstedt, C., Lorioux-Chevalier, U., McClure, M.,

McLellan, C. F., … Exnerová, A. (2023). The evolution and ecology of multiple antipredator

defences. *Journal of Evolutionary Biology*, *36*(7), 975–991.

Kitamura, T., & Imafuku, M. (2015). Behavioural mimicry in flight path of Batesian intraspecific

polymorphic butterfly Papilio polytes. *Proceedings. Biological Sciences / The Royal*

Society, *282*(1809), 20150483.

Kugler, H. (1950). Flower visitation of Eristalomyia tenax. *Zeitschrift Für Vergleichende*

Physiologie, *3*(4), 328–347.

Laitly, A., Callaghan, C. T., Delhey, K., & Cornwell, W. K. (2021). Is color data from citizen

science photographs reliable for biodiversity research? *Ecology and Evolution*, *11*(9),

4071–4083.

Leavey, A., Taylor, C. H., Symonds, M. R. E., Gilbert, F., & Reader, T. (2021). Mapping the

evolution of accurate Batesian mimicry of social wasps in hoverflies. *Evolution;*

International Journal of Organic Evolution, *75*(11), 2802–2815.

Leighton, G. R. M., Hugo, P. S., Roulin, A., & Amar, A. (2016). Just Google it: assessing the

use of Google Images to describe geographical variation in visible traits of organisms.

Methods in Ecology and Evolution / British Ecological Society, *7*(9), 1060–1070.

Lenth, R. V. (2023). *emmeans: Estimated Marginal Means, aka Least-Squares Means*.

https://CRAN.R-project.org/package=emmeans

Lubbock, J. (1882). *Ants. Bees, and wasps*. Kegan Paul, Trench & Company.

Lunau, K. (2014). Visual ecology of flies with particular reference to colour vision and colour

preferences. *Journal of Comparative Physiology. A, Neuroethology, Sensory, Neural, and*

Behavioral Physiology, *200*(6), 497–512.

Lunau, K., An, L., Donda, M., Hohmann, M., Sermon, L., & Stegmanns, V. (2018). Limitations of

learning in the proboscis reflex of the flower visiting syrphid fly Eristalis tenax. *PloS One*,

13(3), e0194167.

Lunau, K., Papiorek, S., Eltz, T., & Sazima, M. (2011). Avoidance of achromatic colours by bees

provides a private niche for hummingbirds. *The Journal of Experimental Biology*, *214*(Pt 9),

1607–1612.

Mappes, J., & Alatalo, R. V. (1997). Batesian Mimicry and Signal Accuracy. *Evolution;*

International Journal of Organic Evolution, *51*(6), 2050.

Moore, C. D., & Hassall, C. (2016). A bee or not a bee: an experimental test of acoustic mimicry

by hoverflies. *Behavioral Ecology: Official Journal of the International Society for*

Behavioral Ecology, *27*(6), 1767–1774.

Moore, M. P., Lis, C., Gherghel, I., & Martin, R. A. (2019). Temperature shapes the costs,

benefits and geographic diversification of sexual coloration in a dragonfly. *Ecology Letters*,

22(3), 437–446.

Morse, D. H. (1979). Prey capture by the crab spider Misumena calycina (Araneae:

Thomisidae). *Oecologia*, *39*(3), 309–319.

Penney, H. D., Hassall, C., Skevington, J. H., Abbott, K. R., & Sherratt, T. N. (2012). A

comparative analysis of the evolution of imperfect mimicry. *Nature*, *483*(7390), 461–464.

Penney, H. D., Hassall, C., Skevington, J. H., Lamborn, B., & Sherratt, T. N. (2014). The

relationship between morphological and behavioral mimicry in hover flies (Diptera:

Syrphidae). *The American Naturalist*, *183*(2), 281–289.

Raine, N. E., & Chittka, L. (2007). The adaptive significance of sensory bias in a foraging

context: floral colour preferences in the bumblebee Bombus terrestris. *PloS One*, *2*(6),

e556.

Raine, N. E., Ings, T. C., Dornhaus, A., Saleh, N., & Chittka, L. (2006). Adaptation, genetic drift,

pleiotropy, and history in the evolution of bee foraging behavior. In *Advances in the Study*

of Behavior (Vol. 36, pp. 305–354). Elsevier.

R Core Team. (2023). *R: A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.R-project.org/

Rettenmeyer, C. W. (1970). Insect mimicry. *Annual Review of Entomology*, *15*(1), 43–74.

Robertson, C. (1928). Flowers and insects: XXV. *Ecology*, *9*(4), 505–526.

Rodríguez-Gasol, N., Avilla, J., Alegre, S., & Alins, G. (2019). Sphaerophoria rueppelli adults

change their foraging behavior after mating but maintain the same preferences to flower

traits. *BioControl (Dordrecht, Netherlands)*, *64*(2), 149–158.

Rowe, C. (1999). Receiver psychology and the evolution of multicomponent signals. *Animal*

Behaviour, *58*(5), 921–931.

Ruxton, G. D. (2009). Non-visual crypsis: a review of the empirical evidence for camouflage to

senses other than vision. *Philosophical Transactions of the Royal Society of London.*

Series B, Biological Sciences, *364*(1516), 549–557.

Sherratt, T. N. (2002). The evolution of imperfect mimicry. *Behavioral Ecology: Official Journal*

of the International Society for Behavioral Ecology, *13*(6), 821–826.

Skevington, J. H., Locke, M. M., Young, A. D., Moran, K., Crins, W. J., & Marshall, S. A. (2019).

Field guide to the flower flies of northeastern north America. Princeton University Press.

Smith, J., & Joost, R. (2012). Digital Imaging Projects. In *GIMP for Absolute Beginners* (pp.

131–166). Apress.

Speed, M. P., & Turner, J. R. G. (1999). Learning and memory in mimicry: II. Do we understand

the mimicry spectrum? *Biological Journal of the Linnean Society. Linnean Society of*

London, *67*(3), 281–312.

Ssymank, A. (1989). *Das Vegetationsmosaik eines Waldgebietes der Schwarzwaldvorbergzone*

und seine funktionale Bedeutung für blütenbesuchende Insekten-unter besonderer

Berücksichtigung der Syrphidae (Diptera): Text: mit 12 pflanzensoziologischen Tab., 22

phänologischen Tab., 7 biozönologischen und blütenbiologischen Tab. und weiteren 33

Tab. im Text.

Ssymank, A. (2001). Vegetation and flower-visiting insects in cultivated landscapes. *Bonn Bad*

Godesb, *64*, 513.

Stevens, M., Párraga, C. A., Cuthill, I. C., Partridge, J. C., & Troscianko, T. S. (2007). Using

digital photography to study animal coloration. *Biological Journal of the Linnean Society.*

Linnean Society of London, *90*(2), 211–237.

Sutherland, J. P., Sullivan, M. S., & Poppy, G. M. (1999). The influence of floral character on

the foraging behaviour of the hoverfly, Episyrphus balteatus. *Entomologia Experimentalis et*

Applicata, *93*(2), 157–164.

Szucsich, N. U., & Krenn, H. W. (2000). Morphology and function of the proboscis in

Bombyliidae (Diptera, Brachycera) and implications for proboscis evolution in Brachycera.

Zoomorphology, 120(2), 79–90.

Thompson, F. (1991). The flower fly genus Ornidia (Diptera: Syrphidae). *Proceedings of the*
Entomological Society of Washington, *93*(2), 248–260.

Tkalcic, M., & Tasic, J. F. (2003). *Colour spaces: perceptual, historical and applicational*
background. The IEEE Region 8 EUROCON 2003. Computer as a Tool, Ljubljana,
Slovenia. https://ieeexplore.ieee.org/abstract/document/1248032

Valkonen, J. K., Nokelainen, O., Niskanen, M., Kilpimaa, J., Björklund, M., & Mappes, J. (2012).
Variation in predator species abundance can cause variable selection pressure on warning
signaling prey. *Ecology and Evolution*, *2*(8), 1971–1976.

Dlusskii GM (1984) [Are Diptera protected by their similarity to stinging Hymenoptera?]
Byulleten Moskovskogo Obshchestva Ispytatelei Prirody Otdel Biologicheskii 89(5): 25-40
(in Russian)

Gilbert FS (1985) Morphometric patterns in hoverflies (Diptera, Syrphidae). *Proceedings of the*
Royal Society of London B Biological Sciences 224: 79-90

Szucsich NU & Krenn HW (2000) Morphology and function of the proboscis in Bombyliidae
(Diptera, Brachycera) and implications for proboscis evolution in Brachycera.
Zoomorphology 120: 79-90

555