Title: CDK9 interacts with a RanGTP-NEMP1-Importin β complex to regulate erythroid enucleation

Authors: Lucas M. Newton[1,2,3,4], Krystle Y. B. Lim[1,2], Donia Y. Abeid[1,2], Christina B. Wölwer[5], Chad J. Johnson[6], Sarah M. Russell[3,4], Edwin D. Hawkins[7,8], Patrick O. Humbert[1,2,9,10]*

Affiliations:

[1] Department of Biochemistry & Chemistry, La Trobe University, Melbourne, VIC, 3073, Australia.
[2] La Trobe Institute for Molecular Science, La Trobe University, Melbourne, VIC, 3073, Australia.
[3] Optical Sciences Centre, Swinburne University of Technology, Hawthorn, VIC, 3072, Australia.
[4] Division of Cancer Research, Peter MacCallum Cancer Centre, Parkville, VIC, 3010, Australia.
[5] Institute of Pathology, University Hospital Cologne, Cologne, 50937, Germany.
[6] La Trobe BioImaging Platform, La Trobe University, Melbourne, VIC, 3073, Australia.
[7] The Walter and Eliza Hall Institute of Medical Research, Parkville, VIC, 3010, Australia.
[8] Department of Medical Biology, University of Melbourne, Parkville, VIC, 3010, Australia.
[9] Department of Biochemistry & Pharmacology, University of Melbourne, Parkville, VIC, 3010, Australia.
[10] Department of Clinical Pathology, University of Melbourne, Parkville, VIC, 3010, Australia.

*Corresponding author:
Patrick O. Humbert
p.humbert@latrobe.edu.au
La Trobe University
Melbourne, Victoria, Australia 3073

Running title: CDK9 and importin β regulate erythroid enucleation

Key words: Enucleation, erythropoiesis, CDK9, importin β

Text word count (incl. in-text references and acknowledgements, excl. methods): 5428
Abstract word count: 170
Number of figures: 6 (+ 5 supplementary)
Number of tables: 2 supplementary
Number of references: 80

Key Points

- Importin β physically interacts with CDK9 in erythroid cells and is a novel regulator of erythroid enucleation
- Functional assays show that CDK9 and importin β act prior to calcium-dependent actin-mediated nuclear extrusion

Abstract

Erythroid enucleation is the final stage of erythroid terminal differentiation and involves the separation of an orthochromatic erythroblast into two daughter cells; a pyrenocyte containing the extruded nucleus, and the other a reticulocyte that will become a red blood cell. Our previous work identified CDK9 as a regulator of erythroid enucleation that appears to act independently of its known role in regulating RNA polymerase II transcription, suggesting the potential for a new CDK9 role. Using a co-immunoprecipitation and mass spectrometry approach, we have identified the interactome of CDK9 in differentiating erythroblasts. We show that CDK9 interacts with a RanGTP-NEMP1-Importin β complex during erythroid terminal differentiation, and inhibition of importin β in erythroblasts blocks erythroid enucleation. Using imaging analysis and functional assays of enucleating erythroblasts, we show that CDK9 and importin β co-locate at a critical site of activity opposite to the nucleus before nuclear extrusion and we describe a novel finding that physically links CDK9 and importin β activity prior to CaM/Ca^{2+} signalling and subsequent F-actin activity to achieve enucleation.

Introduction

Erythroid enucleation is the asymmetrical separation of the future erythrocyte from its nucleus and is the final stage of erythropoiesis before the developing erythroblast enters the peripheral blood stream (Palis, 2014). Enucleation is a rate limiting step in efforts to produce red blood cells *ex vivo* for transfusion medicine (Christaki et al., 2019; Gallego-Murillo et al., 2022), necessitating a clearer understanding of its regulation. Several recent reviews describe what is currently known of the molecular and cellular steps required for enucleation (Menon & Ghaffari, 2021; Moras et al., 2017; Newton et al., 2024). At a cellular level, chromatin compaction begins at the polychromatic erythroblast stage, one division prior to the orthochromatic erythroblast, which subsequently undergoes nuclear polarisation and nuclear extrusion; the anucleate cell, termed a reticulocyte, continues developing in the bone marrow for a short time while it undergoes further morphological changes and organelle clearance before moving into the circulation (Menon & Ghaffari, 2021; Moras et al., 2017). Importantly, the extruded nucleus, termed pyrenocyte, is contained within a separate cell membrane and enclosed in a thin layer of cytoplasm, suggesting the mechanism is not purely exocytosis of nuclear material (Yoshida et al., 2005). Enucleation has been compared to both cytokinesis, and the apoptosis-driven denucleation events observed in lens epithelia and keratinocytes (Rogerson et al., 2018; Wride, 2011), however the mechanisms that underpin enucleation appear quite distinct from either process (Newton et al., 2024). The molecular regulation of this unique cellular process is poorly understood, but we recently made the surprising finding that CDK9 is essential for erythroid enucleation (Wölwer et al., 2015). Interestingly, our findings also suggested that CDK9 did not act through the canonical RNA Pol II-dependent pathway (Anshabo et al., 2021; Paparidis et al., 2017; Wölwer et al., 2015). Despite its importance, the precise mechanisms that govern enucleation remain relatively poorly understood.

Cytoskeletal elements such as F-actin and non-muscle myosin IIB (NMIIB) are required for the nuclear extrusion process, and while the formation of a contractile actomyosin ring (CAR) has been proposed (Ji et al., 2008; Konstantinidis et al., 2012; Koury et al., 1989; Li et al., 2017), other models suggest that enucleation is achieved primarily through the activity of an actin-mediated enucleosome and by membrane reorganisation by trafficking of endocytic vesicles (An et al., 2021; Keerthivasan et al., 2010; Liang et al., 2015; Nowak et al., 2017). Calcium signalling through the calmodulin (CaM) pathway is critical to achieve enucleation following uptake of extracellular calcium prior to nuclear extrusion (Wölwer et al., 2016). This is likely due to the ability for CaMKII to modulate the actin cytoskeleton by altering F-actin cross-linking and the rate of F-actin polymerisation (Hoffman et al., 2013; Ohta et al., 1986; Sanabria et al., 2009).

Although many of the above factors are well established, the use of standard genetic approaches to identify new regulators of enucleation in differentiation systems can be problematic in terms of their interpretation. This is because of the possible indirect effects of genetically mediated perturbation, when using CRISPR or RNAi approaches on upstream processes in terminal differentiation, which can indirectly result in reduced downstream enucleation. To circumvent this, we have developed and used a chemical genetics approach to identify novel regulators of enucleation. We initially identified Cyclin-dependent kinase-9 (CDK9) as a novel regulator of erythroid enucleation (Wölwer et al., 2015). CDK9 is predominantly known for its role as the catalytic subunit of the positive-transcription and elongation factor-B (P-TEFb), which alongside its cyclin partner cyclin T1 (CycT1), cyclin T2a/b (CycT2a/b) or cyclin K, plays a fundamental role in the regulation of RNA polymerase II (RNA Pol II) transcriptional activity (Paparidis et al., 2017; Qin et al., 2017). However, our previous study had shown that blocking the activity of RNA Pol II in late erythropoiesis does not arrest enucleation (Wölwer et al., 2015), suggesting a new role for CDK9 in enucleation independent of RNA Pol II.

Here, using mouse and human erythroid culture models, we have characterised the role of CDK9 in erythroid enucleation. We have conducted imaging analysis of enucleating erythroblasts and found that CDK9 and Cyclin T1 localise nearby F-actin prior to nuclear extrusion. Additionally, we show that CDK9 localisation is dependent on F-actin activity. To identify the pathways CDK9 may be regulating in enucleation, we carried out a proteomics screen to identify relevant CDK9 interactors during erythroid enucleation. We show that CDK9 associates with a RanGTP-NEMP1-Importin β complex and that CDK9 and importin β function upstream of calcium signalling and F-actin activity prior to nuclear extrusion. Our study provides new insights into the function of CDK9 in enucleation, and identifies a key link between the new regulator of enucleation Importin β and previously described regulators CDK9 (Wölwer et al., 2015) and NEMP1 (Hodzic et al., 2022). As CDK9 inhibitors are being tested for clinical use against a variety of cancers, our study also highlights potential side-effects of CDK9 inhibitors on erythropoiesis and enucleation.

Material and Methods

Materials
Antibodies, inhibitors and other reagents are listed in supplemental Table 1. pBABE-Flag-Cdk9-IRES-eGFP, pBABE-Flag-Cdk9-T186A-IRES-eGFP and pBABE-Flag-Cdk9-D167N-IRES-eGFP were gifts from Andrew Rice (Addgene plasmid #28096, RRID: Addgene_28096; Addgene plasmid #28097, RRID: Addgene_28097; Addgene plasmid #28098, RRID: Addgene_28098). pBABE GFP was a gift from William Hahn (Addgene plasmid #10668, RRID: Addgene_10668).

Isolation of orthochromatic erythroblasts
All animal procedures were approved by the La Trobe University Animal Ethics Committee. Splenic orthochromatic erythroblasts were isolated from C57Bl/6 mice between 6-14 weeks of age by phenhylhydrazine treatment and flow cytometry as previously described (Wölwer et al., 2015). Briefly, we sorted for viable (propidium iodide negative) $Ter119^+CD44^{low}$ cells, which have been previously validated to be primarily orthochromatic erythroblasts (Wölwer et al., 2015). In normal conditions, we regularly observed ~60% enucleation 5 hours after flow cytometric sorting, which remained stable for up to 16 hours.

HUDEP-2 and BEL-A cell culture
HUDEP-2 and BEL-A cells were cultured as previously described (Trakarnsanga et al., 2017).

Flow cytometry
Enucleation of mouse orthochromatic erythroblasts was quantified using a FACSymphony A3 (Becton Dickson; Franklin Lakes, NJ) running FACS Diva software (Becton Dickson; Franklin Lakes, NJ) and analysed using FlowJo v10.8.1 (Becton Dickson; Franklin Lakes, NJ). Percentage of enucleation was obtained by dividing the number of enucleated cells ($Ter119^+$ Hoechst $^-$) by the sum of enucleated cells and erythroblasts ($Ter119^+$ Hoechst$^+$) and multiplying by 100. Propidium iodide was used to eliminate dead cells from the analysis.

Cytospins
6×10^4 cells were centrifuged onto slides at 320 rpm for 4 minutes using a Cytospin 4 Centrifuge (Thermo Fisher; Waltham, MA). Slides were air dried prior to fixation in methanol and subsequently stained using RapidDiff as per the manufacturer's guidelines. HUDEP-2 and BEL-A enucleation rates, and cell morphologies were quantified manually by images captured using an Olympus IX81 microscope using 100x/1.4NA oil objective running cellSens Dimension software (Evident/Olympus Life Sciences; Tokyo, Japan).

Immunofluorescence confocal microscopy
Cells were prepared for immunofluorescence microscopy as previously described (Smith et al., 2018) with minor alterations. Briefly, cells were collected by centrifugation following overnight incubation +/- inhibitors, washed in PBS and immediately fixed in 4% paraformaldehyde in PBS overnight at 4°C. Cells were subsequently washed 3 times in PBS and permeabilised in PBS containing 0.3% Triton X-100 for 15 minutes at RT, before blocking with PBS containing 3% BSA, 1% donkey serum for 2 hours at RT. Cells were incubated with primary antibodies in blocking buffer overnight at 4°C, washed 3 times in PBS and incubated with secondary antibodies and 1 µg/mL DAPI in blocking buffer for 2 hours. Cells were washed 3 times in PBS before centrifugation onto poly-D-lysine treated microscopy slides using a cytospin centrifuge, and subsequently mounted in ProLong Gold antifade reagent. Slides were imaged using a Zeiss LSM800 confocal microscope using 63x/1.4NA oil objective running Zeiss Zen Blue software (Zeiss; Oberkochen, Germany). Nuclear to cytoplasmic signal ratios were calculated using Fiji (ImageJ) (Schindelin et al., 2012).

Retroviral transduction of HUDEP-2 cells

Retrovirus was generated by calcium phosphate transfection of HEK293T cells as previously described (Jordan et al., 1996). At 12 hours post-transfection, transfection media was replaced with HUDEP-2 expansion media. Viral media was collected after 24 hours, and HUDEP-2 cells were seeded at 3×10^5 cells/mL in viral growth media for 24 hours. Transduced cells were isolated using FACS Aria Fusion (Becton Dickson; Franklin Lakes, NJ) running FACS Diva software (Becton Dickson; Franklin Lakes, NJ) by sorting for GFP[+] cells.

CDK9 co-immunoprecipitation, mass spectrometry and protein database search

Detailed methodology is available in the Supplementary Methods. Co-immunoprecipitation studies were performed using the Pierce Classic Magnetic IP/Co-IP Kit (Thermo Fisher; Waltham, MA) as per the manufacturer's instructions with minor modifications. Protein extracts were quantified using the DC Protein Assay (BioRad; Hercules, CA). Protein samples were dried and desalted before liquid chromatography tandem mass spectrometry (LC MS/MS) using a Thermo Ultimate 3000 RSLC nano UHPLC system and analysed on a Thermo Q-Exactive HF Orbitrap mass-spectrometer (Thermo Fisher Scientific, Waltham, MA). On-bead trypsin digestion was carried out as previously described (Antonicka et al., 2020).

The protein database searches were conducted using Sequest HT search engine via the Proteome Discoverer 2.4 software suite (Thermo Fisher Scientific, Waltham, MA). Spectra were matched against the Homo sapiens reference proteome downloaded from Uniprot (The UniProt Consortium, 2023). Proteins that were present in all three replicates with a minimum 3-fold enrichment compared to the IgG controls were considered relevant. Results, including generation of Venn diagrams and heatmaps, were analysed using FunRich v3.1.4 (Fonseka et al., 2021). Interaction and gene ontology (GO) analysis was performed using the STRING database (Szklarczyk et al., 2023).

Western blot

Cell lysates were isolated using NETN lysis buffer (250 mM NaCl, 5 mM EDTA (pH 8.0), 50 mM Tris-HCl (pH 8.0), 0.5% NP-40 in distilled water) containing Complete mini protease inhibitor (Roche; Basel, Switzerland) and PhosSTOP phosphatase inhibitor (Roche; Basel, Switzerland). Protein levels were quantified using a DC protein assay (BioRad; Hercules, CA), denatured using SDS loading buffer and subsequently resolved on standard sodium dodecyl-sulfate polyacrylamide gels before western transfer onto polyvinylidene difluoride membrane (Merck Millipore; Burlington, MA). Membranes were blocked in 3% BSA in 0.2% Tween-20 in PBS for 1 hour, then incubated overnight with primary and secondary antibodies respectively. Membranes were visualised using an Odyssey infrared imager (LiCor Biosciences; Lincoln, NE). Antibodies and concentrations used are listed in Supplementary Table 1.

Statistical analysis

Statistical calculations and analysis were performed using GraphPad Prism version 9.2.0 (GraphPad Software, San Diego, CA). Statistical differences were analysed using ANOVA tests, with Dunnett's or Tukey's tests for multiple comparisons. Detailed statistics are described in the figure legends. n = number of cells counted, or number of replicates. P-values below 0.05 were considered statistically significant.

Results

Phosphorylated CDK9 (Thr186) and Cyclin T1 co-localise with F-actin during nuclear extrusion
To better understand how CDK9 might regulate enucleation, we investigated the localisation of CDK9 and key binding partner cyclin T1 during erythroid differentiation and enucleation. We used mouse and human erythroid *in vitro* systems to compare, for two stages of enucleation **(Figure 1A)**, the localisation and phosphorylation (Thr186) to reflect CDK9 activation **(Figure 1)**. CDK9 activity is highly regulated by phosphorylation, particularly the T-Loop region site Thr-186, where phosphorylation opens the ATP binding pocket and substrate binding site, thereby enabling enzymatic activity (Nekhai et al., 2014; Russo et al., 1996). Therefore, we examined phospho-CDK9 (p-CDK9) as a surrogate of CDK9 activation where possible, alongside F-actin staining with phalloidin and the nucleus with DAPI.

In mouse orthochromatic erythroblasts p-CDK9 was observed in smaller clusters within the nucleus during the nuclear condensation phase, before localising to larger clusters within the cytoplasm that are also found in the future reticulocyte during extrusion; these clusters also seem to localise closely with F-actin during nuclear extrusion **(Figure 1B)**. p-CDK9 and CDK9 signal can be seen to co-localise, however total CDK9 protein is more spread throughout the reticulocyte **(Figure 1B)**. We then examined cyclin T1, the main binding partner and regulator of CDK9. Cyclin T1 protein was localised to the cytoplasm of mouse orthochromatic cells during nuclear condensation **(Figure 1C)**. Cyclin T1 remains primarily cytosolic before colocalising with p-CDK9 in clustered regions near F-actin during nuclear extrusion **(Figure 1C)**. These findings were recapitulated during enucleation in the human erythroblast cell line HUDEP-2 **(Figure 1D)**. Using p-CDK9 as a surrogate marker of CDK9 activity, we conclude CDK9 remains active in the nucleolus very late into terminal differentiation, and subsequently localises with F-actin and cyclin T at the at the rear of the extruding nucleus before relocating to the future reticulocyte in both mouse and human enucleating erythroblasts.

Highly selective inhibition of CDK9 activity arrests enucleation at the same stage as F-actin induced enucleation block
Our previous study revealed that pharmacological blockage of CDK9 activity leads to an arrest in enucleation, and considering the observed co-localisation of activated CDK9-Cyclin T1 with F-actin, we believed that CDK9 may be regulating the extrusion event. To examine this potential functional significance, we initially compared the effects of CDK9 inhibition with inhibition of actin polymerisation. With the advent of next generation high selectivity CDK9 inhibitors, we first confirmed the effect of specific CDK9 inhibition on enucleating cells in comparison to the F-actin inhibitor cytochalasin D (CytoD), which consistently arrests cells at a late-stage of enucleation (Konstantinidis et al., 2012; Wang et al., 2012; Wölwer et al., 2015) **(Figure 2)**. Treatment of orthochromatic mouse erythroblasts **(Figure 2A)**, with CDK9 inhibitors NVP-2 (Olson et al., 2017) **(Figure 2B)** and AZD4573 (Barlaam et al., 2020) **(Figure 2C)** arrested enucleation in a dose dependent manner, with a mean reduction of 51% and 64% against the vehicle (DMSO) control at 10 nM respectively. Phenotype analysis of cytospun cells reveal similar arrest patterns for both inhibitors consistent with the phenotypes reported previously for less specific CDK9 inhibitors (Wölwer et al., 2015), with an increased percentage of cells with polarised or partially extruded nucleus **(Figure 2B, C)** compared to the control **(Figure 2A)**. NVP-2 also reduced enucleation of human cells compared to the vehicle (DMSO) control, with a similar increase in polarised nuclei **(Supp. Figure 1A-1B)**. This data confirms that the action of CDK9 on erythroid enucleation is conserved between mouse and human, and further validates our previous findings by alleviating concerns around inhibitor off-target effects on other CDKs described here (Wölwer et al., 2015).

All other methods of chemical inhibition of CDK9 provided similar results. Atuveiclib (BAY 1143572) (Lücking et al., 2017), and the thalidomide-conjugate PROTAC degrader THAL-SNS-032 (Olson et al., 2017) both arrested enucleation compared to the vehicle (DMSO) control **(Supp. Figure 2)**. To compare the effects of CDK9 inhibition to a cytochalasin D (CytoD)-induced late-stage enucleation blockage, we treated mouse orthochromatic erythroblasts to a range of concentrations and quantified the phenotype of cytospins using morphology indicators **(Figure 2)**. Examination of cytospins indicate that CDK9 inhibitors phenocopy closely a blockage by the actin polymerisation inhibitor cytochalasin D (CytoD) **(Figure 2D)**, with quantitation of morphology indicating that cytochalasin D causes a slightly later arrest compared to CDK9 inhibitors, with more cells showing highly polarised and bulging nuclei **(Figure 2D)**. We conclude that CDK9 must act just prior to the final stages of enucleation, with the observed blockage closely resembling that of F-actin inhibition.

CDK9 is active upstream of calcium signalling and F-actin polymerisation prior to nuclear extrusion

Our localisation studies showed CDK9 co-localisation with F-actin at late stages of enucleation and suggested that CDK9 may be required just upstream of F-actin, and therefore may play a role in regulating the nuclear extrusion event. To confirm the order in which activated CDK9, and F-actin assembly may be required for enucleation, we first examined the reciprocal localisation of activated CDK9 and F-actin in the presence of NVP-2 and cytochalasin D (CytoD) in mouse orthochromatic erythroblasts using immunofluorescence imaging **(Figure 3A)**. Cytochalasin D (CytoD), which consistently arrests the cells with a partially extruded nucleus, resulted in accumulation of p-CDK9 at the site of enucleation **(Figure 3A)**. In addition, p-CDK9 signal is diminished in cells treated with NVP-2, as might be expected **(Figure 3A)**. F-actin appears to accumulate and co-localise with p-CDK9 in cytochalasin D (CytoD) treated cells but appears dispersed near the cell membrane in NVP-2 treated cells, where F-actin would normally localise toward the site of enucleation **(Figure 3A)**. Although these experiments indicate a close relationship between CDK9 and actin in the context of enucleation, they did not clarify the signalling hierarchy between CDK9 and actin.

Calcium signalling is required for effective enucleation, but the positioning of calcium signalling components is not known (Wölwer et al., 2016). We found that in orthochromatic erythroblasts calmodulin kinase II (CaMKII), a key component in CaM/Ca^{2+} signalling, localises primarily in the cytoplasm during nuclear polarisation, and co-localises with CDK9 at the rear of the nucleus during nuclear extrusion **(Figure 3B)**. To better observe the relationship between CDK9 and CaMKII, we examined their localisation in orthochromatic erythroblasts treated with either 100 nM NVP-2 or 5 µM KN-62, which inhibits CaMKII activity (Okazaki et al., 1994) **(Figure 3C)**. CDK9 and CaMKII appear to strongly colocalise under NVP-2 and KN-62 induced blockage, possibly suggesting the existence of a checkpoint dependent on the activity of CDK9 and/or CaMKII. However, we again observed an accumulation of CDK9 near the membrane opposite the polarised nucleus under KN-62 blockage **(Figure 3C)**, suggesting CaMKII signalling may be dependent on upstream CDK9 signalling to facilitate downstream F-actin-mediated nuclear extrusion.

To assess the order in which CDK9, actin and CaMKII act during enucleation, we applied reversible inhibitors in sequence **(Figure 3D-G)**. We reasoned that the kinetics of recovery would differ depending on whether downstream events were inhibited before or after upstream events. If before, they should not affect recovery, but if after, recovery would be slower. NVP-2 acts to inhibit CDK9 by reversibly blocking the ATP binding site (Olson et al., 2017), and cytochalasin D reversibly disrupts actin polymerisation by forcing actin dimerization (Friederich et al., 1993; Goddette & Frieden, 1986; Stevenson & Begg, 1994). Additionally, KN-62 can reversibly inhibit CaMKII (Okazaki et al., 1994). As NVP-2, cytochalasin D and KN-62 are reversible inhibitors, we used these inhibitors to study the

order of events prior to nuclear extrusion. A schematic describing the workflow of these experiments is provided **(Figure 3D)**. Enucleation rates were assessed by flow cytometry following 12 hours incubation with 1 μM cytochalasin D (CytoD) or 10 nM NVP-2 and compared to the vehicle (DMSO) control directly after washout, and at hour 14, 15 and 16 after plating cells **(Figure 3E)**. Enucleation rates following washout of cytochalasin D or NVP-2 into media containing DMSO carrier are shown to recover close to the rate of the DMSO vehicle control at hour 16, confirming the reversibility of both competitive inhibitors **(Figure 3E, Panel 1)**. Exchanging cytochalasin D (CytoD) for NVP-2 at the washout point resulted in a recovery of enucleation like the control, however, exchanging NVP-2 for cytochalasin D (CytoD) resulted in impaired recovery of enucleation to a statistically significant degree compared to both the control and the reverse inhibitor exchange at hour 16 **(Figure 3E, Panel 2)**. These experiments indicate that cells arrested during F-actin polymerisation inhibition by cytochalasin D are not affected by later CDK9 inhibition. Together these results indicate that CDK9 acts upstream of F-actin for nuclear extrusion.

Next, we examined the relationship between actin and CDK9 with CaMKII. Enucleation rates are shown to recover following washout of cytochalasin D (CytoD) and KN-62, with cytochalasin D (CytoD)-treated cells recovering with non-significant variance to the control, and KN-62 washout partially recovering, with enucleation rates still significantly lower than the control at hour 16 **(Figure 3F, Panel 1)**. Inhibition of F-actin (CytoD) after CaMKII signalling (KN-62) results in a complete recovery of enucleation rates by hour 16 compared to the control, but the reverse sequence did not **(Figure 3F, Panel 2)**. This indicates that CaMKII activity is required upstream of F-actin polymerisation prior to nuclear extrusion. Similarly, enucleation rates were assessed following 12 hours incubation with KN-62 or NVP-2 and compared to the vehicle (DMSO) control, where enucleation rates recovered equally in both the KN-62 and NVP-2 treatment to non-significantly different levels compared to the control at hour 16 **(Figure 3G, Panel 1)**. Although washout of KN-62 into NVP-2 allowed for a slight recovery of enucleation, it appeared that no sequence could be conferred, which may suggest that CDK9 and CaM signalling occurs concurrently or independently **(Figure 3G, Panel 2)**. Taken together, these results confirm that both CDK9 activity and CaM signalling is required before F-actin during enucleation.

CDK9 is associated with a Ran-NEMP1-Importin β complex in terminally differentiating erythroid cells

Having established that CDK9 acts closely with calcium signalling and upstream of F-actin in its regulation of enucleation, and that our previous work showed that the typical downstream effector of CDK9 being RNA Pol II is dispensable for enucleation (Wölwer et al., 2015), we wanted to assess how CDK9 might modulate actin activity and enucleation in general. To achieve this, we undertook a comprehensive proteomics screen to identify new interactors of CDK9 in the context of erythroid differentiation using CDK9 co-immunoprecipitation mass spectrometry. We used the human HUDEP-2 cell line, as this provided both the sufficient material and the ability to manipulate the system genetically. The results are summarised in **Figure 4**. First, we investigated the interactome of endogenous CDK9 in undifferentiated (Day 0) and differentiated (Day 6) HUDEP-2 cells. We used day 6 differentiated HUDEP-2 cells due to better cell viability and overall protein availability compared to more highly differentiated cells. The original protein database search is available in Supplementary File 2, however, for our analysis we took a stringent cut-off approach, including proteins that were identified in all 3 replicates and ≥3 fold enriched compared to the IgG co-immunoprecipitation controls. 65 unique proteins were identified in undifferentiated HUDEP-2 cells, and 114 unique proteins identified in differentiated HUDEP-2 cells, with 13 proteins identified in both data sets **(Figure 4A)**. We performed a STRING interactome analysis on these 13 proteins. Of note, we identified CDK9 and cyclin T1 (CCNT1) as common to both data sets, validating the presence of a CDK9-Cyclin T1 complex in

differentiating human erythroid cells **(Figure 4B)**. Other known interactors of CDK9, including vimentin (VIM) and the DNA-activated protein kinase subunit PRKDC were also found in both data sets (Yang et al., 2015; Zhang et al., 2014). Surprisingly, we also found 5 members of the Ran GTPase nuclear import complex, including nuclear envelope membrane protein 1 (NEMP1/TMEM194A) and importin β (KPNB1) **(Figure 4B)**. Of note, NEMP1/TMEM194A has recently been described as an essential regulator of enucleation in the mouse (Hodzic et al., 2022), confirming our proteomics screen using human cells can identify functionally relevant proteins for enucleation.

We next explored whether CDK9 activity impacted the interactome by comparing exogenous CDK9, a constitutively active kinase mutant CDK9-T186A, and a dominant negative kinase mutant CDK9-D168N (Dow et al., 2010). We isolated a mixed population of transduced cells by FACS **(Supp. Figure 3)**, and subsequently performed CDK9 co-immunoprecipitation mass spectrometry in day 6 differentiated cells. With the cut-off, described above, we identified a total of 243 proteins, with 98 uniquely identified in the CDK9 expression line, 65 unique to the CDK9-T186A line and 28 unique to the CDK9-D168N line, with 9 proteins identified in all 3 sets **(Figure 4C)**. A STRING analysis of these 9 intersecting proteins identified again 3 members of the Ran GTPase nuclear import complex, including importin β, confirming our previous results **(Figure 4D)**. We then performed STRING interactome analysis on all 243 identified proteins and clustered the results using k-means clustering to understand the variety of processes and pathways that CDK9 may regulate in differentiating human erythroblasts **(Supp. Figure 4)**. 11 clusters included processes related to Ran-mediated nuclear export and import (Cluster 1), minichromosomal maintenance complex (MCM) and DNA transcription (Cluster 2), actin filament capping and binding (Cluster 3), and as mRNA splicing and polyadenylation (Cluster 4). These four clusters were selected for further analysis based on overall higher abundance in mass spectrometry protein identification **(Figure 4E)**. The abundance of proteins in each cluster were compared across the CDK9, CDK9-T186A and CDK9-D168N datasets **(Figure 4E)**. Within the nuclear trafficking cluster (Cluster 1), importin β (KPNB1), which drives nuclear import, was enriched in the constitutively active kinase mutant (T186A), but exportin-1, which drives nuclear export, was depleted **(Figure 4E, Panel 1)**. This is compatible with the notion that CDK9 phosphorylation may regulate its nucleocytoplasmic shuttling as has been previously reported (Napolitano et al., 2002).

CDK9 and cyclin T1 were more abundant in the dominant negative (D168N) CDK9 mutant lines, indicative of a relationship between CDK9 phosphorylation state and turnover of CDK9 and cyclin T1 **(Figure 4E, Panel 2)**. Additionally, CDK9, cyclin T1 and other interactors within the minichromosomal maintenance (MCM) complex & DNA transcription cluster (Cluster 2) were enriched in the dominant negative kinase mutant (D168N), suggesting CDK9 kinase activity may impact the expression of various components involved in DNA transcription, including cyclin T1 and CDK9 itself **(Figure 4E, Panel 2)**. Interestingly, CDK9 has been shown to regulate DNA damage responses through MCM complex members through interaction with cyclin K (Yu & Cortez, 2011), which was not detected in our assay. CDK9 was found to interact with several components involved in actin filament capping & binding, which may carry implications for how CDK9 affects actin-related processes during enucleation **(Figure 4E, Panel 3)**. Finally, examination of the mRNA splicing & polyadenylation complex reveals shifts in how CDK9 kinase functionality may impact on its previously reported role in mRNA splicing **(Figure 4E, Panel 4)** (Hu et al., 2021). Here, our identification of a physical association between CDK9 and the Ran-Importin β nuclear import complex ties together our original finding of an RNA Pol II-independent role for CDK9, and recent findings that NEMP1 regulates nuclear envelope openings during erythroid enucleation.

Importin β is required for erythroid enucleation upstream from Calcium signalling and F-actin polymerisation

We next tested the function of importin β in the context of enucleation using the reversible inhibitor of importin β-mediated nuclear import, importazole (Soderholm et al., 2011). Inhibition of importin β using importazole arrested enucleation in a dose-dependent manner in mouse orthochromatic erythroblasts, demonstrating that importin β function is critical for enucleation **(Figure 5A)**. NEMP1, which has been implicated in maintain nuclear envelope openings which are essential for enucleation (Hodzic et al., 2022), was identified in complex with CDK9 and importin β, and so we aimed to assess the localisation of importin β during both normal and arrested enucleation as a comparison. Morphological analysis of orthochromatic erythroblasts treated with importazole indicated an increased number of cells arrested with a polarised nucleus **(Figure 5A)**, phenocopying both CDK9 and F-actin inhibitors and suggesting that importin β is acting prior to nuclear extrusion. Furthermore, inhibition of importin β supports the observed phenotype in NEMP1 knockout mice (Hodzic et al., 2022). Additionally, we treated day 12 differentiated HUDEP-2 cells with importazole for 24 hours and counted enucleated cells versus non-enucleated and found a significant and dose-dependent reduction in enucleation, indicating conserved function in human erythroid enucleation **(Figure 5B)**. To test whether importin β regulated the nucleocytoplasmic import of CDK9, we imaged fixed, undifferentiated HUDEP-2 cells following treatment with NVP-2, importazole or DMSO to assess the nuclear to cytoplasmic ratio of active CDK9, cyclin T1 and importazole. Blocking CDK9 activity did not alter the nucleocytoplasmic ratio of CDK9, cyclin T1 or importin β **(Supp. Figure 5)**. Interestingly, importazole significantly altered the localisation of cyclin T1, but not CDK9 **(Supp. Figure 5)**. This result, taken together with observed co-immunoprecipitation of CDK9 and importin β **(Figure 4)**, suggests that importin β may interact with CDK9 for reasons other than nucleocytoplasmic transport in the context of enucleation.

As importazole appeared to block enucleation at a similar stage to CDK9 and F-actin inhibitors, we assessed which of these proteins might regulate each other during enucleation. We first examined importin β localisation using fluorescence confocal microscopy. Importin β appears to localise to the nuclear envelope in pre-polarised erythroblasts with a similar phenotype reported for NEMP1 (Hodzic ref) **(Figure 5C)** and is observed in the cytoplasm and near folded regions at the nuclear envelope, before colocalising with p-CDK9 in the future reticulocyte during nuclear extrusion **(Figure 5C)**. Cells arrested by treatment of either cytochalasin D (CytoD) or importazole resulted in an accumulation of p-CDK9 at the site of enucleation, with importin-β localising around the membrane opposite to the polarised nucleus **(Figure 5D)**. Because p-CDK9 accumulates to the extrusion site in the same way under both actin and importin β inhibitors, we postulate that p-CDK9, similarly to actin, may be required for enucleation upstream of importin β before nuclear extrusion.

To test this functionally, we performed the previously described inhibitor washout, exchange, and recovery enucleation assays to determine an order of events using importazole, cytochalasin D (CytoD) and NVP-2. Enucleation rates were assessed by flow cytometry following 12 hours incubation with importazole, cytochalasin D (CytoD) or NVP-2 and compared to the vehicle (DMSO) control directly after washout, and at hour 14, 15 and 16 after plating cells **(Figure 5E, F)**. Enucleation rates in the vehicle (DMSO) treated control group remain steady, and enucleation rates following importazole washout were able to recover, confirming the reversibility of the blockage by importazole **(Figure 5E, Panel 1)**. Inhibition of importin β (importazole) before F-actin (CytoD) led to an initial recovery of enucleation rates but an overall reduction by hour 16 **(Figure 5E, Panel 2)**. Conversely, blocking F-actin (CytoD) before importin β (importazole) resulted in a gradual recovery of enucleation rates **(Figure 5E, Panel 2)**. This indicates that importin-β is required at an earlier stage of enucleation than F-actin polymerisation. Blocking importin β (importazole) before CDK9(NVP-2) resulted in a

consistent recovery of enucleation rates, but the reverse sequence did not **(Figure 5F)**. We repeated this by inhibiting CaMKII (KN-62) and importin β (importazole); however, we did observe altered recovery rates for KN-62 in this experiment, where recovery was significantly reduced compared to the control at hour 16 **(Figure 5G, Panel 1)**. Nonetheless, we could observe a clear shift in recovery when washing out importazole into KN-62, which resulted in a reduction of enucleation, as opposed to washout of KN-62 into importazole which allowed for a partial recovery in enucleation **(Figure 5G, Panel 2)**. Taken together, this supports our previous findings that CaMKII acts upstream of F-actin and indicates CaM signalling is necessary downstream of CDK9 but may occur dynamically or simultaneously to importin β activity. We can conclude that importin β is essential for enucleation downstream of CDK9, and upstream of F-actin, and that both importin β and CDK9 are required near the membrane opposite the polarised nucleus prior to nuclear extrusion. A proposed order of the molecular events regulating enucleation is summarised in **Figure 6**, alongside non-mutually exclusive potential models for the role of CDK9 and importin β in erythroid enucleation.

.

Discussion

Characterisation of the CDK9 interactome in terminally differentiating erythroblasts

This study provides new insights into the CDK9 interactome during erythroblast development **(Figure 4)**. Importantly, we identified several components of the Ran nuclear import pathway, including importin β, that are associated directly with CDK9 in both undifferentiated and differentiated HUDEP-2 cells. How may importin β and CDK9 coordinate nuclear extrusion? RanGTP acts as a master controller for nucleocytoplasmic transport through the nuclear pore complex (NPC) by interactions with importins and exportins (Cavazza & Vernos, 2016). During nuclear import, importin α will bind directly to nuclear localisation signals (NLS) on cargo and facilitate binding to RanGTP via importin β (Percipalle et al., 1997). CDK9 is known to shuttle between the nucleus and cytoplasm, and other interactors of CDK9 including cyclin T and HEXIM have been shown to interact with importin α to facilitate transport through the nuclear pore complex (Czudnochowski et al., 2010; Napolitano et al., 2002). CDK9 has previously been shown to interact with nuclear pore components RANGAP1, RANBP2 and RGPD3 (Ramakrishnan et al., 2012), which we also identified as part of the RanGTP complex that interacts with CDK9 in erythroid cells. However, the previous study only focused on localisation of CDK9-Cyclin T1 to the nuclear pore and did not examine the role of Ran and importin β in CDK9-Cyclin T1 nuclear import. Of note, exportin 7 (Xpo7), which can facilitate nuclear export through the RanGTP pathway, is essential for histone export into the cytoplasm during the nuclear condensation phase of enucleation (Hattangadi et al., 2014; Mingot et al., 2004). We were able to block enucleation by treating cells with importazole **(Figure 5)**, which can disrupt interactions between importin β and RanGTP, thereby inhibiting Ran-dependent nuclear import, without disrupting other pathways for nucleocytoplasmic transport (Soderholm et al., 2011). Whilst this implies that nuclear import mechanisms are essential for enucleation, phenotypic analysis of importazole-treated orthochromatic erythroblasts indicates a late-stage blockage in enucleation **(Figure 5)**, possibly suggesting a different role for importin β. It will be interesting to examine the relationship between the RanGTP pathway and exportin 7 (Xpo7) in the control of histone export and how CDK9 may be involved in this process. While this study offers some insight into the potential regulation of CDK9 and cyclin T1 nucleocytoplasmic transport via importin β interactions **(Supp. Figure 5)**, this should be further investigated in contexts outside of erythroid enucleation.

Besides nuclear import, importin β has been shown to modulate the permeability and structure of the nuclear pore complex (NPC) through direct interaction with nuclear pore complex protein Nup153 (Lowe et al., 2015), highlighting a potential role in modifying nuclear structure to facilitate enucleation. Additionally, importin β is implicated in wider transport roles, where it can regulate kinesin-4 motor activity in plant cells (Ganguly et al., 2018), and can directly bind to certain cargo, such as the parathyroid hormone (PTH)-related protein (PTHrP), to facilitate microtubule-dependent transport (Lam et al., 2002). Importantly, importin β is implicated in the regulation of anillin during cytokinesis (Beaudet et al., 2020); anillin acts to scaffold components of the actomyosin ring, including F-actin, myosin and RhoA (Piekny & Glotzer, 2008), all of which are essential for nuclear extrusion during enucleation. Examining whether anillin is required for enucleation and how this may relate to importin β function will be key to investigating this link. Finally, we observed nuclear envelope protein 1 (NEMP1/TMEM194A) across all sample sets, which forms part of the Ran nuclear import pathway, by recruiting Ran to the nuclear envelope (Shibano et al., 2015).

In addition to the Ran import pathway complex, we also identified several other new interactors of CDK9 **(Figure 4)**. Golgin A2 (GOLGA2), which was observed across all sample sets and was enriched in CDK9 pulldowns, acts to maintain Golgi structure through stacking of Golgi cisternae and has been investigated as a therapeutic target in cancer due to its relationship with autophagy (Chang et al., 2012;

Nakamura et al., 1995). As autophagy is essential for erythroid terminal differentiation, it will be important to examine the relationship between CDK9 and golgin A2-mediated autophagy mechanisms during enucleation. Vimentin (VIM), which is an intermediate filament protein implicated in the function of many cells types (Danielsson et al., 2018) was also identified across all sample sets, and has been previously identified in a similar CDK9 immunoprecipitation mass spectrometry interactome set (Yang et al., 2015). Yang et al. (2015) identified a CDK9 interactome in human A549 pulmonary epithelial cells, and some of these proteins are also identified in our data (Yang et al., 2015). However, most proteins in our identified interactome are new, likely reflecting the specialised role CDK9 may play in erythroid development.

Of note, we identified a large complex involved in minichromosome maintenance (MCM) and DNA damage response. Although CDK9 has been shown to play a role in maintaining genome integrity in partnership with cyclin K (Yu & Cortez, 2011), we did not detect cyclin K in our assay, and it is therefore possible that this mechanism extends to erythroblasts through a different cyclin. This should be investigated further. Furthermore, a role for CDK9 in mRNA splicing has been described (Hu et al., 2021), and with the identification of mRNA splicing mechanisms, polyadenylation factors and several members of the messenger ribonucleoprotein complex assembly, also known as mRNP granules (Buchan, 2014) in human erythroblasts, new insights could be gained on the role of CDK9 in mRNA processing in erythropoiesis. Importantly, we also identified a large complex involved in actin filament capping and binding, including CAPZA1/2, CAPZB, amongst others, which may point to how CDK9 might interact with actin during enucleation processes. Additionally, we identified tropomodulin-1, which has been observed to co-localise with F-actin and non-musical myosin IIB during enucleation (Nowak et al., 2017), and Cdc42, which regulates essential polarisation of the nucleus prior to nuclear extrusion (Ubukawa et al., 2020). The relationship between CDK9 and the regulation of actinomyosin polymerisation should be investigated further, using genetic models for proximity labelling and functional assays to determine how CDK9 activity may be involved in modulating actinomyosin activity in the context of erythroid terminal differentiation.

CDK9 and importin β facilitate cellular reorganisation prior to CaM/Ca^{2+} signalling and F-activity to achieve enucleation

Our previous work had identified CDK9 as a regulator of enucleation (Wölwer et al., 2015), and this study aimed to delineate the mechanisms involved. Here, we propose three non-mutually exclusive models for the role of CDK9 in addition to roles for the newly identified regulator of enucleation importin β **(Figure 6)**. The typical biological function of CDK9 is well-characterised, where it acts as the kinase component and small subunit of P-TEFb, the major regulator of RNA polymerase II (RNA Pol II) transcriptional elongation (Anshabo et al., 2021; Paparidis et al., 2017). P-TEFb phosphorylates the RNA Pol II carboxyl terminal domain (CTD) directly, in addition to two negative regulators of RNA Pol II activity, DRB sensitivity-inducing factor (DSIF) and negative elongation factor (NELF) (Gilchrist et al., 2008; Wada et al., 1998). Our previous study had shown that blocking the activity of RNA Pol II in late erythropoiesis does not arrest enucleation (Wölwer et al., 2015), suggesting a new role for CDK9 in enucleation independently of RNA Pol II. Neither DSIF or NELF have been shown to interact with or modulate the activity of anything other than RNA Pol II, therefore we did not investigate their potential role in enucleation but at this stage cannot rule out a possible involvement (Aoi et al., 2020; Decker, 2021; Deng et al., 2022). Considering the localisation of CDK9 during enucleation and the phenotype of CDK9 inhibitor-induced arrest, we conclude that CDK9 must be acting within the cytoplasm prior to nuclear extrusion, however it remains plausible that CDK9-mediated transcription via RNA Pol II is essential for enucleation, and importin β may play a role in the nucleocytoplasmic location of CDK9 **(Figure 6B: Model 1)**.

NEMP1, which is implicated in enucleation, functions to modulate nuclear envelope openings that are essential for enucleation, and NEMP1 knockout mice show anaemia due to apoptosis of polychromatic erythroblasts (Hodzic et al., 2022). Considering both NEMP1 and importin β are key components of the Ran nuclear import pathway (Shibano et al., 2015), it is likely that the function of importin β in enucleation is related to that of NEMP1. Interestingly, NEMP1 has been shown to support the mechanical stiffness of the nuclear envelope in some settings (Tsatskis et al., 2020), potentially implicating NEMP1 (and by association importin β) in the mechanical processes involved in extrusion of the nucleus during enucleation. Considering CDK9 and importin β were found to physically interact with NEMP1 in the context of erythropoiesis, we postulate that importin β, NEMP1 and possibly other members of the Ran nucleocytoplasmic import complex may regulate changes to the nuclear envelope to assist in the shuttling of key enucleation regulators, including CDK9, between the cytoplasm and nucleus prior to nuclear extrusion **(Figure 6B, Model 2)**.

Finally, our previous work has identified a role for calcium signalling in enucleation via the calmodulin (CaM) pathway, and the role of F-actin is well documented (Newton et al., 2024; Wölwer et al., 2016). More recently, Zhang et al (2015) describe and support the notion of finely tuned CaM/Ca2+ signalling during erythropoiesis and enucleation (Zhang et al., 2025). Here, we examined the order of which CDK9 and importin β may act to regulate enucleation in respect to calcium signalling and F-actin activity and found both to function upstream **(Figure 6A)**. The calmodulin (CaM) pathway is implicated in both myosin and F-actin activation, through the activity of myosin light chain kinase (MLCK) and Ca^{2+}/CaM-dependent kinase II (CaMKII) respectively (Mizuno et al., 2008; Okamoto et al., 2007). Calmodulin signalling has been shown to regulate CDK9 T-loop phosphorylation, which is required for its activation, which may suggest a coordinated role for calmodulin signalling in separate steps in enucleation involving CDK9 and the cytoskeleton (Ramakrishnan & Rice, 2012). Additionally, calcium signalling, and intracellular calcium levels have been shown to block nuclear import by altering the activity and localisation of RanGTP and importin β (Kaur et al., 2014; Sweitzer & Hanover, 1996). Strikingly, blocking enucleation with an actin inhibitor or importin β inhibitor both caused a reproducible and consistent accumulation of phosphorylated CDK9 near the junction of enucleation. This may be due to the aggregation of activated CDK9 at a critical site of activity, therefore priming the cells for nuclear extrusion following release of F-actin or importin β inhibition. Conversely, blocking CDK9 activity with NVP-2 appears to result in diminished p-CDK9, potentially due to blocking CDK9 autophosphorylation, which when recovered after washout still requires F-actin polymerisation to achieve nuclear extrusion. We propose that CDK9 and importin β may function as part of a checkpoint in the final stages of enucleation, in conjunction with an influx of calcium and subsequent calmodulin (CaM) signalling to achieve nuclear extrusion **(Figure 6B, Model 3)**.

To fully understand how CDK9 and importin β work to drive erythroid enucleation, the role of Ran GTPase nuclear import should be further examined, using genetic models and live imaging to decipher how importin β, NEMP1 and other components regulate the movement of molecules between the cytoplasm and nucleus prior to enucleation, or act to drive essential changes to the nuclear pore complex or envelope. Although we don't fully understand how CDK9, importin β and downstream effectors of enucleation directly interact to coordinate enucleation, our studies provide insight into a series of events that must occur before F-actin mediated nuclear extrusion and presents possible new interactors of CDK9 and importin β in the context of erythropoiesis and erythroid enucleation.

Acknowledgements

This work was completed on Wurundjeri land of the Kulin nation, and we pay our respects to the Elders past and present. We acknowledge Dr. Margaret Veale and the La Trobe Bioimaging Platform for flow cytometry and optical microscopy. We acknowledge Dr. Pierre Faou, Dr. Rohan Lowe, Dr. Keshava Datta and the La Trobe Proteomics and Metabolomics Platform for mass spectrometry and proteomics. We acknowledge the La Trobe Animal Research & Training Facility (LARTF) for supporting animal work. We thank Dr. Ryo Kurita from the Riken BioResource Research Center (Tsukuba, Japan) for providing HUDEP-2 cell lines, and Prof. Jan Frayne from the University of Bristol (Bristol, United Kingdom) for providing BEL-A cell lines. LMN was supported by an Australian Government Research Training Program (RTP) scholarship. POH was supported by NHMRC Fellowship APP1079133. This work was supported by NHMRC Project grant APP1103858 (POH) and ARC Discovery project grant DP190103634 (POH and EDH).

Authorship

Contributions: POH conceived of the project. LMN designed and performed experiments. LMN, KYBL, DYA, CBW, EDH and POH contributed to data analysis and optimisation of experimental protocols. SMR provided resources. CJ contributed to image acquisition and analysis. LMN, SMR and POH drafted the manuscript. All authors contributed to editing the manuscript. EDH and POH supervised the study.

ORCID profiles: LMN, 0000-0001-5984-5778; KYBL, 0000-0002-4008-6529; DYB, 0009-0005-0686-0495; CBW, 0000-0003-1190-2408; SMR, 0000-0001-5826-9641; EDH, 0000-0002-3686-8261; POH, 0000-0002-1366-6691.

Conflict of interest disclosure: There are no conflicts of interest to disclose.

Correspondence: Patrick Humbert, Department of Biochemistry and Chemistry, La Trobe Institute for Molecular Science, La Trobe University, Kingsbury Drive, Bundoora, Victoria, Australia, 3073; email: p.humbert@latrobe.edu.au.

References

An, C., Huang, Y., Li, M., Xue, F., Nie, D., Zhao, H., Chen, L., Yazdanbakhsh, K., Sun, L., Jiang, Z., Mohandas, N., & An, X. (2021). Vesicular formation regulated by ERK/MAPK pathway mediates human erythroblast enucleation. *Blood Advances*, *5*(22), 4648-4661. https://doi.org/10.1182/BLOODADVANCES.2021004859

Anshabo, A. T., Milne, R., Wang, S., & Albrecht, H. (2021). CDK9: A Comprehensive Review of Its Biology, and Its Role as a Potential Target for Anti-Cancer Agents. *Frontiers in oncology*, *11*. https://doi.org/10.3389/FONC.2021.678559

Antonicka, H., Lin, Z.-Y., Janer, A., Aaltonen, M. J., Weraarpachai, W., Gingras, A.-C., & Shoubridge, E. A. (2020). A High-Density Human Mitochondrial Proximity Interaction Network. *Cell Metabolism*, *32*(3), 479-497.e479. https://doi.org/10.1016/j.cmet.2020.07.017

Aoi, Y., Smith, E. R., Shah, A. P., Rendleman, E. J., Marshall, S. A., Woodfin, A. R., Chen, F. X., Shiekhattar, R., & Shilatifard, A. (2020). NELF Regulates a Promoter-Proximal Step Distinct from RNA Pol II Pause-Release. *Molecular cell*, *78*(2), 261-274.e265. https://doi.org/10.1016/J.MOLCEL.2020.02.014

Barlaam, B., Casella, R., Cidado, J., Cook, C., De Savi, C., Dishington, A., Donald, C. S., Drew, L., Ferguson, A. D., Ferguson, D., Glossop, S., Grebe, T., Gu, C., Hande, S., Hawkins, J., Hird, A. W., Holmes, J., Horstick, J., Jiang, Y., . . . Yao, T. (2020). Discovery of AZD4573, a Potent and Selective Inhibitor of CDK9 That Enables Short Duration of Target Engagement for the Treatment of Hematological Malignancies. *Journal of medicinal chemistry*, *63*(24), 15564-15590. https://doi.org/10.1021/ACS.JMEDCHEM.0C01754

Beaudet, D., Pham, N., Skaik, N., & Piekny, A. (2020). Importin binding mediates the intramolecular regulation of anillin during cytokinesis. *Molecular Biology of the Cell*, *31*(11), 1124-1139. https://doi.org/10.1091/MBC.E20-01-0006/ASSET/IMAGES/LARGE/MBC-31-1124-G006.JPEG

Buchan, J. R. (2014). mRNP granules. Assembly, function, and connections with disease. *RNA biology*, *11*(8), 1019-1030. https://doi.org/10.4161/15476286.2014.972208

Cavazza, T., & Vernos, I. (2016). The RanGTP pathway: From nucleo-cytoplasmic transport to spindle assembly and beyond. *Frontiers in Cell and Developmental Biology*, *3*(JAN), 82. https://doi.org/10.3389/FCELL.2015.00082/BIBTEX

Chang, S.-H., Hong, S.-H., Jiang, H.-L., Minai-Tehrani, A., Yu, K.-N., Lee, J.-H., Kim, J.-E., Shin, J.-Y., Kang, B., Park, S., Han, K., Chae, C., & Cho, M.-H. (2012). GOLGA2/GM130, cis-Golgi Matrix Protein, is a Novel Target of Anticancer Gene Therapy. *Molecular Therapy*, *20*(11), 2052-2063. https://doi.org/10.1038/mt.2012.125

Christaki, E.-E., Politou, M., Antonelou, M., Athanasopoulos, A., Simantirakis, E., Seghatchian, J., & Vassilopoulos, G. (2019). Ex vivo generation of transfusable red blood cells from various stem cell sources: A concise revisit of where we are now. *Transfusion and Apheresis Science*, *58*(1), 108-112. https://doi.org/https://doi.org/10.1016/j.transci.2018.12.015

Czudnochowski, N., Vollmuth, F., Baumann, S., Vogel-Bachmayr, K., & Geyer, M. (2010). Specificity of Hexim1 and Hexim2 complex formation with cyclin T1/T2, importin alpha and 7SK snRNA. *Journal of Molecular Biology*, *395*(1), 28-41. https://doi.org/10.1016/J.JMB.2009.10.055

Danielsson, F., Peterson, M. K., Caldeira Araújo, H., Lautenschläger, F., & Gad, A. K. B. (2018). Vimentin Diversity in Health and Disease. *Cells*, *7*(10), 147. https://doi.org/10.3390/cells7100147

Decker, T. M. (2021). Mechanisms of Transcription Elongation Factor DSIF (Spt4-Spt5). *Journal of Molecular Biology*, *433*(14). https://doi.org/10.1016/J.JMB.2020.09.016

Deng, N., Zhang, Y., Ma, Z., Lin, R., Cheng, T. H., Tang, H., Snyder, M. P., & Cohen, S. N. (2022). DSIF modulates RNA polymerase II occupancy according to template G + C content. *NAR genomics and bioinformatics*, *4*(3). https://doi.org/10.1093/NARGAB/LQAC054

Dow, E. C., Liu, H., & Rice, A. P. (2010). T-loop phosphorylated Cdk9 localizes to nuclear speckle domains which may serve as sites of active P-TEFb function and exchange between the Brd4 and 7SK/HEXIM1 regulatory complexes. *Journal of Cellular Physiology*, *224*(1), 84-93. https://doi.org/10.1002/jcp.22096

Fonseka, P., Pathan, M., Chitti, S. V., Kang, T., & Mathivanan, S. (2021). FunRich enables enrichment analysis of OMICs datasets. *Journal of Molecular Biology*, *433*(11), 166747. https://doi.org/10.1016/j.jmb.2020.166747

Friederich, E., Kreis, T. E., & Louvard, D. (1993). Villin-induced growth of microvilli is reversibly inhibited by cytochalasin D. *Journal of Cell Science*, *105 (Pt 3)*, 765-775. https://doi.org/10.1242/jcs.105.3.765

Gallego-Murillo, J. S., Iacono, G., van der Wielen, L. A. M., van den Akker, E., von Lindern, M., & Wahl, S. A. (2022). Expansion and differentiation of ex vivo cultured erythroblasts in scalable stirred bioreactors. *Biotechnology and bioengineering*. https://doi.org/10.1002/BIT.28193

Ganguly, A., DeMott, L., Zhu, C., McClosky, D. D., Anderson, C. T., & Dixit, R. (2018). Importin-β Directly Regulates the Motor Activity and Turnover of a Kinesin-4. *Developmental Cell*, *44*(5), 642-651.e645. https://doi.org/10.1016/J.DEVCEL.2018.01.027

Gilchrist, D. A., Nechaev, S., Lee, C., Ghosh, S. K. B., Collins, J. B., Li, L., Gilmour, D. S., & Adelman, K. (2008). NELF-mediated stalling of Pol II can enhance gene expression by blocking promoter-proximal nucleosome assembly. *Genes & Development*, *22*(14), 1921-1933. https://doi.org/10.1101/gad.1643208

Goddette, D. W., & Frieden, C. (1986). Actin polymerization. The mechanism of action of cytochalasin D. *The Journal of Biological Chemistry*, *261*(34), 15974-15980. http://www.ncbi.nlm.nih.gov/pubmed/3023337

Hattangadi, S. M., Martinez-Morilla, S., Patterson, H. C., Shi, J., Burke, K., Avila-Figueroa, A., Venkatesan, S., Wang, J., Paulsen, K., Görlich, D., Murata-Hori, M., & Lodish, H. F. (2014). Histones to the cytosol: exportin 7 is essential for normal terminal erythroid nuclear maturation. *Blood*, *124*(12), 1931-1940. https://doi.org/10.1182/BLOOD-2013-11-537761

Hodzic, D., Wu, J., Krchma, K., Jurisicova, A., Tsatskis, Y., Liu, Y., Ji, P., Choi, K., & McNeill, H. (2022). The inner nuclear membrane protein NEMP1 supports nuclear envelope openings and enucleation of erythroblasts. *PLOS Biology*, *20*(10), e3001811. https://doi.org/10.1371/journal.pbio.3001811

Hoffman, L., Farley, M. M., & Waxham, M. N. (2013). Calcium-calmodulin-dependent protein kinase II isoforms differentially impact the dynamics and structure of the actin cytoskeleton. *Biochemistry*, *52*(7), 1198-1207. https://doi.org/10.1021/BI3016586/SUPPL_FILE/BI3016586_SI_005.PDF

Hu, Q., Poulose, N., Girmay, S., Helevä, A., Doultsinos, D., Gondane, A., Steele, R. E., Liu, X., Loda, M., Liu, S., Tang, D. G., Mills, I. G., & Itkonen, H. M. (2021). Inhibition of CDK9 activity compromises global splicing in prostate cancer cells. *RNA biology*, *18*(sup2), 722-729. https://doi.org/10.1080/15476286.2021.1983287

Ji, P., Jayapal, S. R., & Lodish, H. F. (2008). Enucleation of cultured mouse fetal erythroblasts requires Rac GTPases and mDia2. *Nature Cell Biology*, *10*(3), 314-321. https://doi.org/10.1038/ncb1693

Jordan, M., Schallhorn, A., & Wurm, F. M. (1996). Transfecting Mammalian Cells: Optimization of Critical Parameters Affecting Calcium-Phosphate Precipitate Formation. *Nucleic Acids Research*, *24*(4), 596-601. https://doi.org/10.1093/nar/24.4.596

Kaur, G., Ly-Huynh, J. D., & Jans, D. A. (2014). Intracellular calcium levels can regulate Importin-dependent nuclear import. *Biochemical and Biophysical Research Communications*, *450*(1), 812-817. https://doi.org/10.1016/J.BBRC.2014.06.047

Keerthivasan, G., Small, S., Liu, H., Wickrema, A., & Crispino, J. D. (2010). Vesicle trafficking plays a novel role in erythroblast enucleation. *Blood*, *116*(17), 3331-3340. https://doi.org/10.1182/blood-2010-03-277426

Konstantinidis, D. G., Pushkaran, S., Johnson, J. F., Cancelas, J. A., Manganaris, S., Harris, C. E., Williams, D. A., Zheng, Y., & Kalfa, T. A. (2012). Signaling and cytoskeletal requirements in erythroblast enucleation. *Blood*, *119*(25), 6118-6127. https://doi.org/10.1182/BLOOD-2011-09-379263

Koury, S. T., Koury, M. J., & Bondurant, M. C. (1989). Cytoskeletal distribution and function during the maturation and enucleation of mammalian erythroblasts. *Journal of Cell Biology*, *109*(6 I), 3005-3013. https://doi.org/10.1083/jcb.109.6.3005

Lam, M. H. C., Thomas, R. J., Loveland, K. L., Schilders, S., Gu, M., Martin, T. J., Gillespie, M. T., & Jans, D. A. (2002). Nuclear transport of parathyroid hormone (PTH)-related protein is dependent on microtubules. *Molecular endocrinology (Baltimore, Md.)*, *16*(2), 390-401. https://doi.org/10.1210/MEND.16.2.0775

Li, X., Mei, Y., Yan, B., Vitriol, E., Huang, S., Ji, P., & Qiu, Y. (2017). Histone deacetylase 6 regulates cytokinesis and erythrocyte enucleation through deacetylation of formin protein mDia2. *Haematologica*, *102*(6), 984-994. https://doi.org/10.3324/haematol.2016.161513

Liang, R., Campreciós, G., Kou, Y., McGrath, K., Nowak, R., Catherman, S., Bigarella, C. L., Rimmelé, P., Zhang, X., Gnanapragasam, M. N., Bieker, J. J., Papatsenko, D., Ma'ayan, A., Bresnick, E., Fowler, V., Palis, J., & Ghaffari, S. (2015). A Systems Approach Identifies Essential FOXO3 Functions at Key Steps of Terminal Erythropoiesis. *PLOS Genetics*, *11*(10), 1-27. https://doi.org/10.1371/journal.pgen.1005526

Lowe, A. R., Tang, J. H., Yassif, J., Graf, M., Huang, W. Y. C., Groves, J. T., Weis, K., & Liphardt, J. T. (2015). Importin-β modulates the permeability of the nuclear pore complex in a Ran-dependent manner. *eLife*, *4*(4). https://doi.org/10.7554/ELIFE.04052

Lücking, U., Scholz, A., Lienau, P., Siemeister, G., Kosemund, D., Bohlmann, R., Briem, H., Terebesi, I., Meyer, K., Prelle, K., Denner, K., Bömer, U., Schäfer, M., Eis, K., Valencia, R., Ince, S., von Nussbaum, F., Mumberg, D., Ziegelbauer, K., . . . Brands, M. (2017). Identification of Atuveciclib (BAY 1143572), the First Highly Selective, Clinical PTEFb/CDK9 Inhibitor for the Treatment of Cancer. *Chemmedchem*, *12*(21), 1776-1793. https://doi.org/10.1002/cmdc.201700447

Menon, V., & Ghaffari, S. (2021). Erythroid enucleation: a gateway into a "bloody" world. *Experimental Hematology*, *95*, 13-22. https://doi.org/10.1016/J.EXPHEM.2021.01.001

Mingot, J. M., Bohnsack, M. T., Jäkle, U., & Görlich, D. (2004). Exportin 7 defines a novel general nuclear export pathway. *The EMBO Journal*, *23*(16), 3227. https://doi.org/10.1038/SJ.EMBOJ.7600338

Mizuno, Y., Isotani, E., Huang, J., Ding, H., Stull, J. T., & Kamm, K. E. (2008). Myosin light chain kinase activation and calcium sensitization in smooth muscle in vivo. *American journal of physiology. Cell physiology*, *295*(2). https://doi.org/10.1152/AJPCELL.90645.2007

Moras, M., Lefevre, S. D., & Ostuni, M. A. (2017). From erythroblasts to mature red blood cells: Organelle clearance in mammals. *Frontiers in Physiology*, *8*(DEC), 1-9. https://doi.org/10.3389/fphys.2017.01076

Nakamura, N., Rabouille, C., Watson, R., Nilsson, T., Hui, N., Slusarewicz, P., Kreis, T. E., & Warren, G. (1995). Characterization of a cis-Golgi matrix protein, GM130. *The Journal of Cell Biology*, *131*(6 Pt 2), 1715-1726. https://doi.org/10.1083/jcb.131.6.1715

Napolitano, G., Licciardo, P., Carbone, R., Majello, B., & Lania, L. (2002). CDK9 has the intrinsic property to shuttle between nucleus and cytoplasm, and enhanced expression of cyclin T1 promotes its nuclear localization. *Journal of cellular physiology*, *192*(2), 209-215. https://doi.org/10.1002/JCP.10130

Nekhai, S., Petukhov, M., & Breuer, D. (2014). Regulation of CDK9 Activity by Phosphorylation and Dephosphorylation. *BioMed Research International*, *2014*, 964964. https://doi.org/10.1155/2014/964964

Newton, L. M., Fowler, V. M., & Humbert, P. O. (2024). Erythroblast enucleation at a glance. *Journal of Cell Science*, *137*(19). https://doi.org/10.1242/jcs.261673

Nowak, R. B., Papoin, J., Gokhin, D. S., Casu, C., Rivella, S., Lipton, J. M., Blanc, L., & Fowler, V. M. (2017). Tropomodulin 1 controls erythroblast enucleation via regulation of F-actin in the enucleosome. *Blood*, *130*(9), 1144-1155. https://doi.org/10.1182/BLOOD-2017-05-787051

Ohta, Y., Nishida, E., & Sakai, H. (1986). Type II Ca2+/calmodulin-dependent protein kinase binds to actin filaments in a calmodulin-sensitive manner. *FEBS letters*, *208*(2), 423-426. https://doi.org/10.1016/0014-5793(86)81061-4

Okamoto, K. I., Narayanan, R., Lee, S. H., Murata, K., & Hayashi, Y. (2007). The role of CaMKII as an F-actin-bundling protein crucial for maintenance of dendritic spine structure. *Proceedings of the National Academy of Sciences of the United States of America*, *104*(15), 6418-6423. https://doi.org/10.1073/PNAS.0701656104/ASSET/28E890F6-80E2-4EEE-A931-8B6792E74485/ASSETS/GRAPHIC/ZPQ0150759230006.JPEG

Okazaki, K., Ishikawa, T., Inui, M., Tada, M., Goshima, K., Okamoto, T., & Hidaka, H. (1994). KN-62, a specific Ca++/calmodulin-dependent protein kinase inhibitor, reversibly depresses the rate of beating of cultured fetal mouse cardiac myocytes. *The Journal of Pharmacology and Experimental Therapeutics*, *270*(3), 1319-1324. http://www.ncbi.nlm.nih.gov/pubmed/7932185

Olson, C. M., Jiang, B., Erb, M. A., Liang, Y., Doctor, Z. M., Zhang, Z., Zhang, T., Kwiatkowski, N., Boukhali, M., Green, J. L., Haas, W., Nomanbhoy, T., Fischer, E. S., Young, R. A., Bradner, J. E., Winter, G. E., & Gray, N. S. (2017). Pharmacological perturbation of CDK9 using selective CDK9 inhibition or degradation. *Nature Chemical Biology 2017 14:2*, *14*(2), 163-170. https://doi.org/10.1038/nchembio.2538

Palis, J. (2014). Primitive and definitive erythropoiesis in mammals. *Frontiers in Physiology*, *5*. https://doi.org/10.3389/fphys.2014.00003

Paparidis, N. F. d. S., Durvale, M. C., & Canduri, F. (2017). The emerging picture of CDK9/P-TEFb: more than 20 years of advances since PITALRE. *Molecular BioSystems*, *13*(2), 246-276. https://doi.org/10.1039/c6mb00387g

Percipalle, P., Clarkson, W. D., Kent, H. M., Rhodes, D., & Stewart, M. (1997). Molecular interactions between the importin alpha/beta heterodimer and proteins involved in vertebrate nuclear protein import. *Journal of Molecular Biology*, *266*(4), 722-732. https://doi.org/10.1006/jmbi.1996.0801

Piekny, A. J., & Glotzer, M. (2008). Anillin Is a Scaffold Protein That Links RhoA, Actin, and Myosin during Cytokinesis. *Current Biology*, *18*(1), 30-36. https://doi.org/10.1016/J.CUB.2007.11.068

Qin, W., Eick, D., Schwalb, B., Leonhardt, H., Decker, T. M., Gressel, S., & Cramer, P. (2017). CDK9-dependent RNA polymerase II pausing controls transcription initiation. *eLife*, *6*, 1-24. https://doi.org/10.7554/elife.29736

Ramakrishnan, R., Liu, H., Donahue, H., Malovannaya, A., Qin, J., & Rice, A. P. (2012). Identification of novel CDK9 and Cyclin T1-associated protein complexes (CCAPs) whose siRNA depletion enhances HIV-1 Tat function. *Retrovirology*, *9*(1), 90. https://doi.org/10.1186/1742-4690-9-90

Ramakrishnan, R., & Rice, A. P. (2012). Cdk9 T-loop phosphorylation is regulated by the calcium signaling pathway. *Journal of cellular physiology*, *227*(2), 609-617. https://doi.org/10.1002/JCP.22760

Rogerson, C., Bergamaschi, D., & O'shaughnessy, R. F. L. (2018). Uncovering mechanisms of nuclear degradation in keratinocytes: A paradigm for nuclear degradation in other tissues. *Nucleus (Austin, Tex.)*, *9*(1), 56-64. https://doi.org/10.1080/19491034.2017.1412027

Russo, A. A., Jeffrey, P. D., & Pavletich, N. P. (1996). Structural basis of cyclin-dependent kinase activation by phosphorylation. *Nature Structural Biology*, *3*(8), 696-700. https://doi.org/10.1038/nsb0896-696

Sanabria, H., Swulius, M. T., Kolodziej, S. J., Liu, J., & Waxham, M. N. (2009). βCaMKII Regulates Actin Assembly and Structure. *The Journal of Biological Chemistry*, *284*(15), 9770. https://doi.org/10.1074/JBC.M809518200

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature Methods*, *9*(7), 676-682. https://doi.org/10.1038/nmeth.2019

Shibano, T., Mamada, H., Hakuno, F., Takahashi, S.-I., & Taira, M. (2015). The Inner Nuclear Membrane Protein Nemp1 Is a New Type of RanGTP-Binding Protein in Eukaryotes. *PLOS ONE*, *10*(5), e0127271. https://doi.org/10.1371/journal.pone.0127271

Smith, A. S., Nowak, R. B., & Fowler, V. M. (2018). High-resolution fluorescence microscope imaging of erythroblast structure. *Methods in Molecular Biology*, *1698*, 205-228. https://doi.org/10.1007/978-1-4939-7428-3_12/COVER

Soderholm, J. F., Bird, S. L., Kalab, P., Sampathkumar, Y., Hasegawa, K., Uehara-Bingen, M., Weis, K., & Heald, R. (2011). Importazole, a small molecule inhibitor of the transport receptor importin-β. *ACS Chemical Biology*, *6*(7), 700-708. https://doi.org/10.1021/cb2000296

Stevenson, B. R., & Begg, D. A. (1994). Concentration-dependent effects of cytochalasin D on tight junctions and actin filaments in MDCK epithelial cells. *Journal of Cell Science*, *107 (Pt 3)*, 367-375. https://doi.org/10.1242/jcs.107.3.367

Sweitzer, T. D., & Hanover, J. A. (1996). Calmodulin activates nuclear protein import: A link between signal transduction and nuclear transport. *Proceedings of the National Academy of Sciences of the United States of America*, *93*(25), 14574-14579. https://doi.org/10.1073/PNAS.93.25.14574/ASSET/E4E6112C-EF89-4990-8ADF-A083A90B42B9/ASSETS/GRAPHIC/PQ2462935004.JPEG

Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable, A. L., Fang, T., Doncheva, N. T., Pyysalo, S., Bork, P., Jensen, L. J., & von Mering, C. (2023). The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids Research*, *51*(D1), D638-D646. https://doi.org/10.1093/nar/gkac1000

The UniProt Consortium. (2023). UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Research*, *51*(D1), D523-D531. https://doi.org/10.1093/nar/gkac1052

Trakarnsanga, K., Griffiths, R. E., Wilson, M. C., Blair, A., Satchwell, T. J., Meinders, M., Cogan, N., Kupzig, S., Kurita, R., Nakamura, Y., Toye, A. M., Anstee, D. J., & Frayne, J. (2017). An immortalized adult human erythroid line facilitates sustainable and scalable generation of functional red cells. *Nature Communications 2017 8:1*, *8*(1), 1-7. https://doi.org/10.1038/ncomms14750

Tsatskis, Y., Rosenfeld, R., Pearson, J. D., Boswell, C., Qu, Y., Kim, K., Fabian, L., Mohammad, A., Wang, X., Robson, M. I., Krchma, K., Wu, J., Gonçalves, J., Hodzic, D., Wu, S., Potter, D., Pelletier, L., Dunham, W. H., Gingras, A.-C., . . . McNeill, H. (2020). The NEMP family supports metazoan fertility and nuclear envelope stiffness. *Science Advances*, *6*(35), eabb4591. https://doi.org/10.1126/sciadv.abb4591

Ubukawa, K., Goto, T., Asanuma, K., Sasaki, Y., Guo, Y. M., Kobayashi, I., Sawada, K., Wakui, H., & Takahashi, N. (2020). Cdc42 regulates cell polarization and contractile actomyosin rings during terminal differentiation of human erythroblasts. *Scientific Reports 2020 10:1*, *10*(1), 1-11. https://doi.org/10.1038/s41598-020-68799-1

Wada, T., Takagi, T., Yamaguchi, Y., Watanabe, D., & Handa, H. (1998). Evidence that P-TEFb alleviates the negative effect of DSIF on RNA polymerase II-dependent transcription in vitro. *The EMBO journal*, *17*(24), 7395-7403. https://doi.org/10.1093/emboj/17.24.7395

Wang, J., Ramirez, T., Ji, P., Jayapal, S. R., Lodish, H. F., & Murata-Hori, M. (2012). Mammalian erythroblast enucleation requires PI3K-dependent cell polarization. *Journal of Cell Science*, *125*(Pt 2), 340-349. https://doi.org/10.1242/JCS.088286

Wölwer, C. B., Pase, L. B., Pearson, H. B., Gödde, N. J., Lackovic, K., Huang, D. C. S., Russell, S. M., & Humbert, P. O. (2015). A Chemical Screening Approach to Identify Novel Key Mediators of Erythroid Enucleation. *PLOS ONE*, *10*(11), e0142655. https://doi.org/10.1371/JOURNAL.PONE.0142655

Wölwer, C. B., Pase, L. B., Russell, S. M., & Humbert, P. O. (2016). Calcium signaling is required for erythroid enucleation. *PLOS ONE*, *11*(1), 1-12. https://doi.org/10.1371/journal.pone.0146201

Wride, M. A. (2011). Lens fibre cell differentiation and organelle loss: many paths lead to clarity. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *366*(1568), 1219. https://doi.org/10.1098/RSTB.2010.0324

Yang, J., Zhao, Y., Kalita, M., Li, X., Jamaluddin, M., Tian, B., Edeh, C. B., Wiktorowicz, J. E., Kudlicki, A., & Brasier, A. R. (2015). Systematic determination of human cyclin dependent kinase (CDK)-9 interactome identifies novel functions in RNA splicing mediated by the DEAD Box (DDX)-5/17 RNA helicases. *Molecular and Cellular Proteomics*, *14*(10), 2701-2721. https://doi.org/10.1074/mcp.M115.049221

Yoshida, H., Kawane, K., Koike, M., Mori, Y., Uchiyama, Y., & Nagata, S. (2005). Phosphatidylserine-dependent engulfment by macrophages of nuclei from erythroid precursor cells. *Nature*, *437*(7059), 754-758. https://doi.org/10.1038/nature03964

Yu, D. S., & Cortez, D. (2011). A role for cdk9-cyclin k in maintaining genome integrity. *Cell Cycle*, *10*(1), 28-32. https://doi.org/10.4161/cc.10.1.14364

Zhang, S.-M., Zhang, H., Yang, T.-Y., Ying, T.-Y., Yang, P.-X., Liu, X.-D., Tang, S.-J., & Zhou, P.-K. (2014). Interaction between HIV-1 Tat and DNA-PKcs modulates HIV transcription and class switch recombination. *International Journal of Biological Sciences*, *10*(10), 1138-1149. https://doi.org/10.7150/ijbs.10366

Zhang, S., Zhang, Y., Li, Y., Zhang, Z., Li, H., Xu, M., Lu, Z., Li, Y., & Zhao, B. (2025). Fine-tuned calcium homeostasis is crucial for murine erythropoiesis. *The FEBS Journal*, *n/a*(n/a). https://doi.org/https://doi.org/10.1111/febs.17401

Figure Legends

Figure 1. Phosphorylated CDK9 (Thr186) and Cyclin T1 are excluded from the nucleus during nuclear extrusion and localise to the cytoplasm and future reticulocyte following enucleation. (A) Diagram depicting the stages of erythroid enucleation described in this study. **(B)** Immunofluorescence confocal microscopy of mouse orthochromatic erythroblasts at the pre-polarisation and extrusion phases of erythroid enucleation stained for phospho-CDK9(Thr186), CDK9 (F-6), phalloidin for F-actin and DAPI for nuclei. Brightfield (BF) images and actin are excluded from the merge. Yellow colour on the merge indicates co-localisation of p-CDK9 and CDK9. All scale bars = 5 μm. **(C)** Immunofluorescence confocal microscopy of mouse orthochromatic erythroblasts at the pre-polarisation and extrusion phases of erythroid enucleation stained for phospho-CDK9(Thr186), Cyclin T1, phalloidin for F-actin and DAPI for nuclei. Brightfield (BF) images and actin are excluded from the merge. Yellow colour on the merge indicates co-localisation of p-CDK9 and Cyclin T1. All scale bars = 5 μm. **(D)** Immunofluorescence confocal microscopy of HUDEP-2 cells at early (day 0) and late (day 12) of differentiation stained for phospho-CDK9(Thr186), CDK9 (F-6), phalloidin for F-actin and DAPI for nuclei. Brightfield (BF) images and actin are excluded from the merge. Yellow colour on the merge indicates co-localisation of p-CDK9 and CDK9. All scale bars = 5 μm. **(E)** Immunofluorescence confocal microscopy of HUDEP-2 cells at early (day 0) and late (day 12) of differentiation stained for phospho-CDK9(Thr186), Cyclin T1, phalloidin for F-actin and DAPI for nuclei. Brightfield (BF) images and actin are excluded from the merge. Yellow colour on the merge indicates co-localisation of p-CDK9 and Cyclin T1. All scale bars = 5 μm.

Figure 2. CDK9 inhibitors cause an enucleation arrest that phenocopies cytochalasin D-mediated enucleation blockage. (A) Gating strategy used for FACS isolation of orthochromatic erythroblasts from mouse spleen following phenylhydrazine treatment alongside cytospin rapid diff staining and phenotype analysis following 12-hour incubation of isolated cells. **(B)** Quantification of enucleation of mouse orthochromatic erythroblasts following treatment with NVP-2 for 12 hours. DMSO (vehicle control) and cytochalasin D (positive control) are included, in addition to cytospin rapid diff staining and phenotype analysis. n = 4 replicates across 3 independent experiments. (**$p < 0.01$, ****$p < 0.0001$; one-way ANOVA with Dunnett's multiple comparisons test). Scale bar = 10μm. **(C)** Quantification of enucleation of mouse orthochromatic erythroblasts following treatment with AZD4573 for 12 hours. DMSO (vehicle control) and cytochalasin D (positive control) are included, in addition to cytospin rapid diff staining and phenotype analysis. n = 4 replicates across 3 independent experiments. (**$p < 0.01$, ****$p < 0.0001$; one-way ANOVA with Dunnett's multiple comparisons test). Scale bar = 10 μm. **(D)** Quantification of enucleation of mouse orthochromatic erythroblasts following treatment with cytochalasin D (CytoD) for 12 hours. DMSO (vehicle control) is included, in addition to cytospin rapid diff staining and phenotype analysis. n = 4 replicates across 2 independent experiments. (ns = not significant, ****$p < 0.0001$; one-way ANOVA with Dunnett's multiple comparisons test). Scale bar = 10 μm.

Figure 3. F-actin is required downstream of CDK9 to achieve enucleation, and blocking enucleation with cytochalasin D causes a build-up of active CDK9 that co-localises with F-actin. (A) Immunofluorescence confocal microscopy of mouse orthochromatic erythroblasts treated with 1 μM cytochalasin D (CytoD) or 100 nM NVP-2 for 12 hours and stained for phospho-CDK9(Thr186) phalloidin for F-actin, Ter119 and DAPI for nuclei. Brightfield (BF) images and Ter119 are excluded from the merge. Arrows indicate accumulation of CDK9. Yellow colour on the merge indicates co-localisation of p-CDK9 and F-actin. All scale bars = 5 μm. **(B)** Immunofluorescence confocal microscopy of mouse orthochromatic erythroblasts at the pre-polarisation and extrusion phases of erythroid enucleation stained for CDK9 (F-6), CaMKII, Ter119 and DAPI for nuclei. Brightfield (BF)

and Ter119 are excluded from the merge. Scale bars = 5 μm. **(C)** Immunofluorescence confocal microscopy of mouse orthochromatic erythroblasts following 12 hours treatment with 100 nM NVP-2 or 5 μM KN-62, stained for CDK9 (F-6), CaMKII, Ter119 and DAPI for nuclei. Brightfield (BF) and Ter119 are excluded from the merge. Arrows indicate accumulation of CDK9. All scale bars = 5 μm. **(D)** Schematic diagram describing workflow for inhibitor washout and exchange functional assays used in this study. **(E)** Enucleation rates of mouse orthochromatic erythroblasts following inhibitor washout into fresh media containing DMSO (left) or inhibitors (cytochalasin D or NVP-2; right) at 12 hours post-plating. Enucleation rates were measured at 12-, 14-, 15- and 16-hours post-plating. DMSO (black; negative control) is included. Colours change to match either DMSO (grey) or inhibitors (coloured) after washouts. Enucleation rates at hour 16 were statistically compared. n = 5 replicates across 2 independent experiments. (ns = not significant, $*p < 0.1$, $***p < 0.001$; two-way ANOVA with Tukey's multiple comparisons test). **(F)** Enucleation rates of mouse orthochromatic erythroblasts following inhibitor washout into fresh media containing DMSO (first plot) or inhibitors (cytochalasin D or KN-62) at 12 hours post-plating. Enucleation rates were measured at 12-, 14-, 15- and 16-hours post-plating. DMSO (black; negative control) is included. Colours change to match either DMSO (grey) or inhibitors (coloured) after washouts. Enucleation rates at hour 16 were statistically compared (ns = not significant, $*p < 0.1$, $***p < 0.001$; two-way ANOVA with Tukey's multiple comparisons test). **(G)** Enucleation rates of mouse orthochromatic erythroblasts following inhibitor washout into fresh media containing DMSO (first plot) or inhibitors (KN-62 or NVP-2) at 12 hours post-plating. Enucleation rates were measured at 12-, 14-, 15- and 16-hours post-plating. DMSO (black; negative control) is included. Colours change to match either DMSO (grey) or inhibitors (coloured) after washouts. Enucleation rates at hour 16 were statistically compared (ns = not significant, $*p < 0.1$, $***p < 0.001$; two-way ANOVA with Tukey's multiple comparisons test).

Figure 4. Summary of CDK9 co-immunoprecipitation mass spectrometry results in HUDEP-2 cells. (A) Venn diagram of identified interactors of CDK9 in undifferentiated (Day 0) and differentiated (Day 6) HUDEP-2 cells. Proteins were considered an interactor when present in all 3 replicates and ≥3 fold enriched compared to the IgG controls. See the supplementary file for complete list of identified proteins. **(B)** STRING analysis of the 13 common proteins identified in both undifferentiated (Day 0) and differentiated (Day 6) HUDEP-2 cells. STRING analysis was performed using medium confidence (0.4) for interaction score. Aqua lines represent known interactions from curated databases, purple lines represent experimentally determined known interactors. Green, red, and blue lines represent predicted interactors, yellow lines represent text mining identification and black lines represent known co-expression. **(C)** Venn diagram of identified interactors of CDK9 overexpression, CDK9-T186A and CDK9-D168N lines in differentiated (Day 6) HUDEP-2 cells. Proteins were considered an interactor when present in all 3 replicates and ≥3 fold enriched compared to the IgG controls. See the supplementary file for a complete list of identified proteins. **(D)** STRING analysis of the 9 common proteins identified in all 3 HUDEP-2 CDK9 overexpression cell lines. STRING analysis was performed using medium confidence (0.4) for interaction score. Aqua lines represent known interactions from curated databases, purple lines represent experimentally determined known interactors. Green, red, and blue lines represent predicted interactors, yellow lines represent text mining identification and black lines represent known co-expression. **(E)** Heatmaps of clusters 1-4 described in Supplementary Figure 7. Proteins identified in each cluster were compared to the average abundance found in CDK9, CDK9-T186A and CDK9-D168N overexpression HUDEP-2 cell lines. Top biological processes (GO) and molecular functions (GO) identified in STRING are listed below for each cluster (strength >2, false discovery <0.001).

Figure 5. Importin β activity is required for erythroid enucleation processes downstream of CDK9. (A) Quantification of enucleation of mouse orthochromatic erythroblasts following treatment

with importazole for 12 hours. DMSO (vehicle control) and cytochalasin D (positive control) are included, in addition to cytospin rapid diff staining and phenotype analysis. n = 4 replicates across 3 independent experiments. (ns = not significant, **$p < 0.01$, **$p < 0.01$, ****$p < 0.0001$; one-way ANOVA with Dunnett's multiple comparisons test). **(B)** Quantification of enucleation of day 12 differentiated HUDEP-2 cells following treatment with importazole for 12 hours. DMSO (vehicle control) and cytochalasin D (positive control) are included, in addition to cytospin rapid diff staining and phenotype analysis. n = 4 replicates across 3 independent experiments. (ns = not significant, **$p < 0.01$, **$p < 0.01$, ****$p < 0.0001$; one-way ANOVA with Dunnett's multiple comparisons test). **(C)** Immunofluorescence confocal microscopy of mouse orthochromatic erythroblasts at the pre-polarisation and extrusion phases of erythroid enucleation stained for phospho-CDK9(Thr186), importin β, phalloidin for F-actin and DAPI for nuclei. Brightfield (BF) and F-actin are excluded from the merge. Scale bars = 5 μm. **(D)** Immunofluorescence confocal microscopy of mouse orthochromatic erythroblasts following 12 hours treatment with 1 μM cytochalasin D or 10 μM importazole, stained for phospho-CDK9(Thr186), importin β, phalloidin for F-actin and DAPI for nuclei. Brightfield (BF) and F-actin are excluded from the merge. Arrows indicate accumulation of CDK9. All scale bars = 5μm. **(E)** Enucleation rates of mouse orthochromatic erythroblasts following inhibitor washout into fresh media containing DMSO (first plot) or inhibitors (cytochalasin D or importazole; second plot) at 12 hours post-plating. Enucleation rates were measured at 12-, 14-, 15- and 16-hours post-plating. DMSO (black; negative control) is included. Colours change to match either DMSO (grey) or inhibitors (coloured) after washouts. Enucleation rates at hour 16 were statistically compared. (ns = not significant, **$p < 0.01$, ****$p < 0.0001$; two-way ANOVA with Tukey's multiple comparisons test). **(F)** Enucleation rates of mouse orthochromatic erythroblasts following inhibitor washout into fresh media containing DMSO (first plot) or inhibitors (NVP-2 or importazole; second plot) at 12 hours post-plating. Enucleation rates were measured at 12-, 14-, 15- and 16-hours post-plating. DMSO (black; negative control) is included. Colours change to match either DMSO (grey) or inhibitors (coloured) after washouts. Enucleation rates at hour 16 were statistically compared (ns = not significant, **$p < 0.01$, ****$p < 0.0001$; two-way ANOVA with Tukey's multiple comparisons test. **(G)** Enucleation rates of mouse orthochromatic erythroblasts following inhibitor washout into fresh media containing DMSO (first plot) or inhibitors (KN-62 or importazole) at 12 hours post-plating. Enucleation rates were measured at 12-, 14-, 15- and 16-hours post-plating. DMSO (black; negative control) is included. Colours change to match either DMSO (grey) or inhibitors (coloured) after washouts. Enucleation rates at hour 16 were statistically compared (ns = not significant, *p < 0.1, ***p < 0.001; two-way ANOVA with Tukey's multiple comparisons test).

Figure 6. Proposed non-mutually exclusive models for the role of CDK9 and importin β in regulating erythroid enucleation prior to CaM/Ca^{2+} signalling. (A) Order of action model of CDK9 activty to achieve nuclear extrusion by downstream activation of the calmodulin (CaM) pathway through direct, or indirect activity of importin β. CaM activation results in CaMKII and MLCK activation which in turn results in F-actin polymerisation and myosin IIB contraction respectively to achieve nuclear extrusion. Inhibitors shown in colours correspond to drugs used throughout this study. Inhibitors shown in grey have been previously described in this context (Wölwer et al., 2016). **(B)** Non-mutually exclusive models depicting potential roles for CDK9 and importin β in enucleation. NPC = Nuclear pore complex. Model 1: Importin β regulates nuclear import of essential regulators of enucleation. CDK9 facilitates transcription of key enucleation genes. Model 2: Importin β acts alongside NEMP1 to facilitate essential nuclear envelope openings, allowing nucleocytoplasmic transport of CDK9 and other key enucleation regulators. Model 3: CDK9 regulates an enucleation checkpoint in the cytoplasm. Importin β, NEMP1 and Ran facilitate key signalling mechanisms between the nucleus, nuclear envelope and cytoskeleton.

Supplementary Figure Legends

Supplementary Figure 1. NVP-2 blocks enucleation of HUDEP-2 and BEL-A cells. (A) Quantification of enucleation of differentiated HUDEP-2 cells (Day 12) following 24-hour treatment with NVP-2. DMSO (vehicle control) and cytochalasin D (positive control) are included, in addition to cytospin rapid diff staining of DMSO control cells and phenotype analysis of DMSO control and NVP-2 treated cells. n = 205 cells counted (DMSO) and 186 cells counted (NVP-2). (ns = not significant, **$p < 0.01$, ****$p < 0.0001$; one-way ANOVA with Dunnett's multiple comparisons test). Scale bar = 10 μm. **(B)** Quantification of enucleation of differentiated BEL-A cells (Day 12) following 24-hour treatment with NVP-2. DMSO (vehicle control) and cytochalasin D (positive control) are included, in addition to cytospin rapid diff staining of DMSO control cells and phenotype analysis of DMSO control and NVP-2 treated cells. n = 235 cells counted (DMSO) and 168 cells counted (NVP-2). (ns = not significant, **$p < 0.01$, ****$p < 0.0001$; one-way ANOVA with Dunnett's multiple comparisons test). Scale bar = 10 μm.

Supplementary Figure 2. CDK9 inhibitors Atuveciclib and THAL-SNS-032 cause a dose dependent arrest of mouse erythroblast enucleation. (A) Quantification of enucleation of mouse orthochromatic erythroblasts following treatment with Atuveciclib (BAY1143572) for 12 hours. DMSO (vehicle control) is included. n = 4 replicates across 2 independent experiments. (****$p < 0.0001$; one-way ANOVA with Dunnett's multiple comparisons test). **(B)** Quantification of enucleation of mouse orthochromatic erythroblasts following treatment with THAL-SNS-032 for 12 hours. DMSO (vehicle control) is included. n = 4 replicates across 2 independent experiments. (**$p < 0.01$, ****$p < 0.0001$; one-way ANOVA with Dunnett's multiple comparisons test).

Supplementary Figure 3. Characterisation of HUDEP-2 CDK9 overexpression cell lines. (A) Western blot of HUDEP-2 cell lines, including the GFP vector control, CDK9, CDK9-T186A and CDK9-D168N overexpression lines. Bands can be detected for Flag-CDK9 in the three CDK9 overexpression lines. GAPDH is shown as a loading control. **(B)** Western blot of HUDEP-2 cell lines, including the GFP vector control, CDK9, CDK9-T186A and CDK9-D168N overexpression lines. Bands can be detected for Flag in the three CDK9 overexpression lines. GAPDH is shown as a loading control. **(C)** Flow cytometry histogram showing GFP expression in HUDEP-2 cell lines, including the GFP vector control, CDK9, CDK9-T186A and CDK9-D168N overexpression lines compared to negative control HUDEP-2 cells. Propidium iodide was used to gate for only viable cells. **(D)** Immunofluorescence confocal microscopy of undifferentiated HUDEP-2 cells transduced to overexpress CDK9, CDK9-T186A, CDK9-D168N or GFP only and stained for CDK9, Flag, GFP and DAPI for nuclei. Transduced cells expressing low levels of eGFP are shown in comparison to the wildtype. Brightfield (BF) and eGFP are excluded from the merge. Yellow colour depicts co-localisation of CDK9 and Flag. All scale bars = 5 μm.

Supplementary Figure 4. Interactome of identified proteins in HUDEP-2 CDK9 overexpression cell lines. STRING analysis of proteins identified across all HUDEP-2 overexpression lines. Proteins were considered an interactor of CDK9 if identified in all 3 replicates and were ≥3 fold enriched compared to the IgG control. Clusters were identified using STRING analysis of all proteins identified in overexpression lines (243 total proteins), with high confidence (0.7) for interaction score, with unconnected nodes removed. Clustering was performed using kmeans clustering into 11 groups. Aqua lines represent known interactions from curated databases, purple lines represent experimentally determined known interactors. Green, red, and blue lines represent predicted interactors, yellow lines represent text mining identification and black lines represent known co-expression. Dotted lines

represent connections between clusters. See the supplementary file for a complete list of identified proteins and corresponding abundances.

Supplementary Figure 5. Importazole alters the nucleocytoplasmic ratio of Cyclin T1 but not phosphorylated CDK9. (A) Immunofluorescence confocal microscopy of undifferentiated (Day 0) HUDEP-2 cells stained for phospho-CDK9(Thr186), cyclin T1, F-actin and DAPI. Merges showing pCDK9 and cyclin T1 alongside F-actin are shown. All scale bars = 5 μm. **(B)** Analysis of nuclear to cytoplasmic ratio of pCDK9, cyclin T1 and importin β in fixed undifferentiated (Day 0) HUDEP-2 cells following overnight treatment with 10μM NVP-2, 10μM Importazole or DMSO vehicle control. Each point represents an individual cell. pCDK9 DMSO n = 40, NVP-2 n = 38, Importazole n = 41; CycT1 DMSO n = 18, NVP-2 n = 19, Importazole n = 19; Importin β DMSO n = 22, NVP-2 n = 19, Importazole n = 23. (p values are shown for each comparison; one-way ANOVA with Dunnett's multiple comparisons test).

Figure 1
Newton et al (2025)

Figure 2
Newton et al (2025)

Figure 3
Newton et al (2025)

Figure 4
Newton et al (2025)

A CDK9 Co-IP Mass Spec

B CDK9 Co-IP Common Proteins
(Day 0 vs Day 6 HUDEP-2)

65 | 13 | 114

HUDEP-2 Day 0   HUDEP-2 Day 6

C CDK9 Overexpression Co-IP Mass Spec

CDK9(WT)

98
26 | 13
9
65 | 2 | 28

CDK9-T186A   CDK9-D168N

D CDK9 Co-IP Common Proteins
Day 6 HUDEP-2 Overexpression Lines

Column Z score: -3 -2 -1 0 1 2 3

E

Cluster 1
Ran nuclear import & export

RANBP2
RAN
TERT
NEMP1
MAP2K2
CSNK1A1
RGPD8
KPNB1
LRPPRC
DDX1
KPNA1
RANGAP1
CSE1L
YWHAE
XPO1
PABPC1

Biological Process (GO)
GO:0000056 Ribosomal small subunit export from nucleus
GO:0000055 Ribosomal large subunit export from nucleus
GO:0006607 NLS-bearing protein import into nucleus
Molecular Function (GO)
GO:0008536 Ran GTPase binding
GO:0140142 Nucleocytoplasmic carrier activity

Cluster 2
MCM complex & DNA transcription

TMPO
LMNB1
SSRP1
MCM7
EIF3A
MCM2
RUVBL2
CDK1
TRAPPC3
GOLGA2
CCNT1
RPA2
PRKDC
MCM5
AARS1
MCM3
PRKAG1
PPP2R2D
DARS1
MCM4
RACGAP1
YARS1
GARS1
PPP1CA
CALML5
NCAPD2
FARSA

Biological Process (GO)
GO:1902975 Mitotic DNA replication initiation
GO:0006267 Pre-replicative complex assembly involved in nuclear cell cycle
GO:0000727 Double-strand break repair via break induced replication
Molecular Function (GO)
GO:0097322 7SK snRNA binding
GO:0003688 DNA replication origin binding

Cluster 3
Actin filament capping & binding

VIM
MYL6B
AHSA1
UBAC1
ATP2B4
UBR4
TOLLIP
ACTR3
CAPZA1
CD2AP
TOM1
FKBP4
MDC1
DYNC1H1
CAPZB
CALM3
IGLL5
CAPZA2
FKBP5
KCNH1

Biological Process (GO)
GO:0051016 Barbed-end actin filament capping
GO:0030835 Negative regulation of actin filament depolymerization
GO0032272 Negative regulation of protein polymerisation
Molecular Function (GO)
GO:0030235 Nitric oxide synthase regulator activity
GO:0051015 Actin filament binding

Cluster 4
mRNA splicing & polyadenylation

DHX9
HNRNPL
HNRNPDL
APEX1
SFPQ
HNRNPA0
CPSF6
TARDBP
RBMX
CPSF7
U2AF1
CLK3
U2AF2
ILF2
SRSF7
HNRNPK
TOP1
SRSF1
MATR3
SNRPB
NXF1

Biological Process (GO)
GO:1990120 Messenger ribonucleoprotein complex assembly
GO:0110104 mRNA alternative polyadenylation
GO:0046833 Positive regulation of RNA export from nucleus
Molecular Function (GO)
GO:0030628 pre-mRNA 3-splice site binding
GO:0097157 pre-mRNA intronic binding

Figure 5
Newton et al (2025)

Figure 6
Newton et al (2025)

A

B

Model 1: Importin β regulates nuclear import of essential regulators of enucleation. CDK9 facilitates transcription of key enucleation genes.

Model 2: Importin β acts alongside NEMP1 to facilitate essential nuclear envelope openings, allowing nucleocytoplasmic transport of CDK9 and other key enucleation regulators.

Model 3: CDK9 regulates an enucleation checkpoint in the cytoplasm. Importin β, NEMP1 and Ran facilitate key signalling mechanisms between the nucleus, nuclear envelope and cytoskeleton.

Supplemental Figure 1
Newton et al (2025)

A
HUDEP-2 (Day 12) +DMSO

HUDEP-2 (Day 12) +NVP-2

HUDEP-2 (Day 12) +DMSO

HUDEP-2 (Day 12) +100µM NVP-2

B
BEL-A (Day 12) +DMSO

BEL-A (Day 12) +NVP-2

HUDEP-2 (Day 12) +DMSO

HUDEP-2 (Day 12) +100µM NVP-2

Supplemental Figure 2
Newton et al (2025)

A Atuveciclib
(BAY-1143572)

B THAL-SNS-032

Supplemental Figure 3
Newton et al (2025)

Supplemental Figure 4
Newton et al (2024)

Supplemental Figure 5
Newton et al (2024)