Metagenomic Profiling of Drinking Water Microbiomes: Insights into Microbial Diversity and Antimicrobial Resistance

Shradha Sharma[1,2†], Vikas Kumar[3†], Kaomud Tyagi[3], Barathi Lenin[2,4], Aarti Ravindran[1,2,4], Karthik Raman[2,4*], Inderjeet Tyagi[3*]

[1]Bhupat and Jyoti Mehta School of Biosciences, Indian Institute of Technology Madras, Chennai, 600036, Tamil Nadu, India.
[2]Centre for Integrative Biology and Systems mEdicine (IBSE), Indian Institute of Technology Madras, Chennai, 600036, Tamil Nadu, India.
[3]Centre for DNA Taxonomy, Molecular Systematics Division, Zoological Survey of India, Kolkata, 700053, West Bengal, India.
[4]Wadhwani School of Data Science and AI, Indian Institute of Technology Madras, Chennai, 600036, Tamil Nadu, India.

*Corresponding author(s). E-mail(s): kraman@dsai.iitm.ac.in;
indertyagi011@gmail.com;
[†]These authors contributed equally to this work.

Abstract

Monitoring microbial components in drinking water is as essential as tracking its chemical composition. Although traditional culture-based methods can offer valuable insight into the morphology, behaviour, and phenotype of an organism, their scope is limited to culturable species. With the advent of high-throughput sequencing, we can now detect a wider range of microbes in any ecosystem, gaining more efficient insights into their functional potential and metabolic capabilities. In this study, we performed metagenomic analyses to fully understand the microbiome of drinking water supplied through public distribution systems in an Indian city. Our findings identified bacteria from the phyla Pseudomonadota, Planctomycetota, Bacteroidota, and Actinomycetota, consistent with previous studies of drinking water microbiomes of other countries. We observed that the temperature of the water samples, even as little as a 5°C increase, influenced the composition and diversity of the microbial communities. No significant correlation was detected between the abundance of microbial species and the metal concentration in the sample. In addition, we traced the distribution of antibiotic resistance genes (ARGs), finding widespread resistance to aminoglycosides, tetracyclines, and macrolides across samples. However, the risk associated with these ARGs was beyond the scope of this study. In summary, this research provides a preliminary analysis of the drinking water microbiome, providing a foundation for future management strategies.

Keywords: Metagenomic analysis, Antimicrobial resistance genes, Clean drinking water, Diversity analysis Urban water management

1 Introduction

Providing safe drinking water to all is a fundamental objective of public health agencies of nations worldwide. The global monitoring of drinking water, sanitation, and hygiene, collectively known as WASH, is a critical focal point of the World Health Organization (WHO)[1]. Ensuring the availability of safe and clean drinking water is a crucial priority, especially for a populous country such as India.

Robust drinking water treatment processes (DWTP) are employed to effectively treat water so that it becomes fit for consumption. The design of DWTPs involves managing the physical, chemical, and biological contaminants in untreated water. Briefly, physical processes such as sedimentation and filtration facilitate the removal of particulate matter. Chemical treatments eliminate pollutants such as dissolved harmful substances like Polycyclic Aromatic Hydrocarbons (PAHs), Per- and polyfluoroalkyl substances (PFAs), heavy metals like lead and mercury, and synthetic dyes like malachite green and azo dyes. Biological contaminants comprise organic debris and a host of organisms, typically bacteria, viruses, protozoans, fungi, and algae [2?]. Methods such as adsorption, disinfection, filtration, flocculation, chemical treatment, and UV irradiation are commonly used to effectively remove pathogenic microorganisms that cause waterborne diseases [3]. However, the microbiological component of water consists of more than just the live organisms, making it essential to address and neutralize associated elements that adversely impact human health. However, the microbiological profile of water extends beyond live pathogens and also requires the neutralization of microbial byproducts that pose health risks.

Both free-living and biofilm-forming microbes can contain antibiotic-resistance genes (ARG). Moreover, horizontal gene transfer (HGT) plays a critical role in disseminating AMR in drinking water microbiome [4]. HGT can occur through mechanisms like transformation, transduction, and conjugation, leading to the spread of ARGs even in the absence of live pathogens. Additionally, the microbial load also comprises spores, endotoxins, exotoxins, mycotoxins, and phycotoxins. The toxins are metabolites produced by certain microorganisms and can contribute to long-term health risks if ingested [?].

Addressing the microbial safety of water also aligns with the Sustainable Development Goals (SDGs) adopted by the United Nations in 2015. SDG 6 was formulated to make water universally accessible and sustainable, while SDG 3, was created to promote good health and well-being, and is closely intertwined with water quality and equitable sanitation. Hence, while it is evident that water is the core connecting element across all SDGs achieving these goals depends on the practical implementation and continuous monitoring of DWTP and supply systems [5, 6]

In India, the Bureau of Indian Standards (BIS) regulates drinking water quality, with the latest amendment to the standards made in 2015 (Bureau of Indian Standards, 2012; Chidiac et al., 2023). Despite substantial improvements in water infrastructure, such as those driven by the Swachh Bharat Mission and Jal Jeevan mission (Ministry of Jal Shakti), gaps remain in especially understanding the microbial components of treated drinking water [7, 8].

Notably, metagenomic surveillance and analysis have not been carried out to monitor the quality of drinking water in India. Our study is the first to carry out metagenomic sequencing of samples collected from drinking water outlets in a metropolitan city to understand the microbial component of treated water. This approach will enhance our understanding of potential risks and inform future improvements in water safety monitoring.

2 Materials and Methods

2.1 Collection of water samples and DNA extraction

Drinking water samples (municipal tap water) of approximately 10 litres each were collected in triplicate from 21 locations (n=63) selected based on a grid map approach in a metropolitan city (exact location masked) located in India during March 2022. The samples were collected in High-Density Polyethylene (HDPE) autoclaved sterile bottles marked W1-W21 and kept at 4^{o} C until transportation (3 h) to the laboratory. The samples were then filtered immediately using 0.22 µm MCE membrane (Merck Life Science Pvt. Ltd., Bangalore) with the help of a vacuum pump, and the biomass-containing membranes were placed in DNA extraction tubes and stored at -20 $^{\text{o}}$C until DNA extraction. Extraction of DNA from drinking water samples was performed using a DNeasy PowerWater DNA isolation kit (Qiagen Cat No.: 14900- 100-NF) following the standard manufacturer protocol and pooled into 21 drinking water samples for sequencing. The pooling was carried out due to the limited resources available for this study. The extracted DNA was qualitatively and quantitatively checked using agarose gel electrophoresis (Cell BioScience Alphalmager MINI) and a Qubit® Fluorometer (Invitrogen, USA), respectively, as per our previous studies [9]. In addition to the drinking water samples, a negative control was filtered, as mentioned above; however, we did not obtain amplifiable DNA for sequencing from this sample.

2.2 Library preparation, Whole genome sequencing (WGS), and preprocessing of sequencing reads.

Using a Covaris ultrasonic disruptor, a 1 µg genomic DNA sample was randomly fragmented into segments of approximately 350 bp to build the library. End repair, A-tail addition, sequencing adapter ligation, purification, and PCR amplification were the final steps in library construction. Following library construction, AATI analysis was used to evaluate the size and integrity of the inserted pieces and the library as a whole. The quality of the library was confirmed by accurately quantifying its effective library concentration using qPCR, which was greater than 3 nM. Once the library passed the quality check, it was combined with other libraries on the basis of their effective concentrations and data output targets. Then, PE150 sequencing was performed. Further, the Illumina sequencing platform's (Novaseq 6000 with 150*2 chemistry) raw data was preprocessed using fastp [10] to produce clean data for further analysis. Adapters were trimmed, and the paired end reads were discarded in the following situations: 1) when there were more than 10% unknown nucleotides in each read; 2) when there were more than 50% low-quality nucleotides (base quality less than 5). The generated sequences were submitted to NCBI under the BioProject ID: PRJNA1008921 with accession number SRR25754722 to SRR25754742.

2.3 Metagenomic assembly, binning, and quality control

To remove human contamination from adapter-trimmed reads, BMTagger v3.101 aligned the reads with the hg38 reference genome. Read pairs in which only one read aligned with the human genome are also excluded. Quality checks were performed before and after filtering using FASTQC v0.12.1 [11]. The filtered reads were assembled using the MEGAHIT v1.1.1 [12] assembler within the MetaWRAP pipeline[13]. MEGAHIT employs a multiple k-mer-based strategy, where smaller k-mers fill gaps in low-coverage regions while larger k-mers resolve repeat sequences. This approach leads to a highly accurate assembly. We used the default k-mers of lengths 21, 29, 39, 59, 79, 99, 119, and 141.

Assembly was performed per sample, and contigs shorter than 1,000 bp were removed from the analysis. The assembled contigs were then binned using three algorithms: CONCOCT v0.4.0 [14], MaxBin2 v2.2.4 [15], and MetaBAT2 v2.9.1 [16], each run independently with default parameters. These algorithms leverage tetranucleotide frequencies, coverage information, and sequence composition to achieve precise binning. The consensus approach integrates the strengths of each algorithm, maximising the capture of genomic diversity while minimising potential errors that could arise from using a single method. Each of these algorithms produced separate bins that were then consolidated using the MetaWRAP bin refinement module. Subsequently, the refined bins were reassembled using SPades v3.10.1 [17]. CheckM v1.0.7 [18] was run on the final reassembled bins to prepare a report of completeness and contamination. Metagenome-assembled genomes (MAGs) with more than 90% completion and less than 5% contamination were selected for further analysis. For the estimation of the abundance of each bin, we used the Quant_bins module from the MetaWRAP pipeline with default parameters.

2.4 Diversity and differential abundance analysis

After removing reads that aligned to the human chromosomes, the filtered reads were assigned taxonomy using KRAKEN2 v2.1.3 [19] using the Kraken nt database (downloaded in May 2023). Species-level taxonomy was used for subsequent analysis. The abundance estimation used BRACKEN v2.9 [20]. The abundance table, taxa table, and metadata were integrated into a single object using the phyloseq package in R v4.3.3 for further diversity analysis [21].

CCA and CAP ordinations were used to understand the effect of heavy metals, temperature, pH, and the zone from which the sample was isolated on the microbial composition of the water samples. Beta diversity was evaluated using the Principal Coordinate Analysis (PCoA) technique. The analysis was performed using the ordinate function from the phyloseq package. Temperatures were categorized into Low (25.61 0.11) and High (31.55 0.24). Beta dispersion analysis using the betadisper function, followed by ANOVA, was used to infer differences in the variability of the composition of the microbial community between the temperature groups.

To further investigate the influence of temperature on microbial composition, ANOSIM was applied to assess within- and between-group similarities. The percentage of microbiome variation explained by temperature was quantified using Adonis2. Feature selection using Boruta identified taxa at the species level that significantly differed between high and low temperatures [22]. Additionally, ANCOM-BC was employed to detect taxa with differential abundance across the two temperature conditions [23].

Shannon and observed alpha diversity indices were measured using the estimate_richness function of phyloseq. The Wilcoxon test was used to test the significance of differences between temperature groups and the Kruskal–Wallis test for the significance of differences between zones.

2.5 Taxonomy and functional classification

The MAGs that were generated were classified using the classic workflow (classify_wf) from GTDB-Tk v2.3.2 [24]. The reference database used for the same was R214. To generate a phylogenetic tree, we used GToTree v1.8.4, a command line tool based on the Hidden Markov Model (HMM) that builds a tree using a set of single-copy marker genes [25]. The default cutoff is the presence of at least 50% of the marker genes. However, since some MAGs did not have 50% of the marker genes, we lowered this cutoff to 20%. The output tree from GToTree was passed to IQ-TREE for bootstrapping for 1000 iterations, and the output from IQ-TREE was visualized using iTOL v7 [26, 27]. We used PROKKA v1.14.6, which uses Prodigal for gene prediction [28]. The PROKKA output was subsequently used to identify Clusters of Orthologous Genes (COGs) using cogclassifier v1.0.5 [29].

2.6 Taxonomy and functional classification

The identification of antimicrobial resistance genes was performed using two approaches. First, using the filtered reads in each sample, the RGI (Resistance Gene Identifier) tool's BWT (Burrows-Wheeler Transform) feature in RGI v6.0.3 with the default kma aligner was used [30, 31]. The reads were mapped to the CARD (Comprehensive Antibiotic Resistance Database) v3.2.7 [32]. To ensure robust gene selection, the results were filtered to include only genes with at least 100 mapped reads and an average percentage coverage of 80% or greater. When a particular read was mapped to multiple reference sequences, the median reference sequence was used to ensure a more accurate

representation. RPKM was calculated for genes that meet these criteria for normalisation across samples.

$$\text{Reads per million kilobase (RPKM)} = \frac{\text{Reads mapped to gene}}{\text{Gene length (Kbp)} \times \text{Total mapped reads (millions)}}$$

The prediction of ARGs was also implemented by screening the MAGs obtained against the CARD database using the RGI main feature [31]. ORFs with 60% identity with CARD references were further selected for analysis. Furthermore, potential virulence factor genes (VFGs) in MAGs were identified using BLASTp v2.12.0 against the VFDB database (core protein dataset), with a query coverage threshold of 80%, an identity cutoff of 50%, and an e-value cutoff of 1e-5 [33].

2.7 Taxonomy and functional classification

To investigate water quality, analysis of heavy metals such as magnesium (Mg), manganese (Mn), copper (Cu), arsenic (As), cadmium (Cd), mercury (Hg) and lead (Pb) along with pH and temperature was carried out using AAS ElementAS Model AAS4129 and Hanna 9829 multiparameter respectively. The findings were compared with the guidelines prescribed by the WHO, 2022, and IS 10500:2012 in guidelines for guidelines for drinking water quality (**Table 1**).

3 Results

3.1 Water quality (heavy metals, pH, and temperature)

The pH and temperature of the drinking water samples (W1–W21) ranged from 6.8 to 7.5 and 25.4°C to 32°C, respectively. The pH levels were within the acceptable limits of IS:10500:2012 for potable water. Furthermore, toxic heavy metals with known carcinogenic potential: arsenic (As), cadmium (Cd), lead (Pb), copper (Cu), and mercury (Hg) were detected at concentrations of 0.56–3.05, 0–0.11, 0–1.72, 0.25–11.6, and 0–0.11 µg/ml (ppb), respectively. These values comply with the drinking water quality standards set by WHO and IS:10500:2012 (**Table 1**). I, essential minerals such as magnesium (Mg), manganese (Mn), iron (Fe), and zinc (Zn) were found to be within the permissible limits outlined by these standards. Thus, it can be inferred that in terms of physicochemical contaminants, the drinking water obtained from the metropolitan city falls under the criteria "Safe for drinking purposes".

Table 1: Heavy metal analysis and their comparison with permissible limits prescribed by the WHO, 2022 and IS:10500:2012.

S.no.	Samples	Metals/Parameters	Observed value(ppb)	WHO Limit (2022) (ppb)	IS:10500:2012 Limit (ppb)
1.	W1-W21	Arsenic (As)	0.56-3.05	10	NP[1]
2.	W1-W21	Cadmium (Cd)	0-0.11	3	3
3.	W1-W21	Copper (Cu)	0.25-11.6	2000	50
4.	W1-W21	Lead (Pb)	0-1.72	10	10
5.	W1-W21	Manganese (Mn)	0.56-105.19	80	100
6.	W1-W21	Mercury (Hg)	0-0.11	6	1
7.	W1-W21	Magnesium (Mg)	4391.08-28792.6	NP[1]	30000
8.	W1-W21	Zinc (Zn)	0-306.99	NP[1]	5000
9.	W1-W21	Iron (Fe)	0-277.42	NP[1]	300
10.	W1-W21	pH	6.8-7.5	NP[1]	6.5-8.5
11.	W1-W21	Temperature °C	25.4-32	NP[1]	NP[1]

[1]NP: Not provided

3.2 Diversity of drinking water metagenome

The taxonomy assignment of the filtered reads showed that, on average, $59.19 \pm 6.34\%$ of the reads in a sample were successfully assigned a taxonomy, with $40.85 \pm 6.35\%$ classified at the species level (Supplementary_Table_1). Only reads assigned to Bacteria, Archaea, Fungi, and Viruses were included in the diversity analyses. Bacteria dominated the microbial composition, comprising 98.41% of the identified taxa, while the other domains accounted for 0.38%, 0.99%, and 0.21%, respectively. We tried to identify the core taxa present in all samples. By calculating the prevalence of the species at a read cutoff of more than 10,000, five species were found to have 100% prevalence: *Afipia carboxidovorans, Klebsiella pneumoniae, Pseudomonas aeruginosa, Sphingopyxis macrogoltabida,* and *Variovorax paradoxus.* When the read cutoff was reduced to 1,000, the number of species with a prevalence of 100% increased to more than 500. The composition of Bacteria and Archaea at the Class level followed by a plot of the abundances of the 15 most abundant bacterial/ archaeal species is shown in (**Supplementary Fig. 1**). The composition of Fungi at the order level followed by a plot of the abundances of the 15 most abundant fungal species is shown in (**Supplementary Fig. 2**).

3.3 Drivers of microbial diversity

The effects of covariates on the composition are shown in figure (**Supplementary Fig. 3**). We observe that the temperature and the mercury content are correlated. Since mercury levels have a small range and are well within permissible limits, we infer that temperature appears to drive the microbial composition. Although pH also appears to influence the microbial composition, its range is limited and the results are inconsistent, with PERMANOVA findings contradicting those of CCA. Similar inconsistencies are observed for heavy metals. Across all analyses, changes in temperature consistently correlate with shifts in the microbial composition. Therefore, we focus on analysing the effects of temperature on microbial diversity in more detail.

Principal Coordinates Analysis (PCoA) shows the clustering of samples from the low and high temperature groups (**Fig.1(B)**). The sample sizes were balanced, with 10 samples from the lower temperature group (25°C) and 11 from the higher temperature group (31°C). The homogeneity of the variances was assessed using Beta Dispersion with the betadisper function. ANOVA revealed significant differences in beta dispersion between groups (p = 0.002413), indicating variability in the composition of the microbial community under different temperature conditions.

PERMANOVA, performed using the Adonis2 package, also identified temperature as a significant contributor to microbial diversity, with an R^2 value of 0.4265 and p = 0.001 (**Fig.1(B)**). However, since PERMANOVA results can be influenced by differences in beta dispersion, we performed an ANOSIM analysis. ANOSIM confirmed greater similarity within the temperature groups than between them (p = 0.001, R = 0.4513) (**Supplementary Fig. 4**), further supporting the role of temperature in shaping the composition of the microbial community.

Alpha diversity was measured using both the Shannon index (which measures richness and evenness) and the observed species index (which estimates richness) to estimate the microbial richness and evenness across zones **Fig. 1(A)**. No significant differences were observed, which aligns with expectations, as the water supplied to the zones was subjected to uniform treatment. Although the PCoA plot shows a separation between the North and East zones, their species richness and evenness were not significantly different. In contrast, the north and west zones were clearly separated in the PCoA plot, with a significant difference in alpha diversity (p = 0.024) (**Supplementary Fig. 5**).

To identify differentially abundant taxa under different temperature conditions, we used a random forest-based wrapper, Boruta, to the filtered phyloseq object (prevalence = 0.1, detection = 50). We also used ANCOM-BC on the filtered data, and the results were compiled with a significance cutoff of 0.05 with the p adjustment method - "holm." The taxa picked up by ANCOM-BC are presented in **Fig.1(C)**. In particular, both ANCOM-BC and Boruta identified common taxa that were more abundant at high temperatures, including *Acinetobacter indicus, Acinetobacter junii, Acinetobacter sp.* WCHA45, *uncultured bacterium* contig00142, *Hyphomicrobium zavarzinii*, and *Acinetobacter sp.* SCLZS86. A detailed list of taxa identified by BORUTA and ANCOM-BC is presented in **Supplementary Table 2**

Fig. 1: Effect of temperature on the diversity of the microbial community in the samples **A.** Alpha diversity was estimated between high and low-temperature groups using the Shannon and Observed diversity indices. The p-values for Shannon (0.085) and Observed (0.17) suggest no significant difference in species richness or evenness between the two groups. **B.** Beta diversity comparisons were visualized using Principal Coordinates Analysis (PCoA). These plots reveal that, while species richness and evenness do not differ significantly between temperature groups, temperature plays a crucial role in determining the species composition of the samples. **C.** Differential abundance analysis of taxa at high and low temperatures. Red represents taxa more abundant in low temperature

3.4 Community structure of drinking water

After applying a stringent filter (over 90% completion and less than 5% contamination), 358 bacterial MAGs were obtained for further analysis, with 233 bins from high temperature and 125 from low temperature. We chose not to run dREP to retain information about the samples and the genomic content of the individual strains for COG analysis. Classification of the MAGs revealed that Pseudomonadota was the most dominant phylum in 68.44 ± 14.35% of the samples in various zones. This was followed by Planctomycetota at 10.74 ± 8.68%, Bacteroidota at 6.35 ± 6.28%, and Actinomycetota at 6.92 ± 6.80%. Pseudomonadota was observed to be the dominant phylum in all zones, and Actinomycetota was not observed in the north zone. For a deeper understanding of the microbial composition, the MAGs were quantified to find the most abundant groups. Sphingomonadaceae was the most abundant family in the community. The abundance of different families is shown in **Supplementary Fig. 6(A)**.

We used GRiD to identify genomes with higher replication rates in the samples, as a substitute for viability analysis. GRiD scores identify actively dividing bacterial cells by computing the ratio of the origin of replication to termination sites. Among the families analysed, Burkholderiaceae_B exhibited the highest growth rate. Interestingly, species belonging to the Sphingomonadaceae and the Burkholderiaceae_B families demonstrated high growth rates and substantial abundance within the community (**Fig. 3 , Supplementary Fig. 7(A)**).

Fig. 2: Phylogenetic tree of the MAGs. Phylum-level classification of the MAGs obtained from the samples, showing their completion levels and the respective zones from which the samples were collected.

Our primary objective was to gain a broader understanding of the microbes present in drinking water. We annotated Open Reading Frames (ORFs) to identify coding sequences (CDS) to achieve this. A total of 969,369 distinct Clusters of Orthologous Genes (COGs) were identified, belonging to 25 functional categories. To ensure specificity and sensitivity, we applied a 30% identity cutoff, which refined the number of COGs to 661,322. A comparative analysis of these COGs in all zones revealed

that the number of genes associated with different functional categories did not differ significantly. The abundance of COG categories is represented in **Supplementary Fig. 8**.

Fig. 3: Replication rate of families identified in the samples. Scatter plot depicting bacterial families with a GRiD score greater than 1.3 (indicating actively dividing bacteria) and a log-transformed abundance exceeding 0.75. Only families meeting these thresholds are labeled.

Although the concentrations of all metals in the samples were within the permissible limits, the concentration of iron in one of the samples was quite different from the mean values. Our analysis identified three COGs that are associated with iron metabolism: COG0735 (Fe^{2+} or Zn^{2+} uptake regulation protein Fur/Zur), COG1914 (Mn^{2+} or Fe^{2+} transporter, NRAMP family), and COG0672 (High-affinity Fe^{2+}/Pb^{2+} permease). Interestingly, the presence of COG0735 was also high in the Sphingomonadaceae family and samples at high temperatures (**Fig. 4, Supplementary Fig. 8**). A complete presence/absence list of all COGs identified in the MAGs has been presented in **Supplementary Table 2**.

Fig. 4: Distribution of COGs involved in iron metabolism among the bacterial families. COG0735 (Fe^{2+} or Zn^{2+} uptake regulation protein Fur/Zur), COG1914 (Mn^{2+} or Fe^{2+} transporter, NRAMP family), and COG0672 (High-affinity Fe^{2+}/Pb^{2+} permease). Only families with more than 10% relative abundance are shown in the figure.

3.5 Distribution of ARGs in the drinking water

Antimicrobial Resistance Genes (ARGs) were characterised from metagenomic reads using RGI-bwt, following the filtration criteria described in the Methods section. We identified 82 unique ARGs that span 43 different families of AMR genes after applying these filters. Normalisation was carried out by calculating the RPKM (reads per million kilobases) for each gene identified to quantify the abundance of antibiotic resistance genes and comparing them between different samples. In particular, genes associated with resistance to aminoglycoside, elfamycin, fluoroquinolone, macrolide, and tetracycline drug classes were consistently detected in all samples and zones (**Fig. 5**). Among all drug classes, aminoglycoside and macrolide antibiotics had the highest diversity of ARGs, with 22 and 25 unique genes, respectively.

Multidrug-resistant genes comprised 14% of all the genes identified. The *Pseudomonas aeruginosa* CpxR gene, detected exclusively in the W18 sample, showed resistance to 15 drug classes, the highest among all genes identified. In general, the W18 sample contained resistance genes for 28 of the 29 drug classes identified. Resistance to the drug classes aminocoumarin, monobactam, and sulfonamide was also detected exclusively in this sample. **Supplementary Table 2** provides a detailed list of drug classes, their associated resistance genes, and their corresponding RPKM values.

Fig. 5: Stacked bar plot showing the abundance of antimicrobial resistance genes (ARGs) conferring resistance to different drug classes across samples. The x-axis represents individual samples, while the y-axis indicates the normalized abundance of ARGs expressed in RPKM (Reads Per Kilobase of transcript per Million mapped reads). Each stack corresponds to a specific drug class, with the height proportional to the total abundance of ARGs identified for that class. Only the top five drug classes for which ARGs are identified in all samples are shown here.

The mechanisms by which genes in this study confer resistance include antibiotic efflux, antibiotic inactivation, target replacement, target alteration, and resistance by absence. Although most genes conferred resistance through target alteration, most multidrug resistance genes conferred resistance primarily through the efflux mechanism.

To better understand antibiotic resistance genes (ARGs) in the metagenome-assembled genomes (MAGs), we used the rgi-main tool with the CARD database. The adeF gene, which confers resistance to fluoroquinolone and tetracycline, was the most prevalent ARG in the samples, detected in 15 bacterial families, with the highest representation in the Pseudomonadaceae family. Focussing on multidrug resistant genes, YajC was exclusively identified in species of Pseudomonadaceae. It exhibited resistance to ten drug classes, including fluoroquinolones, cephalosporins, glycylcyclines, penams, tetracyclines, oxazolidinones, glycopeptides, rifamycins, phenols, disinfectant agents, and antiseptics. Furthermore, the identified Pseudomonas aeruginosa soxR gene was resistant to 8 drug classes and was found exclusively in Xanthobacteraceae (**Fig. 6, 7**). Among drug classes, fluoroquinolones had the highest prevalence of resistance-associated genes in the MAGs. A detailed list of genes identified in MAGs, along with their mechanism of action and drug classes to which they confer resistance, is presented in **Supplementary Table 2**. In general, most ARGs were found in the Pseudomonadaceae and Mycobacteriaceae families. The distribution of ARGs across bacterial families is shown in **Fig. 7**.

Fig. 6: Chord diagram representing the distribution of genes identified in the MAGs across different Families. In the chord diagram, the gene sectors are displayed from left to right in the same order as they appear in the legend. Similarly, the Family sectors are displayed from right to left, maintaining the order specified in the legend.

Gene
- *Acinetobacter baumannii* AbaQ
- MtB intrinsic murA conferring resistance to fosfomycin
- MtB rpsL mutations conferring resistance to streptomycin
- BJP-1
- ErmR
- FosI
- MAB
- RbpA
- SGM-6
- YajC
- aadA10
- adeF
- dfrA5
- rsmA
- sul2
- *Pseudomonas aeruginosa* soxR
- *Streptomyces venezuelae* rox

Family
- *Azospirillaceae*
- *Beijerinckiaceae*
- *Burkholderiaceae*
- *Burkholderiaceae_B*
- *Hyphomicrobiaceae*
- *Ilumatobacteraceae*
- *Moraxellaceae*
- *Mycobacteriaceae*
- *Nevskiaceae*
- *Nocardioidaceae*
- *Pseudomonadaceae*
- *Rhodocyclaceae*
- *SXQI01*
- *SXTU01*
- *Sphingomonadaceae*
- *TH1-2*
- *UBA2020*
- *UBA4765*
- *UBA9464*
- *Verrucomicrobiaceae*
- *Xanthobacteraceae*

Fig. 7: Genes detected in the MAGs, their mechanism of action, and the drug classes to which they are resistant. Amongst all ARGs represented here, Pa soxR, MAB, YajC, rsmA, adeF, and ErmR are multidrug resistant. The abbreviations used in the figure are as follows: Pa soxR: Pseudomonas aeruginosa soxR, Ab AbaQ: Acinetobacter baumannii AbaQ, murA: Mycobacterium tuberculosis intrinsic murA conferring resistance to fosfomycin, rpsL: Mycobacterium tuberculosis rpsL mutations conferring resistance to streptomycin, D.A.: Disinfecting agents, Sv: Streptomyces venezuela rox.

3.6 Virulence Factors identified in Drinking water

The virulence factors obtained after filtration belonged to 14 categories whose distribution is shown in **Supplementary Fig. 9**. Our analysis indicates that nutritional / metabolic factors are prevalent in all families, closely followed by immune modulation and motility genes. Given that the Sphingomonadaceae family was the most abundant, as determined from our MAG classification, it also exhibited the highest number of virulence factors. Consequently, no definitive conclusions can be drawn about the enrichment of specific genes in one family over another.

4 Discussion

Periodic monitoring of drinking water supplied through the public water distribution system post-treatment is essential. This ensures that quality is maintained and that public health is not adversely impacted by persistent abiotic or biotic contaminants. The emergence of metagenomics offers a comprehensive and holistic understanding of the microbial community that will help develop effective management and mitigation strategies to ensure safe and high-quality drinking water [34].

In the present study, we sampled water from 21 locations from four zones of an Indian city. To the best of our knowledge this is the first metagenomic study of drinking water from India. The most abundant phyla in our analysis were Pseudomonadota (Proteobacteria), Planctomycetota, Bacteroidota, and Actinomycetota. This finding is consistent with previous reports [35].

Diversity analyses showed that temperature was the main factor influencing microbial composition, followed by the zone, to a lesser extent. The impact of metal concentration on microbial diversity

could not be assessed due to the narrow range of values for some metals and the limited sample size, which makes it difficult to draw any conclusions. Additionally, there was at least one outlier for each metal that would have affected the analysis, so we focused our study on temperature and zone. The levels were also within acceptable limits, making further analysis unnecessary.

Among the microbial families identified in our samples, Sphingomonadaceae was the most dominant, with 45 MAGs assigned to this family. Members of this family have been identified in various ecosystems, including soil, corals, clinical samples, and drinking water[36]. They are also known to be strictly aerobic chemoautotrophs and play a crucial role in initiating biofilm formation and contribute significantly to biofilm maturation [37, 38]. Furthermore, the higher prevalence of the Fur gene within this family highlights intriguing avenues for future research. The Fur gene regulates cellular iron homeostasis, with the iron-Fur complex activating in the presence of elevated ferrous iron levels, thus repressing iron metabolism [39]. However, no correlation was found between the abundance of the Sphingomonadaceae family and the concentration of iron in our study. Although there was a trend suggesting higher levels of COG0735 in the high temperature group, the relationship remains inconclusive due to the limited sample size.

In addition to members of the Sphingomonadaceae family, members of the Rhodobacteraceae and Beijerinckiaceae families, although they had half the abundance of Sphingomonadaceae, showed a significant presence of COG0735 (**Fig. 4**). A higher abundance of nutritional/metabolic virulence factors in Rhodobacteraceae and Beijerinckiaceae highlights their potential roles in competitive nutrient acquisition and survival under trace metal-limited conditions (**Supplementary Fig. 9**). Rhodobacteraceae, in particular, are known for their dependence on trace metals, which supports their diverse metabolic pathways and ecological strategies [40]. The profiles of virulence factors and COGs in these two families raise important questions about the role of iron in their potential synergistic interactions or competition, which could be explored in future studies.

The analysis of antibiotic resistance genes revealed the presence of the ermR gene in both metagenomic reads and MAGs. The sole carrier of this gene is Aeromicrobium erythreum, which was also binned from our samples. The gene was found in samples W11, W18, and W19 (**Supplementary Table 2**). This gene, previously known as ermA, has been reported to confer resistance to macrolide-lincosamide-streptogramin (MLS) antibiotics, consistent with our findings. MAGs for Aeromicrobium erythreum were also binned from these three samples: W18_bin.26.orig, W19_bin.11.orig, and W11_bin.8.orig (**Supplementary Table 1**). This bacterium has not been previously reported in drinking water [41, 42]). It should be noted that the presence of a gene is not an indication of its expression. The expression of a gene depends on environmental conditions and regulatory mechanisms.

5 Conclusion

A comprehensive understanding of the microbial community of drinking water is crucial to ensure its quality and safety. In this study, we used metagenomics to analyse the microbial diversity of drinking water supplied through the public distribution systems of an Indian city. Our analysis revealed that among all the measured physicochemical parameters, temperature was the most significant factor influencing the composition of the microbial community. The influence of other factors could not be assessed due to the limited sample size, which was insufficient to draw any statistically significant patterns. Furthermore, since propidium monoazide (PMA) treatment was not performed on the samples, we could not determine whether the microbes identified were live or dead. Our analysis also identified antimicrobial resistance genes (ARGs). Although the presence of ARGs does not directly indicate a risk, it underscores the importance of future research focused on horizontal gene transfer and risk assessment of these ARGs. This study reinforces previous observations of drinking water microbiomes while also providing new insights into its composition. It also improves our understanding of the drinking water ecosystem and reiterates the importance of monitoring to protect public health.

Supplementary information. The following are the Supplementary data for this article.
Supplementary Figures
Supplementary Table 1
Supplementary Table 2

Acknowledgements. V.K., K.T., and I.T. are thankful to the Director of Zoological Survey of India and the RAMC committee for funding the project titled "Drinking Water Microbiome: Monitoring of the Water Quality and Associated Health Implications." S.S. acknowledges the half-time research assistantship from the Ministry of Education, Government of India. B.L. acknowledges support from the Centre for Integrative Biology and Systems mEdicine (IBSE). A.R. acknowledges support from the project RB22231279BTHINT008481.

Data availability statement

The generated sequences were submitted to NCBI under the bio project ID: PRJNA1008921 with accession number SRR25754722 to SRR25754742.

References

[1] Organization, W.H.: WHO Global Water, Sanitation and Hygiene: Annual Report 2022. World Health Organization, Geneva, Switzerland (2023)

[2] Tyagi, I., Kumar, V., Karri, R.R.: Environmental Metagenomics, Water Quality and Suggested Remediation Measures of Polluted Waters: A Combined Approach. Elsevier, ??? (2024)

[3] Obayomi, O.V., Olawoyin, D.C., Oguntimehin, O., Mustapha, L.S., Kolade, S.O., Oladoye, P.O., Oh, S., Obayomi, K.S.: Exploring emerging water treatment technologies for the removal of microbial pathogens. Current Research in Biotechnology, 100252 (2024)

[4] Dias, M.F., Rocha Fernandes, G., Paiva, M.C., Matos Salim, A.C., Santos, A.B., Nascimento, A.M.A.: Exploring the resistome, virulome and microbiome of drinking water in environmental and clinical settings. Water Research **174**, 115630 (2020)

[5] Alcamo, J.: Water quality and its interlinkages with the sustainable development goals. Current opinion in environmental sustainability **36**, 126–140 (2019)

[6] Bhaduri, A., Bogardi, J., Siddiqi, A., Voigt, H., Vörösmarty, C., Pahl-Wostl, C., Bunn, S.E., Shrivastava, P., Lawford, R., Foster, S., *et al.*: Achieving sustainable development goals from a water perspective. Frontiers in Environmental Science **4**, 64 (2016)

[7] Chidiac, S., El Najjar, P., Ouaini, N., El Rayess, Y., El Azzi, D.: A comprehensive review of water quality indices (wqis): history, models, attempts and perspectives. Reviews in Environmental Science and Bio/Technology **22**(2), 349–395 (2023)

[8] Water–Specification, I.S.D.: Bureau of indian standards. New Delhi, India, 1–12 (2012)

[9] Tyagi, I., Tyagi, K., Gupta, V., Dutta, R., Singhvi, N., Kumar, V., Bhutiani, R., Prakash, O.: Microbial diversity characterizations, associated pathogenesis and antimicrobial resistance profiling of najafgarh drain. Environmental Research **238**, 117140 (2023)

[10] Chen, S.: Ultrafast one-pass fastq data preprocessing, quality control, and deduplication using fastp. Imeta **2**(2), 107 (2023)

[11] Brown, J., Pirrung, M., McCue, L.A.: Fqc dashboard: integrates fastqc results into a web-based, interactive, and extensible fastq quality control tool. Bioinformatics **33**(19), 3137–3139 (2017)

[12] Li, D., Luo, R., Liu, C.-M., Leung, C.-M., Ting, H.-F., Sadakane, K., Yamashita, H., Lam, T.-W.: Megahit v1. 0: a fast and scalable metagenome assembler driven by advanced methodologies and community practices. Methods **102**, 3–11 (2016)

[13] Uritskiy, G.V., DiRuggiero, J., Taylor, J.: Metawrap—a flexible pipeline for genome-resolved metagenomic data analysis. Microbiome **6**, 1–13 (2018)

[14] Alneberg, J., Bjarnason, B.S., De Bruijn, I., Schirmer, M., Quick, J., Ijaz, U.Z., Lahti, L., Loman, N.J., Andersson, A.F., Quince, C.: Binning metagenomic contigs by coverage and composition.

Nature methods **11**(11), 1144–1146 (2014)

[15] Wu, Y.-W., Simmons, B.A., Singer, S.W.: Maxbin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets. Bioinformatics **32**(4), 605–607 (2016)

[16] Kang, D.D., Froula, J., Egan, R., Wang, Z.: Metabat, an efficient tool for accurately reconstructing single genomes from complex microbial communities. PeerJ **3**, 1165 (2015)

[17] Bankevich, A., Nurk, S., Antipov, D., Gurevich, A.A., Dvorkin, M., Kulikov, A.S., Lesin, V.M., Nikolenko, S.I., Pham, S., Prjibelski, A.D., *et al.*: Spades: a new genome assembly algorithm and its applications to single-cell sequencing. Journal of computational biology **19**(5), 455–477 (2012)

[18] Parks, D.H., Imelfort, M., Skennerton, C.T., Hugenholtz, P., Tyson, G.W.: Checkm: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. Genome research **25**(7), 1043–1055 (2015)

[19] Wood, D.E., Lu, J., Langmead, B.: Improved metagenomic analysis with kraken 2. Genome biology **20**, 1–13 (2019)

[20] Lu, J., Breitwieser, F.P., Thielen, P., Salzberg, S.L.: Bracken: estimating species abundance in metagenomics data. PeerJ Computer Science **3**, 104 (2017)

[21] McMurdie, P.J., Holmes, S.: phyloseq: an r package for reproducible interactive analysis and graphics of microbiome census data. PloS one **8**(4), 61217 (2013)

[22] Kursa, M.B., Rudnicki, W.R.: Feature selection with the boruta package. Journal of statistical software **36**, 1–13 (2010)

[23] Lin, H., Peddada, S.D.: Analysis of compositions of microbiomes with bias correction. Nature communications **11**(1), 3514 (2020)

[24] Chaumeil, P.-A., Mussig, A.J., Hugenholtz, P., Parks, D.H.: GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. Oxford University Press (2020)

[25] Lee, M.D.: Gtotree: a user-friendly workflow for phylogenomics. Bioinformatics **35**(20), 4162–4164 (2019)

[26] Letunic, I., Bork, P.: Interactive tree of life (itol) v5: an online tool for phylogenetic tree display and annotation. Nucleic acids research **49**(W1), 293–296 (2021)

[27] Nguyen, L.-T., Schmidt, H.A., Von Haeseler, A., Minh, B.Q.: Iq-tree: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. Molecular biology and evolution **32**(1), 268–274 (2015)

[28] Seemann, T.: Prokka: rapid prokaryotic genome annotation. Bioinformatics **30**(14), 2068–2069 (2014)

[29] Shimoyama, Y.: COGclassifier: A Tool for Classifying Prokaryote Protein Sequences Into COG Functional Category. https://github.com/moshi4/COGclassifier

[30] Clausen, P.T., Aarestrup, F.M., Lund, O.: Rapid and precise alignment of raw reads against redundant databases with kma. BMC bioinformatics **19**, 1–8 (2018)

[31] Alcock, B., Huynh, W., Chalil, R., Smith, K., Raphenya, A., Wlodarski, M., Edalatmand, A., Petkau, A., Syed, S., Tsang, K., Baker, S., Dave, M., McCarthy, M., Mukiri, K., Nasir, J., Golbon, B., Imtiaz, H., Jiang, X., Kaur, K., Kwong, M., Liang, Z., Niu, K., Shan, P., Yang, J., Gray, K., Hoad, G., Jia, B., Bhando, T., Carfrae, L., Farha, M., French, S., Gordzevich, R., Rachwalski, K., Tu, M., Bordeleau, E., Dooley, D., Griffiths, E., Zubyk, H., Brown, E., Maguire, F., Beiko, R., Hsiao, W., Brinkman, F., Van, G., McArthur, A.: CARD 2023: Expanded Curation, Support for Machine Learning, and Resistome Prediction at the Comprehensive Antibiotic Resistance

Database. https://doi.org/10.1093/nar/gkac920 . https://github.com/arpcard/rgi

[32] Alcock, B.P., Huynh, W., Chalil, R., Smith, K.W., Raphenya, A.R., Wlodarski, M.A., Edalatmand, A., Petkau, A., Syed, S.A., Tsang, K.K., *et al.*: Card 2023: expanded curation, support for machine learning, and resistome prediction at the comprehensive antibiotic resistance database. Nucleic acids research **51**(D1), 690–699 (2023)

[33] Liu, B., Zheng, D., Zhou, S., Chen, L., Yang, J.: Vfdb 2022: a general classification scheme for bacterial virulence factors. Nucleic acids research **50**(D1), 912–917 (2022)

[34] Mahajna, A., Dinkla, I.J., Euverink, G.J.W., Keesman, K.J., Jayawardhana, B.: Clean and safe drinking water systems via metagenomics data and artificial intelligence: state-of-the-art and future perspective. Frontiers in microbiology **13**, 832452 (2022)

[35] Sudarshan, A.S., Dai, Z., Gabrielli, M., Oosthuizen-Vosloo, S., Konstantinidis, K.T., Pinto, A.J.: New drinking water genome catalog identifies a globally distributed bacterial genus adapted to disinfected drinking water systems. Environmental Science & Technology **58**(37), 16475–16487 (2024)

[36] Molina, C.A., Quiroz-Moreno, C., Jarrín-V, P., Díaz, M., Yugsi, E., Pérez-Galarza, J., Baldeón-Rojas, L.: Bacterial community assessment of drinking water and downstream distribution systems in highland localities of ecuador. Journal of Water and Health **22**(3), 536–549 (2024)

[37] Vries, H.J., Beyer, F., Jarzembowska, M., Lipińska, J., Brink, P., Zwijnenburg, A., Timmers, P.H., Stams, A.J., Plugge, C.M.: Isolation and characterization of sphingomonadaceae from fouled membranes. npj Biofilms and Microbiomes **5**(1), 6 (2019)

[38] Vaz-Moreira, I., Nunes, O.C., Manaia, C.M.: Diversity and antibiotic resistance patterns of sphingomonadaceae isolates from drinking water. Applied and environmental microbiology **77**(16), 5697–5706 (2011)

[39] Fujita, M., Sakumoto, T., Tanatani, K., Yu, H., Mori, K., Kamimura, N., Masai, E.: Iron acquisition system of sphingobium sp. strain syk-6, a degrader of lignin-derived aromatic compounds. Scientific Reports **10**(1), 12177 (2020)

[40] Kong, Y., Zhang, R., Blain, S., Obernosterer, I.: Seasonal dynamics in microbial trace metals transporters during phytoplankton blooms in the southern ocean. Environmental Microbiology **26**(10), 16695 (2024)

[41] Brikun, I.A., Reeves, A.R., Cernota, W.H., Luu, M.B., Weber, J.M.: The erythromycin biosynthetic gene cluster of aeromicrobium erythreum. Journal of Industrial Microbiology and Biotechnology **31**(7), 335–344 (2004)

[42] Miller, E.S.: Cloning vectors, mutagenesis, and gene disruption (ermr) for the erythromycin-producing bacterium aeromicrobium erythreum. Applied and environmental microbiology **57**(9), 2758–2761 (1991)