Separable Dorsal Raphe Dopamine Projections Mimic the Facets of a Loneliness-like State

Christopher R. Lee[1,2,3,6], Gillian A. Matthews[1,4,6], Mackenzie E. Lemieux[1,4,6], Elizabeth M. Wasserlein[4], Matilde Borio[1,4], Raymundo L. Miranda[1,2,4], Laurel R. Keyes[1,3], Gates P. Schneider[1,3], Caroline Jia[1,2,3], Andrea Tran[1], Faith Aloboudi[1], May G. Chan[1,3], Enzo Peroni[4], Grace S. Pereira[4], Alba López-Moraga[4], Anna Pallé[1,4], Eyal Y. Kimchi[4], Nancy Padilla-Coreano[1,4], Romy Wichmann[1,3,4], Kay M. Tye[1,2,3,4,5,*]

[1]Salk Institute for Biological Studies, 10010 N Torrey Pines Rd, La Jolla, CA 92037, USA.

[2]Neurosciences Graduate Program, University of California San Diego, La Jolla, CA 92093, USA

[3]Howard Hughes Medical Institute, 10010 N Torrey Pines Rd, La Jolla, CA 92037, USA.

[4]The Picower Institute for Learning and Memory, Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Cambridge, MA 02139, USA.

[5]Kavli Institute for Brain and Mind, La Jolla, CA, USA

[6]These authors contributed equally.

*Address Correspondence to: tye@salk.edu (K.M.T.).

Abstract

Affiliative social connections facilitate well-being and survival in numerous species. Engaging in social interactions requires positive or negative motivational drive, elicited through coordinated activity across neural circuits. However, the identity, interconnectivity, and functional encoding of social information within these circuits remains poorly understood. Here, we focus on downstream projections of dorsal raphe nucleus (DRN) dopamine neurons (DRN[DAT]), which we previously implicated in social motivation alongside an aversive affective state. We show that three prominent DRN[DAT] projections – to the bed nucleus of the stria terminalis (BNST), central amygdala (CeA), and posterior basolateral amygdala (BLP) – play separable roles in behavior, despite substantial collateralization. Photoactivation of the DRN[DAT]-CeA projection promoted social behavior and photostimulation of the DRN[DAT]-BNST projection promoted exploratory behavior, while the DRN[DAT]-BLP projection supported place avoidance, suggesting a negative affective state. Downstream regions showed diverse receptor expression, poising DRN[DAT] neurons to act through dopamine, neuropeptide, and glutamate transmission. Furthermore, we show *ex vivo* that the effect of DRN[DAT] photostimulation on downstream neuron excitability depended on region and baseline cell properties, resulting in excitatory responses in BNST cells and diverse responses in CeA and BLP. Finally, *in vivo* microendoscopic cellular-resolution recordings in the CeA with DRN[DAT] photostimulation revealed a correlation between social behavior and neurons excited by social stimuli– suggesting that increased dopamine tone may recruit different CeA neurons to social ensembles. Collectively, these circuit features may facilitate a coordinated, but flexible, response in the presence of social stimuli that can be flexibly guided based on the internal social homeostatic need state of the individual.

INTRODUCTION

A close social network confers a survival advantage, both in the wild and in the laboratory[1–3]. Indeed, our brains have evolved to adapt to many changing conditions, including when we are with others and when we are alone. Many neuromodulatory systems and neural circuits engaged in social behaviors may serve a distinct function when social stimuli are not present. In non-social contexts, dopamine transporter-expressing dorsal raphe nucleus (DRN^{DAT}) neurons can promote incentive memory expression[4], antinociception[5–7], fear response[8], and arousal[9–11] – showing a clear role in many functions essential for survival. Moreover, DRN^{DAT} neurons undergo synaptic strengthening after social isolation and increase responsiveness to social stimuli, and stimulation of these neurons induces a prosocial state[12]. Strikingly, a functional imaging study in humans similarly revealed that 10 hours of social isolation heightened midbrain responses to social stimuli[13]. In mice, we further demonstrated that photostimulation of DRN^{DAT} neurons not only promoted social preference, but also induced place avoidance, suggesting an aversive internal state[12]. This led us to infer a role for these neurons in motivating social approach, driven by the desire to quell a negative state[14], and playing a role in social homeostasis[15,16].

Taken together, this suggests a broad functional role for DRN^{DAT} neurons in motivating adaptive, survival-promoting behaviors under both social and non-social conditions. While the multi-functional role of dopamine neurons in the DRN seems clear, it is yet unclear how these cells exert their influence at a circuit level, and the question remains: how do DRN^{DAT} neurons simultaneously motivate social approach while also inducing a negative state consistent with place avoidance? What downstream targets receive this signal, and how do they respond?

There are several circuit motifs and neural encoding strategies that could enable DRN^{DAT} neurons to simultaneous regulate these behavioral states and motivate adaptive responses. In a drive-state sequence model, if these DRN^{DAT} neurons were the control center in the social homeostat[15,16], the unpleasant state of being isolated could then feed-forward in a sequential chain to induce motivation to rectify this social deficit. However, in an effector state activation model, many parallel actions may be taken to address the challenge, and a pervasive behavioral state may be triggered by a neuromodulatory broadcast signal. In a parallel circuit model, distinct functional roles may be associated with projection-defined subpopulations in parallel (e.g.[17–23]), and neurons may simultaneously encode multiple types of information (i.e. exhibit 'mixed selectivity'[24,25]) or behavioral output may be governed by context- or state-dependency (e.g.[26–29]). Yet, the mechanisms through which DRN^{DAT} neurons exert their influence over social behavior has yet to be unraveled.

Here, we addressed the question of how DRN^{DAT} neurons modulate both sociability and valence, by exploring the functional role and anatomical targets of distinct DRN^{DAT} projections in mice. We show that parallel DRN^{DAT} projections to different targets play separable roles in behavior, in spite of their heavily-collateralizing anatomical arrangement. Downstream, we find that within DRN^{DAT} terminal fields, there is spatial segregation of dopamine and neuropeptide receptor expression. Furthermore, photostimulation of DRN^{DAT} inputs can modulate downstream neuronal excitability depending on their baseline cell properties. Lastly, we find that DRN^{DAT} input enables a shift in central amygdala dynamics that allows it to predict social preference. These findings highlight the anatomical and functional heterogeneity that exists at multiple levels within the DRN^{DAT} system. We suggest this organization may underlie the capacity of the DRN^{DAT} system to exert a broad influence over different

forms of behavior: allowing coordinated control over downstream neuronal activity and across the brain
to signal a behavioral state that mimics a loneliness-like phenotype.

RESULTS

DRN^{DAT} neurons project to and exhibit dense collateralization to distinct subregions of the
amygdala and extended amygdala.

To explore the circuit motifs[30] and computational implementation[31] through which the DRN^{DAT}
system might operate, we examined whether discrete DRN^{DAT} projections underlie distinct features of
behavior. Prominent DRN^{DAT} projections were identified by quantifying downstream fluorescence
following Cre-dependent expression of eYFP in dopamine transporter (DAT)::IRES-Cre mice[12,32–
34](Figure 1—figure supplement 1). We observed a distinct pattern of innervation arising from ventral
tegmental area (VTA)^{DAT} and DRN^{DAT} subpopulations (Figure 1A-D), with DRN^{DAT} projections most
densely targeting the oval nucleus of the BNST (ovBNST) and lateral nucleus of the central amygdala
(CeL). We also observed weaker, but significant, input to the posterior part of the basolateral amygdala
(BLP), consistent with previous tracing studies[4,33,35–37]. Given that the extended amygdala and
basolateral amygdala complex have been implicated in aversion-[38–41] and reward-related
processes[21,42–47], and connect with hindbrain motor nuclei to elicit autonomic and behavioral changes,
we focused on these DRN^{DAT} projections (Figure 1D).

We next considered the anatomical organization of these projections to determine whether form
gives rise to function. In other words, we investigated whether DRN^{DAT} outputs exhibit a circuit
arrangement that facilitates a coordinated behavioral response. Axonal collateralization is one circuit
feature that facilitates coordinated activity across broadly distributed structures[48]. Although VTA^{DAT}
projections to striatal and cortical regions typically show little evidence of collateralization[49–53], in
contrast, DRN serotonergic neurons collateralize heavily to innervate the prefrontal cortex, striatum,
midbrain, and amygdala[54–56]. However, it has yet to be determined whether DRN^{DAT} neurons are
endowed with this property.

To assess whether DRN^{DAT} neurons exhibit axon collaterals, we performed dual retrograde
tracing with fluorophore-conjugated cholera toxin subunit B (CTB)[57]. We injected each tracer into two of
the three downstream sites (BNST, CeA, and/or BLP) (Figure 1E-F and Figure 1—figure supplement
2A-C) and, after 7 days for retrograde transport, we examined CTB-expressing cells in the DRN that
were co-labelled with tyrosine hydroxylase (TH; Figure 1G). CTB injections into the BNST and CeA
resulted in numerous TH+ cells labelled with both CTB-conjugated fluorophores, but fewer dual-labelled
cells were observed when injections were placed in the BNST and BLP, or CeA and BLP (Figure 1H
and Figure 1—figure supplement 2D-E). These data suggest significant collateralization to the
extended amygdala, which includes the BNST and CeA[4,40]. To confirm the presence of axon collaterals
we employed an intersectional viral strategy to selectively label CeA-projecting DRN^{DAT} neurons with
cytoplasmic eYFP (Figure 1—figure supplement 2F-G). This resulted in eYFP-expressing terminals
both in the CeA and in the BNST (Figure 1—figure supplement 2H-I).

DRN^{DAT}-BLP photostimulation promotes place avoidance.

We next considered whether DRN^{DAT} projections to the BNST, CeA, and BLP play separable or overlapping functional roles in modulating behavior. VTA dopaminergic input to the BNST and CeA has been implicated in threat discrimination [58,59], anxiety-related behavior [60], and drug-induced reward [61–64], while in the BLA complex, dopamine signaling supports both fear [65–68] and appetitive learning [46,69]. However, the question remains: do the same DRN^{DAT} projection neurons mediate different facets of a loneliness-like state, such as aversion, vigilance, and social motivation?

To test the hypothesis that distinct DRN^{DAT} projections promote sociability, vigilance, and place avoidance[12], we performed projection-specific ChR2-mediated photostimulation. We injected an AAV enabling Cre-dependent expression of ChR2 into the DRN of DAT::Cre male mice, and implanted optic fibers over the BNST, CeA, or BLP (Figure 2A and Figure 2—figure supplement 1A-F). Given that we previously observed that behavioral effects of DRN^{DAT} photostimulation were predicted by an animal's social rank[12], we also assessed relative social dominance using the tube test[70–72] prior to behavioral assays and photostimulation (Figure 2A and Figure 2—figure supplement 1G-H).

We first assessed whether photostimulation was sufficient to support place preference using the real-time place-preference (RTPP) assay. Here, we found that photostimulation of the DRN^{DAT}-BLP projection, but not the projection to the BNST or CeA, produced avoidance of the stimulation zone, relative to eYFP controls (Figure 2B-G). However, we did not find a significant correlation between social dominance and the magnitude of this effect (Figure 2H-J). Importantly, we did not detect an effect of photostimulation of DRN^{DAT} projections on operant intracranial self-stimulation (Figure 2—figure supplement 2).

DRN^{DAT}-BNST photostimulation promotes non-social exploration.

Next, we considered whether DRN^{DAT} projections to the BNST, CeA, or BLP play a role in increasing vigilance, a common behavioral marker in individuals experiencing loneliness[73,74]. To assess how projection-specific photostimulation of DRN^{DAT} terminals affected exploratory behavior, we used the open field test (OFT) and elevated plus maze (EPM). While we found no effect of optical stimulation of DRN^{DAT} terminals on locomotion or time in center in the OFT (Figure 3—figure supplement 1), we found stimulation of DRN^{DAT} terminals in the BNST (but not in the CeA or BLP) resulted in a weak trend toward increased time spent in the open arm of the EPM (Figure 3A-C), which can be interpreted as exploratory behavior linked with a vigilant state[75]. However, we found no correlation between social dominance and open arm time (Figure 3D-F). Strikingly, during social interaction with a novel juvenile in the home-cage, we found that photoactivation of the DRN^{DAT}-BNST projection increased rearing behavior (a form of nonsocial exploration[76,77]; Figure 3G-L), an effect that was not previously observed with cell body photostimulation[12]. However, we did not find a significant correlation between social dominance and the expression of optically-induced rearing behavior (Figure 3J-L).

DRN^{DAT}-CeA photostimulation promotes sociability.

To assess how projection-specific photostimulation of DRN^{DAT} terminals affected social preference, we used the three chamber sociability task[78], where group-housed mice freely-explored a

chamber containing a novel juvenile mouse and a novel object at opposite ends (Figure 4A-C). This
revealed that optical stimulation of the DRN[DAT]-CeA projection increased social preference, but no
significant effect was observed with photostimulation of either the DRN[DAT]-BNST or DRN[DAT]-BLP
projections (Figure 4D-F; Figure 4—figure supplement 1). Furthermore, we found that the optically-
induced change in social preference in DRN[DAT]-CeA mice was positively correlated with social
dominance, suggesting that photostimulation elicited a greater increase in sociability in dominant mice
(Figure 4G-I). This emulates the previous association found with photostimulation at the cell body level
and social dominance[12].

Next, to gain further insight into the functional divergence of DRN[DAT] projections in ethological
behaviors, we assessed the effects of photostimulation on social interaction with a novel juvenile in the
home-cage. Here, photoactivation of the DRN[DAT]-CeA projection modestly increased face sniffing of the
juvenile mouse, consistent with a pro-social role for this projection (Figure 4J-L), although no correlation
between optically-induced change in face sniffing and social dominance was observed (Figure 4M-O).
When we plotted the difference score (ON-OFF) for face sniffing against rearing (ON-OFF) (Figure 4—
figure supplement 2A-C), we observed that DRN[DAT]-BNST mice tended to engage in more rearing and
less face sniffing during photostimulation (i.e. located in the upper left quadrant) whereas DRN[DAT]-CeA
mice tended to exhibit less rearing and more face sniffing during photostimulation (i.e. located in the
lower right quadrant).

To explore the relationship between social dominance and baseline behavioral profile, we
applied a data-driven approach by examining behavioral measures obtained from different assays in a
correlation matrix (Figure 4—figure supplement 2D). This showed a weak, negative correlation between
social dominance and open arm time in the elevated plus maze (EPM) – consistent with a previous
report of higher trait anxiety in dominant mice[79]. However, social dominance did not correlate
significantly with any other behavioral variable. Additionally, our analysis of baseline behavioral profile
revealed a robust negative correlation between the time spent engaged in social sniffing and time spent
rearing (Figure 4—figure supplement 2D). Furthermore, following dimensionality reduction on baseline
behavioral variables, we did not find clearly differentiated clusters of high- and low-ranked mice (Figure
4—figure supplement 2E), suggesting that the variation governing these latent behavioral features is
not related to social rank.

Finally, to determine whether DRN[DAT]-CeA photostimulation affected the probability of
behavioral state transition[80,81], we examined the sequential structure of behavior using a First-order
Markov model [81,82]. Considering a 2-state model consisting of 'social' and 'nonsocial' behaviors (Figure
4P), we found that photostimulation in DRN[DAT]-CeA mice did not significantly change the probability of
transitioning within social and nonsocial state (Figure 4Q), but did significantly change the probability of
transitioning between social and nonsocial states (Figure 4R). This suggests that the DRN[DAT]-CeA
projection may increase engagement in social behavior by altering the overall structure of behavioral
transitions.

DRN[DAT] terminal fields contain spatially-segregated dopamine and neuropeptide receptor
populations.

Our data suggest that DRN[DAT] projections exert divergent effects over behavior, despite
substantial overlap in their upstream cells of origin. Given this overlap, we reasoned that one

mechanism through which these projections might achieve distinct behavioral effects is via differential
recruitment of downstream signaling pathways. We, therefore, next considered whether the pattern of
receptor expression differed within the DRN[DAT] terminal field of these downstream regions.

Subsets of DRN[DAT] neurons co-express vasoactive intestinal peptide (VIP) and neuropeptide-W
(NPW)[83–85], and so we examined both dopamine (*Drd1* and *Drd2*) and neuropeptide (*Vipr2* and
Npbwr1) receptor expression within DRN[DAT] terminal fields. To achieve this, we performed single
molecule fluorescence *in situ* hybridization (smFISH) using RNAscope (Figure 5—figure supplement
1A-B). In the BNST and CeA we observed a strikingly similar pattern of receptor expression with dense
neuropeptide receptor expression in the oval BNST and ventromedial CeL, and a high degree of co-
localization (Figure 5A-H and Figure 5—figure supplement 1C-H). In the BNST and CeA subregions
containing the highest density of DRN[DAT] terminals, dopamine receptor expression was relatively more
sparse, with *Drd2* more abundant than *Drd1*, as previously described[44,58,60,86,87] (Figure 5A-H). The
DRN[DAT] terminal field of the BLP displayed a markedly different receptor expression pattern, dominated
by *Drd1* (Figure 5I-L and Figure 5—figure supplement 1I-K), consistent with previous reports[60,69,86].
Thus, in contrast to the BNST and CeA, the effects of DRN[DAT] input to the BLP may be predominantly
mediated via D$_1$-receptor signaling. Collectively, this expression pattern suggests that the dopamine-
and neuropeptide-mediated effects of DRN[DAT] input may be spatially-segregated within downstream
regions – providing the infrastructure for divergent modulation of cellular subsets.

224 DRN[DAT] input has divergent effects on downstream cellular excitability.

Our data suggest that DRN[DAT] projections exert divergent effects over behavior, despite
substantial overlap in their upstream cells of origin. One mechanism through which these projections
might achieve distinct behavioral effects is via differential modulation of activity in downstream neurons.
The multi-transmitter phenotype of DRN[DAT] neurons[83,84,88,89], regionally-distinct downstream receptor
expression, and the observed pre- and post-synaptic actions of exogenously applied dopamine[90–97]
provides optimal conditions for diverse modulation of neural activity. However, it remains unknown how
temporally-precise activation of DRN[DAT] terminals influences excitability at the single-cell level.

We, therefore, next examined how DRN[DAT] input affects downstream excitability. To achieve
this, we expressed ChR2 in DRN[DAT] neurons, and used *ex vivo* electrophysiology to record from
downstream neurons (Figure 6A-C and Figure 6—figure supplement 1A-C). Optical stimulation at the
resting membrane potential evoked both excitatory and inhibitory post-synaptic potentials (EPSPs and
IPSPs) in downstream cells (Figure 6D-F), which were typically monosynaptic (Figure 6—figure
supplement 1D-E). During spontaneous firing, BNST cells were universally excited whereas more
diverse responses were observed with the BLP and CeA (Figure 6G-K and Figure 6—figure
supplement 1F-G). The fast rise and decay kinetics of the EPSP suggest an AMPAR-mediated
potential, resulting from glutamate co-release[5,12], whereas the slow IPSP kinetics are consistent with
opening of GIRK channels, which can occur via D$_2$-receptor[98,99] or GABA-$_B$ receptor signaling[100–102].

Given the diversity of responses observed in the CeA and BLP, we next examined these
downstream cells in more detail. To assess whether baseline electrophysiological properties predicted
the optically-evoked response, we used unsupervised agglomerative hierarchical clustering to classify
downstream cells (Figure 6L-M). This established approach has been successfully applied to
electrophysiological datasets to reveal distinct neuronal subclasses[103–105]. The resulting dendrograms

yielded two major clusters in the CeA and BLP, with distinct electrophysiological characteristics (Figure
6N-Q and Figure 6—figure supplement 1H-K). CeA cells in cluster 1 represented 'late-firing' neurons,
whereas cluster 2 were typical of 'regular-firing' neurons[106–108]. Strikingly, these clusters exhibited
dramatically different response to DRN[DAT] photostimulation, with cluster 1 'late-firing' neurons excited
and cluster 2 'regular-firing' neurons mostly inhibited (Figure 6O). Similarly, BLP cells delineated into
two major clusters, with properties characteristic of pyramidal neurons (cluster 1) and GABAergic
interneurons (cluster 2) (Figure 6P-Q). These clusters showed remarkably different responses to
DRN[DAT] input, with 93% of putative pyramidal neurons showing an inhibitory response, and 62% of
putative GABAergic interneurons showing an excitatory response (Figure 6Q). In addition, clustering
CeA and BLP cells together yielded a very similar result (Figure 6—figure supplement 1L-N). Thus,
while photoactivation of DRN[DAT] terminals elicits heterogeneous responses in downstream neurons,
baseline cell properties strongly predict their response, suggesting robust synaptic organization. The
opposing nature of these responses, in different neuronal subsets, suggests that – rather than inducing
an overall augmentation or suppression of activity – DRN[DAT] input may adjust the *pattern* of
downstream activity, in order to exert a functional shift in behavior.

DRN[DAT] input enables a functional shift in CeA dynamics to predict social preference.

Our data thus far suggest that photostimulation of DRN[DAT] projections to downstream extended
amygdala targets elicits divergent behaviors that are, together, congruent with a loneliness-like state,
with the DRN[DAT]-CeA projection promoting sociability. Considering the diversity of responses in the
CeA elicited by DRN[DAT] input *ex vivo*, we next wondered how DRN[DAT] input into the CeA *in vivo* during a
behaviorally-relevant task may modify how the CeA represents social information. Neuromodulatory
input has been previously shown to alter responses to salient stimuli—for instance, stimulation of VTA
dopamine terminals increases the signal-to-noise ratio to aversive stimuli in projection-specific
populations of prefrontal cortex neurons[109]. However, how DRN[DAT] input modifies the coding scheme of
CeA neurons for social information remains unknown.

Therefore, to test the hypothesis that DRN[DAT] input alters the responses of CeA neurons to
functionally-relevant stimuli, we examined the dynamics of CeA neurons while simultaneously
stimulating DRN[DAT] terminals during a three chamber sociability task. To achieve this, we expressed the
calcium indicator GCaMP7f nonspecifically in the CeA and either the red-shifted opsin ChrimsonR or a
control fluorophore (TdTomato) in the DRN of DAT::Cre mice, and additionally implanted a gradient
index (GRIN) lens over the CeA (Figure 7A and Figure 7—figure supplement 1A). This allowed us to
stimulate DRN[DAT] terminals in the CeA while resolving single-cell calcium dynamics in the CeA *in vivo*
(Figure 7B). We confirmed *ex vivo* that blue light delivery alone onto DRN[DAT] terminals did not elicit a
ChrimsonR-mediated postsynaptic potential in CeA neurons (Figure 7—figure supplement 1B-F), and
that red light delivery was still capable of eliciting ChrimsonR-mediated EPSPs and IPSPs during
continuous delivery of blue light (Figure 7—figure supplement 1G-I).

We then performed microendoscopic epifluorescent calcium imaging during the three-chamber
sociability task where mice freely-explored a chamber containing a novel juvenile mouse and a novel
object at opposite ends. Given that social isolation produces changes in long-term potentiation of
synapses onto DRN[DAT] neurons[12], we were limited to a single manipulation of social isolation for each
mouse. We hypothesized that stimulation of DRN[DAT] inputs to CeA would mimic a loneliness-like state,

consistent with our ChR2 manipulations with group-housed mice. Thus, we compared three conditions
—group-housed without DRN[DAT] stimulation (GH off), group-housed with DRN[DAT] stimulation (GH on),
and 24 hours socially isolated without DRN[DAT] stimulation (SI off) (Figure 7C-D) to allow for within-
subjects comparisons.

In contrast to the photostimulation experiments in Figure 4, here we aimed to investigate the
impact of DRN[DAT] neuron stimulation on neural dynamics within the CeA without inducing robust
behavioral changes that could introduce sensorimotor confounds to changes in neural activity due to
stimulation. We successfully optimized viral expression and illumination parameters to minimize
changes in social preference with DRN[DAT]-CeA with ChrimsonR to prioritize comparison of the neural
dynamics (Figure 7E) and also did not observe any behavioral effects of illumination in TdTomato
expressing mice (Figure 7—figure supplement 1J-K).

We then aligned the recorded CeA calcium traces with social cup and object cup interactions
and found a striking diversity of neuronal responses to these stimuli under the three experimental
conditions (Figure 7—figure supplement 2A-B). We next determined the response strength of individual
CeA neurons to either stimulus under the three conditions (Figure 7F-G) using an area under ROC
curve-based approach[110,111] to determine responsiveness of CeA neurons to social and object stimuli
(Figure 7—figure supplement 1M). At a single-cell level, we did not observe significant changes in CeA
response strength or proportion of neurons significantly responding to social or object stimuli across the
three conditions (Figure 7G-H). However, we did find a trend indicating stronger responses toward
social stimuli compared to object stimuli in the GH on condition compared to the GH off condition in
mice expressing ChrimsonR (Figure 7J), but not TdTomato (Figure 7—figure supplement 1L) in DRN[DAT]
neurons. Importantly, in co-registered neurons, we found little overlap between CeA neurons excited
by the social stimulus in both GH on and GH off conditions (Figure 7I), suggesting that DRN[DAT] terminal
stimulation may recruit separate ensembles of CeA neurons to represent social stimuli. Considering the
variability in social preference behavior across mice and the diverse effects of photostimulation
depending on the mouse's social history, we next considered the responses of CeA neurons to social
and object stimuli on an animal-by-animal basis. While we did not observe significant changes in the
proportion of excitatory or inhibitory responses to social or object stimuli across the three conditions
(Figure 7K-L), we did find a significant positive correlation between the proportion of socially-excited
CeA neurons and social preference in the GH on condition, but not in the GH off or SI off conditions
(Figure 7M). Importantly, we do not observe a correlation between social preference and object-excited
CeA neurons (Figure 7—figure supplement 1N) or socially-inhibited CeA neurons (Figure 7—figure
supplement 1O). This result may suggest that DRN[DAT] input in the CeA in a behaviorally-relevant task
allows for a functional shift in its dynamics that enables it to predict the amount of social preference the
mouse exhibits.

DRN[DAT]-CeA photoinhibition blocks isolation-induced sociability

Finally, considering that the DRN[DAT]-CeA projection is sufficient in promoting sociability, we next
assessed whether activity in the DRN[DAT]-CeA projection is necessary for the rebound in sociability that
occurs following acute social isolation[12]. We injected an AAV enabling Cre-dependent expression of
NpHR into the DRN of DAT::Cre male mice, and implanted optic fibers over the BNST, CeA, or BLP
(Figure 8A). We allowed 7 weeks for adequate terminal expression, after which we inhibited DRN[DAT]

terminals in the BNST, CeA, and BLP while mice performed the three-chamber sociability task (Figure
8B). Inhibition of DRN[DAT] terminals in downstream regions while mice were group-housed did not
change social preference (Figure 8—figure supplement 1A-B). However, inhibition of DRN[DAT] terminals
in CeA, but not BNST or BLP, blocked the rebound in sociability associated with acute social isolation
(Figure 8C). Additionally, we found that optically-inhibited changes in social preference in DRN[DAT]-CeA
mice were negatively correlated with social dominance (Figure 8D), suggesting that the DRN[DAT]-CeA
projection is necessary for the expression of isolation-induced social rebound in a rank-dependent
manner.

DISCUSSION

Neural circuits that motivate social approach are essential in maintaining social connections and
preventing isolation. Here we show that DRN[DAT] neurons can exert a multi-faceted influence over
behavior, with the pro-social effects mediated by the projection to the CeA, the avoidance effects
mediated by the projection to the BLP, and the pro-exploratory effects mediated by the projection to the
BNST. Our data suggest these effects are enabled via separable functional projections, dense
collateralization, co-transmission, and precisely-organized synaptic connectivity. Notably, our
experiments were conducted in male mice; future work should investigate whether similar circuit
mechanisms operate in females and explore the biological basis of sex-specific responses to social
isolation[112,113]. In addition, while terminal photostimulation allowed projection-specific manipulation, it
may have inadvertently activated fibers of passage—a limitation that could be addressed in future
studies using intersectional genetic strategies. Despite these caveats, these observed circuit features
may facilitate a coordinated, but flexible, response in the presence of social stimuli, that can be flexibly
guided based on internal social homeostatic need state.

DRN[DAT] circuit arrangement enables a broadly distributed, coordinated response.

Our findings revealed several features of the DRN[DAT] circuit which might facilitate a concerted
response to novel social and non-social situations. Firstly, we observed dissociable roles for discrete
downstream projections – a common motif of valence-encoding neural circuits[30]. Biased recruitment of
these 'divergent paths'[30] to the BNST, CeA, and BLP by upstream inputs may serve to fine-tune the
balance between social investigation and environmental exploration: facilitating behavioral flexibility
with changing environmental conditions or internal state. Secondly, we demonstrate extensive
collateralization of DRN[DAT] neurons In other populations, collateralization is proposed to aid temporal
coordination of a multifaceted response: enabling synchronous activation of distributed regions[48,56].
This feature may, therefore, facilitate coordinated recruitment of the BNST and CeA, allowing these
regions to work in concert to promote social approach while also maintaining vigilance to salient
environmental stimuli. Thirdly, we find precise synaptic organization in the DRN[DAT] modulation of
downstream neuronal activity that allows for qualitatively distinct response profiles of downstream
targets.   Combined with the spatially-segregated downstream receptor expression pattern, this
organization may allow DRN[DAT] neurons to elicit broad, yet finely-tuned, control over the pattern of
neuronal activity, on multiple timescales—perhaps explaining the diverse behavior effects between the
BNST- and CeA- projecting DRN[DAT] populations, despite heavy collateralization.

Although we hypothesized that stimulating DRN[DAT] inputs to the CeA in group-housed mice
would mimic a state similar to that of isolation, we did not observe that the isolation OFF condition
produced neural responses more similar to the group-housed ON condition. This suggests two
possibilities: 1) that the photostimulation impacted neural activity beyond the endogenous dopamine
innervation that may occur with social isolation because it is more potent of a change or 2) that the
timing of endogenous dopamine innervation is different and is partially quenched upon exposure to a
social agent. To completely understand the temporal dynamics of dopamine signaling with isolation
and the firing of DRN[DAT] neurons upon isolation, further experiments will require exploration of DRN[DAT]
stimulation parameter space, endogenous neural activity in the DRN[DAT]-CeA circuit during social
isolation, and the effects of DRN[DAT] stimulation timing.

Separable projections mediate social behavior and valence.

Our data support the hypothesis that separable DRN[DAT] projections mediate distinct functional
roles: a feature which has been previously observed in other neuronal circuits (e.g.[17–21]). The DRN[DAT]
circuit attributes we describe above may further enable this system to modulate other diverse forms of
behavior (e.g. arousal[9], fear/reward associations[4,8], and antinociception[5–7]). These could be mediated
via other downstream projections and/or via these same projections under different environmental
contexts, testing conditions, social histories, and/or internal states. Further work is required to
determine how this system is able to exert a broad influence over multiple forms of behavior. Indeed, a
recent study examined DRN[DAT] projection to the nucleus accumbens and its role in promoting
sociability[114], suggesting a parallel circuit to that described in the current study. Collectively, however,
our data and others support a role for the DRN[DAT] system in exerting a coordinated behavioral
response to novel situations – both social and non-social.

The CeA has been implicated in mediating the response to threats – orchestrating defensive
behavioral responses and autonomic changes via efferents to subcortical[38,115,116] and brainstem
nuclei[117]. One possible interpretation, therefore, is that DRN[DAT] input to the CeA suppresses fear-
promoting neuronal ensembles in order to facilitate social approach. In the maintenance of social
homeostasis, suppression of fear in the presence of social stimuli may represent an adaptive response
– preventing salient social stimuli from being interpreted as a threat. Indeed, other need states, such as
hunger, are associated with fear suppression and higher-risk behavior[118], suggesting a conserved
response to homeostatic imbalance[16]. However, the motivation to attend to social stimuli may also be
driven by territorial defense (interacting with social rank), highlighting a need to further understand how
internal states can play into the output of this system. A more comprehensive knowledge of the
functional cell-types modulated by DRN[DAT] activity will facilitate our understanding of how this input can
shape the downstream neuronal representation of social and non-social stimuli.

In contrast to the CeA, photoactivation of the DRN[DAT]-BLP projection produced avoidance of the
stimulation zone, suggesting an aversive state. This differs from the valence-independent role of VTA
dopamine input to the greater BLA complex, wherein dopamine signaling gates synaptic plasticity for
associative learning of both positive and negative valence[46] and responds to salient stimuli predicting
both positive and negative outcomes[69]. However, DRN and VTA axonal fields differ within the BLA
complex, with DRN[DAT] terminals being more concentrated within the BLP, and VTA[DAT] inputs traversing
the LA, BLA and intercalated cells more densely.

While there have been seemingly contradictory reports on the effect of dopamine on excitability
in the BLA[46,65,69], our observations using photostimulation of DRN[DAT] terminals (in short phasic bursts)
are consistent with *in vivo* extracellular recordings combined with electrical stimulation of the
midbrain[95]. One unifying hypothesis is that dopamine induces an *indirect* GABA-mediated suppression
of pyramidal neurons, which may attenuate their response to weak inputs, while *directly* exciting
pyramidal neurons to augment their response to large inputs[92,95]. In this way, amygdala dopamine may
underlie a similar role to cortical dopamine[109,119] – enhancing signal-to-noise ratio to facilitate behavioral
responses to salient stimuli [109].

A similar complexity surrounds dopamine's effects in the BNST. Photostimulation of DRN[DA]-
BNST projections have been shown to have antinociceptive effects on male mice upon formalin
injection into the paw, whereas females do not show this antinociceptive effect and instead display
contextual hyperlocomotion[7]. Given that our study does not include females, there are likely behavioral
and physiological sex differences that begs further investigation. Additionally, prior studies using
optogenetic stimulation[5,7] or exogenously applied dopamine[91,120] report a wide range of outcomes on
BNST excitability, including a higher frequency of inhibitory responses that we observed. Notably, one
study found that higher doses of exogenously applied dopamine decreased the amplitude of GABA$_A$-
IPSC in ovBNST neurons[91], potentially consistent with our results. Interestingly, animals given
intermittent access to sucrose displayed a significant increase in GABA$_A$-IPSCs in ovBNST neurons
following dopamine application[120], raising the possibility that experience-dependent factors shape
dopamine sensitivity in the BNST, paralleling how dopaminergic modulation in the cortex may be state-
dependent[109]. Considering that our *ex* vivo recordings were performed in group housed mice, future
work should test whether social isolation alters ovBNST responses to dopamine, providing a framework
for understanding how environmental states influence dopaminergic neuromodulatory tone.

Multi-transmitter phenotype of DRN[DAT] neurons may permit modulation on different timescales.

DRN[DAT] neurons possess an impressive repertoire of signaling molecules: alongside dopamine
and glutamate subsets of DRN[DAT] neuron express VIP and NPW[83–85]. While there is some partial
segregation of VIP- and NPW-expressing neurons[84], our receptor expression analyses suggest that
these neuropeptides converge on the same neurons in the BNST and CeA. This co-localization is
intriguing, given that *Vipr2* is typically coupled to the excitatory G$_s$-protein[121], while *Npbwr1* is coupled to
the inhibitory G$_i$-protein[122,123]. Therefore, signaling through these receptors may exert opposing actions
on downstream cells. Recruitment of neuropeptidergic signaling pathways may support slower,
sustained downstream modulation, for example, in hunger-mediating hypothalamic Agouti-Related
Peptide (AgRP) neurons, neuropeptide co-release is essential for sustaining feeding behavior[124].
Therefore, a delayed, persistent neuropeptide-mediated signal might enable downstream modulation to
outlive phasic DRN[DAT] activity: promoting behavioral adjustments over longer timescales.

While the functional role of these neuropeptides remains to be elucidated, studies on knockout
mice suggest a role for NPW in social behavior and stress responding[122] [85]. Furthermore, in humans
with a single-nucleotide polymorphism (SNP) of the *NPBWR1* gene (which impairs receptor function)
the perception of fearful/angry faces was more positive and less submissive[125], suggesting a possible
role for NPW signaling in interpreting social signals. Similarly, the function of DRN VIP+ neurons has
received little attention in rodent models, but there has been more focus on the role of VIP in avian

social behavior[126]. Of particular interest, in the rostral arcopallium (a homolog of mammalian
amygdala[127]), VIP binding density is elevated in birds during seasonal flocking[128]. This suggests that
elevated VIP receptor expression may encourage affiliative social grouping behavior in birds[128]. Thus,
NPW and VIP may act in concert with fast glutamate-mediated and slow dopamine-mediated
neurotransmission in the central extended amygdala, to modulate behavior on different timescales.

Together, these findings suggest that NPW and VIP may act in concert with fast glutamate and
slower dopamine signaling to shape social behavior across multiple timescales within the extended
amygdala. Although we observed relatively low expression of *Drd1* in the BNST and CeA compared to
Vipr2 and *Npbwr1*, it remains possible that dopamine's effects are mediated by other receptors, such
as *Drd3, Drd4,* and *Drd5*—warranting further investigation.

CONCLUSION

Together, these findings reveal that DRN^{DAT} projections exhibit substantial functional
specialization, with anatomically distinct pathways modulating different facets of behavior. The DRN^{DAT}-
CeA projection promotes sociability, DRN^{DAT}-BLP input drives avoidance, and DRN^{DAT}-BNST enhances
vigilant exploration, highlighting the diverse roles of this neural circuit in coordinating adaptive
responses to social and environmental contexts. These findings uncover a circuit mechanism through
which DRN^{DAT} projections orchestrate distinct behavioral features of a loneliness-like state, providing a
framework for understanding how neuromodulatory systems guide complex social and emotional
behaviors and suggesting potential targets for therapeutic intervention in affective disorders.

Acknowledgements

K.M.T. is the Wylie Vale Chair at the Salk Institute for Biological Studies, a New York Stem Cell Foundation - Robertson Investigator, and a McKnight Scholar. This work was supported by funding from the JPB Foundation, Alfred P Sloan Foundation, New York Stem Cell Foundation, Klingenstein Foundation, McKnight Foundation, Clayton Foundation, Dolby Family Fund, R01-MH115920 (NIMH), the NIH Director's New Innovator Award DP2-DK102256 (NIDDK), and Pioneer Award DP1-AT009925 (NCCIH). G.A.M was supported by a Postdoctoral Research Fellowship from the Charles A. King Trust. R.L.M. was funded through the MSRP program in the Brains & Cognitive Sciences Department at MIT, supported by the Center for Brains, Minds and Machines (CBMM), and funded by NSF STC award CCF-1231216. E.M.W was supported by a summer scholarship from Johnson & Johnson. We thank C. Leppla, J. Olsen, P. Namburi, V. Barth, J. Wang, K. Batra, A. Brown, and A. Libster for technical advice, all members of the Tye Lab for helpful discussion, and advice from the CellProfiler team at the Broad Institute. We also thank Rachel Neve for the HSV construct, and Charu Ramakrishnan & Karl Deisseroth for AAV_5-fDIO-eYFP.

Author Contributions

K.M.T., G.A.M, and C.R.L. conceptualized the project, designed experiments, supervised experiments and directed data analyses. M.E.L and M.B. performed and analyzed smFISH experiments. G.A.M., M.E.L., C.R.L., C.J., E.M.W., E.P., G.S.P., A.L.-M., and A.P. performed stereotaxic surgeries. G.A.M., C.R.L, E.M.W, E.P, G.S.P, M.E.L., C.J., F.A., A.T., M.B., A.L.-M., R.M., and A.P. ran optogenetic manipulation experiments and analyzed behavioral data. L.R.K. performed Markov model analysis. G.A.M., C.R.L., E.M.W., G.P.S., M.G.C., E.P., G.S.P., A.L.-M., and A.P. performed immunohistochemistry and analyzed images. G.A.M performed *ex vivo* electrophysiology, and N.P.-C., E.Y.K, and R.W. contributed to experimental design and data interpretation. C.R.L performed *in vivo* calcium imaging and analysis. G.A.M., R.W., C.R.L., and L.R.K. reviewed, organized, and prepared the data for data sharing. G.A.M., C.R.L., and K.M.T. wrote the manuscript with review and editing from M.E.L., E.M.W., M.B., R.M., L.R.K., G.P.S., C.J., A.T., F.A., M.G.C., E.P., G.S.P., A.L.-M., A.P., E.Y.K., N.P.-C., and R.W.

1. Yee, J.R., Cavigelli, S.A., Delgado, B., and McClintock, M.K. (2008). Reciprocal Affiliation Among
Adolescent Rats During a Mild Group Stressor Predicts Mammary Tumors and Lifespan.
Psychosomatic Medicine *70*, 1050. https://doi.org/10.1097/PSY.0b013e31818425fb.

2. Koto, A., Mersch, D., Hollis, B., and Keller, L. (2015). Social isolation causes mortality by disrupting
energy homeostasis in ants. Behavioral Ecology and Sociobiology *69*, 583–591.

3. Silk, J.B., Beehner, J.C., Bergman, T.J., Crockford, C., Engh, A.L., Moscovice, L.R., Wittig, R.M.,
Seyfarth, R.M., and Cheney, D.L. (2010). Strong and Consistent Social Bonds Enhance the
Longevity of Female Baboons. Current Biology *20*, 1359–1361.
https://doi.org/10.1016/j.cub.2010.05.067.

4. Lin, R., Liang, J., Wang, R., Yan, T., Zhou, Y., Liu, Y., Feng, Q., Sun, F., Li, Y., Li, A., et al. (2020).
The Raphe Dopamine System Controls the Expression of Incentive Memory. Neuron *106*, 498-
514.e8. https://doi.org/10.1016/j.neuron.2020.02.009.

5. Li, C., Sugam, J.A., Lowery-Gionta, E.G., McElligott, Z.A., McCall, N.M., Lopez, A.J., McKlveen,
J.M., Pleil, K.E., and Kash, T.L. (2016). Mu Opioid Receptor Modulation of Dopamine Neurons in
the Periaqueductal Gray/Dorsal Raphe: A Role in Regulation of Pain. Neuropsychopharmacology
41, 2122–2132. https://doi.org/10.1038/npp.2016.12.

6. Meyer, P.J., Morgan, M.M., Kozell, L.B., and Ingram, S.L. (2009). Contribution of dopamine
receptors to periaqueductal gray-mediated antinociception. Psychopharmacology (Berl.) *204*, 531–
540. https://doi.org/10.1007/s00213-009-1482-y.

7. Yu, W., Pati, D., Pina, M.M., Schmidt, K.T., Boyt, K.M., Hunker, A.C., Zweifel, L.S., McElligott, Z.A.,
and Kash, T.L. (2021). Periaqueductal gray/dorsal raphe dopamine neurons contribute to sex
differences in pain-related behaviors. Neuron *109*, 1365-1380.e5.
https://doi.org/10.1016/j.neuron.2021.03.001.

8. Groessl, F., Munsch, T., Meis, S., Griessner, J., Kaczanowska, J., Pliota, P., Kargl, D., Badurek, S.,
Kraitsy, K., Rassoulpour, A., et al. (2018). Dorsal tegmental dopamine neurons gate associative
learning of fear. Nature Neuroscience *21*, 952–962. https://doi.org/10.1038/s41593-018-0174-5.

9. Cho, J.R., Treweek, J.B., Robinson, J.E., Xiao, C., Bremner, L.R., Greenbaum, A., and Gradinaru,
V. (2017). Dorsal Raphe Dopamine Neurons Modulate Arousal and Promote Wakefulness by
Salient Stimuli. Neuron *94*, 1205-1219.e8. https://doi.org/10.1016/j.neuron.2017.05.020.

10. Lu, J., Jhou, T.C., and Saper, C.B. (2006). Identification of wake-active dopaminergic neurons in
the ventral periaqueductal gray matter. J. Neurosci. *26*, 193–202.
https://doi.org/10.1523/JNEUROSCI.2244-05.2006.

11. Cho, J.R., Chen, X., Kahan, A., Robinson, J.E., Wagenaar, D.A., and Gradinaru, V. (2021). Dorsal
Raphe Dopamine Neurons Signal Motivational Salience Dependent on Internal State, Expectation,
and Behavioral Context. J. Neurosci. *41*, 2645–2655. https://doi.org/10.1523/JNEUROSCI.2690-
20.2021.

12. Matthews, G.A., Nieh, E.H., Vander Weele, C.M., Halbert, S.A., Pradhan, R.V., Yosafat, A.S.,
Glober, G.F., Izadmehr, E.M., Thomas, R.E., Lacy, G.D., et al. (2016). Dorsal Raphe Dopamine
Neurons Represent the Experience of Social Isolation. Cell *164*, 617–631.
https://doi.org/10.1016/j.cell.2015.12.040.

13. Tomova, L., Wang, K.L., Thompson, T., Matthews, G.A., Takahashi, A., Tye, K.M., and Saxe, R.
(2020). Acute social isolation evokes midbrain craving responses similar to hunger. Nat Neurosci
23, 1597–1605. https://doi.org/10.1038/s41593-020-00742-z.

14. Hull, C.L. (1943). Principles of Behavior: An Introduction to Behavior Theory (D. Appleton-Century
Company, Incorporated).

15. Lee, C.R., Chen, A., and Tye, K.M. (2021). The neural circuitry of social homeostasis:
Consequences of acute versus chronic social isolation. Cell *184*, 1500–1516.
https://doi.org/10.1016/j.cell.2021.02.028.

16. Matthews, G.A., and Tye, K.M. (2019). Neural mechanisms of social homeostasis. Annals of the
New York Academy of Sciences *1457*, 5–25. https://doi.org/10.1111/nyas.14016.

17. Han, W., Tellez, L.A., Rangel, M.J., Motta, S.C., Zhang, X., Perez, I.O., Canteras, N.S., Shammah-
Lagnado, S.J., Pol, A.N. van den, and Araujo, I.E. de (2017). Integrated Control of Predatory
Hunting by the Central Nucleus of the Amygdala. Cell *168*, 311-324.e18.
https://doi.org/10.1016/j.cell.2016.12.027.

18. Kim, S.-Y., Adhikari, A., Lee, S.Y., Marshel, J.H., Kim, C.K., Mallory, C.S., Lo, M., Pak, S., Mattis,
J., Lim, B.K., et al. (2013). Diverging neural pathways assemble a behavioural state from separable
features in anxiety. Nature *496*, 219–223. https://doi.org/10.1038/nature12018.

19. Kohl, J., Babayan, B.M., Rubinstein, N.D., Autry, A.E., Marin-Rodriguez, B., Kapoor, V., Miyamishi,
K., Zweifel, L.S., Luo, L., Uchida, N., et al. (2018). Functional circuit architecture underlying
parental behaviour. Nature *556*, 326–331. https://doi.org/10.1038/s41586-018-0027-0.

20. Lammel, S., Ion, D.I., Roeper, J., and Malenka, R.C. (2011). Projection-specific modulation of
dopamine neuron synapses by aversive and rewarding stimuli. Neuron *70*, 855–862.
https://doi.org/10.1016/j.neuron.2011.03.025.

21. Namburi, P., Beyeler, A., Yorozu, S., Calhoon, G.G., Halbert, S.A., Wichmann, R., Holden, S.S.,
Mertens, K.L., Anahtar, M., Felix-Ortiz, A.C., et al. (2015). A circuit mechanism for differentiating
positive and negative associations. Nature *520*, 675–678. https://doi.org/10.1038/nature14366.

22. Senn, V., Wolff, S.B.E., Herry, C., Grenier, F., Ehrlich, I., Gründemann, J., Fadok, J.P., Müller, C.,
Letzkus, J.J., and Lüthi, A. (2014). Long-Range Connectivity Defines Behavioral Specificity of
Amygdala Neurons. Neuron *81*, 428–437. https://doi.org/10.1016/j.neuron.2013.11.006.

23. Tye, K.M., Prakash, R., Kim, S.-Y., Fenno, L.E., Grosenick, L., Zarabi, H., Thompson, K.R.,
Gradinaru, V., Ramakrishnan, C., and Deisseroth, K. (2011). Amygdala circuitry mediating
reversible and bidirectional control of anxiety. Nature *471*, 358–362.
https://doi.org/10.1038/nature09820.

24. Rigotti, M., Barak, O., Warden, M.R., Wang, X.-J., Daw, N.D., Miller, E.K., and Fusi, S. (2013). The
importance of mixed selectivity in complex cognitive tasks. Nature *497*, 585–590.
https://doi.org/10.1038/nature12160.

25. Tian, J., Huang, R., Cohen, J.Y., Osakada, F., Kobak, D., Machens, C.K., Callaway, E.M., Uchida,
N., and Watabe-Uchida, M. (2016). Distributed and Mixed Information in Monosynaptic Inputs to
Dopamine Neurons. Neuron *91*, 1374–1389. https://doi.org/10.1016/j.neuron.2016.08.018.

26. Krzywkowski, P., Penna, B., and Gross, C.T. (2020). Dynamic encoding of social threat and spatial
context in the hypothalamus. eLife *9*, e57148. https://doi.org/10.7554/eLife.57148.

27. Kyriazi, P., Headley, D.B., and Pare, D. (2018). Multi-dimensional Coding by Basolateral Amygdala
Neurons. Neuron *99*, 1315-1328.e5. https://doi.org/10.1016/j.neuron.2018.07.036.

28. Lemos, J.C., Wanat, M.J., Smith, J.S., Reyes, B.A.S., Hollon, N.G., Van Bockstaele, E.J., Chavkin,
C., and Phillips, P.E.M. (2012). Severe stress switches CRF action in the nucleus accumbens from
appetitive to aversive. Nature *490*, 402–406. https://doi.org/10.1038/nature11436.

29. Seo, C., Guru, A., Jin, M., Ito, B., Sleezer, B.J., Ho, Y.-Y., Wang, E., Boada, C., Krupa, N.A.,
Kullakanda, D.S., et al. (2019). Intense Threat Switches Dorsal Raphe Serotonin Neurons to a
Paradoxical Operational Mode. Science *363*, 538–542. https://doi.org/10.1126/science.aau8722.

30. Tye, K.M. (2018). Neural Circuit Motifs in Valence Processing. Neuron *100*, 436–452.
https://doi.org/10.1016/j.neuron.2018.10.001.

31. Lockwood, P.L., Apps, M.A.J., and Chang, S.W.C. (2020). Is There a 'Social' Brain?
Implementations and Algorithms. Trends in Cognitive Sciences *24*, 802–813.
https://doi.org/10.1016/j.tics.2020.06.011.

32. Bäckman, C.M., Malik, N., Zhang, Y., Shan, L., Grinberg, A., Hoffer, B.J., Westphal, H., and Tomac,
A.C. (2006). Characterization of a mouse strain expressing Cre recombinase from the 3'
untranslated region of the dopamine transporter locus. Genesis *44*, 383–390.
https://doi.org/10.1002/dvg.20228.

33. Cardozo Pinto, D.F., Yang, H., Pollak Dorocic, I., de Jong, J.W., Han, V.J., Peck, J.R., Zhu, Y., Liu,
C., Beier, K.T., Smidt, M.P., et al. (2019). Characterization of transgenic mouse models targeting
neuromodulatory systems reveals organizational principles of the dorsal raphe. Nature
Communications *10*, 4633. https://doi.org/10.1038/s41467-019-12392-2.

34. Lammel, S., Steinberg, E.E., Földy, C., Wall, N.R., Beier, K., Luo, L., and Malenka, R.C. (2015).
Diversity of transgenic mouse models for selective targeting of midbrain dopamine neurons. Neuron
85, 429–438. https://doi.org/10.1016/j.neuron.2014.12.036.

35. Hasue, R.H., and Shammah-Lagnado, S.J. (2002). Origin of the dopaminergic innervation of the
central extended amygdala and accumbens shell: a combined retrograde tracing and
immunohistochemical study in the rat. J. Comp. Neurol. *454*, 15–33.
https://doi.org/10.1002/cne.10420.

36. Meloni, E.G., Gerety, L.P., Knoll, A.T., Cohen, B.M., and Carlezon, W.A. (2006). Behavioral and
Anatomical Interactions between Dopamine and Corticotropin-Releasing Factor in the Rat. J.
Neurosci. *26*, 3855–3863. https://doi.org/10.1523/JNEUROSCI.4957-05.2006.

37. Oh, S.W., Harris, J.A., Ng, L., Winslow, B., Cain, N., Mihalas, S., Wang, Q., Lau, C., Kuan, L.,
Henry, A.M., et al. (2014). A mesoscale connectome of the mouse brain. Nature *508*, 207–214.
https://doi.org/10.1038/nature13186.

38. Davis, M., Walker, D.L., Miles, L., and Grillon, C. (2010). Phasic vs Sustained Fear in Rats and
Humans: Role of the Extended Amygdala in Fear vs Anxiety. Neuropsychopharmacology *35*, 105–
135. https://doi.org/10.1038/npp.2009.109.

39. Goode, T.D., and Maren, S. (2017). Role of the bed nucleus of the stria terminalis in aversive
learning and memory. Learn. Mem. *24*, 480–491. https://doi.org/10.1101/lm.044206.116.

40. Janak, P.H., and Tye, K.M. (2015). From circuits to behaviour in the amygdala. Nature *517*, 284–
292. https://doi.org/10.1038/nature14188.

41. Lebow, M.A., and Chen, A. (2016). Overshadowed by the amygdala: the bed nucleus of the stria
terminalis emerges as key to psychiatric disorders. Molecular Psychiatry *21*, 450–463.
https://doi.org/10.1038/mp.2016.1.

42. Douglass, A.M., Kucukdereli, H., Ponserre, M., Markovic, M., Gründemann, J., Strobel, C., Alcala
Morales, P.L., Conzelmann, K.-K., Lüthi, A., and Klein, R. (2017). Central amygdala circuits
modulate food consumption through a positive-valence mechanism. Nature Neuroscience *20*,
1384–1394. https://doi.org/10.1038/nn.4623.

43. Jennings, J.H., Sparta, D.R., Stamatakis, A.M., Ung, R.L., Pleil, K.E., Kash, T.L., and Stuber, G.D.
(2013). Distinct extended amygdala circuits for divergent motivational states. Nature *496,* 224–228.
https://doi.org/10.1038/nature12041.

44. Kim, J., Zhang, X., Muralidhar, S., LeBlanc, S.A., and Tonegawa, S. (2017). Basolateral to Central
Amygdala Neural Circuits for Appetitive Behaviors. Neuron *93*, 1464-1479.e5.
https://doi.org/10.1016/j.neuron.2017.02.034.

45. Tye, K.M., Stuber, G.D., De Ridder, B., Bonci, A., and Janak, P.H. (2008). Rapid strengthening of
thalamo-amygdala synapses mediates cue–reward learning. Nature *453*, 1253–1257.

46. Tye, K.M., Tye, L.D., Cone, J.J., Hekkelman, E.F., Janak, P.H., and Bonci, A. (2010).
Methylphenidate facilitates learning-induced amygdala plasticity. Nature neuroscience *13*, 475–
481.

47. Bayless, D.W., Davis, C.O., Yang, R., Wei, Y., Carvalho, V.M. de A., Knoedler, J.R., Yang, T.,
Livingston, O., Lomvardas, A., Martins, G.J., et al. (2023). A neural circuit for male sexual behavior
and reward. Cell *186*, 3862-3881.e28. https://doi.org/10.1016/j.cell.2023.07.021.

48. Rockland, K.S. (2018). Axon Collaterals and Brain States. Front Syst Neurosci *12*.
https://doi.org/10.3389/fnsys.2018.00032.

49. Aransay, A., Rodríguez-López, C., García-Amado, M., Clascá, F., and Prensa, L. (2015). Long-
range projection neurons of the mouse ventral tegmental area: a single-cell axon tracing analysis.
Front. Neuroanat. *9*. https://doi.org/10.3389/fnana.2015.00059.

50. Beier, K.T., Steinberg, E.E., DeLoach, K.E., Xie, S., Miyamichi, K., Schwarz, L., Gao, X.J., Kremer,
E.J., Malenka, R.C., and Luo, L. (2015). Circuit Architecture of VTA Dopamine Neurons Revealed
by Systematic Input–Output Mapping. Cell *162*, 622–634. https://doi.org/10.1016/j.cell.2015.07.015.

51. Lerner, T.N., Shilyansky, C., Davidson, T.J., Evans, K.E., Beier, K.T., Zalocusky, K.A., Crow, A.K.,
Malenka, R.C., Luo, L., Tomer, R., et al. (2015). Intact-Brain Analyses Reveal Distinct Information
Carried by SNc Dopamine Subcircuits. Cell *162*, 635–647.
https://doi.org/10.1016/j.cell.2015.07.014.

52. Matsuda, W., Furuta, T., Nakamura, K.C., Hioki, H., Fujiyama, F., Arai, R., and Kaneko, T. (2009).
Single Nigrostriatal Dopaminergic Neurons Form Widely Spread and Highly Dense Axonal
Arborizations in the Neostriatum. J Neurosci *29*, 444–453.
https://doi.org/10.1523/JNEUROSCI.4029-08.2009.

53. Moore, R.Y., and Bloom, F.E. (1978). Central catecholamine neuron systems: anatomy and
physiology of the dopamine systems. Annual Review of Neuroscience *1*, 129–169.
https://doi.org/10.1146/annurev.ne.01.030178.001021.

54. Gagnon, D., and Parent, M. (2014). Distribution of VGLUT3 in Highly Collateralized Axons from the
Rat Dorsal Raphe Nucleus as Revealed by Single-Neuron Reconstructions. PLOS ONE *9*, e87709.
https://doi.org/10.1371/journal.pone.0087709.

55. van der Kooy, D., and Hattori, T. (1980). Dorsal raphe cells with collateral projections to the
caudate-putamen and substantia nigra: a fluorescent retrograde double labeling study in the rat.
Brain Research *186*, 1–7. https://doi.org/10.1016/0006-8993(80)90250-4.

56. Waselus, M., Valentino, R.J., and Van Bockstaele, E.J. (2011). Collateralized dorsal raphe nucleus
projections: a mechanism for the integration of diverse functions during stress. J. Chem.
Neuroanat. *41*, 266–280. https://doi.org/10.1016/j.jchemneu.2011.05.011.

57. Beyeler, A., Chang, C.-J., Silvestre, M., Lévêque, C., Namburi, P., Wildes, C.P., and Tye, K.M.
(2018). Organization of Valence-Encoding and Projection-Defined Neurons in the Basolateral
Amygdala. Cell Rep *22*, 905–918. https://doi.org/10.1016/j.celrep.2017.12.097.

58. De Bundel, D., Zussy, C., Espallergues, J., Gerfen, C.R., Girault, J.-A., and Valjent, E. (2016).
Dopamine D2 receptors gate generalization of conditioned threat responses through mTORC1
signaling in the extended amygdala. Mol Psychiatry *21*, 1545–1553.
https://doi.org/10.1038/mp.2015.210.

59. Jo, Y.S., Heymann, G., and Zweifel, L.S. (2018). Dopamine Neurons Reflect the Uncertainty in
Fear Generalization. Neuron *100*, 916-925.e3. https://doi.org/10.1016/j.neuron.2018.09.028.

60. Perez de la Mora, M., Gallegos-Cari, A., Crespo-Ramirez, M., Marcellino, D., Hansson, A.C., and
Fuxe, K. (2012). Distribution of dopamine D(2)-like receptors in the rat amygdala and their role in
the modulation of unconditioned fear and anxiety. Neuroscience *201*, 252–266.
https://doi.org/10.1016/j.neuroscience.2011.10.045.

61. Eiler, W.J.A., Seyoum, R., Foster, K.L., Mailey, C., and June, H.L. (2003). D1 dopamine receptor
regulates alcohol-motivated behaviors in the bed nucleus of the stria terminalis in alcohol-preferring
(P) rats. Synapse *48*, 45–56. https://doi.org/10.1002/syn.10181.

62. Epping-Jordan, M.P., Markou, A., and Koob, G.F. (1998). The dopamine D-1 receptor antagonist
SCH 23390 injected into the dorsolateral bed nucleus of the stria terminalis decreased cocaine
reinforcement in the rat. Brain Res. *784*, 105–115. https://doi.org/10.1016/s0006-8993(97)01190-6.

63. Rezayof, A., Zarrindast, M.-R., Sahraei, H., and Haeri-Rohani, A.-H.-R. (2002). Involvement of
dopamine D2 receptors of the central amygdala on the acquisition and expression of morphine-
induced place preference in rat. Pharmacol. Biochem. Behav. *74*, 187–197.
https://doi.org/10.1016/s0091-3057(02)00989-9.

64. Thiel, K.J., Wenzel, J.M., Pentkowski, N.S., Hobbs, R.J., Alleweireldt, A.T., and Neisewander, J.L. (2010). Stimulation of dopamine D2/D3 but not D1 receptors in the central amygdala decreases cocaine-seeking behavior. Behav. Brain Res. *214*, 386–394. https://doi.org/10.1016/j.bbr.2010.06.021.

65. Bissière, S., Humeau, Y., and Luthi, A. (2003). Dopamine gates LTP induction in lateral amygdala by suppressing feedforward inhibition. Nature neuroscience *6*, 587–592.

66. Fadok, J.P., Dickerson, T.M.K., and Palmiter, R.D. (2009). Dopamine Is Necessary for Cue-Dependent Fear Conditioning. J Neurosci *29*, 11089–11097. https://doi.org/10.1523/JNEUROSCI.1616-09.2009.

67. Guarraci, F.A., Frohardt, R.J., and Kapp, B.S. (1999). Amygdaloid D1 dopamine receptor involvement in Pavlovian fear conditioning. Brain Res. *827*, 28–40. https://doi.org/10.1016/s0006-8993(99)01291-3.

68. de Oliveira, A.R., Reimer, A.E., de Macedo, C.E.A., de Carvalho, M.C., Silva, M.A. de S., and Brandão, M.L. (2011). Conditioned fear is modulated by D2 receptor pathway connecting the ventral tegmental area and basolateral amygdala. Neurobiol Learn Mem *95*, 37–45. https://doi.org/10.1016/j.nlm.2010.10.005.

69. Lutas, A., Kucukdereli, H., Alturkistani, O., Carty, C., Sugden, A.U., Fernando, K., Diaz, V., Flores-Maldonado, V., and Andermann, M.L. (2019). State-specific gating of salient cues by midbrain dopaminergic input to basal amygdala. Nat Neurosci *22*, 1820–1833. https://doi.org/10.1038/s41593-019-0506-0.

70. Lindzey, G., Winston, H., and Manosevitz, M. (1961). Social Dominance in Inbred Mouse Strains. Nature *191*, 474–476. https://doi.org/10.1038/191474a0.

71. Wang, F., Zhu, J., Zhu, H., Zhang, Q., Lin, Z., and Hu, H. (2011). Bidirectional Control of Social Hierarchy by Synaptic Efficacy in Medial Prefrontal Cortex. Science *334*, 693–697. https://doi.org/10.1126/science.1209951.

72. Zhou, T., Sandi, C., and Hu, H. (2018). Advances in understanding neural mechanisms of social dominance. Curr Opin Neurobiol *49*, 99–107. https://doi.org/10.1016/j.conb.2018.01.006.

73. Cacioppo, S., Bangee, M., Balogh, S., Cardenas-Iniguez, C., Qualter, P., and Cacioppo, J.T. (2016). Loneliness and implicit attention to social threat: A high-performance electrical neuroimaging study. Cogn Neurosci *7*, 138–159. https://doi.org/10.1080/17588928.2015.1070136.

74. Cacioppo, J.T., and Hawkley, L.C. (2009). Perceived Social Isolation and Cognition. Trends Cogn Sci *13*, 447–454. https://doi.org/10.1016/j.tics.2009.06.005.

75. Rodgers, R.J., and Dalvi, A. (1997). Anxiety, defence and the elevated plus-maze. Neuroscience & Biobehavioral Reviews *21*, 801–810. https://doi.org/10.1016/S0149-7634(96)00058-9.

76. Bailey, K.R., and Crawley, J.N. (2009). Anxiety-Related Behaviors in Mice. In Methods of Behavior Analysis in Neuroscience Frontiers in Neuroscience., J. J. Buccafusco, ed. (CRC Press/Taylor & Francis).

77. Lever, C., Burton, S., and O'Keefe, J. (2006). Rearing on hind legs, environmental novelty, and the
hippocampal formation. Rev Neurosci *17*, 111–133. https://doi.org/10.1515/revneuro.2006.17.1-
2.111.

78. Moy, S.S., Nadler, J.J., Perez, A., Barbaro, R.P., Johns, J.M., Magnuson, T.R., Piven, J., and
Crawley, J.N. (2004). Sociability and preference for social novelty in five inbred strains: an
approach to assess autistic-like behavior in mice. Genes Brain Behav. *3*, 287–302.
https://doi.org/10.1111/j.1601-1848.2004.00076.x.

79. Larrieu, T., Cherix, A., Duque, A., Rodrigues, J., Lei, H., Gruetter, R., and Sandi, C. (2017).
Hierarchical Status Predicts Behavioral Vulnerability and Nucleus Accumbens Metabolic Profile
Following Chronic Social Defeat Stress. Current Biology *27*, 2202-2210.e4.
https://doi.org/10.1016/j.cub.2017.06.027.

80. Füzesi, T., Daviu, N., Wamsteeker Cusulin, J.I., Bonin, R.P., and Bains, J.S. (2016). Hypothalamic
CRH neurons orchestrate complex behaviours after stress. Nat Commun *7*, 11937.
https://doi.org/10.1038/ncomms11937.

81. Lee, W., Fu, J., Bouwman, N., Farago, P., and Curley, J.P. (2019). Temporal microstructure of
dyadic social behavior during relationship formation in mice. PLoS One *14*.
https://doi.org/10.1371/journal.pone.0220596.

82. Tejada, J., Bosco, G.G., Morato, S., and Roque, A.C. (2010). Characterization of the rat exploratory
behavior in the elevated plus-maze with Markov chains. Journal of Neuroscience Methods *193*,
288–295. https://doi.org/10.1016/j.jneumeth.2010.09.008.

83. Dougalis, A.G., Matthews, G.A.C., Bishop, M.W., Brischoux, F., Kobayashi, K., and Ungless, M.A.
(2012). Functional properties of dopamine neurons and co-expression of vasoactive intestinal
polypeptide in the dorsal raphe nucleus and ventro-lateral periaqueductal grey. European Journal
of Neuroscience *36*, 3322–3332. https://doi.org/10.1111/j.1460-9568.2012.08255.x.

84. Huang, K.W., Ochandarena, N.E., Philson, A.C., Hyun, M., Birnbaum, J.E., Cicconet, M., and
Sabatini, B.L. (2019). Molecular and anatomical organization of the dorsal raphe nucleus. eLife *8*,
e46464. https://doi.org/10.7554/eLife.46464.

85. Motoike, T., Long, J.M., Tanaka, H., Sinton, C.M., Skach, A., Williams, S.C., Hammer, R.E.,
Sakurai, T., and Yanagisawa, M. (2016). Mesolimbic neuropeptide W coordinates stress responses
under novel environments. Proc Natl Acad Sci U S A *113*, 6023–6028.
https://doi.org/10.1073/pnas.1518658113.

86. McCullough, K.M., Daskalakis, N.P., Gafford, G., Morrison, F.G., and Ressler, K.J. (2018). Cell-
type-specific interrogation of CeA Drd2 neurons to identify targets for pharmacological modulation
of fear extinction. Transl Psychiatry *8*. https://doi.org/10.1038/s41398-018-0190-y.

87. McCullough, K.M., Morrison, F.G., Hartmann, J., Carlezon, W.A., and Ressler, K.J. (2018).
Quantified Coexpression Analysis of Central Amygdala Subpopulations. eNeuro *5*.
https://doi.org/10.1523/ENEURO.0010-18.2018.

88. Dougalis, A.G., Matthews, G.A.C., Liss, B., and Ungless, M.A. (2017). Ionic currents influencing
spontaneous firing and pacemaker frequency in dopamine neurons of the ventrolateral

periaqueductal gray and dorsal raphe nucleus (vlPAG/DRN): A voltage-clamp and computational
modelling study. J Comput Neurosci *42*, 275–305. https://doi.org/10.1007/s10827-017-0641-0.

89. Poulin, J.-F., Caronia, G., Hofer, C., Cui, Q., Helm, B., Ramakrishnan, C., Chan, C.S., Dombeck,
D., Deisseroth, K., and Awatramani, R. (2018). Mapping projections of molecularly defined
dopamine neuron subtypes using intersectional genetic approaches. Nat Neurosci *21*, 1260–1271.
https://doi.org/10.1038/s41593-018-0203-4.

90. Kash, T.L., Nobis, W.P., Matthews, R.T., and Winder, D.G. (2008). Dopamine Enhances Fast
Excitatory Synaptic Transmission in the Extended Amygdala by a CRF-R1-Dependent Process. J.
Neurosci. *28*, 13856–13865. https://doi.org/10.1523/JNEUROSCI.4715-08.2008.

91. Krawczyk, M., Georges, F., Sharma, R., Mason, X., Berthet, A., Bézard, E., and Dumont, É.C.
(2010). Double-Dissociation of the Catecholaminergic Modulation of Synaptic Transmission in the
Oval Bed Nucleus of the Stria Terminalis. Journal of Neurophysiology *105*, 145–153.
https://doi.org/10.1152/jn.00710.2010.

92. Kröner, S., Rosenkranz, J.A., Grace, A.A., and Barrionuevo, G. (2005). Dopamine Modulates
Excitability of Basolateral Amygdala Neurons In Vitro. Journal of Neurophysiology *93*, 1598–1610.
https://doi.org/10.1152/jn.00843.2004.

93. Marowsky, A., Yanagawa, Y., Obata, K., and Vogt, K.E. (2005). A Specialized Subclass of
Interneurons Mediates Dopaminergic Facilitation of Amygdala Function. Neuron *48*, 1025–1037.
https://doi.org/10.1016/j.neuron.2005.10.029.

94. Naylor, J.C., Li, Q., Kang-Park, M., Wilson, W.A., Kuhn, C., and Moore, S.D. (2010). Dopamine
attenuates evoked inhibitory synaptic currents in central amygdala neurons. Eur. J. Neurosci. *32*,
1836–1842. https://doi.org/10.1111/j.1460-9568.2010.07457.x.

95. Rosenkranz, J.A., and Grace, A.A. (1999). Modulation of basolateral amygdala neuronal firing and
afferent drive by dopamine receptor activation in vivo. J. Neurosci. *19*, 11027–11039.

96. Rosenkranz, J.A., and Grace, A.A. (2002). Cellular mechanisms of infralimbic and prelimbic
prefrontal cortical inhibition and dopaminergic modulation of basolateral amygdala neurons in vivo.
J. Neurosci. *22*, 324–337.

97. Silberman, Y., and Winder, D.G. (2013). Corticotropin releasing factor and catecholamines
enhance glutamatergic neurotransmission in the lateral subdivision of the central amygdala.
Neuropharmacology *70*, 316–323. https://doi.org/10.1016/j.neuropharm.2013.02.014.

98. Beckstead, M.J., Grandy, D.K., Wickman, K., and Williams, J.T. (2004). Vesicular dopamine
release elicits an inhibitory postsynaptic current in midbrain dopamine neurons. Neuron *42*, 939–
946. https://doi.org/10.1016/j.neuron.2004.05.019.

99. Marcott, P.F., Gong, S., Donthamsetti, P., Grinnell, S.G., Nelson, M.N., Newman, A.H., Birnbaumer,
L., Martemyanov, K.A., Javitch, J.A., and Ford, C.P. (2018). Regional heterogeneity of D2-receptor
signaling in the dorsal striatum and nucleus accumbens. Neuron *98*, 575-587.e4.
https://doi.org/10.1016/j.neuron.2018.03.038.

100.   Bettler, B., Kaupmann, K., Mosbacher, J., and Gassmann, M. (2004). Molecular structure and
814   physiological functions of GABA(B) receptors. Physiol Rev *84*, 835–867.
815   https://doi.org/10.1152/physrev.00036.2003.

816   101.   Destexhe, A., and Sejnowski, T.J. (1995). G protein activation kinetics and spillover of gamma-
817   aminobutyric acid may account for differences between inhibitory responses in the hippocampus
818   and thalamus. Proc Natl Acad Sci U S A *92*, 9515–9519.

819   102.   Mackay, J.P., Bompolaki, M., DeJoseph, M.R., Michaelson, S.D., Urban, J.H., and Colmers,
820   W.F. (2019). NPY2 Receptors Reduce Tonic Action Potential-Independent GABAB Currents in the
821   Basolateral Amygdala. J Neurosci *39*, 4909–4930. https://doi.org/10.1523/JNEUROSCI.2226-
822   18.2019.

823   103.   Cauli, B., Porter, J.T., Tsuzuki, K., Lambolez, B., Rossier, J., Quenet, B., and Audinat, E.
824   (2000). Classification of fusiform neocortical interneurons based on unsupervised clustering. Proc
825   Natl Acad Sci U S A *97*, 6144–6149. https://doi.org/10.1073/pnas.97.11.6144.

826   104.   Guthman, E.M., Garcia, J.D., Ma, M., Chu, P., Baca, S.M., Smith, K.R., Restrepo, D., and
827   Huntsman, M.M. (2020). Cell-type-specific control of basolateral amygdala neuronal circuits via
828   entorhinal cortex-driven feedforward inhibition. eLife. https://doi.org/10.7554/eLife.50601.

829   105.   Hou, W.-H., Kuo, N., Fang, G.-W., Huang, H.-S., Wu, K.-P., Zimmer, A., Cheng, J.-K., and Lien,
830   C.-C. (2016). Wiring Specificity and Synaptic Diversity in the Mouse Lateral Central Amygdala. J.
831   Neurosci. *36*, 4549–4563. https://doi.org/10.1523/JNEUROSCI.3309-15.2016.

832   106.   Chieng, B.C.H., Christie, M.J., and Osborne, P.B. (2006). Characterization of neurons in the rat
833   central nucleus of the amygdala: cellular physiology, morphology, and opioid sensitivity. The
834   Journal of Comparative Neurology *497*, 910–927. https://doi.org/10.1002/cne.21025.

835   107.   Dumont, E.C., Martina, M., Samson, R.D., Drolet, G., and Paré, D. (2002). Physiological
836   properties of central amygdala neurons: species differences. The European Journal of
837   Neuroscience *15*, 545–552. https://doi.org/10.1046/j.0953-816x.2001.01879.x.

838   108.   Lopez de Armentia, M., and Sah, P. (2004). Firing properties and connectivity of neurons in the
839   rat lateral central nucleus of the amygdala. Journal of Neurophysiology *92*, 1285–1294.
840   https://doi.org/10.1152/jn.00211.2004.

841   109.   Vander Weele, C.M., Siciliano, C.A., Matthews, G.A., Namburi, P., Izadmehr, E.M., Espinel,
842   I.C., Nieh, E.H., Schut, E.H.S., Padilla-Coreano, N., Burgos-Robles, A., et al. (2018). Dopamine
843   enhances signal-to-noise ratio in cortical-brainstem encoding of aversive stimuli. Nature *563*, 397.
844   https://doi.org/10.1038/s41586-018-0682-1.

845   110.   Kingsbury, L., Huang, S., Wang, J., Gu, K., Golshani, P., Wu, Y.E., and Hong, W. (2019).
846   Correlated Neural Activity and Encoding of Behavior across Brains of Socially Interacting Animals.
847   Cell *178*, 429-446.e16. https://doi.org/10.1016/j.cell.2019.05.022.

848   111.   Li, Y., Mathis, A., Grewe, B.F., Osterhout, J.A., Ahanonu, B., Schnitzer, M.J., Murthy, V.N., and
849   Dulac, C. (2017). Neuronal Representation of Social Information in the Medial Amygdala of Awake
850   Behaving Mice. Cell *171*, 1176-1190.e17. https://doi.org/10.1016/j.cell.2017.10.015.

112.   Yang, Y.C., McClintock, M.K., Kozloski, M., and Li, T. (2013). Social Isolation and Adult
Mortality: The Role of Chronic Inflammation and Sex Differences. J Health Soc Behav *54*, 183–203.
https://doi.org/10.1177/0022146513485244.

113.   Zilkha, N., Sofer, Y., Kashash, Y., and Kimchi, T. (2021). The social network: Neural control of
sex differences in reproductive behaviors, motivation, and response to social isolation. Curr Opin
Neurobiol *68*, 137–151. https://doi.org/10.1016/j.conb.2021.03.005.

114.   Choi, J.E., Choi, D.I., Lee, J., Kim, J., Kim, M.J., Hong, I., Jung, H., Sung, Y., Kim, J., Kim, T., et
al. (2022). Synaptic ensembles between raphe and D1R-containing accumbens shell neurons
underlie postisolation sociability in males. Science Advances *8*, eabo7527.
https://doi.org/10.1126/sciadv.abo7527.

115.   Fadok, J.P., Markovic, M., Tovote, P., and Lüthi, A. (2018). New perspectives on central
amygdala function. Curr Opin Neurobiol *49*, 141–147. https://doi.org/10.1016/j.conb.2018.02.009.

116.   Gungor, N.Z., and Paré, D. (2016). Functional Heterogeneity in the Bed Nucleus of the Stria
Terminalis. J Neurosci *36*, 8038–8049. https://doi.org/10.1523/JNEUROSCI.0856-16.2016.

117.   Tovote, P., Esposito, M.S., Botta, P., Chaudun, F., Fadok, J.P., Markovic, M., Wolff, S.B.E.,
Ramakrishnan, C., Fenno, L., Deisseroth, K., et al. (2016). Midbrain circuits for defensive
behaviour. Nature *534*, 206–212. https://doi.org/10.1038/nature17996.

118.   Padilla, S.L., Qiu, J., Soden, M.E., Sanz, E., Nestor, C.C., Barker, F.D., Quintana, A., Zweifel,
L.S., Rønnekleiv, O.K., Kelly, M.J., et al. (2016). AgRP Neural Circuits Mediate Adaptive Behaviors
in the Starved State. Nat Neurosci *19*, 734–741. https://doi.org/10.1038/nn.4274.

119.   Gulledge, A.T., and Jaffe, D.B. (2001). Multiple effects of dopamine on layer V pyramidal cell
excitability in rat prefrontal cortex. J Neurophysiol *86*, 586–595.
https://doi.org/10.1152/jn.2001.86.2.586.

120.   Maracle, A.C., Normandeau, C.P., Dumont, É.C., and Olmstead, M.C. (2019). Dopamine in the
oval bed nucleus of the stria terminalis contributes to compulsive responding for sucrose in rats.
Neuropsychopharmacology *44*, 381–389. https://doi.org/10.1038/s41386-018-0149-y.

121.   White, C.M., Ji, S., Cai, H., Maudsley, S., and Martin, B. (2010). Therapeutic potential of
vasoactive intestinal peptide and its receptors in neurological disorders. CNS & neurological
disorders drug targets *9*, 661.

122.   Nagata-Kuroiwa, R., Furutani, N., Hara, J., Hondo, M., Ishii, M., Abe, T., Mieda, M., Tsujino, N.,
Motoike, T., Yanagawa, Y., et al. (2011). Critical Role of Neuropeptides B/W Receptor 1 Signaling
in Social Behavior and Fear Memory. PLoS One *6*. https://doi.org/10.1371/journal.pone.0016972.

123.   Tanaka, H., Yoshida, T., Miyamoto, N., Motoike, T., Kurosu, H., Shibata, K., Yamanaka, A.,
Williams, S.C., Richardson, J.A., Tsujino, N., et al. (2003). Characterization of a family of
endogenous neuropeptide ligands for the G protein-coupled receptors GPR7 and GPR8. PNAS
100, 6251–6256. https://doi.org/10.1073/pnas.0837789100.

124.   Chen, Y., Essner, R.A., Kosar, S., Miller, O.H., Lin, Y.-C., Mesgarzadeh, S., and Knight, Z.A.
(2019). Sustained NPY signaling enables AgRP neurons to drive feeding. Elife *8*.
https://doi.org/10.7554/eLife.46348.

125.   Watanabe, N., Wada, M., Irukayama-Tomobe, Y., Ogata, Y., Tsujino, N., Suzuki, M., Furutani,
N., Sakurai, T., and Yamamoto, M. (2012). A single nucleotide polymorphism of the neuropeptide
B/W receptor-1 gene influences the evaluation of facial expressions. PLoS ONE *7*, e35390.
https://doi.org/10.1371/journal.pone.0035390.

126.   Kingsbury, M.A., and Wilson, L.C. (2016). The Role of VIP in Social Behavior: Neural Hotspots
for the Modulation of Affiliation, Aggression, and Parental Care. Integr Comp Biol *56*, 1238–1249.
https://doi.org/10.1093/icb/icw122.

127.   Reiner, A., Perkel, D.J., Bruce, L.L., Butler, A.B., Csillag, A., Kuenzel, W., Medina, L., Paxinos,
G., Shimizu, T., Striedter, G., et al. (2004). Revised nomenclature for avian telencephalon and
some related brainstem nuclei. J Comp Neurol *473*, 377–414. https://doi.org/10.1002/cne.20118.

128.   Wilson, L.C., Goodson, J.L., and Kingsbury, M.A. (2016). Seasonal Variation in Group Size Is
Related to Seasonal Variation in Neuropeptide Receptor Density. BBE *88*, 111–126.
https://doi.org/10.1159/000448372.

129.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-
image analysis. Nature Methods *9*, 676–682. https://doi.org/10.1038/nmeth.2019.

130.   McQuin, C., Goodman, A., Chernyshev, V., Kamentsky, L., Cimini, B.A., Karhohs, K.W., Doan,
M., Ding, L., Rafelski, S.M., Thirstrup, D., et al. (2018). CellProfiler 3.0: Next-generation image
processing for biology. PLOS Biology *16*, e2005970. https://doi.org/10.1371/journal.pbio.2005970.

131.   Paxinos, G., and Franklin, K.B.J. (2004). The mouse brain in stereotaxic coordinates 2nd
ed.,Compact ed. (Academic).

132.   Paxinos, G., and Franklin, K.B.J. (2019). Paxinos and Franklin's The mouse brain in stereotaxic
coordinates Fifth edition. (Academic Press, an imprint of Elsevier).

133.   Bogovic, J.A., Hanslovsky, P., Wong, A., and Saalfeld, S. (2016). Robust Registration of
Calcium Images by Learned Contrast Synthesis. 2016 IEEE 13th International Symposium on
Biomedical Imaging (ISBI), 1123–1126. https://doi.org/10.1109/ISBI.2016.7493463.

134.   Conte, W.L., Kamishina, H., and Reep, R.L. (2009). Multiple neuroanatomical tract-tracing using
fluorescent Alexa Fluor conjugates of cholera toxin subunit B in rats. Nat Protoc *4*, 1157–1166.
https://doi.org/10.1038/nprot.2009.93.

135.   Hunter, J.D. (2007). Matplotlib: A 2D Graphics Environment. Computing in Science Engineering
9, 90–95. https://doi.org/10.1109/MCSE.2007.55.

136.   Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., Dubourg, V., et al. (2011). Scikit-learn: Machine Learning in Python. J.
Mach. Learn. Res. *12*, 2825–2830.

137.   Gottman, J.M., and Roy, A.K. (1990). Sequential Analysis: A Guide for Behavioral Researchers
(Cambridge University Press).

138.   Novák, P., and Zahradník, I. (2006). Q-Method for High-Resolution, Whole-Cell Patch-Clamp
Impedance Measurements Using Square Wave Stimulation. Ann Biomed Eng *34*, 1201–1212.
https://doi.org/10.1007/s10439-006-9140-6.

139.   Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski,
E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental algorithms for
scientific computing in Python. Nature Methods *17*, 261–272. https://doi.org/10.1038/s41592-019-
0686-2.

140.   Ward, J.H. (1963). Hierarchical Grouping to Optimize an Objective Function. null *58*, 236–244.
https://doi.org/10.1080/01621459.1963.10500845.

141.   Michael Waskom, Olga Botvinnik, Maoz Gelbart, Joel Ostblom, Paul Hobson, Saulius
Lukauskas, David C Gemperline, Tom Augspurger, Yaroslav Halchenko, Jordi Warmenhoven, et al.
(2020). mwaskom/seaborn: v0.11.0 (Sepetmber 2020) (Zenodo)
https://doi.org/10.5281/zenodo.4019146.

142.   Stamatakis, A.M., Schachter, M.J., Gulati, S., Zitelli, K.T., Malanowski, S., Tajik, A., Fritz, C.,
Trulson, M., and Otte, S.L. (2018). Simultaneous Optogenetics and Cellular Resolution Calcium
Imaging During Active Behavior Using a Miniaturized Microscope. Front. Neurosci. *12*.
https://doi.org/10.3389/fnins.2018.00496.

143.   Pereira, T.D., Tabris, N., Matsliah, A., Turner, D.M., Li, J., Ravindranath, S., Papadoyannis,
E.S., Normand, E., Deutsch, D.S., Wang, Z.Y., et al. (2022). SLEAP: A deep learning system for
multi-animal pose tracking. Nat Methods *19*, 486–495. https://doi.org/10.1038/s41592-022-01426-1.

144.   Pnevmatikakis, E.A., and Giovannucci, A. (2017). NoRMCorre: An online algorithm for
piecewise rigid motion correction of calcium imaging data. J Neurosci Methods *291*, 83–94.
https://doi.org/10.1016/j.jneumeth.2017.07.031.

145.   Zhou, P., Resendez, S.L., Rodriguez-Romaguera, J., Jimenez, J.C., Neufeld, S.Q.,
Giovannucci, A., Friedrich, J., Pnevmatikakis, E.A., Stuber, G.D., Hen, R., et al. (2018). Efficient
and accurate extraction of in vivo calcium signals from microendoscopic video data. eLife *7*,
e28728. https://doi.org/10.7554/eLife.28728.

146.   Sheintuch, L., Rubin, A., Brande-Eilat, N., Geva, N., Sadeh, N., Pinchasof, O., and Ziv, Y.
(2017). Tracking the Same Neurons across Multiple Days in Ca2+ Imaging Data. Cell Rep *21*,
1102–1115. https://doi.org/10.1016/j.celrep.2017.10.013.

Methods

Animals and housing

All procedures involving animals were conducted in accordance with NIH guidelines and
approved by the MIT Committee on Animal Care or the Salk Institute Institutional Animal Care and Use
Committee. DAT::IRES-Cre (B6.SJL-Slc6a3$^{tm1.1(cre)Bkmn}$/J)[32] were purchased from the Jackson
Laboratory (stock no. 006660; the Jackson Laboratory, ME, USA) and bred in-house to generate
heterozygous male offspring for experiments. Wild-type C57BL/6J male mice were purchased from
Charles River Laboratories (MA, USA). Mice were housed on a 12h:12h reverse light dark cycle (MIT:
lights off 9am-9pm; Salk Institute: lights off 9.30am-9.30pm) with food and water available *ad libitum*.
Mice were housed in groups of 2-4 with same-sex siblings. For photoinhibition and CeA calcium
imaging experiments, mice were additionally tested following 24 hours of social isolation. Only mice
with acceptable histological placements were included in final datasets.

Surgery and viral constructs

Mice (>7 weeks of age) were anaesthetized with isoflurane (inhalation: 4% for induction, ~2%
for maintenance, oxygen flow rate 1 L/min) before being placed in a digital small animal stereotax
(David Kopf Instruments, CA, USA). Surgeries were performed under aseptic conditions with body
temperature maintained by a heating pad throughout. Injections of recombinant adeno-associated viral
(AAV) vectors, herpes simplex virus (HSV), or cholera toxin subunit-B (CTB) were performed using a
beveled 33 gauge microinjection needle with a 10 µL microsyringe (Nanofil; WPI, FL, USA). Virus or
CTB was delivered at a rate of 0.1 µL/min using a microsyringe pump (UMP3; WPI, FL, USA)
connected to a Micro4 controller (WPI, FL, USA). Following injection, the needle was maintained in
place for ~2 min, then raised up by 0.05 mm and held for ~10 min (to permit diffusion from the injection
site) before being slowly withdrawn. Skull measurements were made relative to Bregma for all
injections and implants. Implants were secured to the skull by a layer of adhesive cement (C&B
Metabond; Parkell Inc., NY, USA) followed by a layer of black cranioplastic cement (Ortho-Jet; Lang, IL,
USA). Mice were given pre-emptive analgesia (1 mg/kg buprenorphine slow-release; sub-cutaneous;
delivered concurrent with warmed Ringer's solution to prevent dehydration), supplemented with
meloxicam (1.5 mg/kg; sub-cutaneous) where necessary, and were monitored on a heating pad until
recovery from anesthesia.

AAV$_5$-EF1α-DIO-ChR2-eYFP, AAV$_5$-EF1α-DIO-eYFP, AAV$_5$-EF1 α -fDIO-eYFP , and AAV$_8$-Syn-
ChrimsonR-tdTomato were packaged by the University of North Carolina Vector Core (NC, USA) and
received the AAV$_5$-EF1a-fDIO-eYFP construct from Karl Deisseroth and Charu Ramakrishnan. HSV-
LS1L-mCherry-IRES-flpo was packaged by Dr. Rachael Neve at the Viral Gene Transfer Core Facility
at MIT (now located at Massachusetts General Hospital). AAV$_5$-EF1α-DIO-eNpHR3.0-eYFP and AAV$_1$-
syn-jGCaMP7f was packaged by Addgene (MA, USA), and AAV$_1$-CAG-TdTomato was packaged by the
UPenn vector core (PA, USA).

Immunohistochemistry and confocal microscopy

Mice were deeply anaesthetized with sodium pentobarbital (200 mg/kg, intraperitoneal; IP) or
euthasol (150 mg/kg; IP) followed by transcardial perfusion with 10 mL ice-cold Ringer's solution and
15 mL ice-cold 4% paraformaldehyde (PFA). The brain was carefully dissected from the cranial cavity
and immersed in 4% PFA for ~6-18 h before transfer to 30% sucrose solution in phosphate-buffered

saline (PBS) at 4°C. After at least 48 hr, brains were sectioned at 40 μm on a freezing sliding microtome (HM430; Thermo Fisher Scientific, MA, USA) and sections stored at 4°C in 1X PBS. For immunohistochemistry, sections were blocked in PBS containing 0.3% Triton X-100 (PBS-T; Sigma-Aldrich, MO, USA) with 3% normal donkey serum (NDS; Jackson Immunoresearch, PA, USA) for 30-60 min at room temperature. This was followed by incubation in primary antibody solution chicken anti-TH (1:1000; AB9702; EMD Millipore, MA, USA) in 0.3% PBS-T with 3% NDS) overnight at 4°C. Sections were then washed in 1X PBS four times (10 min each) before incubation in secondary antibody solution containing donkey anti-chicken 488 or 647 (1:1000; Jackson Immunoresearch, PA, USA) and a DNA-specific fluorescent probe (DAPI; 1:50000; Invitrogen, Thermo Fisher Scientific, MA, USA) in 0.2% PBS-T with 3% NDS for 1.5-2 hr at room temperature. Sections were again washed four times in 1X PBS (10 min each) before being mounted on glass slides and coverslipped using warmed PVA-DABCO (Sigma-Aldrich, MO, USA).

Images were captured on a laser scanning confocal microscope (Olympus FV1000, Olympus, PA, USA) using Fluoview software version 4.0 (Olympus, PA, USA). Images were collected through a 10X/0.40 NA objective for injection site and optic fiber placement verification, a 20X/0.75 objective for terminal fluoresence quantification, and an oil-immersion 40X/1.30 NA objective for neurobiotin-filled neurons and RNAscope analysis (see individual Methods sections for more detail). FIJI[129], CellProfiler 3.1 (Broad Institute, MA, USA)[130], and Adobe Photoshop CC (Adobe Systems Incorporated, CA, USA) were used for subsequent image processing and analysis.

Downstream fluorescence quantification

In DAT::Cre mice, AAV_5-EF1α-DIO-ChR2-eYFP (300 nL) was injected into the DRN (ML:1.20, AP:-4.10, DV:-2.90; needle at a 20° angle from the midline, bevel facing medial) or VTA (ML:0.85, AP:-2.70, DV:-4.50), and after 8 weeks mice underwent perfusion-fixation. Brains were subsequently sectioned at 40 μm, processed with immunohistochemistry for TH and DAPI, and serial z-stack images (3 μm optical thickness) collected at 20X on a confocal microscope. (See *'Immunohistochemistry and confocal microscopy'* section above for details). A maximum projection was generated in FIJI and background subtraction based on the 'rolling ball' algorithm (radius = 50 pixels) was applied to correct for uneven illumination. The appropriate brain atlas slice[131,132] was overlaid onto the fluorescent image using the BigWarp plugin (https://imagej.net/BigWarp)[133] in FIJI, by designating major anatomical landmarks based on DAPI staining and TH expression. Regions of interest (ROIs) were then annotated from the overlaid atlas, and mean fluorescence within each ROI quantified using FIJI. The PFC was examined from AP: 2.22 to 1.34, the striatum from AP 1.70 to 0.74, the BNST from AP 0.37 to -0.11, the CeA from AP -0.82 to -1.94, and the amygdala from AP -0.82 to -2.92. Average images in Figure 1D, 5A, 5E, and 5I were created by aligning individual images (from the middle AP region of the BNST, CeA, or BLP), using the line ROI registration plugin (https://imagej.net/Align_Image_by_line_ROI) in FIJI. An average projection was then performed across all images and the 'royal' LUT applied to visualize relative fluorescence intensity.

Retrograde tracing and intersectional viral expression

C57BL/6 mice were injected with 150-250 nL CTB conjugated to Alexa Fluor-555 (CTB-555) or Alexa Fluor-647 (CTB-647; Molecular Probes, OR, USA[134]) in two of three locations: the BNST (ML:1.10, AP:0.50, DV: -4.30; needle bevel facing back), CeA (ML:2.85, AP: -1.20; DV:-4.75; needle

bevel facing back), or BLP (ML:3.35, AP:-2.20, DV:-5.25; needle bevel facing back). To assess
retrograde CTB co-expression following injection of both fluorophore-conjugates of CTB into the same
region (Figure 1—figure supplement 2A-C), injections were either performed sequentially, or CTB-555
and CTB-647 were mixed prior to a single injection. After 7 days to allow for retrograde transport, mice
were deeply anaesthetized with sodium pentobarbital (200 mg/kg) and perfused-fixed for subsequent
histology. Brain sections containing injection sites and the DRN were prepared at 40 µm and processed
with immunohistochemistry for TH and DAPI. (See *'Immunohistochemistry and confocal microscopy'*
section above for details). CTB injection sites were verified with images acquired on a confocal
microscope through a 10X objective (serial z-stack with 5 µm optical thickness) and images of the DRN
were acquired through a 40X objective (serial z-stack with 3 µm optical thickness). DRN cells co-
expressing CTB and TH were counted manually using the ROI 'point' tool in Fluoview software version
4.0 (Olympus, PA, USA). Counted files were imported into FIJI, and images overlaid onto the
appropriate brain atlas image of the DRN using the BigWarp plugin (https://imagej.net/BigWarp)[133]. The
x-y coordinates of counted/marked CTB+/TH+ cells were extracted using the 'Measure' function in FIJI.
These coordinates were then used to generate heatmaps of cell location (Figure 1—figure supplement
2D-E) by creating a 2D histogram using the Matplotlib package[135] in Python .

Intersectional labelling of the dopaminergic projection from the DRN to the CeA was achieved
by injecting HSV-LS1L-mCherry-IRES-flpo (300 nL) into the CeA (ML:2.85, AP:-1.45, DV:-4.55; needle
bevel facing medial) and AAV$_5$-fDIO-eYFP (300 nL) into the DRN (ML:1.20, AP:-4.10, DV:-2.90; needle
at a 20° angle from the midline, bevel facing medial) of a DAT::Cre mouse. After 8 weeks, mice were
perfused-fixed with 4% PFA, and the brain sectioned on a freezing microtome at 40 µm before
immunohistochemical processing with TH and DAPI. Images of eYFP-expressing cells in the DRN and
terminals in the CeA and BNST were captured on a confocal microscope through a 20X objective with a
serial z-section thickness of 3 µm.

1065   Behavioral assays and optogenetic manipulations

DAT::Cre mice were injected with 300 nL AAV$_5$- EF1α-DIO-ChR2-eYFP or AAV$_5$-EF1α-DIO-
eYFP in the DRN (ML:1.20; AP:-4.10; DV:-2.90; needle at a 20° angle from the right side, bevel facing
medial) and optic fibers (300 µm core, NA=0.37; Thorlabs, NJ, USA), held within a stainless steel
ferrule (Precision Fiber Products, CA, USA), were implanted unilaterally or bilaterally over the BNST
(unilateral: ML:1.10, AP:0.40, DV:-3.50; bilateral: ML:1.65, AP:0.40, DV:-3.35; 10° angle from midline),
CeA (ML:2.85, AP:-1.35, DV:-4.00), or BLP (ML:3.30, AP:-2.20, DV:-4.30). Behavioral experiments
commenced 7-8 weeks following surgery. Mice were handled and habituated to patch cable connection
once per day for at least 3 days before beginning optical manipulations. Behavioral testing was
performed in dimly-lit soundproofed room during the mice's active dark phase (~10am-5pm). On each
testing day, mice were given at least 1 hr to acclimate to the testing room before experiments began.
For optical manipulations, optic fiber implants were connected to a patch cable via a ceramic sleeve
(Precision Fiber Products, CA, USA), which itself was connected to a commutator (rotary joint; Doric,
Québec, Canada) using an FC/PC adapter, to permit uninhibited movement. The commutator, in turn,
was connected via a second patch cable (with FC/PC connectors) to a 473 nm diode-pumped solid
state (DPSS) laser (OEM Laser Systems, UT, USA). To control the output of the laser, a Master-8
pulse stimulator (AMPI, Israel) was used, and the light power set to 10 mW.

<u>Tube test:</u> Cages of mice (same-sex groups of 2-4) were assayed for social dominance using
the tube test[70,71]. Mice were individually trained to pass through a clear Plexiglas tube (30 cm length,

3.2 cm inner diameter) over 4 days. Each training trial involved releasing the mouse into the tube from
one end, and ensuring it traveled through and out the other side. Mice which attempted to reverse, or
were reluctant to exit at the other end of the tube, were gently encouraged forwards by light pressure
from a plastic stick pressing on their hind region. Between trials mice freely explored the open arena
outside tube (76 x 60 cm) for ~30-60 s. Mice performed 8 training trials (4 from each end) on days 1
and 2, and 3 trials (alternating ends) on days 3 and 4. On days 5-8 mice competed against cagemates
in a round-robin design. For each contest, mice were released simultaneously into opposite ends of the
tube so that they met face-to-face in the center of the tube. The mouse which retreated from the
confrontation was designated as the 'loser' and his opponent designated the 'winner'. Across testing
days, the side from which animals were released and the order in which they were tested against
cagemates was counterbalanced. An animal's 'relative dominance' score reflected their proportion of
'wins' across all contests from 3-4 days of testing.

<u>Open field test (OFT):</u> The open field was composed of a square arena (51 x 51 cm) made of
transparent Plexiglas with 25 cm high walls. Mice freely explored the arena for 15 min, and blue light (8
pulses with 5 ms pulse-width, at 30 Hz, every 5 s) was delivered during the middle 5 min epoch of the
session. Animals were recorded using a video camera positioned above the arena, and Ethovision XT
software used to track mouse location (Noldus, Netherlands). To assess anxiety-related behavior, for
analysis, the chamber was divided into a 'center' square region and a 'periphery', with equal area.

<u>Three chamber sociability assay</u>: The apparatus consisted of a 57.5l x 22.5w x 16.5h cm
chamber, with transparent Plexiglas walls and opaque grey plastic floors. The chamber was divided into
unmarked left and right compartments (each 23 x 22.5 cm) and a smaller center compartment (11.5 x
22.5 cm). An upturned wire mesh cup was placed in the left and right compartments. Each mouse first
underwent a habituation session (10 min) where they freely explored the chamber. They were then
briefly (~1 min) confined to the center compartment by the insertion of clear Plexiglas walls, while a
novel object was placed under one of the two upturned cups, and a juvenile C57BL/6 mouse (3.5-5
weeks of age) was placed under the other upturned cup. The mice were then allowed to freely explore
the chamber for a further 10 min. The task was repeated on the second day, with the chamber rotated
by 90˚ relative to external spatial cues, and with a different novel object and novel juvenile mouse. The
10 min test epoch was paired with blue light delivery (8 pulses with 5 ms pulse-width, at 30 Hz, every 5
s) on one of the two days, counterbalanced across animals. Mice were excluded if they showed a
strong preference (>70% time spent) for one side of the chamber in the habituation phase, or if they
spent more than 1 min on top of the upturned cups during any session. For photoinhibition experiments,
the protocol was exactly the same as the ChR2 experiment, except the 10 min test epoch was paired
with constant yellow light (589nm) delivery for one day in the 'group-housed' and also during the
additional 'isolated' condition. Animals were recorded using a video camera positioned above the
chamber and movement tracked using Ethovision XT (Noldus, Netherlands). The social:object ratio
reflected the time spent in the 'social' side of the chamber (containing a novel juvenile mouse) divided
by the time spent in the 'object' side of the chamber (containing a novel object).

<u>Juvenile intruder assay:</u> Mice were tested individually in their home cage. They freely explored
alone for 5 min after which a novel juvenile mouse was placed in the cage for a further 3 min. The task
was repeated on the second day with a different novel juvenile mouse. One of the two sessions was
paired with blue light delivery (8 pulses with 5 ms pulse-width, at 30 Hz, every 5 s) which commenced
after 2 min and continued until the end of the task (6 min total). The behavior of the mouse during the 3

min with the juvenile was scored manually using ODLog software (Macropod Software, Australia).
Video files were scored twice (by two different observers, blinded to the experimental conditions) and
the average of their counts was used for analysis. (See also *'First order Markov analysis'* section).

<u>Elevated plus maze (EPM)</u>: The EPM was made of grey plastic and consisted of two closed
arms (30l x 5w x 30h cm) and two open arms (30l x 5w cm), radiating at 90° from a central platform (5 x
5 cm) and raised from the ground by 75 cm. Mice freely explored for 15 min, with blue light (8 pulses
with 5 ms pulse-width, at 30 Hz, every 5 s) delivered during the middle 5 min epoch of the session. A
video camera position above the EPM was used to record animals, and movement was tracked using
Ethovision XT (Noldus, Netherlands).

<u>Real-time place preference (RTPP)</u>: Mice were placed in a 52l x 52w x 26.5h cm transparent
Plexiglas chamber, with clear panels separating left and right sides to leave a 11.5 cm gap for mice to
pass through. Mice freely explored for 30 min, during which entry into one side of the chamber resulted
in delivery of blue light (15 pulses with 5 ms pulse-width, at 30 Hz, every 5 s), which continued until
mice exited the zone. Entry into the opposite side did not result in blue light delivery. The side paired
with blue light delivery was counterbalanced across animals. A video camera positioned above the
arena recorded animals, and mouse movement was tracked using Ethovision XT (Noldus,
Netherlands).

<u>Intra-cranial self-stimulation (ICSS)</u>: Mice were food deprived for 16-20 hr prior to each day of
ICSS, in order to encourage behavioral responding. Testing was conducted in an operant chamber
(Med Associates, VT, USA) within a custom sound-attenuating outer box. The operant chamber
contained two illuminated nose-poke ports, each with an infrared beam, and a cue light positioned
above each port. White noise was delivered continuously throughout the session, and successful nose-
pokes (signaled by a beam break) resulted in an auditory tone (1 s duration, 1 or 1.5 kHz) and
illumination of the respective cue light. A nose-poke at the 'active' port also triggered delivery of blue
light (90 pulses with 5 ms pulse-width, at 30 Hz) while a nose-poke at the 'inactive' port did not trigger
light delivery. The physical location of the active and inactive nose-poke ports, and the auditory tone
frequency associated with each port, was counterbalanced across animals. On day 1 (training), mice
completed a 2 hr session in the operant chamber in which both nose-poke ports were baited with a
small amount of palatable food, in order to encourage investigation. On day 2 (testing), mice completed
an identical 2 hr session, except the nose-poke ports were not baited. Nose-poke activity was recorded
with MedPC software (Med Associates, VT, USA) and subject-averaged cumulative distribution plots
were generated using MATLAB (Mathworks, MA, USA). Only data from day 2 was used for analysis.

<u>Analysis of baseline behavior</u>: The baseline behavior of all mice (i.e without stimulation) was
evaluated to uncover any relationships between specific types of behavior assessed in different tasks.
These analyses used relative dominance from the tube test, the first 5 min of the OFT and EPM, and
the OFF trial from the three-chamber sociability and juvenile intruder assays. Correlation matrices were
generated in GraphPad Prism 8 (GraphPad Software, CA, USA) to show the Pearson's correlation
coefficient for each pair of variables.

Dimensionality reduction was performed on baseline behavior data using principal component
analysis (PCA) with the scikit-learn module[136] in Python. The eight input measures from behavioral
assays were (1) percent time moving in the OFT, (2) time in the center of the OFT, (3) time in the open
arms of the EPM, (4) social:object ratio in the three chamber assay, and (5) time spent face sniffing, (6)

anogenital sniffing, (7) rearing, and (8) grooming in the juvenile intruder assay. The data was first
normalized to generate a covariance matrix and then the first 5 PCs were extracted. Relative
dominance was concatenated with the resulting PC values for each mouse to color-code individual
points in the PC1 vs PC2 plot.

First order Markov analysis: Behavioral videos from the juvenile intruder assay were manually
annotated so that each second of the 180 s session was assigned a code(s) from 15 behavioral
categories:

- Social behaviors: face sniff (reciprocated), face sniff (non-reciprocated), flank sniff,
anogenital sniff (reciprocated), anogenital sniff (non-reciprocated), close follow, approach,
dominant climb, attack.
- Nonsocial behaviors: groom, dig, rear, climb, still, ambulate.

We designed a 2-state Markov model, in which behaviors were assigned to either the 'social' or
'nonsocial' categories. For each animal, we created a transition probability matrix from each sequence
by counting the number of transitions that occurred and dividing by the total number of occurrences of
that behavior. To compute the overall transition probability matrix for the eYFP and ChR2 groups, we
took the mean of the transition probability across all individuals in that group. Difference scores
between the stimulation OFF and ON sessions were calculated by taking the difference across pairs of
transition probability matrices corresponding to each individual, then calculating the mean across eYFP
or ChR2-expressing mice.

To verify that a first order Markov model was an appropriate fit for our data we computed the log
likelihood chi squared statistic[137]:

$$G = 2 \sum_j \sum_i O_{ij} \ln \frac{O_{ij}}{E_{ij}},$$

where $O_{ij} \geq 0$ is the observed number of transitions from state i to j, $E_{ij} \geq 0$ is the expected number of
transitions from state i to state j assuming a zeroth order Markov (i.e., no time dependence). We found
that G was statistically significant for all subjects in both the 15 state and 2 state models, thus rejecting
the null hypothesis of randomly transitioning between states.

We also tested whether a non-stationary model was a better fit for the data than a stationary model. To
do this, we divided each subject's behavioral sequence into two segments of equal duration and
computed transition probability matrices for each segment. We then computed a variation on the
likelihood ratio chi square statistic[137]:

$$LRX = 2 \sum_s \sum_j \sum_i f_{ijs} \ln \frac{\bar{p}_{ijs}}{p_{ij}},$$

1198

where *s* represents the segment, p_{ij} is the probability of transition from state *i* to *j* taken over the entire
sequence, $\bar{p}_{ijs}$ is the probability of transition from *i* to *j* for each segment, and f_{ijs} is the number of
transitions from state *i* to *j* for each segment. Since not all subjects had a significant difference, we
determined that a stationary model was the most appropriate model to fit all our data.

Ex vivo electrophysiology

DAT::Cre mice received an injection of 300 nL AAV_5-DIO-ChR2-eYFP or AAV_9-FLEX-
ChrimsonR-TdTomato in the DRN (ML:1.20, AP:-4.10, DV:-2.90; needle at a 20° angle from the
midline, bevel facing medial), and after at least 8 weeks for transgene expression, mice were deeply
anaesthetized with sodium pentobarbital (200 mg/kg) or euthasol (150 mg/kg; IP). They were then
transcardially perfused with ice-cold (~4°C) modified artificial cerebrospinal fluid (ACSF; composition in
mM: NaCl 87, KCl 2.5, NaH2PO4*H20 1.3, MgCl2*6H2O 7, NaHCO3 25, sucrose 75, ascorbate 5,
CaCl2*2H2O 0.5, in ddH20; osmolarity 320-330 mOsm, pH 7.30-7.40), saturated with carbogen gas
(95% oxygen, 5% carbon dioxide) before the brain was rapidly and carefully extracted from the cranial
cavity. Thick coronal (300 µm) slices containing the BNST, CeA, BLP, and DRN were prepared on a
vibrating blade vibratome (VT1200; Leica Biosystems, Germany), in ice-cold modified ACSF saturated
with carbogen gas. Brain slices were hemisected with a scalpel blade before transfer to a holding
chamber containing ACSF (composition in mM: NaCl 126, KCl 2.5, NaH2PO4*H20 1.25, MgCl2*6H2O
1, NaHCO3 26, glucose 10, CaCl2*H2O 2.4; osmolarity 298-302 mOsm, pH 7.30-7.40) saturated with
carbogen, in a warm water bath (~30°C).

Electrophysiological recordings were commenced after the slices had rested for at least 45 min.
During recording, the brain slice was maintained in a bath with continuously perfused ACSF, saturated
with carbogen, at 31±1°C using a peristaltic pump (Minipuls3; Gilson, WI, USA). Slices were visualized
through an upright microscope (Scientifica, UK) equipped with infrared-differential interference contrast
(IR-DIC) optics and a Q-imaging Retiga Exi camera (Q Imaging, Canada). In the BNST, CeA, and BLP,
recordings were performed in the region containing fluorescent DRN^{DAT} terminals (expressing ChR2-
eYFP or Chrimson-TdTomato) with neurons visualized through a 40X/0.80 NA water immersion
objective. Terminal expression was confirmed by brief illumination from a 470 nm LED light source (pE-
100; CoolLED, NY, USA) for ChR2-eYFP, or a metal halide lamp (Lumen 200; Prior Scientific Inc., UK),
for ChrimsonR-TdTomato, combined with the appropriate filter set. Borosilicate glass capillaries were
shaped on a P-97 puller (Sutter Instrument, CA, USA) to produce pipettes for recording that had
resistance values of 3.5-5 MOhm when filled with internal solution (composition in mM: potassium
gluconate 125, NaCl 10, HEPES 20, MgATP 3, and 0.1% neurobiotin, in ddH20 (osmolarity 287 mOsm;
pH 7.3). Whole-cell patch-clamp recordings were made using pClamp 10.4 software (Molecular
Devices, CA, USA), with analog signals amplified using a Multiclamp 700B amplifier, filtered at 3 kHz,
and digitized at 10 kHz using a Digidata 1550 (Molecular Devices, CA, USA). A 5 mV, 250 ms
hyperpolarizing step was used to monitor cell health throughout the experiment, and recordings were
terminated if significant changes (>20%) occurred to series resistance (R_s), input resistance (R_{in}), or
holding current.

Passive cell properties (capacitance, membrane resistance) were estimated from the current
response to hyperpolarizing 5 mV, 250 ms steps, delivered in voltage-clamp from a holding potential of
-70 mV, using custom MATLAB code written by Praneeth Namburi, based on MATLAB implementation
of the Q-Method[138]. To examine the membrane potential response to current injection, cells were
recorded in current-clamp mode, and a series of 1 s steps were delivered, in 20 pA increments, from -
120 pA to 260 pA. The voltage sag amplitude (attributable to the hyperpolarization-activated cation
current; I_h) was measured as the difference between the peak instantaneous and steady-state
membrane potential elicited during a -120 pA step (see Figure 6L). The ramp ratio was calculated by
dividing the average membrane potential between 900-1000 ms by the membrane potential between
100-200 ms following step onset, using the largest current step that elicited a subthreshold response

(i.e. did not evoke action potentials). The firing delay was taken as the time between current step onset
and the first elicited action potential, on delivery of the first current step that was elicited a
suprathreshold response (i.e. rheobase current). The max instantaneous firing frequency (max freq.$_{inst}$)
was taken as the maximum firing frequency attained during the first 100 ms of the depolarizing current
steps.

1252   To photostimulate ChR2-expressing DRN^{DAT} terminals in the BNST, CeA, and BLP, 470 nm
light was delivered through the 40X/0.8 NA objective from an LED light source (pE-100; CoolLED, NY,
USA). Neurons were recorded at their resting membrane potential in current-clamp mode, and 470 nm
light (8 pulses at 30 Hz, 5ms pulse-width) was delivered every 30 s. In a minority of cells which showed
spontaneous activity at the resting potential, negative current was injected to hold the cell at a
subthreshold potential (typically ~-60 mV). The peak amplitude of the optically-evoked excitatory post-
synaptic potential (EPSP) or trough amplitude of the inhibitory post-synaptic potential (IPSP) was
measured from the average trace using Clampfit 10.7 (Molecular Devices, CA, USA), using the 5 s prior
to stimulation as baseline. Tau for the decay phase of the IPSP was estimated by fitting the IPSP with a
single exponential, from the IPSP trough until return to baseline. Total voltage area was calculated from
0-5.5 s following the onset of the first light pulse. In cells where optical stimulation evoked only an
EPSP the response was classed as an 'excitation', only an IPSP was classed as an 'inhibition', and a
combined optically-evoked EPSP and IPSP was classed as 'mixed'. To assess the effect of
photostimulation on firing activity, constant positive current was injected to elicit consistent spontaneous
action potentials, and 470 nm light (8 pulses at 30 Hz, 5 ms pulse-width) was delivered every 30 s. The
interevent interval (IEI) between action potentials was calculated for 5 s before and 5 s after the first
pulse of blue light using Clampfit 10.7 (Molecular Devices, CA, USA). A decrease in IEI (indicating an
increase in firing rate) was classed as an 'excitation' and an increase in IEI (indicating a decrease in
firing rate) was classed as an 'inhibition'.

1271   Following recording, images showing the location of the recording pipette within the slice were
1272   captured through a 4X/0.10 NA objective. Images were subsequently overlaid onto the appropriate
1273   brain atlas image[131,132], recorded cell locations were annotated, and then converted into x-y coordinates
1274   in FIJI. Python was used to generate a scatter plot of cell location, with points color-coded by the
1275   overall membrane potential response to photostimulation (Figure 6—figure supplement 1A-C).

1276   Unsupervised agglomerative hierarchical clustering was used to classify cells according to their
baseline electrophysiological properties. This approach organizes objects (in this case cells) into
clusters, based on their similarity. The electrophysiological properties used as input features for
clustering CeA cells were ramp ratio, max firing frequency, firing delay, and voltage sag, which are
characteristics that have been previously shown to distinguish between subtypes of CeA neuron[105]. For
clustering BLP cells, we replaced ramp ratio with capacitance, as this measure is often used to
distinguish between pyramidal neurons and GABAergic interneurons, which are the two main cell types
in this region. Data for each cell property was max-min normalized to produce a 4 x n matrix of input
features (where n = total number of cells). Clustering was performed using the 'linkage' function of
SciPy[139] in Python, using Ward's linkage method[140] and Euclidean distance. Briefly, this approach
begins with each cell assigned to a single cluster. Cells that are in closest proximity (i.e. have highest
similarity) are then linked to form a new cluster. Then the next closest clusters are linked, and so on.
This process is repeated until all cells are included in a single cluster. The output of this analysis is
plotted as a hierarchical tree (dendrogram), in which each cell is a 'leaf' and the Euclidean distance on

the y-axis indicates the linkage between cells (larger distance indicates greater dissimilarity). To
annotate the photostimulation response of cells on the dendrogram (Figure 6N, 6P, and Figure 6—
figure supplement 1M), the response was designated as 'excitation' if action potential IEI decreased
with optical stimulation and 'inhibition' if action potential IEI increased on stimulation. If firing data was
not available, cells were designated as showing an 'excitation' if only an EPSP was evoked on optical
stimulation, and 'inhibition' if only an IPSP was evoked. In cells where a mixed EPSP/IPSP was elicited,
the response was designated as an 'excitation' if the overall voltage area (0-5.5 s following light onset)
was positive, and an 'inhibition' if the overall voltage area was negative.

At the end of recording, brain slices were fixed in 4% PFA overnight and then washed in 1X
PBS (4 x 10 min each). Slices were blocked in 0.3% PBS-T (Sigma-Aldrich, MO, USA) with 3% NDS
(Jackson Immunoresearch, PA, USA) for 30-60 min at room temperature. They were then incubated in
PBS-T 0.3% with, 3% NDS, and CF405- or CF633-conjugated streptavidin (1:1000; Biotium, CA, USA)
for 90 min at room temperature to reveal neurobiotin labelling. Slices were finally washed four times in
1X PBS (10 min each) before being mounted on glass slides and coverslipped using warmed PVA-
DABCO (Sigma-Aldrich, MO, USA).

Single molecule fluorescent in situ hybridization (smFISH) with RNAscope

C57BL/6 mice were deeply anesthetized with 5% isoflurane and brains were rapidly extracted
and covered with powdered dry ice for ~2 min. Frozen brains were stored in glass vials at -80°C before
sectioning at 20 µm using a cryostat (CM3050 S; Leica Biosystems, Germany) at -16°C. Coronal
sections were thaw-mounted onto a glass slide, by gentle heating from the underside using the tip of a
finger to encourage adhesion of the section to the slide. They were then stored at -80°C until
processing.

Fluorescent *in situ* hybridization (FISH) was performed using the RNAscope Multiplex
Fluorescent assay v2 (Advanced Cell Diagnostics, CA, USA). The following products were used:
RNAscope Multiplex Fluorescent Reagent Kit V2 (Catalog #323110), Fluorescent Multiplex Detection
Reagents (#323110), target probes for *Mus musculus* genes – *Drd1a* (#406491-C1), *Drd2* (#406501-
C3), *Npbwr1* (#547181-C1), and *Vipr2* (465391-C2) – and the Tyramide Signal Amplification (TSA) Plus
Fluorescence Palette Kit (NEL760001KT; PerkinElmer Inc., MA, USA) with fluorophores diluted to
1:1000-1:5000. The protocol was performed as recommended by the manufacturer, with some
modifications to prevent tissue degradation and optimize labelling specificity in our regions of interest.
Fresh frozen slices were fixed in 4% PFA for 1 hr at 4°C. Slices were dehydrated in an ethanol series
(50%, 70%, 100%, and 100% ethanol, 5 min each) and then incubated in hydrogen peroxide for 8 min
at room temperature. Protease treatment was omitted in order to prevent tissue degradation. Slides
were then incubated with the desired probes (pre-warmed to 40°C and cooled to room temperature) for
2 hr at 40°C in a humidified oven. Following washing (2 x 30 s in 1X RNAscope wash buffer), signal
amplification molecules (Amp 1, 2, and 3) were hybridized to the target probes in sequential steps, with
30 min incubation for Amp 1 and 2 and 15 min incubation for Amp 3 at 40°C, all in a 40°C humidified
oven followed by washing (2 x 30 s in wash buffer). For fluorescent labelling of each amplified probe,
slides were incubated in channel-specific HRP for 10 min, followed by incubation with TSA fluorophore
(PerkinElmer, MA, USA) for 20 min, and then incubation in HRP-blocker for 10 minutes (with 2 x 30 s
washes between each step). Probes for *Drd1a*, *Drd2*, *Npbwr1*, and *Vipr2* were each labelled with green
(TSA Plus Fluorescein), red (TSA Plus Cyanine 3), or far red (TSA Plus Cyanine 5) fluorophores in
counterbalanced combinations. Slides were then incubated in DAPI (Advanced Cell Diagnostics, CA,

USA) for 10 min, washed in 1X RNAscope wash buffer, dried for 20 min, coverslipped with warmed PVA-DABCO, (Sigma-Aldrich, St. Louis, MO) and left to dry overnight before imaging.

Images were captured on a confocal laser scanning microscope (Olympus FV1000; Olympus, PA, USA) using a 40X/1.30NA oil immersion objective. Serial Z-stack images were acquired using FluoView software version 4.0 (Olympus, PA, USA) at an optical thickness of 1.5 µm. All images were acquired with identical settings for laser power, detector gain, and amplifier offset. A maximum Z-projection was performed in FIJI followed by rolling ball background subtraction to correct for uneven illumination. Image brightness and contrast were moderately adjusted using FIJI, with consistent adjustments made across images for each probe-fluorophore combination. Regions of interest were annotated on each image by overlaying the appropriate brain atlas image[131,132] with guidance from DAPI staining and using the BigWarp plugin (https://imagej.net/BigWarp)[133] in FIJI. These ROI outlines were used to generate binary masks in order to regionally-restrict subsequent image analysis. Automated cell identification and analysis of fluorescent mRNA labelling was performed in CellProfiler[130] using a modified version of the 'Colocalization' template pipeline (https://cellprofiler.org/examples). The pipeline was optimized to identify DAPI labelling (20-40 pixels in diameter), in order to define cell outlines. This was followed by identification of fluorescent mRNA puncta (2-10 pixels in diameter) for each probe. Puncta that were localized within DAPI-identified cells (classified using the 'relate objects' module) were assigned to that cell for subsequent analysis. Quantification and further analysis/data visualization was performed using a custom-written Python code. Violin plots were made using the violin plot function in the Seaborn library[141] of Python (with smoothing set to 0.2), and colocalization matrices were generated using the Seaborn heatmap function.

In vivo microendoscopic calcium imaging

DAT::Cre mice received an injection of 300 nL AAV$_9$-Syn-FLEX-ChrimsonR-TdTomato in the DRN (ML:1.20, AP:-4.10, DV:-2.90; needle at a 20° angle from the midline, bevel facing medial), and 250 nL AAV$_1$-Syn-GCaMP6f or AAV$_1$-Syn-GCaMP67f in the CeA (ML:2.85, AP:-1.20, DV:-4.75, needle bevel facing posterior). After ~4 weeks mice underwent a second surgery to implant an integrated 0.6 mm diameter, 7.3 mm long gradient refractive index (GRIN) lens with attached baseplate (Inscopix, CA, USA) over the CeA (ML: 2.85, AP:-1.50, DV:-4.60). The lens was lowered slowly into the cleaned craniotomy by hand. The GRIN lens was adhered to the skull by a layer of adhesive cement (C&B Metabond; Parkell Inc., NY, USA) followed by a layer of black cranioplastic cement (Ortho-Jet; Lang, IL, USA), and protected by a small PCR tube cap, held in place by cement. The nVoke miniaturized microscope (Inscopix, CA, USA) consists of a 455±8 nm blue LED for GCaMP excitation, and a 620±30 nm red LED for simultaneous optogenetic manipulation[142].

Behavioral experimentation commenced at least 1 week after baseplate surgery. Mice were first habituated to handling and connection of the microscope for a minimum of 3 consecutive days. For recording, mice were connected to the nVoke miniature microscope by tightening a small set screw on the baseplate. The microscope data cable was connected to a commutator (Inscopix, CA, USA), to allow unrestricted movement, and the commutator was itself connected to a data acquisition (DAQ) box. Grayscale images were acquired at a rate of 20 frames/s (fps; ~50 ms exposure time) with the blue LED delivering 0.2-0.3 mW light power and analog gain on the image sensor set to 2. For the social approach task, mice were placed in the three-chamber apparatus (57.5l x 22.5w x 16.5h chamber with clear walls and grey floors). Following microscope connection, mice freely explored the chamber for 5 min. They were then confined to the center portion of the chamber, by the insertion of

clear Plexiglas panels, during which a novel juvenile mouse was placed under one cup and a novel object was placed under the other cup. The panels were removed, and the test mouse allowed to freely explore for a further 10 min. One 'group-housed' session was conducted without red-light delivery, and another 'group-housed' session was conducted with red 620 nm light delivery (8 pulses with 5 ms pulse-width, at 30 Hz, every 1 s; 10 mW) through the objective lens of the microscope, to activate ChrimsonR-expressing DRN[DAT] terminals. The order of 'group-housed' sessions was counterbalanced. 2-3 weeks later, mice were isolated for 24 hours, and another session ('isolated') commenced without red-light delivery. A top-down SLEAP[143] (v1.3.1) model was trained using 774 labeled frames, annotating a skeleton composed of 15 keypoints (comprised of (1) nose, (2) head, (3) left ear, (4) right ear, (5) neck, (6) left forelimb, (7) right forelimb, (8) trunk, (9) left hindlimb, (10) right hindlimb, (11) tail base, (12-14) points along the length of the tail, and (15) tail tip). Pose estimation using the trained model was then performed on the behavior videos to determine if there was a social or object cup interaction. To determine if there was a social or object cup interaction, the nose of the mouse must be within 1.3x the diameter of the cup, and the cup must be within a 90° cone in front of the mouse's head.

Raw videos of GCaMP fluorescence were first pre-processed in Inscopix Data Processing Software 1.3.0 (Inscopix, CA, USA) by cropping the region outside the GRIN lens, applying 2x spatial downsampling, and a 3x3 median filter to fix defective pixels. A spatial band-pass filter was applied (0.005 to 0.5 oscillations/pixel) to remove high and low spatial frequency content, and rigid motion correction was performed (to account for small lateral displacements) by registering to a stable reference frame with a prominent landmark (e.g. blood vessel). Processed recordings were then exported as TIFF stacks for additional piecewise non-rigid motion correction using the NoRMCorre algorithm[144] in overlapping 64 x 64 pixel grids using a MATLAB implementation. Constrained non-negative matrix factorization for endoscopic recordings (CNMF-E) was then used to extract the spatial shapes and calcium signals from individual cells in the imaging field of view[145] using a MATLAB implementation (key parameters: minimum local correlation for seeding pixels = 0.9, minimum peak-to-noise ratio for seeding pixels =12). The extracted calcium signals were inspected, and non-neuronal objects were manually excluded. All following downstream analyses used raw CNMF-E traces.

Calcium traces were aligned to detected behavioral events (interaction to the social and object cups, as determined by feature thresholds extracted from SLEAP keypoints). Single cell responses to social and object cup interaction were determined using an ROC (receiver operating characteristic) analysis, which has been previously been described to determine neural responses to social behavior[110,111]. A binary behavior vector of social or object cup interaction (calculated in 40ms time bins) was compared to a binary neural activity vectors generated by applying thresholds that span 100 steps from the minimum to maximum z-score value of each calcium trace to determine a true positive rate (TPR) and false positive rate (FPR) at each step. From these values yielded an ROC curve for each neuron that corresponded to the performance of that single neuron in predicting social or object cup interactions. The area under the ROC curve (auROC) was used to determine how strongly modulated each neuron was to the social and object stimuli. To determine the significance of single-cell responses, a null distribution of 1,000 auROC values was generated by randomly circularly shifting the binary behavior vectors and again comparing it to the binary neural activity signal. A neuron was considered having a significant excitatory response to the social or object stimulus if the auROC value exceeded the 97.5[th] percentile of the 1,000 shuffled auROC values, and was considered having a

significant inhibitory response if the auROC value was less than the 2.5th percentile of the 1,000
shuffled auROC values.

Co-registration of active neurons during imaging sessions was performed using CellReg[146]. In
short, the spatial footprint matrices from each imaging session (as determined by CNMF-E) were used
to align different imaging sessions within each animal to a reference session through translational and
rotational shifts. Spatial correlation and centroid distance between cells were used to probabilistically
register active cells across sessions.

Agglomerative hierarchical clustering was performed by averaging each neuron's response to
the onset of social or object cup interaction throughout the trial. A social or object cup interaction was
classified a trial if it (a) lasted a minimum of 1s, (b) if there had been at least 5s that elapsed since the
last interaction, and (c) if there was less than 1.5s pause in interaction with the social or object cups.
The z-scored averaged traces (5s before and after the onset of social or object cup interaction) were
concatenated, such that each row corresponds to one neuronal unit. Agglomerative hierarchical
clustering was performed using MATLAB's "cluster" function. Each neuron was initially designated as
an individual cluster. Those that were in closest proximity were merged to form a new cluster, then the
next closest were merged etc. until a hierarchical tree was formed with all neurons contained within a
single cluster. A threshold at $0.770 \times$ max(linkage) was set to prune branches from the hierarchical
tree, so that all neurons below each cut were assigned to a single cluster. After the dendrogram was
constructed, the average traces (were displayed as a heatmap alongside their corresponding leaf. The
traces of all neurons belonging to a single cluster were then averaged, and the number of neurons that
corresponded to each behavior group was calculated for each cluster.

1439

Statistical analyses

Statistical tests were performed using GraphPad Prism 8 (GraphPad Software, CA, USA). Normality was evaluated using the D'Agostino-Pearson test, and data are expressed as mean±standard error of the mean (SEM), unless otherwise noted. Data which followed a Gaussian distribution were compared using a paired or unpaired t-test (non-directional) for two experimental groups, and a one-way or two-way ANOVA with repeated measures for three or more experimental groups. Data for two experimental groups which did not follow a Gaussian distribution were compared using a Mann-Whitney U test. Correlation between two variables was assessed using the Pearson's product-moment correlation coefficient. Threshold for significance was set at *p<0.05, **p<0.01 and ***p<0.001.

 Figure Legends

Figure 1. DRN^{DAT} and VTA^{DAT} afferents target distinct downstream regions.

(A) Example images of downstream regions showing TH expression from immunohistochemistry.

(B) eYFP expression in the prefrontal cortex (PFC), nucleus accumbens (NAc), bed nucleus of the stria terminalis (BNST), central amygdala (CeA), and posterior basolateral amygdala (BLP) following injection into the DRN (upper panels) and the VTA (lower panels).

(C) Quantification of mean eYFP fluorescence in subregions from each structure (PFC: n=18 and 14 sections, striatum: n=20 and 21 sections, BNST: n=14 and 13 sections, CeA: n=24 and 27 sections, amygdala: n=45 and 51 sections from DRN and VTA injections, respectively, from 6 mice). eYFP fluorescence was significantly greater following VTA injection in all striatal subregions (unpaired t-test: CPu: t_{39}=13.23, p<0.0001; NAc core: t_{39}=13.56, p<0.0001; NAc lateral shell: t_{31}=13.01, p<0.0001; NAc medial shell: t_{37}=4.49, p<0.0001), and significantly greater following DRN injection in the BNST oval nucleus (unpaired t-test: t_{22}=3.95, p=0.0007) and CeA lateral division (unpaired t-test: t_{34}=3.18, p=0.0031).

(D) Images from three selected downstream targets showing average terminal density in the middle anteroposterior (AP) region following eYFP expression in DRN^{DAT} (left) or VTA^{DAT} (right) neurons.

(E) The retrograde tracer cholera toxin subunit-B (CTB) conjugated to Alexa Fluor 555 (CTB-555, pseudo-colored magenta) or Alexa-Fluor 647 (CTB-647, pseudo-colored cyan) was injected into two downstream targets.

(F) Confocal images showing representative injection sites for dual BNST and CeA injections (left panels), BNST and BLP (center panels), and CeA and BLP (right panels).

(G) High magnification images of DRN cells expressing CTB-555 (magenta), CTB-647 (cyan), and TH (green) following injection into the BNST and CeA. White arrows indicate triple-labelled cells.

(H) Venn diagrams showing the proportion of CTB+/TH+ cells in the DRN following dual injections placed in the BNST and CeA (left), BNST and BLP (center), or CeA and BLP (right). When injections were placed in the BNST and CeA, dual CTB-labelled TH+ cells constituted 46% of all BNST projectors and 55% of all CeA projectors. In contrast, when injections were placed in the BNST and BLP, or CeA and BLP, the proportion of dual-labelled cells was considerably lower (7.6% of BNST projectors and 9.7% of CeA projectors).

Bar graphs show mean ±SEM. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. PFC: Cg=cingulate cortex, PL=prelimbic cortex, IL= infralimbic cortex; striatum: CPu=caudate putamen, NAc core=nucleus accumbens core, NAc l.sh.=nucleus accumbens lateral shell, NAc m.sh.=nucleus accumbens medial shell; BNST: oval nuc.= BNST oval nucleus, lat.=BNST lateral division, med.=BNST medial division, vent.=BNST ventral part; CeL=central amygdala lateral division, CeM=central amygdala medial division, CeC=central amygdala capsular division; amygdala: LA=lateral amygdala, BLA=basolateral amygdala, BLP=basolateral amygdala posterior.

Source data 1. Mean DRN^{DAT} eYFP fluorescence in downstream regions, as shown in *Figure 1C*.

Source data 2. Colocalization counts of CTB+/TH+ cells in the DRN, as shown in *Figure 1H*.

 Figure supplement 2—source data 1. Colocalization counts of CTB+/TH+ cells in the DRN, as shown
1490 in *Figure 1—figure supplement 2C*.

1491

Figure 1—figure supplement 1. DRN^{DAT} and VTA^{DAT} eYFP virus injection sites.

(A-D) Confocal images at different AP locations through the VTA and DRN showing the typical spread
of eYFP expression (green) following an injection of AAV$_5$-DIO-ChR2-eYFP into (A) the DRN and (C)
the VTA. Tyrosine hydroxylase (TH; the ratelimiting enzyme in dopamine synthesis) expression from
immunohistochemistry is shown in red. (b,d) Insets showing high magnification images of the
substantia nigra pars compacta (SNc), VTA, rostral linear nucleus (RLi), caudal linear nucleus (CLi),
and DRN. Viral injection in the DRN typically resulted in eYFP-expressing cells within the DRN,
ventrolateral periaqueductal grey (vlPAG), and CLi nuclei, with minimal expression in the RLi, and none
in the VTA or substantia nigra pars compacta (SNc). In contrast, viral injection in the VTA produced
robust eYFP expression in SNc and VTA cell bodies, with some RLi expression, and none in the CLi,
vlPAG, or DRN.

Figure 1—figure supplement 2. Verification of dual-retrograde tracing strategy and
intersectional approach to reveal axon collaterals.

(A) Two retrograde tracers (CTB-555 and CTB-647) were injected into the same location, followed by
sectioning and immunohistochemistry after 7 days. Right panels show example injection site for CTB-
555 and CTB-647 in the BNST.

(B) CTB-expressing cells in the DRN with TH (green) revealed by immunohistochemistry. White arrows
indicate triple-labelled cells.

(C) Within the TH+ cells in the DRN, injection of both retrograde tracers into the same location resulted
in 97% CTB-647+ cells co-labelled with CTB-555, and 100% CTB-555+ cells co-labelled with CTB-647.

(D) Heatmaps indicating the relative density of TH+ CTB+ cells throughout the DRN/CLi for each
projector population and (E) dual-labelled cells. Color intensity represents average number of cells per
slice. The total number of TH+ BNST and CeA projectors per slice was similar (n=27.9 BNST projectors
and n=27.2 CeA projectors per slice), whereas TH+ BLP projectors were significantly fewer in number
(n=6.4 BLP projectors per slice; Kruskal-Wallis statistic = 83.5, p<0.0001; Dunn's posthoc tests: BNST
vs. CeA p>0.05, BNST vs BLP p<0.001, CeA vs BLP p<0.001). TH+ BNST and CeA projectors, and
dual-labelled cells, were broadly distributed throughout the DRN, vlPAG, and CLi, with a higher
concentration in the dorsal aspect of the DRN, whereas BLP projectors tended to be relatively denser in
ventral DRN/CLi.

(F) Injection strategy to enable eYFP expression selectively in the DRN^{DAT}-CeA projection. A
retrogradely-travelling HSV construct encoding mCherry-flpo, expressed in a Cre-dependent manner
(HSV-LS1L-mCherry-IRES-flpo), was injected into the CeA of a DAT::Cre mouse, and an AAV,
expressed in a flpo-dependent manner, encoding eYFP (AAV$_5$-fDIO-eYFP) was injected into the DRN.

(G) After 7 weeks, this resulted in eYFP-expressing TH+ cells in the DRN, and (H) eYFP-expressing
processes in both the CeA (upper panels) and BNST (lower panels).

(I) Injection of only AAV$_5$-fDIO-eYFP into the DRN of a DAT::Cre mouse did not result in eYFP
expression.

Figure 2. DRN^{DAT}-BLP (but not DRN^{DAT}-BNST or DRN^{DAT}-CeA) photostimulation promotes place avoidance.

(A) AAV$_5$-DIO-ChR2-eYFP or AAV$_5$-DIO-eYFP was injected into the DRN of DAT::Cre mice and optic fibers implanted over the BNST, CeA, or BLP to photostimulate DRN^{DAT} terminals. After >7 weeks for viral expression cages of mice were assayed for social dominance using the tube test, prior to other behavioral tasks.

(B-D) Left panels: example tracks of DRN^{DAT}-BNST:ChR2, DRN^{DAT}-CeA:ChR2, and DRN^{DAT}-BLP:ChR2 mice in the real-time place preference (RTPP) assay. Right panels: bar graphs showing the difference in % time spent in the stimulated ('ON') and unstimulated ('OFF') zones. There were no significant RTPP differences detected in (B) DRN^{DAT}-BNST:ChR2 (DRN^{DAT}-BNST:ChR2: N=29 mice, DRN^{DAT}-BNST:eYFP: N=14 mice; unpaired t-test: t_{41}=1.44, p=0.156) and (C) DRN^{DAT}-CeA:ChR2 mice (DRN^{DAT}-CeA:ChR2: N=28 mice, DRN^{DAT}-CeA:eYFP: N=13 mice; unpaired t-test: t_{39}=0.828, p=0.413) compared to their respective eYFP control mice groups. However, (D) DRN^{DAT}-BLP:ChR2 mice spent proportionally less time in the stimulated zone relative to DRN^{DAT}-BLP:eYFP mice (DRN^{DAT}-BLP:ChR2: N=14 mice, DRN^{DAT}-BLP:eYFP: N=8 mice; unpaired t-test: t_{20}=2.13, p=0.0455).

(E-G) Time spent in the ON zone across the 30 min session. (G) DRN^{DAT}-BLP:ChR2 mice spent significantly less time in the ON zone relative to DRN^{DAT}-BLP:eYFP mice (DRN^{DAT}-BLP:ChR2: N=14 mice, DRN^{DAT}-BLP:eYFP: N=8 mice; repeated measures two-way ANOVA: $F_{1,20}$ = 4.53, main effect of opsin p=0.046).

(H-J) Scatter plots showing relative dominance plotted against the difference in zone time (insets show mean values for subordinate, intermediate, and dominant mice) for (H) DRN^{DAT}-BNST, (I) DRN^{DAT}-CeA, or (J) DRN^{DAT}-BLP mice.

Bar and line graphs display mean ±SEM. *p<0.05.

Source data 1. DRN^{DAT}-BNST:ChR2 RTPP percent time difference (ON-OFF), as shown in *Figure 2B*.

Source data 2. DRN^{DAT}-CeA:ChR2 RTPP percent time difference (ON-OFF), as shown in *Figure 2C*.

Source data 3. DRN^{DAT}-BLP:ChR2 RTPP percent time difference (ON-OFF), as shown in *Figure 2D*.

Source data 4. DRN^{DAT}-BNST:ChR2 RTPP percent time in ON (binned), as shown in *Figure 2E*.

Source data 5. DRN^{DAT}-CeA:ChR2 RTPP percent time in ON (binned), as shown in *Figure 2F*.

Source data 6. DRN^{DAT}-BLP:ChR2 RTPP percent time in ON (binned), as shown in *Figure 2G*.

Source data 7. DRN^{DAT}-BNST:ChR2 RTPP percent time difference (ON-OFF) x relative dominance, as shown in *Figure 2H*.

Source data 8. DRN^{DAT}-CeA:ChR2 RTPP percent time difference (ON-OFF) x relative dominance, as shown in *Figure 2I*.

Source data 9. DRN^{DAT}-BLP:ChR2 RTPP percent time difference (ON-OFF) x relative dominance, as shown in *Figure 2J*.

Figure supplement 1—source data 1. Social rank stability, as shown in *Figure 2—figure supplement 1H*.

Figure supplement 2—source data 1. DRN^{DAT}-BNST:ChR2 ICSS number of nose pokes, as shown in *Figure 2—figure supplement 2A*.

Figure supplement 2—source data 2. DRN^{DAT}-CeA:ChR2 ICSS number of nose pokes, as shown in *Figure 2—figure supplement 2B*.

Figure supplement 2—source data 3. DRN^{DAT}-BLP:ChR2 ICSS number of nose pokes, as shown in *Figure 2—figure supplement 2C*.

Figure 2—figure supplement 1. Fiber placement in DRN^{DAT} downstream regions and stability of social dominance within cages.

(A-C) Example confocal images showing ChR2-expressing DRN^{DAT} terminals in (A) the BNST, (B) CeA, and (C) BLP.

(D-F) Fiber placement over (D) the BNST, (E) CeA, and (F) BLP. Colored lines indicate ChR2 subjects whereas grey colored lines indicate eYFP subjects. Lighter shade lines indicate unilateral implants, whereas darker shade lines indicate bilateral implants.

(G) The tube test for social dominance was performed prior to optogenetic manipulations.

(H) Proportion of wins for an individual cage tested across four days, and average for all cages used in optogenetic manipulation experiments, separated by number of mice per cage (red=dominant, orange=intermediate, yellow=subordinate).

 Graphs show mean ±SEM.

Figure 2—figure supplement 2. Photostimulation of DRN^{DAT} projections does not modify operant intra-cranial self-stimulation behavior.

(A-C) Photostimulation of (A) the DRN^{DAT}-BNST, (B) DRN^{DAT}-CeA, or (C) DRN^{DAT}-BLP projection did not support intra-cranial self-stimulation (ICSS) as shown by a lack of preference for the active nosepoke (paired with blue light delivery) over the inactive nosepoke (unpaired t-test: DRN^{DAT}-BNST: DRN^{DAT}-BNST:ChR2: *N*=28 mice, DRN^{DAT}-BNST:eYFP: *N*=16 mice; t_{42}=0.225, p=0.823; DRN^{DAT}-CeA: DRN^{DAT}-CeA:ChR2: *N*=26 mice, DRN^{DAT}-CeA:eYFP: *N*=17 mice; t_{41}=0.225, p=0.823; DRN^{DAT}-BLP: DRN^{DAT}-BLP:ChR2: *N*=14 mice, DRN^{DAT}-BLP:eYFP: *N*=8 mice; t_{20}=0.152, p=0.881).

Figure 3. DRN^{DAT}-BNST (but not DRN^{DAT}-CeA or DRN^{DAT}-BLP) photostimulation promotes non-
social exploratory behavior.

(A-C) Left panels: example tracks in the elevated plus maze (EPM) from a (A) DRN^{DAT}-BNST:ChR2,
(B), DRN^{DAT}-CeA:ChR2, and (C), DRN^{DAT}-BLP:ChR2 mouse. Upper right panels: time spent in the open
arms of the EPM across the 15-minute session. Photostimulation had no significant effect on time spent
in the open arms of the EPM (two-way ANOVA, light x group interaction, BNST – $F_{2,50}$=2.008, p=0.145,
CeA – $F_{2,72}$=0.118, p=0.889, BLP – $F_{2,40}$=0.354, p=0.704) for (A) DRN^{DAT}-BNST, (B), DRN^{DAT}-CeA, or
(C) DRN^{DAT}-BLP mice. Bottom right panels: difference in time spent in open arms of the EPM between
the stimulation ON and first OFF epochs. Photostimulation had no significant effect on time spent in the
open arms of the EPM for (A) DRN^{DAT}-BNST (DRN^{DAT}-BNST:ChR2: N=19 mice, DRN^{DAT}-BNST:eYFP:
N=10 mice; unpaired t-test: t_{27}=1.39, p=0.177), (B) DRN^{DAT}-CeA (DRN^{DAT}-CeA:ChR2: N=23 mice,
DRN^{DAT}-CeA:eYFP: N=14 mice; unpaired t-test: t_{35}=0.639, p=0.527), or (C) DRN^{DAT}-BLP mice (DRN^{DAT}-
BLP:ChR2: N=14 mice, DRN^{DAT}-BLP:eYFP: N=8 mice; unpaired t-test: t_{20}=0.759, p=0.457).

(D-F) Scatter plots showing relative dominance plotted against the difference in the open arm zone time
(insets show mean values for subordinate, intermediate, and dominant mice) for (D) DRN^{DAT}-BNST, (E)
DRN^{DAT}-CeA, or (F) DRN^{DAT}-BLP mice.

(G-I) Home-cage behavior was assessed in the juvenile intruder assay across two counterbalanced
sessions, one paired with photostimulation ('ON') and one without ('OFF') for (G) DRN^{DAT}-BNST, (H)
DRN^{DAT}-CeA, or (I) DRN^{DAT}-BLP mice. DRN^{DAT}-BNST photostimulation increased time spent rearing
(DRN^{DAT}-BNST:ChR2: N=24 mice, DRN^{DAT}-BNST:eYFP: N=13 mice; paired t-test: t_{23}=2,32, p=0.0298),
but DRN^{DAT}-CeA and DRN^{DAT}-BLP photostimulation did not.

(J-L) Scatter plots showing relative dominance plotted against the difference in rearing time with optical
stimulation (ON-OFF) (insets show mean values for subordinate, intermediate, and dominant mice) for
(J) DRN^{DAT}-BNST, (K) DRN^{DAT}-CeA, or (L) DRN^{DAT}-BLP mice.

Bar and line graphs display mean ±SEM. *p<0.05

Figure 3—figure supplement 1. Photostimulation of DRN^{DAT} projections does not modify
locomotor or anxiety-like behavior.

(A-C) Example tracks in the open field test from a (A) DRN^{DAT}-BNST:ChR2, (B) DRN^{DAT}-CeA:ChR2,
and (C) DRN^{DAT}-BLP:ChR2 mouse. Photostimulation had no significant effect on time spent in the
center of the open field (two-way ANOVA, light x group interaction, BNST – $F_{2,90}$=0.2105, p=0.811; CeA
– $F_{2,92}$=0.528, p=0.592; BLP – $F_{2,40}$=0.181, p=0.835) or distance travelled (two-way RM ANOVA, light x
group interaction, BNST – $F_{2,90}$=0.209, p=0.812; CeA – $F_{2,92}$=0.108, p=0.898; BLP – $F_{2,40}$=0.252,
p=0.771) for DRN^{DAT}-BNST, DRN^{DAT}-CeA, or DRN^{DAT}-BLP mice.

Line and bar graphs show mean±SEM.

1630

Figure 4. DRN^{DAT}-CeA (but not DRN^{DAT}-BNST or DRN^{DAT}-BLP) photostimulation promotes sociability in a rank-dependent manner.

(A-C) Heatmaps showing the relative location of ChR2-expressing mice in the three chamber sociability assay, with optic fibers located over the (A) BNST, (B) CeA, or (C) BLP. The task was repeated across two days, with one session paired with photostimulation ('ON') and one without ('OFF').

(D-F) Bar graphs showing social preference in three chamber sociability assay. (D) Photostimulation of DRN^{DAT}-BNST terminals (8 pulses of 5 ms pulse-width 473 nm light, delivered at 30 Hz every 5 s) in ChR2-expressing mice (DRN^{DAT}-BNST:ChR2) had no significant effect on time spent in the social zone relative to the object zone (DRN^{DAT}-BNST:ChR2: N=27 mice, DRN^{DAT}-BNST:eYFP: N=14 mice; 'social:object ratio'; paired t-test: t_{26}=0.552, p=0.586), (E) but increased social:object ratio for DRN^{DAT}-CeA:ChR2 mice (DRN^{DAT}-CeA:ChR2: N=29 mice, DRN^{DAT}-CeA:eYFP: N=13 mice; paired t-test: t_{28}=2.91; corrected for multiple comparisons: p=0.021) (F) and had no significant effect for DRN^{DAT}-BLP:ChR2 mice (DRN^{DAT}-BLP:ChR2: N=14 mice, DRN^{DAT}-BLP:eYFP: N=7 mice; paired t-test: t_{13}=1.62, p=0.130).

(G-I) Scatter plots displaying relative dominance plotted against the change in social zone time with optical stimulation (ON-OFF) for (G) DRN^{DAT}-BNST, (H) DRN^{DAT}-CeA, or (I) DRN^{DAT}-BLP mice, showing significant positive correlation in DRN^{DAT}-CeA:ChR2 mice (Pearson's correlation: r=0.549, p=0.002, N=29 mice). Inset bar graphs show mean values for subordinate, intermediate, and dominant mice.

(J-L) Home-cage behavior was assessed in the juvenile intruder assay across two counterbalanced sessions, one paired with photostimulation ('ON') and one without ('OFF') for (J) DRN^{DAT}-BNST, (K) DRN^{DAT}-CeA, or (L) DRN^{DAT}-BLP mice. DRN^{DAT}-CeA photostimulation in ChR2-expressing mice increased time spent engaged in face investigation with the juvenile mouse (DRN^{DAT}-CeA:ChR2: N=22 mice, DRN^{DAT}-CeA:eYFP: N=14 mice; paired t-test: t_{22}=2.36, p=0.027).

(M-O) Scatter plots showing relative dominance plotted against the difference in face investigation time with optical stimulation (ON-OFF) (insets show mean values for subordinate, intermediate, and dominant mice) for (M) DRN^{DAT}-BNST, (N) DRN^{DAT}-CeA, or (O) DRN^{DAT}-BLP mice.

(P) A two-state Markov model was used to examine behavioral transitions during the juvenile intruder assay for DRN^{DAT}-CeA mice.

(Q-R) Bar graphs showing the difference in transition probability (ON-OFF) for (Q) within-state transitions and (R) across-state transitions, for DRN^{DAT}-CeA:ChR2 and DRN^{DAT}-CeA:eYFP mice. There was no significant difference between ChR2 and eYFP groups for the change in within-state transition probability (DRN^{DAT}-CeA:ChR2: N=22 mice, DRN^{DAT}-CeA:eYFP: N=14 mice; two-way ANOVA: opsin x transition interaction, $F_{1,68}$=3.385, p=0.0702), (R) but there was a significant interaction between opsin and across-state transition probability (DRN^{DAT}-CeA:ChR2: N=22 mice, DRN^{DAT}-CeA:eYFP: N=14 mice; two-way ANOVA: opsin x transition interaction, $F_{1,68}$=4.452, p=0.0385) with photostimulation.

Bar and line graphs display mean ±SEM. *p<0.05, **p<0.01.

Source data 1. DRN^{DAT}-BNST:ChR2 three-chamber social:object ratio, as shown in *Figure 4D.*

Source data 2. DRN^{DAT}-CeA:ChR2 three-chamber social:object ratio, as shown in *Figure 4E.*

Source data 3. DRN^{DAT}-BLP:ChR2 three-chamber social:object ratio, as shown in *Figure 4F.*

Source data 4. DRN^{DAT}-BNST:ChR2 time spent in social zone (ON-OFF) x relative dominance, as
shown in *Figure 4G.*

Source data 5. DRN^{DAT}-CeA:ChR2 time spent in social zone (ON-OFF) x relative dominance, as
shown in *Figure 4H.*

Source data 6. DRN^{DAT}-BLP:ChR2 time spent in social zone (ON-OFF) x relative dominance, as
shown in *Figure 4I.*

Source data 7. DRN^{DAT}-BNST:ChR2 juvenile intruder time spent in face investigation, as shown in
Figure 4J.

Source data 8. DRN^{DAT}-CeA:ChR2 juvenile intruder time spent in face investigation, as shown in
Figure 4K.

Source data 9. DRN^{DAT}-BLP:ChR2 juvenile intruder time spent in face investigation, as shown in
Figure 4L.

Source data 10. DRN^{DAT}-BNST:ChR2 juvenile intruder time spent in face investigation (ON-OFF) x
relative dominance, as shown in *Figure 4M.*

Source data 11. DRN^{DAT}-CeA:ChR2 juvenile intruder time spent in face investigation (ON-OFF) x
relative dominance, as shown in *Figure 4N.*

Source data 12. DRN^{DAT}-BLP:ChR2 juvenile intruder time spent in face investigation (ON-OFF) x
relative dominance, as shown in *Figure 4O.*

Source data 13. DRN^{DAT}-CeA:ChR2 juvenile intruder markov model (transition within state), as shown
in *Figure 4Q.*

Source data 14. DRN^{DAT}-CeA:ChR2 juvenile intruder markov model (transition across states), as
shown in *Figure 4R.*

Figure supplement 1—Source data 1. DRN^{DAT}-BNST:ChR2 three-chamber social zone time, as
shown in *Figure 4—figure supplement 1A.*

Figure supplement 1—Source data 2. DRN^{DAT}-CeA:ChR2 three-chamber social zone time, as shown
in *Figure 4—figure supplement 1B.*

Figure supplement 1—Source data 3. DRN^{DAT}-BLP:ChR2 three-chamber social zone time, as shown
in *Figure 4—figure supplement 1C.*

Figure supplement 2—source data 1. DRN^{DAT}-BNST:ChR2 juvenile intruder rearing time x face
investigation time (ON-OFF), as shown in *Figure 4—figure supplement 2A.*

Figure supplement 2—source data 2. DRN^{DAT}-BNST:ChR2 juvenile intruder rearing time x face
investigation time (ON-OFF), as shown in *Figure 4—figure supplement 2B.*

Figure supplement 2—source data 3. DRN^{DAT}-BNST:ChR2 juvenile intruder rearing time x face
investigation time (ON-OFF), as shown in *Figure 4—figure supplement 2C.*

Figure supplement 2—source data 4. Baseline behavioral measures correlation matrix (r-values), as
shown in *Figure 4—figure supplement 2D.*

Figure supplement 2—source data 5. Baseline behavioral measures correlation matrix (p-values), as shown in *Figure 4—figure supplement 2D*.

Figure supplement 2—source data 6. Baseline behavioral measures (raw values), as shown in *Figure 4—figure supplement 2D-E*.

Figure 4—figure supplement 1. Photostimulation of DRN^{DAT} terminals in CeA (but not in BNST or BLP) increases time spent in three-chamber social zone.

(A-C) Bar graphs showing time spent in the social zone of the three chamber apparatus. (A) Photostimulation of DRN^{DAT}-BNST terminals (8 pulses of 5 ms pulse-width 473 nm light, delivered at 30 Hz every 5 s) in ChR2-expressing mice (DRN^{DAT}-BNST:ChR2) had no significant effect on time spent in the social zone (DRN^{DAT}-BNST:ChR2: N=27 mice, DRN^{DAT}-BNST:eYFP: N=14 mice; 'social:object ratio'; paired t-test: t_{26}=0.165; corrected for multiple comparisons: p>0.999) (B) but increased time spent in the social zone for DRN^{DAT}-CeA:ChR2 mice (DRN^{DAT}-CeA:ChR2: N=29 mice, DRN^{DAT}-CeA:eYFP: N=13 mice; paired t-test: t_{28}=2.88; corrected for multiple comparisons: p=0.015) (C) and had no significant effect for DRN^{DAT}-BLP:ChR2 mice (DRN^{DAT}-BLP:ChR2: N=14 mice, DRN^{DAT}-BLP:eYFP: N=7 mice; paired t-test: t_{13}=1.92; corrected for multiple comparisons: p=0.154).

Figure 4—figure supplement 2. Photostimulation of DRN^{DAT} projections effects on juvenile behavior, and analysis of baseline behavioral traits.

(A-C) Scatter plots displaying the change in face investigation against the change in rearing with photostimulation (ON-OFF) for (A) DRN^{DAT}-BNST:ChR2, (B) DRN^{DAT}-CeA:ChR2, (C) and DRN^{DAT}-BLP:ChR2 mice in the juvenile intruder assay. Outer plots are probability density curves, using kernel density estimation, to show the distribution of each behavior.

(D) Correlation matrix indicating the relationship between baseline behavioral measures for all mice used in Figures 2-4 and associated supplement figures. For the open field test (OFT) and elevated-plus maze (EPM) the first 5 min of the task were used and for the juvenile intruder and 3 chamber assays the data from the 'OFF' session was used.

(E) Principal component analysis (PCA) of behavioral measures with point color representing the social dominance score for each animal. Inset, scree plot showing % variance explained by the first 5 PCs.

Line and bar graphs show mean±SEM.

Figure 5. Spatial segregation of dopamine and neuropeptide receptor populations within DRN^{DAT} terminal fields.

(A) Mean projection of terminal density in the middle anteroposterior (AP) region of the BNST, following eYFP expression in DRN^{DAT} (left) or VTA^{DAT} (right) neurons.

(B) Mean projection showing fluorescent puncta in the BNST indicating detection of *Drd1* (red), *Drd2* (yellow), *Vipr2* (green), or *Npbwr1* (blue) mRNA transcripts.

(C) Line graphs showing the percent of cells expressing each receptor (≥5 puncta) across AP locations for the oval nucleus, dorsolateral BNST, and dorsomedial BNST (two-way ANOVA, oval nucleus: probe x AP interaction, $F_{9,160}$=6.194, p<0.0001, dorsolateral BNST: probe x AP interaction, $F_{12,167}$=3.410, p=0.0002, dorsomedial BNST: probe x AP interaction, $F_{12,161}$=2.268, p=0.0110). *Drd1*: *n*=51,55,53 *Drd2*: *n*=52,55,53 *Vipr2*: *n*=37,39,37 *Npbwr1*: *n*=36,38,38 sections, for oval nucleus, dorsolateral BNST, and dorsomedial BNST, respectively, from 4 mice.

(D) Matrices indicating overlap between mRNA-expressing cells: square shade indicates the percent of cells expressing the gene in the column from within cells expressing the gene in the row.

(E) Mean projection of terminal density in the middle AP region of the CeA, following eYFP expression in DRN^{DAT} (left) or VTA^{DAT} (right) neurons.

(F) Mean projection showing fluorescent puncta in the CeA indicating mRNA expression.

(G) Line graphs showing the % of cells expressing each receptor (≥5 puncta) across AP locations for the CeL, CeM, and CeC (two-way ANOVA, CeL: probe x AP interaction, $F_{12,220}$=8.664, p<0.0001, CeM: main effect of probe, $F_{3,186}$=60.30, p<0.0001, CeC: probe x AP interaction, $F_{12,218}$=4.883, p<0.0001). *Drd1*: *n*=47,40,47 *Drd2*: *n*=70,55,70 *Vipr2*: *n*=65,57,63 *Npbwr1*: *n*=62,50,60 sections, for CeL, CeM, and CeC, respectively, from 4 mice.

(H) Matrices indicating overlap between mRNA-expressing cells.

(I) Mean projection of terminal density in the middle AP region of the BLP, following eYFP expression in DRN^{DAT} (left) or VTA^{DAT} (right) neurons.

(J) Mean projection showing fluorescent puncta in the BLP indicating mRNA expression.

(K) Line graphs showing the percent of cells expressing each receptor (≥5 puncta) across AP locations for the BLP and BMP (two-way ANOVA, BLP: probe x AP interaction, $F_{15,176}$=2.165, p=0.0091, BMP: main effect of probe, $F_{3,141}$=56.92, p<0.0001). *Drd1*: *n*=55,44 *Drd2*: *n*=59,46 *Vipr2*: *n*=41,33 *Npbwr1*: *n*=45,34 sections, for BLP and BMP, respectively, from 4 mice.

(L) Matrices indicating overlap between mRNA-expressing cells. Line graphs show mean ±SEM.

Source data 1. BNST RNAScope sub-regional probe expression (percent), as shown in *Figure 5C*.

Source data 2. BNST RNAScope sub-regional probe co-expression, as shown in *Figure 5D*.

Source data 3. CeA RNAScope sub-regional probe expression (percent), as shown in *Figure 5G*.

Source data 4. CeA RNAScope sub-regional probe co-expression, as shown in *Figure 5H*.

Source data 5. BLA RNAScope sub-regional probe expression (percent), as shown in *Figure 5K*.

Source data 6. BLA RNAScope sub-regional probe co-expression, as shown in *Figure 5L*.

Figure supplement 1—source data 1. Number of puncta x pixels occupied for all RNAScope probes,
as shown in *Figure 5—figure supplement 1B*.

Figure supplement 1—source data 2. BNST RNAScope sub-regional probe expression (percent,
threshold = 1 punctum/cell), as shown in *Figure 5—figure supplement 1D*.

Figure supplement 1—source data 3. BNST RNAScope sub-regional probe expression (percent,
threshold = 3 puncta/cell), as shown in *Figure 5—figure supplement 1E*.

Figure supplement 1—source data 4. CeA RNAScope sub-regional probe expression (percent,
threshold = 1 punctum/cell), as shown in *Figure 5—figure supplement 1G*.

Figure supplement 1—source data 5. CeA RNAScope sub-regional probe expression (percent,
threshold = 3 puncta/cell), as shown in *Figure 5—figure supplement 1H*.

Figure supplement 1—source data 6. BLA RNAScope sub-regional probe expression (percent,
threshold = 1 punctum/cell), as shown in *Figure 5—figure supplement 1J*.

Figure supplement 1—source data 7. BLA RNAScope sub-regional probe expression (percent,
threshold = 3 puncta/cell), as shown in *Figure 5—figure supplement 1K*.

1787

Figure 5—figure supplement 1. Analysis of mRNA expression using different thresholds
qualitatively shows similar spatial pattern of dopamine and neuropeptide receptor expression in
downstream regions.

(A) Workflow for RNAscope and image processing.

(B) Scatter plots showing a linear relationship between fluorescent pixels/cell and number of puncta/cell
for three separate sections for each probe.

(C) Violin plots displaying puncta count per section for each receptor in the BNST (white circle indicates
median; *Drd1*: n=51,55,53 *Drd2*: n=52,55,53 *Vipr2*: n=37,39,37 *Npbwr1*: n=36,38,38 sections, for oval
nucleus, dorsolateral BNST, and dorsomedial BNST, respectively, from 4 mice).

(D) Line graphs for each BNST subregion showing the number of expressing cells when using a
threshold of 1 punctum/cell and (E) 3 puncta per cell.

(F) Violin plots displaying puncta count per section for each receptor in the CeA (white circle indicates
median; *Drd1*: n=47,40,47 *Drd2*: n=70,55,70 *Vipr2*: n=65,57,63 *Npbwr1*: n=62,50,60 sections, for CeL,
CeM, and CeC, respectively, from 4 mice).

(G) Line graphs for each CeA subregion showing the number of expressing cells when using a
threshold of 1 punctum/cell and (H) 3 puncta per cell.

(I) Violin plots displaying puncta count per section for each receptor in the amygdala (white circle
indicates median; *Drd1*: n=55,44 *Drd2*: n=59,46 *Vipr2*: n=41,33 *Npbwr1*: n=45,34 sections, for BLP and
BMP, respectively, from 4 mice).

(J) Line graphs for each amygdala subregion showing the number of expressing cells when using a
threshold of 1 punctum/cell and (K) 3 puncta per cell. These lower thresholds yielded more expressing

cells than using 5 puncta/cell (compare with Fig. 3c,g,k), but with a similar expression pattern across
subregions and AP location.

(L-M) Example images showing expression of *Vipr1* and *Vipr2* within the BNST and CeA. We typically
observed greater *Vipr2* than *Vipr1* expression, and high co-localization, and therefore concentrated our
detailed analyses on *Vipr2*.

Line graphs show mean±SEM.

Figure 6. DRN^{DAT} input distinctly influences downstream activity in each downstream target.

(A-C) In mice expressing ChR2 in DRN^{DAT} neurons, *ex vivo* electrophysiological recordings were made from (A) the BNST, (B), CeA, and (C), BLP.

(D-F) Photostimulation of DRN^{DAT} terminals with blue light (8 pulses delivered at 30 Hz) evoked both excitatory and inhibitory responses at resting membrane potentials in (D) the BNST, (E) CeA, and (F) BLP. Traces show single sweeps and pie charts indicate proportion of cells with no response ('none'), an EPSP only ('excitation'), an IPSP only ('inhibition'), or a mixed combination of EPSPs and IPSPs ('mix'). Recorded cells: BNST *n*=19, CeA *n*=36, BLP *n*=48.

(G-I) When constant current was injected to elicit spontaneous firing, (G) BNST cells responded to photostimulation with an increase in firing ('excitation'), while (H) CeA and (I) BLP cells responded with an increase or a decrease in firing ('inhibition'). Recorded cells: BNST *n*=5, CeA *n*=20, BLP *n*=17.

(J) Properties of the optically-evoked excitatory post-synaptic potential (EPSP) at resting membrane potentials – left: peak amplitude (Kruskal-Wallis statistic = 6.790, p=0.0335; Dunn's posts-hoc tests: CeA vs BLP p=0.0378); middle: change in amplitude across light pulses; right: violin plots showing distribution of onset latencies (white circle indicates median).

(K) Properties of the optically-evoked inhibitory post-synaptic potential (IPSP) at resting membrane potentials – left panel: trough amplitude (one-way ANOVA, $F_{2,31}$=8.150, p=0.0014, CeA vs BLP: **p=0.0014); middle panel: violin plot showing latency to trough peak; right panel: violin plot showing tau for the current decay (white circle indicates median).

(L) Workflow for agglomerative hierarchical clustering of CeA neurons and (M) BLP neurons. Four baseline electrical properties were used as input features (following max-min normalization) and Ward's method used to generate a cluster dendrogram, grouping cells based on Euclidean distance.

(N) Dendrogram for CeA cells indicating two major clusters, with their response to DRN^{DAT} input indicated below each branch (excitatio*n*=black; inhibitio*n*=grey; no response=open).

(O) Upper panels: cluster 1 showed baseline properties typical of 'late-firing' neurons and cluster 2 showed baseline properties typical of 'regular-firing' neurons. Lower panels: pie charts showing the response of cells in each cluster to DRN^{DAT} input.

(P) Dendrogram for BLP cells indicating two major clusters, with their response to DRN^{DAT} input indicated below each branch (excitatio*n*=black; inhibitio*n*=grey; no response=open).

(Q) Upper panels: cluster 1 showed baseline properties typical of pyramidal neurons and cluster 2 showed baseline properties typical of GABA interneurons. Lower panels: pie charts showing the response of cells in each cluster to DRN^{DAT} input.

Bar and line graphs show mean ±SEM. *p<0.05, **p<0.01.

Source data 1. BNST (resting) *ex vivo* responses to DRN^{DAT} optical stimulation, as shown in ***Figure 6D.***

Source data 2. CeA (resting) *ex vivo* responses to DRN^{DAT} optical stimulation, as shown in ***Figure 6E***.

Source data 3. BLP (resting) *ex vivo* responses to DRN^{DAT} optical stimulation, as shown in ***Figure 6F***.

Source data 4. BNST (firing) *ex vivo* responses to DRN^{DAT} optical stimulation, as shown in ***Figure 6G***.

Source data 5. CeA (firing) *ex vivo* responses to DRN[DAT] optical stimulation, as shown in *Figure 6H.*

Source data 6. BLP (firing) *ex vivo* responses to DRN[DAT] optical stimulation, as shown in *Figure 6I.*

Source data 7. BNST/CeA/BLP *ex vivo* EPSP peak amplitude in response to DRN[DAT] optical stimulation, as shown in *Figure 6J.*

Source data 8. BNST/CeA/BLP *ex vivo* EPSP normalized amplitude in response to DRN[DAT] optical stimulation, as shown in *Figure 6J.*

Source data 9. BNST/CeA/BLP *ex vivo* EPSP onset latency in response to DRN[DAT] optical stimulation, as shown in *Figure 6J.*

Source data 10. BNST/CeA/BLP *ex vivo* IPSP trough amplitude in response to DRN[DAT] optical stimulation, as shown in *Figure 6K.*

Source data 11. BNST/CeA/BLP *ex vivo* IPSP trough latency in response to DRN[DAT] optical stimulation, as shown in *Figure 6K.*

Source data 12. BNST/CeA/BLP *ex vivo* IPSP decay tau in response to DRN[DAT] optical stimulation, as shown in *Figure 6K.*

Source data 13. CeA *ex vivo* baseline cell properties used for hierarchical clustering, as shown in *Figure 6L-O.*

Source data 14. BLP *ex vivo* baseline cell properties used for hierarchical clustering, as shown in *Figure 6M-Q.*

Figure supplement 1—source data 1. BNST/CeA/BLP *ex vivo* EPSP/IPSP normalized peak amplitude in response to DRN[DAT] optical stimulation with TTX/4AP application, as shown in *Figure 6—figure supplement 1E.*

Figure supplement 1—source data 2. BNST/CeA/BLP *ex vivo* EPSP/IPSP peak/trough pre-stimulation membrane potential, as shown in *Figure 6—figure supplement 1F.*

Figure supplement 1—source data 3. BNST/CeA/BLP action potential inter-event intervals, as shown in *Figure 6—figure supplement 1G.*

Figure supplement 1—source data 4. CeA baseline cell properties by cluster, as shown in *Figure 6—figure supplement 1H.*

Figure supplement 1—source data 5. BLP baseline cell properties by cluster, as shown in *Figure 6—figure supplement 1I.*

Figure supplement 1—source data 6. Effect of DRN[DAT] input on CeA cell properties by cluster (EPSPs/IPSPs), as shown in *Figure 6—figure supplement 1J.*

Figure supplement 1—source data 7. Effect of DRN[DAT] input on CeA total voltage area by cluster, as shown in *Figure 6—figure supplement 1J.*

Figure supplement 1—source data 8. Effect of DRN[DAT] input on BLP cell properties by cluster (EPSPs/IPSPs), as shown in *Figure 6—figure supplement 1K.*

Figure supplement 1—source data 9. Effect of DRN^{DAT} input on BLP total voltage area by cluster, as
shown in *Figure 6—figure supplement 1K.*

Figure supplement 1—source data 10. CeA/BLP *ex vivo* baseline cell properties used for hierarchical
clustering, as shown in *Figure 6—figure supplement 1L-M.*

Figure 6—figure supplement 1. Effect of DRN^{DAT} photostimulation on downstream cellular
excitability *ex vivo.*

(A-C) Example DIC image, and corresponding eYFP fluorescence, of a brain slice containing (A) the
BNST, (B) CeA, or (C) BLP during *ex vivo* recording. Regional maps show the location of recorded
cells, with color indicating the change in membrane potential elicited by optical stimulation of DRN^{DAT}
terminals.

(D) Example traces showing the optically-evoked EPSP (upper panels) and slow component of the
IPSP (lower panels) was maintained following application of TTX/4AP.

(E) Normalized peak amplitude of the EPSP and IPSP following TTX/4AP (EPSP, *n*=8; IPSP, *n*=3).

(F) Scatter plots showing the amplitude of the optically-evoked EPSP (left) and IPSP (right) recorded in
downstream locations plotted against baseline membrane potential.

(G) Line graphs showing the action potential inter-event interval (IEI) in cells where constant current
was injected to elicit firing. Raw (left) and normalized (right) IEI 5.5 s before and 5 s after optical
stimulation of DRN^{DAT} terminals (blue shading) in BNST, CeA, and BLP cells. Cells which showed a
reduction in IEI with optical stimulation were labelled 'excited' (excit., black) and cells which showed an
increase in IEI with optical stimulation were defined as 'inhibited' (inhib., grey).

(H) Box-and-whisker plots comparing the baseline cell properties (used as input features for
hierarchical clustering; Fig. 4P-U) of the two CeA clusters and (I) the two BLP clusters. Unpaired t-tests
for CeA – ramp ratio: t_{24}=3.502, p=0.0018; max instantaneous firing frequency (max freq$_{inst.}$): t_{24}=4.698,
p<0.0001, firing delay: t_{24}=5.050, p<0.0001, voltage sag: t_{24}=3.983, p=0.0006; unpaired t-tests for BLP
– capacitance: t_{25}=4.803, p<0.0001, max freq$_{inst.}$: t_{25}=15.48, p<0.0001, firing delay: t_{25}=2.743, p=0.0111,
voltage sag: t_{25}=2.705, p=0.0121.

(J) Box-and-whisker plots for the two CeA clusters and (K) the two BLP clusters showing the amplitude
and latency of the EPSP and IPSP, and the combined total voltage area elicited by optical stimulation
of DRN^{DAT} terminals. EPSP peak amplitude, CeA: unpaired t-test, t_{17}=1.40, p=0.180; BLP: unpaired t-
test t_{22}=2.34, p=0.029. EPSP latency, CeA: unpaired t-test, t_{17}=0.673, p=0.510; BLP: Mann-Whitney *U*
= 33.5, p=0.032. Total voltage area, CeA: Mann-Whitney *U* = 22, p=0.0023; BLP: Mann-Whitney *U* =
29, p=0.0019.

(L) Workflow for agglomerative hierarchical clustering of all CeA and BLP neurons combined. Five cell
properties were used as input features, corresponding to the five used in Fig. 4P-U for separate
clustering of CeA and BLP cells.

(M) Dendrogram indicating two major clusters, with the cell location and response to DRN^{DAT} input
indicated by the colored bars below each branch (CeA – pink, BLP – blue; excitation=black;
inhibition=grey; no response=open).

(N) Pie charts showing the response of cluster 1 and cluster 2 CeA cells (upper) and BLP cells (lower)
to optical stimulation DRN^{DAT} input. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001.

1928

Figure 7. Simultaneous calcium imaging of CeA neurons and optogenetic stimulation of DRN^{DAT} terminals in CeA.

(A) AAV$_1$-hSyn-GCaMP7f was injected into the CeA and AAV$_8$-hSyn-FLEX-ChrimsonR-TdTomato or AAV$_1$-CAG-FLEX-TdTomato was injected into the DRN of DAT-Cre mice, and a GRIN lens was implanted over CeA. Experiments were conducted 7 weeks following surgery to allow adequate virus expression in axon terminals.

(B) Example spatial correlation image and extracted ROIs of CeA neurons following calcium imaging processing.

(C) Three chamber sociability paradigm. While group-housed, mice explored a three-chamber apparatus with a novel male juvenile stimulus on one side and a novel object stimulus on the other. During one day of the imaging experiment, DRN^{DAT} terminals were not stimulated, and in another session, DRN^{DAT} terminals were stimulated with red light delivery. Mice underwent a third imaging session, without photostimulation, following 24 hours of social isolation.

(D) Mice first explored the three-chamber apparatus without social or object stimuli for a 5-minute habituation period, then with the social and object stimuli for a 10-minute test period.

(E) Social:object ratio (left) and total social cup interaction time (right) during GH stimulation and no stimulation sessions and 24 hours isolated session in mice expressing ChrimsonR in DRN^{DAT} neurons. Bar and line graphs represent mean ±SEM (N=12 mice; mixed-effects model: $F_{1.897,30.36}$=0.5767, p=0.5591).

(F) Representative traces from CeA calcium imaging during one three chamber imaging session.

(G) Scatter and distribution plots indicating the response strength (auROC) of recorded CeA neurons to social and object cups (GH off: n=429 cells, N=15 mice; GH on: n=441 cells, N=15 mice; SI off: n=484 cells, N=15 mice).

 (H) Difference in response strength (Δ auROC) of CeA neurons to social and object cups (GH off: n=429 cells; GH on: n=441 cells; SI off: n= 484 cells) Kruskal-Wallis test: K-W statistic: 6.172, *p=0.0457; Dunn's multiple comparisons test: GH off vs GH on—p=0.0580, GH off vs SI off—p>0.9999).

(I) Venn diagrams showing overlap of social-encoding neurons (displaying an excitatory response, left, or an inhibitory response, right as defined with auROC) in GH off and GH on sessions (GH off and GH on co-registered neurons: n=202 cells). 16 co-registered GH off cells and 18 GH on cells exhibited an excitatory response to social stimulus with 2 cells having the same response across conditions, whereas 12 co-registered GH off and 11 GH on cells exhibited an inhibitory response with 2 cells having the same response across conditions.

(J) Proportion of CeA neurons responsive to social and object cups, further classified as an excitatory (green) or inhibitory (red) response to the stimulus as defined with auROC.

(K) Proportion of recorded neurons that have an excitatory or inhibitory response to the social cup and (L) to the object cup (N=12 mice).

(M) Correlation between social preference in three chamber task and the proportion of CeA neurons that have an excitatory response to the social cup. The proportion of socially excited neurons is

positively correlated with soc:obj zone ratio only for the GH on condition (pearson correlation: r= 0.6785, p=0.0445, *N*=9 mice). Bar and line graphs show mean ±SEM. *p<0.05.

Source data 1. DRN[DAT]-CeA:ChrimsonR three-chamber social:object ratio and social zone duration, as shown in *Figure 7E.*

Source data 2. CeA response strength to social and object stimuli, as shown in *Figure 7G.*

Source data 3. CeA response strength (change in auROC, social – object), as shown in *Figure 7H.*

Source data 4. CeA response overlap of social-encoding neurons, as shown in *Figure 7I.*

Source data 5. CeA response classification to social and object stimuli, as shown in *Figure 7J.*

Source data 6. Percentage of CeA neurons excited/inhibited by social stimulus, as shown in *Figure 7K.*

Source data 7. Percentage of CeA neurons excited/inhibited by object stimulus, as shown in *Figure 7L.*

Source data 8. Proportion of CeA neurons excited by social stimulus x social:object ratio, as shown in *Figure 7M.*

Figure supplement 1—source data 1. CeA *ex vivo* EPSP/IPSP voltage peak in response to 635nm or 470nm wavelength light, as shown in *Figure 7—figure supplement 1E.*

Figure supplement 1—source data 2. CeA *ex vivo* EPSP/IPSP voltage area in response to 635nm or 470nm wavelength light, as shown in *Figure 7—figure supplement 1F.*

Figure supplement 1—source data 3. CeA *ex vivo* EPSP/IPSP voltage peak in response to just 635nm or simultaneous 635nm and 470nm wavelength light, as shown in *Figure 7—figure supplement 1H.*

Figure supplement 1—source data 4. CeA *ex vivo* EPSP/IPSP voltage area in response to just 635nm or simultaneous 635nm and 470nm wavelength light, as shown in *Figure 7—figure supplement 1I.*

Figure supplement 1—source data 5. DRN[DAT]-CeA:TdTomato three-chamber social:object ratio and social zone duration, as shown in *Figure 7—figure supplement 1J-K.*

Figure supplement 1—source data 6. CeA response strength (change in auROC, social – object), as shown in *Figure 7—figure supplement 1L.*

Figure supplement 1—source data 7. Proportion of CeA neurons excited by object stimulus x social:object ratio, as shown in *Figure 7—figure supplement 1N.*

Figure supplement 1—source data 8. Proportion of CeA neurons inhibited by social stimulus x social:object ratio, as shown in *Figure 7—figure supplement 1O.*

Figure 7—figure supplement 1. Ex vivo validation of simultaneous calcium imaging and photostimulation and behavioral and neural effects of DRN[DAT]-CeA:TdTomato stimulation in CeA.

(A) Representative images of GCaMP-expressing cells in the CeA beneath the GRIN lens, and DRN[DAT]
terminals expressing ChrimsonR.

(B) ChrimsonR was expressed in DRN[DAT] neurons by injection of AAV9-Syn-FLEX-ChrimsonR-
Tdtomato into the DRN of DAT::Cre mice, and (C) after 7 weeks whole-cell patch-clamp
electrophysiological recordings were made from CeA neurons.

(D) Example EPSP and IPSP evoked by delivery of 635 nm red light or 470 nm blue light (8 pulses,
with 5 ms pulse-width, at 30Hz).

(E) Peak amplitude and (F) area of optically-evoked potential elicited by 635 nm (10 mW) and 470 nm
(0.3 and 0.2 mW) light. Inset bar graphs show normalized data. Red light evoked a PSP with a
significantly greater peak amplitude (repeated measures ANOVA, $F_{2,16}=200.1$, $p<0.0001$, red vs 0.3
mW blue: $p<0.0001$, red vs 0.2 mW blue: $p<0.0001$) and area (repeated measures ANOVA, $F_{2,16}=404.2$,
$p<0.0001$, red vs 0.3 mW blue: $p<0.0001$, red vs 0.2 mW blue: $p<0.0001$) than either 0.3 mW or 0.2
mW blue light.

(G) Example EPSP and IPSP evoked by delivery of 635 nm red light alone (8 pulses, with 5 ms pulse-
width, at 30Hz) or during constant 470 nm blue light (0.3 mW) to mimic in vivo recording conditions.

(H) Peak amplitude and (I) area of optically-evoked potential elicited by 635 nm light alone, or during
constant 470 nm light. Inset bar graphs show normalized data. Red light evoked a significantly smaller
PSP in the presence of continuous blue light (peak amplitude: paired t-test: $t_{11}=5.172$, $p=0.0003$;
potential area: paired t-test: $t_{11}=6.431$, $p<0.0001$) similar to a previous report (Stamatakis et al., 2018).
Note that the wavelength of imaging light here (470 nm) is higher than for the nVoke miniature
microscope (455±8 nm), so this experiment may overestimate the constant blue light- induced
suppression of red light-evoked potentials.

(J) Social:object ratio and (K) total social cup interaction time during GH stimulation and no stimulation
sessions and 24 hours isolated session for DRN[DAT]-CeA:TdTomato control animals ($N=2$ mice).

(L) Difference in response strength (Δ auROC) of CeA neurons to social and object cups in DRN[DAT]-
CeA:TdTomato control mice (GH off: $n=39$ cells, $N=2$ mice; GH on: $n=61$ cells, $N=2$ mice; SI off: $n=76$
cells, $N=2$ mice; one-way ANOVA: $F_{2,173}=0.4183$, $p=0.6588$)

(M) (top) ROC curves generated by aligning an example CeA neuron's calcium trace to interaction with
the social (blue) or object (gold) cup in the three-chamber sociability task. This example neuron is
classified as having an excitatory response to the social stimulus while having a neutral response to the
object cup. (bottom) Calcium dynamics of example CeA neuron aligned to mouse's behavior. Blue
shading indicates interaction with the social cup, while gold shading indicates interaction with the object
cup.

(N) No correlation was found between social preference in three chamber task and the proportion of
CeA neurons that have a significantly excitatory response to the object cup (Pearson's correlation: GH
off – r=-0.3231, p=0.6328, $N=9$; GH on – r=-0.1152, p=0.7679, $N=8$; SI off – r=0.3438, p=0.3307, $N=9$).

(O) No correlation was found between social preference in three chamber task and the proportion of
CeA neurons that have a significantly inhibitory response to the social cup (Pearson's correlation: GH
off – r=0.1729, p=0.3625, $N=9$ mice; GH on – r=-0.3941, p=0.2939, $N=8$ mice; SI off – r=0.0116,
p=0.9745, $N=9$ mice).

Bar and line graphs show mean ±SEM. ***$p<0.001$, ****$p<0.0001$.

2045

Figure 7—figure supplement 2. Ex vivo validation of simultaneous calcium imaging and photostimulation and behavioral and neural effects of DRN[DAT]-CeA:TdTomato stimulation in CeA.

2046
2047
2048

2049 (A) Agglomerative hierarchical clustering of trial-averaged CeA traces aligned to interaction with the
2050 social or object cup. The dendrogram (left) reveals 12 functional clusters of neurons, as displayed by
2051 the heatmap of trial-averaged neural activity (right).

2052 (B) Cluster-averaged traces aligned to the onset of social cup (blue) or object cup (gold) interaction for
2053 each cluster. The percentage of neurons per condition is listed in each inset.

2054 Line graphs show mean ±SEM.

2055

2056

Figure 8. DRN^{DAT}-CeA photoinhibition blocks isolation-induced sociability.

(A) AAV$_5$-DIO-NpHR-eYFP or AAV$_5$-DIO-eYFP was injected into the DRN of DAT::Cre mice and optic
fibers implanted over the BNST, CeA, or BLP to photoinhibit DRN^{DAT} terminals.

(B) After >7 weeks for viral expression mice were assayed on the three-chamber sociability assay with
delivery of continuous yellow light for photoinhibition, once when group-housed and once following 24
hours of social isolation (2-3 weeks after the initial session).

(C) Photoinhibition of DRN^{DAT}-BNST terminals in NpHR-expressing mice (DRN^{DAT}-BNST:NpHR) had no
significant effect on time spent in the social zone relative to the object zone (DRN^{DAT}-BNST:NpHR: *N*=7
mice, DRN^{DAT}-BNST:eYFP: *N*=5 mice; 'social:object ratio'; two-way RM ANOVA: light x group
interaction, $F_{1,10}$=1.005, p=0.3397), but reduced social:object ratio for isolated DRN^{DAT}-CeA:NpHR mice
compared to isolated DRN^{DAT}-CeA:eYFP mice (DRN^{DAT}-CeA:NpHR: *N*=20 mice, DRN^{DAT}-CeA:eYFP:
N=12 mice; 'social:object ratio'; two-way RM ANOVA: light x group interaction, $F_{1,30}$=4.909, p=0.0344;
multiple comparisons test, DRN^{DAT}-CeA:NpHR^{SI} vs DRN^{DAT}-CeA:eYFP^{SI} adjusted **p=0.0017). In
addition, terminal photoinhibition had no effect for DRN^{DAT}-BLP:NpHR mice (DRN^{DAT}-BLP:NpHR: *N*=6
mice, DRN^{DAT}-BLP:eYFP: *N*=8 mice; 'social:object ratio'; two-way RM ANOVA: light x group interaction,
$F_{1,12}$=3.346, p=0.0923). Inset bar graphs show the difference in social:object ratio in isolated and
grouped conditions. A significant difference between NpHR^{CeA} and eYFP^{CeA} groups was observed
(unpaired t-test: t_{29}=2.177, p=0.0377).

(D) Scatter plots displaying relative dominance plotted against the change in social zone time (isolated-
grouped), showing significant negative correlation in NpHR^{CeA} mice (Pearson's correlation: r=-0.500,
p=0.0414, *N*=20 mice).

Bar and line graphs show mean ±SEM. *p<0.05, **p<0.01.

Source data 1. DRN^{DAT}-ALL:NpHR three-chamber social:object ratio (GH on and SI on), as shown in
Figure 8C.

Source data 2. DRN^{DAT}-ALL:NpHR three-chamber social:object ratio (SI – GH) x relative dominance,
as shown in ***Figure 8D.***

Figure supplement 1—source data 1. DRN^{DAT}-ALL:NpHR three-chamber social:object ratio (GH off
and GH on), as shown in ***Figure 8—figure supplement 1B***.

2085

2086 **Figure 8—figure supplement 1. Photoinhibition of DRN^{DAT}-BNST:NpHR, DRN^{DAT}-CeA:NPHR,**
2087 **DRN^{DAT}-BLP:NpHR terminals does not affect social preference in group housed mice.**

2088 (A) Schematic showing three-chamber behavior paradigm in group-housed DRN^{DAT}:NpHR or
2089 DRN^{DAT}:eYFP mice, with and without yellow light for photoinhibition.

2090 (B) Photoinhibition had no significant effect on social:object ratio in group-housed mice (two-way
2091 ANOVA, light x group interaction, BNST – $F_{1,11}$ p=0.4571, DRN^{DAT}-BNST:NpHR: N=7 mice, DRN^{DAT}-
2092 BNST:eYFP: N=5 mice; CeA – $F_{1,31}$=0.1353, p=0.7154, DRN^{DAT}-CeA:NpHR: N=20 mice, DRN^{DAT}-
2093 CeA:eYFP: N=12 mice; BLP – $F_{1,14}$=2.517, p=0.1349, DRN^{DAT}-BLP:NpHR: N=6 mice, DRN^{DAT}-
2094 BLP:eYFP: N=8 mice).

2095 Line and bar graphs show mean±SEM.

2096

Figure 1

A downstream TH expression

B downstream eYFP expression

C PFC · striatum · BNST · CeA · amygdala

D terminal density: selected downstream targets

BNST

CeA

BLP

E

F BNST CeA · BNST BLP · CeA BLP

G DRN

H Proportion of CTB+/TH+ cells in DRN

n=1682 cells, 6 mice · n=534 cells, 3 mice · n=754 cells, 4 mice

injection site expression

A DRN^DAT-BNST
B DRN^DAT-CeA
C DRN^DAT-BLP

Figure 4

Figure 6

A — BNST eYFP fluoresence — voltage area (mV·s)

B — CeA eYFP fluoresence — voltage area (mV·s)

C — BLP eYFP fluoresence — voltage area (mV·s)

D — EPSP baseline / TTX/4AP; IPSP baseline / TTX/4AP

E — normalized peak amplitude TTX/4AP

pre-stimulation membrane potential

action potential inter-event interval (IEI)

F — EPSP: CeA, BNST, BLP; IPSP: CeA, BNST, BLP

G — BNST / CeA / BLP — IEI (s), normalized, excit., inhib.

CeA

BLP

baseline cell properties

H — ramp ratio **; max freq.$_{inst}$ (Hz) ****; firing delay (s) ****; voltage sag (mV) ***; cluster 1, 2

I — capacitance (pF) ****; max freq.$_{inst}$ (Hz) ****; firing delay (s) *; voltage sag (mV) *; cluster 1, 2

effect of DRN^{DAT} input

J — EPSP: peak amplitude (mV), latency (ms); IPSP: trough amplitude (mV), latency (s); combined: total voltage area (mV·s) **; cluster 1, 2

K — EPSP: peak amplitude (mV) *, latency (ms) *; IPSP: trough amplitude (mV), latency (s); combined: total voltage area (mV·s) **; cluster 1, 2

hierarchical clustering with CeA and BLP cells combined

L — combined cell properties → input features → hierarchical clustering

(1) ramp ratio
(2) firing delay
(3) voltage sag
(4) capacitance
(5) max freq.$_{inst}$

cell properties
(1) (2) (3) (4) (5)
cell 1 16.5 9.4 7.4 24.7 5.3
cell 2 22.6 57.2 32.7 83.8 72.1
max-min normalized values

cluster dendrogram — distance / individual cells

M — Euclidean distance; cluster 1, cluster 2

N — cluster 1, cluster 2; CeA: excitation, inhib., excit., none, inhibition; BLP: inhibition, none, excitation, inhib.

A three chamber sociability

group-housed (GH)

OFF
social   object
day 1

counter-balanced

ON
social   object
day 2

B

DRN^DAT-BNST
■ DRN^DAT-BNST:NpHR
■ DRN^DAT-BNST:eYFP

social:object ratio

OFF ON   OFF ON
GH   GH

DRN^DAT-CeA
■ DRN^DAT-CeA:NpHR
■ DRN^DAT-CeA:eYFP

social:object ratio

OFF ON   OFF ON
GH   GH

DRN^DAT-BLP
■ DRN^DAT-BLP:NpHR
■ DRN^DAT-BLP:eYFP

social:object ratio

OFF ON   OFF ON
GH   GH

Terminal inhibition