The hepatitis E virus capsid protein ORF2 counteracts cell-intrinsic antiviral responses to enable persistence in hepatocytes

Ann-Kathrin Mehnert[1,2], Sebastian Stegmaier[2,3#], Carlos Ramirez[4#], Vladimir Gonçalves Magalhães[3], Carla Siebenkotten[1], Jungen Hu[1,2], Ana Luisa Costa[4], Daniel Kirrmaier[5,6], Michael Knop[5,6], Xianfang Wu[7], Thibault Tubiana[8], Carl Herrmann[4], Marco Binder[3], and Viet Loan Dao Thi[1,9]*

[1]Schaller Research Group, Department of Infectious Diseases, Virology, Heidelberg University, Medical Faculty Heidelberg, Heidelberg, Germany; [2]Faculty of Biosciences, Heidelberg University, Heidelberg, Germany; [3]Research Group "Dynamics of Early Viral Infection and the Innate Antiviral Response", Division Virus-Associated Carcinogenesis (D430), German Cancer Research Center (DKFZ), Heidelberg, Germany; [4]Institute for Pharmacy and Molecular Biotechnology, Heidelberg University & BioQuant, Heidelberg, Germany; [5]Center for Molecular Biology of Heidelberg University (ZMBH), Heidelberg, Germany; [6]Division Cell Morphogenesis and Signal Transduction (A260), German Cancer Research Center (DKFZ), Heidelberg, Germany; [7]Infection Biology Program and Department of Cancer Biology, Lerner Research Institute, Cleveland Clinic Foundation, Cleveland, OH, USA; [8]Institute for Integrative Biology of the Cell (I2BC), Université Paris-Saclay, CEA, CNRS, Gif-sur-Yvette, France; [9]German Center for Infection Research (DZIF), Partner Site Heidelberg, Heidelberg, Germany

[#]Authors contributed equally
*Corresponding author: VietLoan.DaoThi@med.uni-heidelberg.de

Classification: Biological Sciences, Microbiology

Keywords: Hepatitis E virus, cell-intrinsic antiviral response, interferon response, viral antagonism, single-cell RNA-sequencing, viral persistence

Electronic word count: 11.441

Number of figures and tables: 4 figures

Abstract

Hepatitis E virus (HEV) is a significant human pathogen causing both acute and chronic infections

worldwide. The cell-intrinsic antiviral response serves as the initial defense against viruses and

has been shown to be activated upon HEV infection. HEV can replicate in the presence of this

response, but the underlying mechanisms remain poorly understood.

Here, we investigated the role of HEV structural proteins ORF2 and ORF3 in immunocompetent

cells. Mechanistically, we found that ectopic ORF2, but not ORF3, interfered with inflammatory

signaling pathways as well as with antiviral signaling downstream of pattern recognition receptors.

The latter was at least in part mediated through direct interaction with the central adaptor protein

TANK binding kinase 1. In addition, we discovered a protective mechanism mediated by ORF2

that shielded viral replication from antiviral effectors. Using single-cell RNA-sequencing, we

confirmed that the ORF2-mediated antagonism in infected cells resulted in dampened antiviral

responses in both HEV-infected and bystander cells. As a consequence, we found that early in

the infection process, the progression of authentic HEV infection relied on the presence of ORF2,

facilitating a balance between viral replication and the antiviral response within immunocompetent

cells. Altogether, our findings shed new light on the multifaceted role of ORF2 in the HEV life cycle

and improve our understanding of the determinants that may contribute to HEV persistence.

Significance statement

Hepatitis E virus (HEV) is an important yet often underestimated pathogen. Depending on the

genotype, HEV infections can progress to chronicity, but the underlying mechanisms remain

poorly understood. To gain insight into potential determinants, we investigated how HEV evades

the body's first line of defense, the cell-intrinsic antiviral response. We discovered that the HEV

capsid protein ORF2 is crucial in limiting this response by interfering with antiviral signaling

pathways and shielding viral replication from immune effectors. This balance between viral

replication and the antiviral response contributes to persistent HEV infection in immunocompetent

cells. Our findings reveal a new role for the HEV capsid protein in the viral life cycle and highlight

it as an important target for novel therapeutic approaches.

Introduction

Hepatitis E virus (HEV) is one of the major causes of acute hepatitis across the globe, affecting an estimated 20 million people every year (reviewed in[1,2]). As part of the *Hepeviridae* family, the *Paslahepevirus balayani* genus comprises eight genotypes of which HEV-1 to HEV-4, and more recently HEV-7, have been associated with human infections. HEV-1 and HEV-2 are transmitted fecal-orally and are restricted to humans. In contrast, HEV-3 and HEV-4 spread to humans zoonotically through consumption of meat products from domestic pigs or wild boar. Although HEV-1/-2 infections are usually self-limiting, they are associated with high mortality in pregnant women. Further, HEV-3 and -4 infection of immunocompromised individuals can result in chronicity and development of liver fibrosis and cirrhosis.

HEV is ingested and excreted as a naked virus but circulates in the bloodstream wrapped in a host membrane-derived quasi-envelope, which is acquired during budding from cells (reviewed in[3]). It has a positive-sense, single-stranded RNA genome of approximately 7.2 kilobases, encompassing three open reading frames (ORFs) that give rise to three viral proteins (reviewed in[2]). ORF1 contains the domains involved in viral replication, such as the RNA-dependent RNA polymerase (RdRp). ORF3 is a small phosphoprotein that mediates secretion of viral progeny.

Three distinct isoforms of ORF2 have been described in HEV-infected cells[4-7]. Owing to its N-terminal signal peptide, ORF2 can be secreted along the secretory pathway where it gives rise to the glycosylated (ORF2g) and cleaved (ORF2c) isoforms. ORF2g is secreted as a dimer[5,6] and serves as an immunological decoy *in vivo*[5]. It has been postulated that positively charged residues within an arginine-rich motif (ARM; five arginine residues: RRRGRR) downstream of the signal peptide assist ORF2 in retaining a cytosolic orientation at the endoplasmic reticulum membrane[4]. This intracellular ORF2 isoform (ORF2i) assembles into infectious progenies. ORF2i is mainly located in the cytosol but can translocate to the nucleus, and is likely involved in many virus-host interactions[4,7]. Another study highlighted the importance of two ORF2 start codons. The first start codon can give rise to the secreted ORF2g, while the second gives rise to ORF2i with intracellular localization[5]. Although it remains unclear whether differential usage of these start codons is relevant in HEV infection, they can be used to abrogate expression of individual or all ORF2 isoforms.

Cell-intrinsic defense strategies are initiated by recognition of pathogen-associated molecular patterns (PAMPs) through pattern recognition receptors (PRRs) (reviewed in[8]). In epithelial cells, double-stranded (ds)RNA, the replication intermediate of RNA viruses, is detected by the family of retinoic acid-inducible gene I (RIG-I)-like receptors (RLRs) in the cytosol, including RIG-I and melanoma differentiation-associated protein 5 (MDA5), which are additionally regulated by laboratory of genetics and physiology 2 (LGP2). In the endosome, Toll-like receptor 3 (TLR3) recognizes dsRNA. Different adaptor proteins mediate downstream signaling: Toll/interleukin-1 receptor (TIR) domain-containing adaptor-inducing interferon-β (TRIF) is recruited by TLR3, while RIG-I and MDA5 induce polymerization of mitochondrial antiviral signaling protein (MAVS). Both result in activation of TANK binding kinase 1 (TBK1) and the inhibitor of nuclear factor-κB (IκB) kinase (IKK) complex. TBK1 phosphorylates interferon regulatory factor 3 (IRF3), which translocates into the nucleus and induces expression of type I and type III interferons (IFNs) and IFN-stimulated genes (ISGs). Autocrine and paracrine IFN signaling through Janus kinase/signal transducers and activators of transcription (JAK/STAT) results in expression of hundreds of ISGs

with antiviral functions (reviewed in[9]). Activation of the IKK complex leads to nuclear translocation
of nuclear factor-κB (NF-κB) and expression of inflammatory cytokines.

Despite growing interest and advances in recent years, the cell-intrinsic antiviral response
to HEV infection remains poorly described. HEV-infected chimpanzees had a similar but weaker
ISG response compared to hepatitis C virus (HCV) infection, likely owing to the lower HEV
viremia[10]. *In vitro*, the HEV RNA has been identified as a potent agonist for PRR-mediated IFN
induction[11]. As a result, infection of innate immunocompetent hepatocyte models with HEV
induced robust type III IFN responses[12,13]. The induced IFN response appeared to be persistent
but was unable to eliminate viral replication within the observed time frames[12,13]. Blunting of IFN
induction did not result in enhanced viral replication, suggesting that HEV has developed
mechanisms to persistently replicate in the presence of a sustained antiviral response[13].
Corroborating this, several studies have shown that HEV replication is relatively resistant to
exogenous IFN treatment, as compared to, for example, HCV[12,14-16]. Yin and colleagues proposed
that this is the result of an internal IFN refractoriness due to persistent activation of JAK/STAT1
signaling and retention of phosphorylated STAT1 in the cytosol rather than a direct viral
antagonism[12].

However, all HEV proteins have previously been suggested to antagonize the cell-intrinsic
antiviral response (reviewed in[17]). Within ORF1, the X domain, the putative papain-like cysteine
protease (PCP) domain, and the methyltransferase (MeT) domain were shown to interfere with
type I IFN induction and NF-κB-mediated signaling[18-21]. A combined MeT-Y-PCP polyprotein was
further found to interfere with the JAK/STAT pathway[22]. The ORF2 protein has been reported to
interact with TBK1[23,24] and hinder activation of NF-κB[4,25]. Studies of potential ORF3-mediated
antagonisms have been controversial: while some studies suggested that ORF3 interferes with
type I IFN production, NF-κB-, and IFN-mediated signaling[26-28], one study showed that ORF3
could enhance IFN induction[29]. However, most investigations have relied on overexpression of
individual HEV proteins or subdomains of ORF1 and the physiological relevance of each
antagonism for HEV replication remains elusive.

In the present study, we sought to clarify the role of HEV ORF2 and ORF3 proteins in
antagonizing antiviral responses and their impact on HEV infection. To address this, we made
use of viral mutants and assessed differences in IFN and ISG responses upon authentic infection
of different immunocompetent hepatocellular systems in bulk and at single-cell resolution. We
identified a replication-limiting bottleneck mediated by the antiviral response early in infection that
is overcome by the capsid protein ORF2, but not ORF3, allowing for the establishment of
persistent viral replication.

Results

The HEV ORF2 protein antagonizes both antiviral and inflammatory signaling pathways and
interacts with TBK1

Several studies have suggested that HEV ORF2 and ORF3 antagonize different steps of
the cell-intrinsic antiviral response[23-28]. We aimed to validate these findings and first analyzed their
interference with the sensing pathway. For this mechanistic investigation, we made use of readily
available A549-derived cell lines, which harbor a double knockout of RIG-I and MDA5 and show
negligible TLR3 expression[30]. In addition to reconstitution with weak transgene overexpression of
either RIG-I[31], MDA5[31], or TLR3, we ectopically expressed HEV ORF2, ORF3, or GFP in these

cells. In GFP-expressing cells, stimulation of the respective PRR with Sendai virus (SeV), Mengo-Zn virus, and poly(I:C) supernatant feeding resulted in the induction of *IFNB1* expression (Fig. 1A-C). Ectopic expression of HEV ORF3 did not significantly alter *IFNB1* expression downstream of RIG-I and MDA5 (Fig. 1A-B) but resulted in a minor enhancement downstream of TLR3 compared to the GFP control (Fig. 1C). In cells expressing ectopic ORF2, *IFNB1* induction was significantly lower downstream of all PRRs compared to GFP-expressing cells (Fig. 1A-C). Not only ORF2 from a chronic HEV genotype (HEV-3) but also from an acute genotype (HEV-1) dampened this response (Fig. 1A-C). By ELISA, we confirmed that ectopic ORF2 expression also decreased IFNβ protein secretion (Suppl. fig. 1A-C). While the presence of ectopic ORF2 dampened the strength of *IFNB1* induction, it did not alter the dynamics of RLR signaling, which we tested by time-resolved RT-qPCR analysis of MDA5-expressing cells electroporated with poly(I:C) (Suppl. fig. 1D-E).

We further investigated the impact of ORF2 expression on cytokine-dependent antiviral and inflammatory signaling. Upon TNF stimulation, the presence of ectopic ORF2, but not ORF3, reduced the induction of the inflammatory cytokine *TNFAIP3* significantly (Fig. 1D), thus interfering with NF-κB-dependent signaling. On the other hand, we did not observe significant differences in expression of the ISG *IFIT1* upon stimulation with IFNβ in the presence of either ectopic ORF2 or ORF3, suggesting that neither interfered with JAK/STAT signaling (Fig. 1E).

In agreement with previous studies, these results confirmed that ORF2 interferes with the sensing pathway upstream of IFN induction[23,24] as well as with NF-κB-dependent inflammatory cytokine induction downstream of the TNF receptor. In contrast, ORF3 did not affect antiviral and inflammatory responses as strongly as proposed previously[26-29]. As NF-κB-dependent cytokine induction is particularly important for the crosstalk with immune cells *in vivo*, we continued focusing on the effects of ORF2 on the antiviral IFN/ISG response.

It was previously suggested that ORF2 interacts with the critical adaptor molecule TBK1, likely through the ARM[24], but a direct interaction through this motif was not analyzed. Here, we confirmed the interaction of HEV-3 ORF2 and TBK1 by co-immunoprecipitation (co-IP) (Fig. 1F). We further found that mutation of the last two arginine residues of the ARM to alanine (2R/2A) decreased the interaction of TBK1 with ORF2 (Fig. 1F). However, this particular mutation abrogates nuclear translocation of ORF2i and increases ORF2 secretion and glycosylation along the secretory pathway[4]. Hence, the pool of cytosolic ORF2i is reduced[4] and thus, also the likelihood of interaction with TBK1. In agreement with this, the ORF2-2R/2A protein appeared with a light smear in our Western blot (WB) analysis as compared to WT ORF2, suggesting the presence of post-translational modifications such as glycosylation (Fig. 1F).

In an effort to determine other potential interaction motifs between ORF2 and TBK1, we made use of AlphaFold. Repeated modeling identified the last three residues of the LGSAWRD motif (spanning amino acids 83 to 89) at the N-terminus of ORF2 as the only promising, putative interaction motif (Suppl. fig. 2A, 2C-E). However, mutation of these residues to triple alanine did not result in impaired interaction between ORF2 and TBK1, suggesting that this predicted motif is not essential (Suppl. fig. 2B).

Altogether, we have shown that ORF2 counteracts IFN induction downstream of all relevant PRRs and additionally interferes with NF-κB-mediated signaling. The antagonism of the antiviral response by ORF2 is mediated, at least in part, by interaction with TBK1, although the precise interaction motif remains elusive.

The capsid protein ORF2, but not the ORF3 protein, is pivotal for efficient HEV replication in
immunocompetent cells

HEV replication in hepatocytes persists in the presence of a sustained cell-intrinsic
antiviral response[12,13]. To study how far HEV replication is dependent on ORF2-mediated escape
of the antiviral response, we generated a ΔORF2 and a control ΔORF3 mutant of the HEV-3
Kernow-C1/p6 (wild-type, WT) strain (Fig. 2A). For the ΔORF2 mutant, we mutated the two start
codons of *ORF2* to abrogate protein expression of all ORF2 isoforms without affecting correct
translation of the overlapping *ORF3*, as suggested by Yin *et al*.[5]. Upon electroporation (EPO) into
immunocompetent HepG2/C3A cells[12,32], we observed significantly reduced replication of the
ΔORF2 mutant compared to both WT and the ΔORF3 mutant at day 5 and day 7 post-EPO (Fig.
2B). Of note, at least part of the basal HEV RNA detected up to day 3 post-EPO resulted from
high levels of incoming, electroporated RNA, as revealed by comparison with a replication-
incompetent GNN mutant of the RdRp. Compared to HEV WT, ΔORF2 replication induced a
stronger antiviral response, as evidenced by significantly increased *IFNL1* (Fig. 2C) and *ISG15*
(Fig. 2D) expression at day 5 and day 7 post-EPO (corresponding non-normalized data in Suppl.
fig. 3A-B). Since the ΔORF3 virus replicated at WT levels and ORF3 is the protein responsible for
secretion of progeny, we ruled out effects of ORF2 and ORF3 deletion on secondary virus spread.
In addition, newly produced virus particles released into the supernatant are quasi-enveloped and
therefore, not highly infectious[33].

We hypothesized that the antiviral response induced in the absence of ORF2 was strong
enough to cause, at least in part, the observed decline in viral replication. In order to address this,
we exogenously inhibited IFN and ISG induction by treating the cells with the TBK1 inhibitor
BX795. Prolonged incubation with BX795 can affect cell viability, as TBK1 is involved in many
cellular processes. To minimize such effects (Suppl. fig. 3C-E), we incubated the cells with BX795
for only 48 h before harvesting. Application of BX795 reduced *IFNL1* and *ISG15* induction in
response to viral replication (Suppl. fig. 3F-H). HEV WT replication was not significantly enhanced
(Fig. 2E), while ΔORF3 replication showed only a minor increase (Fig. 2G). In contrast, the BX795
treatment resulted in a significant and stronger increase in ΔORF2 replication (Fig. 2F). This
implied that the antiviral response, which peaked at day 5 post-EPO (Fig. 2C), had a greater effect
on viral replication in the absence of ORF2, in line with the ORF2-mediated antagonism described
above.

To study this observation at the single-cell level, we detected HEV genomes by RNA-
fluorescence *in situ* hybridization (RNA-FISH), allowing us to quantify the percentage of HEV-
infected cells over time, independent of viral antigen expression. As early as day 3 post-EPO,
ΔORF2 RNA signals were weaker compared to WT and ΔORF3 and continued to decline
thereafter (Fig. 2H-I). Analogous to our results in Fig. 2B, the decrease in WT- and ΔORF3-
positive cells from day 3 to day 5 post-EPO was likely due to residual incoming RNA introduced
by the EPO process (Fig. 2I). Overall, the percentages of WT- and ΔORF3-positive cells remained
comparable between day 5 and day 7 post-EPO (Fig. 2I). In contrast, we barely detected ΔORF2
RNA at these days as evidenced by a significant decrease compared to day 3. Interestingly,
ΔORF2 RNA was still detectable in our bulk RT-qPCR analysis at these time points (Fig. 2B).
Since the RNA-FISH assay is based on the hybridization of 20 probe pairs to an *ORF2* target
region spanning approximately 1,000 nucleotides[34], whereas RT-qPCR only amplifies a short

sequence stretch of 70 nucleotides, it is possible that we majorly detected degraded ΔORF2
genomes by RT-qPCR.
Next, we aimed to determine whether the absence of ORF2 resulted in increased
sensitivity of viral replication to the action of antiviral effectors. To focus solely on the effects of
ISGs, we used Huh7.5 cells, which are only weakly responsive to PRR stimulation due to a non-
functional RIG-I and negligible basal expression levels of MDA5[35], TLR3[35], and LGP2[36].
Therefore, HEV replication should not induce an antiviral response. Indeed, EPO of Huh7.5 cells
with WT and ΔORF2 RNA resulted in comparable viral replication (Suppl. fig. 4A), likely due to
the lack of an antiviral response in these cells (Suppl. fig. 4B). Next, we treated WT- and ΔORF2-
electroporated Huh7.5 cells four days post-EPO with either IFNλ1 or IFNα2A for 72 h to induce
ISG expression. Comparable ISG induction between WT and ΔORF2 further confirmed that ORF2
did not interfere with JAK/STAT signaling (Suppl. fig. 4C-D). Importantly, we neither detected
IFNB1 nor *IFNL1* expression upon IFN treatment of WT- and ΔORF2-electroporated cells. Thus,
we could exclude the possibility that the IFN treatment led to an upregulation of PRR expression
and consequently, an enhanced ISG induction in a positive feedback loop due to enhanced
sensing of HEV RNA. Nonetheless, we observed that ΔORF2 replication was more strongly
inhibited by IFN treatment and the ensuing ISG response than WT replication (Fig. 2J). We
therefore concluded that ORF2 replication was more sensitive to the antiviral effects of ISGs
induced by exogenous IFN treatment, even in the absence of a directly virus-induced antiviral
response. This additionally contributed to the strong decrease in viral replication in the absence
of ORF2 (Fig. 2B).
Altogether, we found that the antagonism mediated by ORF2 is critical for efficient HEV
replication by dampening the induced antiviral response. Further, the absence of ORF2 resulted
in increased sensitivity of HEV replication to the action of ISGs, suggesting additional protective
functions of the ORF2 protein, independently of its direct interaction with TBK1.
257
A balance between HEV replication and the antiviral response at an early infection bottleneck is
essential for sustained viral replication
We next sought to identify the time point at which the presence of ORF2 is decisive for
the fate of HEV replication in authentic infection. To this end, we established a trans-
complementation system in hepatoma S10-3 cells ectopically expressing ORF2 to produce
ΔORF2 virus particles (Fig 3A), similar to a previous study[37]. Ectopic ORF2 expression was
comparable to the protein level reached by viral replication after EPO of WT HEV RNA into naïve
S10-3 cells (Fig. 3B). We further confirmed by WB that ORF3 but not ORF2 was expressed upon
EPO of ΔORF2 RNA into naïve cells (Fig. 3B). We then harvested infectious ΔORF2 particles
from ORF2-expressing S10-3 cell lysates to perform HEV infections.
Next, we aimed to study the replication dynamics after cell entry in HepG2/C3A cells in a
time-resolved manner. We synchronized the infection at equal MOI (30 genome equivalents
(GE)/cell) of both WT and ORF2-trans-complemented HEV particles by a binding step for 2 hours
(h) at 4 °C and internalization for 8 h at 37 °C, after which the inoculum was removed. We
analyzed viral replication and the antiviral response every 8 h between day 1 and day 3 post-
internalization and every day (i.e., every 24 h) thereafter by RT-qPCR. Replication of WT and the
ΔORF2 mutant peaked at 48±8 h post-internalization, followed by a decline (Fig. 3C).
Interestingly, while the HEV WT replication leveled off between 56 h and day 7 post-internalization

(2.0-fold decrease), ΔORF2 replication continued to decrease significantly by 8.5-fold between
those two time points. With the onset of viral replication, we observed an increase in *IFNL1* (Fig.
3D) and *ISG15* (Fig. 3E) expression, which was stronger in ΔORF2 infection relative to HEV RNA
(Fig. 3D-E; corresponding absolute HEV RNA genome copies and non-normalized *IFNL1* and
ISG15 expression in Suppl. fig. 5A-C). Following the decrease in HEV WT replication after 56 h,
IFNL1 (Fig. 3D) and *ISG15* (Fig. 3E) expression also decreased, relative to HEV RNA. In contrast,
the expression of antiviral response genes remained high upon ΔORF2 infection until day 7 post-
internalization.
Next, we aimed to validate our findings in a more physiologically relevant hepatocellular
system. To this end, we used human pluripotent stem cell-derived hepatocyte-like cells (HLCs)
which are permissive to HEV infection[13,33] (Fig. 3F). We infected HLCs at equal MOI (= 30 GE/cell)
with HEV WT and ΔORF2 virus particles produced by trans-complementation. Similar to our
results in infected HepG2/C3A cells, ΔORF2 showed reduced replication (Fig. 3G) while inducing
a stronger antiviral response compared to WT HEV RNA, both in absolute numbers as well as
normalized to HEV RNA (Fig. 3H and Suppl. fig. 5D).
Overall, our observations suggest that the antagonism mediated by ORF2 is critical for
establishing an equilibrium between viral replication and the antiviral response early in infection,
allowing HEV to replicate in different immunocompetent hepatocellular systems.
294
scRNA-seq analysis reveals ISG responses in HEV-infected cells and bystanders which are
dampened in the presence of ORF2
Expression of ISGs can be induced immediately downstream of viral recognition by PRRs
(reviewed in[9]). In addition, secreted IFN can lead to amplified ISG induction in an auto- and
paracrine manner. In HEV infection, it remains unclear whether IFN and ISG responses observed
in bulk analyses such as RT-qPCR are derived from infected cells, uninfected bystanders, or
possibly both. ISG responses from uninfected bystanders only could be induced by low levels of
IFN secretion from infected cells. Alternatively, viral RNA transfer to neighboring cells via
extracellular vesicles could lead to IFN and ISG induction in bystanders, similar to what has been
described for HCV and other viruses (reviewed in[38]).
To identify the cellular origin of the antiviral response in HEV infection and assess potential
differences in the response in the absence of the ORF2-mediated antagonism, we performed
single-cell RNA-sequencing (scRNA-seq) using microfluidics-based 3'-targeted 10x Genomics.
To this end, we infected HepG2/C3A cells with HEV WT and trans-complemented ΔORF2 virus
particles in a synchronized manner as described above and analyzed the cells at 56 h post-
internalization. This time point represents the antiviral response-mediated bottleneck we identified
to be essential for persistent replication in the presence of ORF2 (Fig. 3C-E). Since the ΔORF2
virus replicates to almost undetectable levels at later time points, we only analyzed HEV WT-
infected cells at day 7 post-infection.
We first performed a gene set enrichment analysis of the HEV WT- and ΔORF2-infected
samples compared to the uninfected sample at 56 h, using the hallmark gene sets of the Human
MSigDB Collections[39]. Interferon alpha and gamma responses were among the most significantly
upregulated gene sets, in both HEV WT and ΔORF2 infection, highlighting that the global
response was dominated by differential expression of ISGs (Suppl. fig. 6A-B). Enrichment of
oxidative phosphorylation and glycolysis gene sets indicated increased metabolic activity of the

infected samples. Analysis of HEV WT infection at day 7 showed that the enriched gene sets
were comparable over time and interferon-related responses were still prominently enriched
(Suppl. fig. 6C).
Next, we clustered the uninfected, HEV WT-, and ΔORF2-infected samples at 56 h based
on a list of approximately 400 ISGs published previously by Schoggins *et al.*[40] (Fig. 4A-D). This
revealed two major subpopulations in the infected samples (Fig. 4B-C), also at day 7 (Suppl. fig.
6D). We concluded that the cells located in the cluster overlapping with the uninfected sample
were non-responding cells, characterized by unchanged ISG expression. The cells located in the
second cluster showed differential ISG expression compared to the non-responding cells and
were thus classified as responders, from here on called the "active cluster" (Fig. 4E). In agreement
with the globally enhanced metabolic activity (Suppl. fig. 6A-B), this cluster appeared to be more
transcriptionally active, as indicated by an increased number of expressed genes (Fig. 4F) as well
as an overall increased number of transcripts (Suppl. fig. 6F). Within the WT- and ΔORF2-infected
samples, 28.1% and 17.2% of cells were identified as HEV RNA-positive, respectively (Suppl. fig.
6G). Interestingly, we observed that most infected cells of both WT- (Fig. 4G) and ΔORF2-infected
samples (Fig. 4H) were located in the active cluster, together with uninfected bystanders. We
therefore concluded that both actively infected cells and uninfected bystanders responded by
upregulated ISG expression. Although detectable by qRT-PCR (Fig. 3D), we could not detect type
III IFN gene expression in the infected samples via scRNA-seq, likely due to low respective
numbers of transcripts per cell. Therefore, we could not identify whether only infected cells or also
uninfected bystanders were the source of IFN. Nonetheless, we could demonstrate that the ISG
response was not fully suppressed in actively infected cells by the ORF2 antagonism.
Compared with the infected cells in the active cluster, the few infected cells located within
the inactive cluster had lower HEV RNA counts (Suppl. fig. 6H). These cells might be at an earlier
stage of infection due to a delayed onset of replication or due to secondary infection, which,
however, should be limited at this early time point of infection. Finally, not only uninfected
bystanders but also HEV-infected cells still showed enhanced ISG expression and localized to
the active cluster at day 7 post-HEV WT infection (Suppl. fig. 6E). This provided further evidence
that the establishment of an equilibrium between HEV replication and the antiviral response early
in infection is essential for persistent viral replication in an antiviral environment.
Next, we sought to examine the specific ISG signatures (i.e., the set of ISGs) induced in
actively infected cells and uninfected bystanders upon WT and ΔORF2 infection. First, we
observed a globally stronger ISG response in ΔORF2-infected cells and bystanders compared to
WT infection (Fig 4I), indicating overall increased IFN production and secretion, despite remaining
undetectable by scRNA-seq. We then compared the respective lists of differentially expressed
ISGs in the active clusters of WT and ΔORF2 infection and found that out of 243 differentially
expressed ISGs, 79.8% overlapped (Fig. 4J and Suppl. table 2). Based on this overlap and
previous literature on the upregulation of ISGs in HEV infection of different hepatocellular systems
and in chimpanzees[10,12,13,41], we selected a subset of ISGs for further comparison of infected cells
and bystanders. Among these common differentially expressed genes, both infected cells and
uninfected bystanders in the active cluster of the ΔORF2-infected sample showed a globally
stronger expression compared to the respective cell populations of the WT-infected sample (Fig.
4K).

As summarized in Fig. 4L, we identified both actively infected cells and uninfected
bystanders to be the cellular origins of ISG responses at early and later time points of HEV
infection. Early in infection, the ORF2 antagonism likely contributes to dampening the ISG
induction upon sensing in infected cells. In addition, our data suggests that the presence of ORF2
leads to decreased autocrine and paracrine IFN signaling, resulting in dampened ISG responses
in both infected cells and bystanders.
369
Discussion
HEV replication has been shown to persist despite a sustained cell-intrinsic antiviral
response. However, an in-depth understanding of the underlying mechanisms and the active
contribution of a viral antagonism is missing. In this study, we analyzed authentic HEV infection
with mutants lacking expression of individual viral proteins at single-cell resolution. We identified
a replication-limiting bottleneck imposed by the antiviral response where the presence of ORF2
and at least partially, its interaction with the adaptor molecule TBK1 determines whether viral
replication declines or equilibrates with the antiviral response.
378
Single-cell analysis uncovers the role of the multifunctional capsid protein for HEV persistence
amidst an antiviral environment
In the present study, we have shown that the capsid protein ORF2 is a critical viral
determinant of HEV replication efficiency. Investigation of the cell-intrinsic antiviral response at
the single-cell level revealed that both infected cells and uninfected bystanders are the source of
the ISG response in HEV infection. The ISG signatures in both cell types were globally stronger
in the absence of ORF2, but not strikingly different from WT infection (Fig. 4J-K). Thus, the ORF2-
mediated viral antagonism does not completely suppress antiviral responses within infected cells.
Instead, it reduces ISG and presumably IFN induction within infected cells and consequently
further ISG induction via autocrine and paracrine IFN signaling. Overall, antiviral responses are
dampened below a critical threshold, allowing viral replication and the antiviral response to reach
an equilibrium where they can coexist. We propose that this equilibrium is a prerequisite for
persistent HEV infection. Future studies are needed to determine whether IFN is produced only
by infected cells or also by uninfected bystanders, and to characterize the role of secreted ORF2
isoforms in influencing paracrine immune signaling.
HEV ORF2 appears to be a multifaceted protein with a wide range of functions affecting
different host pathways. In addition to its direct involvement in immune evasion downstream of
PRR recognition, we speculate that the capsid protein ORF2 may also exert additional, protective
functions. Here, we demonstrate that ORF2 is essential for maintaining the relative resistance of
HEV to type I and type III IFN treatment, even in the absence of a virus-induced antiviral response
(Fig. 2J). Importantly, we were able to exclude a direct inhibition of IFN signaling by ORF2 (Fig.
1E). Therefore, we concluded that the capsid protein additionally contributes to the shielding of
replicating RNA from the action of ISGs and potentially also from recognition by PRRs.
Most positive-sense RNA viruses replicate in the cytosol and restructure their host cell by
inducing host membrane-derived replication compartments (reviewed in[42]). While such replication
organelles have not yet been reported for HEV, several studies have suggested that different
compartments of the endolysosomal system or the ER-Golgi intermediate compartment (ERGIC)
could be involved[43-46]. ORF2 was frequently detected at these proposed replication sites.

Therefore, it may be involved in either remodeling or redirecting cellular structures to help create
the replication site, thereby potentially contributing to protection of viral RNA from the antiviral
effects of induced ISGs.
Despite the relatively low abundance of ORF1 and ORF3 proteins compared to ORF2
protein in authentic infection[47], we cannot exclude the possibility that these proteins have
additional potential to interfere with antiviral signaling and contribute to viral persistence. In this
study, we did not observe any striking effects of ORF3 on viral replication and antiviral responses.
Ectopically expressed ORF3 may interact with components of the antiviral response as reported
by others[26-28]. Furthermore, the processing of the ORF1 polyprotein in authentic infection remains
controversial (reviewed in[48]). Therefore, physiologically relevant contributions of the previously
reported antagonisms by ORF1 to the overall persistence phenotype remain to be confirmed.
418
The direct interaction of HEV ORF2 with TBK1 is critical for antagonizing replication-limiting
responses
There is no convincing evidence for a viral protease encoded by the HEV genome. Indeed,
unlike HCV[49] or HAV[50], HEV does not cleave MAVS[12], a key adaptor molecule in the induction of
cell-intrinsic antiviral responses. Instead, HEV appears to have developed other counteraction
strategies. In agreement with previous literature[23,24], we found that the HEV ORF2 protein
interacts directly with TBK1 to dampen IFN induction and the establishment of an infection-limiting
antiviral state. Here, we identified an early time point at which the ORF2-mediated antagonism is
critical for the progression of authentic HEV infection, facilitating a balance between viral
replication and the antiviral response within immunocompetent cells.
Lin and colleagues[24] previously suggested that the ARM of ORF2 is responsible for the
inhibition of TBK1-induced phosphorylation of IRF3. Indeed, we could show a reduced interaction
between ORF2 and TBK1 upon mutation of the ARM. However, this motif has also been identified
as the master regulator of the ORF2 isoforms[4]. Therefore, mutating the ARM could also indirectly
affect interference with the antiviral response, as enhanced secretion of glycosylated ORF2 along
the secretory pathway results in decreased cytosolic and nuclear availability of ORF2 to interact
with TBK1 and other unidentified factors. Furthermore, we found that ORF2 also antagonizes NF-
κB-dependent cytokine induction (Fig. 1D) and is hence likely to interfere with additional
components other than TBK1. This observation is in agreement with the previously published
interaction of ORF2 with beta-transducin repeat containing protein (βTRCP)[25], which resulted in
reduced proteasomal degradation of the inhibitory complex of NF-κB. Additionally, nuclear
translocation of ORF2 has been proposed to affect inflammatory cytokine induction[4]. Considering
the numerous suggested ORF2 antagonisms by us and others, we believe that viral persistence
is likely based on combinatorial interference of ORF2 with different pathways.
Due to the apparent multifunctionality of the ORF2 protein, individual functions are difficult
to untangle if specialized domains remain unknown. Indeed, our attempts to predict any putative
interaction sites between ORF2 and TBK1 with AlphaFold remained unsuccessful. The only
promising prediction was localized to the N-terminus of ORF2, however, this portion of the ORF2
protein appears to be intrinsically disordered[51]. While AlphaFold has demonstrated success in
identifying binding regions between other viral proteins and host factors[52], this particular
interaction highlights the challenges of accurately predicting interactions involving disordered
protein regions[53,54]. This complexity could stem from several factors: AlphaFold heavily relies on

co-evolutionary signals, which might be less pronounced in this virus-host interaction.
Furthermore, the ORF2-TBK1 interaction may rather be driven by physico-chemical properties
that are not explicitly considered in AlphaFold's predictions. Finally, the N-terminus of ORF2 may
adopt a specific structure upon interaction with TBK1, as is the case for many intrinsically
disordered proteins (reviewed in[55]). High-resolution structural studies of TBK1-ORF2 complexes
may therefore be required to further identify the interacting regions.
457
Innate immunity as a potential determinant of HEV chronicity and species tropism
Infections with different HEV genotypes result in distinct clinical outcomes. While HEV-1
infections are acute and self-limiting, HEV-3 infections can become chronic in
immunocompromised individuals. The determinants of these intergenotypic differences remain to
be elucidated, but innate and adaptive immune responses are likely critical parameters (reviewed
in[17]).
We have previously shown that infection with an HEV-1 isolate resulted in a stronger IFN
and ISG response than infection with an HEV-3 isolate. Interestingly, HEV-1 but not HEV-3
replication appeared to decrease thereafter[13]. Why HEV-1 induces a stronger response and
whether this contributes to the self-limiting course of infection observed in humans and in HEV-
1-infected chimpanzees[10] should be investigated in the future.
On the other hand, a moderate but prolonged cell-intrinsic IFN response induced by HEV-
3 could help to induce tolerance and dampen the effect of the stronger type I IFN response elicited
by professional immune cells such as plasmacytoid dendritic cells (pDCs). In agreement, we and
others have demonstrated a remarkable resistance of HEV replication to exogenous IFN once
replication is fully established[12,14-16]. Continuous activation of JAK/STAT signaling upon HEV
replication has been shown to render the cells refractory to exogenous IFN stimulation[12]. A recent
publication further suggested that ORF2 isoforms modulate detection of HEV-infected cells by
pDCs, indicating that this cell type might be contributing to HEV control *in vivo*[32]. Not only do pDCs
produce IFN and other cytokines to activate innate immune cells, but they also capture and
process antigens to initiate adaptive T cell responses. Consequently, a reduced T cell activation
could be beneficial for establishing HEV persistence *in vivo*.
Our results showed no difference in the potential of ORF2 from acute and chronic
genotypes to interfere with IRF3- and NF-κB-mediated signaling (Figure 1), which is consistent
with previous studies[23,24]. Therefore, the potency of ORF2 to antagonize these signaling pathways
does not appear to be a critical factor for the potential of HEV-3 to progress to chronic infection.
However, it could play a role in the genotype-dependent species tropism of HEV. While HEV-1 is
limited to primates and human infection, HEV-3 can infect a wide range of animals and is mainly
transmitted zoonotically to humans. Interaction of ORF2 with TBK1 could potentially represent a
prime example of the ongoing arms race between virus and host, driving the co-evolution of viral
evasion strategies and host defenses. Therefore, future research should investigate the
differences in the ability of HEV-1 and HEV-3 ORF2 to counteract TBK1 from different HEV-3 host
species, including pig or rabbit. Such studies will be crucial in identifying the determinants of HEV
species tropism.
Many open questions remain about the mechanisms of HEV persistence. In this study, we
have shown that the HEV capsid protein ORF2 mediates a viral antagonism affecting both antiviral
and inflammatory signaling. ORF2 is essential at a replication-limiting bottleneck early in infection

mediated by the antiviral response, which is decisive for establishment of persistent viral
replication. We have also laid the groundwork for future studies to focus on the role of
intergenotypic differences in the antiviral response and crosstalk with immune cells. This could
lead to a better understanding of the determinants of acute and chronic manifestations of HEV
infection, which will be valuable in identifying novel therapeutic regimens.
500
Materials and Methods
Standard cell culture
The human hepatoma cell lines HepG2/C3A (ATCC HB-8065), Huh7.5 (a kind gift from
Charles Rice, The Rockefeller University), S10-3 (a kind gift from Suzanne Emerson, NIH), and
derived S10-3/ORF2 as well as the human embryonic kidney cell line HEK293T (ATCC CRL-
3216) were cultured in Dulbecco's Modified Eagle Medium (DMEM, Gibco, high glucose,
GlutaMAX supplement) with 10% fetal bovine serum (FBS, Capricorn) and 1%
penicillin/streptomycin (pen/strep), referred to as complete DMEM (cDMEM). HepG2/C3A cells
were grown on collagen-coated cell culture plasticware. A549-derived cell lines were cultured in
DMEM (Gibco, high glucose) with 10% FBS (Pan Biotech), 1% pen/strep, and 1% non-essential
amino acids (NEAA). Cell lines with ectopic protein expression were produced by standard
lentiviral transduction, selected, and continuously cultured under respective antibiotic selection
pressure. Cells were maintained at 37 °C in 95% humidity and 5% CO_2 atmosphere. All cell lines
used in this study routinely tested negative for mycoplasma.
515
Stimulation of PRR-overexpressing A549-derived cell lines
$1x10^5$ A549-derived cell lines were seeded in 24-well plates. The next day, cells were
infected with Sendai virus (SeV, MOI 0.75; prepared from allantoic fluid of embryonated chicken
eggs, a kind gift from Rainer Zawatzky, German Cancer Research Center (DKFZ), Heidelberg,
Germany) for 4 h, Mengo-Zn virus[56] (MOI 1) for 24 h, or supernatant-fed with poly(I:C) (InvivoGen)
at 50 µg/mL for 24 h in DMEM containing 2% FBS, 1% NEAA, and 1% pen/strep. Stimulation with
TNF (Abcam) at 10 ng/mL or IFNβ (R&D Systems) at 200 IU/mL was performed in cDMEM plus
1% NEAA for 8 h. At respective time points, cells were lysed for RNA extraction with Monarch
Total RNA Miniprep Kit (New England Biolabs).
525
Generation of HEV ΔORF2, ΔORF3, and GNN mutants
Mutants were generated in the plasmid pBlueScript SK(+) encoding the HEV-3 Kernow-
C1/p6 sequence (pBSK-HEV-p6, GenBank accession number: JQ679013.1) by site-directed
mutagenesis. Mutations were introduced by overlap extension PCR with Phusion or Q5
polymerase (New England Biolabs) using the primers listed in Table 1, followed by restriction
enzyme digestion (New England Biolabs), ligation with T4 DNA Ligase (New England Biolabs),
transformation of JM109 competent cells (Promega), and validation by Sanger DNA sequencing.
533
In vitro *transcription and EPO of HEV WT, ΔORF2, ΔORF3, and GNN mutants*
Plasmids were linearized with MluI and RNA was *in vitro* transcribed using the
mMESSAGE mMACHINE T7 kit (Invitrogen). $4x10^6$ HepG2/C3A cells were electroporated with
10 µg of IVT HEV RNA in cytomix (120 mM KCl, 0.15 mM $CaCl_2$, 10 mM KPO_4, 25 mM HEPES,
2 mM EGTA, and 5 mM $MgCl_2$), supplemented with adenosine triphosphate (ATP) and glutathione

(GT). After resuspension in cDMEM, HepG2/C3A cells electroporated with HEV WT or mutants
were mixed 1:1 with mock-electroporated cells. $2x10^5$ cells/well were seeded on 24-well plates
for analysis by RT-qPCR and $1x10^5$ cells/well were seeded on 48-well plates for analysis by RNA-
FISH. One day post-EPO, all wells were washed twice with PBS to remove inoculum and cDMEM
was replaced. Cell lysates were harvested for RNA extraction using the Universal RNA kit
(Roboklon) according to manufacturer's instructions. On day 1, day 3, and day 5 post-EPO,
cDMEM was replaced and 6 µM TBK1 inhibitor BX795 (InvivoGen) or corresponding DMSO
vehicle control was added to the wells dedicated to cell lysis 48 h later. Samples for RNA-FISH
were fixed at respective time points with 4% paraformaldehyde (PFA).

EPO of Huh7.5 cells and IFN treatment

Huh7.5 cells were electroporated with HEV WT or ΔORF2 IVT RNA as described above.
For time-course analysis, electroporated Huh7.5 cells were resuspended in cDMEM, seeded in
24-well plates, and harvested for RT-qPCR analysis at respective time points. For IFN treatment,
cells were plated in a T75 cell culture flask. On day 3 post-EPO, cells were reseeded to 24-well
plates at approximately 80% confluency. One day later, cells were treated with 10,000 IU/mL
IFNα2A or 10 ng/mL IFNλ1. IFNs were replenished daily until cells were lysed for RNA extraction
with the Universal RNA kit (Roboklon) and RT-qPCR on day 7 post-EPO.

ΔORF2 virus production by trans-complementation

S10-3/ORF2 cells were electroporated with IVT ΔORF2 RNA and expanded on day 3 post-
EPO. Intracellular nHEV particles were harvested 7 days post-EPO from cell lysates by four
repeated freeze-thaw cycles and pelleted by ultracentrifugation at 28,000 rpm for 3 h through a
20% sucrose cushion. HEV ΔORF2 genome copies were determined by RT-qPCR following
TRIzol extraction from the resuspended virus prep.

Synchronized, time-resolved HEV infection

$6x10^4$ HepG2/C3A cells were seeded on collagen-coated 24-well plates. The next day,
cells were infected with equal genome copies of HEV WT or ΔORF2 virus (30 GE/cell). Virus was
bound at 4 °C for 2 h in Minimum Essential Medium (MEM, Gibco), supplemented with 10% FBS
and 1% pen/strep, followed by internalization at 37 °C for 8 h. After internalization, inoculum was
removed, cells were washed twice with PBS, and cDMEM was added. Samples for RT-qPCR
were lysed at respective time points with TRIzol for RNA extraction.

3'-targeted 10x Genomics and Illumina sequencing

$8x10^4$ HepG2/C3A cells were seeded on collagen-coated 24-well plates, and synchronized
HEV infection was performed as described above. At 56 h and day 7 post-internalization, cells
were trypsinized for 10 min, gently resuspended, and singularized by pipetting thrice through a
70 µm cell strainer. Two wells of each sample were combined. Single-cell suspensions were
washed once with 0.04% BSA in PBS and finally resuspended in 0.04% BSA/PBS. Cell
suspension was counted and inspected for cell death using trypan blue. Single-cell suspensions
were loaded onto the 10x Chromium controller according to manufacturer's instructions of the
Chromium Next GEM Single Cell 3' Kit v3.1 (10x Genomics) with a targeted cell recovery of 5,000.
Sequencing libraries were prepared according to the manufacturer's instructions. Briefly, GEMs

were generated, reverse transcription was performed, GEMs were cleaned up, and cDNA was
amplified and cleaned up with SPRIselect beads (Beckman Coulter). Quantification and quality
control were performed with Qubit (Thermo Fisher Scientific) and the 5200 Fragment Analyzer
System (Agilent Technologies). Fragmentation, end repair, and A-tailing were followed by
SPRIselect-based cleanup, adaptor ligation, sample indexing, and again, SPRIselect cleanup.
Quantification and quality control were performed with 5200 Fragment Analyzer System (Agilent
Technologies) and NEBNext Library Quant Kit for Illumina (New England Biolabs). Resulting
libraries were pooled and sequenced on Illumina NextSeq550 (high-output mode, paired-end, 150
cycles).
592
scRNA-seq data analysis
Fastq files were aligned and counted using the Cell Ranger 7.1.0 pipeline. As a human
genome reference we used GRCh38-3.0.0. For clustering and UMAP projections, we used only
a subset of genes based on a list of ISGs reported by Schoggins *et al.*[40]. Counts were scaled and
log1-normalized. The UMAP projections were calculated over the first 20 dimensions of the PCA,
setting the following parameters: min.dist=0.5 and n.neighbors=200. In order to identify clusters
associated with transcriptionally active/inactive states, we set a resolution of 0.1 in the
FindClusters function and then we inspected the expression of ISGs in the identified clusters. For
the DEA we used the FindMarkers function and filtered out genes expressed in less than 5% of
cells in the compared conditions and with p-adjusted values less than 0.05. For ISG expression
we used AUC values to score cells based on the Schoggins signature mentioned above
implemented with the AUCell 1.24.0 R package. For the comparison of the gene expression
across conditions we used a two-sided Wilcoxon rank sum test. Binarized ORF2 expression was
carried out by defining positive cells as those expressing at least one HEV RNA count. We used
the Seurat 5.0 R package for the entire analysis. The data is accessible under the GEO accession
number GSE288400.
609
610 *Statistical analysis*
611 Graphs were made and statistical analysis was performed using GraphPad Prism 8 or R
612 4.3.3 for the scRNA-seq analysis. Statistical tests and p-values are listed in the respective figure
613 legends.

Acknowledgments

The authors gratefully acknowledge Dr. Suzanne Emerson, Dr. Rainer Ulrich, Dr. Frauke Mücksch, and Dr. Henrik Kaessmann for sharing reagents and Andrew Freistaedter and Céline Schneider for excellent technical support. We acknowledge Dr. Vibor Laketa, head of the Infectious Diseases Imaging Platform (IDIP) at the University Hospital Heidelberg for expert support. The authors thank Dr. Marlène Dreux for critical reading of the manuscript. We thank the BIOI2 platform and the I2BC's IT support team for making AlphaFold 2.3 easily accessible at the I2BC.

Funding

A.M., S.S., C.R., C.H., M.B., and V.L.D.T were supported by grants from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Projektnummer – 272983813 SFB/TRR 179. V.L.D.T. was additionally supported by the Chica and Heinz Schaller Foundation, DFG grant DA1640/3-1, and TTU Hepatitis Project 05.823. J.H. was supported by a fellowship from the China Scholarship Council.

Author contributions

Conceptualization, A.M., and V.L.D.T.; Methodology, A.M., S.S., C.R, V.G.M, C.S., A.L.C., J.H., and T.T.; Investigation, A.M., S.S, C.R., and T.T.; Resources: D.K., M.K., X.W.; Software, A.M., S.S., C.R., A.L.C., T.T.; Data analysis, A.M., S.S., C.R., A.L.C., C.H., J.H.; Writing original draft, A.M., and V.L.D.T.; Final draft, A.M., and V.L.D.T.; Supervision, C.H., B.M., V.L.D.T.; Funding, V.L.D.T.

Declaration of interests

All authors declare no conflict of interest

References

1   Horvatits, T., Schulze Zur Wiesch, J., Lütgehetmann, M., Lohse, A. W. & Pischke, S. The
Clinical Perspective on Hepatitis E. *Viruses* **11**, doi:10.3390/v11070617 (2019).
2   Nimgaonkar, I., Ding, Q., Schwartz, R. E. & Ploss, A. Hepatitis E virus: advances and
challenges. *Nat Rev Gastroenterol Hepatol* **15**, 96-110, doi:10.1038/nrgastro.2017.150
(2018).
3   Das, A., Rivera-Serrano, E. E., Yin, X., Walker, C. M., Feng, Z. & Lemon, S. M. Cell entry
and release of quasi-enveloped human hepatitis viruses. *Nature reviews. Microbiology* **21**,
573-589, doi:10.1038/s41579-023-00889-z (2023).
4   Hervouet, K., Ferrie, M., Ankavay, M., Montpellier, C., Camuzet, C., Alexandre, V.,
Dembele, A., Lecoeur, C., Foe, A. T., Bouquet, P., Hot, D., Vausselin, T., Saliou, J. M.,
Salome-Desnoulez, S., Vandeputte, A., Marsollier, L., Brodin, P., Dreux, M., Rouille, Y.,
Dubuisson, J., Aliouat-Denis, C. M. & Cocquerel, L. An Arginine-Rich Motif in the ORF2
capsid protein regulates the hepatitis E virus lifecycle and interactions with the host cell.
PLoS Pathog **18**, e1010798, doi:10.1371/journal.ppat.1010798 (2022).
5   Yin, X., Ying, D., Lhomme, S., Tang, Z., Walker, C. M., Xia, N., Zheng, Z. & Feng, Z.
Origin, antigenicity, and function of a secreted form of ORF2 in hepatitis E virus infection.
Proc Natl Acad Sci U S A **115**, 4773-4778, doi:10.1073/pnas.1721345115 (2018).
6   Montpellier, C., Wychowski, C., Sayed, I. M., Meunier, J. C., Saliou, J. M., Ankavay, M.,
Bull, A., Pillez, A., Abravanel, F., Helle, F., Brochot, E., Drobecq, H., Farhat, R., Aliouat-
Denis, C. M., Haddad, J. G., Izopet, J., Meuleman, P., Goffard, A., Dubuisson, J. &
Cocquerel, L. Hepatitis E Virus Lifecycle and Identification of 3 Forms of the ORF2 Capsid
Protein. *Gastroenterology* **154**, 211-223 e218, doi:10.1053/j.gastro.2017.09.020 (2018).
7   Ankavay, M., Montpellier, C., Sayed, I. M., Saliou, J. M., Wychowski, C., Saas, L., Duvet,
S., Aliouat-Denis, C. M., Farhat, R., de Masson d'Autume, V., Meuleman, P., Dubuisson,
J. & Cocquerel, L. New insights into the ORF2 capsid protein, a key player of the hepatitis
E virus lifecycle. *Sci Rep* **9**, 6243, doi:10.1038/s41598-019-42737-2 (2019).
8   Wu, J. & Chen, Z. J. Innate immune sensing and signaling of cytosolic nucleic acids. *Annu
Rev Immunol* **32**, 461-488, doi:10.1146/annurev-immunol-032713-120156 (2014).
9   Schneider, W. M., Chevillotte, M. D. & Rice, C. M. Interferon-stimulated genes: a complex
web of host defenses. *Annu Rev Immunol* **32**, 513-545, doi:10.1146/annurev-immunol-
032713-120231 (2014).
10   Yu, C., Boon, D., McDonald, S. L., Myers, T. G., Tomioka, K., Nguyen, H., Engle, R. E.,
Govindarajan, S., Emerson, S. U. & Purcell, R. H. Pathogenesis of hepatitis E virus and
hepatitis C virus in chimpanzees: similarities and differences. *J Virol* **84**, 11264-11278,
doi:10.1128/JVI.01205-10 (2010).
11   Sooryanarain, H., Heffron, C. L. & Meng, X. J. The U-Rich Untranslated Region of the
Hepatitis E Virus Induces Differential Type I and Type III Interferon Responses in a Host
Cell-Dependent Manner. *mBio* **11**, doi:10.1128/mBio.03103-19 (2020).
12   Yin, X., Li, X., Ambardekar, C., Hu, Z., Lhomme, S. & Feng, Z. Hepatitis E virus persists
in the presence of a type III interferon response. *PLoS Pathog* **13**, e1006417,
doi:10.1371/journal.ppat.1006417 (2017).
13   Wu, X., Dao Thi, V. L., Liu, P., Takacs, C. N., Xiang, K., Andrus, L., Gouttenoire, J.,
Moradpour, D. & Rice, C. M. Pan-Genotype Hepatitis E Virus Replication in Stem Cell-
Derived Hepatocellular Systems. *Gastroenterology* **154**, 663-674 e667,
doi:10.1053/j.gastro.2017.10.041 (2018).
14   Dao Thi, V. L., Debing, Y., Wu, X., Rice, C. M., Neyts, J., Moradpour, D. & Gouttenoire, J.
Sofosbuvir Inhibits Hepatitis E Virus Replication In Vitro and Results in an Additive Effect
When Combined With Ribavirin. *Gastroenterology* **150**, 82-85 e84,
doi:10.1053/j.gastro.2015.09.011 (2016).

15   Todt, D., Francois, C., Anggakusuma, Behrendt, P., Engelmann, M., Knegendorf, L.,
Vieyres, G., Wedemeyer, H., Hartmann, R., Pietschmann, T., Duverlie, G. & Steinmann,
E. Antiviral Activities of Different Interferon Types and Subtypes against Hepatitis E Virus
Replication. *Antimicrob Agents Chemother* **60**, 2132-2139, doi:10.1128/AAC.02427-15
(2016).
16   Debing, Y., Emerson, S. U., Wang, Y., Pan, Q., Balzarini, J., Dallmeier, K. & Neyts, J.
Ribavirin inhibits in vitro hepatitis E virus replication through depletion of cellular GTP
pools and is moderately synergistic with alpha interferon. *Antimicrob Agents Chemother*
58, 267-273, doi:10.1128/aac.01795-13 (2014).
17   Bruggemann, Y., Klohn, M., Wedemeyer, H. & Steinmann, E. Hepatitis E virus: from innate
sensing to adaptive immune responses. *Nat Rev Gastroenterol Hepatol*,
doi:10.1038/s41575-024-00950-z (2024).
18   Nan, Y., Yu, Y., Ma, Z., Khattar, S. K., Fredericksen, B. & Zhang, Y. J. Hepatitis E virus
inhibits type I interferon induction by ORF1 products. *J Virol* **88**, 11924-11932,
doi:10.1128/jvi.01935-14 (2014).
19   Kang, S., Choi, C., Choi, I., Han, K. N., Rho, S. W., Choi, J., Kwon, J., Park, M. K., Kim,
S. J. & Myoung, J. Hepatitis E Virus Methyltransferase Inhibits Type I Interferon Induction
by Targeting RIG-I. *J Microbiol Biotechnol* **28**, 1554-1562, doi:10.4014/jmb.1808.08058
(2018).
20   Kim, E. & Myoung, J. Hepatitis E Virus Papain-Like Cysteine Protease Inhibits Type I
Interferon Induction by Down-Regulating Melanoma Differentiation-Associated Gene 5. *J*
Microbiol Biotechnol **28**, 1908-1915, doi:10.4014/jmb.1809.09028 (2018).
21   Myoung, J. & Min, K. Dose-Dependent Inhibition of Melanoma Differentiation-Associated
Gene 5-Mediated Activation of Type I Interferon Responses by Methyltransferase of
Hepatitis E Virus. *J Microbiol Biotechnol* **29**, 1137-1143, doi:10.4014/jmb.1905.05040
(2019).
22   Bagdassarian, E., Doceul, V., Pellerin, M., Demange, A., Meyer, L., Jouvenet, N. & Pavio,
N. The Amino-Terminal Region of Hepatitis E Virus ORF1 Containing a Methyltransferase
(Met) and a Papain-Like Cysteine Protease (PCP) Domain Counteracts Type I Interferon
Response. *Viruses* **10**, doi:10.3390/v10120726 (2018).
23   Hingane, S., Joshi, N., Surjit, M. & Ranjith-Kumar, C. T. Hepatitis E Virus ORF2 Inhibits
RIG-I Mediated Interferon Response. *Front Microbiol* **11**, 656,
doi:10.3389/fmicb.2020.00656 (2020).
24   Lin, S., Yang, Y., Nan, Y., Ma, Z., Yang, L. & Zhang, Y. J. The Capsid Protein of Hepatitis
E Virus Inhibits Interferon Induction via Its N-terminal Arginine-Rich Motif. *Viruses* **11**,
doi:10.3390/v11111050 (2019).
25   Surjit, M., Varshney, B. & Lal, S. K. The ORF2 glycoprotein of hepatitis E virus inhibits
cellular NF-κB activity by blocking ubiquitination mediated proteasomal degradation of
IκBα in human hepatoma cells. *BMC biochemistry* **13**, 7, doi:10.1186/1471-2091-13-7
(2012).
26   Dong, C., Zafrullah, M., Mixson-Hayden, T., Dai, X., Liang, J., Meng, J. & Kamili, S.
Suppression of interferon-alpha signaling by hepatitis E virus. *Hepatology* **55**, 1324-1332,
doi:10.1002/hep.25530 (2012).
27   He, M., Wang, M., Huang, Y., Peng, W., Zheng, Z., Xia, N., Xu, J. & Tian, D. The ORF3
Protein of Genotype 1 Hepatitis E Virus Suppresses TLR3-induced NF-kappaB Signaling
via TRADD and RIP1. *Sci Rep* **6**, 27597, doi:10.1038/srep27597 (2016).
28   Lei, Q., Li, L., Zhang, S., Li, T., Zhang, X., Ding, X. & Qin, B. HEV ORF3 downregulates
TLR7 to inhibit the generation of type I interferon via impairment of multiple signaling
pathways. *Sci Rep* **8**, 8585, doi:10.1038/s41598-018-26975-4 (2018).

29 Nan, Y., Ma, Z., Wang, R., Yu, Y., Kannan, H., Fredericksen, B. & Zhang, Y. J.
Enhancement of interferon induction by ORF3 product of hepatitis E virus. *J Virol* **88**, 8696-
8705, doi:10.1128/jvi.01228-14 (2014).
30 Plociennikowska, A., Frankish, J., Moraes, T., Del Prete, D., Kahnt, F., Acuna, C., Slezak,
M., Binder, M. & Bartenschlager, R. TLR3 activation by Zika virus stimulates inflammatory
cytokine production which dampens the antiviral response induced by RIG-I-like receptors.
J Virol **95**, doi:10.1128/JVI.01050-20 (2021).
31 Magalhães, V. G., Lukassen, S., Drechsler, M., Loske, J., Burkart, S. S., Wüst, S.,
Jacobsen, E. M., Röhmel, J., Mall, M. A., Debatin, K. M., Eils, R., Autenrieth, S., Janda,
A., Lehmann, I. & Binder, M. Immune-epithelial cell cross-talk enhances antiviral
responsiveness to SARS-CoV-2 in children. *EMBO reports* **24**, e57912,
doi:10.15252/embr.202357912 (2023).
32 Joshi, G., Décembre, E., Brocard, J., Montpellier, C., Ferrié, M., Allatif, O., Mehnert, A.-
K., Pons, J., Galiana, D., Dao Thi, V. L., Jouvenet, N., Cocquerel, L. & Dreux, M. Hepatitis
E Virus-induced antiviral response by plasmacytoid dendritic cells is modulated by the
ORF2 protein. 2024.2004.2012.589258, doi:10.1101/2024.04.12.589258 %J bioRxiv
(2024).
33 Dao Thi, V. L., Wu, X., Belote, R. L., Andreo, U., Takacs, C. N., Fernandez, J. P., Vale-
Silva, L. A., Prallet, S., Decker, C. C., Fu, R. M., Qu, B., Uryu, K., Molina, H., Saeed, M.,
Steinmann, E., Urban, S., Singaraja, R. R., Schneider, W. M., Simon, S. M. & Rice, C. M.
Stem cell-derived polarized hepatocytes. *Nat Commun* **11**, 1677, doi:10.1038/s41467-
020-15337-2 (2020).
34 Fu, R., Engels, Z., Weihs, J. A., Mürle, J., Klöhn, M., Todt, D., Hu, J., Steinmann, E.,
Böttler, T., Lozach, P.-Y., Lemon, S. M. & Thi, V. L. D. A high-content RNA-based imaging
assay reveals integrin beta 1 as a cofactor for cell entry of non-enveloped hepatitis E virus.
2023.2010.2027.564362, doi:10.1101/2023.10.27.564362 %J bioRxiv (2023).
35 Sumpter, R., Jr., Loo, Y. M., Foy, E., Li, K., Yoneyama, M., Fujita, T., Lemon, S. M. &
Gale, M., Jr. Regulating intracellular antiviral defense and permissiveness to hepatitis C
virus RNA replication through a cellular RNA helicase, RIG-I. *J Virol* **79**, 2689-2699,
doi:10.1128/JVI.79.5.2689-2699.2005 (2005).
36 Colasanti, O., Burm, R., Huang, H. E., Riedl, T., Traut, J., Gillich, N., Li, T. F., Corneillie,
L., Faure-Dupuy, S., Grünvogel, O., Heide, D., Lee, J. Y., Tran, C. S., Merle, U., Chironna,
M., Vondran, F. F. W., Esser-Nobis, K., Binder, M., Bartenschlager, R., Heikenwälder, M.,
Meuleman, P. & Lohmann, V. Comparison of HAV and HCV infections in vivo and in vitro
reveals distinct patterns of innate immune evasion and activation. *J Hepatol* **79**, 645-656,
doi:10.1016/j.jhep.2023.04.023 (2023).
37 Ding, Q., Heller, B., Capuccino, J. M., Song, B., Nimgaonkar, I., Hrebikova, G., Contreras,
J. E. & Ploss, A. Hepatitis E virus ORF3 is a functional ion channel required for release of
infectious particles. *Proc Natl Acad Sci U S A* **114**, 1147-1152,
doi:10.1073/pnas.1614955114 (2017).
38 Moulin, C., Crupi, M. J. F., Ilkow, C. S., Bell, J. C. & Boulton, S. Extracellular Vesicles and
Viruses: Two Intertwined Entities. *Int J Mol Sci* **24**, doi:10.3390/ijms24021036 (2023).
39 Liberzon, A., Birger, C., Thorvaldsdottir, H., Ghandi, M., Mesirov, J. P. & Tamayo, P. The
Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* **1**, 417-
425, doi:10.1016/j.cels.2015.12.004 (2015).
40 Schoggins, J. W. & Rice, C. M. Interferon-stimulated genes and their antiviral effector
functions. *Curr Opin Virol* **1**, 519-525, doi:10.1016/j.coviro.2011.10.008 (2011).
41 Todt, D., Friesland, M., Moeller, N., Praditya, D., Kinast, V., Brüggemann, Y., Knegendorf,
L., Burkard, T., Steinmann, J., Burm, R., Verhoye, L., Wahid, A., Meister, T. L.,
Engelmann, M., Pfankuche, V. M., Puff, C., Vondran, F. W. R., Baumgärtner, W.,
Meuleman, P., Behrendt, P. & Steinmann, E. Robust hepatitis E virus infection and

transcriptional response in human hepatocytes. *Proc Natl Acad Sci U S A* **117**, 1731-1741,
doi:10.1073/pnas.1912307117 (2020).

42 de Beijer, N. L., Snijder, E. J. & Bárcena, M. A Cool Look at Positive-Strand RNA Virus
Replication Organelles: New Insights from Cryo-Electron Microscopy. *Annual review of*
biochemistry **93**, 163-187, doi:10.1146/annurev-biochem-052521-115736 (2024).

43 Oechslin, N., Da Silva, N., Ankavay, M., Moradpour, D. & Gouttenoire, J. A genome-wide
CRISPR/Cas9 screen identifies a role for Rab5A and early endosomes in hepatitis E virus
replication. *Proc Natl Acad Sci U S A* **120**, e2307423120, doi:10.1073/pnas.2307423120
(2023).

44 Bentaleb, C., Hervouet, K., Montpellier, C., Camuzet, C., Ferrie, M., Burlaud-Gaillard, J.,
Bressanelli, S., Metzger, K., Werkmeister, E., Ankavay, M., Janampa, N. L., Marlet, J.,
Roux, J., Deffaud, C., Goffard, A., Rouille, Y., Dubuisson, J., Roingeard, P., Aliouat-Denis,
C. M. & Cocquerel, L. The endocytic recycling compartment serves as a viral factory for
hepatitis E virus. *Cell Mol Life Sci* **79**, 615, doi:10.1007/s00018-022-04646-y (2022).

45 Metzger, K., Bentaleb, C., Hervouet, K., Alexandre, V., Montpellier, C., Saliou, J. M.,
Ferrie, M., Camuzet, C., Rouille, Y., Lecoeur, C., Dubuisson, J., Cocquerel, L. & Aliouat-
Denis, C. M. Processing and Subcellular Localization of the Hepatitis E Virus Replicase:
Identification of Candidate Viral Factories. *Front Microbiol* **13**, 828636,
doi:10.3389/fmicb.2022.828636 (2022).

46 Glitscher, M., Spannaus, I. M., Behr, F., Murra, R. O., Woytinek, K., Bender, D. & Hildt, E.
The Protease Domain in HEV pORF1 Mediates the Replicase's Localization to
Multivesicular Bodies and Its Exosomal Release. *Cell Mol Gastroenterol Hepatol* **17**, 589-
605, doi:10.1016/j.jcmgh.2024.01.001 (2024).

47 Ankavay, M., Da Silva, N., Pollán, A., Oechslin, N., Dinkelborg, K., Behrendt, P.,
Moradpour, D. & Gouttenoire, J. Monitoring of Hepatitis E Virus Infection and Replication
by Functional Tagging of the ORF2 Protein. *JHEP Reports*,
doi:10.1016/j.jhepr.2024.101293.

48 Karpe, Y. A. Processing of the Hepatitis E virus ORF1 nonstructural polyprotein. *Frontiers*
in Virology **3**, doi:10.3389/fviro.2023.1327745 (2024).

49 Meylan, E., Curran, J., Hofmann, K., Moradpour, D., Binder, M., Bartenschlager, R. &
Tschopp, J. Cardif is an adaptor protein in the RIG-I antiviral pathway and is targeted by
hepatitis C virus. *Nature* **437**, 1167-1172, doi:10.1038/nature04193 (2005).

50 Yang, Y., Liang, Y., Qu, L., Chen, Z., Yi, M., Li, K. & Lemon, S. M. Disruption of innate
immunity due to mitochondrial targeting of a picornaviral protease precursor. *Proc Natl*
Acad Sci U S A **104**, 7253-7258, doi:10.1073/pnas.0611506104 (2007).

51 Shafat, Z., Ahmed, A., Parvez, M. K. & Parveen, S. Intrinsic disorder in the open reading
frame 2 of hepatitis E virus: a protein with multiple functions beyond viral capsid. *Journal,*
genetic engineering & biotechnology **21**, 33, doi:10.1186/s43141-023-00477-x (2023).

52 Ferrie, M., Alexandre, V., Montpellier, C., Bouquet, P., Tubiana, T., Meziere, L., Ankavay,
M., Bentaleb, C., Dubuisson, J., Bressanelli, S., Aliouat-Denis, C. M., Rouille, Y. &
Cocquerel, L. The AP-1 adaptor complex is essential for intracellular trafficking of the
ORF2 capsid protein and assembly of Hepatitis E virus. *Cell Mol Life Sci* **81**, 335,
doi:10.1007/s00018-024-05367-0 (2024).

53 Bret, H., Gao, J., Zea, D. J., Andreani, J. & Guerois, R. From interaction networks to
interfaces, scanning intrinsically disordered regions using AlphaFold2. *Nat Commun* **15**,
597, doi:10.1038/s41467-023-44288-7 (2024).

54 Versini, R., Sritharan, S., Aykac Fas, B., Tubiana, T., Aimeur, S. Z., Henri, J., Erard, M.,
Nüsse, O., Andreani, J., Baaden, M., Fuchs, P., Galochkina, T., Chatzigoulas, A., Cournia,
Z., Santuz, H., Sacquin-Mora, S. & Taly, A. A Perspective on the Prospective Use of AI in
Protein Structure Prediction. *Journal of chemical information and modeling* **64**, 26-41,
doi:10.1021/acs.jcim.3c01361 (2024).

55   Arai, M., Suetaka, S. & Ooka, K. Dynamics and interactions of intrinsically disordered
842   proteins. *Current opinion in structural biology* **84**, 102734, doi:10.1016/j.sbi.2023.102734
843   (2024).
844   56   Hato, S. V., Ricour, C., Schulte, B. M., Lanke, K. H., de Bruijni, M., Zoll, J., Melchers, W.
845   J., Michiels, T. & van Kuppeveld, F. J. The mengovirus leader protein blocks interferon-
846   alpha/beta gene transcription and inhibits activation of interferon regulatory factor 3. *Cell*
847   *Microbiol* **9**, 2921-2930, doi:10.1111/j.1462-5822.2007.01006.x (2007).
848

851
852
Figure 1: The HEV ORF2 protein antagonizes both antiviral and inflammatory signaling pathways
and interacts with TBK1.
(A) A549 cells harboring knockouts of the PRRs RIG-I and MDA5 and ectopically expressing a single PRR
(RIG-I, MDA5, or TLR3) together with either HEV-3 ORF2, HEV-1 ORF2, ORF3, or GFP were challenged
with either Sendai virus (SeV) for 4 h, (B) Mengo-Zn virus for 24 h, or (C) poly(I:C) supernatant feeding for
24 h, as indicated. (D) A549-derived cells ectopically expressing MDA5 were stimulated with TNF or (E)
IFNβ for 8 h. At respective time points, (A-C) *IFNB1*, (D) *TNFAIP3*, and (E) *IFIT1* expression were analyzed
by RT-qPCR, each relative to the housekeeping gene *GAPDH*. Data show mean ± SD of n = 3 independent
biological experiments. Statistical analysis was performed using two-way ANOVA. *: p < 0.05; **: p < 0.01;
***: p < 0.001; ****: p < 0.0001; ns, non-significant. A.U., arbitrary units. (F) HEK293T cells were co-
transfected with ORF2-HA or ORF2-2R/2A-HA and TBK1-V5 and lysed 24 h post-transfection. Anti-HA co-
IP and WB analysis for TBK1 (anti-V5 staining), ORF2 (anti-HA staining), and β-actin were performed.
Shown is a representative blot of n = 3 independent biological experiments.

866
867
Figure 2: The capsid protein ORF2, but not the ORF3 protein, is pivotal for efficient HEV replication
in immunocompetent cells.
(A) Schematic representation of the HEV genome and its replication intermediates. Nucleic acid positions
of GNN mutation and start codon mutations of ΔORF2 and ΔORF3 mutants are indicated. m7G, 7-

methylguanosin cap; A_n, poly(A)-tail. Created in BioRender. Dao Thi, V. (2025)
https://BioRender.com/j85i934 (B) HepG2/C3A cells were electroporated with *in vitro* transcribed (IVT) HEV
WT, ΔORF2, ΔORF3, or replication-incompetent GNN RNA and analyzed by RT-qPCR at indicated time
points post-EPO for HEV RNA, (C) *IFNL1* expression normalized over HEV RNA, and (D) *ISG15* expression
normalized over HEV RNA. Data show mean ± SD of n = 3 independent biological experiments. Statistical
analysis was performed using one-way ANOVA of each time point independently. *: $p < 0.05$; **: $p < 0.01$;
***: $p < 0.001$; ****: $p < 0.0001$; ns, non-significant. A.U., arbitrary units. (E-G) Electroporated HepG2/C3A
cells from (B) were additionally treated with 6 µM of the TBK1 inhibitor BX795 (TBKi) or respective DMSO
vehicle control 48 h prior to the indicated time point post-EPO. HEV RNA was quantified by RT-qPCR and
fold changes over DMSO treatment are shown above the respective graphs. Data show mean ± SD of n =
3 independent biological experiments. Statistical analysis was performed using unpaired two-tailed
Student's t-test of each time point independently. *: $p < 0.05$; **: $p < 0.01$; ns, non-significant. (H)
Electroporated HepG2/C3A cells from (B) were stained for HEV RNA using RNA-FISH at indicated days
post-EPO. Images show exemplary images of n = 3 independent biological experiments. Scale bar, 100
µm. (I) The percentages of electroporated HepG2/C3A cells positive for HEV RNA, stained by RNA-FISH
in (H), were quantified with ilastik and CellProfiler. Data show mean ± SEM of 3 independent biological
experiments and 5 images quantified per experiment. Statistical analysis was performed using unpaired
two-tailed Student's t-test of each condition independently. **: $p < 0.01$; ns, non-significant. (J) Huh7.5 cells
were electroporated with IVT HEV WT or ΔORF2 RNA and treated with 10 ng/mL IFNλ1 or 10,000 IU/mL
IFNα2A from day 4 to day 7 post-EPO. IFNs were replenished every 24 h. HEV RNA was quantified by RT-
qPCR at day 7 post-EPO and normalized over the respective untreated condition. Data show mean ± SD
of 4 biological repeats from 2 independent biological experiments. Statistical analysis was performed using
unpaired two-tailed Student's t-test of IFNλ1 and IFNα2A independently. **: $p < 0.01$; ns, non-significant.
895
896

897
898
Figure 3: A balance between HEV replication and the antiviral response at an early infection
bottleneck is essential for sustained viral replication.
(A) Schematic of ΔORF2 trans-complementation. S10-3 cells stably expressing HEV ORF2 were
electroporated with ΔORF2 HEV RNA to produce virus particles containing the ΔORF2 genome packaged
in capsid ORF2 protein expressed by the producer cell. Trans-complemented ΔORF2 virus was then used
to infect HepG2/C3A cells or stem cell-derived HLCs. Created in BioRender. Dao Thi, V. (2025)
https://BioRender.com/u54x394 (B) WB analysis of Mock, WT-, or ΔORF2-electroporated S10-3 or S10-
3/ORF2 cells for ORF2, ORF3, and β-actin protein expression. (C) Equal genome copies of HEV WT and
ΔORF2 virus particles (30 GE/cell) were bound on HepG2/C3A cells for 2 h at 4 °C prior to internalization
at 37 °C for 8 h, followed by removal of inoculum (= day 0). RT-qPCR was performed at indicated time
points post-internalization to determine fold viral replication over day 0, (D) *IFNL1* expression normalized
to HEV RNA, and (E) *ISG15* expression normalized to HEV RNA. Statistical analysis of fold changes over
time (C) or ΔORF2 over WT (D-E) are indicated above the respective time points in the corresponding color.
A.U., arbitrary units. Data show mean ± SD of n = 3 independent biological experiments. Statistical analysis
was performed using unpaired two-tailed Student's t-test of the respective conditions or days independently.
*: $p < 0.05$; **: $p < 0.01$; ns, non-significant. n/d, non-detectable; A.U., arbitrary units. (F) Exemplary image
of stem cell-derived hepatocyte-like cells (HLCs) infected with HEV WT, fixed, and stained for albumin and
ORF2 by immunofluorescence. Scale bar, 50 µm. (G) Stem cell-derived HLCs were infected with equal
genome copies of HEV WT and ΔORF2 virus particles (30 GE/cell) and analyzed by RT-qPCR for HEV

RNA and (H) *ISG15* expression normalized to HEV RNA over time. Data show mean ± SD of n = 4 biological
replicates of two independent HLC differentiations. Statistical analysis was performed using unpaired two-
tailed Student's t-test of each time point independently. *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$; ****: $p <$
0.0001; ns, non-significant.

922
923
Figure 4: scRNA-seq analysis reveals ISG responses in HEV-infected cells and bystanders which
are dampened in the presence of ORF2.
(A) UMAP projections of uninfected, (B) WT-infected, or (C) ΔORF2-infected HepG2/C3A cells harvested
at 56 h post-synchronized infection and processed by microfluidics-based 3'-targeted 10x Genomics.
UMAPs were generated by clustering of all cells based on a list of approximately 400 ISGs published
previously by Schoggins *et al.*[40] (D) Combined UMAP projections of uninfected, WT-, and ΔORF2-infected

samples, colored by condition. (E) Cells with upregulated ISG expression were highlighted in UMAP projections of uninfected, WT-, and ΔORF2-infected samples in black, representing the active cluster. (F) UMAP projection of uninfected, WT-, and ΔORF2-infected samples, highlighting the number of genes detected per cell in all samples. (G) UMAP projections of the WT-infected and (H) ΔORF2-infected samples with indicated HEV RNA counts. (I) Global changes in the ISG response were analyzed by comparison of normalized ISG expression across all samples (Uninfected, WT, and ΔORF2), based on a list of approximately 400 ISGs. WT- and ΔORF2-infected samples were split into actively infected cells, defined by detection of at least one HEV RNA copy per cell, and uninfected bystanders. Statistical analysis was performed using a two-sided Wilcoxon rank sum test. ****: $p < 0.0001$. (J) Differentially expressed ISGs in the active clusters of WT- and ΔORF2-infected samples were compared and plotted in a Venn diagram. Absolute numbers and percentages of shared and distinct differentially expressed ISGs are shown. (K) Out of the overlapping, differentially expressed ISGs between the active clusters of WT- and ΔORF2-infected samples, 30 ISGs were selected. Normalized expression and the percentage of cells were plotted for only the active clusters (as shown in E) of WT- and ΔORF2-infected samples, split by actively infected cells (at least one HEV RNA count per cell) and uninfected bystanders. (L) Working model depicting direct interference of ORF2 with TBK1 signaling downstream of PRR recognition of HEV RNA in WT-infected cells. Consequently, ISG induction downstream of PRR signaling as well as IFN secretion and thus, autocrine and paracrine ISG induction downstream of the IFN receptor is dampened in infected cells and bystanders, respectively. Additionally, ORF2 shields HEV RNA from the antiviral functions of ISGs. In ΔORF2-infected cells, the lack of the antagonistic functions of ORF2 results in stronger ISG induction downstream of PRR signaling as well as enhanced IFN secretion, resulting in stronger autocrine and paracrine ISG induction downstream of IFN signaling. Viral replication is consequently dampened as a result of the actions of antiviral effectors. Created in BioRender. Dao Thi, V. (2025) https://BioRender.com/h01q004