Psilocybin as a Treatment for Repetitive Mild Head Injury: Evidence from Neuroradiology and Molecular Biology

Eric K. Brengel[1,2], Bryce Axe[2], Ashwath Maheswari[2], Muhammad I. Abeer[3], Richard J. Ortiz[4], Taylor J. Woodward[5], Reagan Walhof[5], Rachel Utama[2], Courtney Sawada[2], Shreyas Balaji[2], Praveen P. Kulkarni[1,2], Heather B. Bradshaw[5], Michael A. Gitcho[3], Craig F. Ferris[1,2,6]

[1]Dept Psychology, Northeastern Univ, Boston, MA, USA; [2]Center for Translational Neuroimaging, Northeastern Univ, Boston, MA; [3]Dept Biological Sciences, Delaware State Univ, Dover, DE; [4]Dept Psychology, Northern Illinois Univ, DeKalb, IL; [5]Psychological & Brain Sciences, Program in Neuroscience, Indiana Univ, Bloomington, IN; [6]Dept Pharmaceutical Sciences, Northeastern Univ, Boston, MA

Address all Correspondence to: Craig Ferris, PhD, Director Center for Translational Neuroimaging, 360 Huntington Ave, Boston, Massachusetts, 02115; Email: c.ferris@northeastern.edu; Phone: 508 259 4908

Conflict of Interest: C.F.F. and P.P.K. have a partnership interest in Ekam Imaging Inc., a company that develops RF electronics and 3D MRI atlases for animal research. None of the other authors have a conflict of interest.

Funding: Ekam Imaging Inc. The Paul H. Boerger Fund of the Delaware Community Foundation. Fellowships for M.I.A.: NIH-NIGMS R25GM122722 and National Science Foundation 2023004

Authors' contributions: All contributed equally to the data generation, while E.K.B., B.A., A.M., R.J.O., R.U., and P.P.K. were responsible for MRI and behavior data analysis. H.B.B. and T.J.W. performed blood analysis for lipidomics. M.A.G. and M.I.A. ran studies on molecular biology. E.K.B., P.P.K., H.B.B., M.A.G., and C.F.F. contributed to the concept, experimental design, manuscript preparation, and interpretation.

Acknowledgements: We thank the National Institute on Drug Abuse Drug Supply Program and the Research Triangle Institute for providing the psilocybin used in these studies. We also thank the Center for Translational NeuroImaging and Ekam Imaging for financial support.

Abstract

Repetitive mild head injuries incurred while playing organized sports, during car accidents and falls, or in active military service are a major health problem. These head injuries induce cognitive, motor, and behavioral deficits that can last for months and even years with an increased risk of dementia, Parkinson's disease, and chronic traumatic encephalopathy. There is no approved medical treatment for these types of head injuries. To this end, we tested the healing effects of the psychedelic psilocybin, as it is known to reduce neuroinflammation and enhance neuroplasticity. Using a model of mild repetitive head injury in adult female rats, we provide unprecedented data that psilocybin can reduce vasogenic edema, restore normal vascular reactivity and functional connectivity, reduce phosphorylated tau buildup, enhance levels of brain-derived neurotrophic factor and its receptor TrkB, and modulate lipid signaling molecules.

Introduction

The Centers for Disease Control and Prevention report around 2.9 million people in the United States suffer from traumatic brain injury (TBI) every year, with 70-90% of these categorized as mild TBI[1, 2, 3]. In 2016, the total yearly healthcare expenses for nonfatal TBI exceeded $40.6 billion. This included $10.1 billion from private insurance, $22.5 billion from Medicare, and $8 billion from Medicaid[3]. There is an expanding literature on the behavioral and neurobiological consequences of mild head injuries that are incurred while playing organized sports, during car accidents and falls, or in active military service. Concussion following a single incident is difficult to detect and any associated cognitive and behavioral problems can resolve within hours of insult[4, 5]. However, a more pernicious, long-lasting condition arises when the brain is exposed to repetitive mild traumatic brain injury (rmTBI)[6, 7]. Repetitive head impacts and rmTBI induce cognitive, motor, and behavioral deficits, which are more severe and protracted, and can last for months and even years[8, 9] with an increased risk of dementia, Parkinson's disease[10-14], and chronic traumatic encephalopathy (CTE)[15, 16]. There are no approved treatments for repetitive head impacts, TBI, or rmTBI.

It has been suggested that the serotonergic hallucinogen psilocybin (PSI) could be used to treat brain injury given its known anti-inflammatory effects and its action as a promoter of neuroplasticity and cell growth[17]. A wide range of psychedelics, including PSI, are being evaluated for their potential therapeutic use in various psychiatric disorders[18] including substance abuse[19], severe depression[20, 21], and anxiety[22]. To date, there are no reports of PSI being used to treat any type of head injury. In a recent functional MRI study, we evaluated the dose-dependent effects of PSI on brain activity in fully awake rats. From these studies we determined a 3.0 mg/kg dose of PSI was most effective in stimulating positive blood-oxygen-level-dependent (BOLD) changes in brain activity. In this study, we tested the efficacy of this dose of PSI in a closed-skull momentum exchange model of rmTBI wherein rats were impacted once each day for three consecutive days[23-27]. To make the model more relevant to the human experience, rats were impacted while fully awake and during the dark phase of the light-dark cycle, when they are normally active. There are no mortalities with this model. Absent are any contusions or damage to the skull. Instead, the only gross radiological evidence of head

injury is swelling of the tissue above the skull at the impact site, i.e. a "bump on the

head." A single dose of PSI after each head impact significantly reduced the

neuroradiological and molecular measures associated with head injury.

Methods and Materials

Animals

Adult female *(N* = 24) Wistar rats were purchased from Charles River Laboratories

(Wilmington, MA, USA). Animals were housed in Plexiglas cages (two per cage) and

maintained in ambient temperature (22–24°C). Animals were maintained on a reverse

light-dark cycle with lights off at 09:00 and studied during the dark phase when they are

normally active. All experiments were conducted between 10:00 and 18:00 to avoid the

transitions between the light-dark cycles. Food and water were provided *ad libitum*. Rats

were ca. nine months of age when head-impacted and imaged. All animals were

acquired and cared for in accordance with the guidelines published in the NIH Guide for

the Care and Use of Laboratory Animals. All methods and procedures described below

were pre-approved by the Northeastern University Institutional Animal Care and Use

Committee under protocol numbers 23-0407R and 24-0517R. Northeastern University's

animal care and use program and housing facilities are fully accredited by AAALAC

International. The protocols used in this study followed the ARRIVE guidelines for

reporting *in vivo* experiments in animal research[28]. Animals were monitored daily over

the duration of the study for general health, including food and water consumption and

body weight. A 15% loss in body weight was set as a humane endpoint. Female rats

were divided into three experimental groups (*n* = 8; determined by *a priori* power

analysis): 1) healthy sham controls injected with saline vehicle but given no head impact

(SHAM-VEH), 2) head-impacted and injected with saline vehicle (rmTBI-VEH), and 3)

head-impacted and injected with psilocybin (rmTBI-PSI).

Overview

The study began with three consecutive days of mild head injury, psilocybin

treatment, and head-twitch response observation. On day three, within an hour of head

injury and treatment, blood plasma samples were collected for lipidomic analysis of

peripheral biomarkers of mild head injury. This was immediately followed by the first

magnetic resonance imaging (MRI) session. Cognitive and motor behaviors were tested

on days 4-10. Acclimation for awake neuroimaging occurred on days 15-19 leading up

to the second MRI session on day 22. The following day, brain tissue was collected for

proteomic analysis. See **Fig 1a** for a timeline of experimental procedures.

Repetitive Mild Head Injury

The momentum exchange model of mild head injury was developed by Viano et

al.[29] and further refined by Mychasiuk et al.[30] and Hightower et al.[31] to simulate the

dynamics of sport-related concussion in the preclinical setting with fully conscious

rodents. Each rat underwent this procedure once per day for three consecutive days.

Prior to impact on the first day, all rats were treated with 0.1 mg/kg extended-release

buprenorphine analgesic via subcutaneous injection to minimize pain for the duration of

the three-day repetitive injury period. Rats were lightly anesthetized via 1-2% isoflurane

inhalation to allow for setup in the momentum exchange apparatus. Under anesthesia,

rats were secured with a bite bar and strapped to a wheeled cradle sitting atop a

chassis to allow for linear and rotational acceleration upon impact. Once fully awake

(typically within 1-2 minutes), a pneumatic pressure system is used to reliably propel a

50 g impactor toward the head at 7.4 m/s for a kinetic energy input of 1.37 J, simulating

the rapid change in head velocity that occurs during concussions in the National

Football League[29, 30]. With the head angled downward into the impact plane, all injuries

were directed to the approximate area of Bregma. All rats demonstrated normal

ambulatory behavior within seconds of being returned to the home cage, and no

mortalities, seizures, loss of consciousness, skull fractures, contusions, or other

complications were observed. SHAM-VEH control rats underwent all of the above

procedures with the exception of head impact; when fully awake, these rats were

removed from the apparatus and the impactor was not launched.

Psilocybin administration and Head Twitch Response

Psilocybin was acquired through the National Institute on Drug Abuse (NIDA) and

distributed by the Research Triangle Institute. PSI was prepared in sterile saline (0.9%

NaCl) at a 3.0 mg/mL concentration for a dose of 3.0 mg/kg via intraperitoneal injection.

Rats were treated within 30 minutes of each head injury (once per day for three days). After each injection, rats were returned to the home cage and recorded from above for 10 minutes for quantification of the head twitch response (HTR), a behavioral indicator of psychoactive dosing. HTR and all subsequent behavioral assays were analyzed without outliers using one-way ANOVA in GraphPad Prism 10.0 software.

Open Field

On day 4, within 24 hours of the last head impact, rats were tested in the Open Field under dim red-light illumination. A detailed description of the Open Field test in rats appears in previous publications[32, 33]. Animals were placed in a lidless black box (60.9 x 69.2 x 70.5 cm) and allowed to explore for 5 minutes while recorded from above. Recordings were processed and data were measured using ANY-maze 7.00 software. In processing, the arena was divided into a peripheral zone (18 cm from the walls) and a central zone (20% of the arena). Agoraphobia (time spent in the center and number of entries to the center), thigmotaxis (time spent along the perimeter), average speed, and total distance traveled were measured and compared.

Novel Object Recognition

On day 5, the Novel Object Recognition task was used to evaluate episodic learning and memory[34-37]. Testing was conducted in the same lidless black box under dim red-light illumination, to which rats were acclimated during Open Field testing the previous day. Rats were first given a 5-minute habituation phase inside the empty box. Rats were then given a 5-minute familiarization phase to explore the box with two identical objects placed in diagonal corners. Lastly, rats returned for the 5-minute test phase, where one familiar object and one new object were presented in the same positions as the familiarization phase. Rats were returned to the home cage to rest for five minutes between each phase. Time spent with the novel and familiar objects during the test phase was measured using ANY-maze 7.00 software and the difference between the two for each subject was analyzed.

Beam Walk

Between days 8-9, rats were trained for motor behavior tasks. A detailed

description of the balance beam has been published[36, 38]. We used a tapered balance

beam equipped with sensors detecting foot faults across all three segments (wide,

middle, thin) to assess fine motor coordination. Animals were acclimated over two days

with three training trials per day. During training, they were placed in a goal box for 60

seconds, then on a start platform to cross the beam. Testing occurred over three trials

on day 10 with identical conditions. Foot faults and goal box latency were recorded and

averaged for each subject.

Rotarod

The Rotarod test is commonly used in Parkinson's disease models to assesses

equilibrium and motor function using a 4 cm diameter rotating rod that gradually

increases in speed[38]. Animals were acclimated over days 8-9, with three training trials

per day. During training, they were placed on the rod rotating at 5 rpm for 3 minutes,

and if they fell, they were immediately returned to the rod. Testing occurred over three

trials on day 10. The rod started at 1 rpm, accelerating at a rate of 0.2 rpm/s to a

maximum of 50 rpm over 245 seconds. The time before animals fell off was recorded

and averaged across trials.

Magnetic Resonance Imaging

Imaging was conducted at two timepoints: day 3 (1-2 hours post-injury and

treatment) and day 22 (three weeks post-injury and treatment). During the first session,

rats were anesthetized with 0.5-1% isoflurane for structural T2-weighted and diffusion

weighted imaging (DWI). The entire anesthetized imaging protocol lasted approximately

60 minutes (10 min setup, 6 min T2-weighted MRI, 44 min DWI). For a week before the

second session, rats were acclimated for awake imaging. The second session included

awake T2-weighted imaging, resting-state functional connectivity (rsFC), and functional

MRI with hypercapnic challenge (fMRI), and concluded with anesthetized DWI. The

entire imaging protocol lasted approximately 90 minutes, including 45 minutes awake

(10 min setup, 6 min T2-weighted MRI, 15 min rsFC, 15 min fMRI) and 45 minutes

anesthetized with 0.5-1% isoflurane (DWI).

Imaging sessions were conducted under dim red-light illumination using a Bruker

Biospec 7.0 T/20-cm USR horizontal magnet (Bruker, Billerica, MA, USA). Both imaging

sessions began with acquisition of a high-resolution T2-weighted anatomical data set

using a rapid acquisition, relaxation enhancement (RARE) pulse sequence to screen for

motion and to ensure there was no skull fracture or neuroanatomical injury as a result of

rmTBI. Due to the length of the imaging procedure, the study was conducted in six

staggered cohorts of four rats, with each experimental group randomly distributed

across all six cohorts. The animal setup and imaging protocol has been described in

detail in previous publications[33, 39-41].

Diffusion Weighted Imaging

DWI was acquired with a 3D spin-echo echo-planar-imaging (3D-EPI) pulse

sequence having the following parameters: TR/TE = 500/20 ms, eight EPI segments,

and 10 non-collinear gradient directions with a single b-value shell at 1000s/mm^2 and

one image with a B-value of 0 s/mm^2 (referred to as B0) as previously described[24, 26, 42].

Geometrical parameters were: 48 coronal slices, each 0.313 mm thick (brain volume)

and with in-plane resolution of 0.313 x 0.313 mm^2 (matrix size 96 x 96; FOV 30 mm^3).

The imaging protocol was repeated two times for signal averaging. For statistical

comparisons among rats, each brain volume was registered to the 3D MRI rat brain

atlas for generation of voxel- and region-based statistics. All image transformations and

statistical analyses were carried out using the in-house EVA software (Ekam Solutions

LLC, Boston, MA, USA). The average value for each region of interest was computed

using map files for indices of apparent diffusion coefficient (ADC) and fractional

anisotropy (FA). Statistical differences in measures of DWI between experimental

groups were determined using a nonparametric Kruskal Wallis multiple comparisons

test (critical value set at <0.05) followed by post hoc analyses using a Wilcoxon rank-

sum test for individual differences.

Acclimation for awake imaging

To mitigate the stress associated with head restraint, rats were acclimated to the

restraining and imaging protocol as previously described[24, 43]. Acclimation sessions

were conducted daily for five consecutive days, progressively increasing in duration up

to 45 minutes, the length of the awake imaging setup and acquisition protocol.

Respiration, heart rate, motor activity, and plasma corticosterone levels significantly

decrease over the course of the acclimation period[44]. This reduction in autonomic and

somatic signs of arousal and stress improves signal resolution and image quality.

Resting-State Functional Connectivity

A detailed description of the data acquisition, preprocessing, registration, and

analysis has been previously described[33, 41, 45]. Scans were collected using a spin-echo

triple-shot EPI sequence with the following parameters: matrix size = 96 x 96 x 20 (H x

W x D), TR/TE = 1000/15 ms, voxel size = 0.312 x 0.312 x 1.2mm, slice thickness = 1.2

mm, 200 repetitions, time of acquisition 15 min. Image preprocessing combined AFNI,

FSL, DRAMMS, and MATLAB software. After manual skull stripping, data underwent

motion correction, outlier removal, slice timing correction, and affine registration to the

rat atlas using DRAMMS. Data were then band-pass filtered (0.01-0.1 Hz), detrended,

and smoothed (FWHM = 0.8 mm). Nuisance regression removed motion parameters,

white matter, and CSF signals. Network analysis was performed in Gephi software

using undirected networks from absolute connectivity matrices. Degree centrality was

calculated as the sum of connections between each node and all other nodes.

Statistical analysis used GraphPad Prism 10.0, with Shapiro-Wilk tests determining

normality. Paired t-tests or Wilcoxon signed-rank tests (for non-normal data) compared

degree centrality between groups.

Functional MRI with Hypercapnic Challenge

Functional images were captured using a multi-slice Half-Fourier Acquisition

Single-Shot Turbo Spin Echo (HASTE) pulse sequence with an in-plane resolution of

312.5 μm². The scanning session for CO_2 challenge lasted 15 minutes with 10

acquisitions per minute. Each scanning session was continuous, starting with a 5-

minute baseline followed by 5 minutes of 5% CO_2 exposure and 5 minutes after

cessation of CO_2.

A detailed description of the data analysis for functional changes in BOLD signal

following CO_2 challenge is published[23, 46]. In brief, the fMRI data analysis consisted of

three main steps: pre-processing, processing, and post-processing. All these steps were

executed using SPM-12 (available at https://www.fil.ion.ucl.ac.uk/spm/) and in-house

MATLAB software. In the pre-processing stage, several operations were performed,

including co-registration, motion correction, smoothing, and detrending. In the

processing stage, all images were aligned and registered to a 3D Rat Brain Atlas©,

which included 173 segmented and annotated brain regions. This alignment was

performed using the GUI-based MIVA software developed by Ekam Solutions LLC

(Boston, MA, USA). All spatial transformations applied were compiled into a matrix for

each subject. Each transformed anatomical pixel location was tagged with its

corresponding brain area, resulting in fully segmented representations of individual

subjects for localization of functional imaging data to precise 3D volumes of interest.

Each scanning session consisted of 150 data acquisitions. The average signal

intensity in each voxel of the first 5 min of baseline (acquisitions 1–50) was compared to

5–10 min (acquisitions 51–100) of CO_2 exposure. We refer to the number of voxels in

each brain area that showed a significant increase in BOLD signal above threshold as

volume of activation. The mean volume of activation is the average of all rats for each

experimental condition for that brain area. Statistical t-tests were performed on each

voxel (~ 36,000 voxels in the whole brain) of each subject within their original coordinate

system. The baseline threshold was set at 1%. The t-test statistics used a 95%

confidence level ($p<0.05$), two-tailed distributions, and heteroscedastic variance

assumptions. As a result of the multiple t-test analyses performed, a false-positive

detection controlling mechanism was introduced using the formula $P(i)≤(i/V)(q/c(V))$

across 173 brain areas, where $q=0.2$ and $c(V)=1$.

Lipid Extraction and Partial Purification of plasma

Blood samples were collected 30 minutes after dosing on day 3 via the lateral tail

vein. Plasma was isolated via centrifugation and stored in 75µL samples at -80°C until

processed as previously described[47, 48]. In brief, methanolic extracts were partially

purified using C^{18} solid phase extraction columns (Agilent, Santa Clara, CA, USA). Final

elutions (i.e. fractions) of 65, 75, and 100 percent methanol were collected and stored at

-80°C until mass spectrometry (MS) analysis.

Lipidomics analysis

Methanolic elutions were analyzed as previously described[49] with the exception
that the API 7500 (Sciex, Framingham, MA 01701, USA) was used for analysis instead
of the API 3000. The API 7500 is coupled to a Shimadzu LC system LC-40DX3 (Kyoto,
Japan). Standard curves were generated by using purchased standards (Cayman
Chemical, Ann Arbor, MI, USA), and those made in-house were validated through NMR
and MS analysis as previously described[50]. Sciex Analyst peak matching software
(Sciex, Framingham, MA 01701, USA) was used to validate standard peaks and sample
peaks.

Statistical analyses for the plasma lipids were completed in IBM SPSS Statistics
29 (Chicago, IL, USA). One-way ANOVAs followed by Fisher's Least Significant
Analysis of individual endogenous lipids in plasma for each experimental group were
analyzed using Student's t-tests set to 2-tails and Type 2. Samples with an endogenous
lipid concentration outside of 2 standard deviations from the group mean were omitted
from statistics for that compound. Statistical significance for all tests was set at $p < 0.05$,
and trending significance at $0.05 < p < 0.10$. Descriptive and inferential statistics were
used to create heatmaps for visualizing changes in the concentration of each lipid
analyte for every condition as previously described[51].

Proteomics

On day 23, rats were deeply anesthetized via isoflurane inhalation for tissue
collection. Brains were rapidly extracted after decapitation, frozen, and stored at -80°C
until proteomic analysis. Rat brain samples of frontal area were isolated and solubilized
with Douce homogenizer followed by sonification in standard RIPA (50 mM Tris pH 8,
150 mM NaCl, 0.5% sodium deoxycholate, 0.1% SDS and 1% NP40) with
protease/phosphatase inhibitor as previously described[52]. The solubility fractionation
was modified and 200µg total protein from the RIPA-soluble fraction was centrifuged at
180,000g for 30 min, the supernatant was removed, and the RIPA-insoluble pellet was
washed in RIPA buffer and solubilized in 7M urea/2M thiourea before Western blot as
previously described[53]. RIPA samples of 20µg each were run on a 4-20% SDS-PAGE
and transferred to PVDF, blocked, probed, and visualized as previously described
(antibody table)[52].

An ANOVA with Tukey post hoc was performed comparing each of the three

groups (n = 8 biological replicates/group) reported as mean ± SD. All blots were

exposed under the same conditions. All raw uncropped data are available upon request.

Results

Behavior

Shown at the top of **Fig 1a** is a timeline for experimental procedures and data

collection. Shown in **Fig 1bc** are the results from the four behavioral assays: Open

Field, Beam Walk, Rotarod and Novel Object Recognition. The data collected from each

assay is organized into Motor Behavior (top row) and Cognitive and Emotional

Behaviors (lower row). The data are shown as dot plots (subjects) and bar graphs

(mean ± 95% confidence interval) and analyzed using a one-way ANOVA. HTR was

collected each of day 1-3 immediately following PSI or VEH treatment with the other

four assays conducted between day 4 and day 10. HTR was observed in all PSI-treated

rats exclusively, but only after the first dose and not after subsequent administration on

day 2 and day 3. This adaptation has been observed in rodents chronically exposed to

hallucinogenic 5-HT2A agonists[54-56]. No significant motor differences were observed

across groups for the Beam Walk (time to reach the goal box and number of slips),

Rotarod (latency to fall), or Open Field (total distance traveled and average speed).

Episodic learning and short-term memory were not significantly different between

groups when tested for object recognition. Measures of anxiety associated with fear of

open spaces (Agoraphobia) or attachment to protected surfaces (Thigmotaxis) were

also not significantly different between experimental groups in the Open Field.

Head Injury and Diffusion Weighted Imaging

Shown in **Fig 2a** are radiograms of each subject for the three experimental

groups characterizing the head injury associated with the site of impact near the

prefrontal cortex. The areas of impact are characterized by swelling and edema to the

soft tissue over the skull, visualized as white contrast in T2-weighted imaging (indicated

by arrows). Importantly, no skull fracture or contusion is observed. In three-week follow-

up scans, all impact-site edema is resolved. **Fig 2b** reports changes in ADC values

collected during DWI 1-2 hours post-injury and treatment on day 3 as a proxy for brain

edema. The box insert shows data from the whole brain (173 brain areas). There is a

pronounced increase in whole-brain ADC values (paired t-test, $p<0.0001$) comparing

SHAM-VEH and rmTBI-VEH rats with a mean difference of 0.06. When comparing all

three experimental groups (matched one-way ANOVA), both SHAM-VEH controls and

rmTBI-PSI treated rats were significantly less than rmTBI-VEH untreated rats

($p<0.0001$). However, it should be noted that PSI treatment did not reduce all of the

edema, as whole-brain measures were still significantly greater than SHAM-VEH. When

the 173 brain areas in the rat atlas are organized into brain regions, it is possible to

delineate those areas that are more or less sensitive to head injury and PSI treatment

as measured by changes in ADC. These data are shown in the bar graphs (mean ± SD)

and dot plots (subregions). For example, the hippocampus, comprised of nine brain

subregions, showed the same whole-brain profile of ADC values with SHAM-VEH

controls and rmTBI-PSI treated rats being less than rmTBI-VEH untreated rats, but with

rmTBI-PSI still greater than SHAM-VEH. This is also true for the somatosensory cortex

(SS cortex) and the prefrontal cortex. PSI treatment reduced edema to SHAM-VEH

levels in the basal ganglia, thalamus, cerebellum, and olfactory system. Specifically, an

inverse ADC:FA correlation is detected, indicating whole-brain vasogenic edema. On

three-week follow-up scans, all ADC and FA differences were resolved; however, lasting

functional changes were detected.

Vascular Reactivity to Hypercapnia

Fig 3 shows the change in vascular reactivity to carbon dioxide (CO_2) challenge

three weeks post-injury and treatment. The box insert shows the number of voxels

activated for positive BOLD for the whole brain for each of the experimental conditions.

A matched one-way ANOVA shows a significant difference between groups ($F_{(1.583, 267.5)}$

$= 90.44$, $p<0.0001$). rmTBI-VEH untreated rats present with significantly higher

($p<0.0001$) voxel numbers than SHAM-VEH controls or rmTBI-PSI treated rats (Tukey's

multiple comparison post hoc test), indicating vascular hyperreactivity. The bar graphs

(mean ± SD) and dot plots (subregions) show the regional differences in vascular

reactivity. In all cases, a matched one-way ANOVA showed a significant difference

between groups. Post hoc tests showed that rmTBI-VEH untreated rats were

significantly higher than SHAM-VEH controls and, in the case of the basal ganglia,

prefrontal cortex, and olfactory system, significantly greater than rmTBI-PSI treated rats.

It should be noted rmTBI-PSI treated rats still showed values significantly higher

(p<0.0001) than SHAM-VEH controls in all brain regions with the exception of the

prefrontal cortex, suggesting the treatment was not totally effective in reducing the

hyperreactivity to CO_2 challenge following head injury.

Resting-State Functional Connectivity

The bar graphs (mean ± SD) in the highlighted box of **Fig 4a** show the

connections to the whole brain for all 173 brain areas three weeks post-injury and

treatment. The global statistics using graph theory analysis for the SHAM-VEH controls

were Average Degree 14.737, Graph Density 0.087 and Average Path Length 2.541.

For rmTBI-VEH untreated rats, the statistics were Average Degree 9.088, Graph

Density 0.053, and Average Path Length 2.909. For rmTBI-PSI treated rats, the Average

Degree was 35.532, Graph Density 0.209, and Average Path Length 1.892. There was

a significant treatment effect using a matched one-way ANOVA [$F_{(1.352,229.8)}$ = 493.6,

p<0.0001)] followed by Tukey's multiple comparison post hoc test (**** p<0.0001). The

significant differences between each experimental group for different brain regions are

shown as dot and bar graphs. The dots represent the different subregions within each

brain region.

Fig 4b shows changes in connectivity to the three midbrain dopaminergic nuclei

(ventral tegmental area, substantia nigra compacta, and substantia nigra reticularis).

The panels above show the nodes (colored circles) and connections, or edges, (red

lines) from the three key dopaminergic nuclei (red circles). This functional connectivity is

displayed for all three experimental groups. The differences are dramatic, as rmTBI-

VEH untreated rats show an extreme loss of connectivity as compared to SHAM-VEH

controls and rmTBI-PSI treated rats. Psilocybin promotes hyperconnectivity, recruiting

nodes and connections to and within the thalamus (blue circles) and sensorimotor

cortices (yellow circles). The network connections between all nodes (black lines) are

shown in the lower panels. rmTBI-VEH untreated rats have very few connections, while

the network connectivity in SHAM-VEH controls, and especially rmTBI-PSI treated rats,

is very pronounced. These differences in degrees, or all connections associated with the

three dopaminergic nuclei, are shown in the top bar graph (mean ± SD). Both SHAM-

VEH controls (p<0.05) and rmTBI-PSI treated rats (p<0.01) were significantly greater

than rmTBI-VEH untreated rats (matched one-way ANOVA, $F_{(1.062, 2.124)}$ = 32.59, p =

0.025). The difference in degrees associated with the entire dopaminergic system is

shown in the bottom bar graph (median, min, and max). Again, SHAM-VEH controls

(p<0.001) and rmTBI-PSI treated rats (p<0.0001) were significantly greater than rmTBI-

VEH untreated rats (Kruskal-Wallis test for non-parametric data p<0.0001). rmTBI-PSI

treated rats were also greater than SHAM-VEH controls (p<0.0001), evidence of

hyperconnectivity.

Phosphorylated Proteins

The RIPA-soluble protein analysis normalized to tubulin shows a significant

increase in phosphorylated tau (PHF-1) when comparing SHAM-VEH controls to rmTBI-

VEH untreated rats. Interestingly, rmTBI-PSI treated rats show a reduction in

phosphorylated tau back to near SHAM-VEH levels (**Fig 5ab**). In addition, GFAP

(astroglia) and CD11b (microglia) levels are significantly increased in rmTBI-PSI treated

rats relative to SHAM-VEH controls. Changes in gliosis may initially be a protective

response to the injury, though it is not significantly different between SHAM-VEH

controls and rmTBI-VEH untreated rats[57] (**Fig 5ab**). When astrocyte activation is

reduced in an APP/PS1 mouse model, accelerated age-related plaque deposition

occurs which implies the need for functional astrocytes to respond to stress[58]. NAD(P)H

quinone dehydrogenase 1 (NQO1), induced by the NRF2 antioxidant response pathway

(review[59]), has been shown to protect against oxidative stress. rmTBI-PSI treated

animals show no significant differences from rmTBI-VEH untreated rats, though there is

an associated response increasing NQO1 compared to SHAM-VEH controls (**Fig 5ab**).

In rmTBI-PSI treated rats, BDNF shows a trend toward a significant increase over

SHAM-VEH controls (p = 0.057) and rmTBI-VEH untreated rats (p = 0.068). BDNF

exerts its effects via the high-affinity tyrosine kinase receptor TrkB[60]. Interestingly, TrkB

(NTRK2) shows a significant increase in rmTBI-PSI treated rats compared to SHAM-

VEH controls, which may be a protective measure increasing downstream factors (**Fig 5

ab**). Other RIPA-soluble proteins evaluated show no significant changes in expression.

These samples were from the frontal area, and the hippocampus and cerebellum may

show additional modulations of these pathways. The RIPA-insoluble (urea-soluble)

phosphorylated tau (PHF-1) also shows a significant increase in rmTBI-VEH untreated

rats but no difference between rmTBI-PSI treated rats and SHAM-VEH controls (**Fig 5**

cd). Though there is no significant decrease in phosphorylated tau comparing rmTBI-

VEH untreated rats to rmTBI-PSI treated rats, there is a distinct clustered trend in

reduced aggregated tau and no significant change when SHAM-VEH controls are

compared to rmTBI-PSI. This reduction in aggregation is consistent to what was

observed in the RIPA-soluble fraction. However, this significant increase in

phosphorylated tau post-injury may be at an early pre-tangle aggregation stage, and

with psilocybin treatment, animals show a significant reduction in PHF-1 to SHAM-VEH

levels. Interestingly, when examining changes in solubility, phosphorylated TDP-43

showed no significant change in aggregation. A soluble fractionation was performed

from the RIPA-soluble fraction (200μg total protein/centrifuge at 160,000g for 30 min,

pellet washed in RIPA and solubilized in 7M urea). Western blotting shows no

significant changes in aggregation most likely due to animal and sample preparation

variability. This variability in aggregation may be consistent with the experimental

dynamics of our rat rmTBI model consistent with the variability of the human disease.

Though phosphorylated TDP-43 in some rats show distinct changes in aggregation,

further evaluation of TDP-43, SNCA, amyloid, and other aggregated proteins will be

investigated.

Plasma signaling lipids

Two levels of analysis reveal important systemic effects of the intersection of our rmTBI

model and repeated PSI treatment. **Fig 6** shows the comparisons of SHAM-VEH versus

rmTBI-VEH where the heatmap illustrates the direction of change with rmTBI. Here we

see that, overall, there were only three changes in circulating levels of signaling lipids;

however, all changes were significant decreases after rmTBI (e.g. stearoyl leucine,

stearoyl taurine, and palmitoyl taurine). By contrast, animals treated with PSI who

underwent rmTBI had six times as many lipids change as a result of the treatment (3

versus 18 respectively), and all significant changes were increases. Key findings were

that PSI treatment caused significant increases in most free fatty acids measured and

caused reversal of levels of stearoyl taurine, suggesting that this signaling lipid may play

a key role in effects of rmTBI.

Discussion

Repetitive mild TBI is very common and can occur over the life span, affecting

adolescents playing organized sports, professional athletes, individuals in service, and

the elderly. rmTBI is a significant risk factor for dementia, Parkinson's, and

Alzheimer's[61]. Neuroinflammation, alterations in gray matter microarchitecture, impaired

cerebral blood flow, disruption in blood-brain barrier (BBB) permeability, and dysfunction

in clearance of unwanted phosphorylated proteins lie at the heart of these

neurodegenerative diseases[62, 63]. The repetitive mild head injury used in this study was

designed to reflect the human experience. Our neuroradiological findings recapitulate

much of the neuropathology observed and measured in the clinic using MRI. Below, we

discuss the "bench to bedside" relevance of these data and the significance of

psilocybin as a putative treatment for head injury and neurodegenerative disease.

Model of Repetitive Mild Head Injury

The guidelines from the Centers for Disease Control and Prevention, World

Health Organization, and American Congress of Rehabilitation Medicine for diagnosing

mild head injuries include self-reports of transient confusion, disorientation, impaired

consciousness, or dysfunction in memory around the time of the injury with no

neuroradiological evidence of structural damage to the brain[64, 65]. To that end, our lab

adopted a head injury model in rats originally developed by Viano and colleagues[29] and

further refined by Mychasiuk et al.[30], controlling for the axis of injury, rotational force,

and head acceleration in different directions. We have added to this model by

performing the mild head impacts while rats are fully awake without the confound of

anesthesia and during the active period of their circadian cycle. As was the case in

previous studies using this model[23, 66, 67], all rats showed normal ambulatory behavior

within seconds of being placed into their home cage after head impact. With this model

there are no mortalities and no evidence of skull damage or contusion as determined by

MRI. These are the "bump on head, ice pack" injuries that show up on radiograms as

edema in the skin at the site of impact as shown in **Fig 2a**. In this model, the

neurobiological effects of a single head impact resolve within 24 hours[25]. However, two

or more head impacts separated by 24 hours each have long-term consequences, as

noted in the clinical literature[68, 69]. In recent publications using this model of rmTBI, our

lab has reported a constellation of neurobiological and neurochemical changes in the

brains of male and female rats. The pathology includes vasogenic edema[25], altered

vascular reactivity[23, 70] and gray matter microstructure[26], disruption in blood-brain barrier

permeability[27], reduction in perivascular clearance[24], increases in microgliosis[24, 26],

altered astrocytic AQP4 expression and polarization[24], changes in brain functional

connectivity[26], and decreased brain-derived neurotrophic factor (BDNF) expression[66]. In

the present study we also report elevated levels of phosphorylated tau in the prefrontal

cortex of rats weeks after rmTBI.

Mild Repetitive Head Injury and Behavior

Measures of motor activity and coordination and learning and memory were not

significantly different across experimental groups. This is not unexpected with mild head

impacts in rodents. Anesthetized rats exposed to a single mild head impact[30, 71] or

anesthetized mice subjected to five mild head impacts spaced 24 hours apart[72] exhibit

minor balance and motor coordination deficits that resolve within a few days. Ren et al.

reported that anesthetized mice experiencing a single mild impact show no cognitive

changes but do display a decrease in motor performance on the rotarod, lasting up to

24 days[73]. In contrast, mice subjected to two mild, repetitive head impacts while fully

awake, evaluated through a series of neurobehavioral tests, show complete recovery

within hours[74]. Whether the early impacts in our study have long-term effects on

behavior as the subjects age remains uncertain. Recurrent mild head injury in humans

can lead to persistent post-concussive symptoms that overlap with other psychiatric

disorders like PTSD and major depression[75].

Vasogenic Edema

Edema plays a major role in the neuropathology of head injuries[76, 77]. Vasogenic

edema results from damage to the BBB, leading to the immediate movement of fluid

into the brain's extracellular space. An increase in apparent diffusion coefficient (ADC),

which measures water mobility, serves as an indicator of this volume change[77]. This

increase in ADC is typically associated with a decrease in fractional anisotropy (FA). In

cases of moderate-to-severe head injury, cytotoxic edema occurs, marked by cellular

swelling due to disrupted osmolarity regulation across the plasma membrane. This type

of edema generally shows a decrease in ADC and an increase in FA[78].

In a previous study we found that a single mild impact, which showed no

neuroradiological evidence of brain damage, led to a temporary increase in vasogenic

edema in the thalamus, basal ganglia, and cerebellum, as indicated by a rise in ADC[25].

This increase in extracellular fluid volume peaked at 6 hours and returned to baseline by

24 hours. In the present study using three mild head impacts, there was a global

increase in ADC values suggestive of vasogenic edema. This increase was shared by

several brain regions including the basal ganglia, hippocampus, thalamus, prefrontal

cortex, and somatosensory cortex. In each of these cases, treatment with PSI

significantly reduced the ADC values. In the basal ganglia and thalamus, these values

were reduced to levels measured in SHAM-VEH controls. How does PSI reduce ADC?

We would suggest two possible mechanisms: 1) Support BBB structural integrity via

endothelial tight junctions to reduce vasogenic edema, and/or 2) enhance astrocytic

activity helping to promote convection of excess extracellular fluid through aquaporin

channels lining the astrocytic endfeet. With respect to the first, there have been no

studies focused solely on PSI and capillary BBB integrity. However, there have been

several studies following changes in gene transcription in the cortex in rodents in

response to exposure to 5-HT2A psychedelics[79-82]. Most relevant to this study, Jefsen

and coworkers showed a significant increase in transcription of select genes in the

prefrontal cortex of rats in response to a single dose of PSI [80]. Several of these genes,

including Cebpb (CCAAT enhancer binding protein beta); Iκβ-α (NFKBIA), a key

regulator of NF-κB signaling; and Nr4a1 (Nur77) are all involved in vascular

inflammation and endothelial cell function.

In terms of enhanced fluid convention, we have hypothesized in a previous

study[83] that astrocytes localized around capillaries form a hydrolytic syncytium

connected by gap junctions[84]. Free water from vasogenic edema would move down

osmotic gradients, promoting swelling of the astrocytes and hydrostatic pressures

favoring convection toward AQP4 water channels at the endfeet surrounding capillaries.

PSI causes an elevation in GFAP (Glial Fibrillary Acidic Protein), a filament protein that

forms a dynamic intracellular scaffold that interacts with other binding proteins like

plectin (cytoskeletal crosslinker) and α-actinin (actin bundling protein) to form a

mechanically integrated system that affects astrocytic cell volume[85]. Changes in

astrocyte calcium fluctuations enhance phosphorylation and reorganization of the

filament structure that could cause mechanical forces, combined with passive osmotic

forces, to promote convection, lowering extracellular fluid and decreasing ADC values.

Vascular Reactivity

Fundamental to brain health is autoregulation of cerebral blood flow in the face of

fluctuations in systemic blood pressure. At the level of the neurovascular unit,

homeostasis is maintained by local changes in vascular reactivity and capillary blood

flow in response to the surrounding metabolic environment. A simple biomarker for

evaluating the health of cerebral blood vessels is a hypercapnic challenge[86] causing a

passive expansion of blood vessels, decreased resistance, and heightened blood flow.

Importantly, when exposed to higher levels of CO_2, there is no change in metabolic

oxygen consumption. Consequently, the alteration in the MRI signal caused by the

increased blood flow in the brain is directly linked to the change in the partial pressure

of CO_2. The change in BOLD signal in response to heightened CO_2 levels is a

straightforward and reliable technique for evaluating the cerebral vascular reactivity

(CVR) in functional imaging studies[87-89]. CO_2-induced changes in BOLD have been

used in the clinic to evaluate the health of cerebral vasculature in several neurological

disorders including Alzheimer's[90-92], stroke[93], multiple sclerosis[94] and TBI[95]. A recent

study by Liu and colleagues evaluated the reliability of hypercapnia-driven CVR as a

biomarker for cerebrovascular function and found it was suitable across different

scanning platforms and imaging sites for use in longitudinal studies and clinical trials[96].

There have been numerous studies using BOLD imaging and hypercapnic

challenge to evaluate cerebrovascular function following TBI. With moderate-to-severe

TBI there is a significant decrease in CVR in response to CO_2 challenge[95]. Subjects with

serious cerebral vascular injury following TBI show reduced global CVR compared to

healthy controls[97-99]. Reports of reduced CVR to CO_2 challenge are also true in

preclinical studies of TBI[100, 101]. However, there are reports of increased CO_2-induced

CVR following mild head injury, [102] particularly in those with sports-related concussions

accompanied by reduced CBF and cerebral metabolism[103, 104]. This would suggest

enhanced regulation of blood flow to affected areas with reduced metabolism[103, 104]. In

this and a previous study on repetitive mild head injury in female rats, [23] we found an

increase in CVR with CO_2 challenge. PSI treatment significantly reduced the increase in

whole-brain CVR as compared to rmTBI-VEH untreated rats but was still significantly

elevated above SHAM-VEH controls. The basal ganglia, prefrontal cortex, and olfactory

system were significantly reduced with PSI treatment as compared to rmTBI-VEH

untreated rats.

Functional Connectivity

This model of rmTBI presents with global functional hypoconnectivity. This result

is not unexpected given the many reports in clinical studies showing a similar decrease

in connectivity following head injury. Patients studied within the first seven days of a

mild TBI and presenting with post concussive syndrome (PCS) show a reduction in

functional connectivity in the sensorimotor and central executive networks as compared

to healthy volunteers[105]. Patients with mild TBI and PCS also show decreases in

connectivity to the thalamus[106] and a decrease in the symmetry of connectivity between

left and right thalamic nuclei[107]. The connectivity between the motor-striatal-thalamic

network is also reduced in mild TBI while the frontoparietal network is increased[108].

Pinky et al. reported a decrease in functional connectivity in the cerebellum and basal

ganglia in sports-related concussions in 12-18-year-olds[109]. Most recently, Fitzgerald

and colleagues ran longitudinal studies on young athletes competing in American

football, collecting functional connectivity prior to, during, and following the season[110].

Patterns of functional connectivity declined during the playing season but recovered after the season ended. They proposed that exposure to multiple head acceleration events may cause changes in brain neurobiology that are similar to concussion but in the absence of any symptomatology. The most robust finding in the present study was the dramatic hyperconnectivity that followed treatment with PSI. Not only did PSI prevent the hypoconnectivity of head injury, it exceeded the normal connectivity of sham controls. This global phenomenon was consistent across multiple brain regions.

Given the fact that head injury is one of the major risk factors for the development of Parkinson's disease, we focused on the connectivity of the midbrain dopaminergic system: the ventral tegmental area (VTA) and the substantia nigra (SN) compacta and reticularis. The efferent connections from these brain areas were significantly increased with PSI over rmTBI-VEH untreated rats with a shift toward greater connectivity to the thalamus and somatosensory cortex when compared to SHAM-VEH controls. The dopaminergic system, as defined by the connectivity within and between all these efferent targets from the VTA and SN, was significantly increased over SHAM-VEH and rmTBI-VEH untreated rats.

Psilocybin, BDNF, TrkB, and Phosphorylated Tau

In a recent study using our head injury model, we reported decreased BDNF levels in the hippocampus and around the midbrain dopaminergic nuclei[66]. Both male and female rats 9 months of age show a significant decrease in BDNF in the hippocampus with head injury while males also showed a decrease in BDNF in the substantia nigra. BDNF is the most prevalent neurotrophin in the brain, crucial for the survival, differentiation, synaptic plasticity, and axonal growth of both peripheral and central neurons during adulthood[111]. BDNF exerts its effects via the high-affinity tyrosine kinase receptor TrkB [60]. After spinal cord injury or TBI there is an increase in TrkB mRNA expression at the site of injury[112]. BDNF has been recognized for its significant involvement in cellular processes related to recovery after TBI, such as promoting neuronal survival, axonal growth, and the formation of new synapses[113]. Given its crucial role, BDNF has been extensively studied in experimental TBI models[114]. In a

recent study, Moliner et al. reported PSI can directly bind to TrkB receptor with a very

high affinity[115]. Further, Zhao et al recently demonstrated the ability of PSI to restore

decreases in prefrontal cortex BDNF expression in a rodent model of major depressive

disorder[116]. Here, we show that PSI elevated BDNF and TrkB proteins above SHAM-

VEH control and rmTBI-VEH untreated rats in the prefrontal cortex. These data provide

a possible mechanism of action for the healing effects of PSI.

Interestingly, the BDNF/TRK2 pathway is altered in neurodegeneration

associated with tauopathies[117], a group of neurodegenerative diseases characterized by

the hyperphosphorylation of tau[118]. Tau proteins bind to and stabilize microtubules in

neurons to help maintain the structure and function of axons[119]. When tau becomes

hyperphosphorylated, it can aggregate into neurofibrillary tangles (NFTs) inside

neurons, disrupting cell function[120]. The accumulation of NFTs and the subsequent loss

of neuronal function are key features in Alzheimer's disease, frontotemporal dementia,

and progressive supranuclear palsy[121-123]. Here, we provide evidence that PSI can

reduce phosphorylated tau following head injury, which may ultimately reduce the risk

for neurodegenerative diseases.

Plasma lipids as biomarkers of CNS effects

Recently we showed that injections of the endogenous signaling lipid, palmitoyl

ethanolamine (PEA) causes significant changes in CNS connectivity and behavior as

well as plasma and CNS signaling lipids[33]. These data demonstrate that the presence of

signaling lipids in plasma have a direct effect on CNS activity. Data here show that

rmTBI alone has a modest but significant effect on signaling lipids by significantly

reducing some of the lipids. By contrast, PSI treatment, which significantly improved

rmTBI outcomes at multiple levels, drove significant increases in a wide range of

circulating signaling lipids. In both cases, these signaling lipids may be key biomarkers

for both injury and recovery.

The dramatic hyperconnectivity we observed in this study most likely reflects the

hyperplastic effects of PSI to enhance synaptogenesis and dendritic growth. Does the

concomitant increase in the blood level of small lipids reflect these morphological

changes occurring in neuronal plasma membranes? Indeed, a majority of the brain

lipidome composition in the prefrontal cortex of humans, chimpanzees, and macaque monkeys occur prior to adulthood, with significant alterations in lipid concentrations during early development[124]. Lipidomic studies in early childhood have shown marked changes in circulating lipid profiles from birth through early childhood, with specific lipid species showing age-dependent changes[125]. These findings collectively indicate that both postnatal and adolescent periods are characterized by enhanced lipidomic activity in plasma and brain, reflecting critical phases of neurodevelopment and metabolic regulation.

Summary

Repetitive mild head injury affects diverse populations across age groups and is a significant risk factor for neurodegenerative diseases. Our rat model replicates clinical mild TBI by delivering impacts to awake animals during their active phase, producing no structural damage but causing measurable neuropathology. While we observed no significant behavioral deficits—consistent with rapid recovery in mild TBI—our model revealed several key pathological changes.

Diffusion imaging showed widespread increases in ADC values, suggestive of transient vasogenic edema. PSI treatment reduced edema, potentially through two mechanisms: 1) strengthening blood-brain barrier integrity and 2) enhancing astrocytic fluid clearance through GFAP-mediated changes in cell volume regulation.

CO_2 challenge revealed lasting vascular hyperreactivity three weeks after injury, contrasting with the hyporeactivity seen in severe TBI. This suggests compensatory regulation in mild injury. PSI partially normalized this response, particularly in the basal ganglia, prefrontal cortex, and olfactory regions.

Functional connectivity analysis demonstrated global hypoconnectivity post-injury, matching clinical observations in head-injured patients. PSI treatment induced dramatic hyperconnectivity, notably in dopaminergic pathways to thalamic and somatosensory regions. This finding is particularly relevant given the role of rmTBI in risk for Parkinson's disease.

Protein analysis showed an increase in BDNF and TrkB, a possible mechanism of PSI action. BDNF is crucial for neuronal survival and plasticity and acts through TrkB receptors, which increase after brain injury. Recent research shows PSI can directly

bind TrkB with high affinity. The ability of PSI to reduce tau phosphorylation suggests

potential therapeutic applications beyond rmTBI, possibly extending to other tau-related

neurodegenerative disorders.

Modulations in plasma lipids with both rmTBI alone and in conjunction with PSI

treatment provide a novel set of biomarkers to exploit for our understanding of how

systemic changes in lipid signaling are associated with both long-term damage of rmTBI

and the therapeutic effects of psilocybin.

This translational model successfully bridges bench-to-bedside by replicating

clinical observations and identifies PSI as a promising therapeutic agent for repetitive

mild head injury and its neurodegenerative consequences.

References

1.	Peterson, A., Xu, L., Daugherty, J. & Breiding, M. Surveillance Report of Traumatic Brain Injury-Related Emergency Department Visits, Hospitalizations, and Deaths. Centers for Disease Control and Prevention. (2014).

2.	Maas, A.I.R.*, et al.* Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. *Lancet Neurol* **16**, 987-1048 (2017).

3.	Centers for Disease Control and Prevention. Economics of Injury and Violence Prevention. (CDC, Atlanta, Georgia, 2024).

4.	Lovell, M.R.*, et al.* Recovery from mild concussion in high school athletes. *Journal of neurosurgery* **98**, 296-301 (2003).

5.	Pellman, E.J., Lovell, M.R., Viano, D.C. & Casson, I.R. Concussion in professional football: recovery of NFL and high school athletes assessed by computerized neuropsychological testing--Part 12. *Neurosurgery* **58**, 263-274; discussion 263-274 (2006).

6.	Guskiewicz, K.M.*, et al.* Cumulative effects associated with recurrent concussion in collegiate football players: the NCAA Concussion Study. *Jama* **290**, 2549-2555 (2003).

7.	Marar, M., McIlvain, N.M., Fields, S.K. & Comstock, R.D. Epidemiology of concussions among United States high school athletes in 20 sports. *Am J Sports Med* **40**, 747-755 (2012).

8.	De Beaumont, L.*, et al.* Brain function decline in healthy retired athletes who sustained their last sports concussion in early adulthood. *Brain : a journal of neurology* **132**, 695-708 (2009).

9.	Gardner, R.C. & Yaffe, K. Epidemiology of mild traumatic brain injury and neurodegenerative disease. *Mol Cell Neurosci* **66**, 75-80 (2015).

10.	Crane, P.K.*, et al.* Association of Traumatic Brain Injury With Late-Life Neurodegenerative Conditions and Neuropathologic Findings. *JAMA Neurol* **73**, 1062-1069 (2016).

11.	Jafari, S., Etminan, M., Aminzadeh, F. & Samii, A. Head injury and risk of Parkinson disease: a systematic review and meta-analysis. *Movement disorders : official journal of the Movement Disorder Society* **28**, 1222-1229 (2013).

12.   Jenkins, P.O.*, et al.* Dopaminergic abnormalities following traumatic brain injury.
Brain : a journal of neurology (2018).

13.   Taylor, K.M.*, et al.* Head injury at early ages is associated with risk of Parkinson's
disease. *Parkinsonism & related disorders* **23**, 57-61 (2016).

14.   Gardner, R.C.*, et al.* Mild TBI and risk of Parkinson disease: A Chronic Effects of
Neurotrauma Consortium Study. *Neurology* **90**, e1771-e1779 (2018).

15.   McKee, A.C.*, et al.* Chronic traumatic encephalopathy in athletes: progressive
tauopathy after repetitive head injury. *Journal of neuropathology and experimental*
neurology **68**, 709-735 (2009).

16.   Konrad, C.*, et al.* Long-term cognitive and emotional consequences of mild
traumatic brain injury. *Psychological medicine* **41**, 1197-1211 (2011).

17.   Allen, J., Dames, S.S., Foldi, C.J. & Shultz, S.R. Psychedelics for acquired brain
injury: a review of molecular mechanisms and therapeutic potential. *Mol Psychiatry* **29**,
671-685 (2024).

18.   Banushi, B. & Polito, V. A Comprehensive Review of the Current Status of the
Cellular Neurobiology of Psychedelics. *Biology (Basel)* **12** (2023).

19.   de Veen, B.T., Schellekens, A.F., Verheij, M.M. & Homberg, J.R. Psilocybin for
treating substance use disorders? *Expert Rev Neurother* **17**, 203-212 (2017).

20.   Vollenweider, F.X. & Preller, K.H. Psychedelic drugs: neurobiology and potential
for treatment of psychiatric disorders. *Nat Rev Neurosci* **21**, 611-624 (2020).

21.   Goodwin, G.M.*, et al.* Single-Dose Psilocybin for a Treatment-Resistant Episode
of Major Depression. *N Engl J Med* **387**, 1637-1648 (2022).

22.   Griffiths, R.R.*, et al.* Psilocybin produces substantial and sustained decreases in
depression and anxiety in patients with life-threatening cancer: A randomized double-
blind trial. *J Psychopharmacol* **30**, 1181-1197 (2016).

23.   Bens, N., Kulkarni, P. & Ferris, C.F. Changes in cerebral vascular reactivity
following mild repetitive head injury in awake rats: modeling the human experience. *Exp*
Brain Res **242**, 2433-2442 (2024).

24.   Cai, X.*, et al.* Mild repetitive head impacts alter perivascular flow in the midbrain
dopaminergic system in awake rats. *Brain Commun* **3**, fcab265 (2021).

25.   Kulkarni, P., *et al.* Evidence of early vasogenic edema following minor head
impact that can be reduced with a vasopressin V1a receptor antagonist. *Brain Res Bull*
165, 218-227 (2020).
26.   Kulkarni, P., *et al.* Neuroradiological Changes Following Single or Repetitive Mild
TBI. *Front Syst Neurosci* **13**, 34 (2019).
27.   Leaston, J., *et al.* Quantitative Imaging of Blood-Brain Barrier Permeability
Following Repetitive Mild Head Impacts. *Front Neurol* **12**, 729464 (2021).
28.   Kilkenny, C., Browne, W.J., Cuthill, I.C., Emerson, M. & Altman, D.G. Improving
bioscience research reporting: the ARRIVE guidelines for reporting animal research.
PLoS Biol **8**, e1000412 (2010).
29.   Viano, D.C., Hamberger, A., Bolouri, H. & Saljo, A. Concussion in professional
football: animal model of brain injury--part 15. *Neurosurgery* **64**, 1162-1173; discussion
1173 (2009).
30.   Mychasiuk, R., Hehar, H., Candy, S., Ma, I. & Esser, M.J. The direction of the
acceleration and rotational forces associated with mild traumatic brain injury in rodents
effect behavioural and molecular outcomes. *J Neurosci Methods* **257**, 168-178 (2016).
31.   Hightower R, Brengel E, Prom S, Kulkarni P & 2024;5(3):, F.C. Testing the
Efficacy of Minocycline Treatment in an Awake, Female Rat Model of Repetitive Mild
Head Injury. *J Exp Neurol.* **5(3)**, 145-161 (2024).
32.   Morrison, T.R., *et al.* Treating head injury using a novel vasopressin 1a receptor
antagonist. *Neurosci Lett* **714**, 134565 (2020).
33.   Balaji, S., *et al.* Palmitoylethanolamide causes dose-dependent changes in brain
function and the lipidome. *Front Neurosci* **18**, 1506352 (2024).
34.   Coleman, J.R., *et al.* Changes in brain structure and function following chronic
exposure to inhaled vaporised cannabis during periadolescence in female and male
mice: A multimodal MRI study. *Addict Biol* **27**, e13169 (2022).
35.   Demaree, J.L., *et al.* Exposure to methylphenidate during peri-adolescence
decouples the prefrontal cortex: a multimodal MRI study. *Am J Transl Res* **13**, 8480-
8495 (2021).

36.   Ferris, C.F.*, et al.* Life without a brain: Neuroradiological and behavioral evidence

of neuroplasticity necessary to sustain brain function in the face of severe

hydrocephalus. *Sci Rep* **9**, 16479 (2019).

37.   Harris-Blum, L.*, et al.* Developmental changes in brain structure and function

following exposure to oral LSD during adolescence. *Sci Rep* **14**, 18632 (2024).

38.   Ferris, C.F.*, et al.* Evidence of Neurobiological Changes in the Presymptomatic

PINK1 Knockout Rat. *Journal of Parkinson's disease* **8**, 281-301 (2018).

39.   Alkislar, I.*, et al.* Inhaled Cannabis Suppresses Chemotherapy-Induced

Neuropathic Nociception by Decoupling the Raphe Nucleus: A Functional Imaging Study

in Rats. *Biol Psychiatry Cogn Neurosci Neuroimaging* **6**, 479-489 (2021).

40.   Brems, B.M.*, et al.* Dose-dependent effects of GAT107, a novel allosteric agonist-

positive allosteric modulator (ago-PAM) for the α7 nicotinic cholinergic receptor: a BOLD

phMRI and connectivity study on awake rats. *Front Neurosci* **17**, 1196786 (2023).

41.   Ghaw, A.*, et al.* Dose-dependent LSD effects on cortical/thalamic and cerebellar

activity: brain oxygen level-dependent fMRI study in awake rats. *Brain Commun* **6**,

fcae194 (2024).

42.   Ferris, C.F.*, et al.* Alterations in brain neurocircuitry following treatment with the

chemotherapeutic agent paclitaxel in rats. *Neurobiology of pain (Cambridge, Mass.)* **6**,

100034 (2019).

43.   Cai, X.*, et al.* Imaging the effect of the circadian light-dark cycle on the

glymphatic system in awake rats. *Proc Natl Acad Sci U S A* **117**, 668-676 (2020).

44.   King, J.A.*, et al.* Procedure for minimizing stress for fMRI studies in conscious

rats. *J Neurosci Methods* **148**, 154-160 (2005).

45.   Colarusso, B.*, et al.* APOE4 rat model of Alzheimer's disease: sex differences,

genetic risk and diet. *BMC Neurosci* **25**, 57 (2024).

46.   R, H., E, B., S, P., P, K. & CF, F. Testing the Efficacy of Minocycline Treatment in

an Awake, Female Rat Model of Repetitive Mild Head Injury. *Journal Experimental*

Neurology **5**, 145-161 (2024).

47.   Bradshaw, H.B. & Johnson, C.T. Measuring the Content of Endocannabinoid-Like

Compounds in Biological Fluids: A Critical Overview of Sample Preparation

Methodologies. *Methods Mol Biol* **2576**, 21-40 (2023).

48.   Leishman, E., Kunkler, P.E., Hurley, J.H., Miller, S. & Bradshaw, H.B. Bioactive
Lipids in Cancer, Inflammation and Related Diseases : Acute and Chronic Mild
Traumatic Brain Injury Differentially Changes Levels of Bioactive Lipids in the CNS
Associated with Headache. *Adv Exp Med Biol* **1161**, 193-217 (2019).

49.   Leishman, E., Mackie, K. & Bradshaw, H.B. Elevated Levels of Arachidonic Acid-
Derived Lipids Including Prostaglandins and Endocannabinoids Are Present Throughout
ABHD12 Knockout Brains: Novel Insights Into the Neurodegenerative Phenotype. *Front*
Mol Neurosci **12**, 142 (2019).

50.   Tan, B*., et al.* Targeted lipidomics: discovery of new fatty acyl amides. *AAPS J* **8**,
E461-465 (2006).

51.   Leishman, E., Mackie, K., Luquet, S. & Bradshaw, H.B. Lipidomics profile of a
NAPE-PLD KO mouse provides evidence of a broader role of this enzyme in lipid
metabolism in the brain. *Biochim Biophys Acta* **1861**, 491-500 (2016).

52.   Gitcho, M.A*., et al.* VCP mutations causing frontotemporal lobar degeneration
disrupt localization of TDP-43 and induce cell death. *J Biol Chem* **284**, 12384-12398
(2009).

53.   Neumann, M*., et al.* Ubiquitinated TDP-43 in frontotemporal lobar degeneration
and amyotrophic lateral sclerosis. *Science* **314**, 130-133 (2006).

54.   Buckholtz, N.S., Zhou, D.F., Freedman, D.X. & Potter, W.Z. Lysergic acid
diethylamide (LSD) administration selectively downregulates serotonin2 receptors in rat
brain. *Neuropsychopharmacology* **3**, 137-148 (1990).

55.   Roberts, B.F*., et al.* Effect of psilocybin on decision-making and motivation in the
healthy rat. *Behav Brain Res* **440**, 114262 (2023).

56.   Smith, D.A., Bailey, J.M., Williams, D. & Fantegrossi, W.E. Tolerance and cross-
tolerance to head twitch behavior elicited by phenethylamine- and tryptamine-derived
hallucinogens in mice. *J Pharmacol Exp Ther* **351**, 485-491 (2014).

57.   Shanaki-Bavarsad, M., Almolda, B., Gonzalez, B. & Castellano, B. Astrocyte-
targeted Overproduction of IL-10 Reduces Neurodegeneration after TBI. *Exp Neurobiol*
31, 173-195 (2022).

58.   Kraft, A.W*., et al.* Attenuating astrocyte activation accelerates plaque
pathogenesis in APP/PS1 mice. *FASEB J* **27**, 187-198 (2013).

59.   Lu, M.C., Ji, J.A., Jiang, Z.Y. & You, Q.D. The Keap1-Nrf2-ARE Pathway As a

Potential Preventive and Therapeutic Target: An Update. *Med Res Rev* **36**, 924-963

(2016).

60.   Klein, R*., et al.* The trkB tyrosine protein kinase is a receptor for brain-derived

neurotrophic factor and neurotrophin-3. *Cell* **66**, 395-403 (1991).

61.   Spillantini, M.G*., et al.* Alpha-synuclein in Lewy bodies. *Nature* **388**, 839-840

(1997).

62.   McKee, A.C*., et al.* The first NINDS/NIBIB consensus meeting to define

neuropathological criteria for the diagnosis of chronic traumatic encephalopathy. *Acta*

Neuropathol **131**, 75-86 (2016).

63.   Bieniek, K.F*., et al.* Chronic traumatic encephalopathy pathology in a

neurodegenerative disorders brain bank. *Acta Neuropathol* **130**, 877-889 (2015).

64.   Cassidy, J.D*., et al.* Incidence, risk factors and prevention of mild traumatic brain

injury: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain

Injury. *Journal of rehabilitation medicine*, 28-60 (2004).

65.   Center for Disease Control and Prevention. National Vital Statistics System,

Mortality File. (ed. National Center for Health Statistics) (2015).

66.   AA, D.S., Kulkarni, P., Ferris, C.F., Amiji, M.M. & Bleier, B.S. Mild repetitive TBI

reduces brain-derived neurotrophic factor (BDNF) in the substantia nigra and

hippocampus: A preclinical model for testing BDNF-targeted therapeutics. *Exp Neurol*

374, 114696 (2024).

67.   R, H., E, B., S, P., P, K. & CF, F. Testing the Efficacy of Minocycline Treatment in

an Awake, Female Rat Model of Repetitive Mild Head Injury. *Journal Experimentlal*

Neurology **5** (2024).

68.   Gaetz, M., Goodman, D. & Weinberg, H. Electrophysiological evidence for the

cumulative effects of concussion. *Brain Inj* **14**, 1077-1088 (2000).

69.   Iverson, G.L., Gaetz, M., Lovell, M.R. & Collins, M.W. Cumulative effects of

concussion in amateur athletes. *Brain Inj* **18**, 433-443 (2004).

70.   Bens, N., Kulkarni, P. & Ferris, C.F. Changes in cerebral vascular reactivity

following mild repetitive head injury in awake rats: modeling the human experience. *Exp*

Brain Res (2024).

71.   Christensen, J.*, et al.* When Two Wrongs Make a Right: The Effect of Acute and
Chronic Binge Drinking on Traumatic Brain Injury Outcomes in Young Adult Female
Rats. *J Neurotrauma* **37**, 273-285 (2020).

72.   Kane, M.J.*, et al.* A mouse model of human repetitive mild traumatic brain injury.
J Neurosci Methods **203**, 41-49 (2012).

73.   Ren, Z.*, et al.* 'Hit & Run' model of closed-skull traumatic brain injury (TBI)
reveals complex patterns of post-traumatic AQP4 dysregulation. *J Cereb Blood Flow*
Metab **33**, 834-845 (2013).

74.   Tagge, C.A.*, et al.* Concussion, microvascular injury, and early tauopathy in
young athletes after impact head injury and an impact concussion mouse model. *Brain :*
a journal of neurology **141**, 422-458 (2018).

75.   Howlett, J.R., Nelson, L.D. & Stein, M.B. Mental Health Consequences of
Traumatic Brain Injury. *Biol Psychiatry* **91**, 413-420 (2022).

76.   Katz, D.I., Cohen, S.I. & Alexander, M.P. Mild traumatic brain injury. *Handb Clin*
Neurol **127**, 131-156 (2015).

77.   Toth, A. Magnetic Resonance Imaging Application in the Area of Mild and Acute
Traumatic Brain Injury: Implications for Diagnostic Markers? in *Brain Neurotrauma:*
Molecular, Neuropsychological, and Rehabilitation Aspects (ed. F.H. Kobeissy) (Boca
Raton (FL), 2015).

78.   Jha, R.M., Kochanek, P.M. & Simard, J.M. Pathophysiology and treatment of
cerebral edema in traumatic brain injury. *Neuropharmacology* **145**, 230-246 (2019).

79.   Gonzalez-Maeso, J.*, et al.* Hallucinogens recruit specific cortical 5-HT(2A)
receptor-mediated signaling pathways to affect behavior. *Neuron* **53**, 439-452 (2007).

80.   Jefsen, O.H., Elfving, B., Wegener, G. & Muller, H.K. Transcriptional regulation in
the rat prefrontal cortex and hippocampus after a single administration of psilocybin. *J*
Psychopharmacol **35**, 483-493 (2021).

81.   Martin, D.A., Marona-Lewicka, D., Nichols, D.E. & Nichols, C.D. Chronic LSD
alters gene expression profiles in the mPFC relevant to schizophrenia.
Neuropharmacology **83**, 1-8 (2014).

82.   Nichols, C.D. & Sanders-Bush, E. A single dose of lysergic acid diethylamide
influences gene expression patterns within the mammalian brain.
Neuropsychopharmacology **26**, 634-642 (2002).

83.   Ferris, C.F. Rethinking the Conditions and Mechanism for Glymphatic Clearance.
Front Neurosci **15**, 624690 (2021).

84.   Giaume, C. & McCarthy, K.D. Control of gap-junctional communication in
astrocytic networks. *Trends Neurosci* **19**, 319-325 (1996).

85.   Ding, M., Eliasson, C., Betsholtz, C., Hamberger, A. & Pekny, M. Altered taurine
release following hypotonic stress in astrocytes from mice deficient for GFAP and
vimentin. *Brain Res Mol Brain Res* **62**, 77-81 (1998).

86.   Kassner, A. & Roberts, T.P. Beyond perfusion: cerebral vascular reactivity and
assessment of microvascular permeability. *Top Magn Reson Imaging* **15**, 58-65 (2004).

87.   Davis, T.L., Kwong, K.K., Weisskoff, R.M. & Rosen, B.R. Calibrated functional
MRI: mapping the dynamics of oxidative metabolism. *Proc Natl Acad Sci U S A* **95**,
1834-1839 (1998).

88.   Brevard, M.E., Duong, T.Q., King, J.A. & Ferris, C.F. Changes in MRI signal
intensity during hypercapnic challenge under conscious and anesthetized conditions.
Magn Reson Imaging **21**, 995-1001 (2003).

89.   Sicard, K.M., Henninger, N., Fisher, M., Duong, T.Q. & Ferris, C.F. Long-term
changes of functional MRI-based brain function, behavioral status, and histopathology
after transient focal cerebral ischemia in rats. *Stroke* **37**, 2593-2600 (2006).

90.   Sur, S.*, et al.* Association of cerebrovascular reactivity and Alzheimer pathologic
markers with cognitive performance. *Neurology* **95**, e962-e972 (2020).

91.   Glodzik, L., Randall, C., Rusinek, H. & de Leon, M.J. Cerebrovascular reactivity
to carbon dioxide in Alzheimer's disease. *J Alzheimers Dis* **35**, 427-440 (2013).

92.   Yezhuvath, U.S.*, et al.* Forebrain-dominant deficit in cerebrovascular reactivity in
Alzheimer's disease. *Neurobiol Aging* **33**, 75-82 (2012).

93.   Pillai, J.J. & Mikulis, D.J. Cerebrovascular reactivity mapping: an evolving
standard for clinical functional imaging. *AJNR Am J Neuroradiol* **36**, 7-13 (2015).

94.   Pelizzari, L.*, et al.* Cerebrovascular reactivity and its correlation with age in
patients with multiple sclerosis. *Brain Imaging Behav* **14**, 1889-1898 (2020).

95. Kenney, K.*, et al.* Cerebral Vascular Injury in Traumatic Brain Injury. *Exp Neurol*
275 Pt 3, 353-366 (2016).

96. Liu, P.*, et al.* Multi-vendor and multisite evaluation of cerebrovascular reactivity
mapping using hypercapnia challenge. *Neuroimage* **245**, 118754 (2021).

97. Amyot, F.*, et al.* Assessment of cerebrovascular dysfunction after traumatic brain
injury with fMRI and fNIRS. *Neuroimage Clin* **25**, 102086 (2020).

98. Amyot, F.*, et al.* Imaging of Cerebrovascular Function in Chronic Traumatic Brain
Injury. *J Neurotrauma* **35**, 1116-1123 (2018).

99. Haber, M.*, et al.* Vascular Abnormalities within Normal Appearing Tissue in
Chronic Traumatic Brain Injury. *J Neurotrauma* **35**, 2250-2258 (2018).

100. Wu, L.*, et al.* Persistent CO_2 reactivity deficits are associated with neurological
dysfunction up to one year after repetitive mild closed head injury in adolescent mice. *J*
Cereb Blood Flow Metab **41**, 3260-3272 (2021).

101. Golding, E.M.*, et al.* Cerebrovascular reactivity to CO_2 and hypotension after
mild cortical impact injury. *Am J Physiol* **277**, H1457-1466 (1999).

102. Mutch, W.A.*, et al.* Longitudinal Brain Magnetic Resonance Imaging CO2 Stress
Testing in Individual Adolescent Sports-Related Concussion Patients: A Pilot Study.
Front Neurol **7**, 107 (2016).

103. Champagne, A.A., Coverdale, N.S., Fernandez-Ruiz, J., Mark, C.I. & Cook, D.J.
Compromised resting cerebral metabolism after sport-related concussion: A calibrated
MRI study. *Brain Imaging Behav* **15**, 133-146 (2021).

104. Champagne, A.A., Coverdale, N.S., Germuska, M. & Cook, D.J. Multi-parametric
analysis reveals metabolic and vascular effects driving differences in BOLD-based
cerebrovascular reactivity associated with a history of sport concussion. *Brain Inj* **33**,
1479-1489 (2019).

105. D'Souza, M.M.*, et al.* Alterations of connectivity patterns in functional brain
networks in patients with mild traumatic brain injury: A longitudinal resting-state
functional magnetic resonance imaging study. *Neuroradiol J* **33**, 186-197 (2020).

106. Messe, A.*, et al.* Specific and evolving resting-state network alterations in post-
concussion syndrome following mild traumatic brain injury. *PLoS One* **8**, e65470 (2013).

107.   Tang, L.*, et al.* Thalamic resting-state functional networks: disruption in patients
with mild traumatic brain injury. *Radiology* **260**, 831-840 (2011).

108.   Shumskaya, E., Andriessen, T.M., Norris, D.G. & Vos, P.E. Abnormal whole-brain
functional networks in homogeneous acute mild traumatic brain injury. *Neurology* **79**,
175-182 (2012).

109.   Pinky, N.N.*, et al.* Multimodal magnetic resonance imaging of youth sport-related
concussion reveals acute changes in the cerebellum, basal ganglia, and corpus
callosum that resolve with recovery. *Front Hum Neurosci* **16**, 976013 (2022).

110.   Fitzgerald, B.*, et al.* Longitudinal changes in resting state fMRI brain self-
similarity of asymptomatic high school American football athletes. *Sci Rep* **14**, 1747
(2024).

111.   Huang, E.J. & Reichardt, L.F. Neurotrophins: roles in neuronal development and
function. *Annu Rev Neurosci* **24**, 677-736 (2001).

112.   Zeiler, F.A.*, et al.* Genetic drivers of cerebral blood flow dysfunction in TBI: a
speculative synthesis. *Nat Rev Neurol* **15**, 25-39 (2019).

113.   Frisen, J.*, et al.* Characterization of glial trkB receptors: differential response to
injury in the central and peripheral nervous systems. *Proc Natl Acad Sci U S A* **90**,
4971-4975 (1993).

114.   Gustafsson, D., Klang, A., Thams, S. & Rostami, E. The Role of BDNF in
Experimental and Clinical Traumatic Brain Injury. *Int J Mol Sci* **22** (2021).

115.   Moliner, R.*, et al.* Psychedelics promote plasticity by directly binding to BDNF
receptor TrkB. *Nat Neurosci* **26**, 1032-1041 (2023).

116.   Zhao, X.*, et al.* Psilocybin promotes neuroplasticity and induces rapid and
sustained antidepressant-like effects in mice. *J Psychopharmacol* **38**, 489-499 (2024).

117.   Rosa, E.*, et al.* Tau downregulates BDNF expression in animal and cellular
models of Alzheimer's disease. *Neurobiol Aging* **48**, 135-142 (2016).

118.   Joachim, C.L., Morris, J.H., Selkoe, D.J. & Kosik, K.S. Tau epitopes are
incorporated into a range of lesions in Alzheimer's disease. *Journal of neuropathology
and experimental neurology* **46**, 611-622 (1987).

119.   Weingarten, M.D., Lockwood, A.H., Hwo, S.Y. & Kirschner, M.W. A protein factor
essential for microtubule assembly. *Proc Natl Acad Sci U S A* **72**, 1858-1862 (1975).

120. Mandelkow, E.M.*, et al.* Tau domains, phosphorylation, and interactions with

microtubules. *Neurobiol Aging* **16**, 355-362; discussion 362-353 (1995).

121. Rippon, G.A.*, et al.* Late-onset frontotemporal dementia associated with

progressive supranuclear palsy/argyrophilic grain disease/Alzheimer's disease

pathology. *Neurocase* **11**, 204-211 (2005).

122. Josephs, K.A.*, et al.* Neuropathological background of phenotypical variability in

frontotemporal dementia. *Acta Neuropathol* **122**, 137-153 (2011).

123. Dickson, D.W.*, et al.* Ballooned neurons in select neurodegenerative diseases

contain phosphorylated neurofilament epitopes. *Acta Neuropathol* **71**, 216-223 (1986).

124. Li, Q.*, et al.* Changes in Lipidome Composition during Brain Development in

Humans, Chimpanzees, and Macaque Monkeys. *Mol Biol Evol* **34**, 1155-1166 (2017).

125. Burugupalli, S.*, et al.* Ontogeny of circulating lipid metabolism in pregnancy and

early childhood - a longitudinal population study. *Elife* **11** (2022).

126. Stuart, J.M., Paris, J.J., Frye, C. & Bradshaw, H.B. Brain levels of prostaglandins,

endocannabinoids, and related lipids are affected by mating strategies. *Int J Endocrinol*

2013, 436252 (2013).

1027

Figure Legends

Fig 1. Experimental Procedures and Behavior

Shown above is a timeline (a.) of experimental procedures. Below (b.) are results from motor behavior on the Beam Walk, Rotarod, and Open Field, and (c.) cognitive behavior from Novel Object Recognition and emotional behaviors agoraphobia and thigmotaxis collected from the Open Field test. All behavioral data were collected within one week of the last head impact. One-way ANOVAs showed no significant (ns) differences between any of the experimental groups for any assay.

Fig 2. Head Impact and Diffusion Weighted Imaging

Shown in (a.) are radiograms of frontal sections of the brain of all subjects taken following the last of three head impacts. The arrows point to the approximate site of impact identified by the T2-weighted enhanced signal showing the edema on the skin above the skull. This is obvious in all hit rats but not in sham controls. Shown in (b.) are whole-brain changes in water diffusivity as measured by apparent diffusion coefficient (ADC) and fractional anisotropy (FA). When comparing SHAM-VEH and rmTBI-VEH there was a global increase in ADC and decrease in FA. There was a significant inverse relationship between ADC and FA values (r = -.5967) as shown in the bottom graph. Each red dot is a brain area taken from the rat 3D brain atlas. A comparison across all three experimental groups shows that PSI treatment significantly reduces ADC and increases FA values, reversing the effects of head injury. Shown in (c.) are the regional changes in ADC values for each of the experimental groups. In all brain regions, rmTBI-VEH rats show a significant increase over SHAM-VEH controls, while rmTBI-PSI treated rats show a significant decrease compared to rmTBI-VEH rats. Error bars show standard deviation. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$

Fig 3. Vascular Reactivity

Shown are changes in vascular reactivity (voxel number) in response to a 5% CO_2 challenge. Whole-brain (box) was significantly elevated above control in rmTBI-VEH rats (gray bars). rmTBI-PSI treated rats were significantly lower than rmTBI-VEH untreated rats. Regional differences in vascular reactivity are shown in the bar and

dot plots (mean ± SD). Each dot is a subregion in that brain region. * p<0.05, **

p<0.01, *** p<0.001, **** p<0.0001

Fig 4. Functional Connectivity

Shown in (a.) are the number of degrees (connections) for the whole brain (red box)

for each experimental group (mean ± SD). Regional differences in degrees are

shown in the bar and dot plots (mean ± SD). Each dot is a subregion of that brain

region. In all cases, rmTBI-PSI treated rats showed greater connectivity than SHAM-

VEH or rmTBI-VEH rats. The connectivity to the dopaminergic system for each

experimental condition is shown below in (b.) The top bar graphs (mean ± SD) to the

left show the number of efferent connections from the three major dopaminergic

nuclei in the midbrain (ventral tegmental area, substantial nigra compacta, and

substantia nigra reticularis) while the radial connectivity maps to the right depict

these connections to color-coded brain regions. The bottom bar graphs to the left

(median and 1^{st} and 3^{rd} quartile) show the number of connections between and

within the dopaminergic system. * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001

Fig 5. Phosphorylated tau reduced to control levels in rmTBI/PSI treated
animals compared to rmTBI

Western blot of RIPA-soluble proteins (a.) Quantitative analysis normalized to tubulin

of RIPA fractions (b.) Western blot RIPA insoluble/urea soluble fractions (c.)

Quantitative analysis normalized to total protein (d.) Statistical analysis: one way

ANOVA with Tukey post hoc (*n* = 8 biological replicates/3 groups) (e.) * p<0.05 and

** p<0.002 (all p vales see table).

Fig 6. Plasma Lipids

The direction of changes for each analysis group relative to SHAM-VEH are

depicted by color and arrow direction, with an increase represented by a green box

and upward arrows and a decrease represented by an orange box with downward

arrows (boxed insert). Level of significance is shown by color shade, wherein

$p < 0.05$ is a dark shade and $0.05 < p < 0.1$ is a light shade. Effect size is represented

by the number of arrows, where 1 arrow corresponds to 1–1.49-fold difference, 2

arrows to a 1.5–1.99-fold difference, 3 arrows to a 2–2.99-fold difference, 4 arrows a

3–9.99-fold difference, and 5 arrows a difference of tenfold or more[126]. An

abbreviation of 'BDL' indicates that the lipid concentration that was present in the

sample was below the detectable levels of our equipment while 'BAL' indicates

below analytical levels.

Fig 1. Experimental Protocol and Behavior Assessment

Fig 2. Head Injury and Diffusion Weighted Imaging

Fig 3. Vascular Reactivity to Hypercapnia

Fig 4. Functional Connectivity

Fig 5. Phosphorylated Proteins

 Fig 6. Plasma Lipids

Small Plasma Lipids	Sham -Vehicle vs Hit -Vehicle	Hit-Vehicle vs Hit-Psilocybin
2-acyl glycerol		
2-palmitoyl glycerol		
2-oleoyl glycerol		
2-linoleoyl glycerol		
2-arachidonoyl glycerol		
N-acyl alanine		
palmitoyl alanine		
stearoyl alanine		
oleoyl alanine		↑↑
linoleoyl alanine		
arachidonoyl alanine	BAL	↑
docosahexaenoyl alanine		
N-acyl ethanolamine		
palmitoyl ethanolamine		
stearoyl ethanolamine		
oleoyl ethanolamine		
linoleoyl ethanolamine		
arachidonoyl ethanolamine		
docosahexaenoyl ethanolamine		
eicosapentaneoyl ethanolamine	BDL	BDL
N-acyl GABA		
palmitoyl GABA	BAL	BAL
stearoyl GABA	BAL	BDL
oleoyl GABA	BDL	BDL
linoleoyl GABA	BDL	BDL
arachidonoyl GABA	BDL	BDL
docosahexaenoyl GABA	BDL	BDL
N-acyl glycine		
palmitoyl glycine		
stearoyl glycine		
oleoyl glycine		↑↑
linoleoyl glycine		↑
arachidonoyl glycine		
docosahexaenoyl glycine		
N-acyl leucine		
palmitoyl leucine		
stearoyl leucine	↓	
oleoyl leucine		
linoleoyl leucine		
arachidonoyl leucine		
docosahexaenoyl leucine		
N-acyl methionine		
palmitoyl methionine		↑
stearoyl methionine		
oleoyl methionine		↑↑
linoleoyl methionine		
docosahexaenoyl methionine		
N-acyl phenylalanine		
palmitoyl phenylalanine		
stearoyl phenylalanine		
oleoyl phenylalanine		
linoleoyl phenylalanine		
arachidonoyl phenylalanine	BAL	BAL
docosahexaenoyl phenylalanine		↑↑
N-acyl proline		
palmitoyl proline		
stearoyl proline		
oleoyl proline	BAL	BAL
linoleoyl proline	BAL	BAL
arachidonoyl proline	BDL	BDL
docosahexaenoyl proline	BDL	BDL

	Sham -Vehicle vs Hit -Vehicle	Hit-Vehicle vs Hit-Psilocybin
N-acyl serine		
palmitoyl serine		
stearoyl serine		
oleoyl serine		↑
linoleoyl serine		↑
arachidonoyl serine	BDL	BDL
docosahexaenoyl serine	BDL	BDL
N-acyl taurine		
palmitoyl taurine	↓	
stearoyl taurine	↓	↑
oleoyl taurine		
arachidonoyl taurine		↑↑
N-acyl tryptophan		
palmitoyl tryptophan	BDL	BDL
stearoyl tryptophan	BDL	BDL
oleoyl tryptophan		
linoleoyl tryptophan	BDL	BDL
arachidonoyl tryptophan	BDL	BDL
docosahexaenoyl tryptophan	BAL	BAL
N-acyl tyrosine		
palmitoyl tyrosine	BAL	BAL
stearoyl tyrosine	BDL	BDL
oleoyl tyrosine		
linoleoyl tyrosine		↑↑
arachidonoyl tyrosine	BDL	BDL
docosahexaenoyl tyrosine	BAL	BAL
N-acyl valine		
palmitoyl valine		
stearoyl valine		
oleoyl valine		
linoleoyl valine		
arachidonoyl valine	BAL	BAL
docosahexaenoyl valine		
Free fatty acids		
oleic acid		↑↑↑
linoleic acid		↑↑↑
arachidonic acid		↑↑
docosahexaenoic acid		↑↑
stearidonic acid		
eicosapentaenoic acid		↑↑

(light orange)	trending (decrease) (p<0.1-0.51)
(orange)	Significant (decrease) (p<0.50)
(green)	Significant (increase) (p<0.50)
(light green)	trending (increase) (p<0.1-0.51)
↓	1 - 1.49 times lower
↓↓	1.5 - 1.99 times lower
↓↓↓	2 - 2.99 times lower
↓↓↓↓	3 - 9.99 times lower
↓↓↓↓↓	10 or more times lower
↑	1 - 1.49 times higher
↑↑	1.5 - 1.99 times higher
↑↑↑	2 - 2.99 times higher
↑↑↑↑	3 - 9.99 times higher
↑↑↑↑↑	10 or more times higher