A dynamic repertoire of wound closure strategies precedes whole-body regeneration

Allison P. Kann and Mansi Srivastava

Department of Organismic and Evolutionary Biology, Museum of Comparative Zoology,

Harvard University, Cambridge, MA 02138

Corresponding Author Emails: allisonkann@fas.harvard.edu, mansi@oeb.harvard.edu

KEYWORDS

Regeneration, wound healing, wound closure, morphogenesis, acoels, epithelium

ABSTRACT

Wound closure is an essential aspect of successful regeneration, often acting as the first morphogenetic

event that precedes downstream cellular events. Despite heavy investment in studying whole-body

regeneration in many invertebrate systems, the steps by which these organisms heal their wounds remain

understudied. Here, we investigate the cellular mechanisms of wound healing in the acoel *Hofstenia*

miamia, an invertebrate worm capable of whole-body regeneration. *H. miamia* have two distinct epithelial

layers, an outer epidermis and the epithelial lining of their pharynx. By labeling regenerating fragments

with an actin dye, we found that *H. miamia* use distinct mechanisms of epithelial wound repair across

different injury contexts. In transverse wounds that don't injure the pharyngeal epithelium, the epidermis

closes by gradual radial constriction. In contrast, injuries that damage both the epidermis and the

pharyngeal epithelium show the formation of long, actin-rich protrusions that cross the wound gap and

form heterotypic bridges prior to re-epithelialization. Muscle contraction is required for the formation of

heterotypic bridges - when animals are anesthetized and immobile, they are unable to form these cellular

bridges, and epidermal cells cannot migrate independently. Global actomyosin contractility also plays a

role in repair mechanisms, and pharmacological perturbation of contractility shifts the dynamics of wound

closure after amputation. In the presence of blebbistatin, heterotypic bridge formation is inhibited but

homotypic re-epithelialization is accelerated through increased cell crawling. Together, this work identifies

mechanisms by which epithelial layers close large wounds *in vivo*, identifying novel heterotypic cellular

bridges in wound closure and demonstrating the conservation of actin-mediated processes in an early-

diverging phylum.

INTRODUCTION

The ability of animals to regenerate varies substantially across phyla, ranging from more limited, tissue-specific or structural regeneration to extreme cases of whole-body regeneration (WBR), where any missing cell or tissue type can be replaced after amputation[1,2]. Highly regenerative animals undergoing WBR face unique healing challenges and must be able to scarlessly repair virtually any injury prior to downstream regeneration. Most mechanistic studies of wound closure have focused on relatively simple injuries (i.e. damaging only the epidermis) in animals that are not capable of substantial regeneration, generating foundational knowledge about cellular behaviors but lacking a strong connection to downstream regeneration[3–8]. The work on wound closure in regenerative animals has primarily focused on either the formation and function of the 'wound epidermis', a specialized epidermis that covers the injury site and acts as an organizing center to induce downstream regeneration[9–12], or large-scale transcriptional atlases detailing the molecular wound response[13–18]. Some studies have demonstrated the presence of conserved cellular mechanisms in regenerative animals[19–25], but it remains ambiguous how much flexibility exists in closure strategies prior to regeneration. Specifically, it is unclear whether highly regenerative animals employ a uniform healing process regardless of injury context (i.e. major vs. minor wounds, varying geometries and tissue organizations) or if they have a range of mechanisms that can be tailored to different injury conditions.

Previous work has shown the presence of shared mechanisms of wound repair across the metazoan tree of life, including in early-branching lineages such as cnidarians[14,16,19,20], and in protostomes[3,4,21,25–27] and deuterostomes[15,18,28–30]. This suggests a deep conservation of cellular behaviors, evolving prior to the emergence of bilateral symmetry. A fundamental aspect of wound closure and subsequent healing is the process of re-epithelialization, in which a damaged epithelial layer undergoes cellular and morphological changes to re-establish an intact tissue[6]. Wound closure through cell crawling is facilitated through the formation of actin-based membrane protrusions (such as lamellipodia or filopodia) in cells at the periphery of a wound that drive migration into the gap[7,31,32]. Actomyosin contractility is essential in this crawling mechanism for both retraction at the rear of migrating cells and the formation of tension between adhesion points[19,33,34]. Purse string closure is similarly driven by contractility; cells around the wound form a supracellular cable, contracting and bringing the edges of

cells inwards to pinch off the gap[31,35,36]. These mechanisms are not mutually exclusive and often work in

tandem, bringing the edges of epithelial cells together across the open wound and restoring tissue

integrity through the reestablishment of cell junctions[7].

Studies both *in vivo* and *in vitro* have revealed that the mechanisms of re-epithelialization vary

substantially between wounding conditions, with factors such as wound size, geometry, tissue type,

extracellular matrix composition, and life stage of the animal all influencing the mechanism of closure[6,7,]

[19,23,26,4133,38–41]. Large injuries typically involve damage to the underlying extracellular matrix, leading to

either wound closure through the formation of actomyosin purse strings, which do not require an intact

substrate, and/or the deposition of a transient matrix that provides a substrate for epithelial cell

crawling[19,22,24,37,42]. These large injuries also require coordination between layers of cells, often relying on

contraction of mesenchymal cells underneath the epithelium to drive closure[29,43,44]. Despite evidence of

variable cellular behaviors across wound conditions, most *in vivo* studies have focused on factors that

determine whether wounds close or not, rather than the flexibility of large-scale closure strategies within

the same system. This is likely due to a few factors. First and foremost, traditional model organisms that

have extensive genetic and imaging tools (e.g. *D. melanogaster, M. musculus*) have distinct physiological

considerations. Not only are these species incapable of whole-body regeneration, restricting the types of

injuries the animal can recover from, but they possess circulatory and immune systems that complicate

and contribute to the healing process[44,45]. Second, visualizing wound closure in a quantifiable and

reproducible manner typically requires live imaging, which is more feasible to set up for minor wounds

(such as a scratch assay, as used in *C. hemisphaerica*[19]) than for large injuries. We therefore have a

detailed understanding of some cellular decision-making, like switching between cell crawling and purse

string formation depending on substrate[3,19,35,40,46], but less is understood about whether animals utilize the

same toolkit of closure strategies across extreme amputations. Studying these cellular processes in

nontraditional animals offers a new approach to understanding fundamental aspects of biology,

capitalizing on their regenerative properties to ask questions that are infeasible in other species.

Hofstenia miamia is a marine invertebrate worm belonging to the enigmatic and understudied

phylum Xenacoelomorpha, which likely represents an early-diverging animal lineage that is a sister group

to all other bilaterians (Fig. 1A-B)[47–49]. *H. miamia* has emerged as a new model system, largely studied for

its ability to undergo whole-body regeneration. Like other animals capable of WBR, such as

platyhelminthes, *H. miamia* rely on a population of adult pluripotent stem cells to drive regeneration [50,51].

However, even in conditions that block regeneration (e.g. functional knockdown of required genes or

irradiation of proliferative cells[47,50]), damaged fragments are still able to visibly close their wounds –

suggesting that healing and regeneration are distinct processes. In addition to their remarkable

regenerative ability, *H. miamia* possess a 'tube within a tube' anatomy, with an outer epidermis and a

centrally located pharynx lined with an epithelium[52]. This dual-epithelial organization prompts both

questions of how complex injuries can heal and how tissues maintain their identity during repair and

regeneration.

In this study, we investigate the healing capacity of *H. miamia*, examining whether wound repair

mechanisms shift in response to different injury conditions. In relatively simple injuries (e.g. transverse

amputations that only damage one epithelial layer, scratch wounds), animals rely on conserved

mechanisms of epithelial contraction and migration. In more complex wounds (e.g. asymmetric sagittal

wounds and other amputations that damage both epithelial layers), *H. miamia* form novel heterotypic

bridges between epithelial layers prior to homotypic re-epithelialization. These bridges are dependent on

both global actomyosin tension and contractility generated by neighboring muscle, but the regenerating

fragments can modify their repair strategies in the face of mechanical perturbation. *H. miamia* therefore

possess an adaptable toolkit of wound closure mechanisms, with flexible strategies to ensure successful

repair and downstream regeneration regardless of injury type.

RESULTS

***Hofstenia miamia* have two distinct, atypical epithelial tissues**

The acoel *Hofstenia miamia* can recover from virtually any injury throughout its lifespan, scarlessly

healing wounds and rebuilding lost tissues[47,48]. Despite wound closure being an essential prerequisite to

this extensive regeneration, the mechanisms of wound closure have not yet been studied in *H. miamia*.

To fully understand the dynamics of re-epithelization, we first needed to characterize the epithelial layers

of *H. miamia*.

H. miamia have two distinct epithelial tissues: (1) an outer epidermis that covers the whole

animal, and (2) the inner lining of the pharynx, a tube-shaped organ that extends ~1/2 down the anterior-

posterior axis (Figure 1C). Although the dynamics of these cells have not yet been studied, previous

transgenic labeling has revealed their cellular morphologies[52]. This transgenic labeling (using *EpiA:RFP*

animals[52]) enables the visualization of detailed cell morphologies (Figure 1D), but the mosaicism prevents

consistent imaging across animals. We found that the cytoskeletal dye SiR-actin robustly labels both

layers of epithelial cells in fixed animals and can therefore be used to visualize entire tissues instead of

individual cells (Figure 1E-F). The apical side of *H. miamia* epithelial cells resemble epithelia in other

systems[53], with the cells organized in a polygonal pattern on the surface of the animal (Figure 1E). As

previously reported, the basal side of both epidermal and pharyngeal epithelial cells form cytoplasmic

protrusions that extend into the body of the worm (Figure 1F-H). The protrusions are irregularly spaced

and heterogeneous in number between cells but extend approximately 10-25µm in length, with the

epidermal protrusions being slightly (but significantly) longer than those in the pharyngeal epithelium

(Figure 1G). The basal protrusions in both cell types end in close proximity to underlying muscle (Figure

1H), but it is unclear whether cell-cell junctions form at the tips of these projections or whether they

adhere to an atypical basement membrane.

The unique morphology and organization of these tissues raise questions of how epithelial cells

mobilize to close wounds upon amputation and facilitate downstream regeneration. Even though *H.*

miamia can recover from virtually any injury configuration, not all injuries require the same cellular

response. Notably, the internal localization of the pharynx means that some amputations will damage

both epithelial layers, adding another layer of complexity to the re-epithelialization process. We therefore

sought to understand the dynamics of re-epithelialization during wound closure, comparing injuries that

damage one or both epithelial layers.

Amputations that damage the outer epidermal layer close through actin-based constriction

Transverse amputations through the middle of the animal, separating the head from tail, have been

utilized as the dominant assay for whole-body regeneration across systems[14,47,48,50,51]. In *H. miamia,* a

typical transverse amputation aims to cut at the middle pigmentation stripe observable on the dorsal

surface of the worm, leaving behind a large gap with open epidermal edges around the wound. This

amputation halves some tissues (e.g. gut, nerve net, and musculature) and completely removes anterior

organs (e.g. pharynx and brain) from the posterior fragment. In addition to the biological process of

wound healing, these fragments therefore face different regenerative challenges: the head fragment

needs to regenerate the posterior tissues, whereas the tail fragment must regenerate anterior-specific

tissues and organs (Figure 2A).

We first amputated juvenile worms transversely below the pharynx, so that only the outer

epidermis (but not the pharyngeal epithelium) would be impacted by the injury. Consistent with previous

reports (cite), each fragment showed a rounded wound edge within 24 hours, suggesting successful

closure. Head fragments formed a pointed tail within 3 days (red arrowhead) and regained their original

shape within 5 days. The tail fragments took longer to regenerate but formed a visible blastema within 5

days (white arrowhead) and regained their original form within 10 days (Figure 2B). This temporal

trajectory reflects the regenerative challenge each fragment faces; the tail fragments need to replace

more cell types than their head counterparts. Although the brightfield imaging suggested that wound sites

had closed within 24 hours, we wanted to assess the cellular dynamics of the re-epithelialization process.

To track the dynamics of the outer epidermis during wound closure, we stained fragments with the

cytoskeletal dye SiR-actin at intermediate time points during the first 10 hours post-amputation (hpa).

Wounds from both head and tail fragments largely closed within 10 hpa, and both showed a strong

anisotropic enrichment of actin at the wound periphery, suggesting the formation of a supracellular

structure (Figure 2C-D). These observations appear consistent with a process of radial constriction,

where actin polarizes towards the wound edge and cells collectively pull towards the center of the wound

to minimize the size of the gap. Some wound edges displayed a smooth, continuous actin cable (yellow

arrowheads), implying the formation of an actomyosin purse-string[31,36], but others appear puckered (red

arrowheads), suggesting more of a general constrictive mechanism. Both head and tail fragments closed

their wounds at a similar rate; head wounds were more open than tail wounds at 3 hpa, but by 6 hpa we

found no difference between the two conditions (Figure 2D). Despite closure of the wound sites by 10

hpa, we observed visible differences in wound edges between 10 and 20 hpa. Closed wounds at 10 hpa

had visible seams where the epidermis was brought together (yellow arrows), but by 20 hpa there was no

detectable seam or wound edge (Figure 2E). This suggests a sustained re-epithelialization process even

after initial gap closure, where the epidermal cells continue to remodel their junctions and re-establish a

seamless epithelium.

To assess whether minor wounds utilize the same mechanisms of actomyosin contractility during

re-epithelialization as the transversely amputated wounds above, we performed 'pinch' wounds, where

fine-tipped forceps were used to pinch through the outer epidermis (Supplemental Figure 1A). These

wounds closed faster than wounds at transverse amputations and seemed to use a different cellular

mechanism. Instead of contracting across the wound gap, we observed the formation of polarized

protrusions within 5 minutes post-amputation, with the wound edges making contact and closing up within

1 hour (Supplemental Figure 1A). This is likely due to the much smaller distance between wounded

epidermal edges; in a pinch wound, the edges remain adjacent to each other, facilitating quick re-

epithelialization, but a transverse amputation causes a comparatively large wound gap that requires more

time to close.

Together, these results show that wounds that disrupt only one epithelial layer use shared

mechanisms of closure seen in other systems across contexts of regeneration, development, and repair.

Minor wounds with small gaps can close quickly and without the formation of supracellular structures,

whereas large amputations require the constriction of epithelial cells across the wound gap. We next

looked at the dynamics of closure in more complex wounds, specifically injuring both the outer epidermis

and the pharyngeal epithelium.

Heterotypic epithelial bridges mediate the first step in multi-epithelial wound closure

192 To investigate mechanisms of wound closure involving multiple epithelial layers, we performed injuries

193 that damage both the outer epidermis and the pharynx. We first cut animals in the sagittal plane,

194 separating the left half from the right, and observed wounds over a 10-day time course of regeneration

195 (Figure 3A-B). Interestingly, these animals were able to mostly restore their original shape within 5 days,

196 rounding themselves up without the formation of a visible blastema (Figure 3B). We therefore asked how

197 the epithelial layers facilitate the morphallactic processes of wound closure. Using SiR-actin labeling, we

198 amputated worms in the sagittal plane and observed the formation of striking, long cytoplasmic

199 protrusions that extend across the wound edge in the first 10 hpa. These protrusions originated from both

200 the outer epidermis and the pharyngeal epithelium, connecting across the middle of the wound (Figure

201 3C). To confirm that these cells were epithelial in nature, we utilized the *EpiA:RFP* transgenic line[52] and

202 detected protrusions in labeled cells within both the outer epidermis and the pharyngeal epithelium

203 (Supplementary Figure 2A-B). We were unable to visualize the basal side of cells forming these bridges,

204 but the extensions protruded from the apical surface of the cells - suggesting that these were new

205 structures and not movement of the homeostatic basal protrusions.

206  In sagittal wounds, epidermal cells showed signs of actin polarization within the first 3 hours post-

207 amputation (hpa), with small filopodial protrusions (5-10 µm) appearing in the outer epidermis oriented

208 towards the wound edge (Figure 3B). Long, thin extensions (>15 µm) emerged around 6 hpa, forming

209 more robustly by 10 hpa and connecting to projections originating from the pharyngeal epithelium (Figure

210 3D-E). This temporal pattern was consistent across animals, and 67% of fragments displayed at least one

211 connected bridge within 10 hpa (Figure 3E). Notably, protrusions were always detected first in the outer

212 epidermis, and protrusions from the pharyngeal epithelium were only visible after their counterparts

213 formed in the epidermis (Figure 3F). Corresponding with the number of bridges increasing between 6 and

214 10 hpa, the average bridge length increased significantly between time points (Figure 3G). This stepwise

215 process (polarization, extension, connection) is consistent with a model of collective cell migration[54,55], but

216 the lengths of these bridges (~30-100 µm) are reminiscent of specialized projections such as cytonemes

217 or tunneling nanotubes (which have been reported to reach up to 200 µm *in vivo*[56–58]). To the best of our

218 knowledge, structures like these heterotypic bridges have not been reported in wound closure literature,

219 and we therefore sought to further investigate this phenomenon.

The difference between transverse, contraction-based closure and sagittal, bridge-based closure

could be attributed to a variety of differences between the amputations other than the dual epithelial

layers, including (but not limited to): angle or orientation of amputation, size of the wound, or the extent of

regeneration required downstream of closure. We next asked whether heterotypic bridges occurred in

transverse amputations further up the anterior-posterior axis of the animal, upon injuries that impact both

epithelial layers by cutting through the pharynx. These wounds require closure in the same plane as the

transverse injuries reported above, but now include damage to the pharyngeal epithelium. When we

performed single and double transverse amputations through the pharynx, we observed robust

heterotypic bridges between the epidermis and the pharyngeal epithelium, similar to those seen in sagittal

amputations (Figure 3H-I). This bridge-based mechanism of closure is therefore not specific to the size or

the irregular wound geometry that comes from a sagittal amputation, but appear to instead be a more

general mechanism by which to close wounds when both epithelial layers are damaged.

Although these heterotypic bridges are unique in presentation, we reasoned that there must be a

substrate below these protrusions acting as a physical track for their extension. The extracellular matrix

(ECM) has been shown to be a key regulator of wound healing, acting as a dynamic substrate on which

cells can crawl and extend to complete re-epithelialization[42,59,60]. Given the known role of collagen IV in

wound closure[43,60–62], we identified a putative ortholog in the *H. miamia* transcriptome and made a custom

antibody (see Methods). Using this antibody, we performed immunofluorescence staining to assess the

localization of collagen IV and detected its expression solely under the pharyngeal epithelium

(Supplemental Figure 2C-D). Next, we asked whether collagen IV was remodeled during wound closure

as the pharyngeal epithelium is extending its protrusions. We performed immunofluorescence on 10 hpa

sagittal fragments and observed strong co-localization of collagen IV specifically underneath pharyngeal

extensions. Collagen signal was never detected in areas without protrusions, suggesting that nascent

collagen is being deposited underneath the pharyngeal epithelium as it is forming wound-induced bridges

(Supplemental Figure 2E). The same pattern was observed after double transverse amputation,

suggesting a shared mechanism regardless of wound geometry (Supplemental Figure 2F). These data

implicate collagen IV as a substrate for epithelial bridges, likely providing physical support for protrusions

as they extend across the wound site. Although our antibody specifically labels the collagen underneath

the pharyngeal epithelium, we speculate that a similar process is occurring underneath protrusions

coming from outer epidermal cells.

Taken together, these findings reveal the presence of heterotypic cellular bridges between

epithelial layers during wound closure, which emerge within the first 10 hours post-amputation. Tissue-

specific collagen is remodeled during the extension of pharyngeal protrusions, and bridges are observed

across different amputation geometries that injure multiple epithelial layers. We therefore hypothesized

that these heterotypic interactions act as an intermediate step in large-scale multi-epithelial wound

closure, possibly minimizing tissue loss and/or maintaining tissue identity prior to full re-epithelialization.

To the best of our knowledge, this phenomenon appears to be a novel healing mechanism, and we

sought to understand how these heterotypic interactions relate to the rest of the wound healing process.

Wound closure involving multiple epithelial tissues utilizes distinct but temporally overlapping

cellular mechanisms

The bridges described above represent a transient phase in wound closure, emerging within the first 10

hours post-amputation (hpa). In order for the animal to heal fully, the sagittal fragments must undergo

continued remodeling to re-establish their original shape. Our regeneration time course demonstrated that

sagittal amputations in *H. miamia* do not form a blastema, but instead slowly round up over the course of

70 hours (Figure 3B). To gain further insight into the cellular processes driving this morphogenetic

movement, we stained sagittal fragments with SiR-actin over the course of 70 hours post-amputation

(hpa) (Figure 4A). We observed that these animals healed through a two-step process: as described

above, the outer epidermis first connects to the pharyngeal epithelium within the first 10 hpa, followed by

the posterior end closing itself up through what appears to be a combination of epithelial crawling and

zippering. Both of these cellular movements are well-described in other systems, and here we see a

similar gradual closure of the outer epidermis as it meets and re-epithelializes with the epidermis from the

other side of the animal.

As the posterior ends are closing, sagittally-cut animals round their bodies up to re-form their

cylindrical shapes. The two processes we observexd (heterotypic bridges and homotypic re-

epithelialization) occur in a stepwise but overlapping sequence, where all fragments first form visible

bridges, then display both bridges and posterior closure, and finally end with just posterior re-

epithelialization (Figure 4B-C). All animals had an open body shape between 0-10 hpa, characterized by

a splayed pharynx and separated dorsal and ventral halves. At 20 hpa, we began to see rounding of the

fragments, with the outer epidermal edges curling over the pharynx towards each other. While some

animals closed as early as 30 hpa, virtually all animals completed this closure by 70 hpa (Figure 4B-D).

Here, closure is defined by the original dorsal and ventral edges completely meeting at the seam of the

wound, forming a cylinder with the pharynx on the inside and the epidermis on the outside. We speculate

that this rounding process is aiding the posterior re-epithelialization by bringing the wound edges

together.

To rule out that this dual closure mechanism was unique to the sagittal amputation, we next

asked whether similar processes occurred in transverse injuries that form heterotypic bridges. We

amputated animals both once (single transverse) and twice (donut) through the pharynx and tracked

regeneration. The single transverse fragments followed a similar regeneration trajectory as their

counterparts amputated below the pharynx (Figure 2B), although the differences in fragment size

correlated with different temporal dynamics (Supplemental Figure 3A). The donut fragments formed

blastemas on both anterior and posterior ends, regenerating all missing structures despite the severity of

the injury (Supplemental Figure 3B). To track the dynamics of wound closure in these amputations, we

fixed animals at intermediate time points and stained fragments with SiR-actin. Unlike the sagittal

fragments, the transverse fragments didn't zipper after the formation of bridges, but instead contracted

their outer epidermis over the wound site in a process that appeared similar to the single-epithelial

transverse amputations (Supplemental Figure 3C-D, Figure 2B).

These observations suggest that while the exact mechanism of wound closure is determined by a

combination of factors, injuries involving both epithelial layers follow a coordinated two-step process.

First, heterotypic protrusions bridge the wound gap between the two epithelial layers, followed by the

outer epidermis re-epithelializing with itself. We next asked how cell and tissue contractility was involved

in these mechanisms of wound closure, seeking to combine the characterization of these processes with

functional perturbations.

304 **Muscle contraction is necessary for both heterotypic wound closure and downstream**

305 **morphogenetic movements**

306 Contraction of muscle at the wound site is an essential component of wound closure in both vertebrates

307 and invertebrates, driving physical and biochemical events through the generation of mechanical

308 force[22,26,29,63]. Because of the close proximity of epithelial cells to underlying muscle in *H. miamia* (Figure

309 1G), we next asked whether muscle contraction was involved in wound closure. *H. miamia* have an

310 extensive network of muscle throughout their bodies (Figure 5A), and we utilized Tropomyosin

311 immunofluorescence to visualize how muscle was reorganized during single and dual epithelial wound

312 closure. In transverse wounds (both head and tail), muscle was largely absent from the wound site itself

313 but was instead present at the periphery of the wound edge as it shrank (Figure 5B). In contrast, sagittal

314 wounds contain parenchymal muscle fibers that link the outer epidermis to the pharyngeal epithelium,

315 coming together as the animal rounds up (Figure 5C). We hypothesized that this difference in muscle

316 organization might influence the closure mechanisms used by each fragment, and asked whether muscle

317 contraction was required for each type of wound closure.

318 To test the functional role of muscle contraction, we anesthetized amputated animals in 0.1%

319 tricaine, which blocks neural action potentials[64] and effectively inhibits muscle contraction in *H. miamia*

320 (Video 1). In the presence of tricaine, single epithelial wound closure in transverse amputations occurs at

321 the same rate as controls (Figure 5C-D), indicating that the epidermis is capable of constricting and re-

322 epithelializing without the aid of muscle contraction. Animals with multi-epithelial wounds, on the other

323 hand, are incapable of forming heterotypic bridges without muscle contraction (Figure 5E-G). We

324 observed some polarization and formation of small protrusions from the epidermis and pharyngeal

325 epithelium, but no bridges ever formed in tricaine-treated animals (Figure 5F). The requirement of muscle

326 contraction for the formation of heterotypic bridges was consistent across all multi-epithelial injuries

327 (Supplemental Figure 4A-B), suggesting a shared mechanism.

328 We performed time courses with tricaine treatment to determine whether muscle contraction was

329 only necessary for the initiation of the heterotypic bridges or whether muscle contraction was needed for

330 both their formation and maintenance (Supplemental Figure 4C). Animals treated with tricaine for 10

331 hours, then placed in sea water for 20 hours, closed wounds at a similar rate to controls. In contrast,

animals that were in sea water for 10 hours and then placed in tricaine for 20 hours were

indistinguishable from animals treated with tricaine for all 30 hours (Supplemental Figure 4D-E). This

indicates both that persistent muscle contraction is necessary for the maintenance of heterotypic bridges,

and that animals can fully recover after anesthetization.

Interestingly, muscle contraction is only required for the participation of the outer epidermis in

forming heterotypic bridges. In the presence of tricaine, the pharyngeal epithelium still formed extensions

and spread to try and cover the wound site (Supplemental Figure 4F-G). While the exact function is not

yet clear, these data implicate muscle contraction as a necessary driver of multi-epithelial wound closure.

This contraction may be involved in generating tension, inducing mechanical or biochemical signals,

and/or physically bringing the wound edges together to allow contact between the epithelia. Without

muscle-generated mechanical force, the outer epidermis cannot generate or maintain long protrusions to

form heterotypic bridges, and the wound closure process stalls and collapses.

Together, these findings highlight the importance of muscle contractility in mediating cellular

events during multi-epithelial wound closure. To distinguish between the role of tissue-scale mechanical

forces generated by muscle and the cell-intrinsic forces required for protrusive activity and wound closure,

we next assessed the role of actomyosin contractility in wound closure.

Types of wound closure are differentially regulated by actomyosin contractility

Actomyosin contractility is generated through the interactions of non-muscle myosin with actin, producing

mechanical forces within cells that enable cytoskeletal rearrangements and cellular movements[65–67].

While actomyosin contractility is broadly implicated in wound closure across species, we wanted to know

how its role varied across injury types in *H. miamia*. We treated amputated animals with the inhibitor

blebbistatin, which inhibits the ATPase activity of myosin and blocks actomyosin contractility[68,69]. In the

presence of blebbistatin, transverse wounds (which only disrupt the outer epidermis) closed at

comparable speeds to control conditions but appeared to switch towards more crawling-dominant

mechanisms (Figure 6A-C): instead of the contractile rings seen in control fragments, we detected

protrusions around the wound edge and saw individual cells crawling across the wound site (Figure 6C).

These data indicate that protrusion-based mechanisms are sufficient to accomplish wound closure in

blebbistatin-treated conditions.

We next asked how actomyosin contractility was involved in multi-epithelial wound closure. In

contrast to the mild phenotype in single epithelial wounds, we observed significant closure defects in the

blebbistatin-treated animals. Blebbistatin treatment eliminated the formation of stable heterotypic bridges

in sagittal fragments at 10 hpa (Figure 6D). We observed short protrusions emerging from both the outer

epidermis and pharyngeal epithelium, but their orientation appeared more random than control fragments,

forming multiple protrusions rather than connected bridges (Figure 6E, Supplemental Figure 5A). These

data indicate that actomyosin contractility is essential for the extension of long protrusions across the

wound edge to form heterotypic bridges. This contractility may be important in the stabilization of

protrusions as they extend, the application of pulling forces to bring the epithelia together, and/or the

stabilization of nascent adhesions at heterotypic interfaces[34,39,66,70–72]. Without this contractility, small

protrusions can form, but they ultimately lack the necessary force to form bridges.

Contrasting with the inhibition of heterotypic bridges, we observed an acceleration of epidermal

re-epithelialization in blebbistatin-treated animals. In control conditions, the zippering up of the posterior

end began around ~20-30 hours after amputation, after the formation of heterotypic bridges at the anterior

end (Figure 4D). In contrast, animals treated with blebbistatin showed posterior closure as early as 10

hpa, even in the absence of heterotypic bridges (Figure 6F-H). This phenotype is consistent with that

seen in other systems, in which the reduction of junctional tension 'loosens' the epithelial sheet and

allows epidermal cells to move more freely relative to each other[30,68,73]. By 20 hpa, blebbistatin-treated

animals showed robust closure of the outer epidermis, but no heterotypic bridges and no rounding of the

pharyngeal epithelium. Prolonged treatment with blebbistatin for 50 hours, however, led to reduced tissue

integrity in the epidermis, manifesting first as holes between cells and leading to eventual epidermal

ruptures (Figure 6H). The presumed decrease in junctional tension that enabled accelerated cell

spreading also prevented the maintenance of strong junctions between cells, resulting in a weakened

epidermal layer that was unable to withstand the mechanical demands of healing and regeneration.

Lastly, we assessed whether this phenotype was reversible and if the epidermis could re-

establish its integrity after accelerated re-epithelialization. We placed blebbistatin-treated animals in fresh

sea water after a 30-hour incubation for either 20 (50 hpa time point) or 40 (70 hpa time point) hours,

comparing directly to control animals and animals left in blebbistatin for the full time course

(Supplementary Figure 5B). Animals that were placed in fresh ASW for 20h were rounder than their

counterparts that remained in blebbistatin, suggesting a re-entry into the typical morphogenetic trajectory

(Supplementary Figure 5C-D). Animals that were placed in fresh ASW for 40 hours were able to

completely recover, resembling the control animals at 70 hpa (Supplementary Figure 5C-D). This

suggests that while prolonged blebbistatin treatment will lead to a loss of epidermal integrity, the animals

will restore junctional tension after removal of the drug, continuing the re- wound closure process.

DISCUSSION

In this study, we demonstrate that the acoel *Hofstenia miamia* uses context-dependent wound healing,

employing complementary and overlapping mechanisms of re-epithelialization to repair wounds prior to

whole-body regeneration. Wounds disrupting only one tissue use rapid and conserved closure

mechanisms, similar to wound closure in other species. In control conditions, these wounds primarily rely

on radial constriction of the epidermis, but in contractility-inhibited conditions the epidermal cells shift

mechanisms and close through crawling. Multi-epithelial wounds close through a combination of novel

heterotypic bridges and homotypic epithelial zippering, requiring both actomyosin-generated tension and

muscle contraction to drive coordinated tissue movements. In conditions of reduced global contractility,

other closure mechanisms are transiently upregulated, but ultimately fail to restore long-term tissue

integrity unless contractility is restored. Mechanisms of wound closure in *H. miamia* are therefore not

uniform or discrete processes, but are instead highly dynamic and cooperative, shifting in response to

both intrinsic and extrinsic factors.

The heterotypic bridges described here are, to the best of our knowledge, a novel wound-induced

response. Heterotypic contacts have been described in other systems as mechanisms by which tissue

boundaries are maintained during development[75–79], but filopodial matching that has been observed[79]

appears distinct from these bridges both in size and duration. Instead of fast filopodial probing between

proximal cells to form the proper adhesive complexes, the bridges observed in *H. miamia* are long (>50

um) and visible for at least 10 hours. At present, we cannot rule out the possibility that these bridges are

dynamic due to a lack of live imaging, but their length paired with the clear progression of formation

(polarization of actin, followed by short protrusions, then connected bridges) suggests some temporal

stability. Although they differ in presentation, it's possible that these heterotypic contacts are functioning

similarly to maintain tissue borders. During regeneration, injuries that damage both epithelial layers must

preserve both tissue identity and orientation, ultimately restoring the 'tube within a tube' anatomy.

Because we do not yet have tissue-specific labeling to differentiate between the epithelial layers, we don't

know how the bridges resolve during secondary closure. However, the framework presented here will set

the stage for future studies, including better visualization of cellular interactions, identification of adhesion

and migration machinery, the role of signaling pathways, and more.

When assessing the dynamics of these heterotypic interactions, we observed that the outer

epidermis requires muscle contraction to form stable protrusions, but the pharyngeal epithelium can

spread across the wound even in the absence of epidermal contributions. This phenotype could be due to

intrinsic differences in the epithelia; for example, perhaps the pharyngeal epithelium has intrinsic cell

motility machinery that is primed for migration regardless of external forces. In the animal's undamaged

configuration, the pharyngeal epithelium is surrounded by rings of pharyngeal muscle. It's possible that

when that muscle is damaged, the epithelial cells are now able to spread and crawl across the wound

site. In contrast, the outer epidermal cells may be more stable and require assistance from muscle

contraction to bring the two layers close enough to form stable adhesions. The differences between the

epithelial responses could also be due to their extrinsic interactions. The epithelial cells in *H. miamia* are

anatomically unique, lacking a flat basal surface as is typically observed in most animal epithelia and

instead forming long projections that end in close proximity to muscle. Electron microscopy in acoels has

failed to identify a visible basement membrane underneath the outer epidermis[74], but the composition of

extracellular matrix (ECM) in *H. miamia* has not been studied. It is therefore unclear whether the ECM is

adhering to protrusions in a punctate fashion, if there is ECM surrounding muscle fibers, if there is a thin

layer of ECM in between cellular protrusions, or any other organization. Using a custom antibody in this

study, we demonstrate that there is tissue-specific collagen underneath the pharyngeal epithelial cells,

and that collagen is present underneath their wound-induced protrusions. We did not observe this

collagen underneath the outer epidermis, but a similar mechanism could be occurring with a different

extracellular matrix protein, perhaps one whose remodeling is mechanosensitive. Differences in ECM

stiffness and composition could affect the mechanical response of each tissue, as well as differences in

regular tensile loads. It has also been shown in other systems that wound-induced muscle contraction

triggers the release of matrix remodeling proteins[42,59], which could be necessary for epidermal cells to

extend far enough to reach their pharyngeal epithelial counterparts. Future studies will need to further

characterize the *H. miamia* ECM and its dynamics to distinguish between these hypotheses.

The difference in wound closure strategies across injury types raises several questions about the

connection between healing and regeneration. It seems probable that different forms of wound closure

correlate with different cellular and molecular environments in regenerating fragments, whether through

biochemical and/or mechanical events associated with re-epithelialization. Future work will be required to

test whether different closure strategies have functional consequences for downstream tissue

regeneration, whether healing and regeneration are distinct and nonoverlapping biological processes, and

whether other species capable of whole-body regeneration utilize distinct mechanisms to heal different

types of wounds.

REFERENCES

1. Bely, A. E. & Nyberg, K. G. Evolution of animal regeneration: re-emergence of a field. *Trends Ecol. Evol.* **25**, 161–170 (2010).

2. Post, Y. & Clevers, H. Defining Adult Stem Cell Function at Its Simplest: The Ability to Replace Lost Cells through Mitosis. *Cell Stem Cell* **25**, 174–183 (2019).

3. Abreu-Blanco, M. T., Verboon, J. M., Liu, R., Watts, J. J. & Parkhurst, S. M. *Drosophila* embryos close epithelial wounds using a combination of cellular protrusions and an actomyosin purse string. *J. Cell Sci.* **125**, 5984–5997 (2012).

4. Fernandez-Gonzalez, R. & Zallen, J. A. Wounded cells drive rapid epidermal repair in the early *Drosophila* embryo. *Mol. Biol. Cell* **24**, 3227–3237 (2013).

5. Rothenberg, K. E. & Fernandez-Gonzalez, R. Forceful closure: cytoskeletal networks in embryonic wound repair. *Mol. Biol. Cell* **30**, 1353–1358 (2019).

6. Begnaud, S., Chen, T., Delacour, D., Mège, R.-M. & Ladoux, B. Mechanics of epithelial tissues during gap closure. *Curr. Opin. Cell Biol.* **42**, 52–62 (2016).

7. Lim, S. E., Vicente-Munuera, P. & Mao, Y. Forced back into shape: Mechanics of epithelial wound repair. *Curr. Opin. Cell Biol.* **87**, 102324 (2024).

8. Jacinto, A., Martinez-Arias, A. & Martin, P. Mechanisms of epithelial fusion and repair. *Nat. Cell Biol.* **3**, E117–E123 (2001).

9. Chablais, F. & Jaźwińska, A. IGF signaling between blastema and wound epidermis is required for fin regeneration. *Development* **137**, 871–879 (2010).

10. Scimone, M. L., Cloutier, J. K., Maybrun, C. L. & Reddien, P. W. The planarian wound epidermis gene equinox is required for blastema formation in regeneration. *Nat. Commun.* **13**, 2726 (2022).

11. Aztekin, C. Tissues and Cell Types of Appendage Regeneration: A Detailed Look at the Wound Epidermis and Its Specialized Forms. *Front. Physiol.* **12**, 771040 (2021).

12. Aztekin, C. *et al.* Identification of a regeneration-organizing cell in the *Xenopus* tail. *Science* **364**, 653–658 (2019).

13. Bideau, L., Kerner, P., Hui, J., Vervoort, M. & Gazave, E. Animal regeneration in the era of transcriptomics. *Cell. Mol. Life Sci.* **78**, 3941–3956 (2021).

14. Amiel, A. *et al.* Characterization of Morphological and Cellular Events Underlying Oral Regeneration

in the Sea Anemone, Nematostella vectensis. *Int. J. Mol. Sci.* **16**, 28449–28471 (2015).

15. Ben Khadra, Y. *et al.* Wound repair during arm regeneration in the red starfish *E chinaster sepositus*.

Wound Repair Regen. **23**, 611–622 (2015).

16. Cheung, S. *et al.* Systemic coordination of whole-body tissue remodeling during local regeneration in

sea anemones. *Dev. Cell* S153458072400666X (2024) doi:10.1016/j.devcel.2024.11.001.

17. Lust, K. *et al.* Single-cell analyses of axolotl telencephalon organization, neurogenesis, and

regeneration. *Science* **377**, eabp9262 (2022).

18. Kassmer, S. H., Nourizadeh, S. & De Tomaso, A. W. Cellular and molecular mechanisms of

regeneration in colonial and solitary Ascidians. *Dev. Biol.* **448**, 271–278 (2019).

19. Kamran, Z. *et al.* In vivo imaging of epithelial wound healing in the cnidarian Clytia hemisphaerica

demonstrates early evolution of purse string and cell crawling closure mechanisms. *BMC Dev. Biol.*

17, 17 (2017).

20. Ramon-Mateu, J., Ellison, S. T., Angelini, T. E. & Martindale, M. Q. Regeneration in the ctenophore

Mnemiopsis leidyi occurs in the absence of a blastema, requires cell division, and is temporally

separable from wound healing. *BMC Biol.* **17**, 80 (2019).

21. Spiegelman, M. & Dudley, P. L. Morphological stages of regeneration in the planarian *Dugesia*

tigrina : A light and electron microscopic study. *J. Morphol.* **139**, 155–183 (1973).

22. Shaw, T. J., Osborne, M., Ponte, G., Fiorito, G. & Andrews, P. L. R. Mechanisms of wound closure

following acute arm injury in Octopus vulgaris. *Zool. Lett.* **2**, 8 (2016).

23. Donaldson, D. J. & Dunlap, M. K. Epidermal cell migration during attempted closure of skin wounds in

the adult newt: Observations based on cytochalasin treatment and scanning electron microscopy. *J.*

Exp. Zool. **217**, 33–43 (1981).

24. Huang, T. *et al.* Re-epithelialization of large wound in paedomorphic and metamorphic axolotls. *J.*

Morphol. **278**, 228–235 (2017).

25. Zattara, E. E., Turlington, K. W. & Bely, A. E. Long-term time-lapse live imaging reveals extensive cell

migration during annelid regeneration. *BMC Dev. Biol.* **16**, 6 (2016).

26. Chandebois, R. THE DYNAMICS OF WOUND CLOSURE AND ITS ROLE IN THE PROGRAMMING

OF PLANARIAN REGENERATION. II - DISTALIZATION. *Dev. Growth Differ.* **22**, 693–704 (1980).

27. Belacortu, Y. & Paricio, N. *Drosophila* as a model of wound healing and tissue regeneration in

vertebrates. *Dev. Dyn.* **240**, 2379–2404 (2011).

28. Brewer, C. M. *et al.* Adaptations in Hippo-Yap signaling and myofibroblast fate underlie scar-free ear

appendage wound healing in spiny mice. *Dev. Cell* **56**, 2722-2740.e6 (2021).

29. Valero, C., Javierre, E., García-Aznar, J. M. & Gómez-Benito, M. J. A Cell-Regulatory Mechanism

Involving Feedback between Contraction and Tissue Formation Guides Wound Healing Progression.

PLoS ONE **9**, e92774 (2014).

30. Tetley, R. J. *et al.* Tissue fluidity promotes epithelial wound healing. *Nat. Phys.* **15**, 1195–1203 (2019).

31. Bement, W. M., Mandato, C. A. & Kirsch, M. N. Wound-induced assembly and closure of an

actomyosin purse string in Xenopus oocytes. *Curr. Biol.* **9**, 579–587 (1999).

32. Nobes, C. D. & Hall, A. Rho, Rac, and Cdc42 GTPases regulate the assembly of multimolecular focal

complexes associated with actin stress fibers, lamellipodia, and filopodia. *Cell* **81**, 53–62 (1995).

33. Henson, J. H. *et al.* Wound Closure in the Lamellipodia of Single Cells: Mediation by Actin

Polymerization in the Absence of an Actomyosin Purse String. *Mol. Biol. Cell* **13**, 1001–1014 (2002).

34. Tang, V. W. Cell–cell adhesion interface: orthogonal and parallel forces from contraction, protrusion,

and retraction. *F1000Research* **7**, 1544 (2018).

35. Da Costa, A. D. S. *et al.* Fibroblasts Close a Void in Free Space by a Purse-String Mechanism. *ACS*

Appl. Mater. Interfaces **14**, 40522–40534 (2022).

36. Rodriguez-Diaz, A. *et al.* Actomyosin purse strings: Renewable resources that make morphogenesis

robust and resilient. *HFSP J.* **2**, 220–237 (2008).

37. Ehrlich, H. P. Wound closure: evidence of cooperation between fibroblasts and collagen matrix. *Eye*

2, 149–157 (1988).

38. Basan, M., Elgeti, J., Hannezo, E., Rappel, W.-J. & Levine, H. Alignment of cellular motility forces

with tissue flow as a mechanism for efficient wound healing. *Proc. Natl. Acad. Sci.* **110**, 2452–2459

(2013).

39. Vedula, S. R. K. *et al.* Mechanics of epithelial closure over non-adherent environments. *Nat.*

Commun. **6**, 6111 (2015).

40. Xu, H. *et al.* Geometry-mediated bridging drives nonadhesive stripe wound healing. *Proc. Natl. Acad.*

Sci. **120**, e2221040120 (2023).

41. Pascolini, R., Tei, S., Vagnetti, D. & Bondi, C. Epidermal cell migration during wound healing in

Dugesia lugubris: Observations based on scanning electron microscopy and treatment with

cytochalasin. *Cell Tissue Res.* **236**, (1984).

42. Chakraborty, S. *et al.* Agrin-Matrix Metalloproteinase-12 axis confers a mechanically competent

microenvironment in skin wound healing. *Nat. Commun.* **12**, 6349 (2021).

43. Brauer, E. *et al.* Collagen Fibrils Mechanically Contribute to Tissue Contraction in an In Vitro Wound

Healing Scenario. *Adv. Sci.* **6**, 1801780 (2019).

44. Yannas, I. V. & Tzeranis, D. S. Mammals fail to regenerate organs when wound contraction drives

scar formation. *Npj Regen. Med.* **6**, 39 (2021).

45. Kimura, S. & Tsuji, T. Mechanical and Immunological Regulation in Wound Healing and Skin

Reconstruction. *Int. J. Mol. Sci.* **22**, 5474 (2021).

46. Nier, V. *et al.* Tissue fusion over nonadhering surfaces. *Proc. Natl. Acad. Sci.* **112**, 9546–9551 (2015).

47. Srivastava, M., Mazza-Curll, K. L., van Wolfswinkel, J. C. & Reddien, P. W. Whole-Body Acoel

Regeneration Is Controlled by Wnt and Bmp-Admp Signaling. *Curr. Biol.* **24**, 1107–1113 (2014).

48. Raz, A. A., Srivastava, M., Salvamoser, R. & Reddien, P. W. Acoel regeneration mechanisms indicate

an ancient role for muscle in regenerative patterning. *Nat. Commun.* **8**, 1260 (2017).

49. Srivastava, M. Studying development, regeneration, stem cells, and more in the acoel Hofstenia

miamia. in *Current Topics in Developmental Biology* vol. 147 153–172 (Elsevier, 2022).

50. Hulett, R. E. *et al.* Acoel single-cell atlas reveals expression dynamics and heterogeneity of adult

pluripotent stem cells. *Nat. Commun.* **14**, 2612 (2023).

51. Reddien, P. W. Positional Information and Stem Cells Combine to Result in Planarian Regeneration.

Cold Spring Harb. Perspect. Biol. a040717 (2021) doi:10.1101/cshperspect.a040717.

52. Ricci, L. & Srivastava, M. Transgenesis in the acoel worm Hofstenia miamia. *Dev. Cell* **56**, 3160-

3170.e4 (2021).

53. Doerr, S., Zhou, P. & Ragkousi, K. Origin and development of primary animal epithelia. *BioEssays* **46**,

2300150 (2024).

54. Friedl, P. & Gilmour, D. Collective cell migration in morphogenesis, regeneration and cancer. *Nat.*

Rev. Mol. Cell Biol. **10**, 445–457 (2009).

55. Macabenta, F. & Stathopoulos, A. Sticking to a plan: adhesion and signaling control spatial

organization of cells within migrating collectives. *Curr. Opin. Genet. Dev.* **57**, 39–46 (2019).

56. Patel, A. *et al.* Cytonemes coordinate asymmetric signaling and organization in the Drosophila

muscle progenitor niche. *Nat. Commun.* **13**, 1185 (2022).

57. Daly, C. A., Hall, E. T. & Ogden, S. K. Regulatory mechanisms of cytoneme-based morphogen

transport. *Cell. Mol. Life Sci.* **79**, 119 (2022).

58. Caviglia, S. & Ober, E. A. Non-conventional protrusions: the diversity of cell interactions at short and

long distance. *Curr. Opin. Cell Biol.* **54**, 106–113 (2018).

59. Sakar, M. S. *et al.* Cellular forces and matrix assembly coordinate fibrous tissue repair. *Nat.*

Commun. **7**, 11036 (2016).

60. Naba, A. Mechanisms of assembly and remodelling of the extracellular matrix. *Nat. Rev. Mol. Cell*

Biol. **25**, 865–885 (2024).

61. Ehrlich, H. P. & Hunt, T. K. Collagen Organization Critical Role in Wound Contraction. *Adv. Wound*

Care **1**, 3–9 (2012).

62. Xue, M. & Jackson, C. J. Extracellular Matrix Reorganization During Wound Healing and Its Impact

on Abnormal Scarring. *Adv. Wound Care* **4**, 119–136 (2015).

63. Li, B. & Wang, J. H.-C. Fibroblasts and myofibroblasts in wound healing: Force generation and

measurement. *J. Tissue Viability* **20**, 108–120 (2011).

64. Attili, S. & Hughes, S. M. Anaesthetic Tricaine Acts Preferentially on Neural Voltage-Gated Sodium

Channels and Fails to Block Directly Evoked Muscle Contraction. *PLoS ONE* **9**, e103751 (2014).

65. Brunet, T. Cell contractility in early animal evolution. *Curr. Biol.* **33**, R966–R985 (2023).

66. Röper, K. Integration of Cell–Cell Adhesion and Contractile Actomyosin Activity During

Morphogenesis. in *Current Topics in Developmental Biology* vol. 112 103–127 (Elsevier, 2015).

67. Weißenbruch, K. & Mayor, R. Actomyosin forces in cell migration: Moving beyond cell body retraction.

BioEssays **46**, 2400055 (2024).

68. Liu, Z. *et al.* Blebbistatin inhibits contraction and accelerates migration in mouse hepatic stellate

cellsbph_477304..315. *Br. J. Pharmacol.* (2010).

69. Holz, O., Apel, D. & Hassel, M. Alternative pathways control actomyosin contractility in

epitheliomuscle cells during morphogenesis and body contraction. *Dev. Biol.* **463**, 88–98 (2020).

70. Li, J. X. H., Tang, V. W., Boateng, K. A. & Brieher, W. M. Cadherin puncta are interdigitated dynamic

actin protrusions necessary for stable cadherin adhesion. *Proc. Natl. Acad. Sci.* **118**, e2023510118

(2021).

71. Padhi, A. *et al.* Force-exerting perpendicular lateral protrusions in fibroblastic cell contraction.

Commun. Biol. **3**, 390 (2020).

72. Miroshnikova, Y. A. *et al.* Adhesion forces and cortical tension couple cell proliferation and

differentiation to drive epidermal stratification. *Nat. Cell Biol.* **20**, 69–80 (2018).

73. Cartagena-Rivera, A. X., Van Itallie, C. M., Anderson, J. M. & Chadwick, R. S. Apical surface

supracellular mechanical properties in polarized epithelium using noninvasive acoustic force

spectroscopy. *Nat. Commun.* **8**, 1030 (2017).

74. Dorey, A. E. The organization and replacement of the epidermis in acoelous turbellarians. *J. Cell Sci.*

S3-106, 147–172 (1965).

75. Batlle, E. & Wilkinson, D. G. Molecular Mechanisms of Cell Segregation and Boundary Formation in

Development and Tumorigenesis. *Cold Spring Harb. Perspect. Biol.* **4**, a008227–a008227 (2012).

76. Moore, R. E., Clarke, J. & Alexandre, P. Protrusion-Mediated Signaling Regulates Patterning of the

Developing Nervous System. *Front. Cell Dev. Biol.* **8**, 579073 (2020).

77. Katsunuma, S. *et al.* Synergistic action of nectins and cadherins generates the mosaic cellular

pattern of the olfactory epithelium. *J. Cell Biol.* **212**, 561–575 (2016).

78. Canty, L., Zarour, E., Kashkooli, L., François, P. & Fagotto, F. Sorting at embryonic boundaries

requires high heterotypic interfacial tension. *Nat. Commun.* **8**, 157 (2017).

79. Zhang, S., Amourda, C., Garfield, D. & Saunders, T. E. Selective Filopodia Adhesion Ensures Robust

Cell Matching in the Drosophila Heart. *Dev. Cell* **46**, 189-203.e4 (2018).

80. Hulett, R. E., Potter, D. & Srivastava, M. Neural architecture and regeneration in the acoel *Hofstenia*

miamia. *Proc. R. Soc. B Biol. Sci.* **287**, 20201198 (2020).

MATERIALS AND METHODS

H. miamia strains and husbandry

Adult worms were housed in artificial sea water (ASW) (37 ppt, pH 7.70-8.00) in plastic boxes within 21°C

incubators. Embryos were collected from adult boxes and placed into petri dishes for development. Once

hatched, juvenile worms were transferred to tanks for post-embryonic growth. All animals were fed twice a

week; adult worms were fed brine shrimp (Artemia, Brine Shrimp Direct) and juvenile hatchlings were fed

rotifers (*Brachionus plicatilis*, Reed Mariculture). *EpiA:RFP* worms were generated as previously

described[52].

Amputations and sample preparation

Juvenile worms were isolated and starved for 1-2 weeks prior to use to minimize autofluorescence of

ingested food. For regeneration experiments, animals were placed in 1mg/mL ethyl 3-aminobenzoate,

methanesulfonic acid ("tricaine", stock solution 5mg/mL of ASW) (Thermo Fisher, AC118000100) in ASW

in a petri dish for 3-5 minutes until they stopped swimming, then amputated using a scalpel. After

amputation, fragments were transferred to a dish containing fresh ASW until their designated time point.

For fixation, animals were placed in 4% paraformaldehyde (PFA) (Thomas Scientific, C993M31) in ASW

for 1 hour at room temperature. The PFA was removed and animals were washed with PBST (PBS +

0.1% Triton-X-100) three times prior to immunofluorescence staining. For pinch wounds, animals were

grasped with fine-tip forceps, then allowed to recover in ASW prior to fixation. For transverse and donut

cuts, animals were cut once or twice transversely, then transferred to fresh ASW until their designated

time point. For single transverse wounds, animals were cut transversely once again, right above the

previous cut, to remove the wound site for imaging. These fragments were then immediately fixed in 4%

PFA in ASW.

Small molecule inhibitors and treatment with anesthetics

For small molecule inhibitor experiments, animals were first amputated as described above, then flushed

with ASW to remove tricaine prior to incubation in designated inhibitors. Animals were left in inhibitor for

3-70 hours, depending on the experiment. Blebbistatin (Sigma-Aldrich, B0560) was dissolved in DMSO

per manufacturer instructions, then diluted in ASW for use at 25µM. DMSO control treatments used an

equal volume of DMSO as Blebbistatin treatments. For tricaine time point experiments, animals were left

in 0.5mg/mL tricaine in ASW for 3-50 hours.

Immunostaining

For antibody staining, all steps were carried out using an orbital shaker. Fixed animals were washed in

PBST for 3x5 minutes, then placed in blocking solution (PBST + 10% goat serum (Thermo Fisher,

16210072)) for one hour at room temperature. Animals were then placed in primary antibody in block

solution for two nights at 4°C. Primary antibody was washed out with PBST, 6x20 minutes. Animals were

then blocked once again for one hour at room temperature, then incubated with secondary antibody

(1:800) (Thermo Fisher, A-11011) in blocking solution overnight at 4°C. The next day, secondary antibody

was washed out using PBST, 6x20 minutes. Finally, animals were incubated in SiR-actin (1:1000) in

PBST (Cytoskeleton, CY-SC001) for 90 minutes at room temperature prior to mounting. All samples were

mounted on double cover slips using Prolong Glass Antifade Mountant (Thermo Fisher, P36980). If

animals were only stained with SiR-actin, they were placed in SiR-actin in PBST for 90 minutes at room

temperature, then mounted immediately. The custom *H. miamia* Tropomyosin antibody[80] (Rabbit IgG)

was used at 1µg/mL Where relevant, Hoechst (1:500) (Invitrogen, H3570) was used to counter-stain

nuclei.

Generation of custom collagen IV antibody

Rabbit polyclonal antibodies were custom-made by GenScript through their PolyExpress immunization

protocol, using a peptide sequence from *H. miamia* Collagen IV (CPPGAKGNDGEPGEKG). This peptide

was expressed in *E. coli* using the provided sequence and then used to immunize two rabbits. Two

antibodies were generated, purified, and validated in-house by Western Blot with the protein immunogen.

Microscopy and image processing

All images were collected using 10x/0.25x (dry), 20x/0.75x (dry), or 63x/1.40 (oil) fluorescent objectives

on a Leica DMI SP8 inverted confocal microscope, equipped with Leica Application Suite software.

Images were processed using Image J/FIJI software.

Quantifications

Length of epithelial basal protrusions

To quantify the average length of basal protrusions in outer epidermal and pharyngeal epithelial cells,

images were taken of labeled cells in *EpiA:RFP* animals. The Measure function in FIJI was used to

quantify the length of each protrusion, tracing from the end of the protrusion to where it meets the body of

the cell. For each cell type (pharyngeal epithelium vs. outer epidermis), n=15 cells from n=5 animals (total

= 45 cells) was measured.

Percentage of closure in transverse amputations

For quantification of wound closure in transverse cuts, images were taken of both the wound edge and

the open edge of each sample, using FIJI to measure the open area of each side. The percentage of

wound closure was determined by dividing the open area of the wound edge by the total open area, then

subtracting that from 1. For each time point and fragment (head vs. tail), measurements were collected

from n=8 animals.

Percentage of closure in sagittal amputations

To quantify the percentage of wound closure after sagittal amputations, images of SiR-actin-stained

fragments were imported into FIJI. Because of mounting imperfections, only one side of each animal was

measured, choosing the side with the clearest signal for the most accurate quantification. 6-12 animals

were quantified per time point and condition. Within FIJI, the Measure function was used to quantify: 1)

the length of the pharynx and 2) the cumulative distance of connected tissue between the epidermis and

the pharyngeal epithelium. The percentage of wound closure was determined by dividing the closed

distance by the total pharynx length. For each time point, measurements were collected from n=12

animals.

Bridge length

To quantify the length of heterotypic bridges, images of SiR-actin-stained fragments were taken at each

time point, then imported into FIJI for analysis. Using the Measure function, bridges were quantified by

drawing a line along the membrane extensions protruding between cell types. Cell bodies were not

included in this measurement. Because of the technical challenge of robustly identifying where one

protrusion met the other, the combined bridge length was reported instead of individual cell contributions.

For each time point, measurements were collected from n=30-40 animals.

Morphological descriptions

To report the morphological characteristics of healing fragments, images of SiR-actin-stained fragments

were taken at each time point. For bridge formation, animals were binned into four categories: none (no

visible polarization of the epithelial layers and no protrusions), polarization (visible polarization of cell

membranes towards the wound edge), protrusions (at least one membrane extension <15um in length),

or bridges (at least one connected heterotypic bridge). For protrusion origin, animals were assessed for

visible protrusions coming from only the outer epidermis, only the pharyngeal epithelium, both, or neither.

Rounding was assessed by looking at the shape of the animal, binning animals into either open (no

visible rounding), rounding (partway through the rounding process), or closed (where the outer edges of

the fragments have met, re-forming a cylindrical body plan). The values plotted for each morphological

trait were the representative percentage of total fragments, ranging from n=10-45 animals (exact values

are reported in accompanying figure legends).

Graphing and statistical analyses

All graphs were generated using GraphPad PRISM. Data with two or more factors were analyzed using

two-way ANOVA with Dunnett's multiple comparisons test. Data comparing changes in one condition over

different time points were analyzed by one-way ANOVA with Dunnett's multiple comparisons test. A two-

tailed unpaired t-test with Welch's correction was used to compare data between two independent

groups. All statistical tests are stated in figure legends. P values were obtained using GraphPad PRISM

and used to determine the level of varying significance between experimental groups; groups were

considered different when $P<0.05$.

ACKNOWLEDGEMENTS

APK is a Fellow of the Jane Coffin Childs Memorial Fund for Medical Research. This project was

supported by grants from the National Institute of General Medical Sciences of the NIH to MS:

R35GM128817 and R35GM153252. Thanks to members of the Srivastava Lab for feedback on the

manuscript.

AUTHOR CONTRIBUTIONS

A.P.K conceived the project, performed experiments, quantified and analyzed data, and wrote the

manuscript under the supervision of M.S. Both authors discussed the results and edited the manuscript.

COMPETING INTERESTS

None declared.

     FIGURES AND FIGURE LEGENDS

Figure 1. *Hofstenia miamia* have two distinct, atypical epithelial tissues

(A) A juvenile *Hofstenia miamia.* (B) A simplified phylogenetic tree showing the position of *Hofstenia*

miamia (red, Xenoacoelomorpha) relative to other clades. mya = million years ago. (C) *EpiA:RFP*

transgenic worms, fluorescently labeling the outer epidermis and the pharyngeal epithelium in a mosaic

manner. (D) Morphology of individual epidermal and pharyngeal (Ph.) epithelial cells, visualized by

EpiA:RFP transgenic labeling. A: apical, B: basal. (E-F) SiR-actin staining showing the apical organization

of *H. miamia* epithelial tissues, particularly the polygonal shape of each cell and the presence of actin at

cell-cell junctions (E) and the presence of actin within epidermal basal protrusions (F). (G) Quantification

of basal protrusions within epidermal and pharyngeal epithelial cells. Data represent n=45 cells per cell

type, pooled from 15 worms. Comparison by unpaired *t*-test. (H) Confocal images showing the proximity

of muscle (visualized through immunofluorescence staining for tropomyosin) to both epithelial layers

(visualized with EpiA:RFP transgenic labeling). Scale bars: 50 µm (A), 100 µm (C), 5µm (D), 20 µm (E,

H), 25µm (F), 10 µm (H, zoomed-in box).

Figure 2. Transverse amputations close through gradual constriction of the epidermis across the

wound edge.

(A-B) Schematic (A) and representative brightfield time course (B) showing regeneration of *H. miamia*

following transverse amputation. Red arrowhead indicates the formation of a pointed tail, white arrowhead

indicates a visible blastema. (C) Representative images of wound closure following transverse

amputation between 0-10 hours post-amputation. Red asterisk indicates the base of the pharynx, visible

in open wounds. The outer epidermis is pseudocolored in pink for clarity. (D) Quantification of wound

closure following transverse amputation, comparing between head and tail fragments from 3-10 hours

post-amputation. Data represent 8 worms per time point, and comparisons were performed by 2-way

ANOVA with Šidák's multiple comparisons test; ns=not significant. (E) High-magnification images

showing continued dynamics of re-epithelialization following initial closure; yellow arrows indicate a visible

seam where the tissue layers were brought together, but seams are no longer visible by 10 hours later.

hpa = hours post-amputation. Scale bars: 50µm (B), 100 µm (C), 25 µm (E).

Figure 3. Injuries that damage both epithelial tissues form long, heterotypic protrusions

connecting the epithelial layers across the wound edge

(A-B) Schematic (A) and representative brightfield time course (B) showing regeneration of *H. miamia*

following sagittal amputation. (C) Representative images of wound closure following sagittal amputation

between 0-10 hours post-amputation. (D) High-magnification images showing polarization, protrusion,

and formation of heterotypic bridges during the first 10 hours of wound closure. (E-G) Quantifications of

heterotypic bridge formation (E), origin of protrusions (F), and combined bridge length (G) over the first 10

hpa. Data represent 8 worms per condition and time point (E-F) and 30-40 bridges per time point (G);

comparison by unpaired *t*-test. (H-I) Representative images demonstrating the presence of heterotypic

bridges in single (H) and double (I) transverse amputations through the pharynx. In all high-magnification

images, the outer epidermis is pseudocolored in pink and the pharyngeal epithelium is pseudocolored in

blue for clarity. Scale bars: 50µm (B-C, H-I), 10 µm (C, zoomed-in box), 5 µm (D), 20 µm (H-I, zoomed-in

box).

Figure 4. Sagittal wounds heal through a two-step process, combining epithelialization with whole-body rounding to restore the fragment's original morphology.

(A) Schematic and time course showing epithelial tissue dynamics in sagittal wound closure during the first 70 hours post-amputation (hpa), using SiR-actin labeling. (B) Higher-magnification images demonstrating the overlapping mechanisms of epithelialization in the anterior and posterior parts of the

fragment. Anterior tissues form heterotypic protrusions described in Figure 3, and the posterior wound

edges gradually close through homotypic re-epithelialization. (C-D) Quantifications of closure

mechanisms (C) and fragment rounding (D). Data represent n=12 animals per time point. hpa = hours

post-amputation. Scale bars: 100 μm (A), 20 μm (B).

Figure 5. Muscle contraction is dispensable for traditional transverse amputations but required for

the formation of heterotypic bridges during sagittal wound closure.

(A-C) Images showing the organization of muscle in uncut worms (A), transverse wound closure (B), and

sagittal wound closure (C), visualized by immunofluorescence staining for tropomyosin. (D-E)

Representative images (D) and quantification (E) of wound closure in control and tricaine-treated animals

after transverse amputation, labeled with SiR-actin. (F) Representative images of wound closure in

control and tricaine-treated animals after sagittal amputation, labeled with SiR-actin.

In D and F, the outer epidermis is pseudocolored in pink and the pharyngeal epithelium is pseudocolored

in blue for clarity. (G-H) Quantification of closure mechanism (G) and fragment rounding (H) in control and

tricaine-treated animals after sagittal amputation. Data represent n=12 animals per condition, per time

point; quantification in (E) by unpaired *t*-test. ns=not significant, hpa = hours post amputation. Scale bars:

100 µm (A,C), 50 µm (B,D,F).

Figure 6. Actomyosin contractility is differentially utilized across wound closure mechanisms.

(A-B) Representative images (A) and quantification (B) of wound closure in control and blebbistatin-

treated animals after transverse amputation, labeled with SiR-actin. (C) High-magnification images

showing the difference between control and blebbistatin-treated wound edges in transverse amputations.

(D) Representative images of wound closure in control and blebbistatin-treated animals after sagittal

amputation. (E) High-magnification images showing heterotypic bridges in control animals and the lack

thereof in blebbistatin-treated animals. (F-G) Quantification of closure mechanism (F) and fragment

rounding (G) in control and blebbistatin-treated animals after sagittal amputation. (H) In high-

magnification images, the outer epidermis is pseudocolored in pink and the pharyngeal epithelium is

pseudocolored in blue for clarity. Data represent n=12-16 animals per condition, per time point;

comparison in (B) by unpaired t-test. ns=not significant, hpa = hours post amputation. Scale bars: 50 µm

(A,D), 10 µm (C,E), 100 µm (H), 20 µm (H, zoomed-in box).

SUPPLEMENTARY FIGURES AND TABLES

Supplementary Figure 1. Pinch wounds heal rapidly, utilizing cell protrusions to bring the wound

edges together.

(A) Schematic and representative images showing wound closure following an epidermal pinch injury,

labeled with SiR-actin. Scale bars: 100 µm (A, top row), 10 µm (A, zoomed-in box). mpa = minutes post

amputation.

Supplementary Figure 2. Collagen IV is visible underneath pharyngeal, but not epidermal,

protrusions during wound closure.

(A-B) *EpiA:RFP* transgenic animals cut in the sagittal plane have mosaic expression of RFP in both the

epidermis and the pharyngeal epithelium (A), and these cells are part of the SiR-actin-labeled population

that is forming protrusions during wound closure (B). (C-F) Immunofluorescence staining using a custom

Collagen IV antibody (see Methods) showing localization underneath the pharyngeal epithelium in uncut

(C) and 10 hours post-amputation (hpa) (D-F) fragments. Scale bars: 50µm (A, C (left), D, E-F (top)),

10µm (B), 20µm (C (right), E-F (bottom)).

Supplementary Figure 3. Transverse amputations that injure the pharyngeal epithelium heal through a similar two-step mechanism.

(A-B) Schematic and representative brightfield time course showing regeneration of *H. miamia* following single (A) and double (B) transverse amputation. (C-D) Representative images showing wound closure following single (C) and double (D) transverse amputation, labeled with SiR-actin. Scale bars: 100μm (A-B), 50μm (C-D).

Supplementary Figure 4. Muscle contraction is required for epidermal protrusions during

heterotypic bridge formation.

(A-B) Representative images showing pharyngeal protrusions after single (A) or double (B) transverse

amputation in tricaine-treated conditions. (C) Schematic of tricaine timing experiment. (D) Representative

images showing wound closure in control, tricaine, 10h tricaine (+ 20h ASW), or 10h ASW (+ 20h tricaine)

conditions. (E) Quantification of wound closure across tricaine conditions. Data represent n=8 worms per

condition and show mean ± s.e.m. Comparisons by one-way ANOVA with Dunnett's multiple comparisons

test; ns=not significant. (F) Representative images showing Collagen IV localization underneath

pharyngeal extensions following transverse amputation in control and tricaine-treated conditions. (G)

Quantification of protrusion origin in control and tricaine-treated animals. Data represent n=7-10 animals

per condition, per time point. Scale bars: 100μm (A,B), 50μm (A,B (zoomed-in boxes), D,F).

Supplementary Figure 5. Blebbistatin treatment is reversible.

(A) Quantification of protrusion origin in control and blebbistatin-treated animals. (B) Schematic of

blebbistatin wash-out experiment. (C-D) Quantification (C) and representative images (D) of fragment

rounding following control and blebbistatin treatments. Data represent n=8 animals per condition, per time

point. Scale bars: 100µm (D).

871   **Supplementary Video 1. Tricaine slows muscle contraction in *H. miamia*.**

872   Animals are anesthetized in 1mg/mL tricaine in ASW.