Figure 1

Figure 1. Lack of pectoral fins in *hoxba;hoxbb* cluster-deleted mutants.
(A-G) Dorsal views of live zebrafish larvae at 3 dpf, obtained from
intercrosses between *hoxba;hoxbb* cluster-deficient hemizygous fish.
Arrowheads indicate the positions of pectoral fins. (H-M) Expression
patterns of *tbx5a* in the pectoral fin bud (arrowhead) at 30 hpf. Dorsal views
are shown capturing images, the genotype of each specimen was determined.
For each genotype, reproducibility was confirmed with at least three
different specimens. Genotyping revealed that all the embryos lacking
pectoral fins (*n*=15) were *hoxba;hoxbb* double homozygotes. Scale bars are
100 μm.

Figure 2. Significantly decreased expression of *tbx5a* in the pectoral fin buds is specific to *hoxba;hoxbb* cluster-deleted mutants.

(A-I) Expression patterns of *tbx5a* in the pectoral fin buds of combinatorial deletion mutants of zebrafish *hox* clusters. Dorsal views of embryos at 30 hpf are displayed. After capturing images, the genotype of each specimen was determined. For each genotype, reproducibility was confirmed with at least three different specimens. Scale bars are 100 μm.

Figure 3. Transformation of pectoral fin progenitor cells into cardiac cells in *hoxba;hoxbb* homozygous embryos. (A-F) Expression patterns of *tbx5a* were compared between sibling wild-type and *hoxba;hoxbb* homozygous embryos during embryogenesis. The range of *tbx5a* expression in the lateral mesoderm is indicated by a bracket. The progenitor cells of pectoral fins are indicated by an arrowhead. (G, H) Expression patterns of *nkx2.5* were compared between sibling and *hoxba;hoxbb* mutants. Enlarged views are shown in (G', H'). The different regions of *nkx2.5* expression between the wild type and the mutant are indicated by brackets. (I, J) Expression patterns of *cmcl2* are shown. All images were captured from the dorsal side. Scale bars indicate 100 μm.

Figure 4. Zebrafish *hoxba;hoxbb* mutants lack a response to RA in fin buds.
(A-D) Exogeneous RA exposure in *raldh2* mutants can rescue *tbx5a* expression
in fin buds. Arrowheads indicate the positions of the pectoral fin buds. (E-H)
RA treatments in *hoxba;hoxbb* cluster-deleted mutants do not rescue *tbx5a*
expression in fin buds. (I, J) Expression patterns of *raldh2* were compared
between sibling and *hoxba;hoxbb* mutants. All images were captured from the
dorsal side. Scale bars represent 100 μm.

Figure 5. The expression patterns of *hox5a* and *hoxb5b* are regulated by RA.
(A) Schematic representation of the 49 *hox* genes organized into seven *hox* clusters in
zebrafish. *hox* genes in *hoxba* and *hoxbb* clusters are highlighted with orange. (B)
Expression profiles of the 49 *hox* genes in wild-type and *raldh2*[-/-] embryos at the 20-
somite stage, analyzed by RNA-seq. The average FPKM of each *hox* gene was
compared between sibling and *raldh2* mutants. The absence of specific *hox* genes is
indicated by a slash. (C-H) Expression patterns of *hoxb5a* and *hoxb5b* in sibling wild-
type, *raldh2*[-/-], and RA-treated *raldh2*[-/-] embryos at the 10-somite stage. Arrowheads
indicate the presumptive positions of pectoral fin buds. All images are captured from
the dorsal side. Scale bars represent 100 μm.

Figure 6

Figure 6. Screening for *hox* genes responsible for the specification of zebrafish pectoral fins.

(A) Schematic representation of the genetic screening for *hox* genes involved in the specification of pectoral fins in zebrafish. (B-G) Dorsal views of live mutant larvae at 3 dpf, obtained during the screening. After capturing images, genotyping and examination of mutations in *hoxb* genes were conducted. For the mutants illustrated in (C-G), *hoxb* genes are shown with frameshift mutations introduced. Detailed information is provided in Supplementary Table S1. (H-J) Dorsal views of l*hoxba*^{-/-};*hoxb5b*^{-/-} and *hoxba*^{-/-};*hoxb5b*^{-/-};*hoxb6b*^{-/-} larvae at 3 dpf. (K-M) Expression patterns of *tbx5a* in *hoxba*^{-/-};*hoxb5b*^{-/-} and *hoxba*^{-/-};*hoxb5b*^{-/-};*hoxb6b*^{-/-} mutants at 30 hpf. Arrowheads indicate the presumptive positions of pectoral fin buds. All images are captured from the dorsal side. Scale bars represent 100 μm.

Figure 7

Figure 7. *hoxb4a-b5a^del/del;hoxb5b^del/del* **partially recapituates the absence of the pectoral fins.**

(A-C) Dorsal views of zebrafish *hoxba^-/-;hoxb5b^del/del* larvae at 3 dpf. (D, E) Expression patterns of *tbx5a* in sibling wild-type and *hoxba^-/-;hoxb5b^del/del* (*n*=4) at 30 hpf. Arrowheads indicate the presumptive positions of pectoral fin buds. (F, G) Dorsal view of frameshift-induced *hoxb5a^-/-;hoxb5b^-/-* and *hoxb5a ^del/del;hoxb5b^del/del* larvae. (H) Expression patterns of zebrafish *hoxb4a* at the 10-somite stage. (I-M) Dorsal view of frameshift-induced *hoxb4a^-/-;hoxb5a^-/-;hoxb5b^-/-*, *hoxb4a-b5a ^del/del;hoxb5b^del/del* , *hoxb4a-b5a ^del/del*, and *hoxb4a ^del/del;hoxb5b^del/del* larvae. All images are captured from the dorsal side. Arrowheads indicate the presumptive positions of pectoral fin buds. Scale bars represent 100 μm.

Genetic evidence: zebrafish *hoxba* and *hoxbb* clusters are essential for

the anterior-posterior positioning of pectoral fins

Morimichi Kikuchi[a,#], Renka Fujii [a,#], Daiki Kobayashi[a,#], Yuki Kawabe[a], Haruna

Kanno[a], Sohju Toyama[a], Farah Tawakkal[a], Kazuya Yamada[a], and Akinori

Kawamura[a,*]

[a] Division of Life Science, Graduate School of Science and Engineering, Saitama

University, Shimo-Okubo 255, Sakura-ku, Saitama 338-8570, Japan

[#]These authors equally contributed to this work.

[*]Author to whom correspondence should be addressed:

Akinori Kawamura, Ph.D.

Division of Life Science, Graduate School of Science and Engineering, Saitama

University, Saitama 338-8570, Japan

Phone: +81-48-858-3808, Fax: +81-48-858-3408

E-mail: akawamur@mail.saitama-u.ac.jp

Abstract

Vertebrate paired appendages, such as the pectoral fins in fish and the

forelimbs in tetrapods, emerge at specific regions along the anterior-posterior axis of the

body. *Hox* genes are considered prime candidates for determining the positioning of

these paired appendages during development. Nevertheless, despite extensive

phenotypic analyses of numerous single and compound *Hox* knockout mice, no genetic

studies have identified substantial defects in limb positioning, leaving questions

unresolved. In a previous study, we generated seven distinct *hox* cluster-deficient

mutants in zebrafish. Here, we provide genetic evidence that zebrafish *hoxba;hoxbb*

cluster mutants specifically exhibit a complete absence of pectoral fins, accompanied by

the absence of *tbx5a* expression in pectoral fin buds. In these mutants, *tbx5a* expression

in the pectoral fin field of the lateral plate mesoderm fails to be induced at an early stage,

suggesting a lack of pectoral fin precursor cells. Furthermore, the competence to

respond to retinoic acid is lost in *hoxba;hoxbb* cluster mutants, indicating that *tbx5a*

expression cannot be induced in the pectoral fin bud. We also identify *hoxb4a, hoxb5a,*

and hoxb5b as pivotal genes underlying this process. Although the frameshift mutations

in these *hox* genes do not recapitulate the absence of pectoral fins, we demonstrate that

deletion mutants at these genomic loci show the absence of pectoral fins, albeit with low

penetrance. Our results suggest that the positioning of zebrafish pectoral fins is

cooperatively determined by *hoxb4a, hoxb5a, and hoxb5b* within *hoxba* and *hoxbb*

clusters, which induce *tbx5a* expression in the restricted pectoral fin field. Our findings

also provide insights into the acquisition of paired appendages in vertebrates.

Introduction

In jawed vertebrates, paired appendages—such as the pectoral and pelvic fins

in fish, and their homologous forelimbs and hindlimbs in tetrapods—develop at precise

locations along the anterior-posterior axis of each species. These paired appendages

arise from progenitor cells located in distinct regions of the lateral plate mesoderm

(Murata et al., 2010; Nishimoto and Logan, 2016; Shimada et al., 2013). The

positioning of these paired appendages has long fascinated researchers, yet our

understanding of the molecular mechanisms underlying this process remains limited.

In bilaterian animals, *Hox* genes—encoding evolutionarily conserved

homeodomain-containing transcription factors—provide positional information and

developmental timing along the anterior-posterior axis (Iimura and Pourquie, 2007;

Izpisua-Belmonte and Duboule, 1992; Krumlauf, 1994). A defining feature of *Hox*

genes is their structural organization into *Hox* clusters, where multiple *Hox* genes are

arranged in a precise order. Additionally, *Hox* clusters exhibit a distinctive phenomenon

known as *Hox* collinearity, where the genomic arrangement of *Hox* genes correlates

with specific developmental regions along the body axes (Dolle et al., 1989; Duboule

and Dolle, 1989; Graham et al., 1989). In vertebrates, *Hox* clusters underwent

divergence due to two rounds of whole-genome duplication early in vertebrate evolution

(Dehal and Boore, 2005; Ohno, 1970), leading to the establishment of four distinct *Hox*

clusters (*HoxA*, *HoxB*, *HoxC*, and *HoxD*), each consisting essentially of 1-13 paralogous

groups in tetrapods. In contrast, teleost fishes experienced an additional teleost-specific

whole-genome duplication, followed by the loss of a partial cluster, resulting in seven

hox clusters in zebrafish (Amores et al., 1998; Woltering and Durston, 2006).

Previous studies in mice and chickens suggest that the initial position of limbs

is likely regulated by *Hox* genes (Tanaka, 2013; Tickle, 2015). Supporting this

hypothesis, earlier research has shown that the anterior boundaries of specific *Hox* gene

expression domains align with the future positions of limbs (Burke et al., 1995; Cohn et

al., 1997). Moreover, experiments with avian embryos, in which several *Hox* genes

were overexpressed or functionally interfered with, demonstrated altered positions of

forelimb buds (Moreau et al., 2019). Additionally, it has been established that Hox

proteins bind to the enhancer of *Tbx5* gene, which is crucial for the initial formation of

forelimbs, thereby inducing *Tbx5* expression in the limb buds (Minguillon et al., 2012;

Nishimoto et al., 2014). However, despite the generation of numerous knockout mice

for single and multiple *Hox* genes, *Hoxb5* knockout mice exhibit only a rostral shift in

the position of forelimb buds, with incomplete penetrance (Rancourt et al., 1995), and

no significant abnormalities have been observed in the early stages of limb formation.

This lack of genetic evidence for limb positioning by *Hox* genes sharply contrasts with

results from genetic analyses of various knockout mice, which have shown that the

paralogous 9-13 *Hox* genes in *HoxA* and *HoxD* clusters are cooperatively essential for

proximal-distal patterning of limbs after limb bud formation (Boulet and Capecchi,

2004; Davis et al., 1995; Fromental-Ramain et al., 1996a; Fromental-Ramain et al.,

1996b; Kmita et al., 2005). Consequently, the precise role of *Hox* genes in defining the

initial position of limb formation along the anterior-posterior axis remains unclear.

Using zebrafish, which belong to the Actinopterygii class, distinct from the

tetrapods of the Sarcopterygii class, we previously generated mutants lacking each of

the seven zebrafish *hox* clusters using the CRISPR-Cas9 method (Yamada et al., 2021).

In our recent genetic analyses of *hoxaa*, *hoxab*, and *hoxda* clusters—homologous to the

mouse *HoxA* and *HoxD* clusters—we demonstrated that, similar to mice, these zebrafish

hox clusters cooperatively play an essential role in the formation of the pectoral fins

(Ishizaka et al., 2024), which are homologous to the forelimbs. In this study, we provide

the first genetic evidence, to our knowledge, that *Hox* genes specify the positions of

paired appendages in the vertebrates. The double-deletion mutants of *hoxba* and *hoxbb*

clusters, derived from the ancient *HoxB* cluster, exhibit a complete absence of pectoral

fins due to the failure to express *tbx5a*. Despite incomplete penetrance, we propose a

model in which *hoxb4a*, *hoxb5a*, and *hoxb5b* cooperatively provide positional cues

along the anterior-posterior axis within the lateral plate mesoderm, thereby specifying

the initial positions for fin bud formation through the induction of *tbx5a* in pectoral fin

field.

Results

The absence of pectoral fins in zebrafish *hoxba;hoxbb* cluster-deleted embryos

In a previous study, we created seven individual *hox* cluster-deficient mutants

in zebrafish using the CRISPR-Cas9 system (Yamada et al., 2021). Among these

mutants, *hoxba* cluster-deleted embryos exhibited morphological abnormalities in their

pectoral fins at 3 dpf (Figure 1A-C). To further investigate the pectoral fin phenotype in

hoxba cluster mutants, we first analyzed the expression patterns of *tbx5a*, which plays a

key role in the initial induction of pectoral fin buds in zebrafish (Ahn et al., 2002;

Garrity et al., 2002; Ng et al., 2002). When comparing with *tbx5a* expression in

wild-type embryos, we found that the *tbx5a* signal was reduced in the pectoral fin buds

of *hoxba* cluster mutants (Figure 1H, I). Given that *hoxba* and *hoxbb* clusters originated

from the ancestral *HoxB* cluster through teleost-specific whole-genome duplication

(Amores et al., 1998), there may be functional redundancy between them. Surprisingly,

we discovered that the simultaneous deletion of both *hoxba* and *hoxbb* clusters resulted

in the complete absence of pectoral fins (Figure 1A, G). In contrast, pectoral fins were

present in *hoxba*$^{-/-}$*;hoxbb*$^{+/-}$ and *hoxba*$^{+/-}$*;hoxbb*$^{-/-}$ mutants (Figure 1E, F), indicating that

an allele from either *hoxba* or *hoxbb* cluster is sufficient for the pectoral fin formation.

Moreover, all embryos lacking pectoral fins were identified as *hoxba;hoxbb* double

homozygous mutants, with the expected penetrance (n=15/252; 5.9%) being consistent

with predictions based on Mendelian genetics (1/16=6.3 %). Furthermore, while

hoxba;hoxbb cluster double homozygous mutants are embryonic lethal around 5 dpf, we

could not detect any trace of pectoral fin development. Alongside this phenotype, the

expression of *tbx5a* was significantly reduced to nearly undetectable levels in

hoxba;hoxbb cluster mutants at 30 hpf (Figure 1H-M). The absence of pectoral fins in

zebrafish *hoxba;hoxbb* cluster mutants sharply contrasts with the results of a previous

study, which showed that mice lacking all *Hoxb* genes except for *Hoxb13* of the *HoxB*

cluster did not exhibit apparent abnormalities in their forelimbs (Medina-Martinez et al.,

2000). Our genetic results suggest that the zebrafish *hoxba* and *hoxbb* clusters

cooperatively play a crucial role in the pectoral fin formation.

Lack of pectoral fins is specific to zebrafish *hoxba;hoxbb* cluster-deleted embryos

Multiple genetic studies in mice have demonstrated the functional redundancy

of *Hox* genes across the four *Hox* clusters (Horan et al., 1995; McIntyre et al., 2007; van

den Akker et al., 2001; Wellik and Capecchi, 2003; Wellik et al., 2002). To investigate

whether the absence of pectoral fins is specific to *hoxba;hoxbb* cluster mutants, we

created double-*hox* cluster mutants by crossing *hoxba* cluster mutants with six other *hox*

cluster mutants in zebrafish. The combinatorial mutations of *hoxba* cluster with five

other *hox* clusters, excluding *hoxbb* cluster, did not enhance the reduced expression of

tbx5a (Figure 2A-H). Furthermore, we examined combined mutants of *hox* clusters,

excluding *hoxba* and *hoxbb*. In *hoxca$^{-/-}$;hoxcb$^{-/-}$* cluster mutants, where the functions of

HoxC genes are completely absent, the expression patterns of *tbx5a* in fin buds were

indistinguishable from those of wild-type embryos (Figure 2I). Additionally, our

previous study showed that *tbx5a* expression is unaffected in *hoxaa$^{-/-}$;hoxab$^{-/-}$;hoxda$^{-/-}$*

cluster mutants (Ishizaka et al., 2024), which lack all *HoxA* and *HoxD*-related genes.

The results of our deletion mutations of zebrafish *hox* clusters in various combinations

emphasize that the diminished expression of *tbx5a* in the pectoral fin buds is evident

only in *hoxba;hoxbb* cluster mutants. Taken together, in contrast to the functional

redundancy of the four *Hox* clusters in mice, our genetic analysis indicates that the *Hox*

clusters responsible for the specification of pectoral fin buds are restricted to the

HoxB-derived *hoxba* and *hoxbb* clusters among the seven *hox* clusters in zebrafish.

Transformation of pectoral fin progenitor cells into cardiac cells in *hoxba;hoxbb*

cluster-deficient embryos

To understand the absence of pectoral fin formation in *hoxba;hoxbb* cluster

mutants, we examined the expression patterns of *tbx5a* during embryogenesis (Figure

3A-F). In wild-type embryos at the 10-somite stage, *tbx5a* expression appeared as

bilateral symmetric stripes in the anterior lateral mesoderm, extending from the

posterior midbrain to the second somite (Figure 3A) (Ahn et al., 2002). These

tbx5a-positive cells subsequently divided into two groups: the anteriorly migrating cells

form the heart primordia, while the posteriorly migrating cells give rise to the future

pectoral fin buds (Figure 3C, E). In contrast, we observed that abnormal expression

patterns of *tbx5a* in the lateral mesoderm were evident at the early stages in

hoxba;hoxbb cluster mutants. Compared to wild-type embryos, the bilateral stripes of

tbx5a expression were shorter along the anterior-posterior axis, and the *tbx5a*-positive

signal was absent from the posterior region where pectoral fin progenitor cells typically

emerge (Figure 3B). Although anterior migration toward the presumptive heart occurred,

posterior migration toward the fin buds was undetectable due to the lack of pectoral fin

progenitor cells in *hoxba;hoxbb* cluster mutants (Figure 3D, F). These results

underscore the early role of the *hoxba* and *hoxbb* clusters in providing positional

information for pectoral fin buds through the induction of *tbx5a* expression. In both

mice and zebrafish, there is a converse relationship between forelimb/pectoral fin and

heart formation in the anterior lateral mesoderm (Keegan et al., 2005; Niederreither et

al., 2001; Niederreither et al., 2002; Waxman et al., 2008). Therefore, we also analyzed

cardiac development (Figure 3G-J). By examining the expression patterns of *nkx2.5*, an

early cardiac progenitor marker (Chen and Fishman, 1996; Serbedzija et al., 1998), we

found that the expression domain of *nkx2.5* in *hoxba;hoxbb* cluster mutants extended

more posteriorly compared to wild-type embryos (Figure 3G, H). Furthermore, the

region positive for the differentiated cardiac cell-specific myosin light chain (*cmlc2*)

was enlarged in *hoxba;hoxbb* cluster mutants (Figure 3I, J) (Yelon et al., 1999). These

results indicate that *hoxba* and *hoxbb* clusters play an essential role in the initial

establishment of the pectoral fin field in zebrafish. In their absence, cells that should

form the pectoral fins instead differentiate into cardiac cells. The transformation of

pectoral fin progenitor cells into cardiac cells observed in zebrafish *hoxba;hoxbb* cluster

mutants is reminiscent of the anterior transformation typical of the loss-of-function

phenotypes of *Hox* genes in *Drosophila* and mice (Le Mouellic et al., 1992; Lewis,

1978).

***hoxba* and *hoxbb* clusters are indispensable for the retinoic acid-mediated**

induction of *tbx5a* expression in the fin buds

Retinoic acid (RA) is a well-known upstream regulator of *Hox* genes in

vertebrates (Boncinelli et al., 1991; Langston and Gudas, 1994; Marshall et al., 1996).

Loss of function of *retinaldehyde dehydrogenase 2* (*raldh2*), an essential regulator in

RA synthesis, leads to the absence of pectoral fins in zebrafish (Begemann et al., 2001;

Grandel et al., 2002), a phenotype similar to that observed in *hoxba;hoxbb* cluster

mutants. Previous studies have shown that exogenous RA treatments can rescue pectoral

fin formation in *raldh2*[-/-] embryos (Gibert et al., 2006; Grandel et al., 2002). It has been

suggested that RA directly or indirectly induces *tbx5a* expression in zebrafish fin buds

(Gibert et al., 2006; Grandel and Brand, 2011; Neto et al., 2012). The phenotypic

similarities between *hoxba;hoxbb* cluster mutants and *raldh2* mutants prompted us to

investigate whether RA exposure could also rescue the absence of pectoral fins in

hoxba;hoxbb cluster mutants. To explore this, we introduced a frameshift mutation in

raldh2 using CRISPR-Cas9, confirming the phenotypic recapitulation of previously

described *raldh2*[-/-] mutants (figure-supplement 1). Consistent with prior results (Gibert

et al., 2006; Grandel et al., 2002), RA treatment in our *raldh2* mutants rescued the

expression of *tbx5a* in pectoral fins (Figure 4A-D). In contrast, RA did not induce *tbx5a*

expression in the fin buds of *hoxba;hoxbb* cluster mutants (Figure 4E-H), and no trace

of pectoral fin formation was observed in the treated embryos even at 5 dpf.

Furthermore, the endogenous expression patterns of *raldh2* in *hoxba;hoxbb* cluster

mutants seem indistinguishable from those in wild-type embryos (Figure 4I, J),

suggesting that the response to RA is absent in the fin buds of *hoxba;hoxbb* cluster

mutants, while RA synthesis appears intact. Although a model proposing that RA

directly induces *tbx5a* expression has been suggested (Gibert et al., 2006; Grandel and

Brand, 2011; Neto et al., 2012), our results favor the model that RA-mediated induction

of *tbx5a* in the fin buds occurs through *hoxba* and *hoxbb* clusters. Given that

hoxba;hoxbb cluster mutants retain the other five intact *hox* clusters, our findings

further support the notion that the *hoxba* and *hoxbb* clusters are specifically responsible

for the RA-mediated specification of pectoral fins.

Expression patterns of *hoxb5a* and *hoxb5b* in pectoral fin buds are dependent on

RA

225   The absence of pectoral fins, commonly observed in *hoxba;hoxbb* cluster

mutants and *raldh2* mutants, suggests that candidate *hox* genes may be downregulated

in *raldh2* mutants. To investigate this, we performed RNA-seq analysis to compare the

expression profiles between sibling and *raldh2* homozygous embryos. As expected, we

found that most of the 3'-located *hox* genes are downregulated in *raldh2*$^{-/-}$ embryos

(Figure 5A, B, figure supplement 2). Notably, the levels of *hoxb5b* transcripts were

significantly reduced, along with those of *hoxb5a* among *hoxba* genes, in *raldh2*

mutants. Previous studies have shown that zebrafish *hoxb5a* and *hoxb5b* are expressed

in the lateral plate mesoderm corresponding to presumptive pectoral fin buds (Waxman

et al., 2008). Additionally, knockout mice for *Hoxb5*, which is homologous to zebrafish

hoxb5a and *hoxb5b*, only exhibit altered positioning of forelimbs (Rancourt et al., 1995),

among various other *Hox* knockouts that have been generated. Therefore, we examined

whether RA regulates the expression of *hoxb5a* and *hoxb5b* in fin buds. In wild-type

embryos, *hoxb5a* expression was observed in the neural tube, somites, and lateral plate

mesoderm (Figure 5C). However, in *raldh2* mutants, *hoxb5a* expression was

significantly reduced, notably in the lateral plate mesoderm, which differentiates into

the fin bud, where it was reduced to the point of being undetectable (Figure 5D). Since

RA treatments in *raldh2* mutants can rescue pectoral fin development, we found that

hoxb5a expression in *raldh2* mutants is also rescued by RA exposure (Figure 5C-E).

Regarding the expression of *hoxb5b*, which diverged from the same ancestral gene as

hoxb5a, RA-dependent expression was also confirmed, mirroring the observations made

with *hoxb5a* (Figure 5F-H). These results suggest that both *hoxb5a* and *hoxb5b* are

involved in the formation of the pectoral fin in zebrafish.

Genetic screening for *hox* genes responsible for the specification of pectoral fin

buds in zebrafish

To understand the molecular mechanisms underlying pectoral fin specification,

we aimed to identify the *hox* genes essential for pectoral fin formation through a genetic

approach. By introducing frameshift mutations into the relevant *hox* genes, we

anticipated that the resulting mutants would replicate the absence of pectoral fins

observed in *hoxba;hoxbb* mutants. Since *hoxbb* cluster contains fewer *hox* genes than

hoxba cluster (Figure 5A), we focused our investigation on *hoxbb* cluster. Although

hoxb5b is expressed in the presumptive fin buds (Figure 5F), other *hoxbb* genes may

also be involved. Therefore, we simultaneously injected three gRNAs targeting *hoxb5b*,

hoxb6b, and *hoxb8b* into *hoxba* hemizygous mutants, expecting various germline

mutations in these *hoxbb* genes in the injected fish (Figure 6A). Subsequently, F1 fish

with a hemizygous deletion of *hoxba* cluster and mosaic mutations were crossed with

hoxba;hoxbb hemizygous mutants. Among the progeny, several embryos lacking

pectoral fins were identified. Following genotyping and DNA sequencing of the target

hoxbb genes, we found that all embryos without pectoral fins possessed frameshift

mutations in *hoxb5b*, with $hoxba^{-/-};hoxbb^{+/-}$ being common among them (Figure 6B-E,

table-supplement 1, figure-supplement 3). Similarly, we injected each crRNA for

hoxb6b and *hoxb8b* into $hoxba^{+/-}$ mutants and confirmed that neither frameshift

mutation in *hoxb6b* nor *hoxb8b* in $hoxba^{-/-};hoxbb^{+/-}$ led to the loss of pectoral fins

(Figure 6F, G, figure-supplement 3). These results suggest that the frameshift mutation

of *hoxb5b* alone can compensate for the deletion of *hoxbb* cluster, recapitulating the

absence of pectoral fins in *hoxba;hoxbb* cluster mutants.

However, we encountered a problem: $hoxba^{-/-};hoxb5b^{-/-}$ embryos, which have a

homozygous frameshift mutation in *hoxb5b* and a homozygous deletion of *hoxba* cluster,

exhibited severely truncated but visibly present pectoral fins (Figure 6H, I). After

analyzing hundreds of embryos from several crossings, we did not find any

$hoxba^{-/-};hoxb5b^{-/-}$ embryos without pectoral fins. We also confirmed that the expression

of *tbx5a* was detectable in the fin buds of $hoxba^{-/-};hoxb5b^{-/-}$ embryos (Figure 6K, L), in

contrast to its absence in *hoxba;hoxbb* cluster mutants (Figure 1M). To examine the

potential contributions of other *hoxbb* genes, we observed the phenotypes of

$hoxba^{-/-};hoxb5b^{-/-}; hoxb6b^{-/-}$ embryos; however, these mutants did not exacerbate the

abnormalities and did not also exhibit the absence of pectoral fins (Figure 6J, M).

The phenotype of $hoxba^{-/-};hoxb5b^{-/-}$ embryos indicates that the frameshift

mutation in $hoxb5b$ cannot fully compensate for the complete deletion of $hoxbb$ cluster,

suggesting that other mechanisms may be involved. A novel genetic compensation

mechanism known as transcriptional adaptation has been identified: mRNA containing a

premature termination codon (PTC) can induce increased expression of structurally

related genes, thereby compensating for the function of mutated genes (El-Brolosy et al.,

2019; Rossi et al., 2015). To avoid producing transcripts with PTC (Sztal and Stainier,

2020), we generated a full locus-deleted allele of $hoxb5b$ using CRISPR-Cas9 with two

crRNAs targeting both ends of the target locus (figure-supplement 4). In contrast to the

phenotype of $hoxba^{-/-};hoxb5b^{-/-}$, we found that $hoxba^{-/-};hoxb5b^{de/dell}$ mutants lack

pectoral fins and exhibit a significant reduction in $tbx5a$ expression when intercrossing

$hoxba^{+/-};hoxb5b^{+/del}$ fish (Figure 7A, B, D, E). However, the occurrence rate of embryos

lacking pectoral fins (n=3/120; 2.5 %) was lower than predicted based on Mendelian

genetics (1/16; 6.3 %). Other $hoxba^{-/-};hoxb5b^{del/del}$ embryos did exhibit shortened

pectoral fins (Figure 7C). Additionally, concerning $hoxba$ cluster, we also generated

frameshift-induced $hoxb5a^{-/-};hoxb5b^{-/-}$ and a full locus-deleted $hoxb5a^{del/del};hoxb5b^{del/del}$

embryos (figure-supplement 4). $hoxb5a^{-/-};hoxb5b^{-/-}$ larvae consistently displayed normal

pectoral fins (Figure 7F). In contrast, $hoxb5a^{del/del};hoxb5b^{del/del}$ mutants exhibited a

slightly more pronounced phenotype of shortened pectoral fins, but no instances of

missing pectoral fins were identified (Figure 7G). Furthermore, due to the functional

redundancy observed between $hoxb4a$ and $hoxb5a$ observed in zebrafish vertebral

patterning and the RA-dependent expression of $hoxb4a$ (Grandel et al., 2002; Maeno et

al., 2024), as well as its expression in fin buds (Figure 7H), we generated

frameshift-induced $hoxb4a^{-/-};hoxb5a^{-/-};hoxb5b^{-/-}$ embryos and an allele lacking the entire

genomic region encompassing *hoxb4a* and *hoxb5a* (figure-supplement 4). Although

hoxb4a$^{-/-}$;hoxb5a$^{-/-}$;hoxb5b$^{-/-}$ embryos did not lack pectoral fins (Figure 7I), some

hoxb4a-b5a$^{del/del}$;hoxb5b$^{del/del}$ embryos did lack pectoral fins, although at a low

penetrance (Figure 7J, *n*=3/397; 0.7%). The majority of *hoxb4a-b5a$^{del/del}$;hoxb5b$^{del/del}$*

embryos displayed truncated pectoral fins (Figure 7K). The lack of pectoral fins was not

detected in *hoxb4a-b5a$^{del/del}$* and *hoxb4a$^{del/del}$;hoxb5b$^{del/del}$* mutants (Figure 7L, M,

(figure-supplement 4). These results suggest that *hoxb4a*, *hoxb5a*, and *hoxb5b* may

cooperatively contribute to anterior-posterior positioning of pectoral fin buds in

zebrafish. Furthermore, our findings reveal that transcriptional adaptation may partially

play a role in genetic compensation for frameshift-induced *hox* mutations during

pectoral fin formation, while also suggesting the potential involvement of other

unknown mechanisms or additional genomic regions in the initial induction of zebrafish

pectoral fin buds.

Discussion

Since the establishment of gene targeting (Mansour et al., 1988), numerous

Hox knockout mice have been generated. However, neither single nor compound *Hox*

mutants, nor *Hox* overexpression approaches, have been reported to cause significant

defects in limb positioning (Jurberg et al., 2013; Moreau et al., 2019). Consequently, the

role of *Hox* genes in the positioning of vertebrate paired appendages remains unclear

due to lack of genetic evidence. In this study, we provide genetic evidence using

zebrafish, demonstrating that specific combinatorial deletions of *hoxba* and *hoxbb*

clusters result in the absence of pectoral fins and a lack of induction of *tbx5a* expression

in the pectoral fin field. Based on our results, we propose a model in which *Hox*

expressions in the lateral plate mesoderm, regulated by RA synthesized in the paraxial

mesoderm, provide positional information and induce *tbx5a* expression in the

presumptive fin buds. Our genetic results shed light on longstanding fundamental

questions regarding the molecular mechanisms by which the positioning of paired

appendages is established in vertebrates.

Zebrafish *hoxba* and *hoxbb* clusters, possibly *hoxb4a*, *hoxb5a*, and *hoxb5b*,

cooperatively define the pectoral fin field in the lateral plate mesoderm.

We showed that *hoxba;hoxba* mutants exhibit a lack of pectoral fins. The

abnormal expression of *tbx5a* in the lateral plate mesoderm, where pectoral fin

precursor cells arise, was evident from its initial expression in the mutants. We presume

that *hox* genes in *hoxba* and *hoxbb* clusters, induced by RA, define the region along the

anterior-posterior axis where pectoral fin formation can occur and promote the

expression of *tbx5a* in the lateral plate mesoderm. Importantly, *tbx5a* expression in the

fin bud was not induced even by RA exposure in *hoxba;hoxbb* mutants (Figure 4E-H).

In zebrafish, it has been suggested that RA synthesized in the paraxial mesoderm acts

directly on the lateral plate mesoderm, or that RA induces *tbx5a* expression via *wnt2b* in

the intermediate mesoderm (Gibert et al., 2006; Grandel and Brand, 2011; Neto et al.,

2012). However, our results suggest that the main pathway integrates these signaling

cascades through *hoxba* and *hoxbb* clusters to induce *tbx5a* expression. Through genetic

studies, despite low penetrance, we showed that *hoxb4a*, *hoxb5a*, and *hoxb5b* play a

significant role in the induction of pectoral fin buds. The homologous Hoxb4 and

Hoxb5 proteins in mice have been shown to bind directly to the enhancer of *Tbx5*

(Minguillon et al., 2012). Furthermore, it has been demonstrated that the *Tbx5*

expression in the forelimb field is restricted by the posterior *Hox* genes, preventing

posterior expansion in mice and chickens (Moreau et al., 2019; Nishimoto et al., 2014).

However, our several combinatorial deletions of zebrafish *hox* clusters did not exhibit

any abnormalities in which the expression of *tbx5a* expanded to the posterior. In

zebrafish, we propose that *hoxb4a*, *hoxb5a*, and *hoxb5b*, which are regulated by RA in

the lateral plate mesoderm, simply induce the expression of *tbx5a*.

Partial recapitulation of genomic locus-deleted *hox* mutants suggests the other

unidentified mechanisms.

In this study, we observed that *hox* mutants with deletions of genomic loci

exhibited more pronounced pectoral fin phenotypes than frameshift-induced *hox*

mutants. The frameshift mutants used in this study are likely to be loss-of-function

alleles, as we noted several defects in vertebral patterning in these mutants (Maeno et al.,

2024). These results suggest that transcriptional adaptation may occur during the

formation of pectoral fins in frameshift-induced *hox* mutants, potentially compensating

for the loss of function. In our previous studies, which focused on dorsal and anal fin

patterning, vertebral patterning, and swim bladder formation, we demonstrated that the

phenotypes observed in zebrafish *hox* cluster-deleted mutants could essentially be

replicated by introducing frameshift mutations into *hox* genes within the cluster (Adachi

et al., 2024; Maeno et al., 2024; Satoh et al., 2024; Yamada et al., 2021). Therefore,

creating a mutant with a deleted locus that does not induce transcriptional adaptation

was unnecessary. One factor that may explain the differences in results between this

study and our prior findings is that even a small amount of *hox* gene products essential

for pectoral fin positioning may be sufficient to induce fin buds, as shown in Figure 1,

potentially leading to the manifestation of gene compensation through transcriptional

adaptation in frameshift-induced mutants. Alternatively, transcriptional adaptation may

function differently depending on various developmental processes or stages.

Interestingly, all the aforementioned phenotypes reproducible by frameshift mutations

were observed at later developmental stages than pectoral fin formation. Additionally,

in *hox* locus-deleted mutants, only some *hoxba$^{-/-}$;hoxb5b$^{de/dell}$* or

hoxb4a-b5a$^{del/del}$;hoxb5b$^{del/del}$ embryos were able to reproduce the loss of the pectoral

fin, suggesting the presence of other compensatory mechanisms. The organized

expression patterns of *Hox* genes are thought to be regulated by complex regulatory

controls, with their expression regulated according to the genomic order within the

cluster. In the absence of *Hox* function, neighboring intact *Hox* genes could compensate

for the loss of function. It is also possible that unidentified non-coding regions in *hoxba*

and *hoxbb* clusters are involved in pectoral fin positioning. These issues should be

clarified in future studies.

***Hox* genes may have contributed to the acquisition of pectoral fins during**

vertebrate evolution.

397   According to fossil records from the Cambrian period, the most primitive

vertebrates did not possess paired fins (Janvier, 1996; Shubin et al., 1997). It is believed

that, during subsequent evolution, vertebrates acquired primitive pectoral and pelvic

fins, which served as evolutionary precursors to tetrapod limbs, before diverging into

ray-finned and lobe-finned fishes. Our results align well with this evolutionary scenario.

Hox genes, which are conserved in bilaterian animals, are thought to have been present

in primitive vertebrates without paired fins. The early fossils of vertebrates with

pectoral fins were discovered in Ordovician and Silurian jawless fish (Janvier, 1996;

Shubin et al., 1997). Since it is estimated that two rounds of whole-genome duplications

occurred early in the Cambrian period (Dehal and Boore, 2005), the first vertebrate with

pectoral fins likely possessed four sets of *Hox* clusters, similar to those found in modern

tetrapods. *Hox* genes have long been fundamental tools for animal development, even

before vertebrates acquired paired fins. The acquisition of new functions in specific *Hox*

genes may have contributed to the innovation of novel morphological features. Our

findings that *hoxba;hoxbb* mutants completely lack pectoral fins suggest that molecular

evolution within the ancestral *HoxB* cluster may have contributed to the development of

pectoral fins in vertebrates. Although the specific molecular modifications remain

unclear, one possible model suggests that *HoxB* acquired a new capacity to induce the

expression of *Tbx5* in the lateral plate mesoderm. This may have played a role in

establishing the RA-*HoxB*-*Tbx5* gene cascade, which is essential for pectoral fin

formation. Notably, the ancestral *Tbx4/5* in ascidians, which do not have paired fins,

already possesses the ability to rescue the loss of forelimbs in mice (Minguillon et al.,

2009). Consistent with our idea, a model has been proposed in which the induction of

Tbx5 in the lateral plate mesoderm may have contributed to the acquisition of pectoral

fins.

Since *hoxba;hoxbb* mutants are embryonic lethal around 5 dpf, we cannot

observe the effects of these clusters on the pelvic fins, which develop around one month

after fertilization in zebrafish (Parichy et al., 2009). However, based on the composition

of zebrafish *hox* genes and their expression patterns in *hoxba* and *hoxbb* clusters, it is

suggested that these clusters have functions specific to the positioning of pectoral fins.

The presence of *hox* genes responsible for the position of pectoral fins implies that there

may be other *hox* genes that determine the position of pelvic fins and possibly

contribute to their acquisition. This remains a critical area for future research.

The positioning of pectoral fins is restricted to specific *hox* clusters in zebrafish, in

contrast to the functional overlap seen in mice.

433   Taking advantage of our availability of seven individual *hox* cluster mutants in

zebrafish, we created mutants with multiple cluster deletions. Through genetic analysis,

we demonstrated that *hox* genes responsible for positioning pectoral fin buds are

restricted to *hoxba* and *hoxbb* clusters. In contrast, analyses of knockout mice reveal

significant functional overlap among the four *Hox* clusters. Mice lacking all *HoxB* genes

except for *Hoxb13* have been generated, yet there are no reports of a phenotype

characterized by absent forelimbs (Medina-Martinez et al., 2000). In *Hoxb5* knockout

mice, the position of the forelimbs shifts rostrally by one vertebra, and forelimbs are not

absent in *Hoxa5;Hoxb5;Hoxc5* triple knockout mice, which lack all *Hox* genes of the

paralog 5 group (Xu et al., 2013). In comparison, *hoxba;hoxbb* mutants in zebrafish

exhibit a striking phenotype of absent pectoral fins. This phenomenon, where function

is confined to specific *hox* clusters, has also been observed in other developmental

systems of teleosts. In both zebrafish and medaka, multiple *hox* genes responsible for

positioning dorsal and anal fins are primarily located in *HoxC*-related clusters (Adachi

et al., 2024). As mentioned earlier, it is believed that paired fins in vertebrates emerged

after the quadruplication of the primitive *Hox* clusters. Therefore, *Hox* genes that

acquired new functions related to pectoral fin formation likely originated from specific

Hox clusters among the four, maintaining restricted functions that diverged after the

teleost-specific whole-genome duplication. The validity of this assumption requires

further verification through a clearer understanding of the overall functions of the 49

hox genes in zebrafish. However, the genetic results thus far suggest that, unlike in mice,

the functions of *hox* genes—especially those related to development uniquely acquired

in vertebrates—may be confined to specific *hox* clusters in zebrafish. While teleosts

have gained functional redundancy through subsequent teleost-specific whole-genome

duplication, it is postulated that, in mice, which have maintained four *Hox* clusters for

over half a billion years, functional overlap among *Hox* genes may ensure the

robustness of the developmental system.

Materials and methods

Zebrafish

Riken Wild-type (RW) zebrafish, obtained from the National BioResource

Project in Japan, were maintained at 27°C with a 14-hour light/10-hour dark cycle.

Embryos were collected from natural spawning, and the larvae were raised at 28.5°C.

Developmental stages of the embryos and larvae were determined based on hours

post-fertilization (hpf), days post-fertilization (dpf), and developmental stage-specific

features (Kimmel et al., 1995; Parichy et al., 2009). The alleles of *hox* cluster-deleted

mutants used in this study were previously generated using CRISPR-Cas9 (Yamada et

al., 2021). The following frameshift-induced *hox* mutants were created using

CRISPR-Cas9: *hoxb5a^{sud125}*, *hoxb5b^{sud136}*, *hoxb6b^{sud137}*, and *hoxb4a;hoxb5a^{sud144}*

(Maeno et al., 2024). All experiments involving live zebrafish were approved by the

Committee for Animal Care and Use of Saitama University.

Generation of mutants by the CRISPR-Cas9 system

All the mutants used in this study were generated using the Alt-R

CRISPR-Cas9 system (Integrated DNA Technologies). To create frameshift-induced

mutants, gene-specific crRNAs were incubated with common tracrRNA, followed by

Cas9 nuclease. For the generation of mutants that delete large genomic regions, two

crRNAs targeting both ends of the regions were incubated with common tracrRNA and

Cas9 nuclease. Approximately one nanoliter of the crRNA:tracrRNA-Cas9

RNA-protein complex was injected into fertilized embryos, which were subsequently

raised to juvenile fish. Candidate founder fish were selected using a heteroduplex

mobility shift assay for frameshift mutants (Ota et al., 2013) and by amplifying genomic

deletions with specific primers. After sexual maturation, candidate founder fish were

mated with wild-type fish to produce heterozygous F1 offspring. Among the F1

offspring, mutant fish carrying the same mutation were identified through PCR-based

genotyping followed by DNA sequencing. The target-specific sequences of the crRNAs

used in this study are listed (table-supplement 2). The frozen sperms from the mutants

used in this study have been deposited in the National BioResource Project Zebrafish in

Japan (https://shigen.nig.ac.jp/zebra/) and are available upon request.

Genotyping of mutants

For the phenotypic analysis of embryos and the maintenance of mutant fish

lines, PCR-based genotyping was performed. Briefly, genomic DNA was extracted from

the analyzed embryos or from the partially dissected caudal fins of anesthetized larvae

or fish using the NaOH method (Meeker et al., 2007), which served as the template for

PCR. The sequences of the primers used for genotyping frameshift-induced mutants are

listed in the supplementary Table 3. For genotyping deletion mutants of *hox* genes, PCR

was conducted using a combination of three primers, with their sequences also provided

(table-supplement 3). Genotyping of *hox* cluster-deleted mutants was performed using

PCR as previously described (Yamada et al., 2021). After the reactions, the PCR

products were separated by electrophoresis. Based on differences in the lengths of PCR

products derived from wild-type and mutated alleles, either a 2 % agarose gel in 0.5x

TBE buffer, a 15 % polyacrylamide gel in 0.5x TBE buffer, or direct DNA sequencing

was utilized to determine the genotype.

Whole-mount *in situ* hybridization

Whole-mount *in situ* hybridization was carried out as described (Thisse and

Thisse, 2014). After the staining, the embryos mounted in 70-80 % glycerol were

captured under the stereomicroscope (Leica M205 FA) with a digital camera (Leica

DFC350F). After taking images, PCR-based genotyping was carried out as described

above.

RNA-seq analysis

For RNA-seq analysis, embryos were obtained by the intercrosses between

raldh2 heterozygous fish. At the 18-20-somite stages, sibling and homozygous embryos

were separated based on the phenotype and 20 embryos per tube were collected. Then,

total RNA was extracted by ISOGENE (Nippon Gene) and followed by DNase I

treatment (Takara). After the quality check of the isolated RNA, RNA-seq analyses (*n*=2

for each) were carried out by Genewiz, Azenta Life Science.

Treatment of retinoic acid on the embryos

Treatment of retinoic acid on the embryos was performed as previously

described, with minor modifications (Gibert et al., 2006; Grandel et al., 2002). Briefly,

embryos were obtained from intercrosses between *raldh2* heterozygous mutants. At the

70 % epiboly stage, the embryos were soaked in a solution containing DMSO or 10^{-8}M

RA and incubated at 28.5°C. Embryos were fixed at 30 hpf for analysis of *tbx5a*

expression and at the 10-somite stage for examination of *hox* gene expression.

Acknowledgments

We thank the NBRP zebrafish for providing fish and preserving the mutant lines generated in this study. We also appreciate Drs. Koji Tamura and Gembu Abe for providing the zebrafish *tbx5a* plasmid. This work was supported by KAKENHI Grants-in-Aid for Scientific Research 18K06177, 23K05790 to AK.

References

References

Adachi, U., Koita, R., Seto, A., Maeno, A., Ishizu, A., Oikawa, S., Tani, T., Ishizaka, M.,
Yamada, K., Satoh, K., et al. (2024). Teleost Hox code defines regional identities
competent for the formation of dorsal and anal fins. *Proc Natl Acad Sci U S A* **121**,
e2403809121.

Ahn, D. G., Kourakis, M. J., Rohde, L. A., Silver, L. M. and Ho, R. K. (2002). T-box gene tbx5 is
essential for formation of the pectoral limb bud. *Nature* **417**, 754-758.

Amores, A., Force, A., Yan, Y. L., Joly, L., Amemiya, C., Fritz, A., Ho, R. K., Langeland, J.,
Prince, V., Wang, Y. L., et al. (1998). Zebrafish hox clusters and vertebrate genome
evolution. *Science* **282**, 1711-1714.

Begemann, G., Schilling, T. F., Rauch, G. J., Geisler, R. and Ingham, P. W. (2001). The zebrafish
neckless mutation reveals a requirement for raldh2 in mesodermal signals that pattern the
hindbrain. *Development* **128**, 3081-3094.

Boncinelli, E., Simeone, A., Acampora, D. and Mavilio, F. (1991). HOX gene activation by
retinoic acid. *Trends Genet* **7**, 329-334.

Boulet, A. M. and Capecchi, M. R. (2004). Multiple roles of Hoxa11 and Hoxd11 in the formation
of the mammalian forelimb zeugopod. *Development* **131**, 299-309.

Burke, A. C., Nelson, C. E., Morgan, B. A. and Tabin, C. (1995). Hox genes and the evolution of
vertebrate axial morphology. *Development* **121**, 333-346.

Chen, J. N. and Fishman, M. C. (1996). Zebrafish tinman homolog demarcates the heart field and
initiates myocardial differentiation. *Development* **122**, 3809-3816.

Cohn, M. J., Patel, K., Krumlauf, R., Wilkinson, D. G., Clarke, J. D. and Tickle, C. (1997).
Hox9 genes and vertebrate limb specification. *Nature* **387**, 97-101.

Davis, A. P., Witte, D. P., Hsieh-Li, H. M., Potter, S. S. and Capecchi, M. R. (1995). Absence of
radius and ulna in mice lacking hoxa-11 and hoxd-11. *Nature* **375**, 791-795.

Dehal, P. and Boore, J. L. (2005). Two rounds of whole genome duplication in the ancestral
vertebrate. *PLoS Biol* **3**, e314.

Dolle, P., Izpisua-Belmonte, J. C., Falkenstein, H., Renucci, A. and Duboule, D. (1989).
Coordinate expression of the murine Hox-5 complex homoeobox-containing genes during
limb pattern formation. *Nature* **342**, 767-772.

Duboule, D. and Dolle, P. (1989). The structural and functional organization of the murine HOX
gene family resembles that of Drosophila homeotic genes. *EMBO J* **8**, 1497-1505.

El-Brolosy, M. A., Kontarakis, Z., Rossi, A., Kuenne, C., Gunther, S., Fukuda, N., Kikhi, K.,
Boezio, G. L. M., Takacs, C. M., Lai, S. L., et al. (2019). Genetic compensation triggered
by mutant mRNA degradation. *Nature* **568**, 193-197.

Fromental-Ramain, C., Warot, X., Lakkaraju, S., Favier, B., Haack, H., Birling, C., Dierich, A.,
Doll e, P. and Chambon, P. (1996a). Specific and redundant functions of the paralogous
Hoxa-9 and Hoxd-9 genes in forelimb and axial skeleton patterning. *Development* **122**,
461-472.

Fromental-Ramain, C., Warot, X., Messadecq, N., LeMeur, M., Dolle, P. and Chambon, P.
(1996b). Hoxa-13 and Hoxd-13 play a crucial role in the patterning of the limb autopod.
Development **122**, 2997-3011.

Garrity, D. M., Childs, S. and Fishman, M. C. (2002). The heartstrings mutation in zebrafish
causes heart/fin Tbx5 deficiency syndrome. *Development* **129**, 4635-4645.

Gibert, Y., Gajewski, A., Meyer, A. and Begemann, G. (2006). Induction and prepatterning of the
zebrafish pectoral fin bud requires axial retinoic acid signaling. *Development* **133**,
2649-2659.

Graham, A., Papalopulu, N. and Krumlauf, R. (1989). The murine and Drosophila homeobox
gene complexes have common features of organization and expression. *Cell* **57**, 367-378.

Grandel, H. and Brand, M. (2011). Zebrafish limb development is triggered by a retinoic acid
signal during gastrulation. *Dev Dyn* **240**, 1116-1126.

Grandel, H., Lun, K., Rauch, G. J., Rhinn, M., Piotrowski, T., Houart, C., Sordino, P., Kuchler,
A. M., Schulte-Merker, S., Geisler, R., et al. (2002). Retinoic acid signalling in the
zebrafish embryo is necessary during pre-segmentation stages to pattern the
anterior-posterior axis of the CNS and to induce a pectoral fin bud. *Development* **129**,
2851-2865.

Horan, G. S., Ramirez-Solis, R., Featherstone, M. S., Wolgemuth, D. J., Bradley, A. and
Behringer, R. R. (1995). Compound mutants for the paralogous hoxa-4, hoxb-4, and
hoxd-4 genes show more complete homeotic transformations and a dose-dependent increase
in the number of vertebrae transformed. *Genes Dev* **9**, 1667-1677.

Iimura, T. and Pourquie, O. (2007). Hox genes in time and space during vertebrate body formation.
Dev Growth Differ **49**, 265-275.

Ishizaka, M., Maeno, A., Nakazawa, H., Fujii, R., Oikawa, S., Tani, T., Kanno, H., Koita, R.
and Kawamura, A. (2024). The functional roles of zebrafish HoxA- and HoxD-related
clusters in the pectoral fin development. *Sci Rep* **14**, 23602.

Izpisua-Belmonte, J. C. and Duboule, D. (1992). Homeobox genes and pattern formation in the
vertebrate limb. *Dev Biol* **152**, 26-36.

Janvier, P. (1996). Early Vertebrates.

Jurberg, A. D., Aires, R., Varela-Lasheras, I., Novoa, A. and Mallo, M. (2013). Switching axial
progenitors from producing trunk to tail tissues in vertebrate embryos. *Dev Cell* **25**,
451-462.

Keegan, B. R., Feldman, J. L., Begemann, G., Ingham, P. W. and Yelon, D. (2005). Retinoic acid
signaling restricts the cardiac progenitor pool. *Science* **307**, 247-249.

Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. and Schilling, T. F. (1995). Stages of
embryonic development of the zebrafish. *Dev Dyn* **203**, 253-310.

Kmita, M., Tarchini, B., Zakany, J., Logan, M., Tabin, C. J. and Duboule, D. (2005). Early
developmental arrest of mammalian limbs lacking HoxA/HoxD gene function. *Nature* **435**,
1113-1116.

Krumlauf, R. (1994). Hox genes in vertebrate development. *Cell* **78**, 191-201.

Langston, A. W. and Gudas, L. J. (1994). Retinoic acid and homeobox gene regulation. *Curr Opin*
Genet Dev **4**, 550-555.

Le Mouellic, H., Lallemand, Y. and Brulet, P. (1992). Homeosis in the mouse induced by a null
mutation in the Hox-3.1 gene. *Cell* **69**, 251-264.

Lewis, E. B. (1978). A gene complex controlling segmentation in Drosophila. *Nature* **276**, 565-570.

Maeno, A., Koita, R., Nakazawa, H., Fujii, R., Yamada, K., Oikawa, S., Tani, T., Ishizaka, M.,
Satoh, K., Ishizu, A., et al. (2024). The Hox code responsible for the patterning of the
anterior vertebrae in zebrafish. *Development* **151**.

Mansour, S. L., Thomas, K. R. and Capecchi, M. R. (1988). Disruption of the proto-oncogene
int-2 in mouse embryo-derived stem cells: a general strategy for targeting mutations to
non-selectable genes. *Nature* **336**, 348-352.

Marshall, H., Morrison, A., Studer, M., Popperl, H. and Krumlauf, R. (1996). Retinoids and
Hox genes. *FASEB J* **10**, 969-978.

McIntyre, D. C., Rakshit, S., Yallowitz, A. R., Loken, L., Jeannotte, L., Capecchi, M. R. and
Wellik, D. M. (2007). Hox patterning of the vertebrate rib cage. *Development* **134**,
2981-2989.

Medina-Martinez, O., Bradley, A. and Ramirez-Solis, R. (2000). A large targeted deletion of
Hoxb1-Hoxb9 produces a series of single-segment anterior homeotic transformations. *Dev*
Biol **222**, 71-83.

Meeker, N. D., Hutchinson, S. A., Ho, L. and Trede, N. S. (2007). Method for isolation of
PCR-ready genomic DNA from zebrafish tissues. *BioTechniques* **43**, 610, 612, 614.

Minguillon, C., Gibson-Brown, J. J. and Logan, M. P. (2009). Tbx4/5 gene duplication and the
origin of vertebrate paired appendages. *Proc Natl Acad Sci U S A* **106**, 21726-21730.

Minguillon, C., Nishimoto, S., Wood, S., Vendrell, E., Gibson-Brown, J. J. and Logan, M. P.
(2012). Hox genes regulate the onset of Tbx5 expression in the forelimb. *Development* **139**,
3180-3188.

Moreau, C., Caldarelli, P., Rocancourt, D., Roussel, J., Denans, N., Pourquie, O. and Gros, J.
(2019). Timed Collinear Activation of Hox Genes during Gastrulation Controls the Avian

647   Forelimb Position. *Curr Biol* **29**, 35-50 e34.

648   **Murata, Y., Tamura, M., Aita, Y., Fujimura, K., Murakami, Y., Okabe, M., Okada, N. and**

649   **Tanaka, M.** (2010). Allometric growth of the trunk leads to the rostral shift of the pelvic fin

650   in teleost fishes. *Dev Biol* **347**, 236-245.

651   **Neto, A., Mercader, N. and Gomez-Skarmeta, J. L.** (2012). The Osr1 and Osr2 genes act in the

652   pronephric anlage downstream of retinoic acid signaling and upstream of Wnt2b to maintain

653   pectoral fin development. *Development* **139**, 301-311.

654   **Ng, J. K., Kawakami, Y., Buscher, D., Raya, A., Itoh, T., Koth, C. M., Rodriguez Esteban, C.,**

655   **Rodriguez-Leon, J., Garrity, D. M., Fishman, M. C., et al.** (2002). The limb identity gene

656   Tbx5 promotes limb initiation by interacting with Wnt2b and Fgf10. *Development* **129**,

657   5161-5170.

658   **Niederreither, K., Vermot, J., Messaddeq, N., Schuhbaur, B., Chambon, P. and Dolle, P.** (2001).

659   Embryonic retinoic acid synthesis is essential for heart morphogenesis in the mouse.

660   *Development* **128**, 1019-1031.

661   **Niederreither, K., Vermot, J., Schuhbaur, B., Chambon, P. and Dolle, P.** (2002). Embryonic

662   retinoic acid synthesis is required for forelimb growth and anteroposterior patterning in the

663   mouse. *Development* **129**, 3563-3574.

664   **Nishimoto, S. and Logan, M. P.** (2016). Subdivision of the lateral plate mesoderm and specification

665   of the forelimb and hindlimb forming domains. *Semin Cell Dev Biol* **49**, 102-108.

666   **Nishimoto, S., Minguillon, C., Wood, S. and Logan, M. P.** (2014). A combination of activation and

667   repression by a colinear Hox code controls forelimb-restricted expression of Tbx5 and

668   reveals Hox protein specificity. *PLoS Genet* **10**, e1004245.

669   **Ohno, S.** (1970). Evolution by gene duplication.

670   **Ota, S., Hisano, Y., Muraki, M., Hoshijima, K., Dahlem, T. J., Grunwald, D. J., Okada, Y. and**

671   **Kawahara, A.** (2013). Efficient identification of TALEN-mediated genome modifications

672   using heteroduplex mobility assays. *Genes Cells* **18**, 450-458.

673   **Parichy, D. M., Elizondo, M. R., Mills, M. G., Gordon, T. N. and Engeszer, R. E.** (2009). Normal

674   table of postembryonic zebrafish development: staging by externally visible anatomy of the

675   living fish. *Dev Dyn* **238**, 2975-3015.

676   **Rancourt, D. E., Tsuzuki, T. and Capecchi, M. R.** (1995). Genetic interaction between hoxb-5 and

677   hoxb-6 is revealed by nonallelic noncomplementation. *Genes Dev* **9**, 108-122.

678   **Rossi, A., Kontarakis, Z., Gerri, C., Nolte, H., Holper, S., Kruger, M. and Stainier, D. Y.** (2015).

679   Genetic compensation induced by deleterious mutations but not gene knockdowns. *Nature*

680   **524**, 230-233.

681   **Satoh, K., Maeno, A., Adachi, U., Ishizaka, M., Yamada, K., Koita, R., Nakazawa, H., Oikawa,**

682   **S., Fujii, R., Furudate, H., et al.** (2024). Physical constraints on the positions and

683   dimensions of the zebrafish swim bladder by surrounding bones. *J Anat*.

684 **Serbedzija, G. N., Chen, J. N. and Fishman, M. C.** (1998). Regulation in the heart field of

685   zebrafish. *Development* **125**, 1095-1101.

686 **Shimada, A., Kawanishi, T., Kaneko, T., Yoshihara, H., Yano, T., Inohaya, K., Kinoshita, M.,**

687   **Kamei, Y., Tamura, K. and Takeda, H.** (2013). Trunk exoskeleton in teleosts is

688   mesodermal in origin. *Nat Commun* **4**, 1639.

689 **Shubin, N., Tabin, C. and Carroll, S.** (1997). Fossils, genes and the evolution of animal limbs.

690   *Nature* **388**, 639-648.

691 **Sztal, T. E. and Stainier, D. Y. R.** (2020). Transcriptional adaptation: a mechanism underlying

692   genetic robustness. *Development* **147**.

693 **Tanaka, M.** (2013). Molecular and evolutionary basis of limb field specification and limb initiation.

694   *Dev Growth Differ* **55**, 149-163.

695 **Thisse, B. and Thisse, C.** (2014). In situ hybridization on whole-mount zebrafish embryos and

696   young larvae. *Methods Mol Biol* **1211**, 53-67.

697 **Tickle, C.** (2015). How the embryo makes a limb: determination, polarity and identity. *J Anat* **227**,

698   418-430.

699 **van den Akker, E., Fromental-Ramain, C., de Graaff, W., Le Mouellic, H., Brulet, P., Chambon,**

700   **P. and Deschamps, J.** (2001). Axial skeletal patterning in mice lacking all paralogous group

701   8 Hox genes. *Development* **128**, 1911-1921.

702 **Waxman, J. S., Keegan, B. R., Roberts, R. W., Poss, K. D. and Yelon, D.** (2008). Hoxb5b acts

703   downstream of retinoic acid signaling in the forelimb field to restrict heart field potential in

704   zebrafish. *Dev Cell* **15**, 923-934.

705 **Wellik, D. M. and Capecchi, M. R.** (2003). Hox10 and Hox11 genes are required to globally

706   pattern the mammalian skeleton. *Science* **301**, 363-367.

707 **Wellik, D. M., Hawkes, P. J. and Capecchi, M. R.** (2002). Hox11 paralogous genes are essential

708   for metanephric kidney induction. *Genes Dev* **16**, 1423-1432.

709 **Woltering, J. M. and Durston, A. J.** (2006). The zebrafish hoxDb cluster has been reduced to a

710   single microRNA. *Nat Genet* **38**, 601-602.

711 **Xu, B., Hrycaj, S. M., McIntyre, D. C., Baker, N. C., Takeuchi, J. K., Jeannotte, L., Gaber, Z.**

712   **B., Novitch, B. G. and Wellik, D. M.** (2013). Hox5 interacts with Plzf to restrict Shh

713   expression in the developing forelimb. *Proc Natl Acad Sci U S A* **110**, 19438-19443.

714 **Yamada, K., Maeno, A., Araki, S., Kikuchi, M., Suzuki, M., Ishizaka, M., Satoh, K., Akama, K.,**

715   **Kawabe, Y., Suzuki, K., et al.** (2021). An atlas of seven zebrafish hox cluster mutants

716   provides insights into sub/neofunctionalization of vertebrate Hox clusters. *Development*

717   **148**.

718 **Yelon, D., Horne, S. A. and Stainier, D. Y.** (1999). Restricted expression of cardiac myosin genes

719 reveals regulated aspects of heart tube assembly in zebrafish. *Dev Biol* **214**, 23-37.

 Figure legends

Figure 1.   Lack of pectoral fins in *hoxba;hoxbb* cluster-deleted mutants. (A-G)

Dorsal views of live zebrafish larvae at 3 dpf, obtained from intercrosses between

hoxba;hoxbb cluster-deficient hemizygous fish. Arrowheads indicate the positions of the

pectoral fins. (H-M) Expression patterns of *tbx5a* in the pectoral fin bud (arrowhead) at

30 hpf. Dorsal views are shown, and the genotype of each specimen was determined.

For each genotype, reproducibility was confirmed with at least three different specimens.

Genotyping revealed that all the embryos lacking pectoral fins (*n*=15) were

hoxba;hoxbb double homozygotes. Scale bars are 100 μm.

Figure 2.   Significantly decreased expression of *tbx5a* in the pectoral fin buds is

specific to *hoxba;hoxbb* cluster-deleted mutants.

(A-I) Expression patterns of *tbx5a* in the pectoral fin buds of combinatorial deletion

mutants of zebrafish *hox* clusters. Dorsal views of embryos at 30 hpf are displayed.

After capturing images, the genotype of each specimen was determined. For each

genotype, reproducibility was confirmed with at least three different specimens. Scale

bars are 100 μm.

Figure 3.   Transformation of pectoral fin progenitor cells into cardiac cells in

***hoxba;hoxbb* homozygous embryos.** (A-F) Expression patterns of *tbx5a* were

compared between sibling wild-type and *hoxba;hoxbb* homozygous embryos during

embryogenesis. The range of *tbx5a* expression in the lateral mesoderm is indicated by a

bracket. The progenitor cells of pectoral fins are indicated by an arrowhead. (G, H)

Expression patterns of *nkx2.5* were compared between sibling and *hoxba;hoxbb* mutants.

Enlarged views are shown in (G', H'). The different regions of *nkx2.5* expression

between wild-type and mutants are indicated by brackets. (I, J) Expression patterns of

cmcl2 are shown. All images were captured from the dorsal side. For each stage,

reproducibility was confirmed with at least three different specimens. Scale bars

indicate 100 μm.

Figure 4.   Zebrafish *hoxba;hoxbb* mutants lack a response to RA in fin buds.

(A-D) Exogenous RA exposure in *raldh2* mutants can rescue *tbx5a* expression in fin

buds. Arrowheads indicate the positions of the pectoral fin buds. (E-H) RA treatments in

hoxba;hoxbb cluster-deleted mutants do not rescue *tbx5a* expression in fin buds. (I, J)

Expression patterns of *raldh2* were compared between sibling and *hoxba;hoxbb* mutants.

For each genotype, reproducibility was confirmed with at least three different specimens.

All images were captured from the dorsal side. Scale bars represent 100 μm.

Figure 5.   The expression patterns of *hox5a* and *hoxb5b* are regulated by RA.

(A) Schematic representation of the 49 *hox* genes organized into seven *hox* clusters in

zebrafish. *hox* genes in *hoxba* and *hoxbb* clusters are highlighted with orange. (B)

Expression profiles of the 49 *hox* genes in wild-type and *raldh2*$^{-/-}$ embryos at the

20-somite stage, analyzed by RNA-seq. The average FPKM of each *hox* gene was

compared between sibling and *raldh2* mutants. The absence of specific *hox* genes is

indicated by a slash. (C-H) Expression patterns of *hoxb5a* and *hoxb5b* in sibling

wild-type, *raldh2*$^{-/-}$, and RA-treated *raldh2*$^{-/-}$ embryos at the 10-somite stage.

Arrowheads indicate the presumptive positions of pectoral fin buds. For each genotype,

reproducibility was confirmed with at least three different specimens. All images are

captured from the dorsal side. Scale bars represent 100 μm.

Figure 6. Screening for *hox* genes responsible for the specification of zebrafish

pectoral fins.

(A) Schematic representation of the genetic screening for *hox* genes involved in the

specification of pectoral fins in zebrafish. (B-G) Dorsal views of live mutant larvae at 3

dpf, obtained during the screening. After capturing images, genotyping and DNA

sequencing in target *hoxb* genes were conducted. For the mutants illustrated in (C-G),

hoxb genes are shown with frameshift mutations introduced. Detailed information is

provided in Supplementary Table S1. (H-J) Dorsal views of *hoxba$^{-/-}$;hoxb5b$^{-/-}$* and

hoxba$^{-/-}$;hoxb5b$^{-/-}$;hoxb6b$^{-/-}$ larvae at 3 dpf. (K-M) Expression patterns of *tbx5a* in

hoxba$^{-/-}$;hoxb5b$^{-/-}$ and *hoxba$^{-/-}$;hoxb5b$^{-/-}$;hoxb6b$^{-/-}$* mutants at 30 hpf. Arrowheads

indicate the presumptive positions of pectoral fin buds. All images are captured from the

dorsal side. Scale bars represent 100 μm.

Figure 7.   *hoxb4a-b5a$^{del/del}$;hoxb5b$^{del/del}$* larvae partially recapitulate the absence of

the pectoral fins.

(A-C) Dorsal views of zebrafish *hoxba$^{-/-}$;hoxb5b$^{del/del}$* larvae at 3 dpf. (D, E) Expression

patterns of *tbx5a* in sibling wild-type and *hoxba$^{-/-}$;hoxb5b$^{del/del}$* (*n*=4) at 30 hpf.

Arrowheads indicate the presumptive positions of pectoral fin buds. (F, G) Dorsal view

of frameshift-induced *hoxb5a$^{-/-}$;hoxb5b$^{-/-}$* and *hoxb5a$^{del/del}$;hoxb5b$^{del/del}$* larvae. (H)

Expression patterns of zebrafish *hoxb4a* at the 10-somite stage. (I-K) Dorsal view of

frameshift-induced *hoxb4a$^{-/-}$;hoxb5a$^{-/-}$;hoxb5b$^{-/-}$* and *hoxb4a-b5a$^{del/del}$;hoxb5b$^{del/del}$*

larvae. All images are captured from the dorsal side. Arrowheads indicate the

presumptive positions of pectoral fin buds. Scale bars represent 100 μm.

Figure S1.   Generation of the zebrafish *raldh2* mutants by CRISPR-Cas9. (A) The

nucleotide sequence surrounding the mutations in *raldh2* is shown on the left. The target

sequence of the crRNA is emphasized with an underline, while the inserted nucleotide

sequence is indicated in orange. Below the DNA sequence, the predicted amino acid is

displayed, with red letters highlighting the abnormal amino acid resulting from the

frameshift mutation. The asterisk represents the termination codon. (B) A schematic

representation of the predicted protein structure is shown, with the red box indicating

the abnormal amino acid sequences resulting from the frameshift mutations. The total

number of predicted amino acids is displayed to the right of the schematic. (C) The

absence of pectoral fins (arrowhead) in *raldh2* homozygous mutants is observed

dorsally at 3 dpf. (D) The expression of *tbx5a* in the pectoral fin buds (arrowhead) is

significantly reduced in *raldh2* mutants. The phenotype of our *raldh2*^{sud118} mutants

closely resembles the phenotype observed in previously isolated *raldh2* mutants.

Figure S2.   Volcano plot of the transcriptome analysis between sibling and *raldh2*

mutants. The volcano plot illustrates both upregulated and downregulated differentially

expressed genes from the comparison between wild-type and *raldh2* mutant embryos.

Dot with more than a 2-fold increase in expression is indicated in orange, while those

with more than a 2-fold decrease are shown in light blue. Among them, dots

corresponding to *hox* genes are specifically highlighted.

Figure S3.   Generation of the frameshift-induced *hox* mutants using

CRISPR-Cas9. Schematic representations show zebrafish *hoxb8b, hoxb5b;hoxb6b,* and

hoxb5b;hoxb6b;hoxb8b mutants. The other frameshift-induced *hox* mutants were

previously described (Maeno et al., 2024). The nucleotide sequence around the

mutations is shown on the left. The target sequence of crRNA is emphasized with an

orange underline. The inserted nucleotide sequence is emphasized in red. Below the

DNA sequence, the predicted amino acid is shown. Red letters indicate the abnormal

amino acid caused by the frameshift mutation. On the right, a schema represents the

predicted protein structure. The blue box indicates the homeodomain, and red box

indicates the abnormal amino acid sequences from the frameshift mutations. The total

number of the predicted amino acids is shown on the right of the schema.

Figure S4. Generation of locus-deletion mutants using CRISPR-Cas9. (A, C, E, G).

Schematic representations show the genomic structure of zebrafish *hoxb4a, hox5a,* and

hoxb5b. Boxes represent the exons, with the blue boxes indicating the coding regions.

Black arrowheads indicate the target sites of crRNA, while red arrowheads represent the

primers used for genotyping. (B, D, F, H) Display genomic locus deletions of *hox4a,*

hox5a, hoxb5b, and *hoxb4a-b5a* deletion mutants. The flanking sequences of the 5'- and

3'- crRNA targets (underline) are shown in blue and green letters, respectively.