1

HoxB-derived hoxba and hoxbb clusters are essential for the

anterior-posterior positioning of zebrafish pectoral fins

Morimichi Kikuchi[a,#], Renka Fujii [a,#], Daiki Kobayashi[a,#], Yuki Kawabe[a], Haruna

Kanno[a], Sohju Toyama[a], Farah Tawakkal[a], Kazuya Yamada[a], and Akinori

Kawamura[a,*]

[a] Division of Life Science, Graduate School of Science and Engineering, Saitama

University, Shimo-Okubo 255, Sakura-ku, Saitama 338-8570, Japan

[#]These authors equally contributed to this work.

[*]Author to whom correspondence should be addressed:

Akinori Kawamura, Ph.D.

Division of Life Science, Graduate School of Science and Engineering, Saitama

University, Saitama 338-8570, Japan

Phone: +81-48-858-3808

E-mail: akawamur@mail.saitama-u.ac.jp

Abstract

Vertebrate paired appendages, such as the pectoral fins in fish and the forelimbs in tetrapods, arise at specific regions along the anterior-posterior axis of the body. Hox genes have long been considered prime candidates for determining the anteroposterior positioning of these paired appendages during development. Evidence from various model organisms, including mouse and chick, supports a role for Hox genes in limb positioning. However, despite extensive phenotypic analyses of numerous single and compound Hox knockout mice, clear genetic evidence for substantial defects in limb positioning has been limited, leaving questions unresolved. In a previous study, we generated seven distinct hox cluster-deficient mutants in zebrafish. Here, we provide genetic evidence that zebrafish *hoxba;hoxbb* cluster-deleted mutants specifically exhibit a complete lack of pectoral fins, accompanied by the absence of *tbx5a* expression in pectoral fin buds. In these mutants, *tbx5a* expression in the pectoral fin field of the lateral plate mesoderm fails to be induced at an early stage, suggesting a loss of pectoral fin precursor cells. Furthermore, the competence to respond to retinoic acid is lost in *hoxba;hoxbb* cluster mutants, indicating that *tbx5a* expression cannot be induced in the pectoral fin buds. We further identify *hoxb4a, hoxb5a, and hoxb5b* as pivotal genes underlying this process. Although the frameshift mutations in these hox genes do not recapitulate the absence of pectoral fins, we demonstrate that deletion mutants at these genomic loci show the absence of pectoral fins with low penetrance. Our results suggest that, by establishing the expression domains along the anteroposterior axis, *hoxb4a, hoxb5a,* and *hoxb5b* within hoxba and hoxbb clusters cooperatively determine the positioning of zebrafish pectoral fins through the induction of *tbx5a* expression in the

45   restricted pectoral fin field. Our findings also provide insights into the evolutionary

46   origin of paired appendages in vertebrates.

Introduction

In jawed vertebrates, paired appendages—such as the pectoral and pelvic fins in fish, and their homologous forelimbs and hindlimbs in tetrapods—develop at precise locations along the anterior-posterior axis of each species. These paired appendages arise from progenitor cells located in distinct regions of the lateral plate mesoderm (Murata et al., 2010; Nishimoto and Logan, 2016; Shimada et al., 2013). The anteroposterior positioning of these paired appendages has long fascinated researchers, yet our understanding of the molecular mechanisms underlying this process remains limited.

In bilaterian animals, Hox genes—encoding evolutionarily conserved homeodomain-containing transcription factors—provide positional information and developmental timing along the anterior-posterior axis (Iimura and Pourquie, 2007; Izpisua-Belmonte and Duboule, 1992; Krumlauf, 1994). A defining feature of Hox genes is their structural organization into Hox clusters, where multiple Hox genes are arranged in a precise order. Additionally, Hox clusters exhibit a distinctive phenomenon known as Hox collinearity, where the genomic arrangement of Hox genes correlates with specific developmental regions along the body axes (Dolle et al., 1989; Duboule and Dolle, 1989; Graham et al., 1989). In vertebrates, Hox clusters underwent divergence due to two rounds of whole-genome duplication early in vertebrate evolution (Dehal and Boore, 2005; Ohno, 1970), leading to the establishment of four distinct Hox clusters (HoxA, HoxB, HoxC, and HoxD), each consisting essentially of 1-13 paralogous groups in tetrapods. In contrast, teleost fishes experienced an additional teleost-specific whole-genome duplication, followed by the loss of a hox cluster,

resulting in seven hox clusters in zebrafish (Amores et al., 1998; Woltering and Durston,

2006).

Multiple studies in chick and mouse have suggested that the initial

anteroposterior position of limbs is regulated by Hox genes (Tanaka, 2013; Tickle,

2015). For example, the anterior boundaries of Hox gene expression domains were

shown to align with future limb positions in chick embryos (Burke et al., 1995; Cohn et

al., 1997). Experimental manipulations in avian embryos, including overexpression or

interference of specific Hox genes, led to altered positions of forelimb buds (Moreau et

al., 2019). In mouse, *Hoxb5* knockout mutants exhibit a rostral shift of forelimb buds

with incomplete penetrance (Rancourt et al., 1995), while other subtle shifts in limb

positioning have also been reported in Hox mutants (Royle et al., 2021). Furthermore,

loss of *Gdf11* in mice, which alters the expression of posterior Hox9–13 genes, causes

posterior displacement of hindlimb buds, whereas ectopic *Gdf11* induces anterior shifts

of hindlimb position (Matsubara et al., 2017). At the molecular level, Hox proteins

directly bind to the Tbx5 limb enhancer and regulate its expression, providing a

mechanistic link between Hox activity and forelimb initiation (Minguillon et al., 2012;

Nishimoto et al., 2014). Together, these studies provide strong evidence across different

model organisms that Hox genes contribute to the regulation of limb positioning.

Nevertheless, despite the generation of numerous single and compound Hox knockout

mice, no severe defects in the initial positioning of limb buds have been documented,

and the precise mechanisms by which Hox genes specify limb position remain

incompletely understood. This stands in contrast to the well-established role of Hox

genes in limb patterning after limb bud formation, where paralogous group 9–13 genes

in the HoxA and HoxD clusters cooperatively control the proximal–distal axis of

developing limbs (Boulet and Capecchi, 2004; Davis et al., 1995; Fromental-Ramain et

al., 1996a; Fromental-Ramain et al., 1996b; Kmita et al., 2005). Consequently, the exact

role of Hox genes in defining the initial position of limb formation along the

anterior-posterior axis remains unclear.

Using zebrafish, which belong to the vertebrate class Actinopterygii and are

distinct from the tetrapods of the class Sarcopterygii, we previously generated mutants

lacking each of the seven zebrafish hox clusters using the CRISPR-Cas9 method

(Yamada et al., 2021). In our recent genetic analyses of hoxaa, hoxab, and hoxda

clusters—orthologous to the mouse HoxA and HoxD clusters—we demonstrated that,

similar to mice, these zebrafish hox clusters cooperatively play an essential role in the

formation of the pectoral fins (Ishizaka et al., 2024), which are homologous to the

forelimbs. In this study, we provide the first genetic evidence, to our knowledge, that

Hox genes specify the positions of paired appendages in vertebrates. The

double-deletion mutants of hoxba and hoxbb clusters, derived from the ancient HoxB

cluster, exhibit a complete absence of pectoral fins due to the failure to express *tbx5a*.

Despite incomplete penetrance, we propose a model in which *hoxb4a*, *hoxb5a*, and

hoxb5b cooperatively provide positional cues along the anterior-posterior axis within

the lateral plate mesoderm, thereby specifying the initial positions for fin bud formation

through the induction of *tbx5a* in the pectoral fin field.

Results

Absence of pectoral fins in *hoxba;hoxbb* cluster-deleted zebrafish

In a previous study, we created seven individual hox cluster-deficient mutants in zebrafish using the CRISPR-Cas9 system (Yamada et al., 2021). Among these mutants, hoxba cluster-deleted embryos exhibited morphological abnormalities in their pectoral fins at 3 dpf (Figure 1A-C). To further investigate the pectoral fin phenotype in hoxba cluster mutants, we first analyzed the expression patterns of *Tbx5* orthologs. Zebrafish possess two paralogs. *tbx5a* and *tbx5b* (Boyle-Anderson et al., 2022). Among them, *tbx5a* plays a predominant role in the initial induction of pectoral fin buds in zebrafish (Ahn et al., 2002; Garrity et al., 2002; Ng et al., 2002). When comparing with *tbx5a* expression in wild-type embryos, we found that the *tbx5a* signal was reduced in the pectoral fin buds of hoxba cluster mutants (Figure 1G, H). Given that hoxba and hoxbb clusters originated from the ancestral HoxB cluster through teleost-specific whole-genome duplication (Amores et al., 1998), there may be functional redundancy between them. Surprisingly, we discovered that the simultaneous deletion of both hoxba and hoxbb clusters resulted in the complete absence of pectoral fins (Figure 1A, F). In contrast, pectoral fins were present in $hoxba^{-/-};hoxbb^{+/-}$ and $hoxba^{+/-};hoxbb^{-/-}$ mutants (Figure 1D, E), indicating that an allele from either hoxba or hoxbb cluster is sufficient for the pectoral fin formation. Moreover, all embryos lacking pectoral fins were identified as *hoxba;hoxbb* double homozygous mutants, with the expected penetrance (n=15/252; 5.9%) being consistent with predictions based on Mendelian genetics (1/16=6.3 %). Furthermore, while *hoxba;hoxbb* cluster double homozygous mutants are embryonic lethal around 5 dpf, we could not detect any trace of pectoral fin

development. Alongside this phenotype, the expression of *tbx5a* was significantly

reduced to nearly undetectable levels in *hoxba;hoxbb* cluster mutants at 30 hpf (Figure

1G-L). The absence of pectoral fins in zebrafish *hoxba;hoxbb* cluster mutants sharply

contrasts with the results of a previous study, which showed that mice lacking all HoxB

genes except for *Hoxb13* of the HoxB cluster did not exhibit apparent abnormalities in

their forelimbs (Medina-Martinez et al., 2000). Our genetic results suggest that the

zebrafish hoxba and hoxbb clusters cooperatively play a crucial role in pectoral fin

formation.

Pectoral fin loss specific to *hoxba;hoxbb* cluster-deleted zebrafish

150   Multiple genetic studies in mice have demonstrated the functional redundancy

of Hox genes across the four Hox clusters (Horan et al., 1995; McIntyre et al., 2007;

van den Akker et al., 2001; Wellik and Capecchi, 2003; Wellik et al., 2002). To

investigate whether the absence of pectoral fins is specific to *hoxba;hoxbb* cluster

mutants, we created double-hox cluster mutants by crossing hoxba cluster mutants with

six other hox cluster mutants in zebrafish. The combinatorial mutations of hoxba cluster

with five other hox clusters, excluding hoxbb cluster, did not enhance the reduced

expression of *tbx5a* (Figure 2A-H). Furthermore, we examined combined mutants of

hox clusters, excluding hoxba and hoxbb. In $hoxca^{-/-};hoxcb^{-/-}$ cluster mutants, where the

functions of HoxC genes are completely absent, the expression patterns of *tbx5a* in fin

buds were indistinguishable from those of wild-type embryos (Figure 2I). Additionally,

our previous study showed that *tbx5a* expression is unaffected in

$hoxaa^{-/-};hoxab^{-/-};hoxda^{-/-}$ cluster mutants (Ishizaka et al., 2024), which lack all HoxA

and HoxD-related genes. The results of our deletion mutations of zebrafish hox clusters

in various combinations emphasize that the diminished expression of *tbx5a* in the

pectoral fin buds is evident only in hoxba;hoxbb cluster mutants. Taken together, in

contrast to the functional redundancy of the four Hox clusters in mice, our genetic

analysis indicates that the Hox clusters responsible for the specification of pectoral fin

buds are restricted to the HoxB-derived hoxba and hoxbb clusters among the seven hox

clusters in zebrafish.

171 Loss of fin progenitors with posterior expansion of cardiac marker expression

To understand the absence of pectoral fin formation in *hoxba;hoxbb* cluster

mutants, we examined the expression patterns of *tbx5a* during embryogenesis (Figure

3A-F). In wild-type embryos at the 10-somite stage, *tbx5a* expression appeared as

bilateral symmetric stripes in the anterior lateral mesoderm, extending from the

posterior midbrain to the second somite (Figure 3A) (Ahn et al., 2002). These

tbx5a-positive cells subsequently divided into two groups: the anteriorly migrating cells

form the heart primordia, while the posteriorly migrating cells give rise to the future

pectoral fin buds (Figure 3C, E). In contrast, abnormal expression patterns of *tbx5a*

were evident at early stages in *hoxba;hoxbb* cluster mutants. Compared to wild-type

embryos, the bilateral stripes of *tbx5a* expression were shortened along the

anterior-posterior axis, and the *tbx5a*-positive signal was absent from the posterior

region where pectoral fin progenitor cells typically emerge (Figure 3B). Although

anterior migration toward the presumptive heart occurred, posterior migration toward

the fin buds was undetectable, consistent with the loss of pectoral fin progenitor cells in

hoxba;hoxbb cluster mutants (Figure 3D, F). These results underscore the early role of

the hoxba and hoxbb clusters in the formation of pectoral fin buds through the induction

of *tbx5a* expression. In zebrafish, studies have shown a reciprocal relationship between

pectoral fin and heart progenitor populations in the anterior lateral mesoderm, where

retinoic acid signaling promotes pectoral fin formation while restricting cardiac

progenitors (Waxman et al., 2008). Therefore, we also analyzed cardiac development

(Figure 3G-J). By examining the expression patterns of *nkx2.5*, an early cardiac

progenitor marker (Chen and Fishman, 1996; Serbedzija et al., 1998), we found that the

expression domain of *nkx2.5* in *hoxba;hoxbb* cluster mutants extended more posteriorly

compared to wild-type embryos (Figure 3G, H). Furthermore, the region positive for the

differentiated cardiac cell-specific myosin light chain (*cmlc2*) showed abnormal

morphology relative to wild-type embryos (Figure 3I, J) (Yelon et al., 1999). Taken

together, these results indicate that hoxba and hoxbb clusters are essential for

establishing the pectoral fin field in zebrafish. In their absence, pectoral fin progenitor

cells fail to emerge, accompanied by a posterior expansion of cardiac marker expression

domain. Our results highlight a shift in the balance between fin and cardiac progenitor

populations within the anterior lateral plate mesoderm.

hoxba/bb clusters required for RA-mediated *tbx5a* induction in the fin buds

Retinoic acid (RA) is a well-known upstream regulator of Hox genes in

vertebrates (Boncinelli et al., 1991; Langston and Gudas, 1994; Marshall et al., 1996).

Loss of function of *retinaldehyde dehydrogenase 2* (*raldh2*), an essential regulator in

RA synthesis, leads to the absence of pectoral fins in zebrafish (Begemann et al., 2001;

Grandel et al., 2002), a phenotype similar to that observed in *hoxba;hoxbb* cluster

mutants. Previous studies have shown that exogenous RA treatments can rescue pectoral

fin formation in *raldh2^-/-* embryos (Gibert et al., 2006; Grandel et al., 2002). It has been

suggested that RA directly or indirectly induces *tbx5a* expression in zebrafish fin buds

(Gibert et al., 2006; Grandel and Brand, 2011; Neto et al., 2012). The phenotypic

similarities between *hoxba;hoxbb* cluster mutants and *raldh2* mutants prompted us to

investigate whether RA exposure could also rescue the absence of pectoral fins in

hoxba;hoxbb cluster mutants. To explore this, we introduced a frameshift mutation in

raldh2 using CRISPR-Cas9, confirming the phenotypic recapitulation of previously

described *raldh2^-/-* mutants (figure-supplement 1). Consistent with prior results (Gibert

et al., 2006; Grandel et al., 2002), RA treatment in our *raldh2* mutants rescued the

expression of *tbx5a* in pectoral fins (Figure 4A-D). In contrast, RA did not induce *tbx5a*

expression in the fin buds of *hoxba;hoxbb* cluster mutants (Figure 4E-H), and no trace

of pectoral fin formation was observed in the treated embryos even at 5 dpf.

Furthermore, the endogenous expression patterns of *raldh2* in *hoxba;hoxbb* cluster

mutants seem indistinguishable from those in wild-type embryos (Figure 4I, J),

suggesting that the response to RA is absent in the fin buds of *hoxba;hoxbb* cluster

mutants, while RA synthesis appears intact. Although a model proposing that RA

directly induces *tbx5a* expression has been suggested (Gibert et al., 2006; Grandel and

Brand, 2011; Neto et al., 2012), our results indicate that RA-dependent induction of

tbx5a does not occur in the absence of hoxba and hoxbb clusters. Given that

hoxba;hoxbb cluster mutants retain the other five intact hox clusters, our findings

further support the notion that the hoxba and hoxbb clusters are specifically responsible

for the RA-mediated specification of pectoral fins.

***hoxb5a/b5b* expression in fin buds is RA-dependent**

The absence of pectoral fins, commonly observed in *hoxba;hoxbb* cluster

mutants and *raldh2* mutants, suggests that candidate hox genes may be downregulated

in *raldh2* mutants. To investigate this, we performed RNA-seq analysis to compare the

expression profiles between sibling and *raldh2* homozygous embryos. As expected, we

found that most of the 3'-located hox genes are downregulated in *raldh2*$^{-/-}$ embryos

(Figure 5A, B, figure supplement 2). Although *hoxc1a* showed the most pronounced

reduction, our analysis focused on *hoxba* and *hoxbb* clusters. Notably, the levels of

hoxb5b transcripts were significantly reduced, along with those of *hoxb5a* among hoxba

genes, in *raldh2* mutants. Previous studies have shown that zebrafish *hoxb5a* and

hoxb5b are expressed in the lateral plate mesoderm corresponding to presumptive

pectoral fin buds (Waxman et al., 2008). Additionally, knockout mice for *Hoxb5*, which

is homologous to zebrafish *hoxb5a* and *hoxb5b*, only exhibit altered positioning of

forelimbs (Rancourt et al., 1995), among various other Hox knockouts that have been

generated. Therefore, we examined whether RA regulates the expression of *hoxb5a* and

hoxb5b in fin buds. In wild-type embryos, *hoxb5a* expression was observed in the

neural tube, somites, and lateral plate mesoderm (Figure 5C). However, in *raldh2*

mutants, *hoxb5a* expression was significantly reduced, notably in the lateral plate

mesoderm, which differentiates into the fin buds, where it was reduced to the point of

being undetectable (Figure 5D). Since RA treatments in *raldh2* mutants can rescue

pectoral fin development, we found that *hoxb5a* expression in *raldh2* mutants is also

rescued by RA exposure (Figure 5C-E). Regarding the expression of *hoxb5b*, which

diverged from the same ancestral gene as *hoxb5a*, RA-dependent expression was also

confirmed, mirroring the observations made with *hoxb5a* (Figure 5F-H). These results

suggest that both *hoxb5a* and *hoxb5b* are involved in the formation of the pectoral fins

in zebrafish.

Genetic screen for hox genes specifying pectoral fin buds

To understand the molecular mechanisms underlying pectoral fin specification,

we sought to identify hox genes essential for fin formation through a genetic approach.

We anticipated that frameshift mutations in relevant hox genes would phenocopy the

absence of pectoral fins observed in *hoxba;hoxbb* mutants. Since hoxbb cluster contains

fewer hox genes than hoxba cluster (Figure 5A), we focused our investigation on hoxbb

cluster. Although *hoxb5b* is expressed in the presumptive fin buds (Figure 5F), other

hoxbb genes may also be involved. Therefore, we simultaneously injected three gRNAs

targeting *hoxb5b*, *hoxb6b*, and *hoxb8b* into hoxba hemizygous mutants, expecting

various germline mutations in these hoxbb genes in the injected fish (Figure 6A).

Subsequently, F1 fish with a hemizygous deletion of hoxba cluster and mosaic

mutations were crossed with *hoxba;hoxbb* hemizygous mutants. Among the progeny,

several embryos lacking pectoral fins were identified. Following genotyping and DNA

sequencing of the target hoxbb genes, we found that all embryos without pectoral fins

possessed frameshift mutations in *hoxb5b*, with *hoxba$^{-/-}$;hoxbb$^{+/-}$* being common among

them (Figure 6B-E, table-supplement 1, figure-supplement 3). Similarly, we injected

each crRNA for *hoxb6b* and *hoxb8b* into *hoxba$^{+/-}$* mutants and confirmed that neither

frameshift mutation in *hoxb6b* nor *hoxb8b* in *hoxba$^{-/-}$;hoxbb$^{+/-}$* led to the loss of pectoral

fins (Figure 6F, G, figure-supplement 3). These results suggest that the frameshift

mutation of *hoxb5b* alone can phenocopy the deletion of hoxbb cluster, recapitulating

the absence of pectoral fins in *hoxba;hoxbb* cluster mutants.

However, we encountered an unexpected result: $hoxba^{-/-};hoxb5b^{-/-}$ embryos,

which have a homozygous frameshift mutation in $hoxb5b$ and a homozygous deletion of

hoxba cluster, exhibited severely truncated but visibly present pectoral fins (Figure 6H,

I). After analyzing hundreds of embryos from several crossings, we did not find any

$hoxba^{-/-};hoxb5b^{-/-}$ embryos without pectoral fins. Consistent with this, the expression of

$tbx5a$ was detectable in the fin buds of $hoxba^{-/-};hoxb5b^{-/-}$ embryos (Figure 6K, L), in

contrast to its absence in $hoxba;hoxbb$ cluster mutants (Figure 1L). To examine the

potential contributions of other hoxbb genes, we observed the phenotypes of

$hoxba^{-/-};hoxb5b^{-/-}; hoxb6b^{-/-}$ embryos; however, these mutants did not exacerbate the

abnormalities and still formed pectoral fins (Figure 6J, M).

The phenotype of $hoxba^{-/-};hoxb5b^{-/-}$ embryos indicates that a frameshift

mutation in $hoxb5b$ does not fully mimic the complete deletion of the entire hoxbb

cluster, suggesting that other mechanisms may be involved. A novel genetic

compensation mechanism known as transcriptional adaptation has been identified:

mRNA containing a premature termination codon (PTC) can induce increased

expression of structurally related genes, thereby compensating for the function of

mutated genes (El-Brolosy et al., 2019; Rossi et al., 2015). To avoid producing

transcripts with PTC (Sztal and Stainier, 2020), we generated a full locus-deleted allele

of $hoxb5b$ using CRISPR-Cas9 with two crRNAs targeting both ends of the target locus

(figure-supplement 4). In contrast to the phenotype of $hoxba^{-/-};hoxb5b^{-/-}$, we found that

$hoxba^{-/-};hoxb5b^{de/dell}$ mutants lack pectoral fins and exhibit a significant reduction in

$tbx5a$ expression when intercrossing $hoxba^{+/-};hoxb5b^{+/del}$ fish (Figure 7A, B, D, E).

However, the occurrence rate of embryos lacking pectoral fins (n=3/120; 2.5 %) was

lower than predicted based on Mendelian genetics (1/16; 6.3 %). Other

hoxba$^{-/-}$;hoxb5b$^{del/del}$ embryos did exhibit shortened pectoral fins (Figure 7C).

Additionally, concerning hoxba cluster, we also generated frameshift-induced

hoxb5a$^{-/-}$;hoxb5b$^{-/-}$ and a full locus-deleted *hoxb5a$^{del/del}$;hoxb5b$^{del/del}$* embryos

(figure-supplement 4). *hoxb5a$^{-/-}$;hoxb5b$^{-/-}$* larvae consistently displayed normal pectoral

fins (Figure 7F). In contrast, *hoxb5a$^{del/del}$;hoxb5b$^{del/del}$* mutants exhibited a slightly more

pronounced phenotype of shortened pectoral fins, but no instances of missing pectoral

fins were identified (Figure 7G). To further clarify which Hox genes could potentially

contribute to fin bud formation, we analyzed the expression of PG4 and PG6–8 genes

from hoxba and hoxbb clusters using whole-mount *in situ* hybridization. Among these,

only *hoxb4a* showed detectable expression in the fin buds, whereas *hoxb6a*, *hoxb6b*,

hoxb7a, *hoxb8a*, and *hoxb8b* were not detectable (Figure 7M-R). Furthermore, due to

the functional redundancy observed between *hoxb4a* and *hoxb5a* observed in zebrafish

vertebral patterning and the RA-dependent expression of *hoxb4a* (Grandel et al., 2002;

Maeno et al., 2024), we generated frameshift-induced *hoxb4a$^{-/-}$;hoxb5a$^{-/-}$;hoxb5b$^{-/-}$*

embryos and an allele lacking the entire genomic region encompassing *hoxb4a* and

hoxb5a (figure-supplement 4). Although *hoxb4a$^{-/-}$;hoxb5a$^{-/-}$;hoxb5b$^{-/-}$* embryos did not

lack pectoral fins (Figure 7H), some *hoxb4a-b5a$^{del/del}$;hoxb5b$^{del/del}$* embryos did lack

pectoral fins, although at low penetrance (Figure 7I, *n*=3/397; 0.7%). The majority of

hoxb4a-b5a$^{del/del}$;hoxb5b$^{del/del}$ embryos displayed truncated pectoral fins (Figure 7J).

The lack of pectoral fins was not detected in *hoxb4a-b5a$^{del/del}$* and

hoxb4a$^{del/del}$;hoxb5b$^{del/del}$ mutants (Figure 7K, L, figure-supplement 4). These results

suggest that *hoxb4a*, *hoxb5a*, and *hoxb5b* may cooperatively contribute to the

anterior-posterior positioning of pectoral fin buds in zebrafish. Furthermore, our

findings reveal that transcriptional adaptation may partially play a role in genetic

compensation for frameshift-induced hox mutations during pectoral fin formation, while

also suggesting the potential involvement of other unknown mechanisms or additional

genomic regions in the initial induction of zebrafish pectoral fin buds.

Discussion

Since the establishment of gene targeting (Mansour et al., 1988), numerous

Hox knockout mice have been generated. However, neither single nor compound Hox

mutants, nor Hox overexpression approaches, have been reported to cause significant

defects in limb positioning (Jurberg et al., 2013; Moreau et al., 2019). Consequently, the

role of Hox genes in the positioning of vertebrate paired appendages remains unclear

due to a lack of direct genetic evidence. In this study, we provide genetic evidence using

zebrafish, demonstrating that specific combinatorial deletions of hoxba and hoxbb

clusters result in the absence of pectoral fins and a lack of induction of *tbx5a* expression

in the pectoral fin field. Based on our results, we propose a model in which Hox

expressions in the lateral plate mesoderm, regulated by RA synthesized in the paraxial

mesoderm, provide positional information and induce *tbx5a* expression in the

presumptive fin buds. Our genetic results shed light on longstanding fundamental

questions regarding the molecular mechanisms by which the positioning of paired

appendages is established in vertebrates.

hoxba/bb **clusters (possibly** *hoxb4a***,** *b5a***,** *b5b***) define the pectoral fin field.**

We showed that *hoxba;hoxba* mutants exhibit a lack of pectoral fins. Abnormal

expression of *tbx5a* in the lateral plate mesoderm, where pectoral fin precursor cells

normally arise, was evident from its initial expression in the mutants. We presume that

hox genes in hoxba and hoxbb clusters, induced by RA, define the region along the

anterior-posterior axis where pectoral fin formation can occur and promote the

expression of *tbx5a* in the lateral plate mesoderm. Importantly, *tbx5a* expression in the

fin bud was not induced even by RA exposure in *hoxba;hoxbb* mutants (Figure 4E-H).

In zebrafish, it has been suggested that RA synthesized in the paraxial mesoderm acts

directly on the lateral plate mesoderm, or that RA induces *tbx5a* expression via *wnt2b* in

the intermediate mesoderm (Gibert et al., 2006; Grandel and Brand, 2011; Neto et al.,

2012). However, our results suggest that intact *hoxba* and *hoxbb* clusters are required

for RA-dependent induction of *tbx5a* expression. One possible explanation is a linear

pathway in which RA induces hox expression, which then activates *tbx5a*. Alternatively,

RA may induce hox expression and, together with Hox proteins, act cooperatively to

activate *tbx5a*. Our data are consistent with both models. Through genetic studies, albeit

with low penetrance, we further showed that *hoxb4a*, *hoxb5a*, and *hoxb5b* play a

significant role in the induction of pectoral fin buds. The homologous Hoxb4 and

Hoxb5 proteins in mice have been shown to bind directly to the enhancer of *Tbx5*

(Minguillon et al., 2012). Moreover, in chickens, *Tbx5* expression in the forelimb field

is restricted by posterior Hox genes, preventing posterior expansion (Moreau et al.,

2019; Nishimoto et al., 2014). In contrast, our combinatorial deletions of zebrafish *hox*

clusters did not produce posterior expansion of *tbx5a* expression. We therefore propose

that, in zebrafish, *hoxb4a*, *hoxb5a*, and *hoxb5b*, induced by RA in the paraxial

mesoderm, act primarily to promote *tbx5a* expression in the pectoral fin field.

Partial phenocopy of locus-deleted hox mutants implies other unidentified

mechanisms.

In this study, we observed that hox mutants with deletions of genomic loci

exhibited more pronounced pectoral fin phenotypes than frameshift-induced hox

mutants. The frameshift mutants used in this study are likely to be loss-of-function

alleles, as we demonstrated several defects in vertebral patterning in these mutants

(Maeno et al., 2024). These results suggest that transcriptional adaptation may occur

during the formation of pectoral fins in frameshift-induced hox mutants, potentially

compensating for the loss of function. In our previous studies, which focused on dorsal

and anal fin patterning, vertebral patterning, and swim bladder formation, we

demonstrated that the phenotypes observed in zebrafish hox cluster-deleted mutants

could essentially be replicated by introducing frameshift mutations into hox genes

within the cluster (Adachi et al., 2024; Maeno et al., 2024; Satoh et al., 2024; Yamada

et al., 2021). Therefore, creating a mutant with a deleted locus that does not induce

transcriptional adaptation was unnecessary. One factor that may explain the differences

in results between this study and our prior findings is that even a small amount of hox

gene products essential for pectoral fin positioning may be sufficient to induce fin buds,

as shown in Figure 1, potentially leading to the manifestation of gene compensation

through transcriptional adaptation in frameshift-induced mutants. Alternatively,

transcriptional adaptation may function differently depending on various developmental

processes or stages. Interestingly, all the aforementioned phenotypes reproducible by

frameshift mutations were observed at later developmental stages than pectoral fin

formation. Additionally, in hox locus-deleted mutants, only some *hoxba$^{-/-}$;hoxb5b$^{de/dell}$*

or *hoxb4a-b5a$^{del/del}$;hoxb5b$^{del/del}$* embryos were able to reproduce the loss of the pectoral

fin, suggesting the presence of other compensatory mechanisms. The organized

expression patterns of Hox genes are thought to be regulated by complex regulatory

controls, with their expression regulated according to the genomic order within the

cluster. In the absence of Hox function, neighboring intact Hox genes could compensate

for the loss of function. It is also possible that unidentified non-coding regions in hoxba

and hoxbb clusters are involved in pectoral fin positioning. These issues should be

clarified in future studies.

Hox genes and the evolutionary origin of pectoral fins.

According to fossil records from the Cambrian period, the earliest vertebrates

did not possess paired fins (Janvier, 1996; Shubin et al., 1997). Later in evolution,

vertebrates acquired primitive pectoral fins, which served as evolutionary precursors to

tetrapod forelimbs, before diverging into ray-finned and lobe-finned fishes. Our results

align well with this evolutionary scenario. Hox genes, which are conserved across

bilaterian animals, were already present in primitive vertebrates without paired fins,

suggesting that their functional diversification may have facilitated the origin of new

appendages. The early fossils of vertebrates with pectoral fins were discovered in

Ordovician and Silurian jawless fishes (Janvier, 1996; Shubin et al., 1997). Our finding

that *hoxba;hoxbb* mutants completely lack pectoral fins implies that molecular

evolution within the ancestral HoxB cluster contributed to the emergence of pectoral

fins. Although the precise molecular modifications remain unclear, one plausible model

is that HoxB genes acquired the capacity to induce *Tbx5* expression in the lateral plate

mesoderm, establishing the RA–HoxB–Tbx5 gene network essential for fin initiation.

We presume that the acquisition of pectoral fins was driven by multiple genomic

changes accumulated over time, among which molecular evolution within the HoxB

cluster may have represented one of the principal factors contributing to the origin of

this novel appendage.

The positioning of pectoral fins is restricted to specific hox clusters in zebrafish, in

contrast to the functional overlap seen in mice.

Taking advantage of our availability of seven individual hox cluster mutants in

zebrafish, we generated multiple cluster deletions and demonstrated that hox genes

required for establishing pectoral fin buds are restricted to hoxba and hoxbb clusters. In

contrast, analyses of knockout mice have revealed extensive functional overlap among

the four Hox clusters. For instance, mice lacking all HoxB genes except for *Hoxb13* do

not lose forelimbs (Medina-Martinez et al., 2000), and forelimbs are still present in

Hoxa5;Hoxb5;Hoxc5 triple knockout mice, which lack all Hox genes of the paralog

group 5 (Xu et al., 2013). Thus, while zebrafish *hoxba;hoxbb* mutants exhibit a striking

phenotype of absent pectoral fins, forelimbs are preserved in comparable mouse mutants.

We speculate that paired fins emerged after the quadruplication of primitive Hox

clusters, and that novel functions related to pectoral fin formation arose within the

HoxB cluster. In zebrafish, these functions remain confined to hoxba and hoxbb clusters,

making them indispensable for pectoral fin development. Interestingly, a similar

cluster-specific confinement has been observed in median fins: in both zebrafish and

medaka, Hox genes responsible for positioning dorsal and anal fins are primarily

located in HoxC-related clusters (Adachi et al., 2024). In contrast, in mice, functional

redundancy among Hox clusters—likely accumulated during over half a billion years of

vertebrate evolution—may mask the requirement of HoxB genes for forelimb formation.

While the precise evolutionary mechanisms remain to be clarified, our results suggest

that cluster-specific specialization in teleosts contrasts with broader redundancy across

Hox clusters in mammals.

Materials and methods

Zebrafish

Riken Wild-type (RW) zebrafish, obtained from the National BioResource Project in Japan, were maintained at 27°C with a 14-hour light/10-hour dark cycle. Embryos were collected from natural spawning, and the larvae were raised at 28.5°C. Developmental stages of the embryos and larvae were determined based on hours post-fertilization (hpf), days post-fertilization (dpf), and developmental stage-specific features (Kimmel et al., 1995; Parichy et al., 2009). The alleles of *hox* cluster-deleted mutants used in this study were previously generated using CRISPR-Cas9 (Yamada et al., 2021). The following frameshift-induced *hox* mutants were created using CRISPR-Cas9: *hoxb5a*[sud125], *hoxb5b*[sud136], *hoxb6b*[sud137], and *hoxb4a;hoxb5a*[sud144] (Maeno et al., 2024). All experiments involving live zebrafish were approved by the Committee for Animal Care and Use of Saitama University.

Generation of mutants by the CRISPR-Cas9 system

All the mutants used in this study were generated using the Alt-R CRISPR-Cas9 system (Integrated DNA Technologies). To create frameshift-induced mutants, gene-specific crRNAs were incubated with common tracrRNA, followed by Cas9 nuclease. For the generation of mutants that delete large genomic regions, two crRNAs targeting both ends of the regions were incubated with common tracrRNA and Cas9 nuclease. Approximately one nanoliter of the crRNA:tracrRNA-Cas9 RNA-protein complex was injected into fertilized embryos, which were subsequently raised to juvenile fish. Candidate founder fish were selected using a heteroduplex

mobility shift assay for frameshift mutants (Ota et al., 2013) and by amplifying genomic

deletions with specific primers. After sexual maturation, candidate founder fish were

mated with wild-type fish to produce heterozygous F1 offspring. Among the F1

offspring, mutant fish carrying the same mutation were identified through PCR-based

genotyping followed by DNA sequencing. The target-specific sequences of the crRNAs

used in this study are listed (table-supplement 2). The frozen sperm from the mutants

used in this study have been deposited in the National BioResource Project Zebrafish in

Japan (https://shigen.nig.ac.jp/zebra/) and are available upon request.

Genotyping of mutants

For the phenotypic analysis of embryos and the maintenance of mutant fish

lines, PCR-based genotyping was performed. Briefly, genomic DNA was extracted from

the analyzed embryos or the partially dissected caudal fins of anesthetized larvae or fish

using the NaOH method (Meeker et al., 2007), which served as the template for PCR.

The sequences of the primers used for genotyping frameshift-induced mutants are listed

in the supplementary Table 3. For genotyping deletion mutants of *hox* genes, PCR was

conducted using a combination of three primers, with their sequences also provided

(table-supplement 3). Genotyping of *hox* cluster-deleted mutants was performed using

PCR as previously described (Yamada et al., 2021). After the reactions, the PCR

products were separated by electrophoresis. Based on differences in the lengths of PCR

products derived from wild-type and mutated alleles, either a 2 % agarose gel in 0.5x

TBE buffer, a 15 % polyacrylamide gel in 0.5x TBE buffer, or direct DNA sequencing

was utilized to determine the genotype.

Whole-mount *in situ* hybridization

Whole-mount *in situ* hybridization was carried out as described (Thisse and Thisse, 2014). After the staining, the embryos mounted in 70-80 % glycerol were captured under the stereomicroscope (Leica M205 FA) with a digital camera (Leica DFC350F). After taking images, PCR-based genotyping was carried out as described above.

RNA-seq analysis

For RNA-seq analysis, embryos were obtained by the intercrosses between *raldh2* heterozygous fish. At the 18-20-somite stages, sibling and homozygous embryos were separated based on the phenotype, and 20 embryos per tube were collected. Then, total RNA was extracted by ISOGENE (Nippon Gene) and followed by DNase I treatment (Takara). After the quality check of the isolated RNA, RNA-seq analyses (n=2 for each) were carried out by Genewiz, Azenta Life Science.

Treatment of retinoic acid in the embryos

Treatment of retinoic acid on the embryos was performed as previously described, with minor modifications (Gibert et al., 2006; Grandel et al., 2002). Briefly, embryos were obtained from intercrosses between *raldh2* heterozygous mutants. At the 70 % epiboly stage, the embryos were soaked in a solution containing DMSO or 10^{-8}M RA and incubated at 28.5°C. Embryos were fixed at 30 hpf for analysis of *tbx5a* expression and at the 10-somite stage for examination of *hox* gene expression.

Acknowledgments

We thank the NBRP zebrafish for providing fish and preserving the mutant

lines generated in this study. We also appreciate Drs. Koji Tamura and Gembu Abe for

providing the zebrafish *tbx5a* plasmid for *in situ* hybridization. This work was

supported by KAKENHI Grants-in-Aid for Scientific Research from the Ministry of

Education, Culture, Sports, Science, and Technology, Japan (18K06177, 23K05790 to

AK).

References

Adachi, U., Koita, R., Seto, A., Maeno, A., Ishizu, A., Oikawa, S., Tani, T., Ishizaka, M., Yamada, K., Satoh, K.*, et al.* (2024). Teleost Hox code defines regional identities competent for the formation of dorsal and anal fins. Proc Natl Acad Sci U S A *121*, e2403809121.

Ahn, D.G., Kourakis, M.J., Rohde, L.A., Silver, L.M., and Ho, R.K. (2002). T-box gene tbx5 is essential for formation of the pectoral limb bud. Nature *417*, 754-758.

Amores, A., Force, A., Yan, Y.L., Joly, L., Amemiya, C., Fritz, A., Ho, R.K., Langeland, J., Prince, V., Wang, Y.L.*, et al.* (1998). Zebrafish hox clusters and vertebrate genome evolution. Science *282*, 1711-1714.

Begemann, G., Schilling, T.F., Rauch, G.J., Geisler, R., and Ingham, P.W. (2001). The zebrafish neckless mutation reveals a requirement for raldh2 in mesodermal signals that pattern the hindbrain. Development *128*, 3081-3094.

Boncinelli, E., Simeone, A., Acampora, D., and Mavilio, F. (1991). HOX gene activation by retinoic acid. Trends Genet *7*, 329-334.

Boulet, A.M., and Capecchi, M.R. (2004). Multiple roles of Hoxa11 and Hoxd11 in the formation of the mammalian forelimb zeugopod. Development *131*, 299-309.

Boyle-Anderson, E.A.T., Mao, Q., and Ho, R.K. (2022). Tbx5a and Tbx5b paralogues act in combination to control separate vectors of migration in the fin field of zebrafish. Dev Biol *481*, 201-214.

Burke, A.C., Nelson, C.E., Morgan, B.A., and Tabin, C. (1995). Hox genes and the evolution of vertebrate axial morphology. Development *121*, 333-346.

Chen, J.N., and Fishman, M.C. (1996). Zebrafish tinman homolog demarcates the heart field and initiates myocardial differentiation. Development *122*, 3809-3816.

Cohn, M.J., Patel, K., Krumlauf, R., Wilkinson, D.G., Clarke, J.D., and Tickle, C. (1997). Hox9 genes and vertebrate limb specification. Nature *387*, 97-101.

Davis, A.P., Witte, D.P., Hsieh-Li, H.M., Potter, S.S., and Capecchi, M.R. (1995). Absence of radius and ulna in mice lacking hoxa-11 and hoxd-11. Nature *375*, 791-795.

Dehal, P., and Boore, J.L. (2005). Two rounds of whole genome duplication in the ancestral vertebrate. PLoS Biol *3*, e314.

Dolle, P., Izpisua-Belmonte, J.C., Falkenstein, H., Renucci, A., and Duboule, D. (1989). Coordinate expression of the murine Hox-5 complex homoeobox-containing genes during limb pattern formation. Nature *342*, 767-772.

Duboule, D., and Dolle, P. (1989). The structural and functional organization of the murine HOX gene family resembles that of Drosophila homeotic genes. EMBO J *8*, 1497-1505.

El-Brolosy, M.A., Kontarakis, Z., Rossi, A., Kuenne, C., Gunther, S., Fukuda, N., Kikhi, K.,

Boezio, G.L.M., Takacs, C.M., Lai, S.L., *et al.* (2019). Genetic compensation triggered by

mutant mRNA degradation. Nature *568*, 193-197.

Fromental-Ramain, C., Warot, X., Lakkaraju, S., Favier, B., Haack, H., Birling, C., Dierich, A.,

Doll e, P., and Chambon, P. (1996a). Specific and redundant functions of the paralogous Hoxa-9

and Hoxd-9 genes in forelimb and axial skeleton patterning. Development *122*, 461-472.

Fromental-Ramain, C., Warot, X., Messadecq, N., LeMeur, M., Dolle, P., and Chambon, P.

(1996b). Hoxa-13 and Hoxd-13 play a crucial role in the patterning of the limb autopod.

Development *122*, 2997-3011.

Garrity, D.M., Childs, S., and Fishman, M.C. (2002). The heartstrings mutation in zebrafish

causes heart/fin Tbx5 deficiency syndrome. Development *129*, 4635-4645.

Gibert, Y., Gajewski, A., Meyer, A., and Begemann, G. (2006). Induction and prepatterning of

the zebrafish pectoral fin bud requires axial retinoic acid signaling. Development *133*,

2649-2659.

Graham, A., Papalopulu, N., and Krumlauf, R. (1989). The murine and Drosophila homeobox

gene complexes have common features of organization and expression. Cell *57*, 367-378.

Grandel, H., and Brand, M. (2011). Zebrafish limb development is triggered by a retinoic acid

signal during gastrulation. Dev Dyn *240*, 1116-1126.

Grandel, H., Lun, K., Rauch, G.J., Rhinn, M., Piotrowski, T., Houart, C., Sordino, P., Kuchler,

A.M., Schulte-Merker, S., Geisler, R., *et al.* (2002). Retinoic acid signalling in the zebrafish

embryo is necessary during pre-segmentation stages to pattern the anterior-posterior axis of the

CNS and to induce a pectoral fin bud. Development *129*, 2851-2865.

Horan, G.S., Ramirez-Solis, R., Featherstone, M.S., Wolgemuth, D.J., Bradley, A., and

Behringer, R.R. (1995). Compound mutants for the paralogous hoxa-4, hoxb-4, and hoxd-4

genes show more complete homeotic transformations and a dose-dependent increase in the

number of vertebrae transformed. Genes Dev *9*, 1667-1677.

Iimura, T., and Pourquie, O. (2007). Hox genes in time and space during vertebrate body

formation. Dev Growth Differ *49*, 265-275.

Ishizaka, M., Maeno, A., Nakazawa, H., Fujii, R., Oikawa, S., Tani, T., Kanno, H., Koita, R.,

and Kawamura, A. (2024). The functional roles of zebrafish HoxA- and HoxD-related clusters

in the pectoral fin development. Sci Rep *14*, 23602.

Izpisua-Belmonte, J.C., and Duboule, D. (1992). Homeobox genes and pattern formation in the

vertebrate limb. Dev Biol *152*, 26-36.

Janvier, P. (1996). Early Vertebrates.

Jurberg, A.D., Aires, R., Varela-Lasheras, I., Novoa, A., and Mallo, M. (2013). Switching axial

progenitors from producing trunk to tail tissues in vertebrate embryos. Dev Cell *25*, 451-462.

Kimmel, C.B., Ballard, W.W., Kimmel, S.R., Ullmann, B., and Schilling, T.F. (1995). Stages of

embryonic development of the zebrafish. Dev Dyn *203*, 253-310.

Kmita, M., Tarchini, B., Zakany, J., Logan, M., Tabin, C.J., and Duboule, D. (2005). Early

developmental arrest of mammalian limbs lacking HoxA/HoxD gene function. Nature *435*,

1113-1116.

Krumlauf, R. (1994). Hox genes in vertebrate development. Cell *78*, 191-201.

Langston, A.W., and Gudas, L.J. (1994). Retinoic acid and homeobox gene regulation. Curr

Opin Genet Dev *4*, 550-555.

Maeno, A., Koita, R., Nakazawa, H., Fujii, R., Yamada, K., Oikawa, S., Tani, T., Ishizaka, M.,

Satoh, K., Ishizu, A.*, et al.* (2024). The Hox code responsible for the patterning of the anterior

vertebrae in zebrafish. Development *151*.

Mansour, S.L., Thomas, K.R., and Capecchi, M.R. (1988). Disruption of the proto-oncogene

int-2 in mouse embryo-derived stem cells: a general strategy for targeting mutations to

non-selectable genes. Nature *336*, 348-352.

Marshall, H., Morrison, A., Studer, M., Popperl, H., and Krumlauf, R. (1996). Retinoids and

Hox genes. FASEB J *10*, 969-978.

Matsubara, Y., Hirasawa, T., Egawa, S., Hattori, A., Suganuma, T., Kohara, Y., Nagai, T.,

Tamura, K., Kuratani, S., Kuroiwa, A.*, et al.* (2017). Anatomical integration of the

sacral-hindlimb unit coordinated by GDF11 underlies variation in hindlimb positioning in

tetrapods. Nat Ecol Evol *1*, 1392-1399.

McIntyre, D.C., Rakshit, S., Yallowitz, A.R., Loken, L., Jeannotte, L., Capecchi, M.R., and

Wellik, D.M. (2007). Hox patterning of the vertebrate rib cage. Development *134*, 2981-2989.

Medina-Martinez, O., Bradley, A., and Ramirez-Solis, R. (2000). A large targeted deletion of

Hoxb1-Hoxb9 produces a series of single-segment anterior homeotic transformations. Dev Biol

222, 71-83.

Meeker, N.D., Hutchinson, S.A., Ho, L., and Trede, N.S. (2007). Method for isolation of

PCR-ready genomic DNA from zebrafish tissues. BioTechniques *43*, 610, 612, 614.

Minguillon, C., Nishimoto, S., Wood, S., Vendrell, E., Gibson-Brown, J.J., and Logan, M.P.

(2012). Hox genes regulate the onset of Tbx5 expression in the forelimb. Development *139*,

3180-3188.

Moreau, C., Caldarelli, P., Rocancourt, D., Roussel, J., Denans, N., Pourquie, O., and Gros, J.

(2019). Timed Collinear Activation of Hox Genes during Gastrulation Controls the Avian

Forelimb Position. Curr Biol *29*, 35-50 e34.

Murata, Y., Tamura, M., Aita, Y., Fujimura, K., Murakami, Y., Okabe, M., Okada, N., and

Tanaka, M. (2010). Allometric growth of the trunk leads to the rostral shift of the pelvic fin in

teleost fishes. Dev Biol *347*, 236-245.

Neto, A., Mercader, N., and Gomez-Skarmeta, J.L. (2012). The Osr1 and Osr2 genes act in the

pronephric anlage downstream of retinoic acid signaling and upstream of Wnt2b to maintain

pectoral fin development. Development *139*, 301-311.

Ng, J.K., Kawakami, Y., Buscher, D., Raya, A., Itoh, T., Koth, C.M., Rodriguez Esteban, C.,

Rodriguez-Leon, J., Garrity, D.M., Fishman, M.C.*, et al.* (2002). The limb identity gene Tbx5

promotes limb initiation by interacting with Wnt2b and Fgf10. Development *129*, 5161-5170.

Nishimoto, S., and Logan, M.P. (2016). Subdivision of the lateral plate mesoderm and

specification of the forelimb and hindlimb forming domains. Semin Cell Dev Biol *49*, 102-108.

Nishimoto, S., Minguillon, C., Wood, S., and Logan, M.P. (2014). A combination of activation

and repression by a colinear Hox code controls forelimb-restricted expression of Tbx5 and

reveals Hox protein specificity. PLoS Genet *10*, e1004245.

Ohno, S. (1970). Evolution by gene duplication.

Ota, S., Hisano, Y., Muraki, M., Hoshijima, K., Dahlem, T.J., Grunwald, D.J., Okada, Y., and

Kawahara, A. (2013). Efficient identification of TALEN-mediated genome modifications using

heteroduplex mobility assays. Genes Cells *18*, 450-458.

Parichy, D.M., Elizondo, M.R., Mills, M.G., Gordon, T.N., and Engeszer, R.E. (2009). Normal

table of postembryonic zebrafish development: staging by externally visible anatomy of the

living fish. Dev Dyn *238*, 2975-3015.

Rancourt, D.E., Tsuzuki, T., and Capecchi, M.R. (1995). Genetic interaction between hoxb-5

and hoxb-6 is revealed by nonallelic noncomplementation. Genes Dev *9*, 108-122.

Rossi, A., Kontarakis, Z., Gerri, C., Nolte, H., Holper, S., Kruger, M., and Stainier, D.Y. (2015).

Genetic compensation induced by deleterious mutations but not gene knockdowns. Nature *524*,

230-233.

Royle, S.R., Tabin, C.J., and Young, J.J. (2021). Limb positioning and initiation: An

evolutionary context of pattern and formation. Dev Dyn *250*, 1264-1279.

Satoh, K., Maeno, A., Adachi, U., Ishizaka, M., Yamada, K., Koita, R., Nakazawa, H., Oikawa,

S., Fujii, R., Furudate, H.*, et al.* (2024). Physical constraints on the positions and dimensions of

the zebrafish swim bladder by surrounding bones. J Anat.

Serbedzija, G.N., Chen, J.N., and Fishman, M.C. (1998). Regulation in the heart field of

zebrafish. Development *125*, 1095-1101.

Shimada, A., Kawanishi, T., Kaneko, T., Yoshihara, H., Yano, T., Inohaya, K., Kinoshita, M.,

Kamei, Y., Tamura, K., and Takeda, H. (2013). Trunk exoskeleton in teleosts is mesodermal in

origin. Nat Commun *4*, 1639.

Shubin, N., Tabin, C., and Carroll, S. (1997). Fossils, genes and the evolution of animal limbs.

Nature *388*, 639-648.

Sztal, T.E., and Stainier, D.Y.R. (2020). Transcriptional adaptation: a mechanism underlying

genetic robustness. Development *147*.

Tanaka, M. (2013). Molecular and evolutionary basis of limb field specification and limb

initiation. Dev Growth Differ *55*, 149-163.

Thisse, B., and Thisse, C. (2014). In situ hybridization on whole-mount zebrafish embryos and

young larvae. Methods Mol Biol *1211*, 53-67.

Tickle, C. (2015). How the embryo makes a limb: determination, polarity and identity. J Anat

227, 418-430.

van den Akker, E., Fromental-Ramain, C., de Graaff, W., Le Mouellic, H., Brulet, P., Chambon,

P., and Deschamps, J. (2001). Axial skeletal patterning in mice lacking all paralogous group 8

Hox genes. Development *128*, 1911-1921.

Waxman, J.S., Keegan, B.R., Roberts, R.W., Poss, K.D., and Yelon, D. (2008). Hoxb5b acts

downstream of retinoic acid signaling in the forelimb field to restrict heart field potential in

zebrafish. Dev Cell *15*, 923-934.

Wellik, D.M., and Capecchi, M.R. (2003). Hox10 and Hox11 genes are required to globally

pattern the mammalian skeleton. Science *301*, 363-367.

Wellik, D.M., Hawkes, P.J., and Capecchi, M.R. (2002). Hox11 paralogous genes are essential

for metanephric kidney induction. Genes Dev *16*, 1423-1432.

Woltering, J.M., and Durston, A.J. (2006). The zebrafish hoxDb cluster has been reduced to a

single microRNA. Nat Genet *38*, 601-602.

Xu, B., Hrycaj, S.M., McIntyre, D.C., Baker, N.C., Takeuchi, J.K., Jeannotte, L., Gaber, Z.B.,

Novitch, B.G., and Wellik, D.M. (2013). Hox5 interacts with Plzf to restrict Shh expression in

the developing forelimb. Proc Natl Acad Sci U S A *110*, 19438-19443.

Yamada, K., Maeno, A., Araki, S., Kikuchi, M., Suzuki, M., Ishizaka, M., Satoh, K., Akama, K.,

Kawabe, Y., Suzuki, K.*, et al.* (2021). An atlas of seven zebrafish hox cluster mutants provides

insights into sub/neofunctionalization of vertebrate Hox clusters. Development *148*.

Yelon, D., Horne, S.A., and Stainier, D.Y. (1999). Restricted expression of cardiac myosin genes

reveals regulated aspects of heart tube assembly in zebrafish. Dev Biol *214*, 23-37.

Figure legends

Figure 1.   Lack of pectoral fins in *hoxba;hoxbb* cluster-deleted mutants. (A-G)

Dorsal views of live zebrafish larvae at 3 dpf, obtained from intercrosses between

hoxba;hoxbb cluster-deficient hemizygous fish. Arrowheads indicate the positions of the

pectoral fins. (H-M) Expression patterns of *tbx5a* in the pectoral fin bud (arrowhead) at

30 hpf. Dorsal views are shown, and the genotype of each specimen was determined.

For each genotype, reproducibility was confirmed with at least three different specimens.

Genotyping revealed that all the embryos lacking pectoral fins (*n*=15) were

hoxba;hoxbb double homozygotes. Scale bars are 100 μm.

**Figure 2.   Significantly decreased expression of *tbx5a* in the pectoral fin buds is

specific to *hoxba;hoxbb* cluster-deleted mutants.**

(A-I) Expression patterns of *tbx5a* in the pectoral fin buds of combinatorial deletion

mutants of zebrafish *hox* clusters. Dorsal views of embryos at 30 hpf are displayed.

After capturing images, the genotype of each specimen was determined. For each

genotype, reproducibility was confirmed with at least three different specimens. Scale

bars are 100 μm.

**Figure 3.   Loss of fin progenitors with posterior expansion of cardiac marker

expression in *hoxba;hoxbb* mutants.** (A-F) Expression patterns of *tbx5a* were

compared between sibling wild-type and *hoxba;hoxbb* homozygous embryos during

embryogenesis. The range of *tbx5a* expression in the lateral mesoderm is indicated by a

bracket. The progenitor cells of pectoral fins are indicated by an arrowhead. (G, H)

Expression patterns of *nkx2.5* were compared between sibling and *hoxba;hoxbb* mutants.

Enlarged views are shown in (G', H'). The different regions of *nkx2.5* expression

between wild-type and mutants are indicated by brackets. (I, J) Expression patterns of

cmlc2 are shown. All images were captured from the dorsal side. For each stage,

reproducibility was confirmed with at least three different specimens. Scale bars

indicate 100 μm.

Figure 4.   Zebrafish *hoxba;hoxbb* mutants lack a response to RA in fin buds.

(A-D) Exogenous RA exposure in *raldh2* mutants can rescue *tbx5a* expression in fin

buds. Arrowheads indicate the positions of the pectoral fin buds. (E-H) RA treatments in

hoxba;hoxbb cluster-deleted mutants do not rescue *tbx5a* expression in fin buds. (I, J)

Expression patterns of *raldh2* were compared between sibling and *hoxba;hoxbb* mutants.

For each genotype, reproducibility was confirmed with at least three different specimens.

All images were captured from the dorsal side. Scale bars represent 100 μm.

Figure 5.   Expression patterns of *hox5a* and *hoxb5b* are regulated by RA.

(A) Schematic representation of the 49 *hox* genes organized into seven *hox* clusters in

zebrafish. *hox* genes in *hoxba* and *hoxbb* clusters are highlighted with orange. (B)

Expression profiles of the 49 *hox* genes in wild-type and *raldh2*$^{-/-}$ embryos at the

20-somite stage, analyzed by RNA-seq. The average FPKM of each *hox* gene was

compared between sibling and *raldh2* mutants. The absence of specific *hox* genes is

indicated by a slash. (C-H) Expression patterns of *hoxb5a* and *hoxb5b* in sibling

wild-type, *raldh2*$^{-/-}$, and RA-treated *raldh2*$^{-/-}$ embryos at the 10-somite stage.

Arrowheads indicate the presumptive positions of pectoral fin buds. For each genotype,

reproducibility was confirmed with at least three different specimens. All images are

captured from the dorsal side. Scale bars represent 100 μm.

Figure 6.   Screening for hox genes responsible for the zebrafish pectoral fin

formation.

(A) Schematic representation of the genetic screening for hox genes involved in the

specification of pectoral fins in zebrafish. (B-G) Dorsal views of live mutant larvae at 3

dpf were obtained during the screening. After capturing images, genotyping and DNA

sequencing in target hoxb genes were conducted. For the mutants illustrated in (C-G),

hoxb genes are shown with frameshift mutations introduced. Detailed information is

provided in Supplementary Table S1. (H-J) Dorsal views of *hoxba^{-/-};hoxb5b^{-/-}* and

hoxba^{-/-};hoxb5b^{-/-};hoxb6b^{-/-} larvae at 3 dpf. (K-M) Expression patterns of *tbx5a* in

hoxba^{-/-};hoxb5b^{-/-} and *hoxba^{-/-};hoxb5b^{-/-};hoxb6b^{-/-}* mutants at 30 hpf. Arrowheads

indicate the presumptive positions of pectoral fin buds. All images are captured from the

dorsal side. Scale bars represent 100 μm.

Figure 7.   *hoxb4a-b5a^{del/del};hoxb5b^{del/del}* larvae partially recapitulate the absence of

the pectoral fins.

(A-C) Dorsal views of zebrafish *hoxba^{-/-};hoxb5b^{del/del}* larvae at 3 dpf. (D, E) Expression

patterns of *tbx5a* in sibling wild-type and *hoxba^{-/-};hoxb5b^{del/del}* (*n*=4) at 30 hpf.

Arrowheads indicate the presumptive positions of pectoral fin buds. (F, G) Dorsal view

of frameshift-induced *hoxb5a^{-/-};hoxb5b^{-/-}* and *hoxb5a^{del/del};hoxb5b^{del/del}* larvae. (I-K)

Dorsal view of frameshift-induced *hoxb4a^{-/-};hoxb5a^{-/-};hoxb5b^{-/-}* and *hoxb4a-b5a*

^{del/del};hoxb5b^{del/del} larvae. All images are captured from the dorsal side. Arrowheads

indicate the presumptive positions of pectoral fin buds. (M-R) Expression patterns of

zebrafish *hoxb4a, hoxb6a, hoxb6b, hoxb7a, hoxb8a,* and *hoxb8b* at the 10-somite stage.

Dorsal views. Arrowheads indicate the presumptive positions of pectoral fin buds. Scale

bars represent 100 μm.

Figure S1.   Generation of the zebrafish *raldh2* mutants by CRISPR-Cas9. (A) The

nucleotide sequence surrounding the mutations in *raldh2* is shown on the left. The target

sequence of the crRNA is emphasized with an underline, while the inserted nucleotide

sequence is indicated in orange. Below the DNA sequence, the predicted amino acid is

displayed, with red letters highlighting the abnormal amino acid resulting from the

frameshift mutation. The asterisk represents the termination codon. (B) A schematic

representation of the predicted protein structure is shown, with the red box indicating

the abnormal amino acid sequences resulting from the frameshift mutations. The total

number of predicted amino acids is displayed to the right of the schematic. (C) The

absence of pectoral fins (arrowhead) in *raldh2* homozygous mutants is observed

dorsally at 3 dpf. (D) The expression of *tbx5a* in the pectoral fin buds (arrowhead) is

significantly reduced in *raldh2* mutants. The phenotype of our *raldh2*^*sud118*^ mutants

closely resembles the phenotype observed in previously isolated *raldh2* mutants.

Figure S2.   Volcano plot of the transcriptome analysis between sibling and *raldh2*

mutants. The volcano plot illustrates both upregulated and downregulated differentially

expressed genes from the comparison between wild-type and *raldh2* mutant embryos.

Dot with more than a 2-fold increase in expression is indicated in orange, while those

with more than a 2-fold decrease are shown in light blue. Among them, dots

corresponding to *hox* genes are specifically highlighted.

Figure S3.   Generation of the frameshift-induced hox mutants using

CRISPR-Cas9. Schematic representations show zebrafish *hoxb8b, hoxb5b;hoxb6b,* and

hoxb5b;hoxb6b;hoxb8b mutants. The other frameshift-induced hox mutants were

previously described (Maeno et al., 2024). The nucleotide sequence around the

mutations is shown on the left. The target sequence of crRNA is emphasized with an

orange underline. The inserted nucleotide sequence is emphasized in red. Below the

DNA sequence, the predicted amino acid is shown. Red letters indicate the abnormal

amino acid caused by the frameshift mutation. On the right, a schema represents the

predicted protein structure. The blue box indicates the homeodomain, and the red box

indicates the abnormal amino acid sequences from the frameshift mutations. The total

number of predicted amino acids is shown on the right of the schema.

Figure S4. Generation of locus-deletion mutants using CRISPR-Cas9. (A, C, E, G).

Schematic representations show the genomic structure of zebrafish *hoxb4a, hox5a,* and

hoxb5b. Boxes represent the exons, with the blue boxes indicating the coding regions.

Black arrowheads indicate the target sites of crRNA, while red arrowheads represent the

primers used for genotyping. (B, D, F, H) Display genomic locus deletions of *hox4a,*

hox5a, hoxb5b, and *hoxb4a-b5a* deletion mutants. The flanking sequences of the 5'- and

3'-crRNA targets (underline) are shown in blue and green letters, respectively.

Figure 1. Lack of pectoral fins in *hoxba;hoxbb* cluster-deleted mutants.
(A-F) Dorsal views of live zebrafish larvae at 3 dpf, obtained from
intercrosses between *hoxba;hoxbb* cluster-deficient hemizygous fish.
Arrowheads indicate the positions of the pectoral fins. (G-L) Expression
patterns of *tbx5a* in the pectoral fin bud (arrowhead) at 30 hpf. Dorsal views
are shown, and the genotype of each specimen was determined. For each
genotype, reproducibility was confirmed with at least three different
specimens. Genotyping revealed that all the embryos lacking pectoral fins
(*n*=15) were *hoxba;hoxbb* double homozygotes. Scale bars are 100 μm.

Figure 2. Significantly decreased expression of *tbx5a* in the pectoral fin buds is specific to *hoxba;hoxbb* cluster-deleted mutants.

(A-I) Expression patterns of *tbx5a* in the pectoral fin buds of combinatorial deletion mutants of zebrafish *hox* clusters. Dorsal views of embryos at 30 hpf are displayed. After capturing images, the genotype of each specimen was determined. For each genotype, reproducibility was confirmed with at least three different specimens. Scale bars are 100 μm.

Figure 3. Loss of fin progenitors with posterior expansion of cardiac marker expression in hoxba;hoxbb mutants. (A-F) Expression patterns of *tbx5a* were compared between sibling wild-type and *hoxba;hoxbb* homozygous embryos during embryogenesis. The range of *tbx5a* expression in the lateral mesoderm is indicated by a bracket. The progenitor cells of pectoral fins are indicated by an arrowhead. (G, H) Expression patterns of *nkx2.5* were compared between sibling and *hoxba;hoxbb* mutants. Enlarged views are shown in (G', H'). The different regions of *nkx2.5* expression between wild-type and mutants are indicated by brackets. (I, J) Expression patterns of *cmcl2* are shown. All images were captured from the dorsal side. For each stage, reproducibility was confirmed with at least three different specimens. Scale bars indicate 100 μm.

Figure 4. Zebrafish *hoxba;hoxbb* mutants lack a response to RA in fin buds. (A-D) Exogenous RA exposure in *raldh2* mutants can rescue *tbx5a* expression in fin buds. Arrowheads indicate the positions of the pectoral fin buds. (E-H) RA treatments in *hoxba;hoxbb* cluster-deleted mutants do not rescue *tbx5a* expression in fin buds. (I, J) Expression patterns of *raldh2* were compared between sibling and *hoxba;hoxbb* mutants. For each genotype, reproducibility was confirmed with at least three different specimens. All images were captured from the dorsal side. Scale bars represent 100 μm.

Figure 5. Expression patterns of *hox5a* and *hoxb5b* are regulated by RA.
(A) Schematic representation of the 49 *hox* genes organized into seven *hox* clusters in zebrafish. *hox* genes in *hoxba* and *hoxbb* clusters are highlighted with orange. (B) Expression profiles of the 49 *hox* genes in wild-type and *raldh2*[-/-] embryos at the 20-somite stage, analyzed by RNA-seq. The average FPKM of each *hox* gene was compared between sibling and *raldh2* mutants. The absence of specific *hox* genes is indicated by a slash. (C-H) Expression patterns of *hoxb5a* and *hoxb5b* in sibling wild-type, *raldh2*[-/-], and RA-treated *raldh2*[-/-] embryos at the 10-somite stage. Arrowheads indicate the presumptive positions of pectoral fin buds. For each genotype, reproducibility was confirmed with at least three different specimens. All images are captured from the dorsal side. Scale bars represent 100 μm.

Figure 6

Figure 6. Screening for hox genes responsible for the zebrafish pectoral fin formation.

(A) Schematic representation of the genetic screening for *hox* genes involved in the specification of pectoral fins in zebrafish. (B-G) Dorsal views of live mutant larvae at 3 dpf, obtained during the screening. After capturing images, genotyping and DNA sequencing in target *hoxb* genes were conducted. For the mutants illustrated in (C-G), *hoxb* genes are shown with frameshift mutations introduced. Detailed information is provided in Supplementary Table S1. (H-J) Dorsal views of *hoxba⁻/⁻;hoxb5b⁻/⁻* and *hoxba⁻/⁻;hoxb5b⁻/⁻;hoxb6b⁻/⁻* larvae at 3 dpf. (K-M) Expression patterns of *tbx5a* in *hoxba⁻/⁻;hoxb5b⁻/⁻* and *hoxba⁻/⁻;hoxb5b⁻/⁻;hoxb6b⁻/⁻* mutants at 30 hpf. Arrowheads indicate the presumptive positions of pectoral fin buds. All images are captured from the dorsal side. Scale bars represent 100 μm.

Figure 7

Figure 7. *hoxb4a-b5a*^{del/del}*;hoxb5b*^{del/del} larvae partially recapitulate the absence of the pectoral fins.

(A-C) Dorsal views of zebrafish *hoxba*^{-/-}*;hoxb5b*^{del/del} larvae at 3 dpf. (D, E) Expression patterns of *tbx5a* in sibling wild-type and *hoxba*^{-/-}*;hoxb5b*^{del/del} (*n*=4) at 30 hpf. Arrowheads indicate the presumptive positions of pectoral fin buds. (F, G) Dorsal view of frameshift-induced *hoxb5a*^{-/-}*;hoxb5b*^{-/-} and *hoxb5a*^{del/del}*;hoxb5b*^{del/del} larvae. (I-K) Dorsal view of frameshift-induced *hoxb4a*^{-/-}*;hoxb5a*^{-/-}*;hoxb5b*^{-/-} and *hoxb4a-b5a*^{del/del}*;hoxb5b*^{del/del} larvae. All images are captured from the dorsal side. Arrowheads indicate the presumptive positions of pectoral fin buds. (M-R) Expression patterns of zebrafish *hoxb4a, hoxb6a, hoxb6b, hoxb7a, hoxb8a,* and *hoxb8b* at the 10-somite stage. Dorsal views. Arrowheads indicate the presumptive positions of pectoral fin buds. Scale bars represent 100 μm.