Nutritionally responsive PMv DAT neurons are dynamically regulated during

pubertal transition

Cristina Sáenz de Miera[a], Nicole Bellefontaine[a], Marina A Silveira[c,d], Chelsea N

Fortin[b], Thais T Zampieri[d], Jose Donato Jr[e], Kevin W Williams[f], Cristiano Mendes-da-

Silva[g], Laura Heikkinen[h], Christian Broberger[h], Renata Frazao[d], Carol F Elias[a,b*]

[a]Department of Molecular and Integrative Physiology, [b]Department of Obstetrics and

Gynecology University of Michigan, Ann Arbor, MI, 48109. [c]Department of

Neuroscience, Developmental and Regenerative Biology, The University of Texas at San

Antonio, San Antonio, TX, 78249. [d]Department of Anatomy and [e]Department of

Physiology and Biophysics, Institute of Biomedical Sciences, University of Sao Paulo,

Sao Paulo, SP, Brazil, 05508. [f]Center for Hypothalamic Research, Department of

Internal Medicine, Peter O'Donnell Jr. Brain Institute, The University of Texas

Southwestern Medical Center at Dallas, Dallas, TX, 75390. [g]Department of

Biosciences, Federal University of Sao Paulo, Santos, SP, Brazil, 11015. [h]Department

of Biochemistry and Biophysics, Stockholm University, Stockholm, Sweden.

*Corresponding Author: Carol F. Elias, Ph.D., North Campus Research Complex B25-

3682, 2800 Plymouth Road, Ann Arbor, MI 48109.

cfelias@umich.edu

ORCID iDs:

Cristina Sáenz de Miera: 0000-0001-8047-035X

Marina Augusto Silveira: 0000-0002-8351-5549

Thais T Zampieri: 0000-0002-1611-354X

Jose Donato Jr: 0000-0002-4166-7608

Kevin W Williams: 0000-0002-8434-8658

Cristiano Mendes-da-Silva: 0000-0001-7027-0127

Christian Broberger: 0000-0002-7050-8809

Renata Frazao: 0000-0002-0877-8453

Carol F Elias: 0000-0001-9878-9203

Key words: dopamine transporter, leptin, puberty, Kiss1, nutrition, hypothalamus

Graphical abstract

The ventral premammillary nucleus of the hypothalamus plays an essential role in the
metabolic control of reproduction. Puberty brings large changes to a subpopulation of
PMv^{LepRb} cells expressing the dopamine transporter (PMv^{DAT}). DAT gene expression is
higher in prepubertal than in adults and is regulated by leptin in prepubertal females.
Dynamic projections from PMv^{DAT} cells contact the kisspeptin and tyrosine hydroxylase
(TH) populations in the AVPV/PeN during puberty, a critical time for the appearance of
these cells in the AVPV/PeN.

Abstract

Pubertal development is tightly regulated by energy balance. The crosstalk between
metabolism and reproduction is orchestrated by complex neural networks and leptin
action in the hypothalamus plays a critical role. The ventral premammillary nucleus
(PMv) leptin receptor (LepRb) neurons act as an essential relay for leptin action on
reproduction. Here, we show that mouse PMv cells expressing the dopamine transporter
(DAT) gene, *Slc6a3* (PMv^{DAT}) form a novel subpopulation of LepRb neurons. Virtually
all PMv^{DAT} neurons expressed *Lepr* mRNA and responded to acute leptin treatment.
Electrophysiological recordings from DAT^{CRE};tdTomato mice showed that PMv^{DAT}
cells in prepubertal females have a hyperpolarized resting membrane potential
compared to diestrous females. *Slc6a3* mRNA expression in the PMv was higher in
prepubertal than in adult females. In prepubertal females *Slc6a3* mRNA expression was
higher in overnourished females from small size litters than in controls. Prepubertal
Lep^{ob} females showed decreased PMv *Slc6a3* mRNA expression, that recovered to

control levels after 3 days of leptin injections. Using a tracer adenoassociated virus in

the PMv of adult DAT^{Cre};$Kiss1^{hrGFP}$ females, we observed PMv^{DAT} projections in the

anteroventral periventricular and periventricular nucleus (AVPV/PeN), surrounding

$Kiss1^{hrGFP}$ neurons, a population critical for sexual maturation and positive estrogen

feedback in females. The DAT^{CRE};tdTomato projections to the AVPV were denser in

adult than in prepubertal females. In adults, they surrounded tyrosine hydroxylase

neurons. Overall, these findings suggest that the DAT expressing PMv^{LepRb}

subpopulation play a role in leptin regulation of sexual maturation via actions on AVPV

kisspeptin/tyrosine hydroxylase neurons.

Significance Statement

Women with excess or low energy stores (*e.g.,* obesity or anorexia) have reproductive

deficits, including altered puberty onset, disruption of reproductive cycles and decreased

fertility. If able to conceive, they show higher risks of miscarriages and preterm birth.

The hypothalamic circuitry controlling the interplay between metabolism and

reproduction is poorly defined. Neurons in the ventral premammillary nucleus express

the leptin receptor and play a key role in the metabolic control of reproduction. Those

neurons are functionally and phenotypically heterogeneous. Here we show that a subset

of leptin-sensitive neurons co-expresses the dopamine transporter (DAT), is

dynamically regulated during pubertal transition and with nutrition and projects to brain

sites relevant for sexual maturation.

Introduction

Pubertal development and the maintenance of reproductive function are disrupted in states of negative energy balance or excess energy reserve (1–3). If energy stores are low, puberty is delayed, the reproductive cycles are prolonged, and sub- or infertility ensues (2–4). High adiposity, on the other end, induces earlier pubertal development and decreased fertility in adult life (5–7). The cross-talk between metabolic and reproductive functions is orchestrated by a complex neuronal network modulated by circulating hormones and metabolic cues (8, 9). Among them, leptin has critical roles (10–12). Leptin signaling-deficient subjects develop obesity and remain in an infertile prepubertal state (13–15). In mice, direct leptin actions only in the brain are sufficient to normalize body weight, induce puberty and maintain fertility (10, 14, 16).

The ventral premammillary nucleus (PMv) contains a dense collection of leptin receptor (LepRb) neurons, and is recognized as an important hypothalamic site in the metabolic control of reproductive function (4, 16–21). Bilateral lesions of the PMv disrupt estrous cycles and the ability of leptin to increase luteinizing hormone secretion after fasting (4). Endogenous restoration of LepRb exclusively in PMv neurons rescues pubertal maturation and fertility in LepRb *null* female mice (16), while activation of PMv LepRb neurons is sufficient to induce LH release even in normally fed female mice (20). The PMv LepRb neurons, however, do not comprise a homogeneous population, *i.e.,* about 75% depolarize and 25% hyperpolarize in response to leptin (22), but their seemingly dissociated nature and function are poorly understood.

The PMv neurons are mostly glutamatergic and innervate brain sites associated with reproductive control, sending direct inputs to kisspeptin and gonadotropin-releasing hormone (GnRH) neurons (16, 18, 19, 21, 23). A subset of PMv neurons also expresses the dopamine transporter (DAT), a membrane protein associated with dopamine reuptake at presynaptic terminals (24–26). PMv^{DAT} neurons are unique in the sense that they show seemingly undetectable levels of tyrosine hydroxylase (TH) and dopamine release to specific brain sites (25, 27). Manipulation of PMv^{DAT} neuronal activity has shown an action in male social behavior, inter-male and maternal aggression and maternal behaviors (25, 26, 28, 29). However, the role of PMv^{DAT} neurons in female

reproductive physiology has not been described, and whether they participate in the
metabolic control of reproductive function is unknown.

In this study, we show that DAT is expressed in a subpopulation of PMv[LepRb] neurons.
Slc6a3 (*DAT*) mRNA expression is higher in prepubertal than in adult females and it is
increased in overnourished prepubertal females. We also show that PMv[DAT] neurons
project to and make apparent contacts with kisspeptin neurons in the the anteroventral
periventricular nucleus (AVPV) in adults, but not in prepubertal mice.

Methods

Experimental animals. All procedures were carried out in accordance with the National
Research Council Guide for the Care and Use of Laboratory Animals, and protocols
were approved by the University of Michigan IACUC (PRO000010420); and in
accordance with the European Community Council directive of November 24, 1986
(86/609/EEC) and had received approval by the local ethical board, *Stockholms*
Djurförsöksetiska Nämnd. Mice were held under a 12h:12h light:dark cycle (lights on at
6 am), temperature-controlled at 21-23 °C, and fed *ad libitum* on a low-phytoestrogen
diet (Envigo 2016 diet) and a higher protein and fat phytoestrogen reduced diet 2019
(Envigo 2019 Teklad diet) when breeding. Strains of mice used were a line expressing
Cre-recombinase under the *Slc6a3* promoter (DAT[Cre], JAX®; Stock 006660) (30), and
(31) only for female electrophysiology, a ROSA26 stop-floxed tdTomato reporter
mouse line (tdTomato, JAX®; Stock 007914), mice expressing GFP under the kiss1
gene promoter: Kiss1[hrGFP] (JAX®, stock 023425) (32), wild type C57B6/J (JAX®;
Stock 000664) and the B6.Cg-Lepob/J strain, homozygous mice with an obese
spontaneous mutation (*Lep[ob]*, JAX®; Stock 000632). Adult animals used were postnatal
(P) age 60-100 days old, unless otherwise specified.

Ovariectomy and estradiol replacement. To assess the effects of estradiol (E2) on
Slc6a3 gene expression we used ovariectomized (OVX, n=5), OVX + E2 (n=5) and
diestrous females (n=4). Females were deeply anesthetized with isoflurane and
underwent bilateral OVX. OVX females received steroid replacement via a Silastic
capsule containing E2 (1 μg, OVX+E2) or oil (OVX) subcutaneously at the time of
surgery. OVX females were perfused 7-14 days following surgery, while OVX+E2
females were perfused two days following E2 replacement. Uterus size was used as

control for the treatment. Only OVX mice with uterine weight below 80 mg and
OVX+E2 mice with uterine weight above 100mg were used. Both groups were perfused
in the morning to avoid time-of-day effects of estradiol feedback.

Leptin treatment. DAT^{Cre};tdTomato adult (P60-70) males and females fasted overnight
and prepubertal (P19) males and females fasted for 4h were intraperitoneally (i.p.)
injected with leptin (2.5 mg/kg, National Hormone and Peptide Program, Harbor-UCLA
Medical Center, CA) or saline (n=5-6 animals/group). Sixty minutes following leptin
injection, mice were perfused with PBS and 10% neutral buffered formalin (NBF,
Sigma), brains were postfixed for 2h in 20% sucrose in 10% NBF and stored with 20%
sucrose in PBS. 30 µm coronal tissue sections 120 µm apart were processed for
pSTAT3 immunohistochemistry as described below.

Two cohorts of adult wild type females in diestrus i.p. injected with saline, Lep^{ob}
animals i.p. injected with saline (Lep^{ob} + saline) or with leptin (3 mg/kg/day, Lep^{ob} +
leptin group, murine leptin, Preprotech), received the treatment for two days at 9 am and
5 pm and one day at 9 am. One hour after the last saline or leptin injection (at 10 am),
females were euthanized by decapitation following anesthesia (isoflurane) and brains
were harvested and snap frozen. Coronal frozen sections (16 µm) were collected on a
cryostat and stored at -80°C until processing for gene expression.

In situ hybridization (ISH) with radioisotopes. Adult wild type (WT) male (n=3) and
female (n=4) mice were used to study sex differences in *Slc6a3* gene expression.
Female mice were also used to determine developmental differences in *Slc6a3* gene
expression, *i.e.,* prepubertal (P19, n=7) *vs.* adult (P60-70, n=5) diestrous mice. To assess
the effects of nutritional factors in development on *Slc6a3* gene expression we used P20
females from small litters (SL 2-3 pups/litter, n=5 females) or normal litter (NL 7-9
pups/litter, n=4 females). To assess the effect of leptin in *Slc6a3* expression we used
diestrous (n=7), Lep^{ob} + saline (n=5) or Lep^{ob} + leptin (n=5) injected females. Coronal
sections were used for radioactive ISH, using an ^{35}S-UTP or ^{33}P-UTP labelled *Slc6a3*
riboprobe. The following primers were used (exon 10-15 of the *Slc6a3* gene): Forward
(5' ACGTCTTGATCACTGGGCTTGTCGATGAGTT 3') and reverse (5'
GCATGGATTGGGTGTGAACAGTC 3') to amplify a 754 base-pair sequence in the
Slc6a3 gene (exons 10-15). A clamp sequence followed by sequences for T7

(CCAAGCCTTCTAATACGACTCACTATAGGGAGA)   and   T3

(CAGAGATGCAATTAACCCTCACTAAAGGGAGA) promoters were added to the

reverse and forward primer sequences, respectively.

Single-labeled ISH was performed on 20 µm fresh frozen or 30 µm fixed brain (120 µm

distance) sections mounted onto SuperFrost Excell or Superfrost Gold slides (Fisher

Scientific). Fixed sections were subjected to a 10-minute microwave sodium citrate (pH

6) pre-treatment and hybridized overnight at 57 °C with ^{35}S-labeled *Slc6a3* riboprobes,

as previously described (17, 33). Frozen sections were fixed in ice-cold 10% NBF,

treated with 0.25% acetic anhydride and underwent dehydration in ethanol, and

hybridized overnight at 57 °C with ^{33}P-labeled *Slc6a3* riboprobes. All slides were then

incubated in 0.002% RNase A followed by stringency washes in sodium chloride-

sodium citrate buffer (SSC). Slides were exposed to film autoradiography (Kodak), for

3-5 days. Slides were dipped in autoradiographic emulsion (Kodak), dried for 3 hours

and stored in light-protected boxes at 4 °C for 2-4 weeks. Slides were developed in D-

19 developer, dehydrated in ethanol, cleared in xylene, and coverslipped with DPX

(Electron Microscopy Sciences). Film images were acquired using a stereoscope

(Zeiss). Darkfield 10x images were captured using a digital camera on an AxioImager

M2 microscope (Zeiss). ISH signals were quantified using integrated optical density

(IOD) in ImageJ software (NIH) using the "freehand" tool to outline the PMv. IOD

from the tissue background of the same area was subtracted.

Fluorescent ISH. We used diestrous WT female mice (n=3) to assess *Slc6a3* and *Lepr*

mRNA co-expression by fluorescent ISH. ISH was performed on fresh frozen 16-µm

thick cryostat sections at 128-µm resolution (8-series). The ISH was performed

following the RNAscope protocol for fresh frozen sections, using Protease III (ACDbio,

RNAscope Multiplex Fluorescent Reagent Kit v2). Briefly, slides were dried at 60 °C

for 15 min, rinsed in PBS for 5 min, fixed in 10% NBF for 15 min at 4 °C, rinsed in

PBS-DEPC 2 times for 3 min, dehydrated through rinses in serial ethanols for 3 min

each, and air-dried for 20 min. A hydrophobic barrier was created around each slide

using the ImmEdge pen (Vector Laboratories). The slides were then incubated in H_2O_2

for 10 min at RT followed by incubation with Protease III for 30 min at 40 °C. ISH was

performed using the RNAscope Protease III (ACDBio). Sections were incubated with

Mm-Slc6a3-C1 (#315441), and Mm-Lepr-C3 (#402731-C3, labeling all *Lepr* isoforms)

RNAscope probes for 2 h at 40°C using the HybEZ Humidifying System (ACDBio).
After all incubation steps following the kit's protocol, slides were incubated in DAPI
solution for 30 s at room temperature, and coverslipped using ProLong Gold Antifade
Mountant (ThermoFisher Scientific).

Quantification of mRNA coexpression within cells was performed on PMv images
acquired with a 40x oil objective on an AxioImager M2 microscope (Zeiss). Based on
observed background outside of the area of interest, a threshold for a minimum number
of 5 puncta per cell was used to consider a cell positive for expression of that gene.
Confocal images were acquired for illustration on a Nikon A1 confocal microscope.

Electrophysiological recordings. Hypothalamic slices from adult DAT^{Cre};tdTomato
male (30) and female (31) mice were prepared and the data analyzed as previously
described (22). Briefly, mice were decapitated following isoflurane anesthesia, and the
entire brain was removed. After removal, the brains were immediately submerged in
ice-cold, carbogen-saturated (95% O_2 and 5% CO_2) artificial cerebrospinal fluid (ACSF,
126 mM NaCl, 2.8 mM KCl, 26 mM NaHCO$_3$, 1.25 mM NaH$_2$PO$_4$, 1.2 mM MgSO$_4$, 5
mM glucose and 2.5 mM CaCl$_2$). Coronal sections (250 µM) from hypothalamic blocks
were cut on a Leica VT1000S vibratome and incubated in oxygenated ACSF at room
temperature for at least 1 hour before the recordings. The slices were transferred to the
recording chamber and allowed to equilibrate for 10–20 min. The slices were bathed in
oxygenated ACSF (32°C) at a flow rate of ~2 mL/min. The pipette solution was in some
cases modified to include an intracellular dye (Alexa Fluor 488) for whole-cell
recording: 120 mM K-gluconate, 10 mM KCl, 10 mM HEPES, 5 mM EGTA, 1 mM
CaCl$_2$, 1 mM MgCl$_2$, 2 mM (Mg)-ATP, and 0.03 mM AlexaFluor 488 hydrazide dye,
pH 7.3. Whole-cell patch-clamp recordings were performed on tdTomato-positive
neurons anatomically restricted to the PMv. Epifluorescence was briefly used to target
the fluorescent cells; at which time the light source was switched to infrared differential
interference contrast imaging to obtain the whole-cell recording (Leica DM6000 FS
equipped with a fixed stage and a fluorescence digital camera). In current-clamp mode,
tdTomato neurons were recorded under zero current injection (I = 0) in whole-cell
patch-clamp configuration. The recording electrodes had resistances of 5-7 MΩ when
filled with the K-gluconate internal solution. The membrane potential values were
compensated to account for the junction potential (-8 mV). In males and females at both

ages, the resting membrane potential (RMP) was monitored for at least 10-20 minutes
(baseline period) before leptin was administered to the bath. Solutions containing leptin
(100 nM) were typically perfused for 15-20 minutes after the baseline period, with a 20-
minute washout with ACSF.

Manipulation of estrous cycles using chemogenetics. Adult virgin DAT-Cre females
(n=14) received bilateral injection (50 nL/side) of an adenoassociated virus (AAV)
expressing a Cre-dependent hM3Dq-mCherry fusion protein,pAAV8-hSyn-DIO-
hM3D(Gq)-mCherry (AAV-hM3Dq, Addgene plasmid # 44361, from Bryan Roth) (34)
in the PMv using the following coordinates: Anteroposterior = -5.4 mm (from rostral
rhinal vein); mediolateral = -0.52 mm (from sagittal sinus); dorsoventral = -5.4 mm
(from dura mater). The stereotaxic protocol is described in detail in a previous study
(35). One month after surgery, we started to follow the reproductive cycle of these
females collecting daily vaginal smears with saline solution. After a period of
adaptation, we continued for 13 days with added DMSO (0.0068%) in water, as a
control. Next, we added CNO (5 mg/kg) dissolved in DMSO in the drinking water and
followed the cycles for 13 more days. The abundant presence of cornified cells in the
smear was considered as estrus/metestrus, a large abundance of leukocytes was
considered as diestrus and a large abundance of nucleated cells with some cornified
cells present was considered as proestrus. At the end of the experiment, the females
received i.p. injection of CNO and 2 h after mice were perfused with PBS and 10%
NBF (Sigma), brains were postfixed for 4 h in 20% sucrose in 10% NBF and stored
with 20% sucrose in PBS. We collected 30 µm coronal tissue sections 120 µm apart.
Sections were cryoprotected and frozen until processed to verify the injection sites for
these females.

Tracing PMv-DAT neuronal projections. DAT-Cre adult females (n=7) received
unilateral stereotaxic injections of an AAV expressing a Cre-dependent
channelrhodopsin-mCherry fusion protein (AAV8-hSyn-double floxed-hChR2(H134R)-
mCherry, UNC Vector Core, from Karl Deisseroth (Addgene 20297), 25-50 nL) in the
PMv. One month after the stereotaxic surgery, mice were perfused with 10% NBF and
brains were harvested and processed for histology as above. Fixed coronal 30 µm
hypothalamic brain sections were processed for immunofluorescence.

Immunohistochemistry. Fixed frozen tissue sections (30 µm at 120 µm distance) from

perfused animals obtained with a freezing microtome (Leica) were rinsed in PBS and

blocked with PBS + Triton-X 0.25% (PBT) and 3% normal donkey serum (NDS).

Primary antibodies were incubated in PBT + 3% NDS overnight at room temperature.

Primary antibodies used were Rabbit dsRed antibody (1:5000, Clontech 632496,

RRID:AB_10013483), rat monoclonal anti-mCherry 16D7 (1:5000, Invitrogen M11217,

RRID:AB_2536611), rabbit anti-cFOS (1:5000, Millipore ABE457,

RRID:AB_2631318), chicken anti-GFP (1:10000, Aves GFP-1010,

RRID:AB_2307313), rabbit polyclonal anti-GnRH (1:5,000, Phoenix Pharmaceuticals

H-003-57, RRID:AB_572248), and sheep anti-TH (1:5000, Millipore AB1542,

RRID:AB_90755). For detection of Fos, endogenous peroxidase was blocked with 0.3%

H_2O_2 for 30 min before the NDS blocking step. For detection of phosphorylation of

signal transducer and activator of transcription 3 (pSTAT3), tissue was pre-treated with

1% H_2O_2 and with 1% sodium hydroxide in water and then with 0.3% Glycine before

blocking with PBT + NDS 3% as before. Tissue was incubated in primary rabbit anti-

pSTAT3 (Tyr705) (D3A7) XP® (1:1,000; Cell Signaling 9145S, RRID:AB_2491009)

for 48 h at 4°C. The corresponding secondary fluorescent antibodies were used for

detection (1:500, Invitrogen). For the Fos antibody we performed immunoperoxidase

detection using a biotinylated-anti rabbit IgG secondary antibody (1:1000, Jackson

Immunoresearch), signal amplification with Avidin Biotin Complex (Vectastain® ABC-

HRP Kit, 1:500, Vector labs) for 1 h and signal development with diaminobenzidine

(DAB, Sigma) 0.05% and 0.01% H_2O_2. Floating sections were then mounted on gelatin-

coated slides, dried overnight and coverslipped with Fluoromount-G (Invitrogen).

Photomicrographs were acquired using Axio Imager M2 (Carl Zeiss Microscopy).

Quantification of pSTAT3, tdTomato and TH positive neurons was performed by an

observer unaware of the images' identity. Dual-labeled tdTomato and pSTAT3

immunoreactive cells were counted in each individual channel and colocalization was

considered where pSTAT3 immunoreactivity (-ir) was clearly nuclear in tdTomato-

positive cells. Two sections at the mid-PMv level were counted (~Bregma: -2.46 mm).

No correction for double counting was performed because sections were 120µm apart.

tdTomato fiber density in the AVPV/PeN was quantified using IOD in ImageJ software

(NIH) on both sides of the ventricle in a representative section for each region. An

elongated rectangle of the same size for all animals was used as region of interest,

placed in contact with the ventricle wall to cover a representative area over the TH-

expressing cells.

Confocal microscopy images were acquired and analyzed using a Nikon A1 microscope

and a Nikon N-SIM + A1R microscope with a resonance scanner.

Data analysis. Data are expressed and represented as mean ± SEM. When data did not

fit a normal distribution or did not have equal variances, they were transformed to fit a

normal distribution and re-analyzed. Unpaired two-tailed Student's t test was used for

comparison between two groups. For comparison between three groups, one-way

ANOVA was used followed by Tukey's *post-hoc* multiple comparison test. For

pSTAT3 and %pSTAT3/tdTomato cells, a two-way ANOVA was used with age and sex

as factors. Correlation was assessed between body weight and *Slc6a3* gene expression

using Pearson R correlation coefficient. A P value less than 0.05 was considered

significant. Data were organized and calculated in Excel software (Microsoft, inc.).

Statistical analyses and graphs were performed using GraphPad Prism v.9.5 (GraphPad

software, inc.). Zen Blue 3.7 software (Carl Zeiss Microscopy GmBH) was used to

acquire and process epifluorescence images. NIS-elements software (Nikon) was used

to acquire and process confocal microscope images. Photoshop 2024 (Adobe, inc.) was

used to integrate graphs and digital images into figures. The graphical abstract was

prepared using BioRender. Only brightness, contrast, and levels were modified to

improve data visualization in the figures.

Results

***Slc6a3* mRNA expression in the PMv is sexually dimorphic and higher in**

prepubertal females

The *Slc6a3* (DAT) gene is expressed in the PMv of male and female mice (24, 25). To

evaluate potential sexual dimorphism or postnatal developmental changes, we assessed

Slc6a3 gene expression in the PMv in adult males and females, and in prepubertal and

adult females. PMv *Slc6a3* mRNA levels were higher in diestrous females compared to

male mice (n=3 female, n=4 male, unpaired t-test p=0.032 Figure 1A, B, D), and higher

in prepubertal females compared to diestrous mice (n=7 prepubertal, n=5 diestrous,

unpaired t-test p=0.013 Figure 1B, C, E).

To assess if the reduction of PMv *Slc6a3* mRNA in adult female mice is a result of

increasing circulating estradiol (E2) during the pubertal transition, hypothalamic

sections from diestrous, ovariectomized (OVX), and OVX + E2 mice were analyzed.

We found no differences between these groups (n=5 diestrous and OVX+E2, n=4 OVX,

one-way ANOVA, $p = 0.54$, Figure 1F).

PMV^{DAT} neurons show heterogenous responses to leptin

We used fluorescent *in situ* hybridization (ISH) to assess transcript coexpression in

adult females in diestrus (n=3). Virtually all *Slc6a3* neurons in the PMv coexpressed

Lepr mRNA (93.6 % ± 2.1), whereas about half of PMv *Lepr* neurons coexpressed

Slc6a3 mRNA (58.6 % ± 4.1, Figure 2A-C).

To investigate the effect of leptin on the membrane excitability of PMv^{DAT} neurons, we

performed current clamp recordings of DAT^{Cre};tdTomato neurons. In males, the average

RMP of recorded neurons was -52.8 ± 2.0 mV (range from -62 to -38 mV, 16 cells from

8 mice). In a separate cohort of females, the RMP was -57.2 ± 5.7 mV (range from -62

to -51 mV, 8 cells from 6 mice). We found that bath application of 100 nM leptin

hyperpolarized 25% of the recorded neurons of male mice (4/16 cells Figure 2D-G).

The RMP of the remaining 75% of the recorded cells was unchanged or showed a

continuous depolarizing trend and were removed from the analysis. In females in

diestrus, ~75% (6/8) of recorded cells hyperpolarized in response to bath application of

100nM leptin (Figure 2G). Two PMv^{DAT} neurons showed a depolarizing response, and

one showed continuous depolarization and was removed from the analysis (Figure 2E).

The leptin-associated hyperpolarization of PMv^{DAT} cells was of similar amplitude in

both sexes: -7.8 ± 0.8 mV in males, and -6.3 ± 2.0 mV in females.

Although most (> 90%) of *Slc6a3* expressing neurons express *Lepr*, only a

subpopulation exhibits a response, including both de- and hyperpolarization, in

electrical properties to the hormone.

Long-term activation of PMv^{DAT} neurons does not alter estrous cycles in adult

virgin females

Given the higher percentage of female's DAT^{Cre};tdTomato neurons that are

hyperpolarized by leptin, we decided to investigate if long-term activation of these

neurons might impair the reproductive cycle of the adult females. We stereotaxically

injected the AAV-hM3Dq virus bilaterally in the PMv of 14 DAT^{Cre};tdTomato virgin

females. Twelve females had bilateral injections and one had a unilateral injection

centered in the PMv, defined by the expression of mCherry and Fos immunoreactivity

that indicates they had been activated by CNO (Figure 3A-B). Two animals had missed

injections with not Cherry expression observed in either PMv, and lack of Fos

confirmed in the PMv of these animals (Figure 3C). Of the twelve animals with bilateral

injection, eleven showed regular cycling (at least two complete cycles) during the

control (DMSO) period and were used for the analyses (Figure 3D). The cycles of the

bilaterally injected females were not altered by the CNO when compared to the DMSO

exposure (paired t-test DMSO *vs*. CNO, days in estrus/metestrus, p=0.59; days in

diestrus, p=0.68; cycle length, p=0.69, n=11, Figure 3E-G). We paid special attention to

the potential virus spread to a nearby population of *Slc6a3* expressing cells, the tubero-

infundibular dopamine (TIDA) neurons in the arcuate nucleus (Arc), (24). Six mice

showed some viral contamination of TIDA neurons, but no differences in cyclicity or

cycle length were noticed when these animals were removed from the analysis. These

results suggest that these cells have no effect on female cyclicity.

372 Prepubertal PMv^{DAT} neurons respond to leptin and show distinct membrane

373 properties compared to adult females.

Due to increased expression of *Slc6a3* in prepubertal females, we explored the

functional response of PMv^{DAT} neurons to exogenous leptin. Adult and prepubertal

DAT^{Cre};tdTomato mice received an i.p. injection of leptin, and one hour after,

colocalization of pSTAT3-ir in tdTomato neurons was quantified. No differences were

observed in the number of pSTAT3-ir cells with age or sex in leptin treated mice (n = 3-

5; Two-way ANOVA, p=0.28 for Sex; p=0.79 for Age; Figure 4A-G). Virtually no

pSTAT3-ir was observed in the PMv of saline treated mice (n=5 per group). As

expected from the *Slc6a3* and *Lepr* coexpression data, 95.3 – 99.3 % of tdTomato

neurons in the PMv colocalized with pSTAT3-ir in adults of both sexes (Figure 4H).

Similar colocalization was observed in prepubertal mice of both sexes (96.4 – 98.9 %,

Figure 4H). About 30% of PMv pSTAT3-ir neurons colocalized with tdTomato in

females (25.9 ± 2.8% prepubertal, and 33 ± 2.4% in diestrous, p=0.14) and males (27.8

± 1.8% prepubertal, and 35.6 ± 4.8% in adults, p=0.18).

When examined by electrophysiology, the PMv^{DAT} neurons of prepubertal (unweaned)

female mice revealed heterogeneous properties. Interestingly, the RMP of prepubertal

PMv^{DAT} neurons was more hyperpolarized compared to adult diestrous females (n=8

prepubertal and n=8 diestrous, unpaired t-test, p <0.0001, Figure 4I). In response to

leptin treatment, three out of eight cells (37.5%) from three mice showed no RMP

change and another three out of eight cells (37.5%) responded by hyperpolarization

(Figure 4J, K). Two out of eight cells (25%) depolarized after acute leptin, but none of

them recovered after washout. Two recorded cells showed continuous depolarization

and were removed from the analysis. The hyperpolarized cells showed a -6.3 ± 2.1 mV

change in the RMP after treatment.

Our findings indicate that the PMv^{DAT} neuron population from prepubertal (unweaned)

female mice is in a less excitable state compared to adult mice.

Postnatal overnutrition increases *Slc6a3* mRNA expression in the PMv of

prepubertal females

We next assessed if the expression of *Slc6a3* mRNA is altered in the PMv of leptin

deficient *Lep^{ob}* infertile female mice, which remain in a prepubertal state. We employed

a paradigm of leptin treatment and pubertal progression in which *Lep^{ob}* females were

injected with saline or leptin twice a day for 2 ½ days are compared to age-matched

wild type diestrous females (16, 36). As expected, the leptin-treated *Lep^{ob}* mice showed

a significant decrease in body weight and displayed signs of pubertal progression

(vaginal opening) following the leptin treatment. We found that *Slc6a3* mRNA

expression is ~40% lower in non-treated *Lep^{ob}* females, compared to wild type females

in diestrus (n=5-7; p=0.0025 one-way ANOVA, Tukey's *post-hoc*, p=0.005 Figure 5A).

The short-duration leptin treatment regimen was sufficient to induce a ~ 40% increase in

Slc6a3 mRNA expression in the PMv of *Lep^{ob}* mice (Tukey's *post-hoc*, p=0.005, Figure

5A), concomitant with the first signs of puberty onset.

Given the complex phenotype of the *Lep^{ob}* mouse (37), we employed a paradigm of

postnatal overnutrition, which leads to high leptin levels and early puberty (38, 39). We

compared *Slc6a3* mRNA expression in the PMv of females raised in normal (NL, n=4

females) versus small (SL, n=5 females) litter sizes. Body weight was higher in SL

offspring as compared to NL (9.67 ± 0.59 g in SL *vs*. 5.65 ± 0.39 g in NL, unpaired t-

test p=0.001). *Slc6a3* mRNA levels were higher in the SL than in those in NL (p=0.007,

Figure 5B-D) and were strongly correlated to body weight (Pearson r=0.78; p=0.01,

Figure 5E).

PMv^{DAT} neurons project to kisspeptin AVPV/PeN neurons

To assess if PMv^{DAT} neurons are part of the circuitry regulating pubertal development,

DAT^{Cre};*Kiss1^{hrGFP}* females were unilaterally injected with a Cre-dependent AAV

expressing a channelrhodopsin-mCherry fusion protein (AAV-ChR2-mCherry) into the

PMv (n=7). Abundant mCherry-ir neurons were observed within the PMv in correctly

targeted animals (n=6). Mice showing virus spread to nearby DAT expressing

populations were removed from the analysis (n=2 were analyzed, Figure 6A). In

accordance with previous studies focused on PMv projections (41, 42), dense mCherry-

ir fibers were found in several hypothalamic regions including the AVPV (Figure 6B),

the periventricular nucleus (PeN) the medial preoptic area (MPA, Figure 6C), and the

ventrolateral subdivision of the ventromedial hypothalamus (VMHvl, not shown). Most

notably, very sparse innervation of the Arc was observed (Figure 6D). No mCherry-ir

projections were observed nearby or in contact with GnRH cell bodies in the medial

septum (MS) or MPA (not shown).

To explore a possible interaction of PMv^{DAT} neurons with kisspeptin, we analyzed

mCherry-ir fibers in proximity to *Kiss1^{hrGFP}* cells using confocal microscopy. As

expected, due to the low innervation of the Arc, kisspeptin/neurokinin 3/dynorphin

(KNDy) neurons did not receive close appositions from PMv^{DAT} neurons (Figure 6E). In

contrast, dense mCherry innervation of kisspeptin cells was observed in the AVPV and

PeN region (AVPV/PeN, a.k.a. rostral periventricular area of the third ventricle, Figure

6F) of the adult female mouse.

PMv^{DAT} innervation of AVPV/PeN is established during the pubertal transition

The AVPV/PeN area contains a sexually dimorphic population of dopaminergic TH and

kisspeptin cells, both denser in females. About 50-90% of the *Kiss1* cells express TH in

mice (40–43), and kisspeptin expression in this region increases during the pubertal

transition (40, 44). In DAT^{Cre};tdTomato mice, we found that AVPV/PeN TH neurons do

not express tdTomato. DAT^{Cre};tdTomato fiber density was about three times higher in

the AVPV (p=0.006, Figure 7A, B and E), and about ten times higher in the PeN of

adult *vs*. prepubertal females (p=0.003, Figure 7C, D and E). As observed for *Kiss1*, the

number of neurons expressing TH was higher in both the AVPV and the PeN of adult

diestrous female mice compared to prepubertal females (p<0.0001 in the AVPV;

p<0.0001 in the PeN, Figure 7A-D, F). DAT^{Cre};tdTomato terminals were in close

apposition to TH neurons and fibers in the AVPV/PeN of adult female mice (Figure 7G-

H).

Discussion

The present study revealed a novel subset of leptin responsive cells within the PMv that

show dynamic regulation of the *Slc6a3* (DAT) mRNA during puberty and specific

projections to hypothalamic sites and neurons involved in puberty and reproductive

control. The overall PMv^{Leprb} population is key in the metabolic regulation of puberty

and fertility, whereas the PMv^{DAT} neurons were previously shown to be involved in

aggression, social and maternal behaviour (25, 26, 28, 29). Our findings indicate that the

PMv^{DAT} population represents a discrete subset of PMv^{LepRb} neurons and a novel

candidate for mediating nutritional modulation of reproduction and pubertal

development.

Within the PMv, leptin-induced pSTAT3 was observed in all DAT-expressing cells.

However, acute leptin exposure had heterogenous sex- and age-dependent effects on the

excitability of DAT^{Cre} neurons. The lack of electrophysiological response of a

subpopulation suggests that these neurons are responsive to leptin in ways not tested in

this study, *e.g.* by means of transcriptional regulation. Acute leptin action in the

PMv^{Leprb} population is also heterogenous, inducing the depolarization of 75% of these

cells, via a putative transient receptor potential channel (TRPC) channel, and

hyperpolarizing 25% cells, via activation of a putative Katp channel, while no cells

were found to be unresponsive to the treatment (22). Importantly, the RMP of

prepubertal females was more hyperpolarized than that of adult females suggesting that

these cells may increase in excitability with maturation and only assume a more active

role within the circuit after puberty. PMv^{DAT} cells are responsive to prolactin and

oxytocin (29, 45) and the influence of these hormones may contribute to switching these

cells from a quiescent to an excitable state in their role in maternal aggression (29).

Further studies are needed to determine the acute and chronic effects of leptin on the

electrical properties of PMv^{DAT} neurons in distinct developmental and physiological

states, as well as the mechanisms associated with changes in intrinsic physiological

properties of PMv^{DAT} neurons from prepubertal to adults.

PMv^{DAT} cells have been described as a non-dopaminergic population (25, 27). These

cells are an active glutamatergic population and send excitatory inputs to projection

sites, in particular the VMHvl (25, 26). Several studies, including ours (not shown) have

observed a lack of TH expression in this population (27, 46). Only one study has shown

an enrichment in TH using Ribotag mice, but to a much lower extent than classical

midbrain dopaminergic cells (25). Still, PMv^{DAT} neurons express other elements of the

dopamine/monoamine regulation pathway, such as *Gucy2c*, *Aadc* and *Vmat2* (25, 47).

In shrews, dopamine and serotonin have been detected in the PMv after L-DOPA and 5-

HTP treatment, respectively (47). However, in mice studies using fast-scan cyclic

voltammetry have suggested that PMv^{DAT} neurons do not produce dopamine, even when

supplemented with L-DOPA (25). Thus, a significant role for dopamine release from

these neurons is viewed as unlikely. Still, here we have discerned differences in the

regulation of the dopamine transporter that merit further attention.

Slc6a3 mRNA expression in the PMv was higher in females than in males, and showed

a developmental decrease after puberty, suggesting a regulation during sexual

maturation. Removal of the ovaries or changes in estrogen levels did not affect *Slc6a3*

mRNA expression, making estrogen an unlikely regulator during the pubertal shift.

Leptin is critical for puberty and fertility. Here, overnourished prepubertal animals from

SL showed increased level of *Slc6a3* mRNA expression in PMv, correlated with

individual's body mass, although other factors, such as altered sex distribution in the

mostly-females SL could also affect gene expression. Furthermore, the rescue of *Slc6a3*

mRNA levels observed in leptin-treated *Lep^{ob}* animals suggests that leptin has an active

role in increasing *Slc6a3* gene expression during pubertal transition. Whether other

hormones (*e.g.* prolactin, oxytocin), different physiological states or social behaviors

affect PMv *Slc6a3* expression is unknown.

Leptin regulation of dopaminergic neurotransmission in the midbrain (substantia nigra

and ventral tegmental area) is involved in motivation for food rewards and locomotion

(48–50). Leptin specifically regulates dopamine-related genes in these populations (51)

and their action in the nucleus accumbens (52). It is important to note though that most

studies investigating the role of PMv neurons have used DAT-Cre mice as a strategy to

assess the PMv's neuronal function and circuitry, not DAT expression and function.

More studies are needed to reveal the role of dopamine-related genes in the PMv, but

our findings suggest that *Slc6a3* and its regulation by leptin have a role in pubertal

maturation. DAT is mostly found in presynaptic terminals (53); so, this role might be

relevant in PMv projection sites, perhaps regulating the neurotransmission of PMv

neurons.

The PMv^{DAT} neurons project to a subset of brain sites innervated by the PMv^{Leprb}

population. The PMv^{Leprb} population innervates VMHvl, and key neuronal populations

in the control of reproduction, namely, GnRH neurons in the OVLT, MPA and medial

septum areas, KNDy neurons in the Arc, and kisspeptin cells in the AVPV/PeN (16, 18,

54). As reported before, PMv^{DAT} neurons project to the VMHvl and the

supramammillary nucleus (25, 26). In the MPA, we observed little density of axons.

However, we observed a very dense collection of terminals in the AVPV/PeN.

Differences with previous studies might arise from the use of a different protein marker

(ChR2 *vs*. synaptophysin). Of the reproductive populations targeted by PMv^{Leprb}

neurons, the PMv^{DAT} neurons seem to specifically target the kisspeptin population in the

AVPV/PeN. These results show that the PMv^{DAT} subpopulation target a group of

neurons essential for pubertal development in females, supporting a function in puberty.

In adults, AVPV/PeN kisspeptin neurons mediate the positive feedback action of

estradiol on LH surge that precedes ovulation on the afternoon of proestrus (55). We

recently showed that acute activation of the PMv^{Leprb} population in females leads to LH

release (20). However, chronic chemogenetic activation of PMv^{DAT} cells had no effect

on estrous cycles (present study). Alterations of the LH surge in proestrus have been

observed in the absence of effects on estrous cycle progression (56). Therefore, we

cannot discard any effects on the LH surge, which can be disrupted after PMv lesions in

rats (4, 57). The lack of direct PMv^{DAT} projections to Arc KNDy or GnRH neurons, the

dynamic changes in *Slc6a3* expression and neuronal properties, the dense projections to

the AVPV/PeN, and the AVPV/PeN's intense chemical remodelling during puberty

maturation, suggest that the PMv^{DAT} population could undergo a functional switch

during pubertal maturation. We hypothesize that the PMv^{DAT} population could play an

important role in the regulation of sexual maturation and later reproductive function,

integrating the nutritional state from leptin signal into the AVPV/PeN.

Similar to gene expression, the innervation of the AVPV/PeN from DAT^{Cre} neurons was

dynamically regulated during female puberty. In adult females DAT projections densely

englobed TH neurons, a trait absent in prepubertal females. This timing probably

reflects an increase in *Slc6a3*-driven *Cre* expression and a delay to observe tdTomato -

ir. The AVPV is a sexually dimorphic population, with higher cell abundance in the

female (58). Similar to kisspeptin neurons, the TH population in the AVPV/PeN is

sexually dimorphic, with more cells present in females (59, 60).

Notably, TH expression in the AVPV/PeN was much lower in prepubertal females than

in the adults, a similar developmental time as kisspeptin appearance (61) and the

increase of DAT terminals in the AVPV/PeN. This is significant because the

AVPV/PeN region is one of the classical dopaminergic regions of the hypothalamus,

also known as A15 area (62). The A15 TH neurons do not express DAT (46, 63, 64).

Prototypical dopaminergic neurons (*i.e.,* in the midbrain), co-express TH and DAT, so

that dopamine is recycled by DAT at the presynaptic terminal. The significance of our

current finding is puzzling. However, a recent report provided evidence that DAT plays

an integral role in managing the dopaminergic micro-circuitry within the ARH TIDA

neurons (65), suggesting that PMv^{DAT} could participate in the AVPV/PeN dopamine

microcircuitry, potentially via expression of Aromatic l-aminoacid decarboxylase

(AADC) (25, 47). We therefore speculate that the presence of DAT at the PMv

presynaptic terminal plays a role in regulating dopaminergic tone in the AVPV/PeN

microcircuits and female reproductive function. In addition, PMv^{DAT} neurons likely act

on AVPV cells via glutamatergic signaling. AVPV TH neurons play a role in maternal

behavior and intermale aggression (66). Whether the PMv^{DAT} neurons' action in the

same social behaviors is associated with AVPV/PeN TH neuronal innervation has not

been determined.

Our present findings demonstrate that the role of PMv neurons in regulating

reproductive physiology is more complex than previously anticipated. While overall

PMv^{Leprb} neurons play a significant role in female reproduction, particularly mediating

leptin's permissive effects in puberty (16), PMv[DAT] cells have been studied primarily in
the context of social behavior and aggression (25, 26, 28, 29). The dynamic changes
observed in *Slc6a3* gene expression during puberty and in response to nutrition, as well
as the developmental difference in the innervation of kisspeptin/TH neurons of
AVPV/PeN, suggest that the PMv[DAT] neurons also play a role in sexual maturation.

Conflict of Interest

579 All the authors declare no conflicts of interest.

Author contributions

581 **CSM, CFE** Conceived and designed the experiments. **NB, JDJr, KWW,** generated
582 preliminary data. **CSM, MAS, CNF, TTZ, CMS, LH,** performed the experiments and
583 acquired the data. **CSM, MAS, CB, RF, CFE** analyzed and interpreted the data. **CSM**
584 wrote the manuscript. All authors were involved in revising and approving the
585 manuscript

Acknowledgements

We thank Dr. Yun-Hee Choi for the design of the DAT riboprobe, and Susan Allen for
expert technical assistance. This work was supported by the National Institutes of
Health (R01-HD-069702 to CFE, CSM and NB, R21 HD109485 to CFE and CSM),
Michigan Nutrition and Obesity Research Center (URM Pilot Grant P30 DK089503 to
CSM) CNPq (Brazilian National Council for Scientific and Technological Development
fellowship to BCB), the Knut and Alice Wallenberg Foundation (2020.0054) and the
Swedish Research Council Distinguished Professorship Grant (2021-00671) to CB, the
Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) –
Finance Code 001 (MAS) and by the São Paulo Research Foundation [FAPESP-Brazil,
grants number: 13/07908-8 (RF), 15/20198-5 (TTZ) and the National Institutes of
Health (R01 DK119169, R56 DK135501, and PO1 DK119130-03 to KWW).

Figure 1. Ventral premammillary nucleus (PMv) _Slc6a3_ gene expression varies

with sex and development. A-C. Darkfield images showing the _Slc6a3_-[35]S

hybridization signal (silver grains) in the PMv of adult male, a diestrous female and a

prepubertal female, respectively. **D-F.** Graphs showing the quantification of the _Slc6a3_

hybridization signal in adult male _vs_. diestrous females, in prepubertal _vs_. diestrous

females and in diestrous _vs_. ovariectomized (OVX) females and OVX females

supplemented with estradiol (E2). All data shown are average ± SEM. * p<0.05. Scale

bar = 100 μm.

Figure 2. A subpopulation of leptin receptor (*Lepr*) expressing neurons in the

ventral premammillary nucleus (PMv) co-expresses dopamine transporter

(*Slc6a3*). A. Fluorescent image showing representative fluorescent *in situ* hybridization

depicting the colocalization of *Lepr* (magenta) and *Slc6a3* (green) in the PMv of a

diestrous female. Arrows point to cells co-expressing *Lepr* and *Slc6a3* mRNA.

Arrowheads point to cells expressing only *Lepr*, but not *Slc6a3*, mRNA. Blue = DAPI.

B-C. Higher magnification of individual cells depicted in A that co-express *Slc6a3* and

Lepr mRNA (B), and of a cell that expresses only *Lepr* mRNA (C). **D.** Fluorescent

image showing the PMv in the brain slices, recognized by tdTomato expression in

DAT^{Cre} neurons. **E.** Merged image showing the colocalization between a recorded

DAT^{Cre};tdTomato neuron (magenta) and the AF488 dye (green), dialyzed during the

recording. **F.** Representative current-clamp recording demonstrating leptin (100 nM)

induced hyperpolarization in a subset of DAT-Cre tdTomato neurons of a male mouse.

The dashed line indicates resting membrane potential (-52 mV). Asterisks indicate

square pulse current injections to assess input and access resistance. **G.** Pie charts

representing the percentage of neurons that hyperpolarized, depolarized or did not

respond to 100nM leptin in adult males (N=16) and in adult diestrous females (N=8).

Scale bars: A and E = 50 µm, B-C: 2 µm, D= 400 µm.

Figure 3. Continuous activation of dopamine-transporter neurons in the ventral

premammillary nucleus (PMv^{DAT}) does not alter estrous cycles in adult DAT^{Cre}

female mice. A. Representative low-magnification fluorescent image of bilateral

injections of adenoassociated virus (AAV) expressing Cre dependent hM3Dq-mCherry

targeted to the PMv of a female DAT^{Cre} mouse. **B.** High magnification image showing

Fos immunoreactivity (Fos-ir) in one PMv side following an intraperitoneal injection of

clozapine-N-oxide (CNO) and corresponding fluorescent image of the PMv showing

mCherry immunofluorescence. **C.** High magnification image showing the lack of Fos-ir

in one PMv side of a missed AAV injection, following an intraperitoneal injection of

clozapine-N-oxide (CNO) and corresponding fluorescent image of the PMv showing the

lack of mCherry immunofluorescence. **D.** Representative estrous cycles of two females

with AAV injections centered in the PMv before treatment (drinking water), during the

treatment with DMSO (vehicle) and CNO in drinking water. E/M: estrus/metestrus; P:

Proestrus; D: Diestrus. **E-G.** Graphs showing the number of days spent in

estrus/metestrus, diestrus and the cycle length (number of days) in the DAT^{Cre} females

with bilateral PMv AAV-hM3Dq injections during the DMSO and the CNO treatment.

Data are average ± SEM. Scale bars: A = 200 µm, B, C = 100 µm

Figure 4. Prepubertal dopamine-transporter neurons in the ventral premammillary nucleus (PMv^{DAT}) are responsive to leptin but are more hyperpolarized than in adult females.

A, B. Fluorescent images showing the colocalization of pSTAT3 (green) and tdTomato (magenta) immunoreactivities in the PMv of prepubertal and adult diestrous DAT^{Cre};tdTomato females 1h after a 2.5 mg/kg intraperitoneal leptin injection. **C-F**: High magnification images of the images depicted in A (**C**, D) and B (E, F). **G, H.**

Graphs showing the number of pSTAT3 neurons per section (G) and the percentage of

tdTomato cells expressing pSTAT3 (H) in the PMv of prepubertal and adult

DAT^{Cre};tdTomato female mice. Prepubertal (n=5) and diestrous (n= 3) females,

prepubertal and adult males (n=4 each). **I.** Graph showing the resting membrane

potential of individual cells from prepubertal and adult females (n=8 each). ***

$p<0.001$. **J.** Representative current-clamp recording demonstrating leptin (100 nM) -

induced hyperpolarization in a DAT-Cre tdTomato neuron of a female prepubertal

mouse. The dashed line indicates resting membrane potential (-67 mV). **K.** Pie chart

representing the percentage of neurons that hyperpolarized or did not show a change in

membrane potential to 100nM of leptin in prepubertal females (N=8). All data shown

are average ± SEM. Scales in A-B = 50 µm. C-F = 20 µm.

Figure 5. *Slc6a3* **(DAT) gene expression in the ventral premammillary nucleus**

(PMv) of prepubertal mice is affected by nutrition and leptin. A. Graph showing the

quantification of *Slc6a3*-^{33}P hybridization signal as relative expression (%) in diestrous

wild type *vs.* prepubertal Lep^{ob} females injected with saline or with leptin. n=7 for

diestrous and for Lep^{ob} + saline, n = 5 for Lep^{ob} + leptin. **B-C.** Dark-field images

showing the $Slc6a3$-^{33}P hybridization signal (silver grains) in the PMv of prepubertal

females (P20) from normal litter (NL) size (B) and from small litter (SL) size (C). **D.**

Graph showing the quantification of the $Slc6a3$ hybridization signal (integrated optic

density) in P20 females from normal and small size litters. n=4 animals for normal litter

and n = 5 animals for small litter. **E.** Graph showing the correlation between female

body mass and the $Slc6a3$ hybridization signal (IOD) in P20 females from normal and

small size litters. Data shown are average ± SEM. ** $p<0.01$. Scale bar in B = 100 µm.

Figure 6. Dopamine-transporter neurons in the ventral premammillary nucleus
(PMv^{DAT}) project to the AVPV and PeN and contact kisspeptin neurons. A-D.
Fluorescent image showing ChR2-mCherry signal at the site of the injection in the PMv

A, and the projections to the Anteroventral periventricular nucleus (AVPV, B), the

Periventricular nucleus (PeN) and the Medial preoptic area (MPA, C), and presence of

few projections in the arcuate nucleus (Arc, D). **E.** Confocal fluorescent maximum

intensity projection image showing the lack of mCherry (magenta) projections close to

kisspeptin (*kiss1*hrGFP, khrGFP) neurons (green) in the Arc. Insets are higher

magnification single z plane images showing detail of mCherry signal in the external

part of the dorsal Arc and the lack of contacts to khrGFP neurons in this area. **F.**

Confocal fluorescent maximum intensity projection image showing mCherry signal

(magenta) in the AVPV/PeN and the intense interaction of these projections to khrGFP

neurons (green) in this area. are higher magnification single z plane images showing

detail of the close contacts between these two populations (arrowheads). f: fornix; 3V:

Third ventricle. Scale bar A-D = 50 µm. E-F = 20 µm.

Figure 7. DAT^{Cre};tdTomato projections to dopaminergic cells in the anteroventral

periventricular and periventricular nuclei (AVPV/PeN) arise after puberty. A-D.

Fluorescent images showing tdTomato (magenta) and tyrosine hydroxylase (TH)

immunoreactivity (green) in prepubertal (A, C) and in diestrous (B, D)

DAT^{Cre};tdTomato females. **E.** Graph showing the quantification of the tdTomato fiber

density (Integrated optical density, IOD) in prepubertal *vs*. adult diestrous females in the

AVPV and the PeN areas. **F.** Graph showing the number of TH neurons per section in

the AVPV and in the PeN in prepubertal *vs*. diestrous female mice. **G.** Confocal

fluorescent maximum intensity projection image showing tdTomato (magenta) and TH

(green) cells and fibers in the lateral region of the AVPV/PeN in an adult diestrous

female. **H.** Higher magnification confocal fluorescent maximum intensity projection

image of a different field from G., showing tdTomato fibers in the PeN (magenta) and

the intense interaction of these projections to TH neurons (green) in this area. All data

shown are average ± SEM. ** p<0.01, *** p<0.001. Scale bars in A-D = 50 µm. G = 10

µm. H = 5 µm.

References

1. Cameron, J. L., Weltzin, T. E., McConaha, C., Helmreich, D. L., and Kaye, W.
H. (1991) Slowing of pulsatile luteinizing hormone secretion in men after forty-
eight hours of fasting. *J. Clin. Endocrinol. Metab.* **73**, 35–41

2. Kennedy, G. and Mitra, J. (1963) Body weight and food intake as initiating
factors for puberty in the rat. *J. Physiol.* **166**, 408–418

3. Cagampang, F., Maeda, K., Yokoyama, A., and Ota, K. (1990) Effect of food
deprivation on the pulsatile LH release in the cycling and ovariectomized female
rat. *Horm. Metab. Res.* **22**, 269–272

4. Donato, J., Silva, R. J., Sita, L. V, Lee, S., Lee, C., Lacchini, S., Bittencourt, J.
C., Franci, C. R., Canteras, N. S., and Elias, C. F. (2009) The ventral
premammillary nucleus links fasting-induced changes in leptin levels and
coordinated luteinizing hormone secretion. *J. Neurosci.* **29**, 5240–5250

5. Biro, F. M., Khoury, P., and Morrison, J. A. (2006) Influence of obesity on
timing of puberty. *Int. J. Androl.* **29**, 272–277

6. Burt Solorzano, C. M. and McCartney, C. R. (2010) Obesity and the pubertal
transition in girls and boys. *Reproduction* **140**, 399–410

7. Mahany, E. B., Han, X., Borges, B. C., Da Silveira Cruz-Machado, S., Allen, S.
J., Garcia-Galiano, D., Hoenerhoff, M. J., Bellefontaine, N. H., and Elias, C. F.
(2018) Obesity and High-Fat Diet Induce Distinct Changes in Placental Gene
Expression and Pregnancy Outcome. *Endocrinology* **159**, 1718–1733

8. Hill, J. W. and Elias, C. F. (2018) Neuroanatomical Framework of the Metabolic
Control of Reproduction. *Physiol. Rev.* **98**, 2349–2380

9. Anderson, G. M., Hill, J. W., Kaiser, U. B., Navarro, V. M., Ong, K. K., Perry, J.
R. B., Prevot, V., Tena-Sempere, M., and Elias, C. F. (2024) Metabolic control of
puberty: 60 years in the footsteps of Kennedy and Mitra's seminal work. *Nat.*
Rev. Endocrinol. **20**, 111–123

10. Quennell, J. H., Mulligan, A. C., Tups, A., Liu, X., Phipps, S. J., Kemp, C. J.,
Herbison, A. E., Grattan, D. R., and Anderson, G. M. (2009) Leptin indirectly
regulates gonadotropin-releasing hormone neuronal function. *Endocrinology* **150**,
2805–2812

11. Ahima, R. S., Prabakaran, D., Mantzoros, C., Qu, D., Lowell, B., Maratos-flier,
E., and Flier, J. S. (1996) Role of leptin in neuroendocrine response to fasting.

Nature **382**, 250–252

12. Ahima, R. S., Dushay, J., Flier, S. N., Prabakaran, D., and Flier, J. S. (1997)
Leptin accelerates the onset of puberty in normal female mice. *J. Clin. Invest.* **99**,
391–395

13. Farooqi, I. S., Wangensteen, T., Collins, S., Kimber, W., Matarese, G., Keogh, J.
M., Lank, E., Bottomley, B., Lopez-Fernandez, J., Ferraz-Amaro, I., Dattani, M.
T., Ercan, O., Myhre, A. G., Retterstol, L., Stanhope, R., Edge, J. A., Mckenzie,
S., Lessan, N., Ghodsi, M., De Rosa, V., Perna, F., Fontana, S., Barroso, I.,
Undlien, D. E., and O'Rahilly, S. (2007) Clinical and Molecular Genetic
Spectrum of Congenital Deficiency of the Leptin Receptor. *N. Engl. J. Med.* **356**,
237–247

14. De Luca, C., Kowalski, T. J., Zhang, Y., Elmquist, J. K., Lee, C., Kilimann, M.
W., Ludwig, T., Liu, S. M., and Chua, S. C. (2005) Complete rescue of obesity,
diabetes, and infertility in db/db mice by neuron-specific LEPR-B transgenes. *J.*
Clin. Invest. **115**, 3484–3493

15. Tartaglia, L. A., Dembski, M., Weng, X., Deng, N., Culpepper, J., Devos, R.,
Richards, G. J., Campfield, L. A., Clark, F. T., Deeds, J., Muir, C., Sanker, S.,
Moriarty, A., Moore, K. J., Smutko, J. S., Mays, G. G., Woolf, E. A., Monroe, C.
A., and Tepper, R. I. (1995) Identification and expression cloning of a leptin
receptor, OB-R. *Cell* **83**, 1263–71

16. Donato, J., Cravo, R. M., Frazão, R., Gautron, L., Scott, M. M., Lachey, J.,
Castro, I. A., Margatho, L. O., Lee, S., Lee, C., Richardson, J. A., Friedman, J.,
Chua, S., Coppari, R., Zigman, J. M., Elmquist, J. K., and Elias, C. F. (2011)
Leptin's effect on puberty in mice is relayed by the ventral premammillary
nucleus and does not require signaling in Kiss1 neurons. *J. Clin. Invest.* **121**,
355–368

17. Scott, M. M., Lachey, J. L., Sternson, S. M., Lee, C. E., Elias, C. F., Friedman, J.
M., and Elmquist, J. K. (2009) Leptin targets in the mouse brain. *J. Comp.*
Neurol. **514**, 518–532

18. Leshan, R. L., Louis, G. W., Jo, Y. H., Rhodes, C. J., Münzberg, H., and Myers,
M. G. (2009) Direct innervation of GnRH neurons by metabolic- and sexual
odorant-sensing leptin receptor neurons in the hypothalamic ventral
premammillary nucleus. *J. Neurosci.* **29**, 3138–3147

19. Louis, G. W., Greenwald-Yarnell, M., Phillips, R., Coolen, L. M., Lehman, M.
N., and Myers, M. G. (2011) Molecular mapping of the neural pathways linking
leptin to the neuroendocrine reproductive axis. *Endocrinology* **152**, 2302–2310

20. Sáenz de Miera, C., Bellefontaine, N., Allen, S. J., Myers, M. G., and Elias, C. F.
(2024) Glutamate neurotransmission from leptin receptor cells is required for
typical puberty and reproductive function in female mice. *Elife* **13**, RP93204

21. Ross, R. A., Leon, S., Madara, J. C., Schafer, D., Fergani, C., Maguire, C. A.,
Verstegen, A. M. J., Brengle, E., Kong, D., Herbison, A. E., Kaiser, U. B.,
Lowell, B. B., and Navarro, V. M. (2018) PACAP neurons in the ventral
premammillary nucleus regulate reproductive function in the female mouse. *Elife*
7, e35960

22. Williams, K. W., Sohn, J.-W., Donato, J., Lee, C. E., Zhao, J. J., Elmquist, J. K.,
and Elias, C. F. (2011) The acute effects of leptin require PI3K signaling in the
hypothalamic ventral premammillary nucleus. *J. Neurosci.* **31**, 13147–13156

23. Boehm, U., Zou, Z., and Buck, L. B. (2005) Feedback loops link odor and
pheromone signaling with reproduction. *Cell* **123**, 683–695

24. Meister, B. and Elde, R. (1993) Dopamine transporter mRNA in neurons of the
rat hypothalamus. *Neuroendocrinology* **58**, 388–395

25. Soden, M. E., Miller, S. M., Burgeno, L. M., Phillips, P. E. M., Hnasko, T. S.,
and Zweifel, L. S. (2016) Genetic Isolation of Hypothalamic Neurons that
Regulate Context-Specific Male Social Behavior. *Cell Rep.* **16**, 304–313

26. Stagkourakis, S., Spigolon, G., Williams, P., Protzmann, J., Fisone, G., and
Broberger, C. (2018) A neural network for intermale aggression to establish
social hierarchy. *Nat. Neurosci.* **21**, 834–842

27. Yip, S. H., York, J., Hyland, B., Bunn, S. J., and Grattan, D. R. (2018)
Incomplete concordance of dopamine transporter Cre (DATIREScre)-mediated
recombination and tyrosine hydroxylase immunoreactivity in the mouse
forebrain. *J. Chem. Neuroanat.* **90**, 40–48

28. Chen, A.-X., Yan, J.-J., Zhang, W., Wang, L., Yu, Z.-X., Ding, X.-J., Wang, D.-
Y., Zhang, M., Zhang, Y.-L., Song, N., Jiao, Z.-L., Xu, C., Zhu, S.-J., and Xu,
X.-H. (2020) Specific Hypothalamic Neurons Required for Sensing Conspecific
Male Cues Relevant to Inter-male Aggression. *Neuron* **108**, 763-774.e6

29. Stagkourakis, S., Williams, P., Spigolon, G., Khanal, S., Ziegler, K., Heikkinen,
L., Fisone, G., and Broberger, C. (2024) Maternal Aggression Driven by the
Transient Mobilisation of a Dormant Hormone-Sensitive Circuit. *bioRxiv,* doi:
10.1101/2023.02.02.526862

30. Bäckman, C. M., Malik, N., Zhang, Y. J., Shan, L., Grinberg, A., Hoffer, B. J.,
Westphal, H., and Tomac, A. C. (2006) Characterization of a mouse strain
expressing Cre recombinase from the 3′ untranslated region of the dopamine
transporter locus. *Genesis* **44**, 383–390

31. Ekstrand, M. I., Terzioglu, M., Galter, D., Zhu, S., Hofstetter, C., Lindqvist, E.,
Thams, S., Bergstrand, A., Hansson, F. S., Trifunovic, A., Hoffer, B., Cullheim,
S., Mohammed, A. H., Olson, L., and Larsson, N. G. (2007) Progressive
parkinsonism in mice with respiratory-chain-deficient dopamine neurons. *Proc.*
Natl. Acad. Sci. U. S. A. **104**, 1325–1330

32. Cravo, R. M., Frazao, R., Perello, M., Osborne-Lawrence, S., Williams, K. W.,
Zigman, J. M., Vianna, C., and Elias, C. F. (2013) Leptin Signaling in Kiss1
Neurons Arises after Pubertal Development. *PLoS One* **8**, e58698

33. Donato, J. J., Lee, C., Ratra, D., Franci, C., Canteras, N., and Elias, C. (2013)
Lesions of the ventral premammillary nucleus disrupt the dynamic changes in
kiss1 and gnrh expression characteristic of the proestrus – estrus transition.
Neuroscience **241**, 67–79

34. Krashes, M. J., Roth, B. L., Lowell, B. B., Koda, S., Ye, C., Rogan, S. C.,
Adams, A. C., Cusher, D. S., and Maratos-flier, E. (2011) Rapid, reversible

activation of AgRP neurons drives feeding behavior in mice. *J. Clin. Invest.* **121**,
1424–1428

35.   Sáenz de Miera, C., Feng, J., Elias, C. F., and Qi, N. (2023) Remote Neuronal
Activation Coupled with Automated Blood Sampling to Induce and Measure
Circulating Luteinizing Hormone in Mice. *J. Vis. Exp.* **198**, e65875

36.   Han, X., Burger, L. L., Garcia-Galiano, D., Sim, S., Allen, S. J., Olson, D. P.,
Myers, M. G., and Elias, C. F. (2020) Hypothalamic and Cell-Specific
Transcriptomes Unravel a Dynamic Neuropil Remodeling in Leptin-Induced and
Typical Pubertal Transition in Female Mice. *iScience* **23**, 101563

37.   Ingalls, A. M., Dickie, M. M., and Snell, G. D. (1950) Obese, a new mutation in
the house mouse. *J. Hered.* **41**, 315–317

38.   Bohlen, T. M., Silveira, M. A., Zampieri, T. T., Frazão, R., and Donato, J. (2016)
Fatness rather than leptin sensitivity determines the timing of puberty in female
mice. *Mol. Cell. Endocrinol.* **423**, 11–21

39.   Castellano, J. M., Bentsen, A. H., Sánchez-Garrido, M. A., Ruiz-Pino, F.,
Romero, M., Garcia-Galiano, D., Aguilar, E., Pinilla, L., Diéguez, C., Mikkelsen,
J. D., and Tena-Sempere, M. (2011) Early metabolic programming of puberty
onset: Impact of changes in postnatal feeding and rearing conditions on the
timing of puberty and development of the hypothalamic kisspeptin system.
Endocrinology **152**, 3396–3408

40.   Semaan, S. J., Murray, E. K., Poling, M. C., Dhamija, S., Forger, N. G., and
Kauffman, A. S. (2010) BAX-dependent and BAX-independent regulation of
Kiss1 neuron development in mice. *Endocrinology* **151**, 5807–5817

41.   Clarkson, J. and Herbison, A. E. (2011) Dual Phenotype Kisspeptin☐Dopamine
Neurones of the Rostral Periventricular Area. *J. Neuroendocrinol.* **23**, 293–301

42.   Kumar, D., Candlish, M., Periasamy, V., Avcu, N., Mayer, C., and Boehm, U.
(2015) Specialized subpopulations of kisspeptin neurons communicate with gnrh
neurons in female mice. *Endocrinology* **156**, 32–38

43.   Stephens, S. B., Rouse, M. L., Tolson, K. P., Liaw, R. B., Parra, R. A., Chahal,
N., and Kauffman, A. S. (2017) Effects of selective deletion of tyrosine
hydroxylase from kisspeptin cells on puberty and reproduction in male and
female mice. *eNeuro* **4**, e0150-17.2017

44.   Clarkson, J., Boon, W. C., Simpson, E. R., and Herbison, A. E. (2009) Postnatal
development of an estradiol-kisspeptin positive feedback mechanism implicated
in puberty onset. *Endocrinology* **150**, 3214–3220

45.   Silveira, M. A., Zampieri, T. T., Furigo, I. C., Abdulkader, F., Donato, J., and
Frazão, R. (2019) Acute effects of somatomammotropin hormones on neuronal
components of the hypothalamic-pituitary-gonadal axis. *Brain Res.* **1714**, 210–
217

46.   Hoffman, B. J., Hansson, S. R., Mezey, É., and Palkovits, M. (1998) Localization
and dynamic regulation of biogenic amine transporters in the mammalian central
nervous system. *Front. Neuroendocrinol.* **19**, 187–231

47.   Karasawa, N., Arai, R., Isomura, G., Yamada, K., Sakai, K., Sakai, M., Nagatsu,

T., and Nagatsu, I. (1994) Phenotypic changes of AADC-only immunopositive
premammillary neurons in the brain of laboratory shrew Suncus murinus by
systemic administration of monoamine precursors. *Neurosci. Lett.* **179**, 65–70

48.   de Vriend, V. A., van 't Sant, L. J., Rozeboom, A., Luijendijk-berg, M. C.,
Omrani, A., and Adan, R. A. (2021) Leptin Receptor Expressing Neurons in the
Substantia Nigra Regulate Locomotion , and in The Ventral Tegmental Area
Motivation and Feeding. *Front. Endocrinol.* **12**, 680494

49.   Opland, D. M., Leinninger, G. M., and Myers, M. G. (2010) Modulation of the
mesolimbic dopamine system by leptin. *Brain Res.* **1350**, 65–70

50.   Xu, Y., Lu, Y., Xu, P., Mangieri, L. R., Isingrini, E., Xu, Y., Giros, B., and Tong,
Q. (2017) VMAT2-mediated neurotransmission from midbrain leptin receptor
neurons in feeding regulation. *eNeuro* **4**, e0083-17.2017

51.   Fulton, S., Pissios, P., Manchon, R. P., Stiles, L., Frank, L., Pothos, E. N.,
Maratos-Flier, E., and Flier, J. S. (2006) Leptin Regulation of the
Mesoaccumbens Dopamine Pathway. *Neuron* **51**, 811–822

52.   Perry, M. L., Leinninger, G. M., Chen, R., Luderman, K. D., Yang, H., Gnegy,
M. E., Myers, M. G., and Kennedy, R. T. (2010) Leptin promotes dopamine
transporter and tyrosine hydroxylase activity in the nucleus accumbens of
Sprague-Dawley rats. *J. Neurochem.* **114**, 666–674

53.   Amara, S. G. and Kuhar, M. J. (1993) Neurotransmitter Transporters: Recent
progress. *Annu. Rev. Neurosci.* **16**, 73–93

54.   Merlino, D. J., Barton, J. R., Charsar, B. A., Byrne, M. D., Rappaport, J. A.,
Smeyne, R. J., Lepore, A. C., Snook, A. E., and Waldman, S. A. (2019) Two
distinct GUCY2C circuits with PMV (hypothalamic) and SN/VTA (midbrain)
origin. *Brain Struct. Funct.* **224**, 2983–2999

55.   Wang, L., Vanacker, C., Burger, L. L., Barnes, T., Shah, Y. M., Myers, M. G.,
and Moenter, S. M. (2019) Genetic dissection of the different roles of
hypothalamic kisspeptin neurons in regulating female reproduction. *Elife* **8**,
e43999

56.   Wagenmaker, E. R. and Moenter, S. M. (2017) Exposure to acute psychosocial
stress disrupts the luteinizing hormone surge independent of estrous cycle
alterations in female mice. *Endocrinology* **158**, 2593–2602

57.   Beltramino, C. and Taleisnik, S. (1985) Ventral premammillary nuclei mediate
pheromonal-induced LH release stimuli in the rat. *Neuroendocrinology* **41**, 119–
124

58.   Bleier, R., Byne, W., and Siggelkow, I. (1982) Cytoarchitectonic sexual
dimorphisms of the medial preoptic and anterior hypothalamic areas in guinea
pig, rat, hamster, and mouse. *J. Comp. Neurol.* **212**, 118–130

59.   Simerly, R. B., Swanson, L. W., and Gorski, R. A. (1985) The distribution of
monoaminergic cells and fibers in a periventricular preoptic nucleus involved in
the control of gonadotropin release: Immunohistochemical evidence for a
dopaminergic sexual dimorphism. *Brain Res.* **330**, 55–64

60.   Simerly, R. B., Zee, M. C., Pendleton, J. W., Lubahn, D. B., and Korach, K. S.

(1997) Estrogen receptor-dependent sexual differentiation of dopaminergic
neurons in the preoptic region of the mouse. *Proc. Natl. Acad. Sci. U. S. A.* **94**,
14077–14082

61.   Clarkson, J. and Herbison, A. E. (2006) Postnatal development of kisspeptin
neurons in mouse hypothalamus; sexual dimorphism and projections to
gonadotropin-releasing hormone neurons. *Endocrinology* **147**, 5817–5825

62.   Björklund, A. and Hökfelt, T., eds. (1984) Handbook of Chemical
Neuroanatomy. Vol 2: Classical Transmitters in the CNS. Part 1. Elsevier
Science Publishers B.V.

63.   Lorang, D., Amara, S. G., and Simerly, R. B. (1994) Cell-type-specific
expression of catecholamine transporters in the rat brain. *J. Neurosci.* **14**, 4903–
4914

64.   Ciliax, B. J., Drash, G. W., Staley, J. K., Haber, S., Mobley, C. J., Miller, G. W.,
Mufson, E. J., Mash, D. C., and Levey, A. I. (1999) Immunocytochemical
localization of the dopamine transporter in human brain. *J. Comp. Neurol.* **409**,
38–56

65.   Stagkourakis, S., Kim, H., Lyons, D. J., and Broberger, C. (2016) Dopamine
Autoreceptor Regulation of a Hypothalamic Dopaminergic Network. *Cell Rep.*
15, 735–747

66.   Scott, N., Prigge, M., Yizhar, O., and Kimchi, T. (2015) A sexually dimorphic
hypothalamic circuit controls maternal care and oxytocin secretion. *Nature* **525**,
519–522

A Prepubertal
B Diestrous
pSTAT3 tdTomato

C **D** **E** **F**

G pSTAT3
Number of cells
Females   Males
● Prepubertal
○ Adult

H pSTAT3/tdTomato
% Colocalization
100
50
0
Females   Males

I Resting Membrane Potential
mV
-50
-55
-60
-65
-70
-75
***
Prepubertal   Diestrous

J Prepubertal
100 nM leptin
-67mV
10mV
1min

K
37.5%
62.5%
Hyperpolarized
Non responding or continuous depolarization

A

Relative expression %

** **

Diestrous Lep^{ob} Lep^{ob}
            + Sal   + Lep

B NL   C SL

PMv   3V

                         $Slc6a3$-^{33}P

D   P20

IOD

**

Normal   Small
litter   litter

E   P20

NL
SL

r=0.78
p=0.01

Body mass (g)

A Prepubertal — AVPV — 3V

B Diestrous

C Prepubertal — 3V — PeN

D Diestrous — MPA

E

Fibers IOD

AVPV ** PeN **

F

#TH neurons

● Prepubertal
○ Diestrous

AVPV *** PeN ***

G 10 μm

H