Bacteriophage purification using CIMmultus monolithic OH-column chromatography for therapeutic purposes

Arne Echterhof[1,2*], Tejas Dharmaraj[1], Patrick Blankenberg[1], Maryam Hajfathalian[1], Francis Blankenberg[1], Jessica Sacher[1], and Paul L. Bollyky[1].

[1]Division of Infectious Diseases and Geographic Medicine, Department of Medicine, Stanford University School of Medicine, Stanford, CA 94305, USA.
[2]Institute of Medical Microbiology, University Hospital of Muenster, Muenster, Germany

Corresponding Author: Paul L. Bollyky, MD, PhD, Division of Infectious Diseases and Geographic Medicine, Department of Medicine, Stanford University School of Medicine, Beckman Center for Molecular and Genetic Medicine, 279 Campus Drive, Stanford, CA 94305, USA. pbollyky@stanford.edu.

Keywords:
Bacteriophage, Purification, FPLC, Chromatography, Endotoxin, CIMmultus Monolith

Abstract

Bacteriophages, viruses that selectively infect bacteria, are increasingly explored as potential therapies to combat antibiotic-resistant infections. Effective phage purification is vital for therapeutic use, particularly to reduce endotoxin (lipopolysaccharide, LPS) and host protein contamination from production processes in gram-negative bacterial lysates. This study investigates the purification efficiency of CIMmultus OH-monolithic chromatography for seven similarly sized tailed phages (Myoviridae and Siphoviridae families) infecting *Pseudomonas aeruginosa*. Using preferential exclusion chromatography with a potassium phosphate gradient, we achieved significant reductions in host proteins and endotoxins, yielding therapeutically compliant samples suitable for human intravenous injection. We observed recovery rates of 85-95% for smaller phages (e.g., PAML-31-1, LPS5), with lower recoveries for larger "giant" phages (e.g., OMKO1, PhiKZ). All purified samples met endotoxin limits, with average reductions of 98.5% for proteins and up to 6.30 log reductions for endotoxins. Despite increased DNA content post-purification in most samples, the approach shows promise for clinical-grade phage purification. This chromatographic strategy is scalable, reproducible, and free from hazardous chemicals, making it suitable for industrial phage production.

Introduction

Bacteriophages (Phages) are viruses that exclusively replicate in bacteria as their host organism. These nano-sized particles comprise proteins encapsulating a DNA or RNA genome [1]. Due to their selective antimicrobial targeting, bacteriophages are a potential treatment option for bacterial infection as an alternative or supplement to conventional antibiotics.

Félix d'Herelle introduced phages as a novel antibacterial treatment and published non-randomized trials from medical experience over 100 years ago [2]. The dawn of the antibiotic age in the 1930s, the lack of reproducibility, and the contradictory results attenuate the enthusiasm for phage therapy in Western countries [3]. However, the emergence of antibiotic-resistant bacteria has restored researchers' interest in reconsidering phages as a potential treatment for antibiotic-resistant pathogens [4,5]. To this day, there are no phage medications approved in Europe or the US hence phage therapy is applied under emergency investigational new drug (eIND) application or under the umbrella of Article 37 of the Declaration of Helsinki [6–8], nonetheless, clinical trials for various phage applications are ongoing [9,10].

Bacteriophages are the most numerous and ubiquitous biological entities on Earth. This is represented in the diversity of different phage morphologies which can be classified as tailed phages, tailless icosahedral phages, filamentous phages, and pleomorphic phages with or without lipid-containing envelopes. Besides diverse morphology, which leads to individual purification challenges among phage entities, phages must be produced in a bacterial host cell [11]. Consequently, key impurities might be host DNA host proteins, media components, and, in the case of gram-negative bacteria, Lipopolysaccharides (LPS). The fine line between removing key impurities while keeping the phages bioactive is the major hurdle in phage purification.

The downstream processing of the infected culture broth should isolate, concentrate, and purify the target phage. Usually, centrifugation and microfiltration steps are used to remove cells and cell debris [12]. Next, dead-end or tangential flow filtration, precipitation, gradient ultracentrifugation, or various chromatography approaches concentrate and purify phages. Phage purification for therapeutic applications requires additional standards since chemicals like CsCl, Triton 100, or chloroform commonly used for virus preparation are considered hazardous by regulatory agencies [11].

LPS remains critical in pharmaceutical production processes due to its ability to cause lethal immune responses in humans when intravenously injected at low-dose concentrations [13]. Lipopolysaccharides are molecules within the membrane of gram-negative bacteria and consist of three distinct regions: a hydrophobic lipid A region, a core oligosaccharide, and a hydrophilic O-specific polysaccharide [14]. As an endotoxin, it is released after cell death, which makes it abandonedly present in gram-negative bacterial lysate therefore, an endotoxin removal step must be conducted in every downstream processing step of gram-negative-based phage production. Due to their amphipathic properties, LPS molecules are difficult to remove from aqueous solutions based on their ability to form aggregates such as micelles (300-1000 kDa) and vesicles (>1000 kDa) stabilized by divalent Calcium and Magnesium cations [15]. The pyrogenicity of LPS requires removal in all biologicals, and the final concentration depends on its use. An intravenous or intrathecal application should not exceed the upper threshold of 5 Endotoxin Units (EU) daily per kilogram of body mass [16].

Phages are nano-sized particles and sensible to shear stress, which makes purification challenging in a conventional porous-solid matrices-based chromatography approach. Convective Interactive Media monolithic columns offer unique advantages regarding virus particle purification. The laminar flow created in monoliths reduces shear stress that leads to a higher recovery of big particles in comparison to bead or membrane-based columns independent of the flow rate, solute size, or viscosity. Additionally, the large channels reduce pressure drop across the column bed The surface of the highly interconnected network can be modified with different

solid phase chemistries such as preferential exclusion columns, ion exchange columns, or
hydrogen bond columns [17,18].
Preferential exclusion describes phenomena where co-solvents like cosmotropic ions are
excluded from the solvation shell of solutes, here bacteriophages. In the presence of cosmotropic
co-solvents like potassium phosphate, a water-cosolvent interaction is enthalpically favorable
resulting in a salt-free area around biomolecules. A hydroxyl-substituted (OH) column is weakly
hydrophobic and the solid phase is preferentially excluded, too. The biomolecules interact with
the solid phase. Column retention of phages eliminates the target particle from the system by
reducing the phage concentration from the solution. This reduces the overall salt-free area and,
therefore, minimizes the free energy to reach a thermodynamically favored state. By reducing the
salt concentration during the elution process, the particles are released from the solid phase and
become soluble again [18–20].
CIMmultus monolithic columns have gained interest in the field of phage purification.
Several papers describe using anion exchange monolithic chromatography to bind the negatively
charged viral particle to the surface. Smrekar et al. used strong anion-exchange quaternary amine
(QA) convective interaction media (CIM) monolith to capture a T4 phage from clarified E. coli
lysate [21]. Adriaenssens et al. reported that eleven morphologically distinct phages can be
purified by either QA or DEAE with a 40-99% recovery rate. However, neither paper includes
measurements regarding LPS concentrations [22]. Van Belleghem et al. investigated the LPS
removal from PNM and Luz 19 phages with an anion-exchange diethyl aminoethyl (DEAE)
convective interaction media (CIM) monolith. They reported a limited reduction of endotoxin level
and a strong titer reduction after chromatography [23]. Lee et al. demonstrated in their study that
a CIMmultus monolithic OH column can be used in steric exclusion chromatography with a
filamentous M13 phage by applying a polyethylene glycol (PEG) gradient. They report a 90%
recovery and a reduction of 98% of host protein [17]. Rebula and colleagues present a two-step
chromatographic purification approach for a PP-01 phage, including a capturing step with a
CIMmultus monolithic OH column and a polishing step using three different column options
CIMmultus with QA, H-Bond, and PrimaS ligands. They reported a 7-log endotoxin reduction by
using H-Bond and PrimaS columns [18].
In our recent study, we investigated the potential of a 1ml CIMmultus monolithic OH
column to purify seven similarly sized and shaped phages with a preferential exclusion approach
using a potassium phosphate salt gradient.
117
118
119 **Results**
120
Phages with similar Morphology and DLS measurements can be purified using OH-monolithic
chromatography
For this study, we investigated seven different *P. aeruginosa* phages regarding their
purification potential using an FPLC chromatography system with a CIMmultus 1ml OH-monolithic
column. All these seven phages share a similar-tailed phage-like morphology and similar
dimensions. Our set of phages includes six members of Myvoridae PAML-31-1, LPS5, OMKO1,
Terra, PhiKZ and DMS3vir (**Fig 1**). Within this group, OMKO1 And PhiKZ can be classified as
giant phages. Furthermore TIVP-H6 belongs to the family of Siphoviridae. PAML-31-1 LPS5…
share a similar head length from …nm to … nm and a tail length from … nm to … An exception
regarding the tail length is TIVP-H6 with … nm. The two included Giant Phages PhiKZ and
OMKO1 have a head length around … nm and a tail length of … nm. We report that those bigger
phages have a lower recovery yield compared to the smaller phages tested in this study reaching
from 42 to 35%.

The z-average measured by DLS does not correlate with the transmission electron
microscopy size……

OH-monolithic chromatography has a high recovery for phages of Myoviridae and Siphoviridae
family

The processed bacterial lysates of the seven lytic *P. aeruginosa* (PAML-31-1, OMKO1,
LPS5, TIVP-H6, Terra, DMS3vir, and PhiKZ) phages were mixed with a 3M KH_2PO_4 buffer (pH
7.0) for a final salt concentration of 1.5M to match the salt potassium phosphate concentration of
the binding buffer (buffer A). After column equilibration, the mixed phage samples were loaded
into the FPLC system by setting the P960 sample pump to a flow rate of 1ml/min. The bound
phages were eluted by linearly decreasing the potassium phosphate concentration to 20mM over
20 CV. Phage quantities of the sample application and the elution phase were assessed by plaque
assay of the pooled collected fractions (**Fig 2**). Most of the phage input for each investigated
phage sample was eluted in the collected elution fraction, while almost no phages were detected
in the sample application phase. CIMmultus monolithic OH-column binds, concentrates, and
elutes the seven investigated phages. The high UV absorbance in the sample application phase
represents unbound impurities like host and media protein, whereas the sharp UV peak during
the linear gradient represents the concentrated phage elution (**Fig.2**). The antipseudomonal
phages PAML-31-1, LPS5 and TIVP-H6 showed the best recovery ranging from 85-95 percent.
PAML-31-1 has a titer of 4.52E+11 PFU/ml in the elution fraction, and TIVP-H6 has a titer of
2.55E+11 PFU/ml. Terra and DMS3vir present with a 61% and a 41% recovery, reaching a final
titer in the pooled elution of 1.14E+11 PFU/ml and a 9.00E+10 PFU/ml. Interestingly, the two giant
phages included in this study, OMKO1 and PhiKZ indicate a lower recovery from the OH column
ranging from 42-35%. OMKO1 has a titer of 2.14E+09 PFU/ml in the elution fraction, and PhiKZ
has a titer of 1.01E+10 PFU/ml (**Table 1**). The lower recovery might be due to increasing shear
stress over the sample application phase induced through already bound giant phages at the
stationary phase. We observed an increased pressure escalation for OMKO1 over the sample
application phase in comparison to the smaller phages like LPS5

OH-monolithic chromatography effectively removes protein and endotoxin impurities from phage
lysates

After removing the bacterial debris with centrifugation and a filtration step, the clarified
bacterial lysates of all seven phages still contain protein and endotoxin impurities in abundance.
The total protein content of all tested phage lysates is, on average, 1880 mg/ml, measured by a
BCA assay. The total average lipopolysaccharide quantity of all tested phages lysates is
1.32E+05 EU/ml, measured with an Endozyme II assay. After performing a washing step of 10
CV to remove unbound samples from the stationary phase and after elution with a linear gradient
over 20 CV, the endotoxin and protein were significantly removed. On average 98.5% of total
protein was removed from the phage lysate. PAML-31-1 elution fraction contains 92.38 mg/ml
total protein, which corresponds to a 1.34 log reduction, while LPS5 elution fraction holds 29.12
mg/ml of total protein, correlating to a 1.79 log reduction, and purified PhiKZ fraction comprises
17.39 mg/ml of total protein, which computes a 2.04 log reduction.
Our phage chromatography approach achieves an average endotoxins removal of 100%
Terra elution fraction holds 66.62 EU/ml total LPS, correlating to a 2.89 log reduction, while LPS5
elution fraction contains 19.83 EU/ml of LPS, which corresponds to a 3.03 log reduction of LPS,
and purified PAML-31-1 fraction holds 0.05 EU/ml of total LPS, which computes a 6.30 log
reduction. For a summary of all purity parameters of all studied phages, see **Table 1**.
The preferential exclusion chromatography conducted with an OH-monolithic column
using a potassium phosphate buffer regimen reliably purifies the seven tested Myoviridae and
Siphovirus from protein and endotoxin contamination. All seven phages meet therapeutical

standards for human intravenous injection regarding the endotoxin level with a given dose of at
least 1E+10 PFU/ml per application applied three times daily for a hypothetical patient of 75kg.
We report a slight increase in DNA molecules after our preferential exclusion
chromatography purification step. On average, the clarified and Benzonase-treated lysate
comprises a total DNA concentration of 246.33 ng/ml. After purification, only the Terra elution
fraction shows a decrease of DNA concentration of 85.83%, correlating with a 0.88 log reduction
The other phage pooled elution fractions contain a higher DNA concentration than their
corresponding pre-chromatography solution.
190
191
192 **Discussion**
193
Here, we report a chromatographic bacteriophage purification approach using a CIMmultus 1 ml
OH column in a preferential exclusion mode as a reliable, fast, and straightforward procedure to
purify and concentrate seven similar-sized tailed bacteriophages (Myoviridae and Siphoviridae)
against *P. aeruginosa*. Our purity assays reveal a significant LPS and protein reduction after
FPLC for all the studied phages. The final LPS concentration of all phage samples meets
therapeutic standards for intravenous injection, considering the maximum threshold of 5
Endotoxin Units (EU) daily per kilogram of body mass. The method can be conducted within three
days without potentially hazardous substances like CsCl, chloroform, or Triton 100.
Convective Media Interaction (CIM) columns offer beneficial properties for purifying nano-
sized particles like phages if binding and elution parameters are optimized. Fast and laminar flow
rates combined with low shear stress, high capacities for large molecules, and flow-unaffected
resolution create optimal phage purification conditions. Additionally, CIMmultus monolithic
columns can be used multiple times and offer the option to scale up the purification production
due to the availability of column volumes reaching from 1 ml to 8000 ml. When optimized, this
chromatography approach can elevate bacteriophage purification to an industrial standard and
would lower production costs if the equipment were regularly. Moreover, CIMmultus monolithic
columns can be combined with versatile column chemistries such as anion-exchange columns
like quaternary amine (QA), Diethyl aminoethyl (DEAE), Hydrogen Bond (H-Bond ADC) columns,
multimodal anion exchange columns with hydrogen bonding potential (PrimaS) and weak
hydrophobic hydroxyl-substituted (OH) column as it is used in this study. The adaptable array of
column chemistry offers the potential to purify a broad spectrum of phages with diverse
morphology. Further studies need to be conducted to optimize chromatographic parameters for
individual phage structures.
Our chromatography approach increases the total DNA content in every Phage sample
except Terra, although all phage lysates were incubated with Benzonase. These results are
contradictory to Rebula et al. [18]. They report a 99% DNA reduction after a chromatographic step
with a CIMmultus monolithic OH-column. It needs to be emphasized that this group does not treat
the bacterial lysate with Benzonase. More investigations are needed to further understand the
optimal way to reduce DNA content from phage lysates. This can be seen as one limitation of our
study.

233
Material and Methods

Bacteriophage propagation and purification

Bacteriophages were propagated in a liquid culture in Erlenmeyer flasks at a suitable
volume. LB medium was inoculated with the *P. aeruginosa* host strain and incubated until an early
to mid-log growth phase was reached (37°C 275 rpm). Planktonic cultures were infected with a
multiplicity of infection (MOI) of 0.01 and incubated overnight. Gross bacterial debris was removed
by two to three cycles of centrifugation (8000xg, 20min 4°C) and the supernatant was filtered
through a 0.22µm polyethersulfone (PES) membrane (Corning, NY, Product No. 4311188). The
lysate was treated with 0.05U/ml Benzonase Nuclease (Sigma-Aldrich, St. Louis, MO, USA; Cat.
No. E8263) overnight to digest free DNA.

Monolithic chromatography phage purification

Benzonase-treated lysates were then mixed with 3M KH_2PO_4 buffer (pH 7.0) in a 1:1 ratio
in preparation for purification by fast protein liquid chromatography (FPLC). The FPLC-based
virus purification was conducted with a CIMmultus OH 1 mL monolithic column with a 2 µm
channel size (Sartorius BIA Separations, Ajdovščina, Slovenia). The columns washing,
regeneration, performance test, and storage were carried out according to manufacturer
recommendations. The column was attached to an ÄKTA Pure FPLC (GE Healthcare
Biosciences, Sweden) equipped with a P960 sample pump and analyzed with UNICORN 5.0
software (Cytiva Life Sciences, Marlborough, MA, USA). All purification protocols were run at
room temperature and all buffers and samples were filtered through a 0.22 µm PES membrane.
10-60 mL of the mixed sample was loaded onto a pre-equilibrated column with 1.5 M KH_2PO_4
loading buffer, pH 7.0. After sample application, the column was washed with 10 column volumes
of loading buffer to remove unbound particles. The elution of bacteriophage was performed using
a linear gradient over 20 column volumes from 1.5M to 20 mM KH_2PO_4. The fraction
corresponding to the eluted phage was collected based on UV A280 measurement. The column
was cleaned by using a 1M NaOH solution followed by a pH neutralization step with 200mM Tris
pH 7.5 and ddH_2O, before Re-equilibration with 1.5 M KH_2PO_4 loading buffer, pH 7.0.

Phage enumeration

Plaque assays were conducted to quantify the number of infectious phage particles per
mL by using the spot dilution double agar overlay method. 100 µL of mid-log phase bacteria was
added to 5 mL of top agar (5 g/L agar, 10 g/L tryptone, 10 g/L NaCl, 20 mM $MgSO_4$, 20 mM $CaCl_2$)
and poured evenly onto the surface of an LB-agar plates to solidify. 10-fold serial dilutions of
phages were prepared in 96-well polystyrene U-bottom tissue culture-treated plates
(ThermoFisher Scientific, Waltham, MA, USA, Cat. No. 168136) in SM+gel buffer (50 mM Tris-
HCl, 100 mM NaCl, 8 mM $MgSO_4$, 0.01% gelatin (Sigma-Aldrich, St. Louis, MO, USA, Cat. No.
G7041), pH 7.50). 10 µL of each dilution was spotted onto the top agar, incubated at 37 °C
overnight, and visible plaques were counted to calculate the PFU/ml values.

Total Protein, total DNA, and Endotoxin quantification

The total protein content of samples was measured using a bicinchoninic acid (BCA)
assay (Thermo Scientific, USA, Pierce BCA Protein Assay kit Cat. No. 23225) according to the
manufacturer's instructions. The total DNA was determined by a Quant-iT PicoGreen assay
(Invitrogen, USA Cat. No. P11496) according to the manufacturer's manual. The endotoxin level
of samples was measured using a recombinant Factor C Endotoxin Detection assay (bioMérieux,
France, Endonext EndoZyme II assay Cat. No. 423129) with an assay range from 0.005 to 50
EU/ml. Samples were diluted with Endotoxin-free water to attain the assay range. All microplate

assays were analyzed using an automated, multifunctional microplate reader (Tecan SPARK
multimode reader, Switzerland).

Transmission electron microscopy (TEM)

TEM imaging was done as previously reported [24]. In brief, the size and morphology of
phages were examined with transmission electron microscopy (TEM) using a JEOL JEM1400
(JEOL USA Inc., Peabody, MA) at 80 kV. 5 µL of diluted phage solution was dropped onto carbon-
coated copper grids (FCF-200-Cu, Electron Microscopy Sciences, Hatfield, PA). After 3 minutes,
the grid was dipped into a ddH$_2$O droplet and then 1% uranyl acetate for staining was dropped at
the sample and was allowed to dry for 15 minutes before performing microscopy.

Dynamic light scattering (DLS)

DLS measurement is well established in our laboratory [24]. Briefly, all DLS size
distributions were conducted with a Nano-Series Zeta Sizer (Nano-NS ZEN3600; Malvern
Instruments, Worcestershire, UK) equipped with a 633-nm laser. Measurements were obtained
at 25°C at a backscattering angle of 173°. Each DLS measurement (single replicate) reported in
this study is the average of 11 10-s measurements obtained after a 1-min vortexing period on
half-speed, 1-min equilibration period in the instrument, and with variable attenuation to generate
count rates >100 kcps. Phage diffusion coefficients were calculated from auto-correlated light
intensity data, and hydrodynamic diameters (DH) were computed using ZetaSizer version 7
software with the Stokes-Einstein equation.

Data availability

Data are available upon reasonable request from the corresponding authors subject to
institutional review and approval. Supporting data values associated with the main manuscript is
available in supplementary materials.

Author Contributions
Conceptualization:PLB, AE, TD. Methodology: AE, TD, MH. Investigation and Data Processing:
AE, TD, MH. Data Analysis: AE, TD, PLB. Writing: PLB, TD, AE.

Acknowledgments
We gratefully acknowledge the following funding: National Institutes of Health grant R01
HL148184-01, R01 AI12492093, R01 DC019965, and grants from the Cystic Fibrosis Foundation
(CFF), the Cystic Fibrosis Research Institute (CFRI) and the Emerson Collective grant (to PLB)
and grants from the Stanford University Medical Scientist Training Program grant T32-GM007365
and the Stanford Interdisciplinary Graduate Fellowship, Gold Family Graduate Fellow (to TD) and
the Merrigan Fellowship (to AE). This project was funded in part with Federal funds from the
National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of
Health and Human Services, under Contract No. HHSN272201700020I, Task Order
75N93021F00003 (A-57).

Competing Interest Statement

References

[1]   Q. Chen, T. Dharmaraj, P.C. Cai, E.B. Burgener, N.L. Haddock, A.J. Spakowitz, P.L. Bollyky, Bacteriophage and Bacterial Susceptibility, Resistance, and Tolerance to Antibiotics, Pharmaceutics 14 (2022). https://doi.org/10.3390/PHARMACEUTICS14071425.

[2]   d'Herelle F, Bacteriophage as a Treatment in Acute Medical and Surgical Infections, Bull N Y Acad Med 7 (1931) 329–48. https://pubmed.ncbi.nlm.nih.gov/19311785/ (accessed October 22, 2024).

[3]   M.D. Eaton, S. Bayne Jones, BACTERIOPHAGE THERAPY: REVIEW OF THE PRINCIPLES AND RESULTS OF THE USE OF BACTERIOPHAGE IN THE TREATMENT OF INFECTIONS, J Am Med Assoc 103 (1934) 1769–1776. https://doi.org/10.1001/JAMA.1934.72750490003007.

[4]   D.G.J. Larsson, C.F. Flach, Antibiotic resistance in the environment, Nature Reviews Microbiology 2021 20:5 20 (2021) 257–269. https://doi.org/10.1038/s41579-021-00649-x.

[5]   W.H. Organisation, 2021 Antibacterial Agents in Clinical and Preclinical Development: an overview and analysis, World Health Organization 2021 (2022). https://www.who.int/publications/i/item/9789240047655 (accessed October 23, 2024).

[6]   A. Khosravi, Q. Chen, A. Echterhof, J.L. Koff, P.L. Bollyky, Phage Therapy for Respiratory Infections: Opportunities and Challenges, Lung 202 (2024) 223–232. https://doi.org/10.1007/S00408-024-00700-7.

[7]   S. Aslam, E. Lampley, D. Wooten, M. Karris, C. Benson, S. Strathdee, R.T. Schooley, Lessons Learned From the First 10 Consecutive Cases of Intravenous Bacteriophage Therapy to Treat Multidrug-Resistant Bacterial Infections at a Single Center in the United States, Open Forum Infect Dis 7 (2020). https://doi.org/10.1093/OFID/OFAA389.

[8]   J.P. Pirnay, S. Djebara, G. Steurs, J. Griselain, C. Cochez, S. De Soir, T. Glonti, A. Spiessens, E. Vanden Berghe, S. Green, J. Wagemans, C. Lood, E. Schrevens, N. Chanishvili, M. Kutateladze, M. de Jode, P.J. Ceyssens, J.P. Draye, G. Verbeken, D. De Vos, T. Rose, J. Onsea, B. Van Nieuwenhuyse, P. Soentjens, R. Lavigne, M. Merabishvili, K.W. Pang, W.J. Metsemakers, D. Van der Linden, O. Chatzis, A. Eskenazi, A. Lopez, A. De Voeght, A.F. Rousseau, A. Tilmanne, D. Vens, J. Gérain, B. Layeux, E. Vlieghe, I. Baar, S. Van Ierssel, J. Van Laethem, J. Guiot, S. De Roock, S. Jennes, S. Uyttebroek, L. Van Gerven, P.W. Hellings, L. Dupont, Y. Debaveye, D. Devolder, I. Spriet, P. De Munter, M. Depypere, M. Vanfleteren, O. Cornu, S. Verhulst, T. Boiy, S. Lamote, T. Van Zele, G. Wieërs, C. Courtin, D. Lebeaux, J. Sartre, T. Ferry, F. Laurent, K. Paul, M. Di Luca, S. Gottschlich, T. Tkhilaishvili, N. Cesta, K. Racenis, T. Barbosa, L.E. López-Cortés, M. Tomás, M. Hübner, T.T. Pham, P. Nagtegaal, J. Ten Oever, J. Daniels, M. Loubert, G. Iheb, J. Jones, L. Hall, M. Young, N. Balarjishvili, M. Tediashvili, Y. Tong, C. Rohde, J. Wittmann, R. Hazan, R. Nir-Paz, J. Azeredo, V. Krylov, D. Cameron, M. Pitton, Y.A. Que, G. Resch, S. McCallin, M. Dunne, S. Kilcher, Personalized bacteriophage therapy outcomes for 100 consecutive cases: a

multicentre, multinational, retrospective observational study, Nature Microbiology
2024 9:6 9 (2024) 1434–1453. https://doi.org/10.1038/s41564-024-01705-x.

[9]   CYstic Fibrosis bacterioPHage Study at Yale (CYPHY) ClinicalTrials.gov identifier:
NCT04684641. Updated 2023-11-18. Accessed October 15, 2024. , (n.d.).

[10] Nebulized Bacteriophage Therapy in Cystic Fibrosis Patients With Chronic
Pseudomonas Aeruginosa Pulmonary Infection. ClinicalTrials.gov identifier:
NCT05010577. Updated 2023-10-18. Accessed October 15, 2024, (n.d.).

[11] J. João, J. Lampreia, D.M.F. Prazeres, A.M. Azevedo, Manufacturing of
bacteriophages for therapeutic applications, Biotechnol Adv 49 (2021).
https://doi.org/10.1016/J.BIOTECHADV.2021.107758.

[12] J. Gill, P. Hyman, Phage choice, isolation, and preparation for phage therapy, Curr
Pharm Biotechnol 11 (2010) 2–14. https://doi.org/10.2174/138920110790725311.

[13] P. de Oliveira Magalhães, A.M. Lopes, P.G. Mazzola, C. Rangel-Yagui, T.C.V. Penna,
A. Pessoa, Methods of endotoxin removal from biological preparations: a review, J
Pharm Pharm Sci 10 (2007) 388–404. https://pubmed.ncbi.nlm.nih.gov/17727802/
(accessed October 22, 2024).

[14] C. Galanos, M.A. Freudenberg, Bacterial endotoxins: biological properties and
mechanisms of action, Mediators Inflamm 2 (1993) S11–S16.
https://doi.org/10.1155/S0962935193000687.

[15] D. Petsch, F.B. Anspach, Endotoxin removal from protein solutions, J Biotechnol 76
(2000) 97–119. https://doi.org/10.1016/S0168-1656(99)00185-6.

[16] K. Regulski, P. Champion-Arnaud, J. Gabard, Bacteriophage Manufacturing: From
Early Twentieth-Century Processes to Current GMP, Bacteriophages: Biology,
Technology, Therapy, Volume 2 (2021) 699–729. https://doi.org/10.1007/978-3-319-
41986-2_25.

[17] J. Lee, H.T. Gan, S.M.A. Latiff, C. Chuah, W.Y. Lee, Y.S. Yang, B. Loo, S.K. Ng, P.
Gagnon, Principles and applications of steric exclusion chromatography, J
Chromatogr A 1270 (2012) 162–170.
https://doi.org/10.1016/J.CHROMA.2012.10.062.

[18] L. Rebula, A. Raspor, M. Bavčar, A. Štrancar, M. Leskovec, CIM monolithic
chromatography as a useful tool for endotoxin reduction and purification of
bacteriophage particles supported with PAT analytics, J Chromatogr B Analyt
Technol Biomed Life Sci 1217 (2023).
https://doi.org/10.1016/J.JCHROMB.2023.123606.

[19] S. Moelbert, B. Normand, P. De Los Rios, Kosmotropes and chaotropes: modelling
preferential exclusion, binding and aggregate stability, Biophys Chem 112 (2004) 45–
57. https://doi.org/10.1016/J.BPC.2004.06.012.

[20] S.N. Timasheff, T. Arakawa, Mechanism of protein precipitation and stabilization by
co-solvents, J Cryst Growth 90 (1988) 39–46. https://doi.org/10.1016/0022-
0248(88)90296-5.

[21] F. Smrekar, M. Ciringer, M. Peterka, A. Podgornik, A. Štrancar, Purification and
concentration of bacteriophage T4 using monolithic chromatographic supports, J
Chromatogr B Analyt Technol Biomed Life Sci 861 (2008) 177–180.
https://doi.org/10.1016/J.JCHROMB.2007.05.048.

[22]   E.M. Adriaenssens, S.M. Lehman, K. Vandersteegen, D. Vandenheuvel, D.L. Philippe,
412   A. Cornelissen, M.R.J. Clokie, A.J. García, M. De Proft, M. Maes, R. Lavigne, CIM(®)
413   monolithic anion-exchange chromatography as a useful alternative to CsCl gradient
414   purification of bacteriophage particles, Virology 434 (2012) 265–270.
415   https://doi.org/10.1016/J.VIROL.2012.09.018.
416   [23]   J.D. Van Belleghem, M. Merabishvili, B. Vergauwen, R. Lavigne, M. Vaneechoutte, A
417   comparative study of different strategies for removal of endotoxins from
418   bacteriophage preparations, J Microbiol Methods 132 (2017) 153–159.
419   https://doi.org/10.1016/J.MIMET.2016.11.020.
420   [24]   T. Dharmaraj, M.J. Kratochvil, J.D. Pourtois, Q. Chen, M. Hajfathalian, A. Hargil, Y.H.
421   Lin, Z. Evans, A. Oromí-Bosch, J.D. Berry, R. McBride, N.L. Haddock, D.R. Holman,
422   J.D. Van Belleghem, T.H. Chang, J.J. Barr, R. Lavigne, S.C. Heilshorn, F.G.
423   Blankenberg, P.L. Bollyky, Rapid assessment of changes in phage bioactivity using
424   dynamic light scattering, PNAS Nexus 2 (2023).
425   https://doi.org/10.1093/PNASNEXUS/PGAD406.
426
427

A

PAML-31-1	Morphology: *Myoviridae* Head length: 75 ± 2 nm Tail length: 147 ± 5 nm Tail width: 21 ± 2 nm
LPS5	Morphology: *Myoviridae* Head length: 76 ± 2 nm Tail length: 143 ± 4 nm Tail width: 24 ± 2 nm
OMKO1	Morphology: *Myoviridae* Head length: 129 ± 9 nm Tail length: 200 ± 19 nm Tail width: 27 ± 3 nm
TIVP-H6	Morphology: *Siphoviridae* Head length: 68 ± 4 nm Tail length: 178 ± 12 nm Tail width: 17 ± 2 nm

@Maryam: TEM pictures of Terra, DMS3vir, and PhiKZ would be a great addition to finalizing this figure. It would be much appreciated

B

PAML-31-1

Z-average: 94.9 nm
PDI: 0.093

LPS5

Z-average: 147.6 nm
PDI: 0.188

OMKO1

Z-average: 218.1 nm
PDI: 0.147

TIVP-H6

Z-average: 113 nm
PDI: 0.163

Terra

Z-average: 80.9 nm
PDI: 0.15

DMS3vir

Z-average: 122.4 nm
PDI: 0.168

PhiKZ

Z-average: 114.3 nm
PDI: 0.188

Figure 1: Size characteristics of 7 lytic *P. aeruginosa* phages used in this study. A. A representative transmission electron microscopy image of every is shown at 80,000x magnification. Head length, tail length, and tail width are measured on at least 5 different transmission electron microscopy examples for each phage**. B.** Dynamic light scattering (DLS) spectra, Z-average and PDI values of purified phages used in this study.

Figure 2: Monolithic Chromatography preparation of PAML-31-1 phage clarified lysate on CIMmultus OH 1ml column on Akta purifier HPLC system. 60 ml of PAML-31-1 phage lysate diluted to 1.5 M KH_2PO_4, pH 7.0 is loaded. Loading Buffer A: 1.5 M KH_2PO_4, pH 7.0, Elution Buffer B: 20mM KH_2PO_4, pH 7.0. Upper lines indicate different phases of the method: Flowthrough phase (**FT**), 10 CV washing the column after sample application with 10 CV (**W$_{0\%}$**), Elution with a linear Gradient over 20 CV to 100% Elution Buffer (**E**), final wash with 100% Buffer B to regenerate the column (**W$_{100\%}$**). Detection: blue: UV-absorbance (280 nm), red: Conductivity (mS/cm), green: Buffer B (%). Created in BioRender.com

Phages	Titer (spot assay)			Endotoxin (Endozyme II)			
	Lysate (PFU/ml)	Elution (PFU/ml)	Recovery (%)	Lysate (EU/ml)	Elution (EU/ml)	depletion (%)	log reduction
PAML-31-1	1.85E+11	4.52E+11	0.94	1.09E+05	0.05	100.00	6.30
OMKO1	2.10E+09	2.14E+09	0.42	4.41E+04	0.47	100.00	4.97
LPS5	6.75E+10	3.10E+11	0.94	3.27E+04	19.83	99.99	3.22
T4P-H6	2.00E+11	2.55E+11	0.85	6.78E+04	35.51	99.96	3.28
Terra	1.25E+11	1.15E+11	0.61	5.15E+04	66.62	99.86	2.89
DMS3vir	1.48E+11	9.00E+10	0.41	4.54E+05	31.58	99.99	4.16
PhiKZ	1.93E+10	1.01E+10	0.35	1.85E+05	19.89	99.98	3.97

Phages	Protein (BCA assay)				DNA (Picogreen assay)			
	Lysate (mg/ml)	Elution (mg/ml)	depletion (%)	log reduction	Lysate (ng/ml)	Elution (ng/ml)	depletion (%)	log reduction
PAML-31-1	2031.69	92.38	98.14	1.34	461.14	1402.92		
OMKO1	1516.66	65.13	95.78	1.37	215.11	926.36		
LPS5	1792.98	29.12	99.65	1.79	277.52	856.66		
T4P-H6	1738.51	29.58	98.67	1.77	264.71	1155.81		
Terra	2031.00	17.12	99.08	2.07	123.08	16.04	85.83	0.88
DMS3vir	2139.04	27.33	97.91	1.89	190.91	226.99		
PhiKZ	1913.44	17.39	98.27	2.04	191.84	130.66		0.17

Table 1: Summary of purification results. Titer (plaque assay), endotoxin (Endozyme II assay), protein (BCA assay), and DNA (Picogreen assay) concentrations before and after chromatography using the CIMmultus OH-column of six lytic Pseudomonas aeruginosa phages. Calculated values are phage recovery (%), impurity depletion (%), and impurity log reduction.