1 **C3H-HeN as a TLR4 competent mouse model of sublethal leptospirosis**

2 Olifan Zewdie Abil[a], Suman Kundu[a], Leonardo Moura Midon[a] and Maria Gomes-Solecki[a*]

3 [a]Department of Microbiology, Immunology and Biochemistry, University of Tennessee Health

4 Science Center, Memphis, Tennessee, United States of America.

5 *Corresponding author: mgomesso@uthsc.edu

Abstract

Abstract

Mice are slowly being accepted as alternative models for investigation of leptospiral infection.

One of the most used strains, the C3H-HeJ model, express a *tlr4* gene in its immune cells that is

hyporesponsive to leptospiral LPS and thus the model is deemed immunocompromised. To help

resolve this valid scientific concern we did a study in which we compared infection with

Leptospira interrogans serovar Copenhageni Fiocruz in mice expressing fully competent *tlr4*

(C57BL6, C3H-HeN) versus hyporesponsive *tlr4* (C3H-HeJ) 15 days post-infection. Overall, we

confirm that the C3H-HeJ mouse is more susceptible to *L. interrogans* infection, as this was the

only strain that sustained lethal infection. However, we found that the two mouse strains with a

fully functional *tlr4* gene (C3H-HeN and C57BL/6) also developed clinical and molecular signs

of sublethal leptospirosis nearly on par with C3H-HeJ, as quantified by weight loss, survival

curves, presence of *Leptospira* 16S rRNA in blood and urine, and burden of viable spirochetes in

kidney. Regarding kidney markers of inflammation and fibrosis, C3H-HeN and C57BL/6

produced less IL-1 beta, iNOS and ColA1 than C3H-HeJ, which is consistent with increased

resilience to infection expected of the *tlr4* competent strains. Regarding Leptospira-specific

antibody, all mouse strains produced IgM and T-cell independent IgG3. However, in contrast to

C57BL/6, both C3H strains produced circulating IgG1 but only C3H-HeN produced IgG2a to *L.*

interrogans at two weeks post infection. Thus, C3H-HeN can be used as a TLR4-competent

mouse model of sublethal leptospirosis.

Keywords: C3H, C3H-HeN, mouse model, leptospirosis, TLR4, *Leptospira interrogans*

Author summary

Previous studies by different laboratories have shown C3H-HeJ to be a susceptible mouse strain

to leptospirosis. The goal of our study was to evaluate if low *tlr4* responsiveness to leptospiral

LPS in C3H-HeJ mice is the factor that drives susceptibility to infection with pathogenic

Leptospira. We did a comparative study using mouse strains immunocompetent for *tlr4* (C3H-

HeN and C57BL/6). The data shows that although *tlr4* does affect the virulence of leptospirosis

(only C3H-HeJ succumbed to infection), the *tlr4* competent strains (C3H-HeN and C57BL/6)

also developed measurable sub-lethal leptospirosis. Thus, competent recognition of leptospiral-

LPS by murine TLR4 is not sufficient to cause susceptibility to leptospirosis. Furthermore, the

data hints that T-cell dependent responses could be involved in susceptibility to leptospirosis in

the C3H models of infection.

Introduction

Leptospirosis, an often overlooked but resurging infectious disease caused by a spirochete

bacteria, is a significant global health concern, impacting millions of individuals worldwide. This

disease carries a high mortality rate, resulting in approximately 65,000 deaths annually [1].

Furthermore, it poses a serious threat to animals of agricultural importance, leading to substantial

economic losses, particularly in tropical and subtropical regions [2]. Although significant efforts

have been made to formulate novel vaccination approaches that confer enduring immunity and

safeguard against various serovars, our understanding of the specific immune factors

contributing to host defense and disease progression remains limited [3].

Leptospirosis research is challenging due to the inconsistent outcomes observed in leptospiral

infections involving small animals, large animals and humans [4] and a plethora of high and low

virulence *Leptospira* serovars [5-8]. Hamsters and guinea pigs have been widely used as small-

animal models of acute disease, as they recapitulate the manifestations of severe disease in

humans. Rats have been used as a model for studying the severity of human leptospirosis as they

become chronically infected and shed *Leptospira* in urine many months after infection [9, 10].

However, the usage of these animals in research is limited in some parts of the world where the

disease is endemic due to stringent animal regulations [4] and the scarcity of accessible reagents

for routine experiments.

Mouse models are extensively utilized in biomedical research because they provide a versatile

approach due to the accessibility of diverse immunological and genetic tools, as well as a broad

range of available reagents [4]. Researchers have studied the outcomes of experimental

leptospiral infection using various mouse models, including studies on lethal, sublethal, and

chronic leptospirosis manifestations [4]. For instance, C57BL/6 and BALB/c, exhibit increased

resilience to acute disease and have the potential to serve as models for persistent infection

caused by *Leptospira interrogans* [11, 12]. C3H-HeJ mice infected with *L. interrogans* develop

disease that can be easily monitored through measurement of clinical scores. This strain produces

valuable models of lethal [13-15] and sublethal disease [16], and has been used for investigating

inflammatory signatures of infection and necroptosis, immunology and efficacy of protection

studies [17-19]. Toll-like receptor (TLR) 4, the LPS receptor, was shown to play a central role in

the control of leptospirosis [20, 21]. One mechanism described how leptospiral-LPS activates

murine, but not human, TLR4 in cultured macrophages [22], and it is associated with resistance

to infection [23]. C3H-HeJ mice have a single amino acid substitution (aa712, P to H) within the

coding region of the *tlr4* gene that makes this molecule hyporesponsive to the atypical

Leptospira LPS [21, 24]. Of note, humans, also believed to be susceptible to leptospirosis

express a TLR4 molecule that does not sense the atypical *Leptospira* LPS [22]. Nevertheless, one

consistent criticism regarding the use C3H-HeJ mice is that its hyporesponsive Toll-Like

Receptor 4 (TLR4) qualifies these mice as immunocompromised. To help resolve this valid

scientific concern we did a study in which we compared infection with *L. interrogans* in mice

expressing fully competent *tlr4* (C57BL6, C3H-HeN) versus hyporesponsive *tlr4* (C3H-HeJ).

115   **Material and methods**

Animals

Male C3H-HeJ, C3H-HeN, and C57BL/6 mice (9-10 weeks old) were used for this study. C3H-HeJ animals were purchased from The Jackson Laboratory (Bar Harbor); C3H-HeN and C57BL/6 were purchased from Charles River (Wilmington, MA) and maintained in specific pathogen-free at the Laboratory Animal Care Unit of the University of Tennessee Health Science Centre, with unrestricted access to food and water. We used male mice because they are more susceptible to leptospirosis [25]. There were 4 animals per group, and the experiment was reproduced once, for a total of 24 mice. All animal procedures were carried out in accordance with the University of Tennessee Health Science Centre Institutional Animal Care and Use Committee protocol number 22-0362.

Bacterial strains and culture

L. interrogans serovar Copenhageni strain Fiocruz L1-130 (henceforth LiC), frozen in -80°C, was passaged in hamster. Their kidneys were harvested and cultivated in 4mL of Hornsby-Alt-Nally (HAN) media [26] with 100 μg/mL 5-fluorouracil (MP Chemicals, CA) at 29°C for better growth. Passage 2 was done in Ellinghausen-McCullough-Johnson-Harris (EMJH) medium supplemented with Difco *Leptospira* enrichment EMJH (Becton, MD) at 28-30°C. EMJH culture passage 2 was allowed to reach the log phase of growth, pelleted by centrifugation at 3,000 $\times g$ for 5 min, and washed and resuspended in sterile 1× phosphate-buffered saline (PBS) (Thermo Fisher Scientific). Next, the cells were used to infect the animals after counted under a dark-field microscope (Zeiss USA, Hawthorne, NY) using a Petroff- Hausser chamber as previously described [25].

Animal infection and collection of specimens

Mice were inoculated with 1×10^8 LiC in 200 μL of endotoxin-free $1\times$ PBS intraperitoneally

(i.p.). Control mice were inoculated with the same volume of endotoxin-free $1\times$ PBS. Survival

and body weight loss were monitored for 15 consecutive days post-infection. Mice were

euthanized 15 days after post-infection or when they reached the endpoint criteria (20% body

weight loss post-infection). Blood, urine, and kidney were collected for further analysis.

Bacterial quantification

The NucleoSpin tissue kit (Clontech, Mountain View, CA) was used to purify genomic DNA

from kidney, blood and urine following the manufacturer's protocol, and the purified DNA was

then stored at −20°C for further analysis. To quantify *Leptospira*, quantitative polymerase chain

reaction (qPCR) was performed using *Leptospira* 16S rRNA primers (Forward:

CCCGCGTCCGATTAG and Reverse: TCCATTGTGGCCGAACAC) and a TAMRA probe

(CTCACCAAGGCGACGATCGGTAGC) obtained from Eurofins (Huntsville, AL). The results

were reported as the number of *Leptospira* genome equivalents. The qPCR mixture consisted of

25 μM of each primer, 250 nM of the specific probe, and 2 μL of DNA sample, with a total

volume of 20 μL. Duplicate reactions were performed. The amplification protocol included an

initial step of 10 min at 95°C, followed by 40 cycles of amplification (15 s at 95°C and 1 min at

60°C). The analysis was conducted using the comparative threshold cycle (CT) method. A

negative result was determined if no amplification occurred.

Measurement of *Leptospira*-specific antibody

Leptospira-specific antibody levels were measured using the enzyme-linked immunosorbent

assay (ELISA). The process of preparing a leptospiral extract for LiC was carried out following

previous instructions [27]. Briefly, LiC was cultured in EMJH media until it reached confluency.

The bacterial cells were then separated by centrifugation to form a pellet. This pellet was

subjected to incubation with BugBuster® solution (1mL) at room temperature in a shaker (100

rpm) for 20 min and mixed thoroughly by vortexing. The resulting mixture was stored at -20°C.

The whole cell extract of *Leptospira* was appropriately diluted in a sodium carbonate coating

buffer with a concentration of 1X. A 96-well flat-bottom ELISA microtiter plate (Nunc-

eBioscience) was coated overnight at $4^{\circ}C$ with 100 µL 1X sodium carbonate coating buffer

whole-cell extract of *Leptospira* (10^7–10^8 bacteria per well). After overnight incubation, the

ELISA plate was washed with 1X PBST. The plate was blocked by a blocking buffer (100

µL/well) containing 1% BSA, followed by incubation for 1 h at $37^{\circ}C$. After washing, serum

samples (1:100) was added, and the plate was incubated for 1 h at $37^{\circ}C$. The unbound primary

antibody was removed by vigorous washing. Next, anti-mouse secondary antibodies for IgM

IgG, IgG3, IgG1 or IgG2a (all from Cell signaling technology, CST) conjugated with horseradish

peroxidase was added, which was incubated for 30 min, followed by standard color development

using TMB Sureblue. Absorbance measurement was carried out at OD 450 nm using Molecular

Devices Spetramax.

Expression of inflammatory and fibrosis markers

Total RNA was extracted using the RNeasy Mini Kit (QIAGEN) according to the manufacturer's

protocol. Complementary DNA (cDNA) was synthesized from the extracted RNA using the

RevertAid First Strand cDNA Synthesis Kit (Thermo Fisher Scientific). The resulting first-strand

cDNA served as the template for reverse transcription PCR (RT-qPCR), which was performed

on a QuantStudio™ 3 Real-Time PCR (Applied Biosystems) using the PowerTrack™ SYBR™

Green Master Mix (Applied Biosystems). Each RT-qPCR reaction (10 μL total volume) included

the cDNA template and specific primers. The cycling conditions were as follows: an initial step

at 50°C for 2 minutes, followed by denaturation at 95°C for 2 minutes, followed by 40 cycles of

95°C for 15 sec for denaturation, and 60°C for 1 min for annealing/extension. A melt curve

analysis was conducted at the end of the amplification to confirm the specificity of the PCR

products. Relative gene expression levels across samples were quantified by the double delta Ct

($2^{-\Delta\Delta Ct}$) method, with glyceraldehyde 3-phosphate dehydrogenase (GAPDH) serving as the

endogenous reference control. The primer sequences used in this study are listed in Table S1 of

the supplemental material.

Statistical analysis

We used GraphPad Prism 10 software. The *P*-value was determined by Mann-Whitney *U* test, or

Two-way ANOVA with Tukey's multiple comparison test.

207   **Results**

Weight loss, burden in blood, shedding in urine and survival after infection with *L.*

interrogans

Following experimental leptospiral infection (10^8 LiC), mice were monitored over a 15-day

period (Fig 1A). In C3H-HeJ, a steady weight loss was observed beginning on day 6, reaching a

peak around day 9, at which point one mouse met the endpoint criteria. The remaining mice

recovered gradually thereafter. The overall weight loss in this group ranged from 2% to 21% (Fig

1D). C3H-HeN mice exhibited daily weight loss from day 1 which decreased gradually until day

14. The overall weight loss in this group ranged from 7% to 16% (Fig 1E). In contrast, C57BL/6

mice experienced low % of weight loss, with an overall decrease of 1% to 8%. (Fig 1F). The

weight loss curves measured for the three strains were statistically significant (p=0.0116,

p<0.0001) as compared to their respective control groups (Fig 1D to F).

Quantitative PCR (qPCR) targeting the *Leptospira* 16S rRNA gene was performed to analyse the

dissemination of *Leptospira* in blood and urine. Blood samples were collected on days 1, 3, 6, 9,

12, and 14 post-inoculation to quantify the *Leptospira* burden. Peak *Leptospira* burdens were

observed on day 3 in C3H-HeJ mice ($\sim$3x10^4), whereas the peak burden occurred on day 1 in

both C3H-HeN and C57BL/6 mice, $\sim$6-9x10^3 (Fig 1B). *Leptospira* shedding in urine was also

assessed using qPCR on alternate days over a 15-day period. *Leptospira* shedding was detected a

low levels ($\sim$$10^2$) during the first week post-infection, peaking on day 10 in the second week for

C3H-HeN and C57BL/6 mice ($\sim$$10^4$), whereas in C3H-HeJ mice it peaked on day 14 at $\sim$$10^6$ (Fig

1C). Regarding survival, the data shows that 75% of C3H-HeJ mice survived at 15 days post-

infection, whereas 100% of C3H-HeN and C57BL/6 mice survived during this period (Fig 2).

The burden of *L. interrogans* and viability in kidney tissue after infection

To assess renal colonization by *L. interrogans*, kidney tissues from infected mice were collected

on the day the mice were euthanized on day 15 post-infection and analyzed for the presence of

Leptospira DNA using qPCR (Fig 3A). The bacterial burden was ~$7.5 \times 10^{\square}$ *Leptospira* per mg

of kidney tissue in C3H-HeJ, ~$4.1 \times 10^{\square}$ in C3H-HeN, and ~$2.1 \times 10^{\square}$ in C57BL/6 infected

mice (Fig 3B). Additionally, the viability of *Leptospira* was assessed by culturing kidney tissues

in EMJH medium at 30°C for 4 days and quantifying viable spirochetes using qPCR. On day 4

of culture, the average number of viable spirochetes per 100 μL of EMJH culture was ~4.3×10^{3}

for C3H-HeJ, ~3.1×10^{3} for C3H-HeN, and ~2.2×10^{3} for C57BL/6 mice (Fig 3C).

Expression of inflammatory cytokines and signatures of fibrosis in kidney of C3H-HeJ,

C3H-HeN and C57BL/6 mice following *L. interrogans* infection

To assess the upregulation of inflammatory cytokines critical for immune responses and fibrosis

markers associated with kidney damage, mice were inoculated with *L. interrogans* as described

in Fig 4. Kidney tissues were collected and processed for mRNA expression analysis 15 days

post-infection. The expression levels of IL-1 beta mRNA, a key cytokine involved in

inflammatory responses was significantly upregulated both in C3H-HeJ and C3H-HeN mice (Fig

4B). Importantly, mRNA of the fibrosis marker, inducible nitric oxide synthase (iNOS), was

significantly increased in both C3H-HeJ and C3H-HeN mice, but not in C57BL/6 (Fig 4C). The

mRNA expression of collagen A1 (ColA1) was also markedly elevated in both C3H-HeJ and

C3H-HeN mice and no ColA1 upregulation was detected in C57BL/6 (Fig 4D). We also

analysed the mRNA expression levels of IFN-gamma, IL-17, IL-6, and IL-23 cytokines in all

mouse strains; however, the differences observed were not statistically significant.

Serological antibody responses in C3H-HeJ, C3H-HeN and C57BL/6 at 15 days post

infection with *L. interrogans*

Terminal blood samples were collected 15 days post-infection to evaluate the levels of antibody

production to *L. interrogans* (Fig 5). As expected, the three strains of mice produced increased

levels of anti-*Leptospira*-specific IgM (Fig 5B), and IgG (Fig 5C) compared to the respective

uninfected controls. IgG isotyping produced the following results: IgG3 was the only antibody

significantly produced by the three mouse strains (Fig 5D) and IgG1/IgG2a were not detected in

C57BL/6 (Fig 5E and F). Significantly higher levels of IgG1 were measured in C3H-HeJ than

C3H-HeN compared to the respective controls (Fig 5E) and IgG2a was measured in both C3H

strains but was only significant in C3H-HeN (Fig 5F).

Discussion

We and others previously used the C3H-HeJ strain to develop a mouse model of experimental

sub-lethal leptospirosis [16, 25, 28]. Since resistance to lethal infection with *L. interrogans* was

associated with *tlr4* [23] and C3H-HeJ is hyporesponsive to *L. interrogans* atypical LPS, there

are valid concerns that this mouse model does not recapitulate a fully competent immune

response to this spirochete. In this study, we did a side-by-side comparative analysis of three

mouse strains (C3H-HeJ, C3H-HeN and C57BL/6). C3H-HeJ was the only strain that sustained

lethal infection, thus confirming that this strain is more susceptible to *L. interrogans* infection

than C3H-HeN and C57BL/6. Nevertheless, the two mouse strains with a fully functional *tlr4*

gene (C3H-HeN and C57BL/6) also developed clinical and molecular signs of sub-lethal

leptospirosis after 2 weeks of infection, nearly on par with C3H-HeJ.

We found weight loss and survival differences between C3H and C57BL/6 in that C3H mice

showed larger weight loss curves (~ 2-21%) than C57BL/6 mice (1-8%) throughout the 15-day

infection period, that in the case of C3H-HeJ, affected survival. For a choice of an animal model

of disease, differences in weight loss in the C3H background produced a clinical score that is

easier to reproduce consistently. qPCR analysis of *Leptospira* 16S rRNA demonstrated

differences in the dynamics of *L. interrogans* dissemination in blood and urinary shedding

among the 3 mouse strains that culminated in a gradient of kidney colonization which was

highest in C3H and lowest in C57BL/6. This is consistent with the severity of systemic

dissemination observed for C3H strains in weight loss and survival, suggesting that C3H mice

are more susceptible to *Leptospira* infection than C57BL/6.

Although there are numerous links between competent recognition of Leptospiral LPS by TLR4

and resistance to infection in mice [20-23], our study shows that susceptibility to infection can

also be observed in C3H-HeN mice, which express a fully competent TLR4. In addition, we

found that infection of potentially susceptible TLR4/MD-2 humanized transgenic mice with *L.*

interrogans did not produce different measurements of disease progression or *Leptospira*

dissemination than wildtype mice [29]. Overall, the data suggests that although TLR4 may play a

part in susceptibility to *Leptospira* infection in humans and certain strains of mice, it is not the

ultimate determinant as other receptors in mammalian cells (ex TLR2) likely compensate for

deficient receptor binding.

While innate immune responses are the main immunological pathway for eliminating *Leptospira*,

the humoral immune response plays a vital role in effectively eradicating the bacteria and

expelling it from the host [30]. Our present observations revealed that LiC infection led to an

increase of *Leptospira*-specific IgM, IgG antibodies in the blood of the three mouse strains. This

was expected as several other groups have reported [16, 23, 31, 32]. However, IgG isotyping in

C3H models of infection produced some interesting, unexpected results for IgG3, IgG1 and

IgG2a. IgG3 was the only antibody significantly produced by the three mouse strains and may

explain high total IgG for C57BL/6 in our study. Two weeks after infection of C57BL/6, IgG1

was extremely low as observed by others [32]. IgG2a antibodies were not detected in C57BL/6

as can be explained by the absence of a gene for IgG2a [33]. However, at this early time point of

infection (2 weeks) significantly higher levels of IgG1 were measured in C3H-HeJ than C3H-

HeN, and IgG2a was measured in both C3H strains but was only significantly higher in C3H-

HeN. The production of Th2-dependent IgG1 antibody in C3H-HeJ and C3H-HeN mice suggests

a Th2-biased activation of the immune response, as shown in our previous studies for C3H-HeJ

[16]. Presence of Th1 dependent IgG2a in C3H-HeN mice at two weeks post infection with *L.*

interrogans suggests that this response may be driven in part by TLR4 which has been associated

with resistance and may explain why disease progression in this mouse is less pronounced than

C3H-HeJ. The data strongly suggests that the mouse genetic background affects the type of IgG

antibody response to pathogenic *Leptospira*; hyporesponsiveness to TLR4 may account for lower

levels of IgG2a in the C3H-HeJ genetic background which result from impaired Th1 responses

and could explain increased susceptibility to leptospirosis.

In conclusion, our data suggests that unlike C57BL/6, the C3H-HeN mouse, a TLR4 competent

strain, can be used in lieu of C3H-HeJ to recapitulate sub-lethal leptospirosis and to investigate

competent Th1/Th2 immune responses to pathogenic *Leptospira* in early time-points of

infection. All mouse strains should be pursued in studies of immunity and host response to

pathogenic *Leptospira* as they all contribute to answer different questions.

Declaration of Competing Interest

The authors do not have competing financial interests or personal relationships that could appear

to influence the work reported in this paper.

Acknowledgments

This work was supported by NIAID grant numbers R21 AI142129 (MGS), R44 AI167605

(MGS) and R01 AI175417 (MPI MGS). The content of this manuscript is totally the

responsibility of the authors and does not involve the official views of NIH NIAID.

Author contributions

MGS designed the study. OZA, SK and LMM performed the experiments and data analysis.

OZA, SK and MGS wrote the paper.

References

367   1.   Costa, F., et al., *Global Morbidity and Mortality of Leptospirosis: A Systematic Review.*

368   PLoS Negl Trop Dis, 2015. **9**(9): p. e0003898.

369   2.   Haake, D.A. and P.N. Levett, *Leptospirosis in humans.* Curr Top Microbiol Immunol,

370   2015. **387**: p. 65-97.

371   3.   McBride, A.J., et al., *Leptospirosis.* Current Opinion in Infectious Diseases, 2005. **18**(5):

372   p. 376-386.

373   4.   Gomes-Solecki, M., I. Santecchia, and C. Werts, *Animal Models of Leptospirosis: Of*

374   *Mice and Hamsters.* Front Immunol, 2017. **8**: p. 58.

375   5.   Silva, E.F., et al., *Characterization of virulence of Leptospira isolates in a hamster*

376   *model.* Vaccine, 2008. **26**(31): p. 3892-6.

377   6.   Lehmann, J.S., et al., *Leptospiral pathogenomics.* Pathogens, 2014. **3**(2): p. 280-308.

378   7.   Philip, N., et al., *In vivo and in silico Virulence Analysis of Leptospira Species Isolated*

379   *From Environments and Rodents in Leptospirosis Outbreak Areas in Malaysia.* Front

380   Microbiol, 2021. **12**: p. 753328.

381   8.   Giraud-Gatineau, A., et al., *Evolutionary insights into the emergence of virulent*

382   *Leptospira spirochetes.* PLoS Pathog, 2024. **20**(7): p. e1012161.

383   9.   Bharti, A.R., et al., *Leptospirosis: a zoonotic disease of global importance.* Lancet Infect

384   Dis, 2003. **3**(12): p. 757-71.

385   10.   Bonilla-Santiago, R. and J.E. Nally, *Rat model of chronic leptospirosis.* Curr Protoc

386   Microbiol, 2011. **Chapter 12**: p. Unit 12E.3.

387   11.   Ratet, G., et al., *Live imaging of bioluminescent leptospira interrogans in mice reveals*

388   *renal colonization as a stealth escape from the blood defenses and antibiotics.* PLoS Negl

389   Trop Dis, 2014. **8**(12): p. e3359.

390   12.   Santos, C.S., et al., *Different outcomes of experimental leptospiral infection in mouse*

391   *strains with distinct genotypes.* J Med Microbiol, 2010. **59**(Pt 9): p. 1101-1106.

392   13.   Pereira, M.M., et al., *Morphological characterization of lung and kidney lesions in*

393   *C3H/HeJ mice infected with Leptospira interrogans serovar icterohaemorrhagiae: defect*

394   *of CD4+ and CD8+ T-cells are prognosticators of the disease progression.* Exp Toxicol

395   Pathol, 1998. **50**(3): p. 191-8.

396   14.   Nally, J.E., et al., *Lethal infection of C3H/HeJ and C3H/SCID mice with an isolate of*

397   *Leptospira interrogans serovar copenhageni.* Infect Immun, 2005. **73**(10): p. 7014-7.

398   15.   Viriyakosol, S., et al., *Toll-like receptor 4 protects against lethal Leptospira interrogans*

399   *serovar icterohaemorrhagiae infection and contributes to in vivo control of leptospiral*

400   *burden.* Infect Immun, 2006. **74**(2): p. 887-95.

401   16.   Richer, L., et al., *Mouse model for sublethal Leptospira interrogans infection.* Infect

402   Immun, 2015. **83**(12): p. 4693-700.

403   17.   Shetty, A., S. Kundu, and M. Gomes-Solecki, *Inflammatory Signatures of Pathogenic*

404   *and Non-Pathogenic Leptospira Infection in Susceptible C3H-HeJ Mice.* Front Cell

405   Infect Microbiol, 2021. **11**: p. 677999.

406   18.   Kundu, S., A. Shetty, and M. Gomes-Solecki, *Necroptosis Contributes to Persistent*

407   *Inflammation During Acute Leptospirosis.* Front Immunol, 2022. **13**: p. 810834.

19.   Kundu, S., A. Shetty, and M. Gomes-Solecki, *Exposure to live saprophytic Leptospira*

before challenge with a pathogenic serovar prevents severe leptospirosis and promotes

kidney homeostasis. Elife, 2024. **13**.

20.   Werts, C., et al., *Leptospiral lipopolysaccharide activates cells through a TLR2-*

dependent mechanism. Nat Immunol, 2001. **2**(4): p. 346-52.

21.   Que-Gewirth, N.L., et al., *A methylated phosphate group and four amide-linked acyl*

chains in leptospira interrogans lipid A. The membrane anchor of an unusual

lipopolysaccharide that activates TLR2. J Biol Chem, 2004. **279**(24): p. 25420-9.

22.   Nahori, M.A., et al., *Differential TLR recognition of leptospiral lipid A and*

lipopolysaccharide in murine and human cells. J Immunol, 2005. **175**(9): p. 6022-31.

23.   Chassin, C., et al., *TLR4- and TLR2-mediated B cell responses control the clearance of*

the bacterial pathogen, Leptospira interrogans. J Immunol, 2009. **183**(4): p. 2669-77.

24.   Qureshi, S.T., et al., *Endotoxin-tolerant mice have mutations in Toll-like receptor 4*

(Tlr4). J Exp Med, 1999. **189**(4): p. 615-25.

25.   Nair, N. and M. Gomes-Solecki, *A Mouse Model of Sublethal Leptospirosis: Protocols*

for Infection with Leptospira Through Natural Transmission Routes, for Monitoring

Clinical and Molecular Scores of Disease, and for Evaluation of the Host Immune

Response. Curr Protoc Microbiol, 2020. **59**(1): p. e127.

26.   Hornsby, R.L., D.P. Alt, and J.E. Nally, *Isolation and propagation of leptospires at*

37°C directly from the mammalian host. Sci Rep, 2020. **10**(1): p. 9620.

27.   Fortes-Gabriel, E., et al., *Enzyme immunoassays (EIA) for serodiagnosis of human*

leptospirosis: specific IgG3/IgG1 isotyping may further inform diagnosis of acute

disease. PLoS Negl Trop Dis, 2022. **16**(2): p. e0010241.

28.   Nair, N., et al., *The route of infection with Leptospira interrogans serovar Copenhageni*

affects the kinetics of bacterial dissemination and kidney colonization. PLoS Negl Trop

Dis, 2020. **14**(1): p. e0007950.

29.   Nair, N., et al., *Role of TLR4 in Persistent Leptospira interrogans Infection: A*

Comparative In Vivo Study in Mice. Front Immunol, 2020. **11**: p. 572999.

30.   Vernel-Pauillac, F. and C. Werts, *Recent findings related to immune responses against*

leptospirosis and novel strategies to prevent infection. Microbes Infect, 2018. **20**(9-10):

p. 578-588.

31.   Krangvichian, P., et al., *Sublethal infection of C3H/HeNJ against Leptospira interrogans*

serovar Pomona. Acta Trop, 2023. **238**: p. 106701.

32.   Vernel-Pauillac, F., et al., *Anti-Leptospira immunoglobulin profiling in mice reveals*

strain specific IgG and persistent IgM responses associated with virulence and renal

colonization. PLoS Negl Trop Dis, 2021. **15**(3): p. e0008970.

33.   Martin, R.M., J.L. Brady, and A.M. Lew, *The need for IgG2c specific antiserum when*

isotyping antibodies from C57BL/6 and NOD mice. J Immunol Methods, 1998. **212**(2): p.

187-92.

Figure 1. Weightloss and bacterial burden in blood and urine after infection. C3H-HeJ, C3H-HeN and C57BL/6 male mice (n=4/group) were inoculated IP with 1×10^8 *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 (LiC) and with PBS (control) (A). Quantification of bacterial load in blood (B) and urine (C) was determined by real-time qPCR of the16S rRNA gene. Body weight (% change) was recorded daily for 15 days after infection (D to F). * $p=0.0116$, **** $p < 0.0001$ by Mann-Whitney *U* test. Data shown represents one of two experiments.

Figure 2. Survival of C3H-HeJ, C3H-HeN and C57BL/6 after infection.
Male mice (n=4/group) were inoculated IP with 1 × 10^8 *L. interrogans* serovar
Copenhageni Fiocruz L1-130 (LiC) and PBS (control), and the percentage of
mice that survived infection for 15 days was recorded. Data represents one of
two experiments.

Figure 3. Burden and viability of _L. interrogans_ in kidney. C3H-HeJ, C3H-HeN and C57BL/6 male mice (n=4/group) were inoculated IP with 1 × 10^8 _L. interrogans_ serovar Copenhageni Fiocruz L1-130 (LiC) and with PBS (control). Kidney tissues were collected 2 weeks post infection for qPCR analysis of 16S RNA from DNA purified from kidney tissue (A) and from DNA purified from kidney culture in EMJH (B). Each dot represents a single mouse. *$p < 0.05$ was determined by Mann-Whitney _U_ test. Data represents one of two experiments.

Figure 4. Inflammatory cytokines and fibrosis markers in kidney of infected mice. The expression of IL-1 beta (B), iNOS (C) and ColA1 (D) in kidney tissue were measured using RT-PCR. Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used as the calibrator gene to normalize the expression of genes. Each dot represents measurement from a single mouse. *p< 0.05, **p< 0.005, ***p<0.0005 or ****p < 0.0001 was determined by two-way ANOVA followed by Tukey's multiple comparison correction test. Data shown represents one of two experiments.

Figure 5. Antibody responses to *L. interrogans* infection. C3H-HeJ, C3H-HeN and C57BL/6 male mice (n=4/group) were inoculated IP with 1×10^8 *L. interrogans* serovar Copenhageni Fiocruz L1-130 (LiC) and with PBS (control) (A). Serologic leptospiral-specific IgM (B), IgG (C), IgG3 (D), IgG1 (E) and IgG2a (F) was measured using ELISA. Each dot represents a single mouse. $*p < 0.05$, $**p < 0.01$, $*** p < 0.001$ or $****p < 0.0001$ was determined by two-way ANOVA followed by Tukey's multiple comparison correction test. Data shown represents one of two experiments.