TLR4 competence and mouse models of leptospirosis

Olifan Zewdie Abil[a+], Suman Kundu[a+], Leonardo Moura Midon[a] and Maria Gomes-Solecki[a*]

[a]Department of Microbiology, Immunology and Biochemistry, University of Tennessee Health

Science Center, Memphis, Tennessee, United States of America.

+ Equal contributions

*Corresponding author: mgomesso@uthsc.edu

Abstract

Mice are slowly being accepted as alternative models for investigation of leptospiral infection. The strain most used to analyse sublethal disease (C3H-HeJ) expresses a *tlr4* gene in its immune cells that is hyporesponsive to leptospiral LPS and thus the model is deemed immunocompromised. To help resolve this valid scientific concern we did a study in which we compared infection with *Leptospira interrogans* serovar Copenhageni Fiocruz in mice expressing fully competent *tlr4* (C3H-HeN, C57BL6) versus *tlr4* hyporesponsive mice (C3H-HeJ) over a period of two weeks. We found that the C3H-HeJ mouse was the only strain that sustained lethal infection, thus confirming the association between *tlr4* function and acute disease. We found that the two mouse strains with a fully functional *tlr4* gene (C3H-HeN and C57BL/6) also developed clinical and molecular signs of sublethal leptospirosis nearly on par with C3H-HeJ, as quantified by weight loss, survival curves, presence of *Leptospira* 16S rRNA in blood and urine, and burden of viable spirochetes in kidney. Regarding kidney markers of inflammation and fibrosis, C3H-HeN and C57BL/6 produced less IL-1 beta, iNOS and ColA1 than C3H-HeJ, which is consistent with increased resilience to infection expected of the *tlr4* competent strains. Regarding Leptospira-specific antibody, all mouse strains produced IgM and T-cell independent IgG3. Interestingly, in contrast to C57BL/6, both C3H strains produced circulating IgG1 but only C3H-HeN produced IgG2a to *L. interrogans* two weeks post infection. Thus, TLR4-competent C3H-HeN and C57BL/6 may be more appropriate mouse models of sublethal leptospirosis, whereas C3H-HeJ can be used to further establish a mouse model of lethal leptospirosis. Our data strongly suggests that TLR4 function is important but not sufficient to cause susceptibility to leptospirosis.

Keywords: C3H, mouse model, leptospirosis, TLR4, *Leptospira interrogans*

Author summary

Previous studies by different laboratories have shown C3H-HeJ to be a susceptible mouse strain

to sublethal and lethal leptospirosis. The goal of our study was to evaluate if low *tlr4*

responsiveness to leptospiral LPS in C3H-HeJ mice is the factor that drives susceptibility to

infection with pathogenic *Leptospira*. We did a comparative study using mouse strains

immunocompetent and hyporesponsive to *tlr4*. The data shows that although *tlr4* does affect the

virulence of leptospirosis (only C3H-HeJ succumbed to infection), the *tlr4* competent strains

(C3H-HeN and C57BL/6) also developed measurable sublethal leptospirosis. Thus, competent

recognition of leptospiral-LPS by murine TLR4 plays a role but it is not sufficient to determine

susceptibility to leptospirosis. Furthermore, the data suggests that T-cell dependent responses

could be involved in susceptibility to leptospirosis in the C3H models of infection.

Introduction

Leptospirosis, an often overlooked but resurging infectious disease caused by a spirochetal bacteria, is a significant global health concern, impacting millions of individuals worldwide. This disease carries a high mortality rate, resulting in approximately 65,000 deaths annually [1]. Furthermore, it poses a serious threat to animals of agricultural importance, leading to substantial economic losses, particularly in tropical and subtropical regions [2]. Although significant efforts have been made to formulate novel vaccination approaches that confer enduring immunity and safeguard against various serovars, our understanding of the specific immune factors contributing to host defense and disease progression remains limited [3].

Leptospirosis research is challenging due to the inconsistent outcomes observed in leptospiral infections involving small animals, large animals and humans [4] and a plethora of high and low virulence *Leptospira* serovars [5-8]. Hamsters and guinea pigs have been widely used as small-animal models of acute disease, as they recapitulate the manifestations of severe disease in humans. Rats have been used as a model for studying the severity of human leptospirosis as they become chronically infected and shed *Leptospira* in urine many months after infection [9, 10]. However, the usage of these animals in research is limited in some parts of the world where the disease is endemic due to stringent animal regulations [4] and the scarcity of accessible reagents for routine experiments.

Mouse models are extensively utilized in biomedical research because they provide a versatile approach due to the accessibility of diverse immunological and genetic tools, as well as a broad range of available reagents [4]. Researchers have studied the outcomes of experimental leptospiral infection using various mouse models, including studies on lethal, sublethal, and chronic leptospirosis manifestations [4]. For instance, C57BL/6 and BALB/c, exhibit increased

resilience to acute disease and have the potential to serve as models for persistent infection

caused by *Leptospira interrogans* [11, 12]. C3H-HeJ mice infected with *L. interrogans* develop

disease that can be easily monitored through measurement of clinical scores. This strain produces

valuable models of lethal [13-15] and sublethal disease [16], and has been used for investigating

inflammatory signatures of infection and necroptosis, immunology and efficacy of protection

studies [17-19]. Toll-like receptor (TLR) 4, the LPS receptor, was shown to play a central role in

the control of leptospirosis [20, 21]. One mechanism described how leptospiral-LPS activates

murine, but not human, TLR4 in cultured macrophages [22], and it is associated with resistance

to infection [23]. C3H-HeJ mice have a single amino acid substitution (aa712, P to H) within the

coding region of the *tlr4* gene that makes this molecule hyporesponsive to the atypical

Leptospira LPS [21, 24]. Of note, humans, also believed to be susceptible to leptospirosis

express a TLR4 molecule that does not sense the atypical *Leptospira* LPS [22]. Nevertheless, one

consistent criticism regarding the use C3H-HeJ mice is that its hyporesponsive Toll-Like

Receptor 4 (TLR4) qualifies these mice as immunocompromised. To help resolve this valid

scientific concern we did a study in which we compared infection with *L. interrogans* in mice

expressing fully competent *tlr4* (C57BL6, C3H-HeN) versus mice that are *tlr4* hyporesponsive

(C3H-HeJ).

Material and methods

Animals. Male, 9-10 week old C3H-HeJ, C3H-HeN, C57BL/6 mice and 7-8 week old C3H-HeJ

were used for this study. C3H-HeJ animals were purchased from The Jackson Laboratory (Bar

Harbor); C3H-HeN and C57BL/6 were purchased from Charles River (Wilmington, MA) and

maintained in specific pathogen-free at the Laboratory Animal Care Unit of the University of

Tennessee Health Science Centre, with unrestricted access to food and water. We used male

mice because they are more susceptible to leptospirosis [25]. We used 4 animals per group, and

experiments were reproduced once. All animal procedures were carried out in accordance with

the University of Tennessee Health Science Centre Institutional Animal Care and Use

Committee protocol number 22-0362.

Bacterial strains and culture. *L. interrogans* serovar Copenhageni strain Fiocruz L1-130

(henceforth LiC), frozen in -80°C, was passaged in hamster. Their kidneys were harvested and

cultivated in 4mL of Hornsby-Alt-Nally (HAN) media [26] with 100 μg/mL 5-fluorouracil (MP

Chemicals, CA) at 29°C for better growth. Passage 2 was done in Ellinghausen-McCullough-

Johnson-Harris (EMJH) medium supplemented with Difco *Leptospira* enrichment EMJH

(Becton, MD) at 28-30°C. EMJH culture passage 2 was allowed to reach the log phase of

growth, pelleted by centrifugation at $3,000 \times g$ for 5 min, and washed and resuspended in sterile

$1\times$ phosphate-buffered saline (PBS) (Thermo Fisher Scientific). Next, the cells were used to

infect the animals after counted under a dark-field microscope (Zeiss USA, Hawthorne, NY)

using a Petroff- Hausser chamber as previously described [25].

Animal infection and collection of specimens. Mice were inoculated with 1×10^8 LiC in 200 μL

of endotoxin-free $1\times$ PBS intraperitoneally (i.p.). Control mice were inoculated with the same

volume of endotoxin-free $1\times$ PBS. Survival and body weight loss were monitored for 15

consecutive days post-infection. Mice were euthanized 15 days after post-infection or when they

reached the endpoint criteria (20% body weight loss post-infection). Blood, urine, and kidney

were collected for further analysis.

Bacterial quantification. The NucleoSpin tissue kit (Clontech, Mountain View, CA) was used

to purify genomic DNA from kidney, blood and urine following the manufacturer's protocol, and

the purified DNA was then stored at −20°C for further analysis. To quantify *Leptospira*,

quantitative polymerase chain reaction (qPCR) was performed using *Leptospira* 16S rRNA

primers (Forward: CCCGCGTCCGATTAG and Reverse: TCCATTGTGGCCGAACAC) and a

TAMRA probe (CTCACCAAGGCGACGATCGGTAGC) obtained from Eurofins (Huntsville,

AL). The results were reported as the number of *Leptospira* genome equivalents. The qPCR

mixture consisted of 25 µM of each primer, 250 nM of the specific probe, and 2 µL of DNA

sample, with a total volume of 20 µL. Duplicate reactions were performed. The amplification

protocol included an initial step of 10 min at 95°C, followed by 40 cycles of amplification (15 s

at 95°C and 1 min at 60°C). The analysis was conducted using the comparative threshold cycle

(CT) method. A negative result was determined if no amplification occurred.

Measurement of *Leptospira*-specific antibody. *Leptospira*-specific antibody levels were

measured using the enzyme-linked immunosorbent assay (ELISA). The process of preparing a

leptospiral extract for LiC was carried out following previous instructions [27]. Briefly, LiC was

cultured in EMJH media until it reached confluency. The bacterial cells were then separated by

centrifugation to form a pellet. This pellet was subjected to incubation with BugBuster® solution

(1mL) at room temperature in a shaker (100 rpm) for 20 min and mixed thoroughly by vortexing.

The resulting mixture was stored at -20°C. The whole cell extract of *Leptospira* was

appropriately diluted in a sodium carbonate coating buffer with a concentration of 1X. A 96-well

flat-bottom ELISA microtiter plate (Nunc-eBioscience) was coated overnight at 4°C with 100 µL

1X sodium carbonate coating buffer whole-cell extract of *Leptospira* (10^7–10^8 bacteria per well).

After overnight incubation, the ELISA plate was washed with 1X PBST. The plate was blocked

by a blocking buffer (100 µL/well) containing 1% BSA, followed by incubation for 1 h at 37°C.

After washing, serum samples (1:100) was added, and the plate was incubated for 1 h at 37°C.

The unbound primary antibody was removed by vigorous washing. Next, anti-mouse secondary

antibodies for IgM IgG, IgG3, IgG1 or IgG2a (all from Cell signaling technology, CST)

conjugated with horseradish peroxidase was added, which was incubated for 30 min, followed by

standard color development using TMB Sureblue. Absorbance measurement was carried out at

OD 450 nm using Molecular Devices Spetramax.

Expression of inflammatory and fibrosis markers. Total RNA was extracted using the

RNeasy Mini Kit (QIAGEN) according to the manufacturer's protocol. Complementary DNA

(cDNA) was synthesized from the extracted RNA using the RevertAid First Strand cDNA

Synthesis Kit (Thermo Fisher Scientific). The resulting first-strand cDNA served as the template

for reverse transcription PCR (RT-qPCR), which was performed on a QuantStudio™ 3 Real-

Time PCR (Applied Biosystems) using the PowerTrack™ SYBR™ Green Master Mix (Applied

Biosystems). Each RT-qPCR reaction (10 µL total volume) included the cDNA template and

specific primers. The cycling conditions were as follows: an initial step at 50°C for 2 minutes,

followed by denaturation at 95°C for 2 minutes, followed by 40 cycles of 95°C for 15 sec for

denaturation, and 60°C for 1 min for annealing/extension. A melt curve analysis was conducted

at the end of the amplification to confirm the specificity of the PCR products. Relative gene

expression levels across samples were quantified by the double delta Ct ($2^{-\Delta\Delta Ct}$) method, with

glyceraldehyde 3-phosphate dehydrogenase (GAPDH) serving as the endogenous reference

control. The primer sequences used in this study are listed in Table S1 of the supplemental

material.

Statistical analysis. We used GraphPad Prism 10 software. The *P*-value was determined by

Mann-Whitney *U* test, or Two-way ANOVA with Tukey's multiple comparison test.

Results

Weight loss, burden in blood, shedding in urine and survival after infection with *L.*

interrogans. Following experimental leptospiral infection (10^8 LiC), mice were monitored over a

15-day period (Fig 1A). In C3H-HeJ, a steady weight loss was observed beginning on day 6,

reaching a peak around day 9, at which point one mouse met the endpoint criteria. The remaining

mice recovered gradually thereafter. The overall weight loss in this group ranged from 2% to

21% (Fig 1D). C3H-HeN mice exhibited daily weight loss from day 1 which decreased gradually

until day 14. The overall weight loss in this group ranged from 7% to 16% (Fig 1E). In contrast,

C57BL/6 mice experienced low % of weight loss, with an overall decrease of 1% to 8%. (Fig

1F). The weight loss curves measured for the three strains were statistically significant

($p=0.0116$, $p<0.0001$) as compared to their respective control groups (Fig 1D to F).

Quantitative PCR (qPCR) targeting the *Leptospira* 16S rRNA gene was performed to

analyse the dissemination of *Leptospira* in blood and urine. Blood samples were collected on

days 1, 3, 6, 9, 12, and 14 post-inoculation to quantify the *Leptospira* burden. Peak *Leptospira*

burdens were observed on day 3 in C3H-HeJ mice (~3×10^4), whereas the peak burden occurred

on day 1 in both C3H-HeN and C57BL/6 mice, ~6-9×10^3 (Fig 1B). *Leptospira* shedding in urine

was also assessed using qPCR on alternate days over a 15-day period. *Leptospira* shedding was

detected a low levels (~10^2) during the first week post-infection, peaking on day 10 in the second

week for C3H-HeN and C57BL/6 mice (~10^4), whereas in C3H-HeJ mice it peaked on day 14 at

~10^6 (Fig 1C). Regarding survival, the data shows that 75% of C3H-HeJ mice survived at 15

days post-infection, whereas 100% of C3H-HeN and C57BL/6 mice survived during this period

(Fig 2).

Survival of C3H-HeJ to lethal infection was further explored by using 7-8 week old mice

infected with the same dose of *L. interrogans* serovar Copenhageni FioCruz (10^8) and weight

was monitored for 15 days. We found that 100% of the mice reached 20% of weight loss

between days 9-12 post infection and met endpoint criteria for euthanasia (Supplemental Fig 1).

The burden of *L. interrogans* and viability in kidney tissue after infection

To assess renal colonization by *L. interrogans*, kidney tissues from infected mice were collected

on the day the mice were euthanized on day 15 post-infection and analyzed for the presence of

Leptospira DNA using qPCR (Fig 3A). The bacterial burden was ~7.5×10^4 *Leptospira* per mg

of kidney tissue in C3H-HeJ, ~4.1×10^4 in C3H-HeN, and ~2.1×10^4 in C57BL/6 infected mice

(Fig 3B). Additionally, the viability of *Leptospira* was assessed by culturing kidney tissues in

EMJH medium at 30°C for 4 days and quantifying viable spirochetes using qPCR. On day 4 of

culture, the average number of viable spirochetes per 100 μL of EMJH culture was ~4.3×10^3 for

C3H-HeJ, ~3.1×10^3 for C3H-HeN, and ~2.2×10^3 for C57BL/6 mice (Fig 3C).

Expression of inflammatory cytokines and signatures of fibrosis in kidney of C3H-HeJ,

C3H-HeN and C57BL/6 mice following *L. interrogans* infection. To assess the upregulation of

inflammatory cytokines critical for immune responses and fibrosis markers associated with

kidney damage, mice were inoculated with *L. interrogans* as described in Fig 4. Kidney tissues

were collected and processed for mRNA expression analysis 15 days post-infection. The

expression levels of IL-1 beta mRNA, a key cytokine involved in inflammatory responses was

significantly upregulated both in C3H-HeJ and C3H-HeN mice (Fig 4B). Importantly, mRNA of

the fibrosis marker, inducible nitric oxide synthase (iNOS), was significantly increased in both

C3H-HeJ and C3H-HeN mice, but not in C57BL/6 (Fig 4C). The mRNA expression of collagen

A1 (ColA1) was also markedly elevated in both C3H-HeJ and C3H-HeN mice and no ColA1

upregulation was detected in C57BL/6 (Fig 4D). We also analysed the mRNA expression levels

of IFN-gamma, IL-17, IL-6, and IL-23 cytokines in all mouse strains; however, the differences

observed were not statistically significant.

Serological antibody responses in C3H-HeJ, C3H-HeN and C57BL/6 at 15 days post

infection with *L. interrogans*. Terminal blood samples were collected 15 days post-infection to

evaluate the levels of antibody production to *L. interrogans* (Fig 5). As expected, the three

strains of mice produced increased levels of anti-*Leptospira*-specific IgM (Fig 5B), and IgG (Fig

5C) compared to the respective uninfected controls. IgG isotyping produced the following

results: IgG3 was the only antibody significantly produced by the three mouse strains (Fig 5D)

and IgG1/IgG2a were not detected in C57BL/6 (Fig 5E and F). Significantly higher levels of

IgG1 were measured in C3H-HeJ than C3H-HeN compared to the respective controls (Fig 5E)

and IgG2a was measured in both C3H strains but was only significant in C3H-HeN (Fig 5F).

Discussion

We previously used the C3H-HeJ strain to develop a mouse model of experimental sublethal leptospirosis [16, 25, 28]. Since resistance to lethal acute infection with *L. interrogans* was associated with *tlr4* [23] and C3H-HeJ is hyporesponsive to *L. interrogans* atypical LPS, there are valid concerns that this mouse model does not recapitulate a fully competent immune response to this spirochete. In this study, we did a side-by-side comparative analysis using three mouse strains (C3H-HeJ, C3H-HeN and C57BL/6). C3H-HeJ was the only strain that sustained lethal infection thus confirming an association between a fully functional *tlr4* gene and susceptibility to acute *L. interrogans* infection. Nevertheless, the two mouse strains with a fully functional *tlr4* gene (C3H-HeN and C57BL/6) also developed clinical and molecular signs of sublethal leptospirosis over two weeks of infection, nearly on par with C3H-HeJ. Our data indicates that TLR4-competent C3H-HeN and C57BL/6 may be more appropriate mouse models of sublethal leptospirosis, whereas C3H-HeJ may be good model of lethal leptospirosis.

We found weight loss and survival differences between C3H and C57BL/6 in that C3H mice showed larger weight loss curves (~ 2-21%) than C57BL/6 mice (1-8%) throughout the 15-day infection period, that in the case of C3H-HeJ, affected survival. For a choice of animal models of disease, differences in weight loss in the C3H background produced clinical scores that are easier to reproduce consistently. qPCR analysis of *Leptospira* 16S rRNA demonstrated differences in the dynamics of *L. interrogans* dissemination in blood and urinary shedding among the 3 mouse strains that culminated in a gradient of kidney colonization which was highest in C3H and lowest in C57BL/6. This is consistent with the severity of systemic dissemination observed for C3H strains in weight loss and survival, suggesting that overall, C3H mice are more susceptible to *Leptospira* infection than C57BL/6.

There are numerous links between competent recognition of Leptospiral LPS by TLR4

and resistance to infection in mice [20-23]. Our study confirms that hyporesponsivenes to *tlr4*

results in lethal infection in C3H-HeJ mice (Fig 1, Fig 2 and Supplemental Figure 1). Our data

also shows that susceptibility to sublethal infection can be observed in C3H-HeN and C57BL/6

mice, which express fully competent TLR4s, although differences in measurements are more

readily obtained in C3H-HeN. Of note, humans, also believed to be susceptible to leptospirosis

express a TLR4 molecule that does not sense the atypical *Leptospira* LPS [22]. In another study

we found that inoculation of potentially susceptible TLR4/MD-2 humanized transgenic mice

with *L. interrogans* did not produce different measurements of disease progression or *Leptospira*

dissemination than wildtype mice [29]. This may have been related to the humanized transgenic

mouse background being C57BL/6. Overall, the data suggests that although TLR4 plays a role

in susceptibility to *Leptospira* infection, it is not the ultimate determinant as other receptors in

mammalian cells (ex. TLR2) likely compensate for deficient receptor binding.

While innate immune responses are the main immunological pathway for eliminating

Leptospira, the humoral immune response plays a vital role in effectively eradicating the bacteria

and expelling it from the host [30]. Our present observations revealed that LiC infection led to an

increase of *Leptospira*-specific IgM, IgG antibodies in the blood of the three mouse strains. This

was expected as several other groups have reported [16, 23, 31, 32]. However, IgG isotyping in

C3H models of infection produced some interesting, unexpected results for IgG3, IgG1 and

IgG2a. IgG3 was the only antibody significantly produced by the three mouse strains and may

explain high total IgG for C57BL/6 in our study. Two weeks after infection of C57BL/6, IgG1

was extremely low as observed by others [32]. IgG2a antibodies were not detected in C57BL/6

as can be explained by the absence of a gene for IgG2a [33]. However, at this early time point of

infection (2 weeks) significantly higher levels of IgG1 were measured in C3H-HeJ than C3H-

HeN, and IgG2a was measured in both C3H strains but was only significantly higher in C3H-

HeN. The production of Th2-dependent IgG1 antibody in C3H-HeJ and C3H-HeN mice suggests

a Th2-biased activation of the immune response, as shown in our previous studies for C3H-HeJ

[16]. Presence of Th1 dependent IgG2a in C3H-HeN mice at two weeks post infection with *L.*

interrogans suggests that this response may be driven in part by TLR4 which has been associated

with resistance and may explain why disease progression in this mouse is less pronounced than

C3H-HeJ. The data strongly suggests that the mouse genetic background affects the type of IgG

antibody response to pathogenic *Leptospira*; hyporesponsiveness to TLR4 may account for lower

levels of IgG2a in the C3H-HeJ genetic background which result from impaired Th1 responses

and could explain increased susceptibility to leptospirosis.

In conclusion, our data suggests C3H-HeN and C57BL/6 mice, both TLR4 competent

strains, can be used to recapitulate sublethal leptospirosis, although the C3H-HeN strain

produces unambiguous differences in clinical and molecular measurements of disease

progression, tissue pathology and bacterial dissemination. Furthermore, and considering that

both C3H-HeJ and humans are hyporesponsive to leptospiral LPS, the C3H-HeJ strain may be

more suitable for analysis of lethal leptospirosis on par with the hamster model. In addition, both

C3H strains can be used to investigate competent Th immune responses to pathogenic *Leptospira*

in early time-points of infection. Each of the mouse strains have different characteristics that can

be leveraged in pursuit of knowledge on immunity and host response to pathogenic *Leptospira*.

Declaration of Competing Interest

Acknowledgments

This work was supported by NIAID grant numbers R21 AI142129 (MGS), R44 AI167605 (MGS) and R01 AI175417 (MPI MGS). The content of this manuscript is totally the responsibility of the authors and does not involve the official views of NIH NIAID.

Author contributions

MGS designed the study. OZA, SK and LMM performed the experiments and data analysis. OZA, SK and MGS wrote the paper.

References

1. Costa, F., et al., *Global Morbidity and Mortality of Leptospirosis: A Systematic Review.*
PLoS Negl Trop Dis, 2015. **9**(9): p. e0003898.
2. Haake, D.A. and P.N. Levett, *Leptospirosis in humans.* Curr Top Microbiol Immunol,
2015. **387**: p. 65-97.
3. McBride, A.J., et al., *Leptospirosis.* Current Opinion in Infectious Diseases, 2005. **18**(5):
p. 376-386.
4. Gomes-Solecki, M., I. Santecchia, and C. Werts, *Animal Models of Leptospirosis: Of*
Mice and Hamsters. Front Immunol, 2017. **8**: p. 58.
5. Silva, E.F., et al., *Characterization of virulence of Leptospira isolates in a hamster*
model. Vaccine, 2008. **26**(31): p. 3892-6.
6. Lehmann, J.S., et al., *Leptospiral pathogenomics.* Pathogens, 2014. **3**(2): p. 280-308.
7. Philip, N., et al., *In vivo and in silico Virulence Analysis of Leptospira Species Isolated*
From Environments and Rodents in Leptospirosis Outbreak Areas in Malaysia. Front
Microbiol, 2021. **12**: p. 753328.
8. Giraud-Gatineau, A., et al., *Evolutionary insights into the emergence of virulent*
Leptospira spirochetes. PLoS Pathog, 2024. **20**(7): p. e1012161.
9. Bharti, A.R., et al., *Leptospirosis: a zoonotic disease of global importance.* Lancet Infect
Dis, 2003. **3**(12): p. 757-71.
10. Bonilla-Santiago, R. and J.E. Nally, *Rat model of chronic leptospirosis.* Curr Protoc
Microbiol, 2011. **Chapter 12**: p. Unit 12E.3.
11. Ratet, G., et al., *Live imaging of bioluminescent leptospira interrogans in mice reveals*
renal colonization as a stealth escape from the blood defenses and antibiotics. PLoS Negl
Trop Dis, 2014. **8**(12): p. e3359.
12. Santos, C.S., et al., *Different outcomes of experimental leptospiral infection in mouse*
strains with distinct genotypes. J Med Microbiol, 2010. **59**(Pt 9): p. 1101-1106.
13. Pereira, M.M., et al., *Morphological characterization of lung and kidney lesions in*
C3H/HeJ mice infected with Leptospira interrogans serovar icterohaemorrhagiae: defect
of CD4+ and CD8+ T-cells are prognosticators of the disease progression. Exp Toxicol
Pathol, 1998. **50**(3): p. 191-8.
14. Nally, J.E., et al., *Lethal infection of C3H/HeJ and C3H/SCID mice with an isolate of*
Leptospira interrogans serovar copenhageni. Infect Immun, 2005. **73**(10): p. 7014-7.
15. Viriyakosol, S., et al., *Toll-like receptor 4 protects against lethal Leptospira interrogans*
serovar icterohaemorrhagiae infection and contributes to in vivo control of leptospiral
burden. Infect Immun, 2006. **74**(2): p. 887-95.
16. Richer, L., et al., *Mouse model for sublethal Leptospira interrogans infection.* Infect
Immun, 2015. **83**(12): p. 4693-700.
17. Shetty, A., S. Kundu, and M. Gomes-Solecki, *Inflammatory Signatures of Pathogenic*
and Non-Pathogenic Leptospira Infection in Susceptible C3H-HeJ Mice. Front Cell
Infect Microbiol, 2021. **11**: p. 677999.
18. Kundu, S., A. Shetty, and M. Gomes-Solecki, *Necroptosis Contributes to Persistent*
Inflammation During Acute Leptospirosis. Front Immunol, 2022. **13**: p. 810834.
19. Kundu, S., A. Shetty, and M. Gomes-Solecki, *Exposure to live saprophytic Leptospira*
before challenge with a pathogenic serovar prevents severe leptospirosis and promotes
kidney homeostasis. Elife, 2024. **13**.

20.   Werts, C., et al., *Leptospiral lipopolysaccharide activates cells through a TLR2-*
389   *dependent mechanism.* Nat Immunol, 2001. **2**(4): p. 346-52.
390   21.   Que-Gewirth, N.L., et al., *A methylated phosphate group and four amide-linked acyl*
391   *chains in leptospira interrogans lipid A. The membrane anchor of an unusual*
392   *lipopolysaccharide that activates TLR2.* J Biol Chem, 2004. **279**(24): p. 25420-9.
393   22.   Nahori, M.A., et al., *Differential TLR recognition of leptospiral lipid A and*
394   *lipopolysaccharide in murine and human cells.* J Immunol, 2005. **175**(9): p. 6022-31.
395   23.   Chassin, C., et al., *TLR4- and TLR2-mediated B cell responses control the clearance of*
396   *the bacterial pathogen, Leptospira interrogans.* J Immunol, 2009. **183**(4): p. 2669-77.
397   24.   Qureshi, S.T., et al., *Endotoxin-tolerant mice have mutations in Toll-like receptor 4*
398   *(Tlr4).* J Exp Med, 1999. **189**(4): p. 615-25.
399   25.   Nair, N. and M. Gomes-Solecki, *A Mouse Model of Sublethal Leptospirosis: Protocols*
400   *for Infection with Leptospira Through Natural Transmission Routes, for Monitoring*
401   *Clinical and Molecular Scores of Disease, and for Evaluation of the Host Immune*
402   *Response.* Curr Protoc Microbiol, 2020. **59**(1): p. e127.
403   26.   Hornsby, R.L., D.P. Alt, and J.E. Nally, *Isolation and propagation of leptospires at 37 °C*
404   *directly from the mammalian host.* Sci Rep, 2020. **10**(1): p. 9620.
405   27.   Fortes-Gabriel, E., et al., *Enzyme immunoassays (EIA) for serodiagnosis of human*
406   *leptospirosis: specific IgG3/IgG1 isotyping may further inform diagnosis of acute*
407   *disease.* PLoS Negl Trop Dis, 2022. **16**(2): p. e0010241.
408   28.   Nair, N., et al., *The route of infection with Leptospira interrogans serovar Copenhageni*
409   *affects the kinetics of bacterial dissemination and kidney colonization.* PLoS Negl Trop
410   Dis, 2020. **14**(1): p. e0007950.
411   29.   Nair, N., et al., *Role of TLR4 in Persistent Leptospira interrogans Infection: A*
412   *Comparative In Vivo Study in Mice.* Front Immunol, 2020. **11**: p. 572999.
413   30.   Vernel-Pauillac, F. and C. Werts, *Recent findings related to immune responses against*
414   *leptospirosis and novel strategies to prevent infection.* Microbes Infect, 2018. **20**(9-10):
415   p. 578-588.
416   31.   Krangvichian, P., et al., *Sublethal infection of C3H/HeNJ against Leptospira interrogans*
417   *serovar Pomona.* Acta Trop, 2023. **238**: p. 106701.
418   32.   Vernel-Pauillac, F., et al., *Anti-Leptospira immunoglobulin profiling in mice reveals*
419   *strain specific IgG and persistent IgM responses associated with virulence and renal*
420   *colonization.* PLoS Negl Trop Dis, 2021. **15**(3): p. e0008970.
421   33.   Martin, R.M., J.L. Brady, and A.M. Lew, *The need for IgG2c specific antiserum when*
422   *isotyping antibodies from C57BL/6 and NOD mice.* J Immunol Methods, 1998. **212**(2): p.
423   187-92.
424

Figure 1. Weightloss and bacterial burden in blood and urine after infection. C3H-HeJ, C3H-HeN and C57BL/6 male mice (n=4/group) were inoculated IP with 1×10^8 *L. interrogans* serovar Copenhageni strain Fiocruz L1-130 (LiC) and with PBS (control) (A). Quantification of bacterial load in blood (B) and urine (C) was determined by real-time qPCR of the 16S rRNA gene. Body weight (% change) was recorded daily for 15 days after infection (D to F). * *p*=0.0116, ****p < 0.0001 by Mann-Whitney *U* test. Data shown represents one of two experiments.

Figure 2. Survival of C3H-HeJ, C3H-HeN and C57BL/6 after infection.
Male mice (n=4/group) were inoculated IP with 1 × 10^8 *L. interrogans* serovar
Copenhageni Fiocruz L1-130 (LiC) and PBS (control), and the percentage of
mice that survived infection for 15 days was recorded. Data represents one of
two experiments.

A LiC burden in kidney tissue

B LiC in Kidney culture (EMJH) at day 4

Figure 3. Burden and viability of *L. interrogans* in kidney. C3H-HeJ, C3H-HeN and C57BL/6 male mice (n=4/group) were inoculated IP with 1×10^8 *L. interrogans* serovar Copenhageni Fiocruz L1-130 (LiC) and with PBS (control). Kidney tissues were collected 2 weeks post infection for qPCR analysis of 16S RNA from DNA purified from kidney tissue (A) and from DNA purified from kidney culture in EMJH (B). Each dot represents a single mouse. *$p < 0.05$ was determined by Mann-Whitney *U* test. Data represents one of two experiments.

Figure 4. Inflammatory cytokines and fibrosis markers in kidney of infected mice. The expression of IL-1 beta (B), iNOS (C) and ColA1 (D) in kidney tissue were measured using RT-PCR. Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) was used as the calibrator gene to normalize the expression of genes. Each dot represents measurement from a single mouse. *$p < 0.05$, **$p < 0.005$, ***$p < 0.0005$ or ****$p < 0.0001$ was determined by two-way ANOVA followed by Tukey's multiple comparison correction test. Data shown represents one of two experiments.

Figure 5. Antibody responses to *L. interrogans* infection. C3H-HeJ, C3H-HeN and C57BL/6 male mice (n=4/group) were inoculated IP with 1 × 10⁸ *L. interrogans* serovar Copenhageni Fiocruz L1-130 (LiC) and with PBS (control) (A). Serologic leptospiral-specific IgM (B), IgG (C), IgG3 (D), IgG1 (E) and IgG2a (F) was measured using ELISA. Each dot represents a single mouse. *$p < 0.05$, **$p < 0.01$, *** $p < 0.001$ or ****$p < 0.0001$ was determined by two-way ANOVA followed by Tukey's multiple comparison correction test. Data shown represents one of two experiments.