Structural Diversity of
Metabotropic Glutamate Receptor/Beta-Arrestin Coupling

Dagan C. Marx[1], Alberto J. Gonzalez-Hernandez[1*], Kevin Huynh[1*], Alexa Strauss[1,2], Carlos Rico[1], Dirk Siepe[3], Pamela Gallo[1], Joon Lee[1], Sheida Sharghi[1], Anisul Arefin[1], Johannes Broichhagen[4], David Eliezer[1,2], Marian Kalocsay[3], George Khelashvili[5,6], Joshua Levitz[1,2,7#]

[1] Department of Biochemistry, Weill Cornell Medicine, New York, NY 10065, USA
[2] Tri-Institutional Program in Chemical Biology, New York, NY 10065, USA
[3] Department of Experimental Radiation Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA
[4] Leibniz-Forschungsinstitut für Molekulare Pharmakologie (FMP), 13125 Berlin, Germany
[5] Department of Physiology and Biophysics, Weill Cornell Medicine, New York, NY 10065, USA
[6] Institute of Computational Biomedicine, Weill Cornell Medicine, New York, NY 10065, USA
[7] Department of Psychiatry, Weill Cornell Medicine, New York, NY 10065, USA

* These authors contributed equally.

Correspondence: jtl2003@med.cornell.edu

Abstract:

Despite the widespread physiological roles of beta-arrestin (β-arr) coupling in G protein-coupled receptor (GPCR) regulation, the molecular basis of GPCR/β-arr interaction has been studied primarily in monomeric family A GPCRs. Here we take an integrative biophysical and structural approach to uncover extreme molecular diversity in β-arr coupling to the neuromodulatory metabotropic glutamate receptors (mGluRs), prototypical, dimeric family C GPCRs. Using a new single molecule pulldown assay, we find that mGluRs couple to β-arrs with a 2:1 or 2:2 stoichiometry via a combination of "tail" and "core" interactions. Single molecule FRET and electron microscopy show that β-arr1 stabilizes active conformations of mGluR8 and a combination of cryo-EM structures and molecular dynamics simulations define the positioning of mGluR8-bound β-arr1, together suggesting a steric mechanism of mGluR desensitization involving interactions with both subunits and the lipid bilayer. Finally, combinatorial mutagenesis enables the identification of a landscape of homo- and hetero-dimeric mGluR/β-arr complexes, including mGluR/β-arr1/β-arr2 megacomplexes, providing a framework for family C GPCR/β-arr coupling and expanding the known range of GPCR/transducer coupling modes.

Introduction:

G protein-coupled receptors (GPCRs) are finely tuned transmembrane signaling molecules that sense extracellular stimuli and activate a wide range of intracellular effector pathways. The precise signaling of GPCRs is central to diverse physiological processes, underscoring their role as major drug targets[1, 2]. Given their biological and therapeutic significance, great effort has been undertaken to understand the regulatory processes that control GPCR function. A major form of GPCR regulation depends on the multifunctional, cytosolic β-arrestins (β-arr1, β-arr2) which bind to activated GPCRs that have been phosphorylated by G protein-coupled receptor kinases (GRKs)[3, 4]. β-arrs desensitize GPCR signaling by sterically blocking G protein access and initiating receptor internalization. β-arr mediated GPCR internalization can be subdivided into two classes: receptors that transiently recruit β-arrs to the plasma membrane but do not co-internalize with β-arrs and typically undergo endosomal recycling (class A), and receptors that form stable β-arr complexes that co-internalize into endosomes and typically undergo lysosomal degradation (class B)[5–9]. In addition to their roles in desensitization, β-arrs can serve as scaffolds to initiate secondary waves of signaling[10]. Biophysical and structural studies have shown how phosphorylated GPCR C-terminal tails mediate β-arr recruitment and activation[11–16] and how β-arrs undergo stepwise recruitment via "tail" and "core" interactions with GPCRs[5, 17–22]. However, the details of β-arr coupling can vary dramatically between GPCR subtypes[5, 23–31], between β-arr subtypes[23, 32–35], and between different ligands for the same GPCR[36–42], motivating ongoing studies of β-arrs across the vast landscape of GPCR signaling and regulation.

β-arr coupling has primarily been studied in the largely monomeric family A and family B GPCRs. Family C GPCRs, including the prototypical metabotropic glutamate receptors (mGluRs), have unique domain and quaternary structures, including large extracellular ligand binding domains (LBDs) and constitutive dimerization[43, 44], that together raise many questions about activation and coupling to G protein and β-arr transducers. For example: what is the stoichiometry of receptor/transducer complexes in homo- or heterodimeric GPCRs? What are the structural determinants of receptor/transducer coupling? How are receptor conformational dynamics altered by transducer coupling? Breakthrough studies of G protein-bound family C GPCRs have begun to provide a framework for understanding G protein-coupling in this family. Strikingly, in all structural studies a 2:1 GPCR: G protein heterotrimer stoichiometry has been observed[45–50]. Spectroscopic studies have shown that G proteins stabilize active receptor conformations[51–53], however family C GPCR/G protein structures do not show canonical signatures of transmembrane domain (TMD) activation, including outward motions of TM6, as observed in family A and B GPCRs[54–57].

While far less is known about family C GPCR/β-arr coupling, we and others have shown recently, using live cell assays, that a subset of mGluRs, including mGluR3, mGluR7, and mGluR8, undergo GRK and β-arr dependent desensitization, endocytosis, and intracellular trafficking(*58–62*). mGluR heterodimerization further diversifies this family, enabling β-arr-dependent endocytosis of otherwise-resistant subtypes (e.g. mGluR2) and altering receptor trafficking fate (e.g. endocytic recycling versus lysosomal degradation)(*59*). Differences in β-arr-dependent internalization and trafficking fates have largely been attributed to distinct patterns of phosphorylatable residues in the receptor C-termini(*58, 59*), although direct analysis of mGluR/β-arr complexes is lacking, limiting our molecular understanding of mGluR/β-arr coupling.

Given the central role of mGluRs to various modes of neuromodulation and their potential as drug targets for neurological and psychiatric disorders(*63, 64*), a more comprehensive biophysical and structural picture of mGluR coupling to β-arrs is needed. In this study, we develop a single molecule pulldown (SiMPull) imaging assay to directly assess GPCR/β-arr complex formation and characterize β-arr coupling across the mGluR family. We find variability in coupling propensity across subtypes, including the formation of both 2:1 and 2:2 mGluR/β-arr1 complexes. Mass spectrometry and mutational scanning, along with single molecule Förster resonance energy transfer (smFRET), produce a mechanistic picture of "tail" and "core" interactions which drive mGluR/β-arr coupling. Negative stain electron microscopy (EM), cryo-EM, and molecular dynamics simulations delineate mGluR8/β-arr1 interaction modes and define β-arr1 positioning in 2:1 core-bound complexes. Finally, structure-guided SiMPull analysis reveal a range of 2:1 and 2:2 mGluR8/β-arr binding modes, including formation of heterodimeric mGluR8/mGluR2/β-arr1 complexes and mGluR8/β-arr1/β-arr2 megacomplexes, greatly expanding the diversity of known GPCR/β-arr assemblies.

Results:

A single molecule assay to assess GPCR/β-arr1 complex formation and stoichiometry

To directly detect and quantitate mGluR/β-arr complexes, we first aimed to establish a single molecule imaging platform based on visualizing constructs bearing self-labeling tags (e.g. SNAP, Halo) for fluorophore conjugation. Live cell imaging controls confirmed that SNAP-mGluR3 and SNAP-mGluR8 recruit β-arr1-Halo upon glutamate addition but that only SNAP-mGluR8 co-internalizes with β-arr1-Halo (**Fig. S1A-D**), in agreement with our prior studies (*59*). We prepared a C-terminally truncated β-arr1 construct ("β-arr1-ΔCTD-Halo") where the autoinhibitory C-terminal domain (aa 383-418) has been

removed. This construct was efficiently recruited to the plasma membrane by both mGluR3 and mGluR8 (**Fig. S1E, F**) and impaired internalization of both receptor subtypes, consistent with the removal of motifs which facilitate clathrin binding within the β-arr1-CTD, (**Fig. S1G, H**)(*65–69*).

To assess the ability of mGluRs to couple to β-arr1 via single molecule pulldown ("SiMPull")(*70, 71*), fresh lysates were prepared from cells transfected with HA-tagged SNAP-mGluRs and Halo-tagged β-arr1 constructs that were labeled with fluorophores (cell impermeable SBG-JF$_{646}$ for SNAP; cell permeable CA-JF$_{549}$ for Halo) and then incubated with saturating concentrations of glutamate. Receptors were isolated on passivated glass coverslips via an anti-HA primary antibody (**Fig. 1A**) and total internal reflection fluorescence (TIRF) microscopy was used to visualize individual receptors and β-arr1 in separate channels and quantify complex formation via colocalization of spots (**Fig. 1B**). In the absence of GRK overexpression, minimal complex formation (<10%) with either WT or β-arr1-ΔCTD was observed for mGluR8, while mGluR3 showed modest colocalization with β-arr1-ΔCTD (~25%) (**Fig. S2A-F**). Co-transfection with either GRK2, GRK5, or a membrane-tethered GRK2 containing a C-terminal CaaX-box motif (GRK2-CaaX), dramatically increased the extent of β-arr1 colocalization for both mGluR3 and mGluR8 (**Fig. S2B, D)**, likely by enhancing the degree of receptor phosphorylation. Both mGluR3 and mGluR8 produced substantial pulldown of β-arr1-ΔCTD with ~60% or 75% of receptors, respectively, showing β-arr1 colocalization under optimal conditions with GRK2-CaaX co-transfected (**Fig. 1B, C**). A striking difference between these two mGluR subtypes was that only mGluR8 showed substantial co-localization with WT β-arr1, across all GRK conditions (**Fig. 1C; Fig. S2E, F**). Importantly, in the absence of receptor co-expression very minimal β-arr1-Halo pulldown was observed (**Fig. S2G**). Overall, these results suggest that mGluR8 has a higher phosphorylation threshold for β-arr1 coupling but can form more stable complexes with β-arr1 than mGluR3, which may explain the ability of mGluR8/β-arr1 complexes to persist following endocytosis.

To assess the generalizability of our assay, we performed the same measurements with SNAP-tagged beta-2 adrenergic receptor (β2AR) or arginine vasopressin 2 receptor (V2R), prototypical family A GPCRs that either interact transiently(*6, 72*) or co-internalize(*6*) with β-arr1, respectively (**Fig. S3A-D**). In the presence of GRK2-CaaX and saturating agonist, each receptor showed pulldown of β-arr1-ΔCTD, but V2R was substantially more efficient (~60% vs. 20% colocalization). Similarly to mGluR8 and mGluR3, respectively, V2R was able to produce strong pulldown of WT β-arr1 while β2AR was not (**Fig. 1C**). These results are consistent with class B (V2R, mGluR8), but not class A (β2AR, mGluR3), GPCRs being capable of stable interactions with WT β-arr1 and indicate that our SiMPull assay can provide a general means of quantitatively probing and classifying GPCR/β-arr interactions.

We next extended our analysis across the mGluR family where we have previously observed variability across subtypes in terms of ligand-induced β-arr desensitization and endocytosis(*58*, *59*). We tested SNAP-tagged mGluR2, mGluR5, and mGluR7 in the presence of GRK2-CaaX and either β-arr1-Halo or β-arr1-ΔCTD-Halo (**Fig. 1B, C**). SNAP-mGluR2 showed negligible levels (<5%) of co-localization with either β-arr1 or β-arr1-ΔCTD (**Fig. 1B, C**), consistent with the lack of β-arr1 recruitment and receptor internalization observed in live cells (**Fig. S3E, F**)(*58*, *59*). In contrast, SNAP-mGluR5 showed modest pulldown (~15-20%) of β-arr1-ΔCTD, but not WT β-arr1 (**Fig. 1C**), that was mirrored in live cell imaging where β-arr1-ΔCTD, but not WT β-arr1, showed detectable glutamate-induced membrane localization (**Fig. S3G, H**). Finally, SNAP-mGluR7 showed weak (~5-10%) pulldown of both β-arr1-ΔCTD, and WT β-arr (**Fig. 1C**), in line with live cell imaging showing modest recruitment of β-arr1-ΔCTD, but not WT β-arr1 (**Fig. S3I, J**).

A major advantage of single molecule imaging is the ability to measure the stoichiometry of protein assemblies via photobleaching step analysis(*73–75*). We asked two key questions about mGluR/β-arr1 complexes: 1) Do mGluRs retain dimerization upon β-arr interaction? And 2) how many β-arr1 molecules can interact with an mGluR at a time? We analyzed the photobleaching step distribution for both the receptor and β-arr1 channels for all β-arr-colocalized mGluR3, mGluR5, and mGluR8 spots (**Fig. 1D-F**). We found that ~50% of spots displayed 2 bleaching steps in the receptor channel for each mGluR subtype (**Fig. 1E, F**), agreeing with recent studies(*59*, *73*) and confirming that mGluRs remain dimeric upon β-arr interaction. As previously established, the subpopulation of mGluRs showing bleaching steps is a function of incomplete fluorophore labeling, incidental bleaching, and other detection issues. Surprisingly, ~25-55% of co-localized complexes also showed 2 bleaching steps in the β-arr1 channel (**Fig. 1D-F**), indicating that a combination of 2:1 and 2:2 mGluR:β-arr1 complexes can form. The proportion of 2-step β-arr1 bleaching steps was lower for mGluR5 than mGluR3 or mGluR8 which may be due to lower overall complex formation for this receptor. As a key control, V2R:β-arr1-ΔCTD complexes showed background levels (~10%) of 2-step bleaching in both channels (**Fig. 1F; Fig. S4A, B**) consistent with formation of a 1:1 complex between monomeric family A GPCRs and a single β-arr1 molecule. To further validate our finding of 2:2 mGluR:β-arr1 complexes we sought to directly image β-arr1 colocalization at a single receptor. To do this, we labeled β-arr1-Halo with a mix of both CA-JF$_{549}$ and CA-JF$_{646}$ fluorophores, while leaving the SNAP-tag on mGluR8 unlabeled (**Fig. S4C**). We observed substantial colocalization (~40%) of β-arr1-ΔCTD when imaging both channels (**Fig. S4D**), in agreement with our photobleaching step analysis implying the existence of mGluR/β-arr1 complexes containing two β-arr1 molecules bound to one receptor dimer.

Mechanistic analysis of mGluR8/β-arr1 coupling

Having observed particularly strong formation and stability of mGluR8/β-arr1 complexes, we decided to focus our analyses on this interaction as a means of illuminating the nature of mGluR/β-arr coupling. Our prior work has pointed to a critical role of phosphorylation sites in the mGluR8 C-terminus(*59*) and mass spectrometry of agonist-treated mGluR8 showed extensive phosphorylation in the distal serine/threonine (S/T)-rich region (aa888-907) of mGluR8, including many double-phosphorylated peptides (**Fig. S5**), consistent with the likelihood that multiple phosphorylation sites simultaneously contribute to β-arr recruitment(*11*, *13*, *76*, *77*).

We performed alanine scanning mutagenesis of the 11 potential phospho-sites in the mGluR8 S/T-rich region to investigate the role of each in β-arr1 complex formation via SiMPull. We tested each mutant using our highest colocalization condition (GRK2-CaaX, β-arr1-ΔCTD-Halo) and a less tightly coupled condition (GRK2, β-arr1-Halo), finding similar trends in both conditions with larger effects for the latter (**Fig. 2A**). All mutants showed impairment with the largest effects seen for T896A, T898A, and T899A, residues that could constitute a recently identified arrestin binding PxPP motif(*13*, *78*). Importantly, we found no pulldown with simultaneous mutation of all 11 site ("mGluR8-11xA"), highlighting the need for phosphorylation in the mGluR8 S/T-rich region for β-arr1 coupling. We then assessed the effects of each alanine variant using an established live cell imaging-based assay of glutamate-induced mGluR8 internalization. We found clear impairment of receptor internalization for most mutations (**Fig. S6A, B**), including a complete loss of internalization for mGluR8-11xA.

A clear correlation across mutations was observed between the extent of impairment of β-arr1 colocalization and the extent of impairment of ligand-induced internalization (**Fig. 2B**). However, three mutations fell outside of the tight linear range. S888A showed a weaker effect in SiMPull compared to internalization while T896A and T899A showed a larger effect in SiMPull compared to internalization (**Fig. 2B**). To better understand the discrepancy between our two assays for these three variants, we used live cell imaging of β-arr1 recruitment and receptor endocytosis. The mGluR8 S888A variant revealed an almost complete loss of β-arr1 recruitment and receptor internalization (**Fig. S6C**). One possible explanation is that S888 serves as a gatekeeper whose phosphorylation helps drive subsequent phosphorylation of other positions. This could explain why SiMPull effects under conditions where GRK2-CAAX is overexpressed are less pronounced. In contrast, T899A showed clear receptor internalization and β-arr1 recruitment to the plasma membrane, but minimal co-localization of β-arr1 with internalized receptors (**Fig. S6F**). A potential explanation is that phosphorylation of T899 is critical for maintaining a conformation of β-arr1 that survives endocytosis but is not required for initial β-arr1 recruitment to the plasma membrane. T896A showed impaired recruitment of β-arr1, but still maintained some weak co-localization in endosomes (**Fig. S6G**).

These results highlight the combined power of single molecule and live cell fluorescence approaches to investigate the complex roles of phospho-sites on GPCR-β-arr complex formation and trafficking. The mutational effects seen across a 19-residue region of the CTD, suggest the potential of multiple β-arr1 binding modes.

We next asked if the mGluR8 TMD core contributes to mGluR8/β-arr1 coupling. We turned to a well-established mutation of a highly conserved phenylalanine (F777 in mGluR8) in intracellular loop 3 that strongly impairs G protein coupling across mGluR subtypes(*75*, *79*, *80*). We hypothesized that, if a similar interface is used for β-arr1 core coupling, mutation to this position would also strongly impair β-arr1 coupling and mGluR8 endocytosis as previously shown for mGluR3(*58*). We found that colocalization of β-arr1 with SNAP-mGluR8-F777D was strongly reduced via SiMPull and glutamate-driven internalization in live cells was abolished (**Fig. 2C; Fig. S7A, B**). While such effects could be due, in part, to impairment of GRK coupling, this result argues that TMD core interactions contribute to mGluR8/β-arr1 coupling. To further test this hypothesis, we asked if receptor core interactions alone are sufficient to produce detectable mGluR8/β-arr1 coupling in cells by producing a mGluR8 construct lacking a CTD (mGluR8-ΔCTD). In line with a role for core coupling, SNAP-mGluR8-ΔCTD showed weak, but clear, internalization (**Fig. S7C**).

As we hypothesize that mGluR8/β-arr1 coupling involves interactions with the TMD core, we asked if β-arr1 coupling alters the conformational dynamics of mGluR8. Prior work using FRET-based assays and cryo-EM has established that G proteins can stabilize mGluRs in their active-like conformation (*45–47*, *51–53*), but whether β-arrs exert a similar or, potentially, opposite effect as part of an inhibitory mechanism, is unknown. We tested this using an inter-LBD single molecule (smFRET) assay that has been used to characterize agonist-induced transitions from inactive, high FRET, to active, low FRET, conformations(*51*, *53*, *81*, *82*) (**Fig. 2D**). As expected, application of saturating (1 mM) glutamate shifted FRET histogram peaks of SNAP-mGluR8 immobilized from lysates of HEK293T cells from a center of ~0.35 to a center of ~0.15 but with a clear shoulder (**Fig. 2E**) indicating incomplete occupancy of the active, low-FRET state. In both the absence (**Fig. S8A**) and presence of glutamate (**Fig. 2D**), a sub-population of molecules showed dynamic transitions between FRET states. This agrees with prior smFRET studies of group II/III mGluRs which showed variable degrees of incomplete occupancy of the low FRET state in response to orthosteric agonists(*81–83*). When SNAP-mGluR8 lysates were prepared from cells co-expressing GRK2 and β-arr1-ΔCTD, the occupancy of the low FRET state in the presence of glutamate was substantially enhanced (**Fig. 2E**). This was accompanied by a reduction in the percentage of molecules showing dynamic transitions (**Fig. 2D; Fig. S8B**), suggesting that β-arr1 helps to "lock" mGluR8 in an active conformation. Hidden Markov modeling of high (0.35), medium (0.25), and low (0.15) FRET states

revealed an increase in the dwell time in the low FRET state, but not the medium or high FRET states (**Fig. S8C**), in the presence of β-arr1, likely owing to decreased transitions between the low FRET active state and medium FRET state (**Fig. S8D**). Together these data show that β-arr1 promotes an active mGluR8 conformation by reducing transitions to inactive states (**Fig. 2F**), suggesting that β-arr1 recognizes a similar active state to G protein. Recent work has shown similar conformational effects of positive allosteric modulators (PAMs) (*36*, *53*, *84–86*), indicating that they likely stabilize a similar active state or active states as β-arr1. Consistent with this, PAMs enhance both agonist-driven G protein and β-arr coupling for mGluR2, mGluR3, mGluR7, and mGluR8(*36*).

Structural Analysis of mGluR8/β-arr1 Complexes

To gain further structural insight into mGluR8/β-arr1 complexes, we first purified either full-length mGluR8 alone or mGluR8/β-arr1ΔCTD complexes in the presence of agonist (L-AP-4) and PAM (VU6005649; "VU600") and performed negative stain EM. The mGluR8 alone sample yielded 2-D classes showing dimeric mGluR8 with clearly resolved LBDs and TMD-containing detergent micelles (**Fig. 3A**). The mGluR8/β-arr1 complex sample contained particles with additional density on the intracellular side of the micelle consistent with complex formation (**Fig. S9**). We identified both "core" and "tail" bound binding modes based on the orientation of the elongated β-arr1 density (**Fig. 3B, C; Fig. S9**), with the core-bound β-arr1 having a position that is consistent with simultaneous interactions with both mGluR8 TMDs. Given our SiMPull evidence for 2:2 mGluR8/β-arr1 complexes, we manually inspected our micrographs and observed a subset of particles containing two separable densities (**Fig. S9**). 2-D classes generated from these particles revealed intracellular density that was larger than calculated for a single β-arr1 molecule and appeared to contain two distinct molecules with different orientations (**Fig. 3D**).

We next turned to cryo-EM to gain deeper insight into the structural basis of mGluR8/β-arr1 coupling. We first determined structures of pre-active mGluR8 bound to L-AP-4 and VU600 (**Fig. 3E; Fig. S10**). Image processing revealed two classes with similar numbers of particles which each showed closed LBDs (**Fig. S11A**) with clear density for L-AP-4 in the inter-domain cleft of the LBDs (**Fig. S11B-D**) and active-like inter-LBD and inter-CRD orientations (**Fig. S11E, F**). The major difference between class 1 and class 2 structures is a lateral offset of the TMD dimer relative to the LBDs (**Fig. S11G**). Both classes show a TM6-TM6 inter-TMD interface (**Fig. S11H**), consistent with prior agonist- and PAM-bound full-length mGluR structures(*36*, *45–47*, *86–89*) (**Fig. S11H, I**). We observed density that is consistent with VU600 at both sides of the interface, suggesting a 2:2 binding stoichiometry (**Fig. S11J,K**). This PAM binding site is similar to a prior mGluR4/G protein structure although in that model only a single PAM per dimer is observed(*47*) (**Fig. S11L**).

We next sought to structurally characterize mGluR8/β-arr1ΔCTD complexes using cryo-EM. Image processing resulted in two major classes with comparable inter-domain orientations and to the classes described above for mGluR8 alone (**Fig. S12; Fig S13A-C**). In both classes, an arrestin-shaped density is observed on the intracellular side of the mGluR8 TMD dimer that is similar to the "core" β-arr1 density observed in our negative stain EM 2D classes (**Fig. 3F; Fig. S13D**). AlphaFold2 prediction of the mGluR8/β-arr1 complex yielded a core-bound β-arr1 orientation that fit the β-arr1 density in our maps to guide an atomistic model of the mGluR8/β-arr1 complex via rigid body docking (**Fig. 3G, H; Fig. S13E**). Importantly, our cryo-EM maps are inconsistent with an alternative model in which β-arr1 has a 180° flipped orientation such that the β-arr1 C-edge is pointing away from the dimer interface (**Fig. S13F**) and our model shows good agreement with negative stain 2-DCs (**Fig. S13G**). Subtle differences were observed in the mGluR8 TMD when bound to β-arr1, including repositioning of the intracellular ends of TM3, TM5, and TM6 and increased resolution of intracellular loops and the initial stretch of the CTD (**Fig. 3I; Fig. S13H, I**). These differences are reminiscent of those observed upon G protein-binding in prior structures of mGluR2 and mGluR4(*45*, *46*) and further support β-arr1 interactions with the mGluR8 TMD core. Notably, these conformational changes do not include the outward motion of TM6 which is canonical for activated family A and B GPCRs(*90*) but has not been seen in family C GPCRs.

The positioning of β-arr1 relative to the homologous seven-helix TMD of various GPCRs shows a range of orientations across reported structures with mGluR8 displaying a unique x-y orientation in the plane parallel to the TMDs that is 90-150º shifted from that of the CB1R, β1AR, 5-HT2BR, and V2R and ~180º opposite of that seen for the NTS1R (**Fig. 3J**). Variability has also been observed in the orientation of β-arr1 relative to the plane of the membrane with some receptors (e.g. NTS1R, V2R) showing a sharp tilt z angle of <70º that may enable interaction between the β-arr1 C-edge and the membrane (**Fig. S13J**). In contrast, in our mGluR8 structural model, the β-arr1 C-edge is largely blocked from accessing the membrane by the other mGluR8 TMD and results in β-arr1 having a z angle of 82.7º (**Fig. 3G**).

To better understand the molecular details of the mGluR8/β-arr1 interaction, we performed molecular dynamics simulations using our cryo-EM based full-length structural model (see Methods). In total, we ran 72 independent replicas of the mGluR8/β-arr1 complex embedded in a model plasma membrane, resulting in over 19 μs of total simulation time. Across all simulations the mGluR8 domains remain in an active-like conformation (**Fig. 3H; Fig. S14A, B**) containing closed-closed/active LBD dimers (**Fig. S14C**) and a TM6-TM6 inter-TMD interface (**Fig. S14A, C, D**). However, we observed a tilting motion of the LBD relative to the TMD (**Fig. 3H**), revealing large-scale conformational flexibility of mGluR ECDs and supporting the two classes of cryo-EM densities reported above.

β-arr1 remained stably bound to the TMD core of mGluR8 in a range of active-like conformations (**Fig. S15A, B**) throughout the simulations but showed positional dynamics in both X-Y and Z planes relative to the mGluR8 TMDs (**Fig. 3H**; **Fig. S15C-H**). This conformational heterogeneity is consistent with the limited resolution of the β-arr1 density in our cryo-EM maps. Importantly, the consensus conformation captured in our cryo-EM model, in terms of the x-y and z-angle represents the dominant orientation of the complex (**Fig. 3K**). We observed a minor subset of simulations (13/72) where the β-arr1 C-edge falls away from the TMD, resulting in a subpopulation of simulation frames where the Z-angle is >100° (**Fig. S15F-H**).

Our structural model and MD simulations of a 2:1 core-bound mGluR8/β-arr1 reveal a primary interaction where the orientation of β-arr1 relative to mGluR8 is such that the cytoplasmic surface of both TMDs is largely occluded (**Fig. 4A, B**). The data suggest a simple steric mechanism for β-arr-mediated desensitization that would provide an efficient mechanism for preventing G protein binding to the receptor dimer, effectively preventing canonical signal initiation. In addition, this orientation would preclude the binding of a second β-arr molecule to the TMD dimer core without either a large rearrangement of the β-arr or a major inter-subunit rotation of the TMD.

A closer analysis of interactions between β-arr1 and the mGluR8 TMD in our model and simulations revealed interactions between the β-arr1 finger loop and residues in ICL1, ICL2, and ICL3 of the receptor chain A (**Fig. 4C; Fig. S16C-E**). The finger loop binding site is comprised primarily of intracellular loops, comparable to what has been observed for the α5 helix of Gα in G protein-bound mGluR structures (**Fig. S16A, B**). Consistent with coupling via these interactions, we observed reduced structural variability in this TMD subunit (chain A) in our simulations (**Fig. S14B**) and a consistent but dynamic interface between the finger loop and mGluR8 (**Fig. S16C, D**). We also observed interactions between the β-arr1 middle and C-loops and mGluR8 chain A TMD (**Fig. S16F**), which may contribute to the stability of the core mGluR8/β-arr1 interaction. Finally, in trajectories with large β-arr1 z angles, we observed that interactions between residues in the β-arr1 N-lobe and the membrane may stabilize this state in a lipid-dependent manner (**Fig S16G, H**).

Owing to the orientation of β-arr1 relative to the mGluR8 dimer, the β-arr1 C-edge comes into proximity with both ICL1 and ICL2 of the mGluR8 chain B TMD and the plasma membrane in many of our simulations, (**Fig. 4D**; **Fig. S16G-J**). ICL1/C-edge interactions are mediated electrostatically and often recruit a negatively charged lipid headgroup (**Fig. 4D**, left) while ICL2 interactions are formed via a long, hydrophobic interface (**Fig. 4D**, right). We also observe C-edge loops embedding into the membrane in some replicas without apparent interactions with mGluR8 (**Fig. S16H)** and with specific lipid preferences (**Fig. S16J**). This observation is in line with recent biophysical studies showing β-arr C-edge/membrane

interactions(*28*, *91*) and further highlights the apparent configurational diversity exhibited by β-arr1 while maintaining the canonical finger loop-based interaction with mGluR8.

To experimentally probe the mGluR8 TMD/β-arr1 core interaction observed in our integrative structural approach, we individually mutated many putative interfacial residues in ICL1, ICL2, and ICL3. Using SiMPull, we found impaired β-arr1-ΔCTD-Halo colocalization with mGluR8 mutants across all loops with the largest effects for F677A and F777A in ICL2 and ICL3, respectively (**Fig. 4E**). Interestingly, the melanoma-associated mutation V773D in ICL3(*92*) also substantially impaired interaction with β-arr1 (**Fig. 4E**). Similar effects of the mutations were observed using our surface labeling internalization assay (**Fig. S17A, B**) with a clear correlation between the two assays (**Fig. 4F**). Results were most variable and subtle for mutations in ICL1, suggesting that these interactions are secondary in importance compared to the primary interactions with ICL2 and ICL3, in line with what has been previously observed in mutational analysis of mGluR/G protein coupling(*46*).

cis and *trans* mGluR/β-arr1 coupling

The involvement of both tail and core interactions in mGluR8/β-arr1 coupling raises the question of whether coupling occurs in *cis*, via interactions of β-arr1 with the tail and TMD core of the same subunit, or in *trans*, via interactions of β-arr1 with the tail of one subunit and the TMD core of the other. Notably, the inter-TM6 interface observed in our structure brings the CTD (e.g. bottom of TM7) to a central site within the receptor dimer, positioning it such that it is conceivable that coupling to either TMD core can occur. To test this, we took advantage of mutations that strongly impair either tail interaction ("11xA") or core interaction (F777D) to prevent mGluR8/β-arr1 co-internalization (**Fig. 5A; Fig. S18A**). We co-expressed SNAP-tagged mGluR8-F777D with untagged mGluR8-11xA and observed clear internalization both in cell imaging and surface labeling internalization assays (**Fig. 5A; Fig. S18A**), supporting the ability of *trans* interactions to mediate β-arr-mediated internalization. Similar results were observed upon co-expression of SNAP-mGluR8-11xA with untagged mGluR8-F777D (**Fig. 5A; Fig. S18A**). Supporting *cis* interactions, co-expression with wild type untagged mGluR8 enabled internalization of SNAP-mGluR8-F777D-11xA (**Fig. 5A; Fig. S18A**). To directly compare the propensities of *cis* and trans mGluR8/β-arr1interactions, we turned to SiMPull. To detect *trans* interactions, we tested the ability of HA-tagged mGluR8-11xA to pulldown with β-arr1-Halo upon co-expression with SNAP-mGluR8-F777D and observed substantial colocalization (**Fig. 18B**). To detect *cis* interactions, we tested the ability of an HA-mGluR8-F777D-11xA construct to pulldown β-arr1-Halo upon co-expression with wild type SNAP-mGluR8 (**Fig. 18C**). In both cases we observed substantial colocalization but with enhanced efficiency in cis (**Fig. 18D**), suggesting a preference for this mode of tail and core interactions.

The ability of β-arr1 to couple in *cis* or *trans* may be particularly relevant for heterodimeric mGluRs which introduce asymmetry into the cytoplasmic face of the receptor and may be regulated in complex ways by subunit-specific pharmacology or post-translational modifications. We focused on mGluR2/8 heterodimers as these combine a strong β-arr coupled mGluR (mGluR8) with a β-arr resistant mGluR (mGluR2) and we have recently shown that these heterodimers undergo β-arr dependent endocytosis and lysosomal degradation(*59*). We first confirmed that co-expression with untagged mGluR8 enabled internalization of SNAP-mGluR2 (**Fig. 5B; Fig. S18E**). Introducing the 11xA mutation into mGluR8 abolished mGluR2 internalization but introducing the F777D mutation into mGluR8 only partially impaired mGluR2 internalization (**Fig. 5B; Fig. S18E**), suggesting that while mGluR8 tail coupling is strictly required, mGluR8 core coupling is not. Internalization was fully abolished when the equivalent mutation, F756D, was also introduced into mGluR2 (**Fig. 5B; Fig. S18E**), suggesting that both *cis* and *trans* mGluR2/8 β-arr coupling can occur via the mGluR8 tail and either the mGluR2 or mGluR8 core. Using SiMPull we found that co-expression of SNAP-mGluR8 and HA-SNAP-mGluR2 enables colocalization with β-arr1-Halo (**Fig. 5C**). Importantly, the vast majority of colocalized β-arr1-Halo showed single-step photobleaching (**Fig. 5C; Fig. S18F**), consistent with a 1:1:1 mGluR2: mGluR8: β-arr1 complex. Finally, we observed a decrease in β-arr1-Halo colocalization when we introduced the F756D mutation into mGluR2, further supporting both *cis* and *trans* β-arr1 coupling to mGluR2/8 heterodimers (**Fig S18G**).

mGluR8 coupling with β-arr2 and mGluR8/β-arr1/β-arr2 megacomplexes

While β-arr1 has most often served as the model β-arr subtype for structural and biophysical studies, β-arr2 also strongly contributes to GPCR regulation and shows structural and functional differences compared to β-arr1(*23*, *32–35*), motivating us to examine mGluR8/β-arr2 coupling (**Fig. S19A, B**). Using SiMPull, we observed clear pulldown of β-arr2-Halo by both SNAP-mGluR8 and V2R (**Fig. 5D, E; S19C-E**), although the efficiency was reduced in comparison to β-arr1. mGluR8-mediated pulldown of β-arr2 was dramatically reduced with the 11xA or F777D mutations (**Fig. S19E**), suggesting contributions from both tail and core interactions. Photobleaching step analysis showed both 2:1 and 2:2 mGluR8/β-arr2 complexes (**Fig. 5E; Fig. S19F**) indicating that the same range of stoichiometries is observed for complexes of mGluR8 with β-arr1 or β-arr2. We further confirmed this by observing substantial colocalization between β-arr2-Halo molecules when labeled with two different colors upon pulldown by HA-mGluR8 (**Fig. S19G**).

We then asked if a single mGluR8 dimer can form 2:1:1 megacomplexes with one molecule each of β-arr1 and β-arr2. We first analyzed cortical single cell RNA sequencing data from the mouse anterior cingulate cortex(*93*) and found that 37.8% of neurons showed co-expression of β-arr1 and β-arr2 (**Fig.

S20A-C). Notably, a wide range of RNA ratios between mGluR8, β-arr1, and β-arr2 is seen across neurons with a large proportion (~24% of the total and ~40% of mGluR8-expressing neurons) expressing all three (**Fig. S20C, D**), raising the possibility that β-arr1 and β-arr2 can either compete for the same GPCR or bind in a combinatorial fashion. Accordingly, live cell imaging revealed endosomal co-localization of CLIP-mGluR8, β-arr1-SNAP, and β-arr2-Halo (**Fig. S20E**). To address this directly, we performed SiMPull with HA-mGluR8 and found a large proportion of spots with colocalized β-arr1-SNAP and β-arr2-Halo (**Fig. 5F-H**). β-arr1-SNAP and β-arr2-Halo channels showed a single bleaching step in each channel, consistent with a 2:1:1 complex (**Fig. 5G**). In contrast, negligible background levels of β-arr1-SNAP and β-arr2-Halo colocalization were observed with pulldown of V2R (**Fig. 5H; Fig. S20F**). This result represents, to our knowledge, the first example of two different arrestin subtypes simultaneously interacting with a single GPCR, further expanding the molecular diversity of mGluR/arrestin coupling.

Discussion:

Our integrative biophysical approach reveals major complexity in mGluR/β-arr coupling, suggesting a high degree of tuneability and flexibility to match the myriad roles of mGluRs in regulating synaptic transmission and plasticity(*63*, *64*, *94*). The ability of mGluRs to couple to either one or two copies of either β-arr1 or β-arr2 in tail or *cis* and *trans* core-binding modes may serve to impart different extents or speeds of desensitization, endocytosis, intracellular trafficking, and/or arrestin-dependent signaling. Combined with the ability of mGluR heterodimers to couple to β-arrs in multiple configurations and the possibilities of mGluR/β-arr1/β-arr2 megacomplex, we propose 20 distinct modes of mGluR/β-arr coupling, contrasting with the limited repertoire of 1:1 family A GPCR/ β-arr coupling (**Fig. 5I**). It is likely that the specific mode of β-arr binding that predominates in each setting depends on the degree or timing of mGluR activation by synaptic glutamate or exogenous ligands and the given cellular context in terms of the β-arr1 and β-arr2 expression levels, subcellular localization, cell morphology, and the presence of other regulatory proteins (e.g. GRKs).

Our study supports an increasing appreciation of the functional specialization of mGluR subtypes. Our data correlate well with prior studies that found that mGluR2 is resistant, while mGluR3 undergoes transient β-arr coupling on the cell surface followed by recycling, and mGluR8 undergoes co-internalization followed by lysosomal degradation(*59*). Our observations that mGluR8 has a steeper dependence on GRK expression (**Fig. S3**) but shows a higher complex stability than mGluR3 (**Fig. 1**), suggest that mGluR3 is specialized for rapid, reversible desensitization while mGluR8 is optimized for longer-lasting downregulation after elevated or extended degrees of activation, potentially allowing each receptor to

regulate distinct forms of synaptic plasticity and metaplasticity. Interestingly, mGluR5 showed complex formation with β-arr1-ΔCTD in SiMPull despite evidence that it undergoes β-arr-independent endocytosis(*58*, *95–97*). This suggests that relatively weak β-arr coupling may play non-trafficking roles in tuning or initiating mGluR5 signaling, in line with studies showing an impairment of mGluR5-mediated long-term depression following β-arr2 knockout(*98*). In contrast, mGluR7 showed weak complex formation with β-arr1 in SiMPull despite evidence that it undergoes β-arr-dependent internalization and degradation in neurons(*60*). This prior work revealed a strong dependence on ubiquitination of mGluR7 for its degradation and, together with our work, suggests that β-arr-dependent GPCR degradation may not strictly require the formation of a tight complex with β-arrs. The distinct modes of β-arr coupling in terms of complex stability and intracellular trafficking uncovered here suggest distinct molecular and/or structural mechanisms across subtypes.

Cryo-EM analysis of mGluR8 allowed us to visualize and model the orientation of β-arr1 on agonist and PAM-bound, full-length, human, wild-type mGluR8, providing a first glimpse into a family C GPCR/β-arr complex. The unique orientation of β-arr1 relative to the TMD of mGluR8 is not well-aligned with existing GPCR/β-arr1 structures, extending our appreciation of the flexibility/dynamics involved in β-arr coupling to GPCRs. Intriguingly, the single core-bound β-arr1 seen with mGluR8 is well-positioned to both occlude G protein binding to the subunit that it directly interacts and to the adjacent subunit via positioning of its C-lobe. The active conformation observed for mGluR8 in the complex structure, with modest repositioning of intracellular loops and the maintenance of a TM6-TM6 interface, supports our smFRET analysis which revealed mGluR8 active-state stabilization by β-arr1. These findings suggest that, analogously to family A GPCRs that have been shown to be stabilized in active states by β-arrs (*28*, *99*, *100*), both G proteins and β-arrs can couple to and stabilize similar active conformations in mGluRs.

The molecular diversity of mGluR/β-arr coupling modes uncovered in this study motivates further structural analyses across mGluR and β-arr subtypes. Recent studies have shown that β-arr2 can adopt distinct family A GPCR binding modes(*23*, *32*), raising the question of how it couples to mGluRs and how mGluR/β-arr1/β-arr2 megacomplexes may be assembled. Intriguingly, our findings raise the possibility of megacomplexes between an mGluR, β-arrs, and G proteins, as variations of this have been seen in a family A GPCRs and may contribute to endosomal signaling(*101–104*). Notably, the core-bound model of mGluR8/β-arr1 we observed here would be incompatible with simultaneous binding of β-arr1 and a G protein heterotrimer without major reconfiguration. However, G protein binding may be compatible with tail-bound β-arr or with alternative conformations, such as the high β-arr1 z-angle binding mode observed in our MD simulations. As we have demonstrated mGluR and GRK subtype dependence on β-arr coupling, further work is also required to understand the structural basis of GRK/mGluR interaction. Finally, the

CTDs of mGluRs are known to interact with myriad other proteins at the synapse, including calmodulin, Munc18, PICK1, and Homer(*63, 105–109*), raising the question of how such interactions may prevent or potentiate β-arr binding and further sculpt mGluR regulation.

Methods:

Molecular Biology

All SNAP- or CLIP-tagged mGluR2 (rat), mGluR3 (rat), mGluR5 (human), mGluR7 (rat), mGluR8 (human), V2R (human), β2AR (human) were generated with an N-terminal signal peptide from rat mGluR5 followed by variable combinations of HA-, SNAP-, FLAG-, or CLIP in the pRK5 vector as previously described(*58*). Untagged bovine GRK2 (#14691), GRK2-CaaX (#166224), and GRK5 (#14690), were purchased from Addgene. Point mutations were introduced using PCR-based site-directed mutagenesis. Rat β-Arr1–Halo, β-Arr1–SNAP, and β-Arr2–Halo were generated by replacing YFP from β-arr1–YFP (Addgene) or β-arr2–YFP (Addgene, #36917) with the Halo or SNAP tag using a Gibson Assembly Cloning Kit. β-arr1-ΔCTD-Halo constructs were created by removing residues 383-418 from the WT β-Arr1–Halo plasmid using In-Fusion cloning (Takara Bio). mGluR8-ΔCTD was created by introducing a STOP codon in position 845 by PCR-based site-directed mutagenesis. For cryo-EM experiments, cDNA for human mGluR8 was cloned into the pEZT-BM BacMam expression vector (*110*) and the N-terminus was fused to an HA and FLAG (DYKDDDD) tag followed by a 3-alanine linker (AAA).

Cell culture and transfection

HEK 293T (ATCC: CRL-3216) cells were grown and maintained in Dulbecco's Modified Eagle's Medium (Corning) supplemented with 10% fetal bovine serum (Thermo Fisher) and cultured at 37ºC/5% CO_2. For live cell imaging and single molecule imaging experiments, cells were seeded on poly-L-lysine (Sigma) coated 18 mm glass coverslips in 12-well plates and transfected using Lipofectamine 2000 (Thermo Fisher Scientific). For cell imaging experiments (e.g. Fig. 1A, B), cells were transfected with 700 ng each of receptor, β-arr, and GRK2 per well. SiMPull experiments also used 700 ng of each plasmid per well. For some SiMPull experiments, GRK2 was replaced with GRK2-CaaX or GRK5 or omitted. For surface labeling assay experiments (e.g. Fig. S3D), only receptor was transfected. For experiments with expression of two receptor constructs, a 1:1 ratio was maintained with 700 ng of each receptor. For β-arr co-expression experiments a 1:2 ratio of β-arr1: β-arr2 was used. Six hours after transfection with mGluRs, media was replaced and supplemented with the antagonist LY341495 until the time of the experiment (5 µM for mGluR2 and mGluR3; 50 µM for mGluR7 and mGluR8) to maintain cell health.

Live cell fluorescence imaging

For live cell imaging of receptor internalization/β-arr localization, 24-48 hours post-transfection cells were labeled for 30 minutes at 37°C with 1 μM of the appropriate SNAP (SNAP-Surface546 from NEB for SNAP-tagged receptors or custom-made permeable SNAP-JF$_{549}$ for β-arr1–SNAP), CLIP (CLIP-Surface488, from NEB), or Halo (permeable JF$_{646}$ from Promega) dyes in extracellular buffer (EX) containing (in mM): 10 HEPES, 135 NaCl, 5.4 KCl, 2 CaCl$_2$, and 1 MgCl$_2$, pH 7.4. After labeling, cells were washed twice with fresh EX to remove excess fluorophore and incubated in EX containing different agonists (1 mM Glu for mGluR2 and mGluR8, 100 μM Glu for mGluR3, 10 mM Glu for mGluR7, 10 μM Vasopressin for V2R, 10 μM Isoproterenol for β2AR) for 30 min. Images were acquired using an Olympus IX83 inverted microscope equipped with scientific Complementary Metal-Oxide Semiconductor (sCMOS) ORCA Flash4v3.0 camera (Hamamatsu) and a 100X immersion oil objective (NA 1.45). Fluorophores were excited with 488 nm, 561 nm, and/or 640 nm lasers in widefield mode.

mGluR internalization was assessed using a surface labeling assay as previously described(*36*, *58*, *59*). Briefly, 24 hours post-transfection of SNAP-mGluR constructs, cells were incubated for 60 min at 37°C in warm DMEM + 10% FBS with either antagonist (50 μM LY341495) or agonist (1 mM Glu) for each condition. Subsequently, cells were labeled with 1 μM SNAP Surface Alexa 546 dye (NEB) in EX for 20 min at RT. Images were captured using an Olympus inverted microscope with a 60X immersion oil objective (NA 1.49). At least 10 images per condition were obtained per day, and the experiment was repeated for a minimum of three independent biological replicates. Quantification of remaining surface receptors was performed using the Li algorithm threshold method in ImageJ (Fiji). Background fluorescence mean intensity was subtracted from the mean intensity of fluorescent cells, and these values were normalized to the antagonist condition per day and construct. Percent internalization was calculated by normalizing the image values of the agonist condition to the antagonist condition, transforming these values into percentages, and averaging among all images per day. Each mean value represented an independent data point.

Single Molecule Pulldown (SiMPull)

SiMPull measurements were performed as previously described(*59*, *74*) with minor modifications. Forty hours post-transfection (see details above), HEK 293T cells were washed with EX buffer, and then labeled for 1 hour at 37° C with 1 μM CA-JF$_{549}$ (permeable; for Halo; Promega) and 1 μM SBG-JF$_{646}$ (impermeable; for SNAP; custom synthesized as previously(*111*)). For experiments with dual labeling of β-arr1-Halo (Fig. S2L, M) or β-arr2-Halo (Fig. S7C), cells were simultaneously labeled with 1 μM CA-JF$_{549}$ and 1 μM CA-JF$_{646}$ (permeable; for Halo; Promega). For experiments with dual labeling of β-arr1-SNAP and β-arr2-Halo (Fig. 7I-K), cells were simultaneously labeled with 1 μM BG-JF$_{646}$ (permeable; for SNAP; custom

synthesized as previously(111)) and 1 µM CA-JF$_{549}$. Following labeling, cells were washed with EX buffer before the incubation of agonist (1 mM Glu for mGluR2, mGluR3, mGluR5, mGluR8; 10 mM Glu for mGluR7; 10 µM Vasopressin for V2R; 10 µM Isoproterenol for β2AR) for 45 minutes at 37° C. Cells were then gently harvested in phosphate-buffered saline (0 Ca^{2+}, 0 Mg^{2+}) and pelleted at 10,000 rpm at 4°C for 5 minutes. Receptors were isolated via gentle lysis using 0.5% Lauryl Maltose Neopentyl Glycol (LMNG) + 0.05% Cholesteryl hemisuccinate (CHS) (Anatrace) plus protease inhibitor cocktail (Thermo Fisher Scientific) in Lysis Buffer (10 mM Tris HCl, 150 mM NaCl, 1 mM EDTA, pH 8.0) at 4°C for 1 hour. After lysis, samples were centrifuged at 17,000g for 20 min at 4°C, and supernatants were collected.

A microflow chamber was prepared using a glass coverslip and quartz slide passivated with mPEG-SVA and biotinylated-PEG (molecular weight: 5000; 50:1 molar ratio; Laysan Bio). Before each experiment, each chamber was washed with T50 buffer (50 mM NaCl and 10 mM tris, pH 8.0), incubated with NeutrAvidin (0.2 mg/mL) in T50 buffer for 5 minutes, followed by incubation with biotinylated anti-HA antibody (0.002 mg/ml; ab26228, Abcam) in T50 buffer for 20 minutes. Flow chambers were washed with T50 buffer after incubations with both NeutrAvidin and anti-HA.

Fresh lysate containing fluorescently labeled receptors was then diluted using buffer containing 0.05% LMNG and 0.005% CHS in EX buffer and added to the flow chamber. When a desired single-molecule spot density (~0.2 spots/µm^2) was obtained, unbound receptors were washed with the dilution buffer. Single-molecule movies were recorded using a 100× oil immersion objective (NA 1.49) on an inverted microscope (Olympus IX83) with TIR mode at 20 Hz with 50-ms exposure time using two scientific sCMOS cameras (same as above, Hamamatsu ORCA-Flash4v3.0). Movies were recorded until >90% of molecules were bleached in the field of each movie. Data were analyzed using a custom-built LabVIEW (National Instruments) program(112). Briefly, movies from both channels were concatenated and loaded on the analysis program to visualize each channel for identification of colocalized molecules. Bleaching steps were assigned by inspecting the fluorescence traces manually for each molecule and plotting to show the bleaching step distribution. As previously established, the subpopulation of molecules showing 2 bleaching steps is a function of incomplete fluorophore labeling, incidental bleaching, and other detection issues with 45-55% 2-step bleaching typically seen for strict dimers(52, 73).

Single Molecule FRET

48 hours post-transfection with a modified SNAP-mGluR8 clone containing a C-terminal HA tag alone or in combination with GRK2 and β-arr1ΔCTD, HEK 293T cells were washed with EX buffer, and then labeled for 45 min at 37° C. with 1 µM of BG-LD555 (donor) and 3 µM of BG-LD655 (acceptor). Labeling solution was removed, cells were washed with EX buffer, and lysates were prepared as described above for SiMPull experiments. After immobilization of receptors, buffer was exchanged with imaging buffer and

oxygen scavenging system with or without 1 mM glutamate as previously described(*81*). FRET measurements were conducted using an Olympus IX83 microscope in TIR mode, using a 1.49 NA 100x oil immersion objective, and solid-state 561-nm laser for excitation. Fluorescence emission was filtered through a 635LP and detected using two synchronized sCMOS cameras (Hamamatsu ORCA-Flash 4v3.0). smFRET time lapses were captured at 30 ms exposure. FRET movies were analyzed using the program SPARTAN (*113*). Briefly, FRET efficiency was calculated using the following equation: $\frac{I_A - 0.1 I_D}{I_D + I_A}$ where I_D is the donor intensity and I_A is the acceptor intensity after subtracting background fluorescence. Hidden Markov analysis was performed on the FRET traces using 3 states (0.15, 0.25, 0.35) to derive transition rates in SPARTAN. FRET histograms were plotted using GraphPad Prism and Gaussian fitting was performed using Origin.

mGluR8 Purification

Expi GNTi- cells (Thermo Fisher Scientific) were grown at 37 °C and 8% CO_2 to a density of 3×10^6 cells/mL in Expi media (Thermo Fisher Scientific) supplemented with 1:500 penicillin/streptomycin (Gibco) prior to transfection. Following manufacturer provided protocols, HA-FLAG-mGluR8 construct was transfected using the Expi293 transfection kit. The suspension was incubated at 37 °C for 20 h, then Expi enhancers were added to the cells as specified by the manufacturer (Thermo Fisher). Cell pellets were collected 72 hours after transfection by centrifugation at 4,000 rpm, flash-frozen in liquid nitrogen, and stored at −80 °C. For the mGluR8/β-arr1 complex, the steps above were followed except for co-transfecting HA-FLAG-mGluR8 with β-arr1-ΔCTD and GRK2-CaaX at a 2:1:1 ratio and adding 30 μM L-AP-4 (Tocris)and 10 μM VU6005649 (Tocris) 1 hour prior to harvesting cells.

Cell pellets were Dounce homogenized in lysis buffer containing 20 mM HEPES (pH 7.5), 150 mM NaCl, 2 mM CaCl2, 30 μM L-AP-4, and 10 μM VU6005649. For mGluR8-β-arr1ΔCTD purifications, lysis buffer was supplemented with 3uM 8:0 PI(4,5)P2 (Avanti Polar Lipids). Protein was extracted via the addition of n-Dodecyl-β-D-maltopyranoside (DDM) with cholesteryl hemisuccinate (CHS) (Anatrace, D310-CH210) to a final concentration of 50 mM DDM/4 mM CHS and nutated for 1 hour at 4 °C. Following ultracentrifugation at 4° C at 96,000 × g for 30 minutes, the supernatant was filtered using a 0.22μm filter (VWR). A column containing 5 mL FLAG resin (Genscript) was then equilibrated with running buffer containing 100 mM HEPES (pH 7.5), 150 mM NaCl, 2 mM CaCl2, 0.1% glyco-diosgenin (GDN), and 30 μM L-AP-4. Supernatant was then added to FLAG column resin and incubated at room temperature for 1 hour while being stirred. Resin was then reapplied to the column and flow through was collected. The column was washed first with 10 mL running buffer supplemented with 10 mM MgCl2 and 5 mM ATP to remove contaminating heat-shock proteins, followed 10 mL of running buffer. This series of

two washes was then repeated to ensure removal of any contaminants and to facilitate detergent exchange. FLAG-mGluR8 was subsequently eluted with running buffer supplemented with 0.2 mg/mL FLAG (DYKDDDDK) peptide (GenScript). Elution fractions were collected, analyzed by SDS-PAGE, and concentrated using a 50 kDa molecular weight cut-off spin concentrators (Amicon). The sample was then loaded onto a Superose 6 Increase 10/300 GL (Cytiva) column equilibrated in SEC buffer containing 100 mM HEPES (pH 7.5), 150 mM NaCl, 2 mM $CaCl_2$, 0.01% GDN, and 30 μM L-AP-4. Elution fractions were collected and analyzed by SDS-PAGE for purity. Key fractions were then concentrated using a 50 kDa molecular weight cut off spin concentrator.

For purifications containing mGluR8/β-arr1-ΔCTD complexes, protein was incubated with 2.5 µM Fab7 for 1 hour at room temperature, followed by addition of Sulfo-LC-SDA (Thermo Scientific) crosslinker to a final concentration of 250 µM. Samples were then incubated for 45 minutes at room temperature in the dark. To quench NHS-crosslinking Tris was added to a final concentration of 3 mM and samples were then irradiated with 365 nm light using an ENF-240C UV lamp (Spectro-UV, 4 watt tube) for 45 minutes at 4 °C. To increase arrestin association, 1uM 8:0 PIP2 was added and allowed to incubate overnight at 4 °C. Crosslinked samples were then purified using the Superose 6 size exclusion column and buffers as described above. Final samples for both mGluR8 and mGluR8/β-arr1ΔCTD complexes were concentrated to 3-5 mg/mL using a 50 kDa molecular weight cut off spin concentrator (as above) and used for negative stain and cryo-EM studies protein was either used immediately or flash frozen in liquid nitrogen and stored at -80 °C.

Mass Spectrometry

Purified mGluR8/β-arr1 preparations were dissolved in SDS lysis buffer (10 mM (tris(2-carboxyethyl)phosphine) (TCEP), 2% SDS, 150 mM NaCl, 50 mM Tris, pH 8.5) and reduced cysteine residues were alkylated with iodoacetamide at a final concentration of 20 mM and incubation at room temperature in the dark for 25 minutes. Alkylation was quenched with 50 mM freshly prepared DTT. SDS was removed by SP3 protein cleanup(114) and proteins were digested with Lys-C protease (Wako, Catalog Number 129–02541) and trypsin (Promega Catalog Number V5113). Peptides were subjected to micro-phospho enrichment as described recently in Susa et al., 2024(115) and analyzed on an Orbitrap Ascend Mass spectrometer (Thermo Fisher Scientific) using 180 or 300 min FAIMS runs (CVs -40V, -55V, -70V, -85V, cycle time 6s) at an MS1 resolution of 120,000 in the orbitrap. After HCD fragmentation (25% collision energy), MS2 scans were collected at 45,000 resolution in the orbitrap with a 1.6 m/z quadrupole isolation window. Data were searched with COMET(116) using additional masses of +15.994914 Da for oxidized methionine modification and + 79.966330 Da for phospho-modification on S, T and Y residues, dynamically. Peptides were identified using MS2 spectra and a false discovery rate (FDR) < 1% and was

achieved by applying a target-decoy database search strategy. Details of phospho-proteomic analysis are provided in Susa et al., 2024(*115*).

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD055267 and 10.6019/PXD055267.

Negative Stain EM

An aliquot of 3.5 µL of mGluR8 or mGluR8/β-arr1-ΔCTD samples was applied onto glow-discharged carbon-coated grids (Electron Microscopy Services, Cu 400 mesh) and allowed to adsorb for 30 s. Excess sample was blotted off using filter paper. Then 5 µL of freshly prepared 0.75% (w/v) uranyl formate stain was added for 30 s and immediately blotted. Another 5 µL of stain was then added to the grid for 5 minutes, followed by blotting and benchtop air-drying for 5 minutes.

Grids were imaged on a Talos L120C electron microscope (FEI) equipped with a 4K x 4K OneView CMOS camera (Gatan, Inc.) at a nominal magnification of 92,000x, corresponding to a pixel size of 1.6 Å. Manual data collection was performed with a defocus ranging from -0.4 µm to -0.8 µm and a dose of approximately 60 electrons per Å2 with 1 s of exposure time. CryoSPARC was used to collect and process micrographs, manually pick particles, and perform reference-free 2D classification.

Cryo-EM

mGluR8 or mGluR8/β-arr1-ΔCTD samples ranging from 3-5 mg/mL were applied at a volume of 3 µl to plasma treated UltrAuFoil 1.2/1.3 300 mesh grids (Quantifoil). Vitrified samples were prepared using a Vitrobot Mk IV (Thermo Fisher), blot time of 2 seconds, blot force of -2, at 100% humidity and 22°C. Grids were imaged with a Titan Krios electron microscope (Thermo Fisher) operated at 300 kV and a nominal magnification of 105,000× and equipped with a K3 camera (Gatan) set in super-resolution mode (0.426 Å pixel size). Images were collected using Leginon.

For mGluR8 alone datasets, beam induced motion was corrected using MotionCor2 in Relion 3.1 using two-fold binning and dose-weighting. These images were then used for contrast transfer function estimation via CTFFIND4.1(*117*). In Relion, particles were picked using a 3D initial model template of the receptor and extracted using a box size of 416. Further processing was then performed in CryoSPARC version 3.3.2(*118*). Ab initio followed by multiple rounds of heterogeneous refinement resulted in two good classes that were further refined via subsequent ab initio and heterogeneous refinement. The best heterogeneous refinement class/classes were then used for homogeneous refinement followed by non-uniform refinement with C1 symmetry. These particles were then imported back into Relion for Bayesian polishing(*119*) followed by non-uniform refinement to obtain the best full-length map. Initial mGluR8

models were built using AlphaFold2(*120*). Each model was then refined via iterative rounds of real-space refinement using the full-length map in Phenix(*121*) and manual refinement in Coot(*122*). Regions with poor resolution were either deleted, as in the case of some loops, or stubbed, as is the case of the TMDs. Visual representations were prepared using ChimeraX(*123*) and Pymol (Schrödinger, LLC 2015).

For datasets containing the mGluR8/β-arr1-ΔCTD complex, micrographs were motion corrected using Warp(*124*) and imported to CryoSPARC v4.5.3 for processing. Contrast transfer function estimation was performed using the PatchCTF job and particles were picked using an initial 3D model template of the complex and extracted with a box size of 588. Four iterative rounds of 2D classification were followed by ab initio and heterogeneous refinement, further 2D classification, and a second round of ab initio and heterogeneous refinement revealed one promising class of particles. This class was then split into two further classes following ab initio, heterogeneous, and non-uniform refinement with C1 symmetry to obtain best full-length maps. These maps contained higher resolution density for mGluR8 and lower resolution density that was assigned to β-arr1ΔCTD. The final models of mGluR8 in the absence of β-arr1 (described above) were used as starting points for modeling mGluR8 in the presence of β-arr1. Each model was then modified using Phenix and Coot as described above. Regions with poor resolution were either deleted, as in the case of some loops, or stubbed, as is the case of the TMDs. To place the β-arr1ΔCTD protomer, AlphaFold2 (AF2) multimer(*125*) was used to generate complexes of mGluR8 TMD dimers with a single copy of β-arr1ΔCTD. The CTDs of mGluR8 were removed for AF2 modeling due to their predicted disorder, the potential presence of "conditionally folded" regions of the CTD(*126*), and the inability of AF2 to phosphorylate residues. We found that AF2 produced a model that fit our density well, and this model was used to initially place the β-arr1 near our full length mGluR8 model. Further rigid-body refinement of the β-arr1 into our density was performed using Coot. Visual representations were prepared using ChimeraX(*123*) and Pymol (Schrödinger, LLC 2015). Structural comparison and alignment with existing GPCR-β-arr1 structures were performed in Chimera X by aligning the receptor using the 'matchmaker' function. Angles were measured as the difference in the middle reference plane change between arrestins (Fig. 5A) or from receptor to arrestin (Fig. 5B).

Molecular Dynamics Simulations

Atomistic molecular dynamics (MD) simulations were performed on the full-length mGluR8/β-arr1 complex embedded in a lipid bilayer with composition mimicking that of the plasma membrane. To build this full-length, lipid embedded mGluR8/βarr1 complex, we combined the mGluR8/β-arr1-ΔCTD model derived from the combination of cryo-EM experiments with AlphaFold2 predictions and rigid body docking as described above. To mimic the biologically relevant system, we docked a peptide containing the fully

phosphorylated mGluR8 CTD ST-rich region (residues 888 to 908) to the β-arr1 in the same orientation as a V2R-CTD/β-arr1 cryo EM structure (PDB ID: 4JQI). Additionally, we replaced the orthosteric agonist L-AP-4 with glutamate and removed the PAM (VU600) in our model to mimic the endogenous form of the mGluR8/β-arr1 complex. Using CHARMM-GUI membrane builder(*127*), the protein complex was embedded in a lipid bilayer containing POPC, POPE, POPS, cholesterol, sphingomyelin, and PIP$_2$ at approximately biological amounts(*128*). There were 20 cholesterol, 196 POPC, 20 POPE, 16 POPS, and 200 sphingomyelin molecules in the upper leaflet and 16 cholesterol, 104 POPC, 172 POPE, 116 POPS, 32 PIP$_2$ molecules in the lower leaflet. The protein-membrane system was then solvated and ionized in CHARMM-GUI using 0.2 M NaCl concentration. Overall, the system contained 792,623 total atoms including water, sodium, and chloride ions. Residues were protonated as recommended by CHARMM-GUI to reflect pH 7.0 (all histidine residues were deprotonated except residue 55).

Using NAMD version 2.14, the initial system was separately equilibrated three times (i.e. three independent replicates) using the standard CHARMM-GUI equilibration protocol. Each of the resulting three equilibrated systems were then used to seed a new set of 24 ~267 ns-long independent simulations (with randomly resetting velocities of all the atoms in the system). Thus, 72 independent replicates were collected with the cumulative MD sampling time of ~19 μs. These production runs were carried out with OpenMM version 7.5.1 using the following run parameters: 4 fs integration timestep enabled with hydrogen mass repartitioning, Particle Mesh Ewald (PME) for electrostatic interactions, under isothermal-isobaric (NPT) ensemble conditions using semi-isotropic pressure coupling, with MonteCarloMembraneBarostat and Langevin thermostats used to maintain constant pressure and temperature, respectively. Additional simulations parameters included: "friction" set to 1.0/picosecond, "EwaldErrorTolerance" 0.0005, "rigidwater" True, and "ConstraintTolerance" 0.000001. The van der Waals interactions were calculated applying a cut-off distance of 12 Å and switching the potential from 10 Å.

Simulations were analyzed using VMD plugins and home-made TCL and python scripts. Statistical tests were performed using GraphPad Prism10. All analyses were performed on the trajectories strided with a timestep of 0.4 ns, except the buried SASA calculation which was calculated at a timestep of 0.8 ns (sampled half as often as the other analyses for computational efficiency). To determine the x-y angles of β-arr1 relative to mGluR8, the principal axes of mGluR8 were determined using the Orient plugin in VMD and compared to a vector connecting the center-of-mass of the β-arr1 N and C lobes. To determine the β-arr1 z-angle, we compared the same β-arr1 vector to a vector along the membrane normal. RMSD values were calculated by aligning individual protomer chains and using the VMD RMSD command. Residue-to-residue and domain center-of-mass distances were calculated by measuring the length of the vector connecting the two points. Lipid hydrogen bond analyses were carried out using the default settings in the

VMD hydrogen bond calculator using default settings for bond lengths and angles and buried SASA analyses were calculated using the VMD "sasa" function using a probe radius of 1.4 Å as previously (*129*).

Single cell RNA sequencing analysis

We analyzed previously obtained single cell RNA sequencing data from the Anterior Cingulate Cortex (ACA SMART-seq, Allen Institute, 2018)(*93*). All cells passed the quality control criteria and underwent subsequent hierarchical clustering according to the similarity of their individual transcriptomes(*130*). A total of 5028 cells included from the mouse ACC cortex were grouped by cell type according to the different brain layers and populations of interest (L2/3, L2/5, L5, L5/6, L6, L6b glutamatergic and NGF, Sst+, PV+ and VIP GABAergic neurons). For co-expression analysis of *Arrb1* and *Arrb2* (Fig. 7 and S9), we took into consideration the low rate of false positives and under-sampling of a cell's total mRNA with this technique, thus imposing a threshold of five copies per million (CPM). Data were analyzed using Microsoft Excel and R Studio for plotting (*ggplot* package).

Acknowledgements:

We thank the entire Levitz lab (Weill Cornell Medicine), Jonathan Javitch (Columbia University), and Kaavya Krishna Kumar (Stanford University) for helpful discussions. We also thank the Weill Cornell Medicine Cryo-EM Core facility, the Weill Cornell Scientific Computing Unit, the NYU Langone Health Microscopy Laboratory, and the NYU Langone Health Cryo–EM Laboratory. This work is supported by NIH grants F31NS129320 (A.S.), F32GM148001 (D.M.), R01NS129904 (J. Levitz), the Charles Revson Fellowship (A.G.-H.), the Rohr Family Research Scholar Award (J. Levitz), and the Monique Weill-Caulier Award (J. Levitz).

Declaration of Interests:

The authors declare no conflicts of interest.

References:

1. A. S. Hauser, M. M. Attwood, M. Rask-Andersen, H. B. Schiöth, D. E. Gloriam, Trends in GPCR drug discovery: New agents, targets and indications. Nat Rev Drug Discov 16, 829–842 (2017).

2. D. Wacker, R. C. Stevens, B. L. Roth, How Ligands Illuminate GPCR Molecular Pharmacology. Cell 170, 414-427 (2017).

3. M. Von Zastrow, A. Sorkin, Mechanisms for Regulating and Organizing Receptor Signaling by Endocytosis. Annual Review of Biochemistry 44, 14 (2024).

4. A. K. Shukla, K. Xiao, R. J. Lefkowitz, Emerging paradigms of β-arrestin-dependent seven transmembrane receptor signaling. TiBS 36, 457-469 (2011).

5. J. Janetzko, R. Kise, B. Barsi-Rhyne, D. H. Siepe, F. M. Heydenreich, K. Kawakami, M. Masureel, S. Maeda, K. C. Garcia, M. von Zastrow, A. Inoue, B. K. Kobilka, Membrane phosphoinositides regulate GPCR-β-arrestin complex assembly and dynamics. Cell 185, 4560-4573.e19 (2022).

6. R. H. Oakley, S. A. Laporte, J. A. Holt, M. G. Caron, L. S. Barak, Differential affinities of visual arrestin, βarrestin1, and βarrestin2 for G protein-coupled receptors delineate two major classes of receptors. Journal of Biological Chemistry 275, 17201–17210 (2000).

7. R. H. Oakley, S. A. Laporte, J. A. Holt, L. S. Barak, M. G. Caron, Molecular Determinants Underlying the Formation of Stable Intracellular G Protein-coupled Receptor-β-Arrestin Complexes after Receptor Endocytosis. Journal of Biological Chemistry 276, 19452–19460 (2001).

8.J. Zhang, L. S. Barak, P. H. Anborgh, S. A. Laporte, M. G. Caron, S. S. G. Ferguson, Cellular trafficking of G protein-coupled receptor/β-arrestin endocytic complexes. Journal of Biological Chemistry 274, 10999–11006 (1999).

9. R. H. Oakley, A. Laporte, J. A. Holt, L. S. Barak, M. G. Caron, Association of-Arrestin with G Protein-coupled Receptors during Clathrin-mediated Endocytosis Dictates the Profile of Receptor Resensitization. Journal of Biological Chemistry 274, 32248–32257 (1999).

10. S. M. DeWire, S. Ahn, R. J. Lefkowitz, S. K. Shenoy, β-Arrestins and cell signaling. Annu. Rev. Physiol. 69:483-510 (2007).

11. N. R. Latorraca, M. Masureel, S. A. Hollingsworth, F. M. Heydenreich, C. M. Suomivuori, C. Brinton, R. J. L. Townshend, M. Bouvier, B. K. Kobilka, R. O. Dror, How GPCR Phosphorylation Patterns Orchestrate Arrestin-Mediated Signaling. Cell 183, 1813-1825.e18 (2020).

12. W. B. Asher, D. S. Terry, G. G. A. Gregorio, A. W. Kahsai, A. Borgia, B. Xie, A. Modak, Y. Zhu, W. Jang, A. Govindaraju, L. Y. Huang, A. Inoue, N. A. Lambert, V. V. Gurevich, L. Shi, R. J. Lefkowitz, S. C. Blanchard, J. A. Javitch, GPCR-mediated β-arrestin activation deconvoluted with single-molecule precision. Cell 185, 1661-1675.e16 (2022).

13. J. Maharana, P. Sarma, M. K. Yadav, S. Saha, V. Singh, S. Saha, M. Chami, R. Banerjee, A. K. Shukla, Structural snapshots uncover a key phosphorylation motif in GPCRs driving β-arrestin activation. Mol Cell 83, 2091-2107.e7 (2023).

14. A. K. Shukla, A. Manglik, A. C. Kruse, K. Xiao, R. I. Reis, W. C. Tseng, D. P. Staus, D. Hilger, S. Uysal, L. Y. Huang, M. Paduch, P. Tripathi-Shukla, A. Koide, S. Koide, W. I. Weis, A. A. Kossiakoff, B. K. Kobilka, R. J. Lefkowitz, Structure of active β-arrestin-1 bound to a G-protein-coupled receptor phosphopeptide. Nature 497, 137–141 (2013).

15. D. Mayer, F. F. Damberger, M. Samarasimhareddy, M. Feldmueller, Z. Vuckovic, T. Flock, B. Bauer, E. Mutt, F. Zosel, F. H. T. Allain, J. Standfuss, G. F. X. Schertler, X. Deupi, M. E. Sommer, M. Hurevich, A. Friedler, D. B. Veprintsev, Distinct G protein-coupled receptor phosphorylation motifs modulate arrestin affinity and activation and global conformation. Nat Commun 10 (2019).

16. A. Sente, R. Peer, A. Srivastava, M. Baidya, A. M. Lesk, S. Balaji, A. K. Shukla, M. M. Babu, T. Flock, Molecular mechanism of modulating arrestin conformation by GPCR phosphorylation. Nat Struct Mol Biol 25, 538–545 (2018).

17. C. Krasel, M. Bünemann, K. Lorenz, M. J. Lohse, β-arrestin binding to the β2-adrenergic receptor requires both receptor phosphorylation and receptor activation. Journal of Biological Chemistry 280, 9528–9535 (2005).

18. Y. Shiraishi, Y. Kofuku, T. Ueda, S. Pandey, H. Dwivedi-Agnihotri, A. K. Shukla, I. Shimada, Biphasic activation of β-arrestin 1 upon interaction with a GPCR revealed by methyl-TROSY NMR. Nat Commun 12 (2021).

19. N. R. Latorraca, J. K. Wang, B. Bauer, R. J. L. Townshend, S. A. Hollingsworth, J. E. Olivieri, H. E. Xu, M. E. Sommer, R. O. Dror, Molecular mechanism of GPCR-mediated arrestin activation. Nature 557, 452–456 (2018).

20. A. K. Shukla, G. H. Westfield, K. Xiao, R. I. Reis, L. Y. Huang, P. Tripathi-Shukla, J. Qian, S. Li, A. Blanc, A. N. Oleskie, A. M. Dosey, M. Su, C. R. Liang, L. L. Gu, J. M. Shan, X. Chen, R. Hanna, M. Choi, X. J. Yao, B. U. Klink, A. W. Kahsai, S. S. Sidhu, S. Koide, P. A. Penczek, A. A. Kossiakoff, V. L. Woods, B. K. Kobilka, G. Skiniotis, R. J. Lefkowitz, Visualization of arrestin recruitment by a G-protein-coupled receptor. Nature 512, 218–222 (2014).

21. Y. Kang, X. E. Zhou, X. Gao, Y. He, W. Liu, A. Ishchenko, A. Barty, T. A. White, O. Yefanov, G. W. Han, Q. Xu, P. W. De Waal, J. Ke, M. H. E. Tan, C. Zhang, A. Moeller, G. M. West, B. D. Pascal, N. Van Eps, L. N. Caro, S. A. Vishnivetskiy, R. J. Lee, K. M. Suino-Powell, X. Gu, K. Pal, J. Ma, X. Zhi, S. Boutet, G. J. Williams, M. Messerschmidt, C. Gati, N. A. Zatsepin, D. Wang, D. James, S. Basu, S. Roy-Chowdhury, C. E. Conrad, J. Coe, H. Liu, S. Lisova, C. Kupitz, I. Grotjohann, R. Fromme, Y. Jiang, M. Tan, H. Yang, J. Li, M. Wang, Z. Zheng, D. Li, N. Howe, Y. Zhao, J. Standfuss, K. Diederichs, Y. Dong, C. S. Potter, B. Carragher, M. Caffrey, H. Jiang, H. N. Chapman, J. C. H. Spence, P. Fromme, U. Weierstall, O. P. Ernst, V. Katritch, V. V. Gurevich, P. R. Griffin, W. L. Hubbell, R. C. Stevens, V. Cherezov, K. Melcher, H. E. Xu, Crystal structure of rhodopsin bound to arrestin by femtosecond X-ray laser. Nature 523, 561–567 (2015).

22. J. Grimes, Z. Koszegi, Y. Lanoiselée, T. Miljus, S. L. O'Brien, T. M. Stepniewski, B. Medel-Lacruz, M. Baidya, M. Makarova, R. Mistry, J. Goulding, J. Drube, C. Hoffmann, D. M. Owen, A. K. Shukla, J. Selent, S. J. Hill, D. Calebiro, Plasma membrane preassociation drives β-arrestin coupling to receptors and activation. Cell 186, 2238-2255.e20 (2023).

23. J. Maharana, F. K. Sano, P. Sarma, M. K. Yadav, L. Duan, T. M. Stepniewski, M. Chaturvedi, A. Ranjan, V. Singh, S. Saha, G. Mahajan, M. Chami, W. Shihoya, J. Selent, K. Young Chung, R. Banerjee, O. Nureki, A. K. Shukla, Molecular insights into atypical modes of b-arrestin interaction with seven transmembrane receptors. Science 383, 101–108 (2024).

24. S. Pandey, P. Kumari, M. Baidya, R. Kise, Y. Cao, H. Dwivedi-Agnihotri, R. Banerjee, X. X. Li, C. S. Cui, J. D. Lee, K. Kawakami, J. Maharana, A. Ranjan, M. Chaturvedi, G. D. Jhingan, S. A. Laporte, T. M. Woodruff, A. Inoue, A. K. Shukla, Intrinsic bias at non-canonical, β-arrestin-coupled seven transmembrane receptors. Mol Cell 81, 4605-4621.e11 (2021).

25. K. Chen, C. Zhang, S. Lin, X. Yan, H. Cai, C. Yi, L. Ma, X. Chu, Y. Liu, Y. Zhu, S. Han, Q. Zhao, B. Wu, Tail engagement of arrestin at the glucagon receptor. Nature 620, 904–910 (2023).

26. Y. Y. Liao, H. Zhang, Q. Shen, C. Cai, Y. Ding, D. D. Shen, J. Guo, J. Qin, Y. Dong, Y. Zhang, X. M. Li, Snapshot of the cannabinoid receptor 1-arrestin complex unravels the biased signaling mechanism. Cell 186, 5784-5797.e17 (2023).

27. J. Bous, A. Fouillen, H. Orcel, S. Trapani, X. Cong, S. Fontanel, J. Saint-Paul, J. Lai-Kee-Him, S. Urbach, N. Sibille, R. Sounier, S. Granier, B. Mouillac, P. Bron, Structure of the vasopressin hormone-V2 receptor-   -arrestin1 ternary complex. Sci. Adv 8, 7761 (2022).

28. D. P. Staus, H. Hu, M. J. Robertson, A. L. W. Kleinhenz, L. M. Wingler, W. D. Capel, N. R. Latorraca, R. J. Lefkowitz, G. Skiniotis, Structure of the M2 muscarinic receptor–β-arrestin complex in a lipid nanodisc. Nature 579, 297–302 (2020).

29. W. Huang, M. Masureel, Q. Qu, J. Janetzko, A. Inoue, H. E. Kato, M. J. Robertson, K. C. Nguyen, J. S. Glenn, G. Skiniotis, B. K. Kobilka, Structure of the neurotensin receptor 1 in complex with β-arrestin 1. Nature 579, 303–308 (2020).

30. Y. Lee, T. Warne, R. Nehmé, S. Pandey, H. Dwivedi-Agnihotri, M. Chaturvedi, P. C. Edwards, J. García-Nafría, A. G. W. Leslie, A. K. Shukla, C. G. Tate, Molecular basis of β-arrestin coupling to formoterol-bound β1-adrenoceptor. Nature 583, 862–866 (2020).

31. S. Nuber, U. Zabel, K. Lorenz, A. Nuber, G. Milligan, A. B. Tobin, M. J. Lohse, C. Hoffmann, β-Arrestin biosensors reveal a rapid, receptor-dependent activation/deactivation cycle. Nature 531, 661–664 (2016).

32. Q. Chen, C. T. Schafer, S. Mukherjee, M. Gustavsson, P. Agrawal, X.-Q. Yao, A. A. Kossiakoff, T. M. Handel, J. J. G. Tesmer, ACKR3-arrestin2/3 complexes reveal molecular consequences of GRK-dependent barcoding. bioRxiv (2023).

33. A. Srivastava, B. Gupta, C. Gupta, A. K. Shukla, Emerging Functional Divergence of β-Arrestin Isoforms in GPCR Function. Trends in Endocrinology and Metabolism 26, 11:628–642 (2015).

34. E. Ghosh, H. Dwivedi, M. Baidya, A. Srivastava, P. Kumari, T. Stepniewski, H. R. Kim, M. H. Lee, J. van Gastel, M. Chaturvedi, D. Roy, S. Pandey, J. Maharana, R. Guixà-González, L. M. Luttrell, K. Y. Chung, S. Dutta, J. Selent, A. K. Shukla, Conformational Sensors and Domain Swapping Reveal Structural and Functional Differences between β-Arrestin Isoforms. Cell Rep 28, 3287-3299.e6 (2019).

35. R. S. Haider, E. S. F. Matthees, J. Drube, M. Reichel, U. Zabel, A. Inoue, A. Chevigné, C. Krasel, X. Deupi, C. Hoffmann, β-arrestin1 and 2 exhibit distinct phosphorylation-dependent conformations when coupling to the same GPCR in living cells. Nat Commun 13 (2022).

36. A. Strauss, A. J. Gonzalez-Hernandez, J. Lee, N. Abreu, P. Selvakumar, L. Salas-Estrada, M. Kristt, A. Arefin, K. Huynh, D. C. Marx, K. Gilliland, B. J. Melancon, M. Filizola, J. Meyerson, J. Levitz, Structural basis of positive allosteric modulation of metabotropic glutamate receptor activation and internalization. Nat Commun 15 (2024).

37. L. M. Wingler, M. A. Skiba, C. Mcmahon, D. P. Staus, A. L. W. Kleinhenz, C.-M. Suomivuori, N. R. Latorraca, R. O. Dror, R. J. Lefkowitz, A. C. Kruse, STRUCTURAL BIOLOGY Angiotensin and biased analogs induce structurally distinct active conformations within a GPCR. Science 367, 888–892 (2020).

38. J. D. Violin, R. J. Lefkowitz, β-Arrestin-biased ligands at seven-transmembrane receptors. TiPS 28, 8:416-422 (2007).

39. C.-M. Suomivuori, N. R. Latorraca, L. M. Wingler, S. Eismann, M. C. King, A. L. W. Kleinhenz, M. A. Skiba, D. P. Staus, A. C. Kruse, R. J. Lefkowitz, R. O. Dror, Molecular mechanism of biased signaling in a prototypical G protein-coupled receptor. Science 367, 881–887 (2020).

40. C. Hoffmann, N. Ziegler, S. Reiner, C. Krasel, M. J. Lohse, Agonist-selective, receptor-specific interaction of human P2Y receptors with β-arrestin-1 and -2. Journal of Biological Chemistry 283, 30933–30941 (2008).

41. A. K. Shukla, J. D. Violin, E. J. Whalen, D. Gesty-Palmer, S. K. Shenoy, R. J. Lefkowitz, Distinct conformational changes in-arrestin report biased agonism at seven-transmembrane receptors. PNAS 105, 29:9988-9993 (2008).

42. K. Eichel, D. Jullié, B. Barsi-Rhyne, N. R. Latorraca, M. Masureel, J. B. Sibarita, R. O. Dror, M. Von Zastrow, Catalytic activation of β-Arrestin by GPCRs. Nature 557, 381–386 (2018).

43. A. Ellaithy, J. Gonzalez-Maeso, D. A. Logothetis, J. Levitz, Structural and Biophysical Mechanisms of Class C G Protein-Coupled Receptor Function. TiBS 45, 12:1049-1064 (2020).

45. A. B. Seven, X. Barros-Álvarez, M. de Lapeyrière, M. M. Papasergi-Scott, M. J. Robertson, C. Zhang, R. M. Nwokonko, Y. Gao, J. G. Meyerowitz, J. P. Rocher, D. Schelshorn, B. K. Kobilka, J. M. Mathiesen, G. Skiniotis, G-protein activation by a metabotropic glutamate receptor. Nature 595, 450–454 (2021).

46. S. Lin, S. Han, X. Cai, Q. Tan, K. Zhou, D. Wang, X. Wang, J. Du, C. Yi, X. Chu, A. Dai, Y. Zhou, Y. Chen, Y. Zhou, H. Liu, J. Liu, D. Yang, M. W. Wang, Q. Zhao, B. Wu, Structures of Gi-bound metabotropic glutamate receptors mGlu2 and mGlu4. Nature 594, 583–588 (2021).

47. X. Wang, M. Wang, T. Xu, Y. Feng, Q. Shao, S. Han, X. Chu, Y. Xu, S. Lin, Q. Zhao, B. Wu, Structural insights into dimerization and activation of the mGlu2–mGlu3 and mGlu2–mGlu4 heterodimers. Cell Res 33, 762–774 (2023).

48. F. He, C. G. Wu, Y. Gao, S. N. Rahman, M. Zaoralová, M. M. Papasergi-Scott, T. J. Gu, M. J. Robertson, A. B. Seven, L. Li, J. M. Mathiesen, G. Skiniotis, Allosteric modulation and G-protein selectivity of the Ca2+-sensing receptor. Nature 626, 1141–1148 (2024).

49. C. Mao, C. Shen, C. Li, D. D. Shen, C. Xu, S. Zhang, R. Zhou, Q. Shen, L. N. Chen, Z. Jiang, J. Liu, Y. Zhang, Cryo-EM structures of inactive and active GABAB receptor. Cell Res 30, 564–573 (2020).

50. C. Shen, C. Mao, C. Xu, N. Jin, H. Zhang, D. D. Shen, Q. Shen, X. Wang, T. Hou, Z. Chen, P. Rondard, J. P. Pin, Y. Zhang, J. Liu, Structural basis of GABAB receptor–Gi protein coupling. Nature 594, 594–598 (2021).

51. E. Doumazane, P. Scholler, L. Fabre, J. M. Zwier, E. Trinquet, J. P. Pin, P. Rondard, Illuminating the activation mechanisms and allosteric properties of metabotropic glutamate receptors. Proc Natl Acad Sci U S A 110 (2013).

52. V. A. Gutzeit, J. Thibado, D. S. Stor, Z. Zhou, S. C. Blanchard, O. S. Andersen, J. Levitz, Conformational dynamics between transmembrane domains and allosteric modulation of a metabotropic glutamate receptor. eLife 2019;8:e45116 doi: 10.7554/eLife.45116.001 (2019).

53. A. M. Cao, R. B. Quast, F. Fatemi, P. Rondard, J. P. Pin, E. Margeat, Allosteric modulators enhance agonist efficacy by increasing the residence time of a GPCR in the active state. Nat Commun 12 (2021).

54. Q. Zhou, D. Yang, M. Wu, Y. Guo, W. Guo, L. Zhong, X. Cai, A. Dai, W. Jang, E. Shakhnovich, Z. J. Liu, R. C. Stevens, N. A. Lambert, M. M. Babu, M. W. Wang, S. Zhao, Common activation mechanism of class a GPCRs. Elife 8 (2019).

55. A. S. Hauser, A. J. Kooistra, C. Munk, F. M. Heydenreich, D. B. Veprintsev, M. Bouvier, M. M. Babu, D. E. Gloriam, GPCR activation mechanisms across classes and macro/microscales. Nat Struct Mol Biol 28, 879–888 (2021).

56. D. Hilger, K. K. Kumar, H. Hu, M. F. Pedersen, E. S. O'Brien, L. Giehm, C. Jennings, G. Eskici, A. Inoue, M. Lerch, J. M. Mathiesen, G. Skiniotis, B. K. Kobilka, Structural insights into differences in G protein activation by family A and family B GPCRs. Science 369, 52 (2020).

57. D. Hilger, M. Masureel, B. K. Kobilka, Structure and dynamics of GPCR signaling complexes. Nat Struct Mol Biol 25, 4–12 (2018).

58. N. Abreu, A. Acosta-Ruiz, G. Xiang, J. Levitz, Mechanisms of differential desensitization of metabotropic glutamate receptors. Cell Rep 35 (2021).

59. J. Lee, A. J. Gonzalez-Hernandez, M. Kristt, N. Abreu, K. Roßmann, A. Arefin, D. C. Marx, J. Broichhagen, J. Levitz, Distinct beta-arrestin coupling and intracellular trafficking of metabotropic glutamate receptor homo-and heterodimers Sci. Adv. 9, eadi8076. (2023)

60. S. Lee, S. Park, H. Lee, S. Han, J.-M. Song, D. Han, Y. H. Suh, Nedd4 E3 ligase and beta-arrestins regulate ubiquitination, trafficking, and stability of the mGlu7 receptor. eLife 2019;8:e44502 (2019).

61. L. Iacovelli, G. Molinaro, G. Battaglia, M. Motolese, L. Di Menna, M. Alfiero, J. Blahos, F. Matrisciano, M. Corsi, C. Corti, V. Bruno, A. De Blasi, F. Nicoletti, Regulation of group II metabotropic glutamate receptors by G protein-coupled receptor kinases: MGlu2 receptors are resistant to homologous desensitization. Mol Pharmacol 75, 991–1003 (2009).

62. L. Iacovelli, M. Felicioni, R. Nisticò, F. Nicoletti, A. De Blasi, Selective regulation of recombinantly expressed mGlu7 metabotropic glutamate receptors by G protein-coupled receptor kinases and arrestins. Neuropharmacology 77, 303–312 (2014).

63. A. Reiner, J. Levitz, Glutamatergic Signaling in the Central Nervous System: Ionotropic and Metabotropic Receptors in Concert. Neuron 98, 6:1080-1098 (2018).

64. F. Nicoletti, J. Bockaert, G. L. Collingridge, P. J. Conn, F. Ferraguti, D. D. Schoepp, J. T. Wroblewski, J. P. Pin, Metabotropic glutamate receptors: From the workbench to the bedside. Neuropharmacology 60, 7-8:1017-1041 (2011).

65. K. L. Pierce, R. J. Lefkowitz, CLASSICAL AND NEW ROLES OF β-ARRESTINS IN THE REGULATION OF G-PROTEIN-COUPLED RECEPTORS Nat Rev. Neurosci. 2,727-733 (2001).

66. B. Jr, J. G. Krupnick, F. Santini, V. V Gurevich, R. B. Penn, A. W. Gagnon, J. H. Keen, J. L. Benovic, "Arrestin/Clathrin Interaction LOCALIZATION OF THE CLATHRIN BINDING DOMAIN OF NONVISUAL ARRESTINS TO THE CARBOXYL TERMINUS*" Journal of Biological Chemistry 272, 23:15011-15016. (1996)

67. R. C. Kern, D. S. Kang, J. L. Benovic, Arrestin2/clathrin interaction is regulated by key N- and C-terminal regions in arrestin2. Biochemistry 48, 7190–7200 (2009).

68. Y. M. Kim, J. L. Benovic, Differential roles of arrestin-2 interaction with clathrin and adaptor protein 2 in G protein-coupled receptor trafficking. Journal of Biological Chemistry 277, 30760–30768 (2002).

69. S. A. Laporte, R. H. Oakley, J. A. Holt, L. S. Barak, M. G. Caron, The interaction of β-arrestin with the AP-2 adaptor is required for the clustering of β2-adrenergic receptor into clathrin-coated pits. Journal of Biological Chemistry 275, 23120–23126 (2000).

70. A. Jain, R. Liu, B. Ramani, E. Arauz, Y. Ishitsuka, K. Ragunathan, J. Park, J. Chen, Y. K. Xiang, T. Ha, Probing cellular protein complexes using single-molecule pull-down. Nature 473, 484–488 (2011).

71. T. Ha, C. Kaiser, S. Myong, B. Wu, J. Xiao, Next generation single-molecule techniques: Imaging, labeling, and manipulation in vitro and in cellulo. Mol. Cell. 82, 2:304-314. (2022)..

72. M. Von Zastrow, B. K. Kobilka, Ligand-regulated internalization and recycling of human β2-adrenergic receptors between the plasma membrane and endosomes containing transferrin receptors. Journal of Biological Chemistry 267, 3530–3538 (1992).

73. J. K. Thibado, J. Y. Tano, J. Lee, L. Salas-Estrada, D. Provasi, A. Strauss, J. M. L. Ribeiro, G. Xiang, J. Broichhagen, M. Filizola, M. J. Lohse, J. Levitz, Differences in interactions between transmembrane domains tune the activation of metabotropic glutamate receptors. Elife 10 (2021).

74. J. Lee, H. Munguba, V. A. Gutzeit, M. Kristt, J. S. Dittman, J. Levitz, Defining the Homo- and Heterodimerization Propensities of Metabotropic Glutamate Receptors. Cell Rep 31 (2020).

75. J. Levitz, C. Habrian, S. Bharill, Z. Fu, R. Vafabakhsh, E. Y. Isacoff, Mechanism of Assembly and Cooperativity of Homomeric and Heteromeric Metabotropic Glutamate Receptors. Neuron 92, 143–159 (2016).

76. X. E. Zhou, Y. He, P. W. de Waal, X. Gao, Y. Kang, N. Van Eps, Y. Yin, K. Pal, D. Goswami, T. A. White, A. Barty, N. R. Latorraca, H. N. Chapman, W. L. Hubbell, R. O. Dror, R. C. Stevens, V. Cherezov, V. V. Gurevich, P. R. Griffin, O. P. Ernst, K. Melcher, H. E. Xu, Identification of Phosphorylation Codes for Arrestin Recruitment by G Protein-Coupled Receptors. Cell 170, 457-469.e13 (2017).

77. H. Chen, S. Zhang, X. Zhang, H. Liu, QR code model: a new possibility for GPCR phosphorylation recognition. Cell Communication and Signaling 20:23 (2022)..

78. P. Isaikina, I. Petrovic, R. P. Jakob, P. Sarma, A. Ranjan, M. Baruah, V. Panwalkar, T. Maier, A. K. Shukla, S. Grzesiek, A key GPCR phosphorylation motif discovered in arrestin2·CCR5 phosphopeptide complexes. Mol Cell 83, 2108-2121.e7 (2023).

79. A. Francesconi, R. M. Duvoisin, Role of the second and third intracellular loops of metabotropic glutamate receptors in mediating dual signal transduction activation. Journal of Biological Chemistry 273, 5615–5624 (1998).

80. J. Kniazeff, A. S. Bessis, D. Maurel, H. Ansanay, L. Prézeau, J. P. Pin, Closed state of both binding domains of homodimeric mGlu receptors is required for full activity. Nat Struct Mol Biol 11, 706–713 (2004).

81. R. Vafabakhsh, J. Levitz, E. Y. Isacoff, Conformational dynamics of a class C G-protein-coupled receptor. Nature 524, 497–501 (2015).

82. C. H. Habrian, J. Levitz, V. Vyklicky, Z. Fu, A. Hoagland, I. McCort-Tranchepain, F. Acher, E. Y. Isacoff, Conformational pathway provides unique sensitivity to a synaptic mGluR. Nat Commun 10 (2019).

83. C. Habrian, N. Latorraca, Z. Fu, E. Y. Isacoff, Homo- and hetero-dimeric subunit interactions set affinity and efficacy in metabotropic glutamate receptors. Nat Commun 14 (2023).

84. B. Wey-Hung Liauw, A. Foroutan, M. Schamber, W. Lu, H. S. Afsari, R. Vafabakhsh, Conformational fingerprinting of allosteric modulators in metabotropic glutamate receptor 2. Elife 11 (2022).

85. N. Lecat-Guillet, R. B. Quast, H. Liu, E. Bourrier, T. C. Møller, X. Rovira, S. Soldevila, L. Lamarque, E. Trinquet, J. Liu, J.-P. Pin, P. Rondard, E. Margeat, "Concerted conformational changes control metabotropic glutamate receptor activity" Sci. Adv. 9:eadf1378 (2023) https://doi.org/10.1126/sciadv.adf1378.

86. K. Krishna Kumar, H. Wang, C. Habrian, N. R. Latorraca, J. Xu, E. S. O'Brien, C. Zhang, E. Montabana, A. Koehl, S. Marqusee, E. Y. Isacoff, B. K. Kobilka, Stepwise activation of a metabotropic glutamate receptor. Nature 629, 951–956 (2024).

87. W. Fang, F. Yang, C. Xu, S. Ling, L. Lin, Y. Zhou, W. Sun, X. Wang, P. Liu, P. Rondard, P. Shi, J. P. Pin, C. Tian, J. Liu, Structural basis of the activation of metabotropic glutamate receptor 3. Cell Research 32:695–698 (2022).

88. C. Nasrallah, G. Cannone, J. Briot, K. Rottier, A. E. Berizzi, C. Y. Huang, R. B. Quast, F. Hoh, J. L. Banères, F. Malhaire, L. Berto, A. Dumazer, J. Font-Ingles, X. Gómez-Santacana, J. Catena, J. Kniazeff, C. Goudet, A. Llebaria, J. P. Pin, K. R. Vinothkumar, G. Lebon, Agonists and allosteric modulators promote signaling from different metabotropic glutamate receptor 5 conformations. Cell Rep 36 (2021).

89. A. Koehl, H. Hu, D. Feng, B. Sun, Y. Zhang, M. J. Robertson, M. Chu, T. S. Kobilka, T. Laermans, J. Steyaert, J. Tarrasch, S. Dutta, R. Fonseca, W. I. Weis, J. M. Mathiesen, G. Skiniotis, B. K. Kobilka, Structural insights into the activation of metabotropic glutamate receptors. Nature 566, 79–84 (2019).

90. A. Manglik, A. C. Kruse, Structural Basis for G Protein-Coupled Receptor Activation. Biochemistry 56 (42), 5628-5634 (2017).

91. J. Grimes, Z. Koszegi, Y. Lanoiselée, T. Miljus, S. L. O'Brien, T. M. Stepniewski, B. Medel-Lacruz, M. Baidya, M. Makarova, R. Mistry, J. Goulding, J. Drube, C. Hoffmann, D. M. Owen, A. K. Shukla, J. Selent, S. J. Hill, D. Calebiro, Plasma membrane preassociation drives β-arrestin coupling to receptors and activation. Cell 186, 2238-2255.e20 (2023).

92. T. D. Prickett, X. Wei, I. Cardenas-Navia, J. K. Teer, J. C. Lin, V. Walia, J. Gartner, J. Jiang, P. F. Cherukuri, A. Molinolo, M. A. Davies, J. E. Gershenwald, K. Stemke-Hale, S. A. Rosenberg, E. H. Margulies, Y. Samuels, Exon capture analysis of G protein-coupled receptors identifies activating mutations in GRM3 in melanoma. Nat Genet 43, 1119–1126 (2011).

93. N. L. Jorstad, J. Close, N. Johansen, A. M. Yanny, E. R. Barkan, K. J. Travaglini, D. Bertagnolli, J. Campos, T. Casper, K. Crichton, N. Dee, S. L. Ding, E. Gelfand, J. Goldy, D. Hirschstein, K. Kiick, M. Kroll, M. Kunst, K. Lathia, B. Long, N. Martin, D. McMillen, T. Pham, C. Rimorin, A. Ruiz, N. Shapovalova, S. Shehata, K. Siletti, S. Somasundaram, J. Sulc, M. Tieu, A. Torkelson, H. Tung, E. M. Callaway, P. R. Hof, C. D. Keene, B. P. Levi, S. Linnarsson, P. P. Mitra, K. Smith, R. D. Hodge, T. E. Bakken, E. S. Lein, Transcriptomic cytoarchitecture reveals principles of human neocortex organization. Science 382 (2023).

94. A. J. Gonzalez-Hernandez, H. Munguba, J. Levitz, Emerging modes of regulation of neuromodulatory G protein-coupled receptors. TiNS 47, 8:635:650 (2024).

95. L. Fourgeaud, A. S. Bessis, F. Rossignol, J. P. Pin, J. C. Olivo-Marin, A. Hémar, The metabotropic glutamate receptor mGluR5 is endocytosed by a clathrin-independent pathway. Journal of Biological Chemistry 278, 12222–12230 (2003).

96. J. R. van Senten, T. C. Møller, E. Von Moo, S. D. Seiersen, H. Bräuner-Osborne, Use of CRISPR/Cas9-edited HEK293 cells reveals that both conventional and novel protein kinase C isozymes are involved in mGlu5a receptor internalization. Journal of Biological Chemistry 298 (2022).

97. N. Scheefhals, L. A. E. Catsburg, M. L. Westerveld, T. A. Blanpied, C. C. Hoogenraad, H. D. MacGillavry, Shank Proteins Couple the Endocytic Zone to the Postsynaptic Density to Control Trafficking and Signaling of Metabotropic Glutamate Receptor 5. Cell Rep 29, 258-269.e8 (2019).

98. A. G. Eng, D. A. Kelver, T. P. Hedrick, G. T. Swanson, Transduction of group I mGluR-mediated synaptic plasticity by β-arrestin2 signalling. Nat Commun 7 (2016).

99. A. B. Kleist, S. Jenjak, A. Sente, L. J. Laskowski, M. Szpakowska, M. M. Calkins, E. I. Anderson, L. M. McNally, R. Heukers, V. Bobkov, F. C. Peterson, M. A. Thomas, A. Chevigné, M. J. Smit, J. D. McCorvy, M. Madan Babu, B. F. Volkman, "Conformational selection guides b-arrestin recruitment at a biased G protein-coupled receptor" Science 377(6602):222-228. (2022).

100. F. Bumbak, J. B. Bower, S. C. Zemmer, A. Inoue, M. Pons, J. C. Paniagua, F. Yan, J. Ford, H. Wu, S. A. Robson, R. A. D. Bathgate, D. J. Scott, P. R. Gooley, J. J. Ziarek, Stabilization of pre-existing neurotensin receptor conformational states by β-arrestin-1 and the biased allosteric modulator ML314. Nat Commun 14 (2023).

101. A. R. B. Thomsen, B. Plouffe, T. J. Cahill, A. K. Shukla, J. T. Tarrasch, A. M. Dosey, A. W. Kahsai, R. T. Strachan, B. Pani, J. P. Mahoney, L. Huang, B. Breton, F. M. Heydenreich, R. K. Sunahara, G. Skiniotis, M. Bouvier, R. J. Lefkowitz, GPCR-G Protein-β-Arrestin Super-Complex Mediates Sustained G Protein Signaling. Cell 166, 907–919 (2016).

102. A. H. Nguyen, A. R. B. Thomsen, T. J. Cahill, R. Huang, L. Y. Huang, T. Marcink, O. B. Clarke, S. Heissel, A. Masoudi, D. Ben-Hail, F. Samaan, V. P. Dandey, Y. Z. Tan, C. Hong, J. P. Mahoney, S. Triest, J. Little, X. Chen, R. Sunahara, J. Steyaert, H. Molina, Z. Yu, A. des Georges, R. J. Lefkowitz, Structure of an endosomal signaling GPCR–G protein–β-arrestin megacomplex. Nat Struct Mol Biol 26, 1123–1131 (2019).

103. J. S. Smith, T. F. Pack, A. Inoue, C. Lee, K. Zheng, I. Choi, D. S. Eiger, A. Warman, X. Xiong, Z. Ma, G. Viswanathan, I. M. Levitan, L. K. Rochelle, D. P. Staus, J. C. Snyder, A. W. Kahsai, M. G. Caron, S. Rajagopal, Noncanonical scaffolding of Gαi and b-arrestin by G protein-coupled receptors. Science (1979) 371 (2021).

104. B. Sokrat, A. H. Nguyen, A. R. B. Thomsen, L. Y. Huang, H. Kobayashi, A. W. Kahsai, J. Kim, B. X. Ho, S. Ma, J. Little, C. Ehrhart, I. Pyne, E. Hammond, M. Bouvier, Role of the V2R–βarrestin–Gβγ complex in promoting G protein translocation to endosomes. Commun Biol 7 (2024).

105. J. Perroy, F. Raynaud, V. Homburger, M. C. Rousset, L. Telley, J. Bockaert, L. Fagni, Direct interaction enables cross-talk between ionotropic and group I metabotropic glutamate receptors. Journal of Biological Chemistry 283, 6799–6805 (2008).

106. V. O'Connor, E. F. Oussama, E. Bofill-Cardona, C. Nanoff, M. Freissmuth, A. Karschin, J. M. Airas, H. Betz, S. Boehm, Calmodulin Dependence of Presynaptic Metabotropic Glutamate Receptor Signaling. Science 286, 1180–1184 (1999).

107. P. R. Brakeman, A. A. Lanahan, O'Brien. R., K. Roche, C. A. Barnes, R. L. Huganir, P. F. Worley, Homer: a protein that selectively binds metabotropic glutamate receptors. Nature 386, 284–288 (1997).

108. C. M. Niswender, P. J. Conn, Metabotropic glutamate receptors: Physiology, pharmacology, and disease. Annu. Rev. Pharmacol. Toxicol. 50:295–322 (2010).

109. Y. Nakajima, S. Mochida, K. Okawa, S. Nakanishi, Ca 2-dependent release of Munc18-1 from presynaptic mGluRs in short-term facilitation. Proc Natl Acad Sci U S A 106, 18385–18389 (2009).

110. C. L. Morales-Perez, C. M. Noviello, R. E. Hibbs, Manipulation of Subunit Stoichiometry in Heteromeric Membrane Proteins. Structure 24, 797–805 (2016).

111. P. Poc, V. A. Gutzeit, J. Ast, J. Lee, B. J. Jones, E. D'Este, B. Mathes, M. Lehmann, D. J. Hodson, J. Levitz, J. Broichhagen, Interrogating surface: Versus intracellular transmembrane receptor populations using cell-impermeable SNAP-tag substrates. Chem Sci 11, 7871–7883 (2020).

112. M. H. Ulbrich, E. Y. Isacoff, Subunit counting in membrane-bound proteins. Nat Methods 4, 319–321 (2007).

113. M. F. Juette, D. S. Terry, M. R. Wasserman, R. B. Altman, Z. Zhou, H. Zhao, S. C. Blanchard, Single-molecule imaging of non-equilibrium molecular ensembles on the millisecond timescale. Nat Methods 13, 341–344 (2016).

114. C. S. Hughes, S. Moggridge, T. Müller, P. H. Sorensen, G. B. Morin, J. Krijgsveld, Single-pot, solid-phase-enhanced sample preparation for proteomics experiments. Nat Protoc 14, 68–85 (2019).

115. K. J. Susa, G. A. Bradshaw, R. J. Eisert, C. M. Schilling, M. Kalocsay, S. C. Blacklow, A. C. Kruse, A spatiotemporal map of co-receptor signaling networks underlying B cell activation. Cell Rep 43 (2024).

116. J. K. Eng, M. R. Hoopmann, T. A. Jahan, J. D. Egertson, W. S. Noble, M. J. MacCoss, A Deeper Look into Comet - Implementation and Features. J Am Soc Mass Spectrom 26, 1865–1874 (2015).

117. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron micrographs. J Struct Biol 192, 216–221 (2015).

118. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, CryoSPARC: Algorithms for rapid unsupervised cryo-EM structure determination. Nat Methods 14, 290–296 (2017).

119. J. Zivanov, T. Nakane, S. H. W. Scheres, A Bayesian approach to beam-induced motion correction in cryo-EM single-particle analysis. IUCrJ 6, 5–17 (2019).

120. P. Bryant, G. Pozzati, A. Elofsson, Improved prediction of protein-protein interactions using AlphaFold2. Nat Commun 13 (2022).

121. D. Liebschner, P. V. Afonine, M. L. Baker, G. Bunkoczi, V. B. Chen, T. I. Croll, B. Hintze, L. W. Hung, S. Jain, A. J. McCoy, N. W. Moriarty, R. D. Oeffner, B. K. Poon, M. G. Prisant, R. J. Read, J. S. Richardson, D. C. Richardson, M. D. Sammito, O. V. Sobolev, D. H. Stockwell, T. C. Terwilliger, A. G. Urzhumtsev, L. L. Videau, C. J. Williams, P. D. Adams, Macromolecular structure determination using X-rays, neutrons and electrons: Recent developments in Phenix. Acta Crystallogr D Struct Biol 75, 861–877 (2019).

122. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. Acta Crystallogr D Biol Crystallogr 66, 486–501 (2010).

123. T. D. Goddard, C. C. Huang, E. C. Meng, E. F. Pettersen, G. S. Couch, J. H. Morris, T. E. Ferrin, UCSF ChimeraX: Meeting modern challenges in visualization and analysis. Protein Science 27, 14–25 (2018).

124. D. Tegunov, P. Cramer, Real-time cryo-electron microscopy data preprocessing with Warp. Nat Methods 16, 1146–1152 (2019).

125. R. Evans, M. O'neill, A. Pritzel, N. Antropova, A. Senior, T. Green, A. Žídek, R. Bates, S. Blackwell, J. Yim, O. Ronneberger, S. Bodenstein, M. Zielinski, A. Bridgland, A. Potapenko, A. Cowie, K. Tunyasuvunakool, R. Jain, E. Clancy, P. Kohli, J. Jumper, D. Hassabis, Protein complex prediction with AlphaFold-Multimer. bioRxiv (Preprint) https//doi.org/10.1101/2021.10.04.463034 (2022).

126. T. Reid Alderson, I. Pritišanac, D. Kolaric, A. M. Moses, J. D. Forman-Kay, Systematic identification of conditionally folded intrinsically disordered regions by AlphaFold2. Proc Natl Acad Sci U S A 120 (2023).

127. S. Jo, T. Kim, V. G. Iyer, W. Im, CHARMM-GUI: A web-based graphical user interface for CHARMM. J Comput Chem 29, 1859–1865 (2008).

128. M. Doktorova, J. L. Symons, I. Levental, Structural and functional consequences of reversible lipid asymmetry in living membranes. Nat. Chem. Bio. 16, 12:1321-1330. (2020).

129. C. D. Mancinelli, D. C. Marx, A. J. Gonzalez-Hernandez, K. Huynh, L. Mancinelli, A. Arefin, G. Khelashvilli, J. Levitz, D. Eliezer, Control of G protein–coupled receptor function via membrane-interacting intrinsically disordered C-terminal domains. Proc Natl Acad Sci U S A 121 (2024).

130. B. Tasic, Z. Yao, L. T. Graybuck, K. A. Smith, T. N. Nguyen, D. Bertagnolli, J. Goldy, E. Garren, M. N. Economo, S. Viswanathan, O. Penn, T. Bakken, V. Menon, J. Miller, O. Fong, K. E. Hirokawa, K. Lathia, C. Rimorin, M. Tieu, R. Larsen, T. Casper, E. Barkan, M. Kroll, S. Parry, N. V. Shapovalova, D. Hirschstein, J. Pendergraft, H. A. Sullivan, T. K. Kim, A. Szafer, N. Dee, P. Groblewski, I. Wickersham, A. Cetin, J. A. Harris, B. P. Levi, S. M. Sunkin, L. Madisen, T. L. Daigle, L. Looger, A. Bernard, J. Phillips, E. Lein, M. Hawrylycz, K. Svoboda, A. R. Jones, C. Koch, H. Zeng, Shared and distinct transcriptomic cell types across neocortical areas. Nature 563, 72–78 (2018).

FIGURE 1

Figure 1. A single molecule assay for detection of GPCR/β-arr complexes reveals a range of efficiencies and stoichiometries, including 2:2 mGluR/β-arr1 complexes.

(A-C) Schematic (A), representative images (B), and summary quantification (C) of SiMPull assay of GPCR/β-arr1 complexes under agonist-treated conditions with GRK2-CaaX co-expression. Yellow circles (B) indicate colocalized spots indicative of mGluR/β-arr1 complexes. **(D-E)** Representative images and bleaching step traces (D) and photobleaching step distributions (F) for mGluR3/β-arr1-ΔCTD complexes. Arrows indicate bleaching steps. **(F)** Summary of % of colocalized molecules showing 2-step bleaching in the receptor (top) and β-arr1 (bottom) channels across receptors. Points represent individual movies (C, E, F). All data shown as mean ± SEM. Scale bar: 5 μm.

FIGURE 2

Figure 2. Defining tail and core interactions that drive mGluR8/β-arr1 coupling. (A) Summary graph of SiMPull data for colocalization of all mGluR CTD point mutations either with wild type β-arr1-Halo (with GRK2 co-expression) or β-arr1-ΔCTD (with GRK2-CaaX co-expression). **(B)** Scatter plot showing correlation between effects of mutations on β-arr1 co-localization in SiMPull (y-axis) and internalization in cell-based surface labeling assay (x-axis). A linear fit to the data excluding outliers (S888A, T896A, T899A) is shown with the 95% confidence interval shaded. **(C)** Summary graphs showing impairment of β-arr1ΔCTD colocalization in SiMPull (left) and mGluR8 internalization (right) with the F777D mutation in ICL3. **(D)** Left, schematic showing inter-LBD FRET assay where glutamate binding decreases FRET levels due to LBD closure and reorientation. Right, representative smFRET traces showing occasional transitions between low, medium, and high FRET states (dotted lines). **(E)** Histograms showing FRET distributions in the absence or presence of 1 mM Glutamate. β-arr1 increases the height of the low FRET peak. Inset shows occupancy of the low FRET 0.15 peak as calculated from Gaussian fits across individual experiments. **(F)** Summary schematic showing that β-arr1 stabilizes the active, low FRET conformation of mGluR8. Points represent individual movies (A, C, E) or separate experimental days (C, right). Unpaired t-test is used in (C, E). **, p<0.01; ***, p<0.001. All data shown as mean ± SEM.

FIGURE 3

Figure 3. Structural analysis of agonist- and PAM-bound mGluR8 and mGluR8/β-arr1 complexes. (A-D) Representative 2-D classes from negative stain analysis of mGluR8 alone (A) or mGluR8/β-arr1 (B) samples. A cartoon representation showing proposed ligand binding domain (LBD), micelle, and β-arr1 organization for each complex. **(E)** Full-length cryo-EM map and model of mGluR8 class 1 bound to the orthosteric agonist L-AP-4 (green) and PAM VU600 (blue). **(F)** Multiple views showing mGluR8/β-arr1 cryo-EM map contoured to two different levels with β-arr1 aligned into the β-arr1 density. **(G)** Model of mGluR8/β-arr1 bound to β-arr1 guided by cryo-EM density and AlphaFold2. The β-arr1 tilt z-angle is shown. **(H)** Snapshot showing overlay of the final mGluR8/β-arr1 conformation from 72 molecular dynamics trajectories. **(I)** Top, extracellular views showing similar TM6-TM6 interface between mGluR8 alone and mGluR8/β-arr1. Bottom, alignment of β-arr1-bound chain A TMD of mGluR8/β-arr1 structure with chain A of the mGluR8 alone structure showing repositioning and increased resolution of intracellular face of mGluR8 TMD to accomodate β-arr1. **(J)** Overlay of the mGluR8/β-arr1 model with β-arr1 bound structures of family A GPCRs (PDB: 8WU1, CB1R-β-arr1; 6TKO, β1AR-β-arr1; 7SRS, 5-HT2BR-β-arr1; 7R0C, V2R-β-arr1; 6UP7, NTSR1-β-arr1). The different β-arr1 orientations are quantified by the x-y angle of rotation relative to that in mGluR8. **(K)** Summary heat map of x-y and z-angle of β-arr1 relative to mGluR8 across all MD simulations.

FIGURE 4

Figure 4. Structural and functional analysis of mGluR8/β-arr1 core coupling. (A) Cytoplasmic view of mGluR8/β-arr1 model showing that anchoring via core interactions with one subunit (light purple, left) enables a β-arr1 (translucent green) orientation that sterically occludes access to both subunits. **(B)** Side views showing mGluR8/β-arr1 interface. Intracellular loop residues are shown as spheres in blue (ICL1), pink (ICL2), and gold (ICL3). **(C)** Zoom in of mGluR8/β-arr1 core interface highlighting key residues from cryo-EM model. **(D)** Snapshots from MD simulations highlighting two different interaction modes between the β-arr1 C-edge loops and the TMD of mGluR8 chain B and membrane lipids (grey spheres). **(E)** Summary graph of SiMPull data for colocalization of all mGluR8 core point mutations with β-arr1-ΔCTD (with GRK2 co-expression). **(F)** Scatter plot showing correlation between effects of mutations on β-arr1-ΔCTD co-localization in SiMPull (y-axis) and internalization in cell-based surface labeling assay (x-axis). A linear fit is shown with the 95% confidence interval shaded. Points represent individual movies (F). 1-way ANOVA with Dunnet's multiple comparisons test (to WT) is shown in (F). ***, p<0.001. All data shown as mean ± SEM.

FIGURE 5

Figure 5. An expanded repertoire of mGluR8/β-arr complexes. (A-B) Surface labeling assay internalization data showing evidence for *trans* and *cis* β-arr1 coupling driving mGluR8 (A) and mGluR2/8 (B) internalization. **(C)** Schematic, top, representative SiMPull image, left, and representative photobleaching step trace, right, showing the formation of 1:1:1 mGluR2:mGluR8:β-arr1 complexes **(D-E)** Schematic (D) and summary graph (E) showing the formation of 2:1 mGluR8:β-arr2 complexes. **(F-H)** Schematic (F), representative image (G, left), photobleaching step traces (G, right), and summary graph (H) showing co-localization of β-arr1 and β-arr2 upon pulldown via mGluR8 but not V2R. (I) Schematic of proposed modes of β-arr coupling for family A GPCRs (2 modes) and mGluR8 as a representative family C GPCR (20 modes). Points represent individual experiments (A, B) or movies (E, H). GRK2-CaaX was co-expressed for all SiMPull experiments shown in this figure. 1-way ANOVA with Tukey's multiple comparisons test is used in (A, B) and unpaired t-test is used in (E, H). *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$. All data shown as mean ± SEM.