Type of article: ORIGINAL ARTICLE

Informative title: Extensive clonal reproduction drives liana proliferation following

forest fragmentation.

Concise title: Liana clonal proliferation in fragmented forests.

Authors: Hideki Mori*[1], Takashi Kamijo[2]

[1] Forestry and Forest Products Research Institute, Matsunosato 1, Tsukuba, Ibaraki, 305-

8687, Japan

[2] Faculty of Life and Environmental Sciences, University of Tsukuba, 1-1-1 Tennodai,

Tsukuba, Ibaraki, 305-8572 Japan

*Corresponding author: Hideki Mori (Email: s1010750@gmail.com)

ABSTRACT

• **Background and Aims:** As forest landscapes become increasingly fragmented worldwide, concerns are growing over the cascading effects on biodiversity and ecosystem functioning. While such fragmentation often promotes liana proliferation, the role of clonal reproduction in this process remains poorly understood. In this study, we investigated how a temperate liana, *Wisteria floribunda*, reproduces and expands its distribution through clonal growth in fragmented forests.

• **Methods:** Using DNA analyses in fragmented forests (21.6 ha) and in a continuous forest plot (6 ha) in central Japan, we examined the distribution and structure of genets to evaluate their contribution to the dominance of *W. floribunda*.

• **Key Results:** Our findings revealed that 94% of the apparent individuals (ramets, N = 273 in total) in the fragmented forests, were derived from clonal reproduction, significantly exceeding the 45% observed in the continuous forest. These clonal genets collectively covered 57% of the study area in the fragmented forests, compared to only 7% in the continuous forest. Notably, three large genets in the fragmented forests spanned multiple patches, with the largest covering 9.5 ha across seven patches. These genets were predominantly located in areas historically covered by forests, suggesting that they likely expanded through clonal reproduction before fragmentation. Furthermore, stem density in fragmented forests was approximately three times higher than in the continuous forest.

• **Conclusions:** These results demonstrate that extensive clonal reproduction drives liana proliferation in fragmented forests. This study underscores the importance of considering clonal strategies when assessing and predicting liana dynamics in fragmented landscapes,

39   where historical and ongoing clonal reproduction play critical roles in their dominance.

40   (250 words)

42   **Keywords:** Woody vines; fragmented forest; anthropogenic disturbance; habitat

43   fragmentation; vegetative reproduction; clonal reproduction; forest fragmentation; lianas;

44   *Wisteria floribunda*

INTRODUCTION

In recent decades, deforestation has escalated rapidly, driving widespread forest fragmentation on a global scale and prompting concerns over its profound impacts on biodiversity and ecosystem functioning (Hansen et al., 2013; Liu et al., 2018). Such fragmentation disrupts habitat structure and connectivity, altering ecological processes across diverse taxa, including plants, animals, and microorganisms (Haddad et al., 2015; Li et al., 2022; Kiesewetter et al., 2023) and ultimately affecting species interactions and ecosystem functions (Fahrig, 2003; Wilson et al., 2016). Among these impacts, increased liana (woody vine) abundance is particularly noteworthy, with fragmented forests often exhibiting higher liana density and dominance (Schnitzer and Bongers, 2011; Campbell et al., 2015; Ladwig and Meiners, 2015). Lianas thrive in disturbed conditions, especially in edge habitats where they benefit from greater light availability and suitable small trees for colonisation (Laurance et al., 2001; Londré and Schnitzer, 2006; Jones et al., 2017; Campbell et al., 2018). This proliferation is considered a major factor in their global rise relative to trees (Schnitzer and Bongers, 2011; Ngute et al., 2024), yet the specific ecological mechanisms—particularly at the species level—that drive liana expansion in fragmented landscapes remain poorly understood.

Clonal reproduction is an important mechanism influencing the abundance and spatial distribution patterns of lianas at the stand scale. In natural continuous forests, the substantial contribution of clonal reproduction to liana abundance and distribution has been reported in previous studies (Yorke et al., 2013; Ledo and Schnitzer, 2014; Mori et al., 2018; Schnitzer et al., 2021). In contrast, under bright-light environments such as gaps and edges, several studies have reported relatively higher occurrence rates (Yuan, 2024) and survival rates (Mori et al., 2020) of seed-origin liana seedlings compared to those

under closed canopies. In addition, forest fragmentation may limit the spatial expansion

of clonal reproduction beyond the edges of fragmented stands, reducing its role in liana

proliferation compared to continuous forests, where clonal growth enables wider

distribution within a stand (Putz, 1984; Mori et al., 2018). Simulation studies also suggest

that fragmentation, which reduces patch size, favours long-distance dispersal strategies,

while clonal reproduction is less effective in small patches (Travis and Dytham, 1998;

Hiebeler, 2004; Fukui and Araki, 2014). However, the extent to which clonal reproduction

contributes to liana proliferation in fragmented forests remains largely unexamined,

leaving a critical gap in understanding the ecological mechanisms underlying liana

proliferation in these altered landscapes.

Wisteria floribunda (Fabaceae) is a liana species that is widely distributed across

temperate forests in the Northern Hemisphere (GBIF Secretariat, 2023). It reproduces

clonally by producing apparent individuals (hereafter referred to as ramets) through the

extension of stolons (Sakai et al., 2002; Mori et al., 2021). This species is known as a

dominant liana species in Japanese temperate forests; In a 6-ha permanent plot established

in natural continuous forests in the same region as the study area (Ibaraki Prefecture,

Japan), *W. floribunda* accounted for 69% of the number of liana stems and 85% of the

total basal area (Mori et al., 2016a, 2024a). In addition, a previous study conducted in the

same 6-ha plot have revealed the clonal structure of this species, showing that more than

half of its stems are derived from clonal reproduction, with ramets up to 180 m apart

belonging to the same clone (Mori et al., 2018). As a representative temperate liana

species with well-documented genetic structure and abundance in natural continuous

forests, *W. floribunda* offers valuable insights for understanding the influence and

mechanisms of clonal reproduction on liana proliferation in fragmented forests.

The primary objective of this study was to investigate how clonal reproduction

contributes to the proliferation of the liana *W. floribunda* in fragmented forests. We

focused on a 21.6-ha area comprised of multiple forest fragments in Ibaraki Prefecture,

Japan (hereafter, "Fragmented Forest Site"), where *W. floribunda* is a dominant species.

Using DNA analyses, we assessed the impact of forest fragmentation on the role of clonal

reproduction. For comparison, we utilised data from a 6-ha permanent plot in a continuous

natural forest covering 103.51 ha (Ogawa Forest Reserve; Nakashizuka and Matsumoto,

2002; Mori et al., 2018, 2024a) approximately 100 km from the Fragmented Forest Site,

within the Ibaraki Prefecture (hereafter, "Continuous Forest Site").

We hypothesised that clonal reproduction plays a more significant role in the

Continuous Forest Site due to its low-light environment, which suppresses the

establishment of seedlings derived from seed reproduction of *W. floribunda* in the

Continuous Forest Site (Mori et al., 2020), and its spatial continuity, which facilitates

clonal growth and expansion. In contrast, seed reproduction is expected to become more

prominent in the Fragmented Forest Site, where higher light availability and spatially

constrained conditions could limit the relative contribution of clonal reproduction.

Specifically, we addressed two key questions: (1) Does clonal reproduction—quantified

by the proportion of multiple-ramet genets and clonal patch size—contribute less to *W.*

floribunda proliferation in the Fragmented Forest Site compared to the Continuous Forest

Site? (2) Could clonal reproduction have contributed to spatial expansion prior to forest

fragmentation? To address the second question, we examined whether single genets

spanned multiple fragments and whether they were predominantly located in areas

historically covered by forests. By answering these questions, this study aims to advance

our understanding of the ecological mechanisms underlying *W. floribunda* proliferation

in fragmented landscapes and to provide broader insights into the ways in which forest

fragmentation may affect liana dynamics and dominance.

MATERIALS AND METHODS

Study Sites

The study site comprises multiple fragmented forests within a 21.6-ha area located on the

campus of the University of Tsukuba (36°06'40"N, 140°06'17"E) located in the Ibaraki

Prefecture, Japan (Figs. S1; S2). The forest area in 1959 was estimated to be 15.1 ha,

representing approximately 70% of the study area (Figs. 1 and S3), based on aerial

photographs (1959–2008) provided by the Geospatial Information Authority of Japan and

Copernicus Sentinel data (2018, 2024). Forest areas were defined as regions with visible

tree canopies. Historically, these forests were likely used as traditional woodlands.

Over 65 years, forests in the region became increasingly fragmented into

multiple smaller patches (Figs. 1, 2, and S4). This fragmentation occurred due to large-

scale development during the 1960s-1980s, which significantly reduced the forest area

(Figs. 1 and S3). By 1975, the forest area had decreased to 4.6 ha (30% of the original

size). Subsequently, the forest began to recover through secondary succession and

afforestation, reaching 12.5 ha by 1998. By that time, the forest area and the number of

forest patches had reached a stable state, with no significant changes observed thereafter

(Fig. 1).

Currently, areas with frequent human access, primarily along roads, are subject

to ongoing management, including 2-3 rounds of mowing annually between May and

November (Fig. S5, hereafter referred to as "managed areas"). In contrast, most other

forest areas remain unmanaged.

Study Species

The focal species of this study is *Wisteria floribunda* (Fabaceae), a temperate deciduous

liana (Fig. S6) (Ohashi, 1989). *Wisteria floribunda* is reported to be a dominant species

with high abundance in old-growth temperate forests in the same region as the study site

(Ibaraki Prefecture, Japan; Suzuki, 2002; Mori et al., 2016a; 2024a), making it a

representative liana species in the region. This species is characterised by its relatively

large stems, with individuals reaching a maximum of approximately 40 cm in diameter

under natural conditions (Mori et al., 2018). *Wisteria floribunda* reproduces clonally

through the extension of stolons (Sakai et al., 2002), which increases the number of ramets

(clonal individuals). Genetic analyses in a 6-ha plot of the Continuous Forest Site revealed

that clonal reproduction contributed up to 71% of the total number of ramets of the study

species (Mori et al., 2018). This species is also a major problematic liana in forestry, as it

entwines around planted trees, reducing the quality of harvested timber (Suzuki, 1989).

Furthermore, many liana species native to East Asia, including *W. floribunda*, are invasive

in regions such as North America, where they pose significant ecological challenges, such

as competition with native species (Trusty et al., 2007; Leicht-Young, 2015).

Given these characteristics, focusing on *W. floribunda* offers valuable insights

that range from fundamental ecological knowledge to applied perspectives on

management and conservation in liana species. As such, *W. floribunda* serves as an ideal

model species for investigating the role of clonal reproduction in the proliferation of

lianas under forest fragmentation.

Liana Inventory

To estimate the abundance and community structure of lianas within the study site, a liana

census was conducted. The study area was divided into 20 × 20 m grid cells (Fig. S7).

Lianas attached to trees (hereafter on-tree stage) were almost absent in the managed areas,

likely due to management activities, although ground-creeping lianas (hereafter on-floor

stage) were present. Therefore, fifteen grid cells (total area: 0.6 ha) were randomly

selected within forested areas outside the managed areas (Fig. S7). Species and diameter

at breast height (DBH) of all lianas (DBH > 0 cm) attached to trees (DBH > 1 cm) in

these grid cells were recorded following Mori et al., (2024a), which is basically the same

method described in Gerwing et al., (2006) except that the diameters were measured at

1.3 m above ground. Species names followed the nomenclature of Yonekura and Kajita,

(2003).

Sampling, DNA extraction, and Genotyping

The study area, divided into 20 × 20 m grid cells as described above, was used for

sampling. One sample of the target species was randomly selected per grid cell. In the

managed areas, few on-tree lianas were present; however, ground-creeping individuals

were occasionally found. Sampling was conducted as follows: if ramets attached to trees

(DBH > 1 cm) were present, they were randomly selected for sampling. If such on-tree

ramets were not available, ramets in the on-floor stage were randomly sampled.

185   Samples were first stored at 4°C and then frozen at −30°C until DNA extraction.

DNA extraction was performed using the modified 2× CTAB (cetyltrimethylammonium

bromide) protocol (Murray and Thompson, 1980). Genotypes were determined through

PCR amplification of 10 microsatellite loci described in Mori et al., (2016b). The PCR

products were analysed using a 3130 Genetic Analyzer (Applied Biosystems, CA, USA).

Electropherograms from each marker were meticulously examined for peak patterns

using GeneMarker v1.95 (https://softgenetics.com/products/genemarker/).

Data Analysis

Identification of clones

Clones were identified based on the methods proposed by Arnaud-Haond et al., (2007).

The ability of the microsatellite markers to distinguish multilocus genotypes (MLGs) was

tested by calculating the number of MLGs for all combinations of a given locus, and the

results were then verified based on the plateaus of the genotype accumulation curve (Fig.

S8). To ascertain whether stems of the same MLG belonged to the same clone, the

probability of a given MLG occurring in a population under Hardy-Weinberg equilibrium

was calculated using the equation $p_{gen} = \prod_{i=1}^{l}(f_i)2^h$ (Parks and Werth, 1993), where

f_i is the frequency of each allele at the *i*-th locus estimated with a round-robin method,

and *h* is the number of heterozygous loci. Then, the probability of obtaining *n* repeated

MLGs from a population more than once by chance in N samples (*p_sex*) was calculated

using the equation $p_{sex} = \sum_{i=n}^{N} \frac{N!}{i!(N-i)!}[p_{gen}]^i[1-p_{gen}]^{N-i}$ (Parks and Werth 1993).

Both *p_gen* and *p_sex* were calculated using the R package "RClone" (Bailleul et al., 2016).

To distinguish each distinct MLG that belonged to a distinct clone, multilocus lineages

(MLLs) were defined based on pairwise genetic distances that have been genotyped in

this study (Fig. S9). This procedure was necessary to prevent the false detection of clones

due to slightly different MLGs resulting from somatic mutation or genotyping errors. The

pairwise genetic distance threshold was determined with the cutoff_predictor function of

the R package "poppr" (Kamvar et al., 2014, 2015). MLLs are equivalent to genets

(clones) in ecological studies, and we will use "genets" hereafter for MLLs.

Comparative analysis of liana abundance

To compare whether the abundance (stem density and basal area) observed in the

Fragmented Forest Site differs from that in the Continuous Forest Site, we performed the

following analysis using a bootstrap method. We used the liana census data of *W.*

floribunda (DBH > 0 cm) attached to trees (DBH > 1 cm), as reported by Mori et al.,

(2024a). These data were collected from a 6-ha plot (300 m × 200 m) of Continuous

Forest Site (Fig. S1). The plot was divided into a grid of 20 × 20 m cells (15 × 10 grid

cells), following the same methodology as this study. We randomly selected 15 grid cells

from the Continuous Forest Site and obtained the stem density and basal area of the study

species.

To account for the effects of random selection, we repeated this process 1,000

times using a bootstrap approach to calculate mean values and 95% confidence intervals

(CIs). We defined the abundance in the Fragmented Forest Site as significantly different

from that in the Continuous Forest Site if it fell outside the 95% CI derived from the

bootstrap for the Continuous Forest Site. In our study, random sampling avoided managed

areas, such that each selected grid cell in the Fragmented Forest Site contained at least

one *W. floribunda* individual. To ensure comparability, the bootstrap procedure for the

Continuous Forest Site also restricted the selection to grid cells that contained at least one

W. floribunda individual.

We then fitted generalised linear models (GLMs) for each bootstrap iteration to

estimate the magnitude of differences in abundance between the Fragmented Forest

and the Continuous Forest Site. Stem density and basal area were used as the response

variables, with site (fragmented or continuous) as the explanatory variable. Stem density

was modelled using a Poisson distribution, whereas basal area was modelled using a

normal distribution. Residual diagnostics via the R package "DHARMa" (Florian, 2022)

revealed non-uniformity and overdispersion ($P < 0.001$) for the Poisson GLM; hence, we

employed a Poisson generalised linear mixed model (GLMM) with grid cell as a random

effect.

Comparative analysis of contribution of clonal reproduction

To test whether the contribution of clonal reproduction observed in the Fragmented Forest

Site differs from that in the Continuous Forest Site, we conducted a bootstrap analysis

using two metrics: (1) the percentage of ramets derived from clonal reproduction (%

Clonal ramet), calculated as the proportion of clonal ramets (multiple-ramet genets)

relative to the total number of ramets, and (2) the percentage of genet area (% Genet area),

calculated as the proportion of the total study area covered by genets. For comparison

between the two sites, both metrics were calculated based on liana individuals in the on-

tree stage. We used clonal structure data for *W. floribunda* from the Continuous Forest

Site (the 6-ha plot), as reported by Mori et al., (2018), which included all *W. floribunda*

individuals (DBH > 0 cm) attached to host trees (DBH > 5 cm) throughout the 6-ha area.

To ensure comparability, the site was divided into 20×20 m grid cells (15×10 grid cells),

and one sample was randomly selected from each grid cell. The contribution of clonal

reproduction was calculated for each site, and the bootstrap method was used to repeat

this process 1,000 times, providing mean values and 95% confidence intervals (CIs). The

observed values from the Fragmented Forest Site were considered significantly different

from those in continuous natural forests if they fell outside the 95% CI derived from the

Continuous Forest Site.

Although the minimum host tree size thresholds differ between the Fragmented

Forest Site (DBH > 1 cm) and the Continuous Forest Site (DBH > 5 cm), a previous

census of *W. floribunda* in the Continuous Forest Site (Mori et al., 2024a) suggests that

this discrepancy has minimal impact on our comparison. In that census, 93.4% of

individuals attached to smaller host trees (DBH 1–5 cm) co-occurred in the same 20 × 20

m grid cells as those on larger trees (DBH > 5 cm). Moreover, individuals on larger trees

appeared in 13 additional grid cells, indicating a broader spatial distribution. The pairwise

nearest-neighbour distance between *W. floribunda* on smaller vs. larger host trees

averaged 5.2 ± 4.4 m, with 95% of occurrences within 14.8 m. These findings (Fig. S9)

demonstrate that, despite using different size thresholds, the spatial overlap is sufficient

to ensure that the lower threshold of 1 cm in the Fragmented Forest Site has minimal

impact on our comparative analysis.

To compare differences in proportions (% Clonal ramet and % Genet area)

between the sites, we used odds ratios, defined as $\{p/(1\text{-}p)\}/\{q/(1\text{-}q)\}$, where p represents

the proportion in the Fragmented Forest Site and q represents the proportion in the

Continuous Forest Site. An odds ratio of 1 indicates no difference in the odds of clonal

reproduction contribution between the two sites. If the 95% CI of the odds ratio was

significantly greater than 1, it indicated higher odds in the Fragmented Forest Site,

whereas a 95% CI significantly below 1 indicated higher odds in the Continuous Forest

Site. The mean odds ratio and its 95% CI were calculated using the bootstrap method.

Previous research in the Continuous Forest Site suggested that large genets of *W.*

floribunda (genets with N ≥ 3 ramets) exhibit mutually exclusive distribution patterns

(Mori et al., 2018). To test whether the genet distribution patterns observed under natural

conditions in the Continuous Forest Site differ from those in the Fragmented Forest Site,

we evaluated the distribution exclusivity in both sites. Large genets (N ≥ 3 ramets) were

randomly placed within the study area, and the total overlapping area of their convex hull

polygons was calculated using the R package "sf" (Edzer 2018; Edzer and Roger 2023).

This process was repeated 1,000 times for each site to obtain 95% CIs. If the observed

value fell significantly below the 95% CI, genets were considered significantly mutually

exclusive, while values above the 95% CI indicated significantly overlapping

distributions.

Distribution of genets across fragmented forests

To examine whether genets expanded their distribution solely through clonal reproduction

after forest fragmentation or whether some genets had already expanded their distribution

before fragmentation, we asked the following questions: (1) Do genets occur in multiple

fragmented forest patches? (2) If so, are such genets disproportionately distributed in

areas covered by forest in 1959, prior to fragmentation? These questions were examined

as follows:

Fragmented forest patches were defined as forest units isolated by roads or other

artificial structures where forest canopies no longer connect, based on aerial photographs

(Fig. S4). If a genet was found exclusively within a single forest patch, it was defined as

having expanded its distribution through clonal reproduction within that patch after

fragmentation. To test whether genets occurring in multiple patches were

disproportionately distributed in areas covered by forest before fragmentation, we

compared their distribution to that of genets confined to a single patch using Fisher's

exact test.

If genets were distributed across multiple forest patches and were significantly

more likely to occur in areas covered by forest before fragmentation, the present study

considered that these genets had most likely expanded their distribution through clonal

reproduction before fragmentation. All analyses were performed using R version 4.3.2 (R

Core Team, 2023).

RESULTS

Stem Density and Basal Area

A total of 9 liana species were recorded during the census conducted in the 0.6-ha (15-

grid cell) area, comprising 573 stems and a basal area of 0.287 m² (Table S1). Among

these, *W. floribunda* was the most abundant species, with 393 stems and a basal area of

0.258 m². This accounted for 69% of the total stem density and 90% of the total basal area

of all liana species. The next most abundant species in terms of stem density and basal

area were *Parthenocissus tricuspidata* (Vitaceae; 75 stems) and *Toxicodendron orientale*

subsp. *orientale* (Anacardiaceae; 0.012 m²), which corresponded to 13% and 4% of the

total stem density and basal area, respectively. These results indicate that *W. floribunda*

is the dominant species, contributing the majority of liana abundance in these fragmented

forests.

In the Continuous Forest Site, subsampling over a comparable area of 0.6 ha (15-

grid cells) yielded a mean stem density of *W. floribunda* of 126.1 (95% CI: 75.0–192.0)

and a basal area of 0.282 m² (95% CI: 0.106–0.504) (Fig. 3). Compared to the Continuous

Forest Site, the Fragmented Forest Site exhibited a statistically significant higher stem

count of *W. floribunda*, with an estimated site effect coefficient mean of 1.14,

corresponding to exp(1.14) = 3.12 times higher stem density (95% CI: 2.07–4.86, $P <$

0.001). In contrast, the difference in basal area between the sites was not statistically

significant (mean estimated site effect coefficient = -0.003, 95% CI: -0.019–0.009, $P =$

0.51).

Clonal Structure in Fragmented Forests

Genetic analysis revealed that the 273 ramets belonged to a total of 47 genets (Fig. 4).

Among these, 257 ramets (94% of all ramets) were part of multiple-ramet genets, which

are definitively derived from clonal reproduction. The probability of the study species

obtaining a given genotype ($p_{gen} < 0.001$) or obtaining repeated genotypes that originated

from distinct sexual reproductive events by chance ($p_{sex} < 0.001$) were low; thus, it was

assumed that errors in the identification of clones were unlikely. When one ramet was

excluded from each multi-ramet genet (under the assumption that each multi-ramet genet

contains a single ramet derived from seed reproduction), the proportion remained 93%.

Of the 273 ramets, 231 were in the on-tree stage and 42 in the on-floor stage. Among

these, 94% of on-tree ramets (217/231) and 95% of on-floor ramets (40/42) were part of

multiple-ramet genets, with no significant difference between the two groups (Fisher's

Exact Test; Odds ratio: 1.29; 95% CI: 0.279–12.12; $P = 1.00$). This indicates that the

proportion of ramets derived from clonal reproduction does not differ significantly

between growth stages (on-tree versus on-floor).

The area of the 47 genets estimated using convex hull polygons was 12.5 ha,

accounting for 58% of the total study area (21.6 ha). When considering only on-tree

ramets, the genet area was 12.3 ha, equivalent to 57% of the study area. The largest genet

(Genet ID = 1; Fig. 4) consisted of 105 ramets (39% of all ramets) and occupied an area

of 9.5 ha, which accounted for 77% of the total genet area (Fig. 5).

Comparison of Contribution of Clonal Reproduction

The proportion of on-tree ramets derived from clonal reproduction (% Clonal ramet) was

94% in the Fragmented Forest Site, compared to 45% (95% CI: 38–52%) in the

Continuous Forest Site (Fig. 6). The odds ratio was 18.2 (95% CI: 13.2–24.0), indicating

that the proportion was significantly higher in the Fragmented Forest Site, by 49% (odds

18.2 times greater), compared to the Continuous Forest Site.

The proportion of the study area covered by on-tree genets (% Genet area) was

57% in the Fragmented Forest Site, compared to 7% (95% CI: 4–10%) in the Continuous

Forest Site, with an odds ratio of 18.6 (95% CI: 11.9–31.6) (Fig. 6). This indicates that

the proportion was significantly higher in the Fragmented Forest Site, by 50% (odds 18.6

times greater), compared to the Continuous Forest Site.

The total area of overlapping genets (N ramet ≥ 3) was 0.46 ha in the Fragmented

Forest Site and 0.05 ha in the Continuous Forest Site (Fig. 5). In both sites, the observed

values were significantly smaller than their respective 95% CIs (0.96–4.97 for the

Fragmented Forest Site and 0.06–0.23 for the Continuous Forest Site). This indicates that

large genets (N ramet ≥ 3) exhibited mutually exclusive distribution patterns in both sites.

Distribution of Genets Across Fragmented Forests

Three genets (Genet IDs: 1, 9, and 10; Fig. 4) were distributed across multiple fragmented

forest patches, while the remaining 44 genets (94% of all genets) were found exclusively

within a single forest patch. The total number of ramets belonging to these three genets

was 117, accounting for 43% of all ramets. Among the nine forest patches where *W.*

floribunda was present, the largest genet (Genet ID = 1) spanned seven patches, while the

other two genets were each distributed across two different patches (Fig. 4).

382   Of the 117 ramets belonging to genets distributed across multiple forest patches,

10 were located in areas not covered by forest prior to fragmentation (6.5 ha), while 107

were in areas covered by forest (15.1 ha). In contrast, among the 156 ramets confined to

a single forest patch, 61 were in areas not covered by forest, and 95 were in areas covered

by forest. The proportion of ramets in areas covered by forest was significantly higher for

genets distributed across multiple patches (91%) compared to those confined to a single

patch (61%) (Fisher's Exact Test; Odds ratio: 6.83; 95% CI: 3.24−15.8; $P < 0.001$).

DISCUSSION

We initially hypothesised that clonal reproduction would be more prominent in the

Continuous Forest Site, whereas seed reproduction would prevail in the Fragmented

Forest Site, due to the differences in light conditions and spatial continuity. Contrary to

this expectation, our genetic analyses revealed that clonal reproduction played a

substantial role in liana proliferation, with large clonal patches evident throughout the

fragmented forests. Moreover, these clonal expansions appeared to have been established

even before forest fragmentation, suggesting that both current and historical clonal

structures underpin the species' proliferation in these landscapes. In the following

sections, we discuss how these findings refine our understanding of liana species'

ecological strategies in response to fragmentation, and consider broader implications for

liana dominance in human-altered forests.

Liana Abundance in Fragmented Forests

In the Fragmented Forest Site, stem density was approximately three times higher. This

is likely due to substantial anthropogenic disturbances that altered forest structure,

increased edge effects, and the presence of young forest patches within the Fragmented

Forest Site. Such increases in liana abundance following forest fragmentation have been

widely reported in temperate (Ladwig and Meiners, 2015) and tropical forests (Laurance

et al., 2001; Jones et al., 2017). The elevated stem density of the study species observed

here is consistent with these findings. Genetic analyses further suggest that this high stem

density in fragmented forests is primarily due to clonal reproduction, which enables rapid

population expansion under forest disturbance.

Despite the higher stem density, the maximum stem diameter of the study species

in the Fragmented Forest Site was about 14 cm (Table S1), whereas stems reaching 40

cm have been documented in the Continuous Forest Site (Mori et al., 2018). This

difference in stem diameter likely reflects variations in forest structure (e.g., tree density)

and successional stage. Although the maximum diameter in the Fragmented Forest Site is

smaller, the overall basal area of lianas remains comparable to that of the Continuous

Forest Site, presumably due to their greater density. Larger lianas are more common in

non-fragmented forests, consistent with findings in tropical forests (Campbell et al., 2018).

However, as succession progresses in the Fragmented Forest Site, it is plausible that lianas

already reaching the canopy will continue to grow, potentially leading to further increases

in the basal area (Ladwig and Meiners, 2010).

Role of Clonal Reproduction in Fragmented Forests

This study hypothesised that in fragmented forests, where edge effects increase light

availability and fragmentation reduces spatial continuity, seed reproduction may play a

more prominent role compared to continuous forests. However, the proportion of clonal

ramets and the genet area in fragmented forests were significantly higher than in

continuous forests (both approximately 50%, with an odds ratio of 18), which indicated

that the proliferation of lianas in fragmented forests cannot be fully explained by

improved light conditions and spatial continuity alone, but is likely driven by other factors

related to large-scale disturbances that enhance clonal growth. One possible explanation

is the inherent characteristics of clonal plants, which allow them to share resources such

as water and nutrients through stolons or rhizomes, enabling rapid recovery from partial

damage and facilitating extensive distribution (Fahrig et al., 1994; Winkler and Fischer,

2002; Saitoh et al., 2006). Moreover, previous studies have shown that improved light

conditions alone cannot account for the abundance of lianas in temperate forests

(Carrasco-Urra and Gianoli, 2009; Mori et al., 2024a). Instead, factors such as clonal

reproduction and host tree availability may play a more critical role in their proliferation.

The exclusive distribution patterns of genets observed in fragmented forests

closely resemble those found under natural conditions in continuous old-growth forests.

This suggests that the clonal spread of genets operates in a similar way in both fragmented

and non-fragmented forests. Clonal reproduction likely promotes the dominance of

certain genets by enabling them to occupy suitable habitats in forest patches, potentially

suppressing the spread of others (Schmid and Harper, 1985; van Groenendael et al., 1996).

As structural parasites, lianas rely on external biomechanical support (i.e., host trees) as

an essential habitat for their growth and survival (Putz and Holbrook, 1991; Sakai and

Suzuki, 1999). Such exclusivity in the distribution of the studied genets may therefore be

a consequence of competition among genets for suitable host trees. Studies in continuous

forest sites (Mori et al., 2016a, 2024a) have shown that lianas, including *W. floribunda*,

exhibit a strong preference for host tree size, while small-scale natural disturbances (e.g.,

canopy gaps caused by one or a few fallen trees) and landforms play only a minor role in

their distribution. Specifically, stem-twining and tendril-climbing lianas are more likely

to establish on small host trees when their stems are small and shift to larger host trees as

they grow. These findings highlight that host-tree availability—and the resulting

competition for host trees—strongly shapes the spatial distribution of lianas by

determining their ability to establish and grow on suitable hosts. This supports our

hypothesis that clonal reproduction and genet exclusivity are tightly linked to host-tree

competition.

Clonal Reproduction Before and After Forest Fragmentation

The proliferation of lianas through clonal reproduction in fragmented forests reflects the

combined influences of both pre- and post-fragmentation processes. Our results indicate

that most genets were confined to single forest patches, suggesting that the expansion of

genet distribution primarily occurred after forest fragmentation. This may be because

clonal offspring face difficulties migrating to fragmented patches (Fukui and Araki, 2014).

However, three specific genets were found across multiple forest patches and were

significantly more abundant in areas historically covered by forest before fragmentation.

This pattern indicated that these genets had already expanded across multiple patches

prior to fragmentation. Notably, these three genets accounted for 43% of all recorded

ramets, highlighting the substantial role of pre-fragmentation clonal structures in driving

the pronounced proliferation of lianas in fragmented forests.

Increased liana abundance following forest fragmentation is a well-known

phenomenon in temperate and tropical forests (Schnitzer and Bongers, 2011; Campbell

et al., 2015; Ladwig and Meiners, 2015). However, our findings suggest that a key factor

underlying this trend may lie in the clonal structures established prior to fragmentation.

As discussed earlier, the adaptive traits and life-history strategies of clonal plants enable

them to thrive under disturbance and environmental change, facilitating the proliferation

of these clonal species in fragmented landscapes (Fahrig et al., 1994; Winkler and Fischer,

2002; Saitoh et al., 2006). It is likely that a small number of successful genets,

characterised by high adaptability or favourable traits, drive the post-fragmentation

proliferation of liana species in response to altered environmental conditions. Indeed, in

warm temperate forests of Japan, the dominant liana *Trachelospermum asiaticum* var.

asiaticum (Apocynaceae) was reported to establish a few genets via seed dispersal during

early successional stages. These genets rapidly expanded through clonal reproduction in

forests recovering from large-scale natural disturbances caused by volcanic eruptions

(Mori et al., 2024b).

Implications of Clonal Reproduction for Liana Management

Wisteria floribunda is a problematic liana species requiring management and control in

forestry (Suzuki, 1989). At our study site, small pre-climbing ramets were found even in

managed areas subjected to multiple rounds of mowing annually. The present study found

that most of these ramets were derived from clonal reproduction. Mature *W. floribunda*

individuals are known to expand their distribution by producing numerous ramets through

stolon growth (Sakai et al., 2002; Mori et al., 2021). However, in managed areas, pre-

climbing small ramets were observed even in the absence of mature individuals in the on-

tree stage. This finding suggests that clones established during periods when the forest

was continuous have persisted underground and remained viable despite repeated

mowing. This finding aligns with previous studies on kudzu (*Pueraria montana* var.

lobata) in North America, which reported that mowing enhances clonal diversity

(Kartzinel et al., 2015), and on *Clematis vitalba* in New Zealand, which reported the

ability to regenerate from severed stems (Jarvis-Lowry et al., 2024). These results suggest

that for lianas with strong clonal growth capabilities, conventional mowing and liana

cutting practices may be insufficient to effectively suppress the number of genets. This

highlights the need for a precise understanding of the life history strategies of lianas at

the species level to effectively manage and conserve them in forests.

CONCLUSIONS

This study elucidates the impact of forest fragmentation on liana proliferation through the

lens of clonal reproduction. Our findings strongly suggest that clonal reproduction serves

as a key mechanism driving the spatial expansion and dominance of lianas in fragmented

forests. Additionally, the distribution patterns of large genets indicate that the proliferation

of lianas following fragmentation heavily depends on pre-existing clonal structures

established before fragmentation. These results clarify that forest fragmentation not only

promotes new clonal growth but also highlights the significant influence of historical

clonal structures on the current distribution and abundance of genets. Furthermore, this

study suggests that the proliferation of lianas in fragmented forests may be driven by a

limited number of successful genets belonging to a few clonally dominant species,

highlighting the critical role of species-specific reproductive strategies and life histories

in facilitating the dynamic population expansion of lianas following forest fragmentation.

This study also underscored the necessity of considering clonality and life history

strategies at the species level when developing management and conservation approaches

for lianas. Overall, the findings of this study indicate that lianas, due to their sensitivity

to environmental changes, serve as valuable ecological indicators (Schnitzer, 2018) for

assessing the impacts of human-induced disturbances, such as deforestation and habitat

fragmentation. While this study provides significant insights, it is limited to a single study

site. Future research with similar designs across multiple sites is necessary to generalise

these findings and gain a more comprehensive understanding of how forest fragmentation

influences liana proliferation.

ACKNOWLEDGEMENTS

This study was supported by JSPS KAKENHI (grant numbers 21K14882; 24K01802).

LITERATURE CITED

Arnaud-Haond S, Duarte CM, Alberto F, Serrao EA. 2007. Standardizing methods to

address clonality in population studies. *Molecular Ecology*, *16*(24), 5115–5139.

https://doi.org/10.1111/j.1365-294X.2007.03535.x

Bailleul D, Stoeckel S, Arnaud-Haond S, Poisot T. 2016. RClone: a package to identify

MultiLocus Clonal Lineages and handle clonal data sets in R. *Methods in Ecology*

and Evolution, *7*(8), 966–970. https://doi.org/10.1111/2041-210x.12550

Campbell M, Laurance WF, Magrach A. (2015). Ecological effects of lianas in

fragmented forests. In Schnitzer SA, Bongers F, Burnham RJ, Putz FE (Eds.),

Ecology of Lianas (pp. 443-450). John Wiley & Sons, Ltd.

https://doi.org/10.1002/9781118392409.ch29

Campbell MJ, Edwards W, Magrach A et al. 2018. Edge disturbance drives liana

abundance increase and alteration of liana-host tree interactions in tropical forest

fragments. *Ecol Evol*, *8*(8), 4237–4251. https://doi.org/10.1002/ece3.3959

Carrasco-Urra F, Gianoli E. 2009. Abundance of climbing plants in a southern

temperate rain forest: host tree characteristics or light availability? *Journal of*

Vegetation Science, *20*(6), 1155–1162. https://doi.org/10.1111/j.1654-

1103.2009.01115.x

Edzer P. 2018. Simple Features for R: Standardized Support for Spatial Vector Data. *The*

R Journal, *10*(1), 439–446. https://doi.org/10.32614/RJ-2018-009

Edzer P, Roger B. 2023. *Spatial Data Science: With applications in R*. Chapman and

Hall/CRC. https://doi.org/10.1201/9780429459016

Fahrig L. 2003. Effects of Habitat Fragmentation on Biodiversity. *Annual Review of*

Ecology, Evolution, and Systematics, *34*(1), 487–515.

https://doi.org/10.1146/annurev.ecolsys.34.011802.132419

Fahrig L, Coffin DP, Lauenroth WK, Shugart HH. 1994. The advantage of long-distance

clonal spreading in highly disturbed habitats. *Evolutionary Ecology*, *8*(2), 172–

187. https://doi.org/10.1007/bf01238248

Florian H. 2022. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level /

Mixed) Regression Models. https://CRAN.R-project.org/package=DHARMa

Fukui S, Araki KS. 2014. Spatial niche facilitates clonal reproduction in seed plants

under temporal disturbance. *PLoS One*, *9*(12), e116111.

https://doi.org/10.1371/journal.pone.0116111

GBIF Secretariat. (2023). Wisteria floribunda *(Willd.) DC. GBIF Backbone Taxonomy.*

Checklist dataset. https://doi.org/10.15468/39omei accessed via GBIF.org on

2025-01-21.

Gerwing JJ, Schnitzer SA, Burnham RJ et al. 2006. A Standard Protocol for Liana

Censuses. *Biotropica*, *38*(2), 256–261. https://doi.org/10.1111/j.1744-

7429.2006.00134.x

Haddad NM, Brudvig LA, Clobert J et al. 2015. Habitat fragmentation and its lasting

impact on Earth's ecosystems. *Sci Adv*, *1*(2), e1500052.

https://doi.org/10.1126/sciadv.1500052

Hansen MC, Potapov PV, Moore R et al. 2013. High-resolution global maps of 21st-

century forest cover change. *Science*, *342*(6160), 850–853.

https://doi.org/10.1126/science.1244693

Hiebeler D. 2004. Competition between near and far dispersers in spatially structured

habitats. *Theor Popul Biol*, *66*(3), 205–218.

https://doi.org/10.1016/j.tpb.2004.06.004

Jarvis-Lowry B, Harrington KC, Ghanizadeh H, Robertson AW. 2024. Potential

importance of vegetative spread and fragment regeneration for invasiveness of

Clematis vitalba. *Weed Research*, *64*(4), 282–289.

https://doi.org/10.1111/wre.12633

Kamvar ZN, Brooks JC, Grünwald NJ. 2015. Novel R tools for analysis of genome-

wide population genetic data with emphasis on clonality. *Front Genet*, *6*, 208.

https://doi.org/10.3389/fgene.2015.00208

Kamvar ZN, Tabima JF, Grünwald NJ. 2014. Poppr: an R package for genetic analysis

of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, *2*,

e281. https://doi.org/10.7717/peerj.281

Kartzinel TR, Hamrick JL, Wang C, Bowsher AW, Quigley BG. 2015. Heterogeneity of

clonal patterns among patches of kudzu, Pueraria montana var. lobata, an invasive

plant. *Annals of Botany*, *116*(5), 739–750. https://doi.org/10.1093/aob/mcv117

Kiesewetter KN, Otano L, Afkhami ME. 2023. Fragmentation disrupts microbial effects

on native plant community productivity. *Journal of Ecology*, *111*(6), 1292–1307.

https://doi.org/10.1111/1365-2745.14097

Ladwig LM, Meiners SJ. 2010. Spatiotemporal dynamics of lianas during 50 years of

succession to temperate forest. *Ecology*, *91*(3), 671–680.

Ladwig LM, Meiners SJ. 2015. The role of lianas in temperate tree communities. In

Schnitzer SA, Bongers F, Burnham RJ, Putz FE (Eds.), *Ecology of Lianas* (pp.

188-202). Wiley. https://doi.org/10.1002/9781118392409.ch15

Laurance WF, Pérez-Salicrup D, Delamônica P et al. 2001. RAIN FOREST

FRAGMENTATION AND THE STRUCTURE OF AMAZONIAN LIANA

COMMUNITIES. *Ecology*, *82*(1), 105–116. https://doi.org/10.1890/0012-

9658(2001)082[0105:rffats]2.0.co;2

Ledo A, Schnitzer SA. 2014. Disturbance and clonal reproduction determine liana

distribution and maintain liana diversity in a tropical forest. *Ecology*, *95*(8), 2169–

2178.

Leicht-Young SA, Pavlovic NB. 2015. Lianas as invasive species in North America. In

Schnitzer SA, Bongers F, Burnham RJ, Putz FE (Eds.), *Ecology of Lianas* (pp.

427-442). Wiley. https://doi.org/10.1002/9781118392409.ch28

Li D, Yang Y, Xia F, Sun W, Li X, Xie Y. 2022. Exploring the influences of different

processes of habitat fragmentation on ecosystem services. *Landscape and Urban*

Planning, *227*, 104544. https://doi.org/10.1016/j.landurbplan.2022.104544

Liu J, Wilson M, Hu G, Liu J, Wu J, Yu M. 2018. How does habitat fragmentation affect

the biodiversity and ecosystem functioning relationship? *Landscape Ecology*,

33(3), 341–352. https://doi.org/10.1007/s10980-018-0620-5

Londré RA, Schnitzer SA. 2006. The distribution of lianas and their change in

abundance in temperate forests over the past 45 years. *Ecology*, *87*(12), 2973–

2978. https://doi.org/10.1890/0012-9658(2006)87[2973:TDOLAT]2.0.CO;2

Mori H, Kamijo T, Masaki T. 2016a. Liana distribution and community structure in an

old-growth temperate forest: the relative importance of past disturbances, host

trees, and microsite characteristics. *Plant Ecology*, *217*(10), 1171–1182.

https://doi.org/10.1007/s11258-016-0641-6

Mori H, Ueno S, Matsumoto A et al. 2016b. Development and characterization of 10

microsatellite markers from *Wisteria floribunda* (Fabaceae). *Silvae Genetica*,

65(1), 55–58. https://doi.org/10.1515/sg-2016-0007

Mori H, Ueno S, Matsumoto A, Kamijo T, Tsumura Y, Masaki T. 2018. Large

contribution of clonal reproduction to the distribution of deciduous liana species

(*Wisteria floribunda*) in an old-growth cool temperate forest: evidence from

genetic analysis. *Annals of Botany*, *121*(2), 359–336.

https://doi.org/10.1093/aob/mcx153

Mori H, Masaki T, Tsunamoto Y, Naoe S. 2020. Survival rate and environmental

response of current-year seedlings of the temperate liana *Wisteria floribunda*

across a heterogeneous environment. *Journal of Plant Research*, *133*, 193–203.

https://doi.org/10.1007/s10265-019-01163-1

Mori H, Ueno S, Kamijo T, Tsumura Y, Masaki T. 2021. Interspecific variation in

clonality in temperate lianas revealed by genetic analysis: Do clonal proliferation

processes differ among lianas. *Plant Species Biology*, *36*(4), 578–588.

https://doi.org/10.1111/1442-1984.12348

Mori H, Kamijo T, Oguro M, Masaki T. 2024a. Liana distribution and small-scale

natural disturbance in an old-growth temperate forest. *European Journal of Forest*

Research. https://doi.org/10.1007/s10342-024-01741-6

Mori H, Kamijo T. 2024b. Clonal reproduction as a driver of liana proliferation

following large-scale disturbances in temperate forests. *bioRxiv*.

https://doi.org/10.1101/2024.08.21.609080

Murray MG, Thompson WF. 1980. Rapid isolation of high molecular weight plant

DNA. *Nucleic Acids Res*, *8*(19), 4321–4325. https://doi.org/10.1093/nar/8.19.4321

Nakashizuka T, Matsumoto Y. 2002. *Diversity and interaction in a temperate forest*

community: Ogawa Forest Reserve of Japan. Springer Japan.

Ngute ASK, Schoeman DS, Pfeifer M et al. 2024. Global dominance of lianas over trees

is driven by forest disturbance, climate and topography. *Glob Chang Biol*, *30*(1),

e17140. https://doi.org/10.1111/gcb.17140

Ohashi H. 1989. Wild Flowers of Japan: woody plants I. Tokyo: Heibonsha, (in

Japanese) Wisteria, p. 247--248. In Satake Y, Hara H, Watari S, Tominari T (Eds.).

Parks JC, Werth CR. 1993. A study of spatial features of clones in a population of

bracken fern, Pteridium aquilinum (Dennstaedtiaceae). *American Journal of Botany*, *80*, 537–544.

Putz FE. 1984. The natural history of lianas on Barro Colorado Island, Panama. *Ecology*, *65*(6), 1713–1724. https://doi.org/https://doi.org/10.2307/1937767

Putz FE, Holbrook NM. 1991. Biomechanical studies of vines. In Putz FE, Mooney HA (Eds.), *The Biology of Vines* (pp. 73-98). Cambridge University Press. https://doi.org/10.1017/CBO9780511897658.005

R Core Team. 2023. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. https://www.R-project.org/

Saitoh T, Seiwa K, Nishiwaki A. 2006. Effects of resource heterogeneity on nitrogen translocation within clonal fragments of Sasa palmata: an isotopic (15N) assessment. *Annals of Botany*, *98*(3), 657–663. https://doi.org/10.1093/aob/mcl147

Sakai A, Nomiya H, Suzuki W. 2002. Horizontal distribution of stolons of a temperate liana *Wisteria floribunda* DC. and its ecological significance. *Journal of Forest Research*, *7*(3), 125–130.

Sakai A, Suzuki W. 1999. Effect of support on the growth of a woody vine, *Wisteria floribunda* DC. (Leguminosae). *Journal of Forest Research*, *4*(2), 183–186.

Schmid B, Harper JL. 1985. Clonal Growth in Grassland Perennials: I. Density and Pattern-Dependent Competition Between Plants with Different Growth Forms. *The Journal of Ecology*, *73*(3), 793. https://doi.org/10.2307/2260147

Schnitzer SA, Bongers F. 2011. Increasing liana abundance and biomass in tropical forests: emerging patterns and putative mechanisms. *Ecol Lett*, *14*(4), 397–406. https://doi.org/10.1111/j.1461-0248.2011.01590.x

Schnitzer SA. 2018. Testing ecological theory with lianas. *New Phytologist*, *220*(2),

366–380. https://doi.org/10.1111/nph.15431

Schnitzer SA, DeFilippis DM, Visser M et al. 2021. Local canopy disturbance as an

explanation for long-term increases in liana abundance. *Ecol Lett*, *24*(12), 2635–

2647. https://doi.org/10.1111/ele.13881

Suzuki W. 1989. Tree damage caused by climbing plants and the mechanism of their

development in Hinoki (*Chamaecyparis obtusa*) plantations (in Japanese with

English summary). *Journal of the Japanese Forestry Society*, *71*(10), 395–404.

https://doi.org/10.11519/jjfs1953.71.10_395

Suzuki W. 2002. Forest Vegetation in and Around Ogawa Forest Reserve in Relation to

Human Impact. In Nakashizuka T, Matsumoto Y (Eds.), *Diversity and Interaction*

in a Temperate Forest Community: Ogawa Forest Reserve of Japan (pp. 27-41).

Springer. https://doi.org/10.1007/978-4-431-67879-3¥_4

Travis JMJ, Dytham C. 1998. The evolution of dispersal in a metapopulation: a spatially

explicit, individual-based model. *Proceedings of the Royal Society of London.*

Series B: Biological Sciences, *265*(1390), 17–23.

https://doi.org/10.1098/rspb.1998.0258

Trusty JL, Goertzen LR, Zipperer WC, Lockaby BG. 2007. Invasive *Wisteria* in the

Southeastern United States: genetic diversity, hybridization and the role of urban

centers. *Urban Ecosystems*, *10*(4), 379–395. https://doi.org/10.1007/s11252-007-

0030-y

van Groenendael JM, Klimeš L, Klimešová J, Hendriks RJJ. 1996. Comparative

ecology of clonal plants. *Philosophical Transactions of the Royal Society of*

London. Series B: Biological Sciences, *351*(1345), 1331–1339.

707   https://doi.org/10.1098/rstb.1996.0116

708 Wilson MC, Chen X-Y, Corlett RT et al. 2016. Habitat fragmentation and biodiversity

709   conservation: key findings and future challenges. *Landscape Ecology*, *31*(2), 219–

710   227. https://doi.org/10.1007/s10980-015-0312-3

711 Winkler E, Fischer M. 2001. The role of vegetative spread and seed dispersal for

712   optimal life histories of clonal plants: a simulation study. *Evolutionary Ecology*,

713   *15*(4-6), 281–301. https://doi.org/10.1023/a:1016080714200

714 Yonekura K, Kajita T. 2003. BG Plants: Japanese name-scientific name index (YList).

715   http://ylist.info

716 Yorke SR, Schnitzer SA, Mascaro J, Letcher SG, Carson WP. 2013. Increasing liana

717   abundance and basal area in a tropical forest: the contribution of long-distance

718   clonal colonization. *Biotropica*, *45*(3), 317–324.

719 Yuan C-M. 2024. Liana regeneration in the treefall gaps, at the gap edges and in the

720   forest interior of a subtropical montane moist evergreen broad-leaved forest in the

721   Ailao Mountains, Southwestern China. *Tropical Ecology*, *65*(1), 133–141.

722   https://doi.org/10.1007/s42965-023-00319-4

FIGURE CAPTIONS

Figure 1. Temporal changes in forest area and the number of forest patches over 65 years

at the study site following forest fragmentation. Forest area and the number of patches

were estimated from aerial photographs (1959-2008) and satellite images (2018-2024).

Figure 2. Approximate forest patch age (in years) of the fragmented forests in the study

site, estimated from the temporal changes in forest area over 65 years.

Figure 3. Stem density and basal area of the study species in 15 quadrats (20 x 20 m grid

cells; 0.6 ha) of the Fragmented Forest Site (this study; vertical red line) and Continuous

Forest Site (Mori et al. 2024a; shaded bars). Stem density and basal area at the Continuous

Forest Site were obtained by subsampling 15 quadrats with 1000 bootstraps.

Figure 4. Clonal structure of the study species in fragmented forests of the study site.

Each number indicate ramets; same numbers indicate genetically identical individuals

(i.e., genets/clones). Orange polygons with solid lines indicate genet area obtained as

convex hull polygons. Filled green background represent area currently covered by forest.

Figure 5. Distribution of genets in the Fragmented Forest Site (this study) and in the

Continuous Forest Site (Mori et al. 2018), shown at the same spatial scale. Genets are

shown as convex hull polygons following Figure 4. For visibility purposes, clonal ramets

distributed within the genet polygons are not shown in this figure.

Figure 6. Proportion of clonal ramets to the total ramets (% Clonal ramets; upper panel)

and proportion of genet area to the total study area (% Genet area; lower panel) of the

study species in the Fragmented Forest Site (this study; vertical red line) and Continuous

Forest Site (Mori et al. 2018; shaded bars). Proportion of clonal ramets and genet area in

the Continuous Forest Site were obtained by subsampling a sample within each 20 x 20

m grid cell with 1000 bootstraps.

APPENDIX

Figure S1. Map of the study sites.

Figure S2. Photographs of the Fragmented Forest Site.

Figure S3. Maps of forest area in the Fragmented Forest Site**.**

Figure S4. Fragmented forest patches in the study site.

Figure S5. Managed zones in the Fragmented Forest Site.

Figure S6. Photographs of the study species.

Figure S7. 20 x 20 m grid cells used in this study.

Figure S8. Genotype accumulation curve.

Figure S9. Pairwise genetic distance.

Figure S10. Distribution map of the study species in the Continuous Forest Site.

Table S1. Liana community structure in the Fragmented Forest Site.

Figure 1. Temporal changes in forest area and the number of forest patches over 65 years

at the study site following forest fragmentation. Forest area and the number of patches

were estimated from aerial photographs (1959-2008) and satellite images (2018-2024).

Figure 2. Approximate forest patch age (in years) of the fragmented forests in the study

site, estimated from the temporal changes in forest area over 65 years.

Figure 3. Stem density and basal area of the study species in 15 quadrats (20 x 20 m grid

cells; 0.6 ha) of the Fragmented Forest Site (this study; vertical red line) and Continuous

Forest Site (Mori et al. 2024a; shaded bars). Stem density and basal area at the Continuous

Forest Site were obtained by subsampling 15 quadrats with 1000 bootstraps.

 Figure 4. Clonal structure of the study species in fragmented forests of the study site.

 Each number indicate ramets; same numbers indicate genetically identical individuals

 (i.e., genets/clones). Orange polygons with solid lines indicate genet area obtained as

 convex hull polygons. Filled green background represent area currently covered by forest.

Figure 5. Distribution of genets in the Fragmented Forest Site (this study) and in the

Continuous Forest Site (Mori et al. 2018), shown at the same spatial scale. Genets are

shown as convex hull polygons following Figure 4. For visibility purposes, clonal ramets

distributed within the genet polygons are not shown in this figure.

Figure 6. Proportion of clonal ramets to the total ramets (% Clonal ramets; upper panel)

and proportion of genet area to the total study area (% Genet area; lower panel) of the

study species in the Fragmented Forest Site (this study; vertical red line) and Continuous

Forest Site (Mori et al. 2018; shaded bars). Proportion of clonal ramets and genet area in

the Continuous Forest Site were obtained by subsampling a sample within each 20 x 20

m grid cell with 1000 bootstraps.