A scoping study of the whole-cell imaging literature: a foundational corpus, potential for
mesoscale data synthesis, and implications for standardization of an emerging field

Mary Mirvis[1], Brooke Weingard[1], Steven N. Goodman[2], Wallace F. Marshall[1]

[1]Department of Biochemistry & Biophysics, University of California, San Francisco, San
Francisco, CA

[2]Department of Epidemiology and Population Health, Stanford University, Stanford, CA

ABSTRACT

The level of cellular organization bridging the mesoscale and whole-cell scale is coming into

focus as a new frontier in cell biology. Great progress has been made in unraveling the complex

physical and functional interconnectivity of organelles, but how the entire organelle network is

spatially arranged within the cytoplasm is only beginning to be explored. Drawing on cross-

disciplinary research synthesis methods, we systematically curated the whole-cell volumetric

imaging literature, resulting in a corpus consisting of 89 studies and 118 image datasets. We

describe the trajectory and current state of the field between 2004 and 2024. A broad

characterization, or "scoping review", of bibliometrics, study design, and reporting practices

shows accelerating technological development and research output. We find high variability in

study design and reporting practices, including imaging modality, model organism, cellular

contexts, organelles imaged, and analyses. Due to the laborious, low-throughput nature of most

volumetric imaging methods, we find trends toward small sample sizes (<10 cells) and small cell

types. We describe common quantitative analyses across studies, including volumetric ratios of

organelles and inter-organelle contact analyses. This work establishes the initial iteration of a

growing dataset of whole-cell imaging literature and data, and motivates a call for standardized

whole-cell imaging study design, reporting, and data sharing practices in the context of an

emerging sub-field of cell biology. Our curated dataset now provides the basis for a plethora of

future aggregate and comparative analyses to reveal larger patterns and generalized

hypotheses about the systems behavior and regulation of whole-cell organelle networks. More

broadly, we showcase the potential of new rigorous secondary research methods to strengthen

cell biology's literature review and reproducibility toolkit, create new avenues for discovery, and

promote open research practices that support secondary data-reuse and integration.

INTRODUCTION

Biological organization spans many physical scales, from the atomic and molecular, to global

ecosystems. In cell biology, the dominant relevant scale is that of the cell as the indivisible unit

of life, where questions of cellular function, behavior, motion, communication, and shape

dominate. At the subcellular scale of molecular biology and biochemistry, the structure, function,

information flow of biological molecules, from ions to large macromolecular complexes such as

nucleic acids and ribosomes reigns. Between these two scales lies the less clearly defined

mesoscale, encompassing all subcellular structures larger than ~100 nm including condensates,

cytoskeletal assemblies, organelles, and the cytoplasm (Sear et al., 2015). Great progress has

been made in identifying the relevant structural components of the mesoscale, but defining the

driving forces and principles governing the complex spatial patterning, dynamics, interactions

and systems properties of mesoscale entities is now a pressing challenge in cell biology.

At this frontier of biological organization, new types of questions arise at the intersection of cell,

systems, and physical biology. How do well-understood molecular mechanisms scale *en masse*

across the entirety of the cell? How do disparate mesoscale structures, all well known to be

highly dynamic and structurally complex, interact with each other? To what extent are

mesoscale structure and function directly governed by genetically programmed or energetically

active mechanisms versus emerging or self-organizing from intrinsic mechanical interactions,

self-organization (Laughlin et al., 2000)? Can the entire set of mesoscale components be

understood as a whole interconnected system, and what conceptual frameworks and

experimental and theoretical approaches are necessary for this understanding?

Organelles, the largest mesoscale structural components, together represent the highest level

of organization within the mesoscale. The fine ultrastructure of organelles has been extensively

characterized through electron and light microscopy since the 1950s, and a rich understanding

of the complex molecular composition, cellular functions, and dynamics of each individual

organelle has been amassed since the 1980s. In the last decade, a new picture of organelles as

interconnected components in a complex and dynamic network mediated by many contact site

proteins and communication mechanisms has emerged, summarized in many reviews (Cohen

et al., 2018; Scorrano et al., 2019; Voeltz et al., 2024). A systems view of organelles spawns

new questions around the structure, dynamics, and regulation of the organelle network; its

responsiveness to perturbations and mechanism of disruption in disease; and the role of each

organelle component and pairwise interaction in network structure and function. Questions of

this nature are being increasingly addressed through systems imaging (Bray et al., 2016;

Rohban et al., 2017; Wang & Mukherji, 2022), spatial proteomics (Cho et al., 2022; Hein et al.,

2023; Lundberg & Borner, 2019; Qin et al., 2021), and computational modeling (Donovan-Maiye

et al., 2022; Earnest et al., 2017; Majarian et al., 2019; Murphy, 2012; Viana et al., 2023; Zhao

& Murphy, 2007).

A new construct: Cell Anatomy

Studying the structure of organelle networks is a challenging task. Highly dynamic and

morphologically heterogeneous organelles interact physically through a combination of protein-

based membrane contact sites (MCS), mechanical forces, spatial constraints, membrane

dynamics, and potentially other unidentified mechanisms. In aggregate, the coalescence of

organelle morphologies, spatial distributions, and inter-relationships produces whole-cell

patterns, which we term "cell anatomy". Cell anatomy is a fundamental, but still incompletely

defined cellular property bridging the mesoscale and whole cell, which is presumably both

unique to specific cell types, as well as variable and dynamic within and across cell states. By

defining and observing cell anatomy across many cells and contexts, we can formulate novel

questions. Which types, features, or variations of cell anatomy are common, and which are

context-specific? Which are genetically regulated versus determined by simple geometry or

intrinsic physics? Answering such questions requires the generation of microscopy images of

multiple organelles in tandem in the context of the entire cellular volume, revealing the

morphology and positioning of each organelle in its native, three-dimensional context.

Whole-cell imaging has been accomplished across a range of cell types using a variety of

imaging modalities, including electron, X-ray, and light microscopy methods. Among image data

types, whole-cell imaging is particularly technically challenging. Images encompassing large

biological volumes[1] must be acquired and reconstructed at resolutions high enough to

accurately capture and segment multiple subcellular structures. Segmentation must be done

either manually, which requires specialist expertise and many hours of labor per image, or

automatically, which requires sufficient amounts of high-quality manually-segmented training

data and rigorous validation (Chen et al., 2024; Ekman et al., 2020; Erozan et al., 2024).

Addressing questions about the variability and regulation of 3D mesoscale patterns requires

data from large numbers of cells and is therefore generally beyond the scope of single whole-

cell imaging studies. Aggregating the existing set of relatively small whole-cell imaging studies

may reveal trends in cell anatomy across cell types and biological contexts, providing an avenue

for novel analyses and insights into cell anatomy and organelle interaction networks.

This goal requires finding and curating relevant whole-cell imaging studies across the literature

as comprehensively as possible, extracting useful information, and analyzing image data and

reported measurements. Under the broad umbrella of secondary research, there exists a wide

array of methods to synthesize and review published work, yielding summaries of existing

knowledge, meta-scientific insights into the state of a field or subset of literature, and

identification of trends and gaps in the research landscape to motivate future work. A large

family of systematic, rigorous, transparent, reproducible approaches to literature review

collectively known as *evidence synthesis* or *research synthesis*, offers an array of rigorously

developed methods for comprehensively and reproducibly curating and analyzing trends and

knowledge across studies (Munn et al., 2018, 2023; Tricco et al., 2016). Here, we present a first

step towards leveraging these tools for synthesizing mesoscale data from whole-cell imaging

studies.

Our evidence synthesis approach ultimately seeks to uncover general and context-specific

trends in cell anatomical patterns and organelle interdependencies. Combining information from

many publications and datasets is a promising avenue to answer larger-scale questions than

could be addressed by the results of individual published works. In order to formulate specific

testable hypotheses, it is necessary to first have a clear picture of the amount and content of

data currently available, and the ways it can be merged and compared across studies.

Therefore, a crucial first step is to produce a thorough characterization of the volume, quality,

and technical and biological features of the whole-cell image data landscape. In this work, we

describe a "scoping study", based on "scoping review" methods previously described and

standardized (Arksey & O'Malley, 2005; Tricco et al., 2018). We present the development of our

scoping methodology, including iterative search strategy development and screening, and

delineate the contours of the whole-cell imaging field. We summarize publication trends and the

range of methodologies, reporting practices, and biological conditions included in the rapidly

growing corpus of whole-cell imaging data. We highlight key points of overlap in the data which

lay the foundation for future systematic review work. Our work produces a foundational corpus

of whole-cell imaging data with rich potential for comprehensive knowledge synthesis, novel

biological insights, and a bird's eye view of the changing trends and needs of this promising and

rapidly-crystallizing sub-field of mesoscale cell biology.

WORKFLOW

We adapted our approach from the well-defined scoping review methodology (Arksey &

O'Malley, 2005; Peters et al., 2022). Our workflow for this study is summarized in Figure 1, and

full details into each step of the workflow, including screening criteria, categorization, and

analysis, are described in Methods. In brief, we developed a search strategy consisting of a set

of microscopy terms and a set of cell- and organelle- scale terms, based on an initial set of 11

target papers. Our search strategy returned over 11,000 results from PubMed. We defined two

main inclusion criteria: 1) generation of original three-dimensional *whole-cell* image data in

eukaryotic cells, including complete cell boundary delineation, and 2) volumetric information for

at least two *organelles* concurrently must be included in the whole-cell volume. After three

screening stages – title and PubMed summary, abstract, and full text – conducted by a team of

three reviewers, 89 studies were fully included. We extracted information from studies to

summarize publication trends and study design, including imaging modality and resolution, cell

types, sample sizes, organelles, comparative conditions, and quantitative analyses,

RESULTS

Metadata and bibliometrics.

The potential for effective study curation and data extraction, and ultimately, novel insights into

whole-cell structure, depends on the overall volume, accessibility, and type of studies screened.

We first aimed to broadly characterize the publishing trends within our corpus. The publication

rate of studies containing whole-cell imaging data has risen steadily since 2005, with a peak at

14 papers in 2023 (the most recent complete year, as our search stopped in May 2024), and

49.4% of the included studies (44/89) published in the last 5 years (2019-2024) (Figure 2A). The

majority are open-access publications, which is important for study amenability to systematic

search and screening approaches (70/89, 78.7%) (Figure 2B).

We reasoned that, while many of the studies we include may be similar in nature to our initial

set, in that they perform extensive data collection and analysis in pursuit of a particular

biological question, whole-cell image data may be found in other publication types. Technical

studies which describe some advance in microscopy, image processing, or analysis, often

provide small proof-of-concept datasets. While such studies may or may not undertake

extensive biologically-driven analysis, the images themselves may include valuable structural

information and be available for further analysis in combination with other datasets. The majority

(57, 64%) were "research" articles, focused on answering a biological question, while 31 articles

were considered "technical", focused on method development, although some biological

observations may be included (Figure 2C). One publication is a technical book chapter, which

would have been otherwise excluded from our dataset as a secondary source, but included

some original image data (Gluenz E et al., 2015).

We asked whether authors share the original imaging data via data repositories or other open-

access platforms, which would enable secondary analysis of this information-rich data. Across

the core included datasets and borderlines, only 11 included studies (12.4%) and 1 borderline

study reported that underlying whole-cell image datasets were publicly available, although those

that did were more recently published, from 2015 onwards (Figure 2D). Of these, seven

datasets are available in their entirety as reported in the study (Cortese et al., 2020; Ezzedine et

al., 2023; Laundon et al., 2019; Mocaer K et al., 2023; Müller et al., 2021; Rudlaff et al., 2020;

Uwizeye et al., 2021a), while five are available in partial form (Antao NV et al., 2023; Hair et al.,

2024; McQuate A et al., 2023; Sakaguchi M et al., 2016; Uwizeye C et al., 2021). A summary of

publicly available datasets is provided in Additional File 3. Overall, we see a progressive

expansion of study and data volume, as well as open science practices, with a pronounced

increase in all of these trends in the past 5 years, suggesting good current and future potential

for meta-scientific studies and secondary re-analysis of whole-cell imaging datasets.

Can whole-cell imaging represent an emerging field? We asked whether the corpus is

scientifically interconnected, showing a substantial degree of scientific collaboration and

scholarly acknowledgement. Analysis of co-authorship networks shows many tightly-linked co-

authorship clusters (Figure S1A), and one large cluster of collaborators among prolific whole-

cell microscopy researchers (authors on 2+ studies) (Figure S1B). A citation network shows that

most (71/89), but not all studies in the corpus cite each other. A few studies (Larabell & Le Gros,

2004; Loconte et al., 2021; Uchida et al., 2009, 2011; Uwizeye et al., 2021b) emerged as the

central, most highly connected studies. The most cited study (Valm et al., 2017), with over 1000

citations, does not cite previous studies in this corpus, but is cited by many subsequent papers

that used other microscopy methods, indicating that there has been a convergence of whole-cell

imaging efforts from disparate research streams. This bibliometric overview illustrates the

growing pace and cohesion of the whole-cell imaging literature.

Imaging modality and resolution

Structural biology through volumetric imaging has made great strides in recent years. The rapid

improvement of image resolution, volumetric and statistical throughput, structure identification,

and integration of cross-modal data, including correlative and dynamics approaches, have

enabled exciting advancements in the overall accessibility and applicability of many imaging

modalities to a wider and more complex array of problems than ever before. Whole-cell imaging

approaches the upper limit of capabilities of most high resolution, 3D microscopy and

reconstruction techniques. Each technique offers a slightly different set of strengths in terms of

resolution, throughput, accessibility, cell type amenability and ultrastructural preservation,

organelle identity and number, potential for quantitative analysis, and so on. Therefore, in

characterizing the whole-cell imaging corpus, it is crucial to understand which imaging methods

are being used, which are more common, and how they data they produce compare and

contrast. This may motivate recommendations for modalities best suited for future whole-cell

imaging studies.

In total, we identified 13 distinct volumetric imaging types of modalities used to produce whole-

cell image data. Focused ion beam SEM (FIB-SEM) (also called ion-abrasion SEM, IA-SEM)

and soft X-ray tomography (SXT) and are the most common modalities, represented in 22 and

21 studies, respectively (Figure 3A). These were followed by serial block-face SEM (SBF-SEM),

serial sectioning TEM (ssTEM), electron tomography (ET), fluorescence (including 3D SIM, and

multispectral lattice light sheet), optical tomography (including optical diffraction and optical

computed tomography/live cell CT), hard X-ray tomography (HXT, X-ray ptychography),

scanning TEM (STEM), and X-ray diffraction (XRD) techniques. Field emission SEM (FE-SEM),

scanning transmission X-ray microscopy (STXM), and SEM array tomography (SEM-AT) were

each found in one study. In total, 55 studies used electron microscopy (EM) techniques (FIB-

SEM, SBF-SEM, ssTEM, ET, STEM, FE-SEM, SEM-AT), 27 (30%) used X-ray techniques

(SXT, HXT, XRD, STXM), and 10 used optical techniques (fluorescence, optical tomography).

Each modality excels within a limited range of spatial resolution, determining the size scale,

volumetric completeness, throughput, and potential for artifacts of anisotropic resolution in the

resulting images. Generally, optical imaging methods are limited by the wavelength of light

(~200nm), typically capable of achieving higher lateral (xy) than axial (z) resolution, but can be

used to produce sample sizes in the hundreds or thousands. Electron microscopy modalities are

capable of achieving resolution in the sub-10nm range, but are typically limited in volumetric

scope and statistical throughput due to the high information density and cumbersome sample

preparation, imaging, and image processing. X-ray based methodologies typically fall

somewhere in between light and electron microscopy in throughput and resolution (Guo &

Larabell, 2019; McDermott et al., 2012). The range of cell types potentially accessible for whole-

cell reconstruction, information content contained in the resulting images, comparability across

datasets and modalities, and capacity for secondary analysis is strongly dependent on image

resolution. Whole-cell imaging requires the acquisition of large volumes, so even the best

resolution possible with a particular modality or microscopy may need to be compromised to

increase the reconstruction volume and/or throughput.

We extracted the technical parameter information for each distinct dataset across the corpus,

where a dataset is considered a distinct set of cells of the same type imaged with the same

modality, such that multiple datasets may appear within one study (e.g. four datasets

comprising two cell types (*C. pepo, N. tabacum*) each imaged by two modalities each (ssTEM,

FIB-SEM) in (Zechmann et al., 2022). We found 118 total datasets across the 89 studies

(Additional File 4). The reported resolutions are summarized as lateral (L) and axial (A)

resolutions for electron microscopy (EM), X-ray microscopy (X-ray), and light microscopy (LM)

methods in Figure 3B. Each dataset is plotted as a separate datapoint in Figure 3C. Overall, 72

datasets (61%) were generated using EM techniques, 29 (24.6%) with X-ray techniques, and 17

(14.4%) with optical techniques. As expected, the highest resolution was achieved by electron

microscopy modalities (ET, FIB-SEM, SBF-SEM, ssTEM, STEM, FE-SEM, SEM-AT), followed

by X-ray based methodologies (SXT, HXT, X-ray diffraction, STXM), and the lowest and most

variable resolution in optical data (fluorescence microscopy, optical tomography). Isotropic

resolution, which is highly advantageous for structural accuracy and morphological analyses

(Chen et al., 2024; Le Gros et al., 2005), was far more common in X-ray based techniques

(26/27 datasets, 92.9%, counting only those reporting both axial and lateral resolution), than in

EM-based (25/56, 53.6%) or light-based (0/16) modalities.

Imaging resolution is therefore unique for each study, chosen within the available range to

optimize feasibility and information density for the particular question or goal, and is a critical

methodological parameter in methods reporting for critical interpretation of results,

reproducibility, and evidence synthesis. In extracting the reported resolution from each study,

we found that imaging resolution was not clearly reported for 18 datasets from 14 studies, in

some cases inconsistently stated (conflicting and/or ambiguous mentions throughout the text),

while in others, resolution was incompletely reported (e.g. axial (z), but not lateral (xy),

resolution clearly stated, Figure 3C, right panel).

In summary, our results show that many modalities are capable of producing whole-cell image

data, but X-ray based techniques, particularly soft X-ray tomography, have so far been most

successfully leveraged to produce high resolution, isotropic, high-throughput whole cell

datasets.

Cell type and sample size

The potential to pool and compare findings across studies depends largely on the biological

similarity of the data, in particular, the cell type or model organism imaged. The same cell type

imaged in multiple studies (with varying sample sizes) offers the potential to cross-reference

and pool data across modalities and experimental conditions. The range of cell types

represented across the corpus will point to appropriate groupings and comparisons for cross-

cell type syntheses.

A wide range of cell types are covered in the corpus, including yeast, protists, plants,

invertebrate animals, and both cultured and *in situ* mammalian cell types from rodents, non-

human primates, and humans (Fig. 4A). In total, we found 80 distinct cell types across 118

datasets. The most strongly-represented cell types (in terms number of separate datasets and

number of cells imaged in total across datasets, which we call "data density"), are highlighted in

Table 1.

Table 1. Cell types with highest data density. A summary of data available for cell
types/species recurring in multiple studies across the corpus.

Cell type	# of studies	Total # cells
S. cerevisiae	11 (13 datasets)	380
P. falciparum	5 (6 datasets)	84
C. reinhardtii	4	30
T. cruzi	4	4
Rat insulinoma pancreatic β cell (INS-1E)	3	228
Mouse pancreatic β cell	3	16
S. pombe	3	8
human platelets	2	94
Trypanosoma brucei	2	80
Candida utilis	2	42
human B lymphocytes	2	18
COS-7	2	12
Toxoplasma gondii	2	5
HEK293T	2	5

Most studies had small sample sizes, defined as the number of cells fully reconstructed in 3D

with multiple organelle volumetric information compatible with our inclusion criteria, with 81

(68.6%) of datasets reporting n<=10 whole cell images, and 36 (30.5%) studies consisting of a

single cell (Figure 4B). Single-cell datasets were more common in earlier papers, and frequently

occurred in technical papers – 16 of the 89 studies were technical papers reporting a single

whole-cell reconstruction, representing half (51.6%) of the 31 technical studies and 66.7% of the

single-cell studies. This suggests that the first ten or so years of our dataset (~2005-2015) could

be described as technical early days, but the advances during this time paved the way for more

biologically-focused research studies with larger datasets in the following decade (~2015-

today). Sample sizes were largest in SXT (up to n=200) and SBF-SEM (up to n=141) studies. In

14 datasets, the study sample size, was either reported inconsistently or incompletely, in which

case we reported a minimum known sample size, or else was not reported at all, resulting in

missing data in our analysis (Figure 4A, white bubbles).

Organelles and anatomical completeness

Our primary inclusion criterion was the presence of at least two organelles within the same

whole-cell reconstruction. We argue that for visualizing cell anatomy, at least two different

structures must be present in order to provide spatial context for one another. Furthermore, as

pairwise spatial and morphological inter-organelle interactions are building blocks of the whole-

cell organelle network, the presence of two organelles reveals at least one pair whose

interaction can be described. We therefore extracted the list of organelles reconstructed in each

dataset to assess the potential for analysis at the single-organelle, inter-organelle, and cell

anatomical levels.

Figure 5A shows the cumulative organelle data per cell type across the corpus. The nucleus

and mitochondria are the most commonly segmented organelles across all datasets (106 and

91 instances respectively, accounting for multiple distinct cell types per paper in some cases),

followed by vacuoles/lysosomes (53 instances) endoplasmic reticulum (28 instances), lipid

droplets (27 instances), Golgi (26 instances), chloroplast/plastid (27 instances), vesicles (18

instances), flagella/cilia (17 instances), peroxisome (7 instances), centrosome/basal body (5

instances), endosome (5 instances), multivesicular bodies (1 instance), and autophagosome (1

instance). Some organisms may appear to have only one or no organelles included despite our

inclusion criterion requiring imaging of two or more organelles . This is due to the fact that the

heatmap only includes "common organelles" present in all or most eukaryotes. The blank lines

represent studies which image and reconstruct only specialized or cell type-specific organelles,

such as the specialized secretory organelles (microneme, rhoptry, dense granules) or conoid

structures in *Toxoplasma gondii* (Koutsogiannis Z *et al.*, 2023; Paredes-Santos TC *et al.*, 2012).

In some cases, we find one common organelle in combination with specialized organelles, such

as the study in C57BL/6 resting mouse platelets (mitochondria, dense granules, alpha-granules,

canalicular system) (Pokrovskaya I *et al.*, 2021). The recurrence of cell types in multiple studies

enables potential examination of organelle morphologies across datasets, experimental

designs, and biological contexts. While not all studies imaging the same cell type segment the

same set of organelles, the most frequently-imaged specific organelles are found among the

best-represented cell types: *S. cerevisiae* vacuoles (12 instances), nuclei (11), mitochondria

(10), ER (7),lipid droplets (7), and vesicles (5); *P. falciparum* nuclei (5) and vacuoles (4), and *T.*

cruzi nuclei (5).

The rationale for requiring at least two organelles per segmented cell in the dataset is 1) the

supposition that a single internal landmark is insufficient to constitute an anatomical pattern, and

2) to reveal spatial interactions between organelles in the 3D context of the entire cell. Overall,

we found 59 pairwise organelle combinations occurring across the corpus, of which 35 had at

least five independent instances, suggesting rich potential for comparison and synthesis of each

organelle-organelle relationship across many cell types and contexts (Figure 5B). The most

common organelle co-occurrences are mitochondria-nucleus (82 instances across 65 cell

types), nucleus-vacuole/lysosome (48 instances across 36 cell types), and mitochondria-

vacuole/lysosome (40 instances across 33 cell types). These cases will enable within-cell type

synthesis of similar data (e.g. analysis of a specific organelle pair, as well as cross-study

comparison, which can potentially isolate biological and technical differences.

The ultimate requirement for an understanding of cell anatomy and modeling whole-cell patterns

and inter-organelle interaction networks is to be able to visualize as many organelles

simultaneously as possible. The number of "common organelles" per reconstruction, termed

"anatomical completeness" is distributed with a peak at 3-4 organelles (Figure 5C). The most

anatomically complete reconstructions are in choanoflagellate *S. rosetta* (Laundon et al., 2019)

with 8 organelles, and 7 organelles in *L. mexicana* (Hair et al., 2024), sponge *O. carmella*

choanocyte (Laundon et al., 2019), *A. thaliana* cortical root cell (Cui Y et al., 2019), and *S.*

cerevisiae (Wei et al., 2012).

Morphological metrics and Quantitative analyses

The unique advantage of high-resolution 3D structural data of whole cells is their capacity to

provide rich and precise quantitative morphological and spatial information. Future meta-

analysis of organelle features relies on the presence of comparable quantitative measurements

across studies. An assessment of quantitative analyses performed across studies revealed a

limited set of organelle geometric features measured (Figure 6A). Volume was the most

frequently measured property (65 studies, 73%), followed by volume fraction (organelle-to-cell

volume ratio, 50 studies, 56%). Also common were surface area (34 studies, 38%), count/copy

number (29 studies, 32.6%) and linear dimensions such as length and diameter (27 studies,

30%). Less frequent, and more variable, were measures of density such as LAC or electron

density and positioning/distance (n=18 each, 20.2%), organelle contacts/associations and

shape (n=14 each, 15.7%). Organelle volume scaling trends were calculated in 7 studies (7.9%)

and three studies measured refractive index. One study devised a method to quantify

subcellular crowding of organelles ("congestion index", Loconte V *et al.*, 2022). These

frequencies roughly reflect the general ease of measurement and conceptual familiarity of these

metrics - volume and surface area are generally trivial to measure, albeit with variable accuracy,

with many standard software tools automatically generating these calculations. Count and

dimensions can be straightforwardly manually or semi-automatically calculated. Distance and

contact frequency, shape, and crowding require more complex calculations and computational

operations, and can be interpreted in variable ways. A sizable fraction of studies (n=14, 15.7%)

performed no quantitative analysis.

Comparators

While whole-cell structural data is largely static, cellular structure is in fact highly dynamic and

variable across cells. We recorded the experimentally compared conditions between samples

("comparators"), reasoning that variety in conditions for a single cell type could suggest

functional organelle and anatomical variability, while overlap in conditions across cell types

could reveal generalizable context-specific organelle and anatomical trends. Most studies (71,

79.8%) included comparator data across at least one cellular perturbation or condition, which

we grouped into broad classes (Figure 6B). The most common of these are cell cycle stages

and metabolic conditions, such as glucose stimulation of pancreatic cells (14 studies each).

Other comparators, in decreasing order of frequency, were cross-cell type comparisons (n=12),

stressors such as drug treatments (n=9), methodology, such as fixation methods or imaging

modalities (n=6), stages of life cycle, development, or differentiation (n=6), environmental

conditions, such as light exposure (n=3), morphology, such as shape or size (n=3) , disease or

infection states (n=2) , functional or behavioral states, such as cell-cell communication (n=1),

and genetic mutation (n=1). The predominance of cell cycle stages and cell type as the primary

comparators highlights the potential of whole-cell imaging to serve as a basis for defining cell

type and functional cell state-related cellular reorganizations.

In summary, the whole-cell multi-organelle imaging corpus includes 89 published studies

between 2004 and 2024 from over 11,000 initial PubMed results. In total, the dataset represents

the whole-cell imaging and reconstruction of at least 1690 individual cells, of 80 eukaryotic cell

types from protists to humans, across 118 independent datasets, 13 imaging modalities, at least

13 common organelles in many different combinations, 71 perturbative or comparative

conditions, and at least 12 types of morphological metrics quantitatively analyzed.

LIMITATIONS

Our scoping study is, to our knowledge, one of the first applications of scoping review methodology to the field of whole-cell microscopy, and to fundamental cell biology more broadly. Our method comes with limitations arising from the iterative nature of proof-of-concept methods development, the breadth and heterogeneity of the literature corpus, and the size of our team. One of the goals of a formal systematic literature synthesis is to be as comprehensive as possible, spanning peer-reviewed and grey literature sources as thoroughly as possible. Our approach focused specifically on PubMed, but ideally, other databases such as Web of Science and EMBASE, would be included as well. Although our PubMed search was extremely broad relative to our final inclusion set, which comprised only less than 1% of the original search results, we cannot be sure how many relevant studies might have been missed by the search entirely. Furthermore, grey literature sources such as dissertations and online image data hosting sites without corresponding publications would be of relevance to a comprehensive set of cell anatomy data, but would be missed by our strategy. For example, the Allen Institute for Cell Science hosts a collection of 3D confocal images in human iPSCs, a subset of which would fit our inclusion criteria, but these datasets have not been directly thoroughly described in a publication, despite their inclusion within a large dataset underlying many impactful publications (*Cell Line Catalog*, 2024). Furthermore, as of January 2025, our search string retrieves 300 new results since our last round of curation in May 2024. The dataset and analysis presented here should be seen only as an initial characterization which will be continually updated as the literature base expands. We intend to establish our growing curated corpus and analysis as a publicly available, living database of cell anatomy data and field trends, inviting community input to identify gaps and curate the corpus as thoroughly as possible.

Another limitation of our approach, as of any systematic research synthesis, is that the choice of

strict inclusion criteria necessarily excludes potentially valuable studies. These boundary cases

help to define the field of whole-cell reconstructions and reveal peripheral sources of knowledge

that can be incorporated into follow-up studies depending on the specific biological question.

One core criterion was the need to define "*whole cell*" strictly, requiring that the full cell

periphery is clearly delineated. The rationale for this is that, first, the full anatomy of the cell

requires that no part of the cell is missing from the image, and second, the intact cell boundary

is needed for accurate measurement of cell volume, which we consider to be the critical

common measurement required for any cross-study quantitative analysis as the denominator for

organelle volume ratio. Therefore, many studies reporting mostly complete cell volumes, in

some cases even describing the data as "whole cell" despite missing small cell edges, had to be

excluded. This was often the case for adherent cultured cells, which spread out flat on culture

dishes with thin, faint, irregular protrusions when sparsely plated, and for very large, irregularly

shaped cell types such as muscle cells and neurons, which are extremely difficult to perfectly

capture in their entirety *in situ* or in culture. In other cases, we found studies of tissue sections

containing partial reconstructions of multiple cells clipped at the section boundary, including

organelle volumes and volume ratios pooled and normalized across the entire section volume,

but no single cell fully reconstructed. Therefore, our strict "whole cell" criterion introduces a cell

type bias into our corpus. Moving forward, a case could be justified for softening this criterion to

estimate rather than directly calculate cell volume. The systematic nature of our screening

process means that steps can be back-tracked to retrieve specific studies previously excluded

for either of these reasons and include them under an adjusted set of criteria.

DISCUSSION

We have created a comprehensive corpus of whole-cell imaging literature spanning a wide breadth of modalities, cell types, organelles, and contexts through a novel application of formal scoping review methodology. There is substantial comparability across studies, highlighting opportunities for further meta-scientific and synthetic studies building on this corpus. Future directions include systematic review and meta-analysis approaches to understand how organelles typically self-organize and interact within the whole cell space across biological contexts, including qualitative categorizations of cell anatomy, and meta-analyses of key morphometric measurements.

Although our search is not exhaustive, this corpus comprises a foundational evidence base revealing an emerging field of significant, and rapidly increasing, volume and scholarly interconnectedness, unified by a specific data type – whole cell, multi-organelle 3D reconstruction (Figures 1 and S1). Our systematic curation and meta-scientific characterization of these remarkably information-dense structural data, from raw data to processed images to derived measurements, enables a range future secondary analysis and synthesis efforts. The substantial methodological and biological overlap opens a wide range of possibilities for within- and cross-group synthetic studies to uncover novel insights that would otherwise be difficult to access through traditional primary research methods. Data in a specific cell type pooled across datasets, modalities, and conditions provide opportunities to query the consistency of similar metrics across studies, integrate disparate findings into an organism-centric model, and compare distinct imaging modalities and parameters applied to the same experimental system. Summarizing quantitative analyses and extending them where possible allows for a more comprehensive quantitative understanding of generalizable principles of organization, an assessment of single organelle, between-organelle, and cell-wide feature variability across contexts, and provide a strong empirical foundation for integrative modeling efforts (Loconte et al., 2023). Publicly shared original data enable novel analyses and the development of new data

integration and secondary analysis workflows. Ultimately, work building on our corpus promises

to reveal emergent patterns of cell anatomy across cell types and contexts, and insights into the

constraints and variations embedded in these patterns as a result of cellular geometry and

organelle interactions.

We intend for our dataset and analysis to become a growing tool for the scientific community. In

addition to generalized, high level synthesis, the corpus can be subdivided in many ways to

enable inquiry into specific cell types, organelles and organelle interactions, biological

perturbations, imaging and image analysis approaches. In the near future, we intend to make

the corpus easily accessible, interactive, and kept up-to-date so that researchers interested in

any component of the corpus can easily contribute to and draw upon it for their own

investigations. We hope this work will galvanize whole-cell imaging and analysis researchers,

synthesis researchers, and anyone else interested in organelle-to-whole cell mesoscale

structural biology to create a community, collaborating to refine the ongoing curation and re-use

of whole-cell data, refine and unify the generation and reporting of novel work, and engaging in

discourse around terminology, best practices, optimal applications for existing imaging

technologies and needs for technical advancement, and a broad vision of the future of the

whole-cell imaging and cell anatomy field.

The range of imaging modalities and technical parameters represented across our corpus

demonstrates the versatility of approaches available to achieve whole-cell anatomy data. It also

provides opportunities to directly compare these approaches and trends in how they are

applied, identify common challenges, and motivate recommendations for future work. We find

that the current whole-cell imaging landscape, on the whole, is biased toward small cell types

and small sample sizes, with a single cell reconstruction reported in nearly one-third of studies.

This is largely due to the time-, cost-, and labor-intensive nature of whole-cell image acquisition,

reconstruction and multi-organelle 3D segmentation, particularly at sub-100nm resolution. The

13 imaging modalities represented each offer particular advantages and disadvantages, with

unique tradeoffs between resolution, throughput, native structure preservation, and overall

tractability depending on the application. For example, the two dominant modalities, SXT and

FIB-SEM, are both capable of achieving high isotropic resolution with some degree of

quantitative throughput. SXT generally achieves isotropic 30-60nm resolution with minimal

sample damage or tomographic artifacts, and is capable of producing throughput in the

hundreds per study. However, cell size is strictly capped by the ~10um diameter of the capillary

used to hold the samples, limiting the cell types one can study by this method, and access is

limited by the requirement for beamline time. FIB-SEM excels for very high-resolution

reconstructions (~5 nm), but is limited by throughput and, as a sample-destructive method, is

more likely to introduce ultrastructural artifacts. Extensive reviews of strengths and weaknesses

across volumetric imaging modalities have been discussed in depth elsewhere (Hanssen et al.,

2012; Loconte et al., 2021; Loconte V et al., 2022), and technical capabilities for all modalities

are constantly being improved.

However, the limiting factor for maximizing the utility of whole-cell image data is perhaps not

data quality or abundance itself, but rather the limited use and reuse of these incredibly rich

datasets, potentially leaving much of the embedded information untapped. We found that the

publication of whole-cell imaging studies is accelerating, with growing trends toward open-

access publication and data sharing, suggesting good current and future potential for meta-

scientific studies and secondary re-analysis of datasets (Fig. 2). However, only a small subset of

our corpus made original data publicly available, and we found inconsistent reporting of key

methodological details, such as imaging resolution and sample size. Such metadata items are

crucial for meta-scientific analysis, including secondary analysis and synthesis research as well

as within-study reproducibility and cross-study replicability assessment. Currently, there is

increasing momentum in the development of open science and data sharing guidelines and

implementation led by international collaborations of scientists (FAIR, DORA), funding bodies

(NIH, NSF), and publishers. Many data type-specific metadata standards have been developed

for bioimage data by community consensus (Global BioImaging, REMBI, foundingGIDE, Open

Microscopy Environment) and the growth of image dataset repositories (EMBD, EMPIAR, IDR,

SSBD, NIH Imaging Data Commons). Image data, particularly high-content volumetric imaging

such as whole-cell reconstruction datasets, present particular challenges for data sharing, such

as the large size and complexity of files and datasets, irregular file formats. Ongoing efforts to

develop novel bioimage file formats (OME-Zarr (Moore et al., 2023); OME-NGFF (Moore et al.,

2021)) and accessibility tools (BioFileFinder) promise to help tackle some of these barriers, and

lead to new solutions as the public image data ecosystem scales up. There may be factors of

particular interest to the emerging scientific community generating and using whole-cell imaging

data with regards to specific imaging modalities and data types, such as segmentation and 3D

reconstruction parameters. As the publication of whole-cell imaging studies accelerates, we

envision that community discourse and establishment of specific standards for methodological

design, reporting, and data sharing could be of broad benefit. We hope to see a coordinated

effort to develop open science practices to maximize the utility of these costly-to-produce data.

While interest in open science and data-reuse is quite mainstream and rapidly growing in the

sphere of cell and molecular biology, the range of approaches currently used are largely

focused on big -omics data informatics. The past two decades have yielded the independent

development of synthesis and integration methods in some specific contexts, such as GWAS

meta-analyses in genomics, computationally-assembled gene- and protein interaction networks

in bioinformatics, and massive consortium collaborations to map cell types and beyond ("Meta-

Analysis in Basic Biology," 2016). There is still a clear need for robust knowledge and data

synthesis in biology, driven by widely recognized challenges associated with "data deluges"

("The Data Deluge," 2012; Zhang, 2024) and the exponential growth of scientific output (Hanson

et al., 2024; Pautasso, 2012). The monumental goal of constructing meaningful holistic

understandings of highly complex biological systems by merging a rich tradition of narrowly

focused mechanistic studies and molecular parts lists, with interdisciplinary systems,

biophysical, evolutionary, synthetic, and engineering perspectives (Rafelski & Theriot, 2024),

requires robust frameworks to comprehensively and rigorously synthesize data, knowledge, and

interpretations across disciplinary boundaries, aided by a high-level understanding of field

trends revealed by meta-scientific analyses.

There is at present limited familiarity in the fundamental biology field of evidence synthesis

methods widely used in many disciplines including the clinical and social sciences. These are

extensive, carefully developed toolkits supported by synthesis researchers and large

international collaborations and initiatives such as Cochrane and the Joanna Briggs Institute.

Systematic research syntheses in fundamental cell biology, particularly intermediate-scale

syntheses to address specific biological questions or hypotheses, are beginning to come on to

the scene in recent years, still rare and methodologically disparate (Gillooly et al., 2015; Opalek

et al., 2023; Rutkowska et al., 2020; Sercel et al., 2023; Trumpff et al., 2021).

The present work illustrates the unique considerations, challenges, and scientific and meta-

scientific potential of developing an evidence synthesis toolkit tailored to fundamental cell

biology. In contrast to translational and applied disciplines, particularly those based on animal or

human subjects, exploratory fields such as fundamental cell biology marked by high

heterogeneity in their technical repertoire, study design, experimental and reporting practices,

philosophical and sociological standards for epistemology and argument construction, and

conceptual and linguistic frameworks. This heterogeneity presents unique and significant

challenges for the application of highly standardized, guideline-driven evidence synthesis

methods. For example, development of a narrow and specific bibliographic database search

strategy is particularly challenging. Firstly, deep and broad technical expertise is required to

understand the scope of research likely to contain a particular data type, specific experiment, or

conceptual theme. Second, highly varied terminology used across the literature, and particularly

in the lack of concrete norms for titles and abstracts results in a tendency toward very large

search result corpuses, preferring comprehensiveness over specificity. The search scope and

resulting corpus may therefore be larger than is the accepted norm in other disciplines.

Therefore, inclusion criteria must be carefully justified and iteratively refined, and robust

stepwise methods for screening and data extraction, ideally carried out by teams of 2-4 expert

investigators working in parallel to minimize bias, are needed. In general, a typical systematic

review process can take 1-1.5 years of painstaking manual work conducted and validated by a

dedicated team.

The prospect of employing AI tools to aid in these intensive steps is attractive, and being

actively explored by many scientific groups and software companies (Bolaños et al., 2024; Yip

et al., 2024). However, due to the goal of maximum comprehensiveness and need for a careful

balance between rigor and pragmatic flexibility to handle the highly heterogeneous and nuanced

nature of the fundamental biology literature, the requirement for human judgement and

validation is unlikely to be replaced by AI in the immediate future. We could envision a manually

curated scoping study, such as this one, forming the foundation for a novel synthesis effort, and

in tandem, creating a robust training dataset for an AI synthesis assistant, while further

refinement, establishment of a living corpus, and updating with novel publications as they are

released, can be accomplished with the aid of the trained AI model and community curation and

validation. We look forward to a bright future in which specialized synthesis researchers, in

collaboration with the expert community and automated tools, help to create a thriving

ecosystem for novel discovery and meta-research to derive the maximal benefit from the

immense volume and richness of biological literature and data.

METHODS

Search strategy:

We developed our search strategy starting with an initial set of 11 target papers (Decelle J et al.,

2021; Heinrich L et al., 2021; Henderson et al., 2007; Laundon et al., 2019; Müller et al., 2021;

Uchida et al., 2009, 2011; Uwizeye et al., 2021a; Valm et al., 2017; Wei et al., 2012; White et

al., 2020)], thoroughly combed for cell attribute- (e.g. "cell organization", "subcellular

architecture"), mesoscale-, and imaging-related terms. These terms were used to devise an

iterative series of search strings for PubMed aimed at retrieving a defined corpus that includes

the full initial set. While an initial attempt focused on cell anatomy-related terms was

unsuccessful due to the heterogeneity and inconsistent usage of such conceptual language,

iterations on search strategies consisting of 3D imaging and mesoscale (cell/subcellular and

organelle-scale) terms successfully retrieved the initial papers.

(("Imaging, Three-Dimensional"[MH] OR "3D imaging"[tiab] OR "three-dimensional

imaging"[tiab] OR "3-D imaging"[tiab] OR "Soft X-ray tomography"[tiab] OR (3-D[tiab] OR

3D[tiab] OR "three dimensional"[tiab] OR "three-dimensional"[tiab]) AND (tomography[tiab]

OR microscopy[tiab] OR Microscopy[MH] OR Tomography[MH] OR imaging[tiab]))

628   AND

((ultrastructure[MeSH Subheading]) OR (Organelles[mh] OR organell*[tiab]) OR

(intracell*[tiab] OR intra-cell*[tiab]) OR (subcell*[tiab] OR sub-cell*[tiab]) OR (((whole[tiab]

OR entire[tiab]) AND cell[tiab]) OR "whole-cell"[tiab] or "whole cell" OR "entire cell"[tiab])))

NOT Review[pt]

We set our search to begin at 2005 due to the emergence of technological capability to perform

whole-cell microscopy and reconstructions around this time (Le Gros et al., 2005). However, this

missed one key earlier study, which was added *post hoc* (Larabell & Le Gros, 2004). The most

recent PubMed search update was performed on May 23, 2024.

Screening

Our search strategy returned over 11000 results from PubMed. Because our search was aimed

at a particular data type (whole-cell volumetric microscopy) rather than a specific research topic,

our target corpus is highly heterogeneous, spanning a wide range of publication venues, key

terminology, and other bibliometric features. Therefore, our primary objective was to retrieve our

initial set as well as any studies with even a marginal potential of containing whole-cell

microscopy data, yielding a highly sensitive but not specific search.

To curate our dataset, we performed three rounds of screening according to the following

strictly-defined inclusion criteria:

1. Cell anatomy is necessarily a whole-cell property. The study must have collected original

three-dimensional *whole-cell* images in eukaryotic systems, described in the primary study

in the form of direct visualizations (still images and/or supplementary movies displaying

volumetric rotations), and/or quantitative analyses necessarily taken from whole-cell data

(e.g. organelle-to-cell volume ratios). "Whole-cell" requires that the entire cellular boundary

(cell membrane or cell wall, or other marker showing the continuous cell periphery) is

captured within the image, such that the cellular volume can be calculated.

2. Volumetric information for at least two *organelles* concurrently must be included in the

whole-cell volume. We reasoned that at least two subcellular structures must be visible in

relation to one another for the concept of "anatomy" to be applicable. This criterion requires

a full volumetric reconstruction of the multi-organelle cell, with organelle identities clearly

annotated (typically in the form of a segmentation). The reconstructed data should either be

directly shown as an image in figures, movie files, or associated data, or necessarily

generated to produce published analysis, such as volume calculations. Although the usage

of the term "organelle" has expanded in recent years, we used a traditional definition,

referring to a structurally and functionally distinct, membrane-bound, intracellular

compartment.

Curating a set of target papers from this large set of search results required a series of

increasingly thorough screening steps – title and PubMed summary, abstract, and full text –

conducted by a team of three reviewers. Search results and screening decisions for each stage,

including exclusion reasons, along with information extracted from initial target papers are

provided in Supporting File 1.

Title/summary. The first screening step was a single-reviewer rapid screen through search

result pages in the web browser (Mozilla Firefox) of titles and short three-line PubMed-

generated summary statements. Studies that were obviously irrelevant, focused on the wrong

biological scale, or purely technical with no biological data were excluded. Studies that clearly

described whole-cell imaging and/or structures of entire cells were included. If unclear, a skim of

the abstract (and PMC figures, if available) was done, and studies that were still ambiguous

were included for further screening. This step was performed by one reviewer (MM), and

excluded 95% of results.

Abstract. A custom Python (v3) script using BioEntrez module of Biopython (Cock et al., 2009)

was used to pull abstracts from PubMed e-utilities (*Entrez Programming Utilities Help*, 2010)

and record screening decisions from one reviewer. Studies were either included, excluded with

exclusion reasoning recorded, or marked for full text screening. A 10% subset was screened in

parallel by a second reviewer (WM) to check inter-rater consistency. A low level of consistency

was initially found due to differing definitions of "organelle". This led to a refinement of our

working definition of "organelle" for the purpose of screening criteria (see above), and the full

set of abstract screening was performed in parallel by two independent reviewers (MM & BW).

This step yielded 60 clear hits (10.8%), excluded 184 (32.3%), and sequestered 326 studies for

full text review.

Full text screening. Full texts, including supplementary data, were screened in parallel by two

reviewers (MM & BW). Screening decisions were recorded in Covidence software (Veritas

Health Innovation, n.d.), along with reasons for exclusion. Exclusion reasons were categorized

as: missing 3D data, including cases where 3D data is textually described but not included as

data in the paper ("Not 3D"); imaging only at subcellular or multicellular/tissue scales ("Wrong

scale"); cells only partially imaged and/or reconstructed, or with cut-off edges such that cell

volume cannot be calculated ("Not whole-cell"); raw data missing organelle segmentation ("Not

segmentation"); fewer than 2 organelles segmented ("Not multi-organelle"); data reused from

other studies ("No original data"); review or perspective articles ("Wrong article type"); and a

general lack of whole-cell, multi-organelle reconstruction data ("No relevant data").

We also designated 39 studies as "Borderlines" if they represented edge cases with regards to

our inclusion criteria, but nevertheless contained relevant data, analysis, or insights, therefore

not warranting complete exclusion from our dataset, summarized in Additional File 1. One

borderline category is "whole-cell data not shown", including studies in which the target data

type is implied or explicitly described in the text but not included in figures or associated data

files, or provided in supplement where files are inaccessible. This includes several studies that

partially reconstructed prohibitively large cell types (Trebichalská et al., 2021), as well as

711 reconstructed tissue volumes from which while relevant metrics are quantified, such as

712 organelle volume/volumetric ratios, calculated across the tissue block rather than in individual

713 cells, as distinct cell boundaries could not be delineated. In some cases, the whole-cell images

714 had slightly clipped cell edges due to nuances such as cells in cultures spreading their edges

715 and protrusions far from the main cell body on a culture substrate, or an elongated cell type

716 such as myocytes. We chose to exclude such examples but include others such as images of

717 protists with flagella partially cut-off, due to the fact a missing edge of the cell body leaves key

718 morphometrics, such as cell volume and shape, impossible to calculate, and creates

719 uncertainties about holistic cell anatomy due missing information, whereas missing part of a

720 cilium or flagellum would only affect analyses of ciliary length, but would preserve the ability to

721 calculate organelle to cell size ratios. Other borderline categories include: organelle identities

722 uncertain; ineligible structures defined as "organelles"; a very minor volume of the cell edge is

723 cut off in the image, technically violating the "whole cell" criterion; cell membrane not clearly

724 delineated; the paper itself does not include the relevant data, but accompanies the

725 establishment of a public repository or resource containing relevant data.

727 Disagreements in inclusion/exclusion decisions were discussed between reviewers, prompting

728 slight adjustments to our protocol, such as establishing new exclusion reasons or borderline

729 categories. One earlier study was added *post hoc* (Larabell & Le Gros 2004). Ultimately, 89

730 studies were fully included.

732 ***Data Extraction and Analysis***

733 We extracted information from studies to summarize study design, including imaging modality

734 and resolution, cell types, sample sizes, organelles, comparative conditions, and quantitative

735 analyses, only as pertains to information within the study directly describing or derived from

736 whole-cell imaging data, as opposed to other types of data within the same paper. Data

extraction was performed manually by two researchers independently (MM & BW), and

subsequently discussed observations to reach consensus.

Data items were annotated on full text PDFs in Zotero (*Zotero | Your Personal Research*

Assistant, n.d.) and entered into the data extraction template in Covidence (Veritas Health

Innovation, n.d.), followed by inter-rater consensus discussions to resolve disagreements and

inconsistencies. Extracted data were exported from Covidence as .csv, and manually processed

and formatted in Excel. The full set of final data extraction items, including 89 included studies

and 39 borderline studies, is provided in Additional File 2. Co-authorship networks were

generated in VOSViewer (van Eck & Waltman, 2010) using PubMed files as input and the

following settings: bibliographic data, all co-authorship, include documents with a large number

of authors, allow unconnected items, and overlay visualization. Citation network was created in

Litmaps (*Litmaps (Version 2025-01-16) [Search Tool].*, 2024) using the following settings: X

axis: Publication Date, compact; Y axis: Map Connectivity; Size: Cite Count, linear interpolation;

disable Position Editing, Show citation arrows, Avoid collisions. All other data analysis and

plotting was performed in Python using Jupyter notebooks and the following packages:

Matplotlib (Hunter, 2007), SciPy (Virtanen et al., 2020), Pandas (The pandas development

team, 2020), Numpy (Harris et al., 2020), Seaborn (Waskom, 2021). Code and source data files

are available at https://github.com/bdwucsf/Cell-Anatomy-Scoping-Review.

Acknowledgements

This project was funded by the UCSF Program for Breakthrough Biomedical Research, funded

in part by the Sandler Foundation, and NIH grant R35 GM130327. We owe a debt of gratitude to

the UCSF librarians, Peggy Tahir, Josephine Tan, and Evans Whitaker, for evidence synthesis

methods consultations and assistance with developing our search strategy. We thank Dr. Jane

Maienschein, Dr. Susanne Rafelski, Dr. Sabina Leonelli, Dr. Matthew Akamatsu, Dr. Shane

Jinson, Dr. Sam Church, and members of the Marshall lab for constructive feedback and

support.

FIGURE LEGENDS

Figure 1. Study workflow to curate whole-cell imaging literature. PRISMA diagram detailing

the PubMed search strategy and screening steps used, with included and excluded study

counts at each stage. In search strategy, asterisk (*) indicates wildcard character, [MH]

indicates Mesh header, [tiab] indicates title/abstract field, [pt] indicates publication type.

Figure 2. Scientometric summary of the whole-cell imaging literature. A) Histogram of total

publication frequency by year. B) Open access (OA) publishing frequency by year. C)

Publication type by year. D) Data availability, as reported, by year. Data are represented

between 2004 and May 2024.

Figure S1. Bibliometric network analysis. A-B) Co-authorship network consisting of authors

with A) any number of publications and B) those with at least two publications represented in the

corpus. Color gradient represents average publication year per author and bubble size indicates

relative number of articles per author. Visualizations created in VOSViewer. C) Citation network,

created in Litmaps, representing most connected papers at the top, and most recent papers on

the right. Bubble size reflects relative citation number, and arrows point to articles cited.

Figure 3. Summary of imaging methodology and resolution. A) The number of studies

using each of the 13 imaging modalities. B) Box-and-whisker plots of reported lateral and axial

imaging resolutions, in nanometers (nm), grouped by modality type: electron microscopy (EM),

X-ray, or light microscopy (LM). A single value was plotted for each dataset. If multiple values or

a range of values were reported, the value plotted is the median of the reported values. C)

Scatterplot of axial vs lateral resolution in nanometers for each dataset, where reported, colored

by modality. Error bars represent the reported range, when applicable, with the datapoint

representing the median. The dotted line represents isotropic resolution (axial = lateral). Right

panel represents datasets for which only axial resolution was reported.

Figure 4. A) Datasets summarized according to cell type and modality. Each bubble represents

one dataset, and bubble size represents reported sample size (refer to legend, top left), colored

according to modality for clearly-reported sample size, and white for minimum known sample

size when not clearly reported. B) Distribution of sample sizes, according to minimum known

sample size. Inset show distribution for studies with sample sizes <=10.

Figure 5. Representation of common organelles across datasets. A) Organelle data collected

for each cell type represented across 118 datasets, excluding specialized or cell type-specific

organelles. Heatmap color represents the number of independent datasets in which the

organelle appears. Summary bar graphs show the total number of organelle instances per cell

type (right) and total number of instances per organelle across cell types (bottom) (Nuc –

nucleus, Mito – mitochondria, Vac/Lyso – vacuole/lysosome, ER – endoplasmic reticulum, LD –

lipid droplet, Golgi – Golgi body, Plast – plastid/chloroplast, Ves – vesicle, Cilia – cilia/flagella,

Perox – peroxisome, Cent – centrosome/centriole/basal body, Endo – endosome, MVB –

multivesicular body, Autophag – autophagosome.) B) Organelle combinations. Heatmap

represents the number of instances of each organelle pair co-imaged in the same dataset. C)

Distribution of anatomical completeness, defined by the number of distinct common organelles

coinciding in a single dataset.

Figure 6. A) Quantitative analyses performed on whole-cell reconstruction data. Dimensions,

density, positioning/distance, contact/association, and shape are all groupings including

different measurements conveying similar properties. (No quant – studies with no quantitative

analysis; RI – refractive index) B) Frequency of comparator classes by study. Within-study

comparable conditions or perturbations were grouped into general classes. ("No comp" –

studies with no comparator; Life Cycle/Dev/Dif – stages of life cycle, development, or

differentiation)

ADDITIONAL FILES

Additional File 1. Step by step details of corpus construction, with each step of searching and

screening listed in a separate tab. 0_Initial Target Papers: list of initial papers and extracted

features exemplifying target article types, used to construct PubMed Search Strategy.

1_Pubmed Search (2005-5.28.24): Complete list of 11,104 PubMed search results. 2_Title

Screening: PubMed metadata for 570 titles remaining after first screening step based on titles

and PubMed summary statements. 3_Abstract Screening: Results of second screening step

based on abstracts with PMID, inclusion decision (Include (I), Exclude (E), and Check Full Text

(C) and brief justification for each study. 4a_Full Text Screening Includes: Metadata for 127

studies included after full text screening. 4b_Full Text Screening Excludes: Metadata for 256

studies excluded after full text screening. 5_Post hoc additions: Metadata for 1 study included

after completion of bulk screening.

Additional File 2. Complete data extraction information for included studies after full text

screening, with full includes (Data Extraction_Includes) and borderlines (Data

Extraction_Borderlines) separated.

Additional File 3. Publicly available datasets. Parameters for 12 studies reporting public data

availability, including dataset completeness, study design, links, amount and format of data.

Additional File 4. Complete information for 118 datasets. Each individual dataset from 89

studies is listed including specifics of imaging modality, resolution, cell type and taxonomy,

sample size, and organelles.

Additional File 5. Data underlying analysis plotted in figures. Specific subsets of data used to

plot figures is separated into tabs.

REFERENCES

Antao NV, Lam C, Davydov A, Riggi M, Sall J, Petzold C, Liang FX, Iwasa JH, Ekiert DC, &

Bhabha G. (2023). 3D reconstructions of parasite development and the intracellular

niche of the microsporidian pathogen Encephalitozoon intestinalis. *Nat Commun*, *14*(1),

7662. https://doi.org/10.1038/s41467-023-43215-0

Arksey, H., & O'Malley, L. (2005). Scoping studies: Towards a methodological framework.

International Journal of Social Research Methodology, *8*(1), 19–32.

https://doi.org/10.1080/1364557032000119616

Bolaños, F., Salatino, A., Osborne, F., & Motta, E. (2024). Artificial intelligence for literature

reviews: Opportunities and challenges. *Artificial Intelligence Review*, *57*(10), 259.

https://doi.org/10.1007/s10462-024-10902-3

Bray, M.-A., Singh, S., Han, H., Davis, C. T., Borgeson, B., Hartland, C., Kost-Alimova, M.,

Gustafsdottir, S. M., Gibson, C. C., & Carpenter, A. E. (2016). Cell Painting, a high-

content image-based assay for morphological profiling using multiplexed fluorescent

dyes. *Nature Protocols*, *11*(9), 1757–1774. https://doi.org/10.1038/nprot.2016.105

Cell Line Catalog. (2024). ALLEN CELL EXPLORER. https://www.allencell.org/cell-catalog.html

Chen, J.-H. F., Mirvis, M., Ekman, A., Vanslembrouck, B., Gros, M. L., Larabell, C., & Marshall,

W. F. (2024). *Automated segmentation of soft X-ray tomography: Native cellular*

structure with sub-micron resolution at high throughput for whole-cell quantitative

imaging in yeast (p. 2024.10.31.621371). bioRxiv.

https://doi.org/10.1101/2024.10.31.621371

Cho, N. H., Cheveralls, K. C., Brunner, A.-D., Kim, K., Michaelis, A. C., Raghavan, P.,

Kobayashi, H., Savy, L., Li, J. Y., Canaj, H., Kim, J. Y. S., Stewart, E. M., Gnann, C.,

McCarthy, F., Cabrera, J. P., Brunetti, R. M., Chhun, B. B., Dingle, G., Hein, M. Y., …

Leonetti, M. D. (2022). OpenCell: Endogenous tagging for the cartography of human

cellular organization. *Science*, *375*(6585), eabi6983.

https://doi.org/10.1126/science.abi6983

Cock, P. J. A., Antao, T., Chang, J. T., Chapman, B. A., Cox, C. J., Dalke, A., Friedberg, I.,

Hamelryck, T., Kauff, F., Wilczynski, B., & de Hoon, M. J. L. (2009). Biopython: Freely

available Python tools for computational molecular biology and bioinformatics.

Bioinformatics, *25*(11), 1422–1423. https://doi.org/10.1093/bioinformatics/btp163

Cohen, S., Valm, A. M., & Lippincott-Schwartz, J. (2018). Interacting organelles. *Current*

Opinion in Cell Biology, *53*, 84–91. https://doi.org/10.1016/j.ceb.2018.06.003

Cortese, M., Lee, J.-Y., Cerikan, B., Neufeldt, C. J., Oorschot, V. M. J., Köhrer, S., Hennies, J.,

Schieber, N. L., Ronchi, P., Mizzon, G., Romero-Brey, I., Santarella-Mellwig, R., Schorb,

M., Boermel, M., Mocaer, K., Beckwith, M. S., Templin, R. M., Gross, V., Pape, C., …

Bartenschlager, R. (2020). Integrative Imaging Reveals SARS-CoV-2-Induced

Reshaping of Subcellular Morphologies. *Cell Host & Microbe*, *28*(6), 853-866.e5.

https://doi.org/10.1016/j.chom.2020.11.003

Cui Y, Cao W, He Y, Zhao Q, Wakazaki M, Zhuang X, Gao J, Zeng Y, Gao C, Ding Y, Wong

HY, Wong WS, Lam HK, Wang P, Ueda T, Rojas-Pierce M, Toyooka K, Kang BH, &

Jiang L. (2019). A whole-cell electron tomography model of vacuole biogenesis in

Arabidopsis root cells. *Nat Plants*, *5*(1), 95–105. https://doi.org/10.1038/s41477-018-

0328-1

Decelle J, Veronesi G, LeKieffre C, Gallet B, Chevalier F, Stryhanyuk H, Marro S, Ravanel S,

Tucoulou R, Schieber N, Finazzi G, Schwab Y, & Musat N. (2021). Subcellular

architecture and metabolic connection in the planktonic photosymbiosis between

Collodaria (radiolarians) and their microalgae. *Environ Microbiol*, *23*(11), 6569–6586.

https://doi.org/10.1111/1462-2920.15766

Donovan-Maiye, R. M., Brown, J. M., Chan, C. K., Ding, L., Yan, C., Gaudreault, N., Theriot, J.

A., Maleckar, M. M., Knijnenburg, T. A., & Johnson, G. R. (2022). A deep generative

model of 3D single-cell organization. *PLOS Computational Biology*, *18*(1), e1009155.

https://doi.org/10.1371/journal.pcbi.1009155

Earnest, T. M., Watanabe, R., Stone, J. E., Mahamid, J., Baumeister, W., Villa, E., & Luthey-

Schulten, Z. (2017). Challenges of Integrating Stochastic Dynamics and Cryo-Electron

Tomograms in Whole-Cell Simulations. *The Journal of Physical Chemistry B*, *121*(15),

3871–3881. https://doi.org/10.1021/acs.jpcb.7b00672

Ekman, A., Chen, J.-H., Dermott, G. M., Le Gros, M. A., & Larabell, C. (2020). Task Based

Semantic Segmentation of Soft X-ray CT Images Using 3D Convolutional Neural

Networks. *Microscopy and Microanalysis*, *26*(S2), 3152–3154.

https://doi.org/10.1017/S1431927620023983

Entrez Programming Utilities Help. (2010). National Center for Biotechnology Information (US).

https://www.ncbi.nlm.nih.gov/books/NBK25501/

Erozan, A., Lösel, P. D., Heuveline, V., & Weinhardt, V. (2024). Automated 3D cytoplasm

segmentation in soft X-ray tomography. *iScience*, *27*(6), 109856.

https://doi.org/10.1016/j.isci.2024.109856

Ezzedine, J. A., Uwizeye, C., Si Larbi, G., Villain, G., Louwagie, M., Schilling, M., Hagenmuller,

P., Gallet, B., Stewart, A., Petroutsos, D., Devime, F., Salze, P., Liger, L., Jouhet, J.,

Dumont, M., Ravanel, S., Amato, A., Valay, J.-G., Jouneau, P.-H., … Maréchal, E.

(2023). Adaptive traits of cysts of the snow alga Sanguina nivaloides unveiled by 3D

subcellular imaging. *Nature Communications*, *14*(1), 7500.

https://doi.org/10.1038/s41467-023-43030-7

Gillooly, J. F., Hein, A., & Damiani, R. (2015). Nuclear DNA Content Varies with Cell Size

across Human Cell Types. *Cold Spring Harbor Perspectives in Biology*, *7*(7), a019091.

https://doi.org/10.1101/cshperspect.a019091

Gluenz E, Wheeler RJ, Hughes L, & Vaughan S. (2015). Scanning and three-dimensional

electron microscopy methods for the study of Trypanosoma brucei and Leishmania

mexicana flagella. *Methods Cell Biol*, *127*, 509–542.

https://doi.org/10.1016/bs.mcb.2014.12.011

Guo, J., & Larabell, C. A. (2019). Soft X-ray tomography: Virtual sculptures from cell cultures.

Current Opinion in Structural Biology, *58*, 324–332.

https://doi.org/10.1016/j.sbi.2019.06.012

Hair, M., Yanase, R., Moreira-Leite, F., Wheeler, R. J., Sádlová, J., Volf, P., Vaughan, S., &

Sunter, J. D. (2024). Whole cell reconstructions of Leishmania mexicana through the cell

cycle. *PLOS Pathogens*, *20*(2), e1012054. https://doi.org/10.1371/journal.ppat.1012054

Hanson, M. A., Barreiro, P. G., Crosetto, P., & Brockington, D. (2024). The strain on scientific

publishing. *Quantitative Science Studies*, 1–21. https://doi.org/10.1162/qss_a_00327

Hanssen, E., Knoechel, C., Dearnley, M., Dixon, M. W. A., Le Gros, M., Larabell, C., & Tilley, L.

(2012). Soft X-ray microscopy analysis of cell volume and hemoglobin content in

erythrocytes infected with asexual and sexual stages of Plasmodium falciparum. *Journal*

of Structural Biology, *177*(2), 224–232. https://doi.org/10.1016/j.jsb.2011.09.003

Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D.,

Wieser, E., Taylor, J., Berg, S., Smith, N. J., Kern, R., Picus, M., Hoyer, S., van Kerkwijk,

M. H., Brett, M., Haldane, A., del Río, J. F., Wiebe, M., Peterson, P., … Oliphant, T. E.

(2020). Array programming with NumPy. *Nature*, *585*(7825), 357–362.

https://doi.org/10.1038/s41586-020-2649-2

Hein, M. Y., Peng, D., Todorova, V., McCarthy, F., Kim, K., Liu, C., Savy, L., Januel, C.,

Baltazar-Nunez, R., Bax, S., Vaid, S., Vangipuram, M., Ivanov, I. E., Byrum, J. R.,

Pradeep, S., Gonzalez, C. G., Aniseia, Y., Wang, E., Creery, J. S., … Leonetti, M. D.

(2023). *Global organelle profiling reveals subcellular localization and remodeling at*

proteome scale. https://doi.org/10.1101/2023.12.18.572249

Heinrich L, Bennett D, Ackerman D, Park W, Bogovic J, Eckstein N, Petruncio A, Clements J,

Pang S, Xu CS, Funke J, Korff W, Hess HF, Lippincott-Schwartz J, Saalfeld S, & Weigel

AV. (2021). Whole-cell organelle segmentation in volume electron microscopy. *Nature*,

599(7883), 141–146. https://doi.org/10.1038/s41586-021-03977-3

Henderson, G. P., Gan, L., & Jensen, G. J. (2007). 3-D Ultrastructure of O. tauri: Electron

Cryotomography of an Entire Eukaryotic Cell. *PLoS ONE*, *2*(8), e749.

https://doi.org/10.1371/journal.pone.0000749

Hunter, J. D. (2007). Matplotlib: A 2D Graphics Environment. *Computing in Science &*

Engineering, *9*(3), 90–95. Computing in Science & Engineering.

https://doi.org/10.1109/MCSE.2007.55

Koutsogiannis Z, Mina JGM, Suman R, & Denny PW. (2023). Assessment of Toxoplasma gondii

lytic cycle and the impact of a gene deletion using 3D label-free optical diffraction

holotomography. *Front Cell Infect Microbiol*, *13*, 1237594.

https://doi.org/10.3389/fcimb.2023.1237594

Larabell, C. A., & Le Gros, M. A. (2004). X-ray Tomography Generates 3-D Reconstructions of

the Yeast, Saccharomyces cerevisiae, at 60-nm Resolution. *Molecular Biology of the*

Cell, *15*(3), 957–962. https://doi.org/10.1091/mbc.e03-07-0522

Laughlin, R. B., Pines, D., Schmalian, J., Stojković, B. P., & Wolynes, P. (2000). The middle

way. *Proceedings of the National Academy of Sciences*, *97*(1), 32–37.

https://doi.org/10.1073/pnas.97.1.32

Laundon, D., Larson, B. T., McDonald, K., King, N., & Burkhardt, P. (2019). The architecture of

cell differentiation in choanoflagellates and sponge choanocytes. *PLOS Biology*, *17*(4),

e3000226. https://doi.org/10.1371/journal.pbio.3000226

Le Gros, M. A., McDermott, G., & Larabell, C. A. (2005). X-ray tomography of whole cells.

Current Opinion in Structural Biology, *15*(5), 593–600.

https://doi.org/10.1016/j.sbi.2005.08.008

Litmaps. (2024). Litmaps. https://app.litmaps.com

Loconte, V., Chen, J.-H., Cortese, M., Ekman, A., Le Gros, M. A., Larabell, C., Bartenschlager,

R., & Weinhardt, V. (2021). Using soft X-ray tomography for rapid whole-cell quantitative

imaging of SARS-CoV-2-infected cells. *Cell Reports Methods*, *1*(7), 100117.

https://doi.org/10.1016/j.crmeth.2021.100117

Loconte, V., Chen, J.-H., Vanslembrouck, B., Ekman, A. A., McDermott, G., Le Gros, M. A., &

Larabell, C. A. (2023). Soft X-ray tomograms provide a structural basis for whole-cell

modeling. *The FASEB Journal*, *37*(1), e22681. https://doi.org/10.1096/fj.202200253R

Loconte V, Singla J, Li A, Chen JH, Ekman A, McDermott G, Sali A, Le Gros M, White KL, &

Larabell CA. (2022). Soft X-ray tomography to map and quantify organelle interactions at

the mesoscale. *Structure*, *30*(4), 510-521.e3. https://doi.org/10.1016/j.str.2022.01.006

Lundberg, E., & Borner, G. H. H. (2019). Spatial proteomics: A powerful discovery tool for cell

biology. *Nature Reviews Molecular Cell Biology*, *20*(5), 285–302.

https://doi.org/10.1038/s41580-018-0094-y

Majarian, T. D., Cao-Berg, I., Ruan, X., & Murphy, R. F. (2019). CellOrganizer: Learning and

Using Cell Geometries for Spatial Cell Simulations. In W. S. Hlavacek (Ed.), *Modeling*

Biomolecular Site Dynamics (Vol. 1945, pp. 251–264). Springer New York.

https://doi.org/10.1007/978-1-4939-9102-0_11

McDermott, G., Le Gros, M. A., & Larabell, C. A. (2012). Visualizing Cell Architecture and

Molecular Location Using Soft X-Ray Tomography and Correlated Cryo-Light

Microscopy. *Annual Review of Physical Chemistry*, *63*(1), 225–239.

https://doi.org/10.1146/annurev-physchem-032511-143818

McQuate A, Knecht S, & Raible DW. (2023). Activity regulates a cell type-specific mitochondrial

phenotype in zebrafish lateral line hair cells. *Elife*, *12*.

https://doi.org/10.7554/eLife.80468

Meta-analysis in basic biology. (2016). *Nature Methods*, *13*(12), 959–959.

https://doi.org/10.1038/nmeth.4102

Mocaer K, Mizzon G, Gunkel M, Halavatyi A, Steyer A, Oorschot V, Schorb M, Le Kieffre C, Yee

DP, Chevalier F, Gallet B, Decelle J, Schwab Y, & Ronchi P. (2023). Targeted volume

correlative light and electron microscopy of an environmental marine microorganism. *J*

Cell Sci, *136*(15). https://doi.org/10.1242/jcs.261355

Moore, J., Allan, C., Besson, S., Burel, J.-M., Diel, E., Gault, D., Kozlowski, K., Lindner, D., Linkert, M., Manz, T., Moore, W., Pape, C., Tischer, C., & Swedlow, J. R. (2021). OME-NGFF: A next-generation file format for expanding bioimaging data-access strategies. *Nature Methods*, *18*(12), Article 12. https://doi.org/10.1038/s41592-021-01326-w

Moore, J., Basurto-Lozada, D., Besson, S., Bogovic, J., Bragantini, J., Brown, E. M., Burel, J.-M., Casas Moreno, X., de Medeiros, G., Diel, E. E., Gault, D., Ghosh, S. S., Gold, I., Halchenko, Y. O., Hartley, M., Horsfall, D., Keller, M. S., Kittisopikul, M., Kovacs, G., … Swedlow, J. R. (2023). OME-Zarr: A cloud-optimized bioimaging file format with international community support. *Histochemistry and Cell Biology*, *160*(3), 223–251. https://doi.org/10.1007/s00418-023-02209-1

Müller, A., Schmidt, D., Xu, C. S., Pang, S., D'Costa, J. V., Kretschmar, S., Münster, C., Kurth, T., Jug, F., Weigert, M., Hess, H. F., & Solimena, M. (2021). 3D FIB-SEM reconstruction of microtubule–organelle interaction in whole primary mouse β cells. *Journal of Cell Biology*, *220*(2), e202010039. https://doi.org/10.1083/jcb.202010039

Munn, Z., Pollock, D., Barker, T. H., Stone, J., Stern, C., Aromataris, E., Schünemann, H. J., Clyne, B., Khalil, H., Mustafa, R. A., Godfrey, C., Booth, A., Tricco, A. C., & Pearson, A. (2023). The Pandora's Box of Evidence Synthesis and the case for a living Evidence Synthesis Taxonomy. *BMJ Evidence-Based Medicine*, *28*(3), 148–150. https://doi.org/10.1136/bmjebm-2022-112065

Munn, Z., Stern, C., Aromataris, E., Lockwood, C., & Jordan, Z. (2018). What kind of systematic review should I conduct? A proposed typology and guidance for systematic reviewers in the medical and health sciences. *BMC Medical Research Methodology*, *18*(1), Article 1. https://doi.org/10.1186/s12874-017-0468-4

Murphy, R. F. (2012). CellOrganizer: Image-Derived Models of Subcellular Organization and Protein Distribution. In *Methods in Cell Biology* (Vol. 110, pp. 179–193). Elsevier. https://doi.org/10.1016/B978-0-12-388403-9.00007-2

Opalek, M., Tutaj, H., Pirog, A., Smug, B. J., Rutkowska, J., & Wloch-Salamon, D. (2023). A

Systematic Review on Quiescent State Research Approaches in S. cerevisiae. *Cells*,

12(12), Article 12. https://doi.org/10.3390/cells12121608

Paredes-Santos TC, de Souza W, & Attias M. (2012). Dynamics and 3D organization of

secretory organelles of Toxoplasma gondii. *J Struct Biol*, *177*(2), 420–430.

https://doi.org/10.1016/j.jsb.2011.11.028

Pautasso, M. (2012). Publication Growth in Biological Sub-Fields: Patterns, Predictability and

Sustainability. *Sustainability*, *4*(12), Article 12. https://doi.org/10.3390/su4123234

Pokrovskaya I, Tobin M, Desai R, Aronova MA, Kamykowski JA, Zhang G, Joshi S, Whiteheart

SW, Leapman RD, & Storrie B. (2021). Structural analysis of resting mouse platelets by

3D-EM reveals an unexpected variation in α-granule shape. *Platelets*, *32*(5), 608–617.

https://doi.org/10.1080/09537104.2020.1799970

Qin, Y., Huttlin, E. L., Winsnes, C. F., Gosztyla, M. L., Wacheul, L., Kelly, M. R., Blue, S. M.,

Zheng, F., Chen, M., Schaffer, L. V., Licon, K., Bäckström, A., Vaites, L. P., Lee, J. J.,

Ouyang, W., Liu, S. N., Zhang, T., Silva, E., Park, J., … Ideker, T. (2021). A multi-scale

map of cell structure fusing protein images and interactions. *Nature*, *600*(7889), 536–

542. https://doi.org/10.1038/s41586-021-04115-9

Rafelski, S. M., & Theriot, J. A. (2024). Establishing a conceptual framework for holistic cell

states and state transitions. *Cell*, *187*(11), 2633–2651.

https://doi.org/10.1016/j.cell.2024.04.035

Rohban, M. H., Singh, S., Wu, X., Berthet, J. B., Bray, M.-A., Shrestha, Y., Varelas, X., Boehm,

J. S., & Carpenter, A. E. (2017). Systematic morphological profiling of human gene and

allele function via Cell Painting. *eLife*, *6*, e24060. https://doi.org/10.7554/eLife.24060

Rudlaff, R. M., Kraemer, S., Marshman, J., & Dvorin, J. D. (2020). Three-dimensional

ultrastructure of Plasmodium falciparum throughout cytokinesis. *PLOS Pathogens*,

16(6), e1008587. https://doi.org/10.1371/journal.ppat.1008587

Rutkowska, J., Lagisz, M., Bonduriansky, R., & Nakagawa, S. (2020). Mapping the past, present

and future research landscape of paternal effects. *BMC Biology*, *18*(1), 183.

https://doi.org/10.1186/s12915-020-00892-3

Sakaguchi M, Miyazaki N, Fujioka H, Kaneko O, & Murata K. (2016). Three-dimensional

analysis of morphological changes in the malaria parasite infected red blood cell by

serial block-face scanning electron microscopy. *J Struct Biol*, *193*(3), 162–171.

https://doi.org/10.1016/j.jsb.2016.01.003

Scorrano, L., De Matteis, M. A., Emr, S., Giordano, F., Hajnóczky, G., Kornmann, B., Lackner,

L. L., Levine, T. P., Pellegrini, L., Reinisch, K., Rizzuto, R., Simmen, T., Stenmark, H.,

Ungermann, C., & Schuldiner, M. (2019). Coming together to define membrane contact

sites. *Nature Communications*, *10*(1), 1287. https://doi.org/10.1038/s41467-019-09253-3

Sear, R. P., Pagonabarraga, I., & Flaus, A. (2015). Life at the mesoscale: The self-organised

cytoplasm and nucleoplasm. *BMC Biophysics*, *8*(1), 4. https://doi.org/10.1186/s13628-

015-0018-6

Sercel, A. J., Sturm, G., Shaulson, E. D., Gallagher, D., St-Onge, M.-P., Kempes, C. P.,

Pontzer, H., Hirano, M., & Picard, M. (2023). *Accelerated physiology and increased*

energy expenditure in animals and humans with mitochondrial defects: A meta-analysis

(p. 2023.09.09.556754). bioRxiv. https://doi.org/10.1101/2023.09.09.556754

The data deluge. (2012). *Nature Cell Biology*, *14*(8), 775–775. https://doi.org/10.1038/ncb2558

The pandas development team. (2020). *pandas-dev/pandas: Pandas* (Version 2.2.3) [Computer

software]. https://doi.org/10.5281/zenodo.3509134

Trebichalská, Z., Javůrek, J., Tatíčková, M., Kyjovská, D., Kloudová, S., Otevřel, P., Hampl, A.,

& Holubcová, Z. (2021). High-Resolution 3D Reconstruction of Human Oocytes Using

Focused Ion Beam Scanning Electron Microscopy. *Frontiers in Cell and Developmental*

Biology, *9*. https://www.frontiersin.org/articles/10.3389/fcell.2021.755740

Tricco, A. C., Lillie, E., Zarin, W., O'Brien, K. K., Colquhoun, H., Levac, D., Moher, D., Peters,

M. D. J., Horsley, T., Weeks, L., Hempel, S., Akl, E. A., Chang, C., McGowan, J.,

Stewart, L., Hartling, L., Aldcroft, A., Wilson, M. G., Garritty, C., … Straus, S. E. (2018).

PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation.

Annals of Internal Medicine, *169*(7), 467–473. https://doi.org/10.7326/M18-0850

Tricco, A. C., Soobiah, C., Antony, J., Cogo, E., MacDonald, H., Lillie, E., Tran, J., D'Souza, J.,

Hui, W., Perrier, L., Welch, V., Horsley, T., Straus, S. E., & Kastner, M. (2016). A

scoping review identifies multiple emerging knowledge synthesis methods, but few

studies operationalize the method. *Journal of Clinical Epidemiology*, *73*, 19–28.

https://doi.org/10.1016/j.jclinepi.2015.08.030

Trumpff, C., Michelson, J., Lagranha, C. J., Taleon, V., Karan, K. R., Sturm, G., Lindqvist, D.,

Fernström, J., Moser, D., Kaufman, B. A., & Picard, M. (2021). Stress and circulating

cell-free mitochondrial DNA: A systematic review of human studies, physiological

considerations, and technical recommendations. *Mitochondrion*, *59*, 225–245.

https://doi.org/10.1016/j.mito.2021.04.002

Uchida, M., McDermott, G., Wetzler, M., Le Gros, M. A., Myllys, M., Knoechel, C., Barron, A. E.,

& Larabell, C. A. (2009). Soft X-ray tomography of phenotypic switching and the cellular

response to antifungal peptoids in *Candida albicans*. *Proceedings of the National*

Academy of Sciences, *106*(46), 19375–19380. https://doi.org/10.1073/pnas.0906145106

Uchida, M., Sun, Y., McDermott, G., Knoechel, C., Le Gros, M. A., Parkinson, D., Drubin, D. G.,

& Larabell, C. A. (2011). Quantitative analysis of yeast internal architecture using soft X-

ray tomography. *Yeast*, *28*(3), 227–236. https://doi.org/10.1002/yea.1834

Uwizeye, C., Decelle, J., Jouneau, P.-H., Flori, S., Gallet, B., Keck, J.-B., Bo, D. D., Moriscot,

C., Seydoux, C., Chevalier, F., Schieber, N. L., Templin, R., Allorent, G., Courtois, F.,

Curien, G., Schwab, Y., Schoehn, G., Zeeman, S. C., Falconet, D., & Finazzi, G.

(2021a). Morphological bases of phytoplankton energy management and physiological

responses unveiled by 3D subcellular imaging. *Nature Communications*, *12*(1), 1049.

https://doi.org/10.1038/s41467-021-21314-0

Uwizeye, C., Decelle, J., Jouneau, P.-H., Flori, S., Gallet, B., Keck, J.-B., Bo, D. D., Moriscot,

C., Seydoux, C., Chevalier, F., Schieber, N. L., Templin, R., Allorent, G., Courtois, F.,

Curien, G., Schwab, Y., Schoehn, G., Zeeman, S. C., Falconet, D., & Finazzi, G.

(2021b). Morphological bases of phytoplankton energy management and physiological

responses unveiled by 3D subcellular imaging. *Nature Communications*, *12*(1), 1049.

https://doi.org/10.1038/s41467-021-21314-0

Uwizeye C, Mars Brisbin M, Gallet B, Chevalier F, LeKieffre C, Schieber NL, Falconet D,

Wangpraseurt D, Schertel L, Stryhanyuk H, Musat N, Mitarai S, Schwab Y, Finazzi G, &

Decelle J. (2021). Cytoklepty in the plankton: A host strategy to optimize the

bioenergetic machinery of endosymbiotic algae. *Proc Natl Acad Sci U S A*, *118*(27).

https://doi.org/10.1073/pnas.2025252118

Valm, A. M., Cohen, S., Legant, W. R., Melunis, J., Hershberg, U., Wait, E., Cohen, A. R.,

Davidson, M. W., Betzig, E., & Lippincott-Schwartz, J. (2017). Applying systems-level

spectral imaging and analysis to reveal the organelle interactome. *Nature*, *546*(7656),

162–167. https://doi.org/10.1038/nature22369

van Eck, N. J., & Waltman, L. (2010). Software survey: VOSviewer, a computer program for

bibliometric mapping. *Scientometrics*, *84*(2), 523–538. https://doi.org/10.1007/s11192-

009-0146-3

Veritas Health Innovation. (n.d.). *Covidence systematic review software*. Covidence. Retrieved

February 3, 2025, from www.covidence.org

Viana, M. P., Chen, J., Knijnenburg, T. A., Vasan, R., Yan, C., Arakaki, J. E., Bailey, M., Berry,

B., Borensztejn, A., Brown, E. M., Carlson, S., Cass, J. A., Chaudhuri, B., Cordes

Metzler, K. R., Coston, M. E., Crabtree, Z. J., Davidson, S., DeLizo, C. M., Dhaka, S., …

Rafelski, S. M. (2023). Integrated intracellular organization and its variations in human

iPS cells. *Nature*, *613*(7943), 345–354. https://doi.org/10.1038/s41586-022-05563-7

Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D.,

Burovski, E., Peterson, P., Weckesser, W., Bright, J., van der Walt, S. J., Brett, M.,

Wilson, J., Millman, K. J., Mayorov, N., Nelson, A. R. J., Jones, E., Kern, R., Larson, E.,

… van Mulbregt, P. (2020). SciPy 1.0: Fundamental algorithms for scientific computing

in Python. *Nature Methods*, *17*(3), 261–272. https://doi.org/10.1038/s41592-019-0686-2

Voeltz, G. K., Sawyer, E. M., Hajnóczky, G., & Prinz, W. A. (2024). Making the connection: How

membrane contact sites have changed our view of organelle biology. *Cell*, *187*(2), 257–

270. https://doi.org/10.1016/j.cell.2023.11.040

Wang, S., & Mukherji, S. (2022). *Uncovering the principles coordinating systems-level organelle*

biogenesis with cellular growth [Preprint]. Systems Biology.

https://doi.org/10.1101/2022.11.01.514705

Waskom, M. L. (2021). seaborn: Statistical data visualization. *Journal of Open Source Software*,

6(60), 3021. https://doi.org/10.21105/joss.03021

Wei, D., Jacobs, S., Modla, S., Zhang, S., Young, C. L., Cirino, R., Caplan, J., & Czymmek, K.

(2012). High-resolution three-dimensional reconstruction of a whole yeast cell using

focused-ion beam scanning electron microscopy. *BioTechniques*, *53*(1), 41–48.

https://doi.org/10.2144/000113850

White, K. L., Singla, J., Loconte, V., Chen, J.-H., Ekman, A., Sun, L., Zhang, X., Francis, J. P.,

Li, A., Lin, W., Tseng, K., McDermott, G., Alber, F., Sali, A., Larabell, C., & Stevens, R.

C. (2020). Visualizing subcellular rearrangements in intact β cells using soft x-ray

tomography. *Science Advances*, *6*(50), eabc8262.

https://doi.org/10.1126/sciadv.abc8262

Yip, R., Sun, Y. J., Bassuk, A. G., & Mahajan, V. B. (2024). *Artificial Intelligence's Contribution*

to Biomedical Literature Search: Revolutionizing or Complicating?

https://doi.org/10.1101/2024.10.07.617112

Zechmann, B., Möstl, S., & Zellnig, G. (2022). Volumetric 3D reconstruction of plant leaf cells

using SEM, ion milling, TEM, and serial sectioning. *Planta*, *255*(6), 118.

https://doi.org/10.1007/s00425-022-03905-3

Zhang, Z. (2024). Expanding bioinformatics: Toward a paradigm shift from data to theory.

Fundamental Research. https://doi.org/10.1016/j.fmre.2024.11.019

Zhao, T., & Murphy, R. F. (2007). Automated learning of generative models for subcellular

location: Building blocks for systems biology. *Cytometry Part A*, *71A*(12), 978–990.

https://doi.org/10.1002/cyto.a.20487

Zotero | Your personal research assistant. (n.d.). Retrieved February 3, 2025, from

https://www.zotero.org/

1180

Figure 1.

Figure S1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.