Arbor function of TRANSPARENT TESTA GLABRA 1 and LIGHT REGULATED WD scaffold

proteins in the Arabidopsis circadian oscillators includes transcriptional repression through PSEUDO

RESPONSE REGULATORS

Authors

Eva Herrero[a], Dora Cano-Ramirez[ab], Beverley J. Glover[a] and Alex A. R. Webb[a]

[a]Department of Plant Sciences, University of Cambridge

[b]Current address Sainsbury Laboratory Cambridge University

Abstract

Arabidopsis circadian oscillators contain DNA-binding proteins that function at specific times of

the day. In contrast, we have discovered a unique continuous function of WD40-repeat scaffold

proteins from the TRANSPARENT TESTA GLABRA-1 (TTG1), LIGHT REGULATED WD1 and

LIGHT REGULATED LWD2 subfamily (TLWD) which is essential to maintain circadian rhythms.

Gene expression analyses indicate multifunctional activity of TLWD in both transcriptional activation

and repression. TLWD proteins interact with an array of circadian oscillator activators and repressors

that act sequentially throughout the diel cycle. While TLWD proteins were known to participate in

activator complexes, our data indicate a novel role of TLWD in transcriptional repression through

complex formation with repressors from the PSEUDO RESPONSE REGULATOR (PRR) family. In

an analogy to mechanical clocks, TLWD scaffold proteins constitute an arbor, or spindle, to which

transcription factors, which represent the cogs, bind to sustain circadian rhythms.

Introduction

Plant circadian oscillators regulate networks of gene activity to respond appropriately to the daily rhythms of the environment, contributing to fitness[1,2]. Consequently, plant circadian oscillators control critical yield traits such as flowering time, growth, and response to environmental stress[3]. Circadian oscillators contain arrays of transcription factors that are sequentially expressed through the diel cycle and orchestrate circadian rhythms of expression of over a third of the Arabidopsis transcriptome[4].

In the morning, Myb-like transcription factors contribute both to transcriptional repression, for example CIRCADIAN CLOCK ASSOCIATED (CCA1)[5] and LATE ELONGATED HYPOCOTYL (LHY)[6,7], and to transcriptional activation, for example REVEILLE8 (RVE8)[8] in Arabidopsis. The members of the PSEUDO RESPONSE REGULATOR (PRR) family PRR9[9], PRR7[10], PRR5[11] and PRR1 (also known as TIMING OF CAB EXPRESSION1 (TOC1))[12] work sequentially as transcriptional repressors from early morning to late night[13]. LUX ARRYTHMO (LUX), a member of the GOLDEN2, ARR-B, PSR1 (GARP) family, binds DNA as part of an Evening Complex of proteins to mediate transcriptional repression[14–16]. Additionally, members of the TEOSINTE BRANCHED 1, CYCLOIDEA, PCF1 (TCP) family TCP20 and TCP22 contribute to the activation of *CCA1*[17,18], whereas CCA1-HICKING EXPEDITION (CHE or TPC21) reduces *CCA1* expression[19]. Overall, oscillations of expression of circadian rhythmic genes (CRGs) are orchestrated by the sequential binding of transcription factors to specific promoter *cis*-elements, resulting in either transcriptional activation or repression.

Plant circadian oscillators also feature a large array of structurally diverse scaffold proteins. Scaffold proteins do not bind directly to DNA, instead they coordinate the formation of protein complexes through their multiple binding surfaces. For instance, EARLY-FLOWERING-3 (ELF3) nucleates the Evening Complex with ELF4 and LUX[16,20]. The scaffold protein GIGANTEA (GI) promotes high amplitude oscillations of PRR5 and TOC1 by modulating the activity of the F-Box protein ZEITLUPE (ZTL)[21–23]. Both ELF3 and GI function as hubs that integrate multiple environmental signals to control growth and development[14,24,25]. In Arabidopsis, scaffold proteins of the TRANSPARENT-TESTA-GLABRA-1 (TTG1) and LIGHT-REGULATED-WD40-repeat (LWD) family (TLWD) are an essential part of the circadian oscillator besides controlling many other aspects of plant biology[26–28]. Functional divergence in the TLWD family arose at the base of the seed plant lineage[27]. In Arabidopsis, TTG1 itself is an integral part of the Myb-bHLH-WDR (MBW) complexes that control pigment biosynthesis and epidermal cell differentiation[28] while LWD1 and LWD2 control circadian periodicity and light entrainment[26]. Importantly, LWD1, LWD2 and TTG1 all redundantly contribute to sustain circadian rhythmicity[27] since the *ttg1 lwd1 lwd2* triple mutant has arrhythmic circadian outputs such as gene expression and leaf movement. LWDs are required for TCP mediated activation of *CCA1* expression[17,18]. However, loss of TCP function does not lead to arrhythmicity but

only changes in amplitude of *CCA1* expression, suggesting additional roles of TLWD in circadian
oscillators.[17,18]

Here we report that TLWD proteins have a unique multifunctional role in the circadian
oscillator. We show that both LWDs and TTG1 interact with transcriptional repressors of the Myb-
like and PRR families, and with transcriptional activators of the TCP family. Specifically, we have
found that transcriptional repression mediated by TLWD-PRR complexes is essential to sustain
circadian oscillations and regulate circadian outputs. TLWD proteins represent a new category of
circadian oscillator component, which, unlike most plant circadian oscillator proteins, is not
temporally restricted in abundance. Because of their continuous presence, and their ability to bind
multiple different transcriptional repressors and activators throughout the 24 h cycle, we describe
TLWD proteins as arbor proteins, an arbor being the spindle of a mechanical clock or watch to which
the cogs and gears are attached.
76
77 **Results**
78
79 ***TLWD function is required for both activation and repression of circadian rhythmic genes***
80
We reported that TLWD function is essential to sustain circadian oscillations of *CCA1* gene
expression and leaf movements in Arabidopsis[27]. To understand more widely the impact of LWDs
and TTG1 in transcriptional regulation, we used RNAseq to evaluate changes in transcript
abundance between *ttg1*, *lwd1 lwd2* and *ttg1 lwd1 lwd2* mutant 10-day-old seedlings under neutral
photoperiod cycles (12L/12D) in the middle of the day (ZT6) and the middle of the night (ZT18). We
considered differentially expressed genes (DEG) those with a $Log2^{FC}$ >1.5 between the mutant and
the wild type at the fixed time point. Only a small number of genes were downregulated in the *ttg1*
mutant compared to wild type at ZT6. These corresponded to known direct targets of MBW
complexes involved in flavonoid biosynthesis[29] (Extended data Fig. 1A, Extended data Table 1),
demonstrating the ability of our approach to identify direct targets of TLWD transcriptional activity.
These direct targets of MBW complexes were also downregulated in the *ttg1 lwd1 lwd2* triple mutant
but not in the *lwd1 lwd2* double mutant (Extended data Fig. 1A, Extended data Table 1), consistent
with LWDs not complementing the TTG1 functions mediated by MBW complexes [27]. The majority of
DEGs in *lwd1 lwd2* (Day: 97% of the down-regulated and 80% of the upregulated genes, Night: 97%
of the upregulated genes) and in *ttg1 lwd1 lwd2* (Day: 57% of the downregulated genes and 75% of
the upregulated genes, Night: 45% of the downregulated genes and 79% of the upregulated genes)
were known circadian rhythmic genes (CRG)[4], indicating that the predominant activity of TLWD
proteins in young seedlings is circadian regulation. Consistent with the lack of circadian phenotypes
in the single *ttg1*[27] mutant, we did not find a defect in circadian regulated gene expression in *ttg1*
(Extended data Table 1).
The trend of misregulation of CRGs was comparable between *lwd1 lwd2* and *ttg1 lwd1 lwd2,*
but was more pronounced in the triple mutant, confirming genetic redundancy between TLWD

proteins (Extended data Fig.1B). At ZT6, from the 97 CRG downregulated in *ttg1 lwd1 lwd2* mutants

(Log2^{FC} >-1.5), 62 were also downregulated in *lwd1 lwd2* (Log2^{FC} >-0.5) (Extended data Table 1).

We used the CAST-R database[30] to analyse the wild type circadian phase of expression of

misregulated CRGs in *lwd1 lwd2* and *ttg1 lwd1 lwd2* mutants (Figure 1A, Extended data Table 1).

Downregulated CRGs at ZT6 in *lwd1 lwd2* and *ttg1 lwd1 lwd2* usually had a morning circadian phase

(ZT0-6). In addition, 47 out of 97 CRGs downregulated in *ttg1 lwd1 lwd2* usually had a late morning

circadian phase ZT6-ZT8 (Fold enrichment 3.89-3.78, p-value <0.05). The lack of activation of CRGs

in the morning in *TLWD* mutants is consistent with the known role of LWDs in activating gene

expression in the morning.[17,18]

Additionally, we found defects of circadian-associated transcriptional repression both in the

day and in the night (Figure 1A, Extended data Table 1). The majority of CRGs upregulated in *ttg1*

*lwd1 lwd2 (*Log2^{FC} >1.5) were also upregulated in *lwd1 lwd2 (*Log2^{FC} >0.5), both at ZT6 (66/86) and

at ZT18 (87/111) (Extended data Table 1). CRGs upregulated in *lwd1 lwd2* at ZT6 usually had a

circadian phase of expression of ZT14 (Figure 1A, Extended data Table 1; Fold enrichment

ZT6=1.92, p value < 0.005). CRGs upregulated in *ttg1 lwd1 lwd2* at ZT6 have a usual circadian

phase of expression of ZT14-ZT20 (Figure 1A, Extended data Table 1; ZT (Fold enrichment): ZT14

(2.4), ZT16 (2.4), ZT18 (2), ZT20 (1.9), p-value < 0.05). Conversely, CRGs upregulated in *lwd1 lwd2*

and *ttg1 lwd1 lwd2* mutants at ZT18 normally have a morning circadian phase of expression (ZT2 to

ZT6) (Figure 1A, Extended data Table1, ZT (Fold enrichment): *lwd1 lwd2* ZT2 (1.8), ZT4 (2.8), ZT6

(4.6); *ttg1 lwd1 lwd2* ZT4 (4.4), ZT6 (6.3), p value < 0.05). These data suggest an important but

previously overlooked role of TLWD in transcriptional repression at different times of the day.

Most components of the Arabidopsis circadian oscillator were misexpressed in *lwd1 lwd2*

and *ttg1 lwd1 lwd2* mutants. Morning genes *CCA1*, *PRR9* and *LNK2* were downregulated at ZT6

and upregulated at ZT18 in comparison to wild type, whereas evening expressed genes *TOC1* and

ELF4 were upregulated at ZT6. *LUX* was upregulated both at ZT6 and ZT18 (Figure 1B, Extended

data Fig. 1C and Table 1). Taken together, we have identified a dual role of TLWD in the circadian

oscillator mediating transcriptional activation in the morning and transcriptional repression in the

morning and at night. To look more closely at the temporal aspects of the effects of TLWD function

we analysed rhythms of activity of *CCA1*, *TOC1* and *PRR7* promoters fused to firefly luciferase (LUC)

both under diel and free-running conditions. In the *ttg1* mutant, we saw no defect in *pCCA1::LUC*

relative to wild type (Figure 1C*)*, confirming that TTG1 function in circadian regulation is redundant

to that of the LWDs[27]. Both *lwd1 lwd2* and *ttg1 lwd1 lwd2* mutants had reduced amplitude of

pCCA1::LUC activity under diel cycles (Figure 1D). In *ttg1 lwd1 lwd2* there was a second peak of

pCCA1::LUC in the second half of the day. This second peak was only apparent under long day

photoperiods (Figure 1D) but not under neutral or short-day photoperiods (Extended data Fig.2),

indicating a lack of repression which allows light to induce *CCA1* promoter activity. Under free-

running conditions, oscillations of *pCCA1::LUC* became arrhythmic from that second peak in *ttg1*

lwd1 lwd2 plants (Figure 1D). These data indicate that TLWD have a dual function in sustaining

CCA1 oscillations. In the morning LWDs activate *CCA1* expression [17,18] and in the second part of

the day both LWDs and TTG1 repress *CCA1* expression. The activity of both *pPRR7* and *pTOC1*

was increased in *ttg1 lwd1 lwd2* plants relative to wild type through the light phase under diel cycles

and arrhythmic under free-running conditions (Figure 1E-F), supporting an essential role of TLWD

in circadian mediated transcriptional repression throughout the diel cycle.

TLWD proteins are present through the diel cycle

The abundance and/or activity of plant circadian oscillator components is restricted to specific

times of the day. However, our data are consistent with a continuous function of TLWD proteins

through the diel cycle. To test whether LWD1 and TTG1 proteins oscillate in abundance through the

diel cycle, we generated complementation lines of *lwd1 lwd2* and *ttg1* mutants with LWD1-Venus or

Venus-TTG1 flanked by their respective promoters and downstream sequence, respectively. For

LWD1, we selected complementation lines that fully complemented *lwd1 lwd2 pCCA1::LUC*

oscillations (Extended data Fig. 3). For TTG1, we selected complementation lines that restored

trichomes and seed coat pigmentation of *ttg1* mutants (Extended data Fig. 4), indicating that the

addition of the Venus tag did not compromise protein function. We confirmed the nuclear localization

of LWD1-Venus and TTG1-Venus in the corresponding complementation lines (Extended data Fig.

3 and 4). We obtained whole protein extracts of these lines every 4 hours during one diel cycle and

detected LWD1-Venus or TTG1-Venus by western blot. To confirm that our approach can detect

oscillations in protein abundance, we included a previously described *pCCA1::CCA1-YFP*[31] line in

our analysis. We used the same anti-GFP antibody for detecting CCA1, LWD1 and TTG1. The

presence of CCA1-YFP protein was restricted to early day with a peak at ZT4 (Figure 2A). In contrast,

both LWD1-Venus and TTG1-Venus were present throughout the day and night at similar levels

(Figure 2 B-C). The presence of two bands with similar molecular weight could indicate either two

isoforms or protein modification. Additionally, we generated complementation lines with LWD1 and

TTG1 fused to LUC in the *lwd1 lwd2* and *ttg1* mutant lines (see complementation data for TTG1-

LUC lines in Extended data Fig.4C). We detected low amplitude oscillations of both LWD1-LUC and

TTG1-LUC with a first peak at dawn and a second peak around dusk under diel cycles (Figure 2D-

E). There was a decrease of both LWD1-LUC and TTG1-LUC during the dark period. Low amplitude

oscillations of both LWD1 and TTG1 persisted under continuous light. Taken together, both LWD1

and TTG1 are present through the diel cycle with low amplitude oscillations under diel cycles, being

slightly more abundant during the day.

TLWD proteins interact with both circadian oscillator activators and repressors

ttg1 lwd1 lwd2 mutants are arrhythmic [27]. However, why this might be the case has been

unclear because the only known interactors with the TLWD proteins were the TCP proteins. The

effects of *ttg1 lwd1 lwd2* cannot be solely due to lack of transcriptional activity of LWD-TCP

complexes at dawn, since *tcp* mutants only have defects in amplitude of *pCCA1* activity and retain

full rhythmicity[17]. Our transcriptome and promoter activity data indicated other potential roles for
TLWD proteins in transcriptional repression (Figure 1), which we hypothesized might explain why
the triple mutant is arrhythmic. We tested whether TLWD proteins interact with additional clock
transcription factors, particularly the transcriptional repressors that are active throughout the day. In
yeast 2 hybrid assays, TCP20, TCP22 and CHE interacted with LWD1 and LWD2, as previously
reported[17]. We also found interaction of TCP20, TCP22 and CHE with TTG1 (Extended data Fig. 5).
Additionally, LWD1, LWD2 and TTG1 all interacted with PRR5 (Figure 3A). TTG1 itself also
interacted with PRR9, PRR7, TOC1 and PRR3 (Figure 3A). Finally, TTG1 interacted with Myb-like
transcriptional repressors CCA1, LHY and RVE1, but not with the activator RVE8 (Extended data
Fig. 6A). These data are consistent with a continuous function of TLWD proteins throughout the
circadian cycle, forming complexes with different transcriptional repressors and activators that act
sequentially through the day and night in a sequence that starts with CCA1/LHY after dawn, moves
through the PRRs and ends with the TCPs just around dawn. To test predictions of the yeast two
hybrid data *in planta* we performed co-immunoprecipitation in *N. benthamiana* using transient
expression. Both LWD1 and TTG1 interacted with PRR9, PRR7 and PRR5. Also, LWD1 interacted
with TOC1 (Figure 3B-C). Differences in the interaction ability between LWD1 and TTG1 with PRR
proteins in our Y2H or Co-IP experiments may be caused by differences in posttranslational
modifications between both systems. In summary, we found that both LWD1 and TTG1 interact with
multiple PRR proteins *in planta,* implying that multiple different TLWD-PRR complexes may be
formed during progress through the diel cycle*.*

Similarly to *ttg1 lwd1 lwd2* mutants, triple *prr* loss of function mutants are arrhythmic[32] .
Hence, it is possible that loss of PRR repression, in addition to the previously identified loss of TCP
activation[17], might account for the arrhythmicity in *ttg1 lwd1 lwd2* triple loss of function mutants. We
reasoned that potential direct targets of TLWD-PRR complexes would be upregulated both in higher
order *tlwd* and *prr* mutants due to a common defect in transcriptional repression. Hence, we
compared CRGs that were upregulated both in our RNAseq experiment at ZT18 in both *lwd1 lwd2*
and *ttg1 lwd1 lwd2* with CRGs upregulated in *prr5 toc1* mutants at a similar time ZT15 (Log2^{FC}>1)[33].
We found that 49 genes were shared between these two datasets (Fold enrichment (p value): *lwd1*
lwd2 62.07 (p 5.9 e^{-63}) and *ttg1 lwd1 lwd2* 36.33 (5.9 e^{-63}) (Figure 3D). Of these 49 genes, 9 were
also direct targets of both PRR5[34] and TOC1[12] repression as established by previous CHIP-seq
experiments (Figure 3E, F). Both *PRR7* and *CCA1* were potential targets of TLWD-PRR complexes
(Figure 3F). Consistently, we found a lack of repression of both *pCCA1* and *pPRR7* in *ttg1 lwd1 lwd2*
mutants (Figure 1D and 1F). Notably, we found that both *prr7 prr5* and *prr7 prr5 toc1* had a second
peak of *pCCA1* activity in the second half of the diel cycle and under free-running conditions *pCCA1*
activity remained high (Figure 3G), phenocopying the characteristic *pCCA1* expression of *ttg1 lwd1*
lwd2 mutants (Figure 1D). The arrhythmic mutant *elf3* did not phenocopy *ttg1 lwd1 lwd2* aberrant
pCCA1 oscillations under diel cycles (Figure 3H), supporting the idea that loss of activity of PRR
complexes leads to the arrhythmicity in *ttg1 lwd1 lwd2* triple mutant plants.

The Y2H data also suggested that TLWD proteins interact with Myb-like transcription factors
(Extended data Fig. 6A). To test whether a defect in CCA1 /LHY activity might contribute to the
arrhythmicity of *ttg1 lwd1 lwd2* mutants, in addition to our analysis of pCCA1 (Figure1C, D), we
compared rhythms of *pTOC1::LUC* under diel cycles and free running conditions in *ttg1 lwd1 lwd2*
and lhy cca1 mutants as *TOC1* is a direct target of CCA1/LHY repression. We also included in the
comparison *elf3* which is arrhythmic and has low levels of *CCA1/LHY* expression[20,35], similar to *ttg1*
lwd1 lwd2 mutants[27] (Figure 1D). Wild type plants had a small transient peak of *pTOC1* at dawn and
a second main peak around dusk. *ttg1 lwd1 lwd2* and *lhy cca1* mutants had similarly high levels of
pTOC1 expression after the dawn peak until dusk where *pTOC1* levels decreased (Extended data
Fig 6B). The *elf3* mutant was different since *pTOC1* was clearly repressed after the dawn peak and
the second peak was only slightly advanced (Extended data Fig. 6B). These data are consistent with
TLWD contributing to CCA1/LHY repression of *TOC1*.
231
232   ***TLWD proteins control growth through repression of* PIF4/5 *expression***
233
Arabidopsis circadian oscillators provide a temporal framework to alter growth in response to
environmental signals by generating circadian rhythms of expression of the growth promoting
transcription factors *PHYTOCHROME INTERACTING FACTOR4* (*PIF4*) and *PIF5* [16,36]. In our
transcriptome analysis, we found that *PIF4* and *PIF5* are upregulated specifically in the *ttg1 lwd1*
lwd2 mutant at night (ZT18), together with various direct targets of PIF4 and PIF5 activation such as
CYCLING DOF FACTOR 5 (CDF5) and *ARABIDOPSIS THALIANA HOMEOBOX PROTEIN 2*
(ATBH2) (Figure 4A, Extended data Fig. 7A and Table 1). Both *PIF4* and *PIF5* are also directly
repressed by PRR7, PRR5 and TOC1, and are consequently potential targets of TLWD-PRR
complexes (Figure 3F). The increase in expression of *PIF4/5* and its targets was accompanied by
an increase in hypocotyl elongation under both short day (SD) and long day (LD) photoperiods in
ttg1 lwd1 lwd2 triple mutants (Figure 4B-C). Importantly, *lwd1 lwd2* double mutants and *ttg1* single
mutants had similar hypocotyl lengths to the wild type, indicating that all three TLWD proteins work
redundantly to repress growth. The hypocotyls of *ttg1 lwd1 lwd2* mutants were longer than wild type
but not as long as those of *prr7 prr5 toc1, prr9 prr7 prr5* and *elf3* mutants (Figure 4B-C, Extended
data Fig. 7B). We tested whether hypocotyl length correlated with expression of *PIFs* and target
genes under short day photoperiods. Consistent with a role of TLWD in transcriptional repression at
night, *PIF4*, *PIF5* and *CDF5* were upregulated in *ttg1 lwd1 lwd2* plants particularly in the second half
of the diel cycle. Expression levels were lower in *ttg1 lwd2 lw2* mutants than in *prr7 prr5 toc1* plants,
which correlated with their respective hypocotyl lengths (Figure 4D). Other well documented
phenotypes of increased *PIF4* expression are elongated petioles and smaller leaves when grown
under short day photoperiods. While *PIF4-OX*, *elf3* and *ttg1 lwd1 lwd2* plants had elongated petioles
and smaller leaves, both *ttg1* and *lwd1 lwd2* plants were similar to wild type (Extended data Fig. 7C).
Taken together, these data show that TLWD contributes to the repression of PIF4/5 to control
hypocotyl elongation.

Discussion

Unique continuous function of TLWD in Arabidopsis circadian oscillators

Our work highlights the importance of scaffold proteins in circadian oscillators. In the context of transcriptional regulation, scaffold proteins mediate the formation of multiprotein complexes which are essential for the function of transcription factors[16,20]. In Arabidopsis, ELF3 protein is restricted to evening between ZT10 and ZT20[16,37] and nucleates the evening complex that represses *CCA1*, *PRR9*, *PRR7* and *GI*. GI protein can be detected around dusk[38,39], when it forms complexes with ZTL and modulates TOC1 abundance[22,23,40]. In contrast to the time-restricted activity of ELF3 or GI scaffolds, we have found a unique continuous presence of TLWD proteins in the circadian oscillator. TLWD proteins are present throughout the diel cycle (Figure 2) and uniquely form different complexes with multiple transcription factors of different families (Figure 3, Extended data Fig. 5-6) that work sequentially to control either the activation or repression of target genes (Figure 5). Consistent with a continuous function of TLWD, we have previously shown that continuous expression LWD1 does not alter circadian rhythms of pCCA1 activity[27]. Therefore, the arrhythmic phenotype of *ttg1 lwd1 lwd2* mutants is a composite of sequential loss of transcriptional repression and activation activities of the binding partners of TLWD proteins throughout the diel cycle.

Novel function of TLWD proteins in circadian oscillator transcriptional repression

Analysing circadian gene expression under diel cycles allowed us to compare circadian arrhythmic mutants and identify specific activation and repression defects. We found that in *ttg1 lwd1 lwd2* mutants lack of repression correlated with Myb-like transcription factor repression during the day and with PRR-mediated repression at night (Figure 1,4 and Extended data Fig. 6). Our data are consistent with the Evening Complex not being compromised by the absence of *tlwd* function because *elf3* mutants have different diel oscillations of both *pCCA1* and *pTOC1* to *ttg1 lwd1 lwd2* mutants (Figure 4 and Extended data Fig. 6).

Until now, the activity of TTG1 and LWDs has been associated with transcriptional activation, through MBW complexes in the case of TTG1 and through LWD-TCP complexes[17,18,28,41]. However, our data indicate that it is the lack of repression that causes arrhythmicity in the absence of TLWD function. Blue light dependent LWD1-TCP-CRY2 association with the *CCA1* promoter at dawn mediates *CCA1* transcriptional activation [17,18]. Notably, in the *tcp20/tcp22* double mutant, LWD1 is still associated with the *CCA1* promoter, indicating that additional transcription factors mediate its binding. Our data support a direct role of TLWD-PRR complexes in repressing *CCA1,* which is most apparent in the second half of the diel cycle. High order *prr* mutants phenocopy the aberrant expression of *pCCA1::LUC* in *ttg1 lwd1 lwd2* mutants (Figure 1D and 4G), including the second *pCCA1::LUC* peak under long day photoperiod. [17,18]

We identified a function of TLWD in controlling hypocotyl growth by repressing *PIF4* and *PIF5*
expression, possibly via TLWD-PRR complexes (Figure 4F and 5, Extended data Fig.7). Notably,
ttg1 lwd1 lwd2 mutants have lower levels of *PIF4* expression and shorter hypocotyls when compared
to *prr5 prr7 toc1* mutants (Figure 4). Since CCA1 has been shown to promote the expression of
PIF4, it is possible that lower *CCA1* levels in *ttg1 lwd1 lwd2* mutants cause lower *PIF4* expression
and hypocotyl elongation. Another scenario is that TLWD complexes control hypocotyl elongation
only at the level of *PIF4/PIF5* expression levels via TLWD-PRR complexes but they do not participate
in controlling PIF4 and PIF5 protein activity, as shown for PRRs and ELF3[36,42].

CCA1/LHY directly repress the expression of *PRR* genes[43]. We found that *lhy cca1* and *ttg1*
lwd1 lwd2 have a similar de-repression of *pTOC1* rhythms during the day (Figure 1, Extended data
Fig. 6). Notably, Wang *et al.* previously found that LWD1 associates with the *TOC1* promoter at ZT0
and at ZT6, when *TOC1* is repressed, but not at ZT12, when *TOC1* expression is highest [26]. This
supports a direct role of TLWD proteins in the repression of *TOC1* via TLWD-CCA1 complexes. In
addition, LWD1 binds to the *PRR9* promoter at ZT0, 6 and 12, which coincides with both high and
low levels of *PRR9* expression[26]. We found that *PRR9* was downregulated at ZT6 in both *lwd1 lwd2*
and *ttg1 lwd1 lwd2* mutants (Figure 1B, Extended data Fig. 1), further supporting a continuous
function of TLWD proteins which involves both activation and repression of circadian regulated
genes.

TLWD complexes as a model for dynamic transcriptional regulation

How TLWD proteins contribute to the function of circadian oscillator transcription factors is
still poorly understood. We propose that TLWD scaffolds might provide a continuous platform that
connects the activity of different transcription factor families that bind sequentially to the same
circadian-regulated promoters such as *CCA1* and *TOC1* (Figure 5). TPC20 protein binds to the
CCA1 promoter in the *lwd1 lwd2* mutant and TT8 and TT2 can bind to target promoters in the *ttg1*
mutant[17,41], indicating that TLWD is not required for TF binding to DNA. However, in the absence of
TLWD function these transcription factors cannot activate target gene expression. Similarly, we
found a common defect of transcriptional repression of PRR-direct targets such as *CCA1*, *PIF4* and
PIF5 in higher order *prr* and *tlwd* mutants, indicating that TLWD is required for PRR-mediated
repression. The essential function of TLWD might be due to simultaneous recruitment of additional
transcriptional regulators and regulatory proteins. For example, it was reported that LWD1 forms
complexes with YET ANOTHER KINASE (YAK1) and that *yak1* mutants have circadian
phenotypes[44]. More recently, LWD1 was implicated in the phase separation of CRY2-TCP22-PKK
complexes. Understanding the composition and regulated assembly and disassembly of TLWD-
clock complexes throughout the diel cycle might uncover novel targets to alter crop yield components
such as growth and flowering time. More broadly, TLWD can be used as a model of the interplay
between repression and activation required for dynamic transcriptional regulation.

A new dynamic in circadian regulation

Rhythm generation in circadian biology has in recent times been considered analogous to a "clock"[45]. Whilst clock analogies have their limitations[46], they have been useful in understanding the mechanistic basis of circadian timing. The circadian clocks are thought to comprise oscillators of transcriptional translational feedback loops (TTL) and separate or coupled feedback loops of metabolism and signalling[46,47]. The Arabidopsis TTL comprises the sequential expression of CCA1/LHY-PRRs-EC transcriptional repressors from dawn to end of night, and TCP and RVE8 transcriptional activators active at dawn and at midday, respectively[17,48,49]. In the clock model the sequential expression and repression of the transcriptional activators are considered analogous to the wheels or cogs of a mechanical clock[46,50]. Essential to the sequential activities of the transcriptional repressors and activators are rhythmic changes in the abundance of the encoding transcripts and the proteins[13,33,48,51]. Here we find that TLWDs contribute to circadian timing by forming complexes with both transcriptional repressors and activators. TLWD proteins are present throughout the circadian cycle and do not appear to undergo large oscillatory changes in abundance, allowing for interactions with their binding partners as those change in abundance through the cycle (Figure 5). Thus, TLWD proteins represent a new category of circadian oscillator component, that are continuously present and essential for the function of the moving parts. In the clock analogy, the TLWD proteins act as arbors or spindles, being attached to the gears or wheels of the clock formed by the transcriptional repressors/activators. The arbor TLWD proteins provide the pivotal centre of the clock, attached to which the wheels of the clock can turn.

Author contributions

A.A.R.W., E.H. and B.J.G. conceived the project and designed the experiments. EH conducted all the experiments. D.C.R. contributed to RNAseq analysis. E.H., B.J.G. and A.A.R.W. wrote the manuscript. All authors commented on the manuscript before submission.

Acknowledgments

We thank M. Stancombe and M. Dorling for laboratory and plant growth support. We thank Prof. David Somers and Dr. Tina Schneider for providing guidance on co-immunoprecipitation experiments. E.H. is supported by a Broodbank Fellowship from the Department of Plant Sciences at the University of Cambridge awarded to E.H., and a UK Space Agency International Bilateral Fund Pathfinder Phase 2 grant 10089325 awarded to A.A.R.W. Research performed by E.H. was also supported by a Leverhulme Trust Award RPG-2018-318 awarded to A.A.R.W and B.J.G. D.C.R. was supported by UKRI BBSRC grant BB/S002251/1 awarded to A.A.R.W.

Competing interests

The authors declare no competing interests

Data availability

All data that support the findings of this study are available at the Apollo repository (https://doi.org/10.17863/CAM.115480). RNAseq Raw data (FASTQ) files has been deposited under BioProject PRJNA1217148.

Methods

Plant materials and growth conditions

See Supplementary Table 1 for details of the *A. thaliana* lines used in this study. Seedlings were surface sterilized by sequential immersion in 100% ethanol, 33% bleach with 0.01% (v/v) Triton X-100 and ultra-pure sterile water. Seeds were plated on Murashige and Skoog (MS) media (Duchefa) supplemented with 0.8% of bactoagar. After 3 days of stratification at 4 °C, they were placed in a controlled cabinet with constant 20 °C and different photoperiods of white light (130 μmol m^{-2} s^{-1}).

Generation of transgenic lines

Col-0 ttg1 lwd1 lwd2 pCCA1::LUC were already described[27]. *ttg1 lwd1 lwd2* plants were crossed with either Col-O *pPRR7::LUC*[52] or *pTOC1::LUC*[53]. F1 heterozygous plants were backcrossed with either *ttg1 lwd1 lwd2* to obtain triple mutants with the relevant LUC reporter in the next generation. Mutants were confirmed by genotyping using previously described primers[27]. In the next generation, homozygous LUC reporter was confirmed by lack of segregation for resistance to phosphinothricin (PPT) for *pPRR7::LUC*, or hygromycin (HYG) for *pTOC1::LUC* when seedlings were grown on agar supplements with the selection media.

For LWD1-Venus and Venus-TTG1 complementation lines, LWD1 promoter (1068 bp) or TTG1 promoter (1328bp), LWD1 or TTG1 coding sequence, mVenus, downstream sequences (dss) of LWD1 (713 bp) or TTG1 (1000bp) were cloned in the pGREEN-BASTA binary vector by Gibson assembly to obtain pLWD1::LWD1-Venus::LWD1dss and pTTG1::Venus-TTG1::TTG1dss. For LWD1-LUC+ and TTG1-LUC+ complementation lines, the plasmids pCsA-BASTA-pLWD1::LWD1-LUC::LWD1dss or pCsA-BASTA-pTTG1::TTG1-LUC+::TTG1-dss were obtained by loop assembly[54]. A flexible linker was inserted between the coding sequence of LWD1 and TTG1 and Venus or LUC+. The corresponding plasmids were transformed by floral dipping into *lwd1 lwd2 pCCA1::LUC* or *ttg1-21 pCCA1::LUC*. T0 seeds were selected on MS PPT plates. In T1, at least two independent lines were selected on their ability to complement mutant phenotypes: circadian periodicity of *pCCA1::LUC* and flowering time for *lwd1 lwd2*, and seed coat pigmentation and trichomes for *ttg1-21*. Then T2 homozygous lines were obtained from selected lines and the presence of the fusion protein was confirmed by western blots in either T2 or T3 seedlings.

Measuring diel and circadian rhythms of LUC reporter lines

Luminescence assays were performed as described in [27]. Clusters of 5 plants with the indicated LUC reporters were grown for 10 days under entrainment photoperiods before luciferin

solution was added to the media and plants were placed in the imaging system (either a Photek
ICCD25 or a Berthold camera system). Every hour, bioluminescence signal was integrated for 800
seconds. Similar light intensity entrainment conditions were maintained for 2-3 days, followed by 3-
5 days of continuous light. Where indicated, traces were normalized to the average bioluminescence
through the experiment.

RNAseq

Seeds were grown for 10 days in ½MS plates supplemented with 0.8% agar under diel cycles
of 12 light / 12 dark. Seedlings were harvested at time ZT6 and time ZT18 and immediately frozen
in liquid nitrogen. Each technical replicate consisted of three clusters of 10 seedlings growing on
different agar plates in the same controlled cabinet. Plant material was ground using a Tissue Lyser
(QIAGEN). Total RNA was extracted with the RNeasy Plant Mini Kit (QIAGEN). After extraction, the
quality of the RNA was determined with a Bioanalyzer. Three biological replicates were obtained for
each time point, by pooling the RNA of three technical replicates. Library preparation and pair-end
sequencing was performed with the Illumina RNA-seq platform with 15-20 million 150bp-reads by
Genewiz. Sequencing quality was assessed by generating FastaQC and MultiQC reports.
Trimmomatics was used to remove adapters and non-Arabidopsis sequences. Sequences were
mapped to the Arabidopsis genome (TAIR10) and counted by RNAstar and Feature Counts,
respectively. Differential gene expression between Col-0 and *ttg1*, *lwd1 lwd2* and *ttg1 lwd1 lwd2*
mutants was calculated with DESeq2. Genes with a fold change Log2 $\geq$ 1.5 with respect to wildtype
were considered DEGs.

qPCR

Seedlings were grown for 10 days under short day photoperiods (8h light /16h dark). For each time
point, three biological replicates (each formed by three pooled clusters of approximately 20
seedlings) were collected, frozen in liquid nitrogen and ground with a Tissue Lyser (QIAGEN). Total
RNA was extracted using the RNeasy Plant Mini Kit (QIAGEN) and RNase-Free DNase on-column
treatment (QIAGEN). cDNA was synthesized from 500 ng RNA with the RevertAid First Strand cDNA
Synthesis Kit (Thermo Scientific) using oligo (dT) primers. The gene-specific products were amplified
using the Quantinova SYBR Green PCR Kit (QIAGEN) on a Biorad CFX384 Real-Time System.
Primers used are detailed in Supplementary Table 2. Relative transcript levels were normalized to
both *PP2A* and *UBQ10* expression.

Yeast 2-hybrid experiments

All coding sequences were amplified from Col-O cDNA and cloned into pDONR201 by
Gateway BP reaction. Coding sequences were cloned into pAS2 and pC-ACT2 (Clontech) vectors

by Gateway LR reaction. Plasmids were verified by Sanger sequencing. pAS2 and pC-ACT2 vectors were introduced into yeast strains AH109 and Y187, respectively, by Li-Ac mediated transformation following Clontech Yeast Protocols Handbook (Clontech). Positive colonies were re-streaked in the appropriate synthetic dropout (SD) media lacking leucine (-L) or tryptophane (-W), respectively. Appropriate pAS+pACT2 diploid strains were obtained by mating and selecting in SD-LW media. Protein interactions were evaluated by the ability of diploid strains to grow on SD media without Histidine and Adenine. Serial dilutions of diploid strains were plated in SD -LW, SD-LHW supplemented with 3-amino-1,2,4-triazole (3AT) and SD-LHWA and grown for 3-4 days before imaging. Y2H results were confirmed in at least 2 independent experiments.

Arabidopsis whole protein extracts and western blot

Approximately 40 15-day old seedlings were collected and frozen in nitrogen at specific times of the diel cycle. Plant materials were finely ground in a Tissue lyser and resuspended in fresh extraction buffer (50 mM Tis-HCl pH 7.5, 150 mM Na CL, 0.5% NP-40, 1mM EDTA, 2mM Na_3VO_4, 10 mM NaF, 2mM β-glycerol phosphate, 1mM PMSF, 50 µM MG132, 50 µM MG115, 50 µM ALLN, 3mM DTT and Protein inhibitor tablet). Protein extracts were cleared by centrifugation at max speed at 4°C. Protein concentration was measured with Pierce™ 660nm Protein Assay Reagent (Thermo Scientific) following manufacturer's instructions. 6X SDS buffer was added to protein extracts. 10 µg of each sample was loaded into precast gels (SurePAGE Bis-Tris, GenScript). Proteins were transferred onto supported nitrocellulose membrane (Biorad) by wet transfer. Membrane was blocked with Western blocking reagent (Roche) and incubated with α-GFP (Roche, 1181446000) and α-mouse-HRP (Proteintech SA00001-1). Membranes were covered with ECL SuperBright (Agrisera) for 2 minutes and briefly rinsed in TBS before imaging with a G:BOX Chemi XRQ imaging system.

Co-Immunoprecipitation

The coding sequences of LWD1 and TTG1 were cloned into pCK1_35S::mTurq-LWD1 and pCK1_p35S::mTurq-TTG1 by Loop assembly cloning[55]. The coding sequences of PRR9, PRR7, PRR5 and TOC1 were cloned into pJCV52[56] to obtain pJCV52_p35S::PRR-HA by Gateway cloning. Plasmids were introduced into Agrobacterium strain AGL1. Appropriate Agrobacterium cultures were diluted and mixed with p19 plasmid to OD_{600}=0.5 in infiltration solution (1/4MS pH 6, 1% sucrose, 100µM acetosyringone and 50 µl/l Silwet L-77) and left for 3-4 hours in the dark before infiltrating leaves of 3-4 week old *Nicotiana benthamiana* plants with a 1mL syringe. Plants were kept in the light until leaves were dried, then covered with a black bag for 24 hours. Infiltrated leaves were collected 48-72 hours after infiltration and immediately frozen in liquid nitrogen. Plant material was finely ground in a Tissue lyser and resuspended in fresh extraction buffer [57] (50 mM Tis-HCl pH 7.5, 150 mM Na CL, 0.5% NP-40, 1mM EDTA, 2mM Na_3VO_4, 10 mM NaF, 2mM β-glycerol phosphate,

1mM PMSF, 50 µM MG132, 50 µM MG115, 50 µM ALLN, 3mM DTT and Protein inhibitor tablet).
Protein extracts were cleared by centrifugation at max speed at 4°C then incubated with µMACS™
HA beads (Miltenyi Biotec) for 2 hours with rotation at 4°C. Columns (Miltenyi Biotec) were used to
wash the beads with extraction buffer following the manufacturer's protocol. To elute protein
complexes, HA beads were incubated for 15 minutes with HA peptide solution at 30°C. Then, 6X
sample buffer was added to the eluate and incubated for 1 min at 90°C before loading into precast
gels (SurePAGE Bis-Tris, GenScript). Electrophoresis was performed at 4°C and low voltage (60-
90V). Proteins were transferred onto supported nitrocellulose membrane (Biorad) by wet transfer at
4°C. Membrane was blocked with Western blocking reagent (Roche) and incubated with primary
antibodies α-HA (Abcam, ab9110) and α-GFP (Roche, 1181446000) and secondary antibodies α-
rabbit-HRP (Proteintech SA00001-2) and α-mouse-HRP (Proteintech SA00001-1), respectively.
Membranes were covered with ECL SuperBright (Agrisera) for 2 minutes and briefly rinsed in TBS
before imaging with a G:BOX Chemi XRQ imaging system.
502
503 *Hypocotyl elongation*
504
505   Seeds were plated on MS media (4.4 g/L with 0.5 g/L MES and 1.5% phytoagar pH 5.7).
506 After stratification plants were placed in a growth cabinet with 100 µmol m^{-2} s^{-1} of white light under
507 short day (8h_L/ 16D) or long day (16h_L/ 8h_D) photoperiods. Hypocotyl length was measured
508 after 7 days using Image J software as previously described[58].
509
510 **Figure Legends**
511
Figure 1. TLWD proteins regulate circadian rhythms of gene expression. (A) Percentage of
genes upregulated (red) or downregulated (blue) that are circadian rhythmic (CRG) in *lwd1 lwd2*
(green) and *ttg1 lwd1 lwd2* (pink) mutants during day (ZT6) or night (ZT18) and distribution of their
usual circadian phase of expression in wild type. RP: representation factor. Asterisks depict
statistically significant enrichment of the circadian phase. Figure data in Extended Data Table 1. (B)
Circadian oscillator components upregulated (red) or downregulated (blue) relative to wild type in
lwd1 lwd2 (green) and *ttg1 lwd1 lwd2* (pink) mutants during day (ZT6) or night (ZT18). Genes are
depicted according to their usual temporally restricted expression pattern through the diel cycle.
Genes with no change in expression are depicted in grey. See Extended data Fig. S1C for
normalized expression levels. (C-D) Oscillations of *pCCA1::LUC* in wild type and *ttg1* mutants (C)
and *lwd1 lwd2* and *ttg1 lwd1 lwd2* mutants (D). Y axis depicts mean bioluminescence (counts in 800
sec) of cluster of seedlings, n=24 per genotype. Note the different maximum Y-axis values. (E-F)
Oscillations of *pTOC1::LUC* (E) and *pPRR7::LUC* (F) in wild type and *ttg1 lwd1 lwd2* mutants. Y axis
depicts mean bioluminescence (counts in 800 sec), of *pTOC1::LUC* (wild type n=8, *ttg1 lwd1 lwd2*
n=8) and *pPRR7::LUC* (wild type n=8, *ttg1 lwd1 lwd2* n=12) . Error bars in C-F depict SEM.
527

Figure 2. TLWD proteins are present throughout the diel cycle. Representative western blots from protein extracts of (A) *pCCA1::CCA1-YFP*, and two independent (B) *lwd1 lwd2 pCCA1::LUC pLWD1::LWD1-YFP* and (C) *ttg1 pCCA1::LUC pTTG1::YFP-TTG1*. Ponceau staining indicates similar protein extract loading between time points. (D) and (E) Rhythms of *pLWD1::LWD1-LUC* and *pTTG1::TTG1-LUC* under long days and continuous light. Error bars depicts SEM. *lwd1 lwd2 pLWD1::LWD1-LUC* line 1 and 2 , n=19. *ttg1 pTTG1::TTG1-LUC* line 1 and 2 , n=20.

Figure 3. TLWD proteins form complexes with PRRs. (A) Y2H protein interaction assay: 3 serial dilutions (1:10) of two representative colonies. Growth in LHW media indicates interaction between protein pair. (B) and (C) Co-immunoprecipitation of LWD1 (B) and TTG1 (C) with PRR proteins from *N. benthamiana* extracts. (D) Venn diagram depicts overlap of CRGs upregulated relative to wildtype in *lwd1 lwd2* (ZT18), *ttg1 lwd1 lwd2* (ZT18), and *prr5 toc1 (ZT15)*[32]. *(E)* Venn diagram depicts overlap between the 49 CRGs upregulated in both *tlwd* and *prr* mutants and the direct targets of TOC1 and PRR5 as established by ChIPseq[12, 33]. (F) Fold change of expression of potential TLWD-PRR complex targets in corresponding transcriptome analysis. (G-H) Comparison of *pCCA1::LUC* oscillations in (G) *tlwd* and *prr* mutants or (H) *tlwd* and *elf3* mutants. Y axis depicts mean normalized bioluminescence of clusters of seedlings. Error bars depict SEM, genotype (n): in (G) wild type (15), *lwd1 lwd2* (12), *prr5 prr7* (16), *prr5 prr7 toc1* (16), *ttg1 lwd1 lwd2* (16) and in (H) wild type (20), *lwd1 lwd2* (11), *ttg1 lwd1 lwd2* (23), *elf3* (20).

Figure 4. TLWD control growth by repressing *PIF4* and *PIF5* expression at night. (A) Normalized expression of *PIF4*, *PIF5* and their gene targets in wildtype, *lwd1 lwd2*, *ttg1* and *ttg1 lwd1 lwd2* mutants as measured by RNAseq. Normalized counts were obtained with Deseq2 (Extended data Table 1). Error bars depict SEM, n=3. (B) and (C) Representative seedlings and mean hypocotyl length of plants grown for 7 days under long day (16L/8D) and short day (8L/16D) photoperiods of white light. Error bars depict SEM; Long day, n= wild type (64), *lwd1 lwd2* (46), *ttg1* (71), *ttg1 lwd1 lwd2* (77) and *prr5 prr7 toc1* (48); Short day, n=wild type (54), *lwd1 lwd2* (47), *ttg1* (55), *ttg1 lwd1 lwd2* (46) and *prr5 prr7 toc1* (58). Letters depict genotypes with statistically different average length with the Tukey Kramer test (D) Expression profile of *PIF4, PIF5* and *CDF5* under short day photoperiods. Expression values are normalized to both *PP2A* and *UBQ10* expression and plotted relative to average wild type expression through the diel cycle. Error bars depict SEM, n=3

Figure 5. TLWD are arbor proteins in the Arabidopsis circadian oscillator. TLWD provides a continuous platform for the sequential formation of activator and repressor complexes in the Arabidopsis circadian oscillator. The model depicts sequential temporal complex formation between TLWD (central arbor) and clock transcription factors (outer wheels). TLWD structure was generated by Alphafold and illustrated with ChimeraX. Interaction surfaces between TLWD and TF are not based on structural information. Outer bars depict timing of activity of TLWD interacting clock transcription factors (red=activators, blue=repressors, pink=TLWD).

Extended Data Figure Legends

Extended data Figure 1. Transcript abundance of MBW complex targets and circadian oscillator components in *tlwd* mutants. (A) Normalized expression of MBW complex direct targets[29] in wild type, *lwd1 lwd2*, *ttg1* and *ttg1 lwd1 lwd2* mutants. (B) Hierarchical clustering of DEGs in *ttg1*, *lwd1 lwd2* and *ttg1 lwd1 lwd2* at Day (ZT6) and Night (ZT18) when compared to wild type expression. (C) Normalized expression of selected circadian oscillator components in wild type, *lwd1 lwd2*, *ttg1* and *ttg1 lwd1 lwd2* mutants. Normalized counts in A and C were obtained with Deseq2. Error bars depict SEM, n=3 See Extended data Table S1 for values.

Extended data Figure 2. Rhythms of *pCCA1:LUC* under different photoperiods in wildtype, *lwd1 lwd2* and *ttg1 lwd1 lwd2* mutants. Note that under long days (left panel) the *ttg1 lwd1 lwd2* mutant has a second peak of *CCA1* in the second half of the diel cycle which is not apparent under neutral days or short days (middle and right panels). Error bars depict SEM, n= (LD) wild type (8), *lwd1 lwd2* (16) and *ttg1 lwd1 lwd2* (12); (ND) wild type (12), *lwd1 lwd2* (12) and *ttg1 lwd1 lwd2* (12); (SD) wild type (24), *lwd1 lwd2* (22) and *ttg1 lwd1 lwd2* (28).

Extended data Figure 3. *LWD1-Venus* complementation lines. (A) Complementation of *lwd1 lwd2* mutant short periodicity phenotype in two independent *pLWD1::LWD1-Venus::LWD1dss* lines. Y-axis depict average normalized bioluminescence, error bars depict SEM, n= wild type (12), *lwd1 lwd2* (12), *lwd1 lwd2 pLWD1::LWD1-Venus::LWD1* line 1(8) and line 2 (8). Confocal microscopy images show nuclear localization of LWD1-Venus in both complementation lines.

Extended data Figure 4. TTG1 complementation lines. (A) Oscillations of *pCCA1::LUC*. Complementation of *ttg1* mutant with *pTTG1::Venus-TTG1::TTG1dss* does not affect circadian rhythms. Note that *ttg1* does not have a circadian phenotype. Y-axis depict average bioluminescence of 8 seedling clusters per genotype, error bars depict SEM. (B) confocal microscopy images show nuclear localization of Venus-TTG1-Venus in complementation lines both in stem and trichomes. (C) Complementation of *ttg1-21* mutant lack of seed coat pigmentation and trichomes in *Venus-TTG1* and *TTG1-LUC* complementation lines.

Extended data Figure 5. Interaction of TLWD proteins with TCPs. All three TLWD proteins interact with TCP20, TCP22 and CHE (TCP21). 3 serial dilutions (1:10) of four independent colonies. Growth in LHW media indicates interaction between protein pairs.

Extended data Figure 6. TLWD proteins interact with Myb-like transcription factors and control *TOC1* expression. (A) TTG1 interacts with CCA1, LHY and RVE1 but not with RVE8. 3 serial dilutions (1:10) of two independent colonies. Growth in LHW media with 1mM 3AT indicates

interaction between protein pair. (B) Oscillations of *pTOC1::LUC* in wild type, *ttg1 lwd1 lwd2, cca1 lhy and elf3* under diel cycles and continuous light. Note that *ttg1 lwd1 lwd2* and *elf3* are in Col-0 background whereas *lhy cca1* is in the Ws background. Col-0 and Ws wild types have similar rhythms. Y-axis depicts average normalized bioluminesce of 8 cluster of seedlings per genotype.

Extended data Figure 7. *ttg1 lwd1 lwd2* mutants recapitulate some *PIF4-OX* phenotypes. (A) Normalized expression of PIF4/5 targets in wild type and *ttg1 lwd1 lwd2* mutants. Error bars depict SEM, n=3. Normalized counts were obtained with Deseq2. See Extended data Table 1 for values. B) Hypocotyl growth of plants grown for 7 days under short day photoperiods. Error bars depict SEM, n= wild type (88), *ttg1* (91), *lwd1 lwd2* (73), *ttg1 lwd1 lwd2* (116), *prr5 prr7 prr9* (82) and *elf3* (103). Letters depict genotypes with statistically different average length with the Tukey Kramer test. (D) Representative rosettes of wild type, *ttg1*, *lwd1 lwd2*, *ttg1 lwd1 lwd2*, *elf3* and *PIF4-OX*[59] mutants grown under short day photoperiods.

Extended data Table 1. RNAseq differentially expressed genes Log2^{FC}, normalized counts and circadian phase of expression.

Supplementary Table 1. Arabidopsis thaliana lines used in this study
Supplementary Table 2. qPCR primers

References

1.   Sanchez, S. E. & Kay, S. A. The plant circadian clock: From a simple timekeeper to a complex developmental manager. *Cold Spring Harb Perspect Biol* **8**, (2016).

2.   Dodd, A. N. *et al.* Plant Circadian Clocks Increase Photosynthesis, Growth, Survival, and Competitive Advantage. *Science (1979)* **309**, 630–633 (2005).

3.   Steed, G., Ramirez, D. C., Hannah, M. A. & Webb, A. A. R. Chronoculture, harnessing the circadian clock to improve crop yield and sustainability. *Science* **372**, (2021).

4.   Romanowski, A., Schlaen, R. G., Perez-Santangelo, S., Mancini, E. & Yanovsky, M. J. Global transcriptome analysis reveals circadian control of splicing events in Arabidopsis thaliana. *Plant Journal* **103**, (2020).

5.   Nagel, D. H. *et al.* Genome-wide identification of CCA1 targets uncovers an expanded clock network in Arabidopsis. *Proceedings of the National Academy of Sciences* **112**, E4802 – E4810 (2015).

6.   Adams, S. *et al.* Circadian control of abscisic acid biosynthesis and signalling pathways revealed by genome-wide analysis of LHY binding targets. *New Phytologist* **220**, 893–907 (2018).

7.   Adams, S., Manfield, I., Stockley, P. & Carré, I. A. Revised Morning Loops of the Arabidopsis Circadian Clock Based on Analyses of Direct Regulatory Interactions. *PLoS One* **10**, (2015).

8.   Hsu, P. Y., Devisetty, U. K. & Harmer, S. L. Accurate timekeeping is controlled by a cycling activator in Arabidopsis. *Elife* **2**, e00473 (2013).

9. Liu, T. L., Newton, L., Liu, M.-J., Shiu, S.-H. & Farré, E. M. A G-Box-Like Motif Is
Necessary for Transcriptional Regulation by Circadian Pseudo-Response Regulators in
Arabidopsis. *Plant Physiol* **170**, 528–539 (2016).
10. Liu, T., Carlsson, J., Takeuchi, T., Newton, L. & Farré, E. M. Direct regulation of abiotic
responses by the Arabidopsis circadian clock component PRR7. *Plant J* **76**, 101–114
(2013).
11. Nakamichi, N. *et al.* Transcriptional repressor PRR5 directly regulates clock-output
pathways. *Proc Natl Acad Sci U S A* **109**, 17123–17128 (2012).
12. W., H. *et al.* Mapping the Core of the Arabidopsis Circadian Clock Defines the
Network Structure of the Oscillator. *Science (1979)* **336**, 75–79 (2012).
13. Nakamichi, N. *et al.* PSEUDO-RESPONSE REGULATORS 9, 7, and 5 Are Transcriptional
Repressors in the Arabidopsis Circadian Clock . *Plant Cell* **22**, 594–605 (2010).
14. Ezer, D. *et al.* The evening complex coordinates environmental and endogenous
signals in Arabidopsis. *Nat Plants* **3**, 17087 (2017).
15. Helfer, A. *et al.* LUX ARRHYTHMO encodes a nighttime repressor of circadian gene
expression in the Arabidopsis core clock. *Current Biology* **21**, (2011).
16. Nusinow, D. A. *et al.* The ELF4-ELF3-"LUX complex links the circadian clock to diurnal
control of hypocotyl growth. *Nature* **475**, (2011).
17. Wu, J.-F. *et al.* LWD–TCP complex activates the morning gene CCA1 in Arabidopsis.
Nat Commun **7**, 13181 (2016).
18. Mo, W. *et al.* Arabidopsis cryptochrome 2 forms photobodies with TCP22 under blue
light and regulates the circadian clock. *Nat Commun* **13**, (2022).
19. Pruneda-Paz, J. L., Breton, G., Para, A. & Kay, S. A. A functional genomics approach
reveals CHE as a component of the Arabidopsis circadian clock. *Science (1979)* **323**,
(2009).
20. Herrero, E. *et al.* EARLY FLOWERING4 Recruitment of EARLY FLOWERING3 in the
Nucleus Sustains the Arabidopsis Circadian Clock. *The Plant Cell Online* **24**, 428–443
(2012).
21. Cha, J. Y. *et al.* GIGANTEA is a co-chaperone which facilitates maturation of ZEITLUPE
in the Arabidopsis circadian clock. *Nat Commun* **8**, (2017).
22. Fujiwara, S. *et al.* Post-translational regulation of the Arabidopsis circadian clock
through selective proteolysis and phosphorylation of pseudo-response regulator
proteins. *Journal of Biological Chemistry* **283**, 23073–23083 (2008).
23. Somers, D. E., Fujiwara, S., Kim, W. Y. & Suh, S. S. Posttranslational photomodulation
of circadian amplitude. in *Cold Spring Harbor Symposia on Quantitative Biology* vol.
72 (2007).
24. Nohales, M. A. *et al.* Multi-level Modulation of Light Signaling by GIGANTEA Regulates
Both the Output and Pace of the Circadian Clock. *Dev Cell* **49**, (2019).
25. Dalchau, N. *et al.* The circadian oscillator gene GIGANTEA mediates a long-term
response of the Arabidopsis thaliana circadian clock to sucrose. *Proc Natl Acad Sci U S*
A **108**, 5104–5109 (2011).
26. Wang, Y. *et al.* LIGHT-REGULATED WD1 and PSEUDO-RESPONSE REGULATOR9 form a
positive feedback regulatory loop in the Arabidopsis circadian clock. *Plant Cell* **23**,
486–498 (2011).
27. Airoldi, C. A., Hearn, T. J., Brockington, S. F., Webb, A. A. R. & Glover, B. J. TTG1
proteins regulate circadian activity as well as epidermal cell fate and pigmentation.
Nat Plants **5**, 1145–1153 (2019).
28. Ramsay, N. A. & Glover, B. J. MYB-bHLH-WD40 protein complex and the evolution of
cellular diversity. *Trends Plant Sci* **10**, 63–70 (2005).

29.   Xu, W., Dubos, C. & Lepiniec, L. Transcriptional control of flavonoid biosynthesis by
698   MYB-bHLH-WDR complexes. *Trends Plant Sci* **20**, 176–185 (2015).
699   30.   Bonnot, T., Gillard, M. B. & Nagel, D. H. CAST-R: An application to visualize circadian
700   and heat stress-responsive genes in plants. *Plant Physiol* **190**, (2022).
701   31.   Gould, P. D. *et al.* Coordination of robust single cell rhythms in the Arabidopsis
702   circadian clock via spatial waves of gene expression. *Elife* **7**, (2018).
703   32.   Yamashino, T. *et al.* Involvement of Arabidopsis clock-associated pseudo-response
704   regulators in diurnal oscillations of gene expression in the presence of environmental
705   time cues. *Plant Cell Physiol* **49**, (2008).
706   33.   Li, N., Zhang, Y., He, Y., Wang, Y. & Wang, L. Pseudo Response Regulators Regulate
707   Photoperiodic Hypocotyl Growth by Repressing PIF4/5 Transcription. *Plant Physiol*
708   **183**, 686–699 (2020).
709   34.   Nakamichi, N. *et al.* Transcriptional repressor PRR5 directly regulates clock-output
710   pathways. *Proc Natl Acad Sci U S A* **109**, 17123–17128 (2012).
711   35.   Kolmos, E. *et al.* A Reduced-Function Allele Reveals That EARLY FLOWERING3
712   Repressive Action on the Circadian Clock Is Modulated by Phytochrome Signals in
713   Arabidopsis. *The Plant Cell Online* **23**, 3230–3246 (2011).
714   36.   Martín, G. *et al.* Circadian Waves of Transcriptional Repression Shape PIF-Regulated
715   Photoperiod-Responsive Growth in Arabidopsis. *Current Biology* **28**, 311–318.e5
716   (2018).
717   37.   Yu, J. W. *et al.* COP1 and ELF3 Control Circadian Function and Photoperiodic
718   Flowering by Regulating GI Stability. *Mol Cell* **32**, (2008).
719   38.   Kim, W. Y. *et al.* ZEITLUPE is a circadian photoreceptor stabilized by GIGANTEA in blue
720   light. *Nature* **449**, (2007).
721   39.   Park, Y. J. *et al.* GIGANTEA Shapes the Photoperiodic Rhythms of
722   Thermomorphogenic Growth in Arabidopsis. *Mol Plant* **13**, (2020).
723   40.   Más, P., Kim, W. Y., Somers, D. E. & Kay, S. A. Targeted degradation of TOC1 by ZTL
724   modulates circadian function in Arabidopsis thaliana. *Nature* **426**, (2003).
725   41.   Baudry, A. *et al.* TT2, TT8, and TTG1 synergistically specify the expression of BANYULS
726   and proanthocyanidin biosynthesis in Arabidopsis thaliana. *The Plant Journal* **39**,
727   366–380 (2004).
728   42.   Nieto, C., López-Salmerón, V., Davière, J. M. & Prat, S. ELF3-PIF4 interaction regulates
729   plant growth independently of the evening complex. *Current Biology* **25**, (2015).
730   43.   Kamioka, M. *et al.* Direct repression of evening genes by CIRCADIAN CLOCK-
731   ASSOCIATED1 in the Arabidopsis circadian clock. *Plant Cell* **28**, 696–711 (2016).
732   44.   Huang, W. Y. *et al.* Plant dual-specificity tyrosine phosphorylation-regulated kinase
733   optimizes light-regulated growth and development in Arabidopsis. *Plant Cell Environ*
734   **40**, (2017).
735   45.   Aviram, R. & Manella, G. A Metaphor That Keeps on Ticking: The 'Clock' as a Driving
736   Force in the History of Chronobiology Research. *Philosophy, Theory, and Practice in*
737   *Biology* **12**, (2020).
738   46.   Webb, A. A. R., Seki, M., Satake, A. & Caldana, C. Continuous dynamic adjustment of
739   the plant circadian oscillator. *Nat Commun* **10**, 550 (2019).
740   47.   Stangherlin, A., Seinkmane, E. & O'Neill, J. S. Understanding circadian regulation of
741   mammalian cell function, protein homeostasis, and metabolism. *Current Opinion in*
742   *Systems Biology* vol. 28
743   48.   Rawat, R. *et al.* REVEILLE8 and PSEUDO-REPONSE REGULATOR5 form a negative
744   feedback loop within the arabidopsis circadian clock. *PLoS Genet* **7**, (2011).

745   49.   Xie, Q. *et al.* LNK1 and LNK2 are transcriptional coactivators in the Arabidopsis
746   circadian oscillator. *Plant Cell* **26**, 2843–2857 (2014).

747   50.   Hsu, P. Y. & Harmer, S. L. Wheels within wheels: The plant circadian system. *Trends in*
748   *Plant Science* vol. 19 (2014).

749   51.   Wang, Z. Y. & Tobin, E. M. Constitutive expression of the CIRCADIAN CLOCK
750   ASSOCIATED 1 (CCA1) gene disrupts circadian rhythms and suppresses its own
751   expression. *Cell* **93**, 1207–1217 (1998).

752   52.   de Melo, J. R. F. *et al.* Magnesium maintains the length of the circadian period in
753   arabidopsis. *Plant Physiol* **185**, (2021).

754   53.   Salomé, P. A. & McClung, C. R. Pseudo-response Regulator 7 and 9 are partially
755   redundant genes essential for the temperature responsiveness of the Arabidopsis
756   circadian clock. *Plant Cell* **17**, (2005).

757   54.   Sauret-Güeto, S. *et al.* Systematic Tools for Reprogramming Plant Gene Expression in
758   a Simple Model, Marchantia polymorpha. *ACS Synth Biol* **9**, (2020).

759   55.   Pollak, B. *et al.* Loop assembly: a simple and open system for recursive fabrication of
760   DNA circuits. *New Phytologist* **222**, 628–640 (2019).

761   56.   Karimi, M., Inzé, D. & Depicker, A. GATEWAY[TM] vectors for Agrobacterium-mediated
762   plant transformation. *Trends in Plant Science* vol. 7 Preprint at
763   https://doi.org/10.1016/S1360-1385(02)02251-3 (2002).

764   57.   Kim, W. Y., Salomé, P. A., Fujiwara, S., Somers, D. E. & McClung, C. R. Characterization
765   of Pseudo-Response Regulators in Plants. in *Methods in Enzymology* vol. 471 (2010).

766   58.   Ronald, J. & Davis, S. J. Measuring Hypocotyl Length in Arabidopsis. in *Methods in*
767   *Molecular Biology* vol. 2398 (2022).

768   59.   Shor, E. *et al.* PIF-mediated sucrose regulation of the circadian oscillator is light
769   quality and temperature dependent. *Genes (Basel)* **9**, (2018).

770
771

Figure 1. TLWD proteins regulate circadian rhythms of gene expression

Figure 2. TLWD proteins are present throughout the diel cycle

Figure 3. TLWD proteins form complexes with PRRs

Figure 4. TLWD control growth by repressing *PIF4* and *PIF5* expression at night

Figure 5. TLWD are arbor proteins in the Arabidopsis circadian oscillator