Mac/Lac-tosylceramide regulates intestinal homeostasis and secretory cell fate commitment by facilitating Notch signaling

Kebei Tang[1, 2#], Xuewen Li[1#], Jiulong Hu[1, 3#], Jingyuan Shi[1], Yumei Li[1], Yansu Chen[1], Chang Yin[1], Fengchao Wang[1, 4], and Rongwen Xi[1, 4*]

[1] National Institute of Biological Sciences, No. 7 Science Park Road, Zhongguancun Life Science Park, Beijing 102206, China

[2] Academy for Advanced Interdisciplinary Studies, Peking University, Beijing 100871, China

[3] School of Life Sciences, Tsinghua University, Beijing 100084, China

[4] Tsinghua Institute of Multidisciplinary Biomedical Research, Tsinghua University, Beijing 102206, China.

[#] These authors contributed equally to this work.

* Correspondence: *xirongwen@nibs.ac.cn*

Abstract

Cell-to-cell communication via Delta-Notch signaling is widely used in various tissues and organs to regulate development and patterning; however, the mechanisms regulating Notch signaling for precise cell fate decisions remain poorly understood. Similar to mammals, the intestinal stem cells (ISCs) in the adult *Drosophila* midgut generate both absorptive and secretory cell progeny, guided by differential levels of Notch activation. Here we performed a forward genetic screen in *Drosophila* and identified glucosylceramide synthase (GlcT), a rate-limiting enzyme for glycosphingolipid (GSL) production, whose mutation causes the development of secretory cell tumors. Genetic analysis of the GSL synthesis pathway, combined with metabolite rescue experiments, revealed that the tumor formation is linked to a deficiency in Mactosylceramide/Lactosylceramide. This deficiency impaired the endocytic recycling of the Delta, subsequently reducing Notch signaling activation. Conditional knockout of *UGCG*, the mammalian ortholog of *GlcT*, in mouse small intestine caused an excessive differentiation of goblet cells, phenotypes similar to these caused by Notch inhibition. Our study suggests an evolutionarily conserved role for a specific GSL metabolite in modulating Notch signaling during stem cell fate decisions and provides a molecular connection between ceramide metabolism and Notch signaling in regulating tissue homeostasis and tumor formation.

 INTRODUCTION

Notch signaling is an evolutionarily conserved pathway in metazoans that mediates local cell-cell interactions through the membrane-tethered ligand Delta (Dl) and the membrane receptor Notch. This pathway transduces signals from the cell surface to the nucleus, regulating the transcription of target genes [1,2]. Dl-Notch signaling controls various cell fates and developmental processes, with mutations in the pathway implicated in numerous human diseases, including congenital defects and cancer [3-5]. Given that Notch can transduce, amplify, and consolidate molecular differences to influence cell fate decisions, it is crucial to understand how the strength of the Notch signaling pathway is precisely regulated to ensure proper cell fate determination and tissue development.

Notch signaling is known as a major regulator of cell fate decisions in both mammalian and *Drosophila* intestinal stem cell (ISC) lineages, as its activity determines the binary fate of intestinal progenitor cells. A high level of Notch activity promotes the specification of absorptive enterocytes, while low or absent Notch activity favors the specification of secretory cell types.[6,7] The *Drosophila* intestinal stem cell (ISC) lineage in the adult midgut is relatively simple, comprising only one type of secretory cell: enteroendocrine cells (EEs). This simplicity makes it an attractive experimental system for studying Notch signaling in cell fate decisions during epithelial renewal [6]. The fly ISCs, which specifically express Dl, periodically generate two types of committed daughter cells: enteroblasts, which experience high levels of Notch activation and are primed for the specification of absorptive enterocytes, and enteroendocrine progenitor cells (EEPs), which receive low levels of Notch activation. Typically, EEPs undergo one round of mitosis before terminal differentiation, resulting in the formation of EE pairs [8-11]. The loss of Notch signaling results in a complete blockage of enteroblast differentiation, leading to the accumulation of ISCs, EEPs and EEs in the intestinal epithelium, producing an "EE tumor" phenotype [8,10]. It is suggested that the expression level of Dl in individual ISCs varies, leading to differential levels of Notch activation in the immediate daughter cells [9]. This variation in Dl expression may result from dynamic bidirectional Dl-Notch signaling between ISCs and EEPs, which in parallel or in conjunction

with an intrinsic feedback mechanism in ISCs, guides a periodic activation of an EE fate

inducer Scute, thereby periodic generation of EEPs from ISCs [10,12,13]. The transient activation

of Scute induces irreversible Prospero expression in EEPs through a Phyl-Sina-Ttk69

regulatory cascade, initiating terminal differentiation into EEs [10,14-16]. During the period of

Scute inactivation, ISCs only generate enterocyte-committed enterobalsts by default via

Dl-Notch mediated lateral inhibition [8,17]. In newly enclosed flies, increased lipid intake from

food can alter the membrane trafficking of Dl, subsequently increasing the frequency of EE

specification from ISCs [18]. Thus, both intrinsic and extrinsic mechanisms are involved in

regulating Dl-Notch signaling to orchestrate stem cell fate decisions.

To better understand fate regulation in the *Drosophila* ISC lineage, here we conducted a

forward mosaic genetic screen in the adult *Drosophila* midgut, focusing on the right arm of

the second chromosome to identify genes whose mutations could disrupt ISC fate decisions.

From this screen, we identified several complementation groups that result in EE tumor

phenotypes. While most of the identified gene loci correspond to known components of the

Notch signaling pathway, we discovered one gene locus not previously associated with Notch

signaling. Further analyses revealed a specific GSL metabolite that regulates Notch signaling

and cell fate decisions in the intestinal epithelium, and this function appears to be conserved

from *Drosophila* to mice.

RESULTS

A mosaic genetic screen identifies *GlcT* as a tumor suppressor in the adult *Drosophila*

midgut

We performed chemical mutagenesis on chromosome 2R (carrying FRTG13/42B) and used

the MARCM system to induce mutant clones in the adult midgut (Supplementary Figure 1).

We screened approximately 10,000 lethal lines and identified totally 12 mutations across six

complementation groups that exhibited a "small cell tumor" phenotype, characterized by a

significant accumulation of diploid cells within the clones (Supplementary Figure 1 and

supplementary Table 1). Co-staining with the ISC marker DI and the EE cell marker Pros

revealed that most of these tumors contained an excessive number of ISCs and EEs, the EE

tumor phenotype commonly observed in Notch mutant clones [11,19]. Further genetic mapping

showed that three complementation groups were associated with genes known to function

in the Notch pathway: *mam*, *Gmer*, and *O-fut1* [20-22]. One complementation group, comprising

two alleles (*EA30* and *E230*), mapped to the *GlcT* gene (see details in Methods), while the

remaining two complementation groups, each consisting of one allele, remain undetermined

(Supplementary Table 1). *GlcT* encodes glucosylceramide synthase, an enzyme that catalyzes

the formation of glucosylceramide, the core component of glycosphingolipids (GSLs). Both

alleles carried one or more missense mutations resulting in the amino acid replacements

E292K (for *EA30*) and L395P/S397T (for *E230*, Figure S1D). The two mutations produced a

similar tumor phenotype in terms of tumor size and the composition of EEs in the tumors

(Figure 1B-D). The large polyploid cells were still present in the tumors, indicating that the

multipotency of ISCs was retained, but the differentiation of ISCs may have been biased

towards the EE fate commitment. Additionally, $GlcT^{\Delta 8}$, a previously characterized null allele of

GlcT [23], exhibited a similar EE tumor phenotype (Figure 1B-D). Expressing a *UAS-GlcT*

transgene in the mutant clones fully prevented the tumor phenotype (Figure 1E, F).

Furthermore, depleting *GlcT* in progenitor cells using RNAi with *esg-GAL4*[ts] also led to

overproliferation of ISCs and excessive generation of EEs in the midgut epithelium (Figure

1G-I). Taken together, these data suggest that *GlcT* acts as a tumor suppressor in adult

Drosophila midgut. Loss of *GlcT* induces ISC over-proliferation and excessive EE

differentiation, resulting in an EE tumor or "neuroendocrine tumor (NET)" phenotype.

Genetic analysis of the GSL synthesis pathway components reveals specific

tumor-suppressive activity for *egh*, in addition to *GlcT*

GlcT is involved in the initial step in the GSL biosynthesis pathway, which synthesizes

glucosylceramide by transferring glucose to ceramide (Figure 2A). The loss of *GlcT* could

result in the accumulation of ceramide or the absence of glucosylceramide and the

downstream GSL metabolites. Ceramide accumulation has been linked to promoting cell

apoptosis [24], which could potentially explain the ISC over-proliferation phenotype since

apoptotic cells can induce compensatory ISC proliferation [25]. However, we did not observe a

significant increase in apoptotic cells within the mutant clones (Figure 2B). Moreover, the

overexpression of p35, an inhibitor of cell apoptosis, failed to suppress or alleviate the tumor

phenotype (Figure 2B). Ceramidase (CDase) catalyzes ceramide into shingosine and free fatty

acid, and overexpressing *CDase* can in theory downregulate ceramide levels. We therefore

introduced *CDase* expression in the mutant clones, but this intervention also did not alleviate

the tumor phenotype (Figure 2B). Hence, the EE tumors developed from *GlcT* mutant clones

is unlikely caused by the accumulation of ceramide.

In the GSL biosynthesis pathway in *Drosophila*, glucosylceramide undergoes sequential

modifications catalyzed by Egghead (Egh) and Brainiac (Brn), which add the second

(Mactosyl-) and third (GlcNAc-) glycosyl residues, respectively (Figure 2A) [26,27]. This leads to

the synthesis of Mactosylceramide (MacCer) and GlcNAcβ1-3Manβ1-4Glcβ1Ceramide.

Complex GSL biosynthesis involves the participation of β4-N-Acetylgalactosyltransferase

(*β4GalNAcTA/TB*) and α1,4-Galactosyltransferase (*α4GT1/2*) in the fourth and fifth steps of

GSL sugar chain elongation [28,29]. These enzymes catalyze the formation of

GalNAcβ1-4GlcNAcβ1-3Manβ1-4Glcβ1Ceramide   and

GalNAc-α1-4GalNAcβ1-4GlcNAcβ1-3Manβ1-4Glcβ1Ceramide, respectively.

To understand the consequences of mutations in these GSL biosynthesis pathway genes, we

generated mutant MARCM clones for each of the above-mentioned genes and examined

their effects. Interestingly, mutations in *egh*, assessed using three different loss-of-function

alleles, consistently resulted in cell over-proliferation and excessive EE phenotypes, similar to

those caused by the loss of *GlcT* (Figure 2E-G). In contrast, none of the mutant clones of *brn*,

β4GalNAcTA, or *α4GT1* exhibited any noticeable abnormalities in clone size or percentages of

EEs, when compared to the control group of wild-type clones (Figure 2E-G). Therefore, GlcT

and Egh, but not other enzymes in the GSL biosynthesis pathway, exhibit specific tumor

suppressive activity in the adult *Drosophila* midgut.

The EEC tumor phenotype is a result of MacCer deficiency

As Egh is necessary for the biosynthesis of MacCer, the specific phenotype resulted from the

loss GlcT and Egh but not Brn could be attributed to the absence of MacCer. To test this

hypothesis, we fed flies carrying *GlcT* mutant clones with Lactosylceramide (LacCer), an

analog of MacCer that functions similarly to MacCer in mammalian cells. Remarkably, LacCer

significantly suppressed the overgrowth and excessive EE phenotype in the anterior midgut

clones (Figure 3B, C). The suppression effect was also significant, though relatively mild, in

the posterior midgut (Figure 3B, C). This difference may be attributed to substantial

absorption or metabolism of LacCer prior to its entry into the posterior midgut. Therefore,

among the various GSL metabolites, MacCer and LacCer appear to possess specific

tumor-suppressive activities in the *Drosophila* midgut.

Loss of *GlcT* leads to reduced activation of Notch signaling in ISC progenies.

In adult *Drosophila* midgut, the proliferation and differentiation of ISCs are regulated by

multiple signaling pathways. Among these, the EGFR/Ras/MAPK and JAK/STAT signaling

pathways play significant roles in promoting ISC proliferation during intestinal renewal and

regeneration following tissue damage or infection [30,31]. Additionally, Notch signaling not only

regulates ISC proliferation but also controls fate decisions between EE/EC during ISC

differentiation [19,32,33]. Loss of Notch leads to increased ISC proliferation and excessive EE

generation, phenotype similar to those caused by *GlcT* mutations. Therefore, we examined

whether the loss of *GlcT* affects the activities of these signaling pathways.

Normally, among the intestinal cells in the midgut epithelium, progenitor cells including ISCs

and enteroblasts, exhibit low but specific activation of EGFR and JAK/STAT activities [30,31,34].

The growth of *GlcT* mutant clones appeared to cause a general increase of EGFR and

JAK/STAT activities within the intestinal epithelium, as many differentiated cells also

exhibited EGFR or JAK/STAT signaling activation (Supplementary Figure 2), which is likely due

to stress responses induced by tumor growth [35]. Nevertheless, we observed that pERK, a

direct indicator of EGFR/Ras/MAPK signaling activity, was not significantly increased in *GlcT*

mutant cells compared to cells outside the clones (Supplementary Figure 2). Similarly, the

nuclear expression of STAT92E, which reflects JAK/STAT pathway activation, did not show

significant changes in *GlcT* mutant cells compared to normal cells outside the clones

(Supplementary Figure 2). However, the expression of *NRE-lacZ*, a reporter for Notch

pathway activation, was specifically down-regulated in *GlcT* mutant clones. Most cells within

the clones exhibited lower LacZ expression levels compared to *NRE-lacZ* positive cells outside

the clones (Figure 4A, B). Previous studies have shown that while the loss of function of

Notch completely blocks EC differentiation from ISCs, a reduction in Notch activity does not

necessarily hinder EC differentiation from ISCs. Instead, it may lead to a dose-dependent

increase in the ratio of differentiated EEs to ECs, depending on the severity of Notch loss [19].

Therefore, the increased EE to EC ratio and the reduced NRE-lacZ expression observed in

GlcT mutant clones collectively suggest that the mutant phenotype may be caused by a

reduction in Notch signaling activity.

In the ISC lineage, Notch signaling not only guides EC/EE fate decisions, but is also

participated in regulating EE subtype diversity. Typically, each EEP undergo one round of

asymmetric cell division to yield a pair of distinctive EEs, an allatostatin C (AstC)$^+$ class I EE

and a tachykinin (Tk)$^+$ class II EE [10]. The loss of *Notch* specifically compromises

Mirr-dependent specification of class II EE generation, resulting in all EEs in *Notch* mutant

clones becoming AstC$^+$ class I EEs [16,36]. We found that in *GlcT* mutant clones, virtually all EEs

were positive for AstC but TK (Figure 4C), further supporting the idea that Notch signaling is

compromised in these clones.

We next tested whether forced activation of Notch in *GlcT* mutant clones could suppress the

EE tumor phenotype. To achieve this, we expressed N^{intra}, an active form of Notch, in *GlcT*

mutant clones. As expected, this resulted in the complete suppression of cell proliferation in

GlcT mutant clones and promoted the differentiation of these cells into ECs, leading all

clones to become single-cell clones that often contained a polyploid cell (Figure 4D). This

finding suggests that *N^{intra}* is epistatic to *GlcT* in the signaling transduction pathway,

indicating that GlcT may regulate Notch signaling at the ligand/receptor level.

GlcT regulates the endocytic trafficking of Delta.

The observations above collectively suggest that full activation of Notch signaling in the ISC

lineage may require a specific GSL: Mac/Lac-Cer. As transmembrane proteins, the proper

internalization and recycling of Dl and Notch are important for the proper activation of Notch

signaling [37,38]. Specific lipids have been implicated in the formation of micro-domains on the

cell membrane, which play a role in regulating cell-cell signaling [39]. Therefore, we asked

whether the loss of MacCer caused by *GlcT* mutation could disrupt Notch signaling by

affecting the stability, endocytosis, or recycling of either Dl or Notch.

Knocking down *GlcT* in ISCs using *Dl-GAL4^{ts}* cased an increase in EEs in the midgut epithelium

(Figure 5 A, B). However, when *GlcT* was specifically knocked down in EBs using *NRE-GAL4^{ts}*,

there was no noticeable increase in EEs (Figure 5 A, B). This suggests that *GlcT* is specifically

required in ISCs and implies its potential role on the regulation of Dl. It is well-known that

the internalization of Dl in the signaling-sending cell is necessary for Notch activation in the

signaling-receiving cell [37]. To investigate whether the endocytosis process of Dl is altered in

GlcT mutant ISCs, we performed an *ex vivo* Dl internalization assay, following the previously

described protocol [40]. In this assay, we cultured and incubated the freshly dissected gut with

anti-Dl antibodies, and the endocytosis of the antibody-labeled Dl was visualized after

fixation and immunostaining with secondary antibodies. Interestingly, during the 3-hour

period, we observed a significant difference in the subcellular localization pattern of Dl

between normal and *GlcT* mutant ISCs (Figure 5 C). In *GlcT* mutant ISCs, the majority of

labeled Dl had already been internalized, while in wild-type ISCs, it remained on the cell

membrane. This observation suggests that the loss of *GlcT* causes either reduced stability of

Dl on the cell membrane or accelerated internalization of Dl from the cell membrane.

To further characterize the endocytic trafficking of Dl in *GlcT* mutant ISCs, we performed

conditional depletion of *GlcT* and examined the co-localization of internalized Dl with the

early endosome marker Rab5 and the late endosome marker Rab7, respectively. In

comparison to control ISCs, *GlcT-RNAi* ISCs exhibited a significant increase in early

endosomes containing Dl (Figure 5D, F). However, the percentage of late endosomes

containing Dl remained similar between wild-type and *GlcT-RNAi* ISCs (Figure 5E, F). These

observations suggest that the internalized Dl in *GlcT-RNAi* ISCs experiences a delay in early

endosomes, which might cause a delay in Dl recycling and consequently a reduction in

Dl-Notch signaling activity (Figure 7E).

GlcT shows tissue specificity in regulating Notch signaling activity

Notch signaling has pleiotropic function and is known to be participated in the development

of diverse tissue and organs. We therefore asked whether the discovered mechanism above

is generally utilized in controlling Notch signaling in diverse tissues and organs. In the wing

disc of third instar larvae, Notch signaling is activated by ligands Dl and Serrate at the

dorsal-ventral boundary to induce the expression of Cut [41]. We found that in *GlcT* mutant

clones spanning the dorsal-ventral boundary region of the wing disc, the expression of Cut

was not significantly altered (Figure 6A). This indicates that *GlcT* is not essential for

regulating Notch-mediated Cut expression in this specific context.

During oogenesis in the *Drosophila* ovary, Notch signaling regulates differentiation and

transition of follicle cells from the mitotic cell cycle to the endocycle during stage 6 [42,43]. We

found that in the developing egg chambers with germline mutations of *GlcT*, the patterning

of the surrounding follicle cells remained largely normal (Figure 6B). However, we observed

an abnormal accumulation of Dl protein in cytoplasmic vesicles of the mutant germline cells

(Figure 6C), indicating that there is a defect in Dl trafficking in *GlcT* mutant germline cells.

However, this defect is not sufficient to impact Dl-Notch signaling to a degree where follicle

cell development is adversely affected. Collectively, these observations suggest that the role

for GlcT in Notch signaling does not appear to be a universal mechanism across all

developmental contexts. Even in situations where this mechanism is involved, the extent to

which GlcT affects Dl-Notch signaling may vary in different developmental scenarios.

Transient loss of *UGCG* in mouse small intestine causes reduced number of ISCs and

increased number of goblet cells

As both the Notch signaling and the metabolic pathway of GSLs are conserved from

Drosophila to mammals, we asked whether this mechanism is conserved in the mouse small

intestine to regulate the proliferation and differentiation of ISCs. Notch is known to control

the absorptive versus secretory cell fate of mammalian intestinal progenitors as well, and the

inhibition of Notch signaling in the intestinal epithelium of mouse small intestine leads

excessive number of goblet cells [44,45]. However, unlike in *Drosophila*, Notch positively

regulates ISC self-renewal in mice. The inhibition of Notch causes ISC loss, and the excessive

activation of Notch promotes the proliferation of intestinal progenitor cells [45,46]. We

therefore performed conditional knockout of *UGCG*, which encodes the mammalian

glucosylceramide synthase, in the mouse small intestine and examined the consequences.

It has been previously reported that in the new born mice with intestine-specific knockout of

UGCG, the proliferation of enterocytes appeared largely normal. However, 3-4 days after

intestinal-specific knockout of UGCG in adult mice, epithelial phenotypes including epithelial

hyperproliferation, impaired lipid absorption and the detachment of enterocytes from the

basal lamina were observed, and these phenotypes are considered to be consequences of

gut barrier disruption [47]. To separate the primary from the secondary phenotypes, we

repeated the experiment by generating *UGCG^flox/flox; Vil-CreERT2* mice, and examined the

phenotype at 48h post tamoxifen administration, an early timepoint with a hope that the

secondary phenotypes have not yet developed.

Consistent with the previous findings, the conditional knockout mice showed a severe

diarrhea phenotype and died several days following tamoxifen induction. Interestingly, we

observed a few phenotypes at the 48h timepoint that was not observed before. We found

that in the mutant intestine, there was a marked increase in crypt depth accompanied by a

marked reduction in villus length (Supplementary Figure 3). In addition, an excess population

of goblet cells was observed, dispersed along the crypt-villus axis (Figure 7C, D). By

examining the expression of Olfm4, which is transcriptionally regulated by Notch in ISCs [48],

we found that the fluorescence intensity of Olfm4 staining was significantly reduced in *UGCG*

mutant ISCs (Figure 7A, B). It is unclear how the crypt-villus morphology phenotype is

developed in *UGCG* mutant epithelium, but the increased goblet cells and the reduced

expression level of Olfm4 are consistent with a Notch phenotype, suggesting a conserved

role for glucosylceramide synthase in modulating Notch signaling from *Drosophila* to

mammals.

Discussion

From an unbiased genetic screen on the right arm of chromosome II, here we have

pinpointed four mutant loci that cause an excessive EE phenotype in the intestinal

epithelium. Three of them encode known components of the Notch signaling (*mam*, *Gmer*,

and *O-fut1*), and the remaining one, *GlcT*, turns out to be a new Notch regulator as well. This

work further reinforces the notion that Notch as the key regulator of the secretory versus

absorptive cell fate decisions from ISCs, and reveals an evolutionarily conserved link between

GSL metabolism and Notch signaling in regulating intestinal homeostasis.

As *GlcT* encodes glucosylceramide synthase, which that catalyzes the formation of

glucosylceramide from ceramide, we further analyzed *egh* and *brn*, which encodes enzymes

for the subsequent sequential modifications of glucosylceramide by adding the second

(Mactosyl-) and third (GlcNAc-) glycosyl residues, respectively. We found that loss of *egh*, but

not *brn*, can give rise to a similar excessive EE phenotype. In addition, the phenotype caused

by *GlcT* can be suppressed by feeding with LacCer, which is analogous to MacCer. Collectively,

these observations demonstrate that the excessive EE phenotype caused by *GlcT* or *egh*

mutation is a result of MacCer deficiency.

GSLs constitute a specific category of glycolipids localized on the outer leaflet of eukaryotic

cell membranes, and studies have implicated their roles in regulating multiple cellular

signaling pathways by serving on the lipid raft to facilitate signaling transductions [49-52]. In

particular, there is an evolutionarily conserved phospholipid binding domain (C2 domain) on

the Notch ligand that is known to interacts with GSLs, and this interaction facilitates

ligand-receptor interactions and thereby promotes robust Notch signaling [53-55]. Our analyses

of *GlcT* mutant cells in the *Drosophila* midgut suggest that MacCer may represent a key

member of GSLs in facilitating Dl-Notch signaling transduction between intestinal stem cells

(ISCs) and their immediate progeny to steer binary cell fate decisions. To do so, MacCer may

help to modulate the endocytic trafficking of Dl, an event that is known to be essential for

Notch activation. We found that Dl on membrane of *GlcT* mutant ISCs tends be rapidly

endocytosed and the resultant endosomes appear to experience a delay in transition from

early endosome to late endosome. Prior genetic investigations have linked glycosphingolipids

synthesized by α4GT1 to the modulation of Dl ligand activity i*n vivo*. While the loss of α4GT1

does not overtly affect Dl-Notch signaling, it can exacerbate the wing phenotype of

haploinsufficient *Notch* mutants. Furthermore, the overexpression of α4GT1 has been shown

to counteract signaling and endocytosis dysfunctions of Dl caused by the inhibition of the E3

ligase Mib1 or Neur activity [50]. Therefore, the robust genetic results presented in this study

provide definitive evidence for a functional role of MacCer and its downstream derivatives in

regulating Notch signaling. Future structural studies focusing on the interactions between

GSLs and Dl should offer valuable insights into the role of GSLs in modulating Dl-Notch interactions and membrane trafficking processes.

One intriguing observation from our study is the tissue-specific regulatory role of *GlcT* in Notch signaling. Its loss in ISCs significantly impairs Notch signaling, leading to the formation of EE tumors. However, when *GlcT* is lost in the germline cyst within the developing egg chamber, it only modestly disrupts the endocytic trafficking of Dl, without conspicuous effects on the differentiation and patterning of surrounding follicle cells. In the context of imaginal discs, the loss of *GlcT* appears to have no discernible impact on Notch signaling activity at all. This tissue-specific modulation of Notch signaling by *GlcT* suggests a potential variability in membrane lipid composition across different cell types. Such variability could render certain cells more susceptible to alterations in specific GSLs and their influence on Dl-Notch signaling transduction, while others remain unaffected. As Dl trafficking in *Drosophila* ISCs can be modulated by dietary lipid content, which impacts the duration and level of Notch activation to control the production of EEs in newly enclosed flies [18], the sensitivity of ISCs to specific GSL content may serve as a mechanism enabling ISCs to govern the fate of their progeny based on nutrient content.

Taken together, our study identified MacCer/LacCer as a tissue-specific regulator of Dl-Notch signaling by regulating the membrane localization and endocytic trafficking of Dl. As Notch signaling has pleiotropic roles in many developmental processes, diseases and cancer, the findings should contribute to our understanding of Notch signaling transduction and may help to open up new approaches for developing prevention and treatments against human diseases and cancer.

ACKNOWLEDGEMENTS

We thank Drs. Feng Yu and Sanduo Zheng for technical assistance, Drs. Dick Nassel,

Jan-Adrianus Veenstra, Zhaohui Wang, and the Developmental Studies Hybridoma Bank

(DSHB) for antibodies, Drs. Sarah Bray, Steven Hou, Akiko Satoh, Tao Wang, the Bloomington

Drosophila Stock Center (BDSC) and Vienna Drosophila Resource Center (VDRC) for fly stocks,

and Dr. Sylvie Robine for the villin-Cre[ERT2] mice. This work was supported by National Key

Research and Development Program of China (2020YFA0803502 and 2017YFA0103602 to

R.X.) from the Chinese Ministry of Science and Technology.

DECLARATION OFINTERESTS

381 The authors declare no competing interests.

MATERIALS AND METHODS

Drosophila Stocks and mice

Drosophila melanogaster stocks were maintained on standard cornmeal molasses food in a

25°C incubator. Flies for all experiments were maintained at a constant 25°C with 15–25 flies

per vial and tossed onto new food every 2–3 days. *Drosophila* stocks used in this study: *hsflp,*

act-GAL4, UAS-GFP; FRT42B tub-GAL80; hsflp, act-GAL4, UAS-GFP; FRT42D tub-GAL80; and

hsflp, Tub-GAL80, FRT19A; act-GAL4,UAS-GFP/Cyo [56]; *hsflp; FRT42D, arm-lacZ/Cyo; esg-GAL4,*

UAS-GFP [56]; *Su(H)GBE (NRE)-GAL4;* and *Dl-Gal4* [57]; *Su(H)GBE-lacZ* (NRE-lacZ, gift from Sarah

Bray); *Glc^{EA30}*(generated in this study); *Glc^{E230}*(generated in this study); *GlcT^{Δ8}* [23]; *Tub-GAL80*^{ts}

[58]; *UAS-GlcT* (generated in this study); *GlcT-RNAi* #1 (VDRC, id:44912); *GlcT-RNAi* #2 (VDRC,

id:108064); *UAS-CDase* (gift from Tao Wang); *UAS-p35* (BDSC, #6298); *egh^A* (BDSC, #52353);

egh^B (BDSC, #52354); *egh^7* (BDSC, #77889); *brn^{228}* (BDSC, #7392); *β4GalNAcTA-RNAi* (VDRC,

id:4867); *α4GT1-RNAi* (VDRC, id:2608); *UAS-N^{intra}* [56]; *UAS-GFP-Rab5* (BDSC, #43336).

Mice used in this study include floxed *UGCG*, which was generated as previously described [47],

and villin-CreERT2 [59]. All the research performance underwent strict ethical review and was

approved by the Ethics Committee of the National Institute of Biological Sciences, Beijing.

Mice were housed at the animal center in the National Institute of Biological Sciences in a

Specific Pathogen Free (SPF) facility.

EMS Mutagenesis in *Drosophila*

3~5 day-old, isogenized y w; FRT42B(G13) males were subjected to a 10-hour starvation

period before being fed with EMS (30 mM in sucrose solution applied on filter papers)

overnight. Following recovery, these males were mated with y w; Sco/Cyo virgin females. In

the F1 offspring, y w; FRT42B */Cyo males were individually mated with several y w; Sco/Cyo

virgin females. The y w; FRT42B */Cyo males and virgin females in the F2 generation were

carefully selected and crossed with each other to establish balanced stocks for further

genetic studies. Altogether, around 10,000 stocks harboring lethal mutations were obtained

and screened.

Deficiency Mapping

The complete Bloomington Deficiency Kit for chromosome 2R

(https://bdsc.indiana.edu/stocks/df/dfkit-info.html) was used for initial mapping, which

helped to localize both the EA30 and E230 lethal mutations to a cytogenetic location

between 58A3 and 58F1. Subsequent mapping with additional deficiency lines encompassing

this region allowed the identification of two small deficiency lines, Df(2R)Exel7170 and

Df(2R)Excel6078, which failed to complement both mutations. This refinement narrowed the

critical region down to 58B2-58C1. There are 13 protein-coding genes within this region, and

genomic sequencing data indicated that both lines carried missense mutations on the *GlcT*

gene, as described in the text.

Genomic Sequencing and Analysis Method

As all mutants exhibited lethality in the adult stage, homozygous larvae were selected using a

green balancer, and the extracted genomic DNAs were sequenced at BGI Genomics using a

library with an insert size of ≤800bp. The sequencing platform employed was the HiSeq 2000.

For sequence analysis, all mutants were compared to a reference genome. Subsequently,

sequencing reads were assigned to each allele based on heterozygote SNPs retained using a

custom Perl script. The candidate heterozygous sites were annotated using snpEff.

Mosaic Analysis with a Repressible Cell Marker (MARCM) Analysis

To generate MARCM clones [60], flies of the desired genotype were raised and maintained at

25°C until they reached 7 days of age. Subsequently, they were subjected to a 60-minute

heat shock in a water bath at 37°C. Following the heat shock, the flies were fed with standard

food supplemented with yeast paste and this feeding regimen was repeated every 2 days

before the flies were dissected for further analysis.

GAL4/GAL80^{ts} System

The GAL4/UAS and GAL80ts systems were utilized as previously described [58,61]. To inhibit the

GAL4 system, the crosses were sustained at 18°C when employing temperature-sensitive

GAL4-mediated RNAi or gene overexpression. Subsequently, F1 adult flies with the correct

genotype were transitioned to 29°C to activate the GAL4 system, thereby triggering RNAi or

gene overexpression.

Tamoxifen Treatment in Mice

Cre recombinase activity was induced via intraperitoneal injection of 2 mg tamoxifen

(Sigma-Aldrich, T5648), and tissues were collected 48 hours post-injection. Cre-negative

littermates served as controls.

Immunofluorescent Staining and Imaging

Adult female midguts were dissected in PBS and fixed for 30 minutes at room temperature in

4% paraformaldehyde. The fixed tissues were then dehydrated in methanol for 5 minutes

followed by rehydration in a PBT solution (PBS containing 0.1% Triton X-100). The tissues

underwent four washes with 0.1% Triton in 1×PBS (PBST) and were subsequently incubated

with PBST and primary antibodies overnight at 4°C. After another round of washing, the

samples were incubated for 2 hours with secondary antibodies in PBST. Finally, the samples

were stained with DAPI for visualization of nuclei.

Mouse small intestine was dissected out and gently flushed with cold 1× PBS to remove fecal

content, and fixed overnight at 4°C in 4% paraformaldehyde (PFA) prepared in 1× PBS. The

tissue was then incubated in a 30% sucrose solution at 4°C for 24 hours, and embedded in

OCT (Tissue-Tek) before frozen. Frozen sections of 15 μm thickness were prepared for further

analysis. For immunostaining, tissue sections were permeabilized with 1% Triton X-100

(Sigma-Aldrich) in PBS for 1 hour at room temperature (RT), followed by blocking with a solution containing 0.2% Triton X-100, 1% bovine serum albumin (BSA), and 3% goat serum in PBS for 1 hour at RT before antibody staining. The sections were mounted in Vectashield mounting media with DAPI (Vector Laboratories) for imaging.

Primary antibodies used in this study: mouse anti-Pros (DSHB #MR1A; 1:300); mouse anti-Dl (DSHB Cat#C594.9B; RRID:AB_528194; 1:300); rabbit anti-AstC (gift from Dr. Dick Nassel; 1:300); rabbit anti-Tk (gift from Dr. Jan-Adrianus Veenstra; 1:300); Rabbit anti-pH3 (CST Cat# 9701; RRID:AB_331535; 1:500); mouse monoclonal anti-β-galactosidase (DSHB, #40-1a; 1:30); mouse anti-Cut (DSHB, #2B10; 1:20); mouse anti-Rab7 (DSHB, 1:300); rabbit anti-pERK (CST, 1:200); rabbit anti-STAT92E (gift from Dr. Zhaohui Wang, 1:300); rabbit anti-DCP-1 (CST Cat#9578S; 1:300); mouse monoclonal anti-EYA (DSHB, #10H6; 1:300); rabbit monoclonal anti-GFP (Invitrogen Cat#G10362; 1:300). Secondary antibodies used in this study include Alexa Fluor 568- or Cy5-conjugated goat anti-rabbit, anti-mouse IgGs (Molecular Probes, A11034-A11036, A10524; 1:300). For nuclei staining, DAPI (Sigma-Aldrich, 1 μg/ml) was used. The preparations were mounted in 70% glycerol, and the slides were stored at -20°C. Imaging was conducted using a Leica SP8 confocal microscope. All captured images were processed and compiled using Adobe Photoshop and Illustrator.

LacCer Treatment

The 2-3 days old female flies of the specified genotype were moved to fresh food layered with a filter paper soaked in a 5% sucrose solution containing 20 μM of LacCer (Sigma-Aldrich CAS#4682-48-8) and this transfer was repeated daily before dissection and analysis.

Antibody Uptake Assay

The anti-Dl antibody Uptake assay was conducted following a previously described protocol [40]. In brief, the *Drosophila* gut was dissected and placed in fresh medium containing anti-Delta. The dissecting dish was then placed in a humidified box and incubated at 25 °C for the necessary duration. Subsequently, the medium containing the antibody was removed, and a single wash with Schneider's *Drosophila* medium was carried out. The midgut was then fixed directly in the dissecting dish for 20 minutes under a chemical fume hood by adding 1 ml of fixative. Following this, samples underwent three washes with PBT and were then

incubated for 60 minutes with fluorescently labeled anti-mouse secondary antibodies in PBT

on a rotating platform. After another three washes with PBT, the samples were mounted in

50% glycerol in PBS for analysis under microscopy.

Fluorescence Intensity Statistics

The Image J software (downloaded from https://imagej.nih.gov/ij/) was utilized to measure

the fluorescence intensity of the captured images. The corrected fluorescence intensity is

determined as the average fluorescence intensity of each cell's nuclear region minus the

average background fluorescence intensity.

Statistical analysis

All quantifications were presented in the form of mean ± SEM. GraphPad Prism 5 software

(GraphPad Software Inc.) was used to calculate p-values using an unpaired Student's t-test.

The number of intestines used for calculations was labeled on the figures or in the related

figure legends.

REFERENCES

1.   Bray, S.J. (2006). Notch signalling: a simple pathway becomes complex. Nat Rev Mol Cell Biol
7, 678-689. 10.1038/nrm2009.
2.   Kopan, R., and Ilagan, M.X. (2009). The canonical Notch signaling pathway: unfolding the
activation mechanism. Cell *137*, 216-233. 10.1016/j.cell.2009.03.045.
3.   Artavanis-Tsakonas, S., Rand, M.D., and Lake, R.J. (1999). Notch signaling: cell fate control and
signal integration in development. Science *284*, 770-776.
4.   Penton, A.L., Leonard, L.D., and Spinner, N.B. (2012). Notch signaling in human development
and disease. Semin Cell Dev Biol *23*, 450-457. 10.1016/j.semcdb.2012.01.010.
5.   Bolos, V., Grego-Bessa, J., and de la Pompa, J.L. (2007). Notch signaling in development and
cancer. Endocr Rev *28*, 339-363. 10.1210/er.2006-0046.
6.   Boumard, B., and Bardin, A.J. (2021). An amuse-bouche of stem cell regulation: Underlying
principles and mechanisms from adult Drosophila intestinal stem cells. Curr Opin Cell Biol *73*,
58-68. 10.1016/j.ceb.2021.05.007.
7.   Koch, U., Lehal, R., and Radtke, F. (2013). Stem cells living with a Notch. Development *140*,
689-704. 10.1242/dev.080614.

8.   Ohlstein, B., and Spradling, A. (2007). Multipotent Drosophila intestinal stem cells specify
daughter cell fates by differential notch signaling. Science *315*, 988-992. 315/5814/988
[pii10.1126/science.1136606 [doi].

9.   Ohlstein, B., and Spradling, A. (2006). The adult Drosophila posterior midgut is maintained by
pluripotent stem cells. Nature *439*, 470-474. nature04333 [pii]10.1038/nature04333 [doi].

10.   Chen, J., Xu, N., Wang, C., Huang, P., Huang, H., Jin, Z., Yu, Z., Cai, T., Jiao, R., and Xi, R. (2018).
Transient Scute activation via a self-stimulatory loop directs enteroendocrine cell pair
specification from self-renewing intestinal stem cells. Nat Cell Biol *20*, 152-161.
10.1038/s41556-017-0020-0.

11.   Micchelli, C.A., and Perrimon, N. (2006). Evidence that stem cells reside in the adult
Drosophila midgut epithelium. Nature *439*, 475-479. nature04371 [pii]10.1038/nature04371
[doi].

12.   Guo, Z., and Ohlstein, B. (2015). Stem cell regulation. Bidirectional Notch signaling regulates
Drosophila intestinal stem cell multipotency. Science *350*. 10.1126/science.aab0988.

13.   Henrique, D., and Schweisguth, F. (2019). Mechanisms of Notch signaling: a simple logic
deployed in time and space. Development *146*. 10.1242/dev.172148.

14.   Yin, C., and Xi, R. (2018). A Phyllopod-Mediated Feedback Loop Promotes Intestinal Stem Cell
Enteroendocrine Commitment in Drosophila. Stem Cell Reports *10*, 43-57.
10.1016/j.stemcr.2017.11.014.

15.   Wang, C., Guo, X., Dou, K., Chen, H., and Xi, R. (2015). Ttk69 acts as a master repressor of
enteroendocrine cell specification in Drosophila intestinal stem cell lineages. Development
142, 3321-3331. 10.1242/dev.123208.

16.   Guo, X., Zhang, Y., Huang, H., and Xi, R. (2022). A hierarchical transcription factor cascade
regulates enteroendocrine cell diversity and plasticity in Drosophila. Nat Commun *13*, 6525.
10.1038/s41467-022-34270-0.

17.   de Navascues, J., Perdigoto, C.N., Bian, Y., Schneider, M.H., Bardin, A.J., Martinez-Arias, A.,
and Simons, B.D. (2012). Drosophila midgut homeostasis involves neutral competition
between symmetrically dividing intestinal stem cells. EMBO J *31*, 2473-2485.
10.1038/emboj.2012.106.

18.   Obniski, R., Sieber, M., and Spradling, A.C. (2018). Dietary Lipids Modulate Notch Signaling
and Influence Adult Intestinal Development and Metabolism in Drosophila. Dev Cell *47*,
98-111 e115. 10.1016/j.devcel.2018.08.013.

19.   Perdigoto, C.N., Schweisguth, F., and Bardin, A.J. (2011). Distinct levels of Notch activity for
commitment and terminal differentiation of stem cells in the adult fly intestine. Development
138, 4585-4595. dev.065292 [pii]10.1242/dev.065292.

20.   Ayukawa, T., Matsumoto, K., Ishikawa, H.O., Ishio, A., Yamakawa, T., Aoyama, N., Suzuki, T.,
and Matsuno, K. (2012). Rescue of Notch signaling in cells incapable of GDP-L-fucose
synthesis by gap junction transfer of GDP-L-fucose in Drosophila. Proc Natl Acad Sci U S A *109*,
15318-15323. 10.1073/pnas.1202369109.

21.   Petcherski, A.G., and Kimble, J. (2000). Mastermind is a putative activator for Notch. Curr Biol
10, R471-473. 10.1016/s0960-9822(00)00577-7.

22.   Sasamura, T., Ishikawa, H.O., Sasaki, N., Higashi, S., Kanai, M., Nakao, S., Ayukawa, T., Aigaki, T.,
Noda, K., Miyoshi, E., et al. (2007). The O-fucosyltransferase O-fut1 is an extracellular

570		component that is essential for the constitutive endocytic trafficking of Notch in Drosophila.
571		Development *134*, 1347-1356. 10.1242/dev.02811.
572	23.	Satoh, T., Inagaki, T., Liu, Z., Watanabe, R., and Satoh, A.K. (2013). GPI biosynthesis is essential
573		for rhodopsin sorting at the trans-Golgi network in Drosophila photoreceptors. Development
574		*140*, 385-394. 10.1242/dev.083683.
575	24.	Selzner, M., Bielawska, A., Morse, M.A., Rudiger, H.A., Sindram, D., Hannun, Y.A., and Clavien,
576		P.A. (2001). Induction of apoptotic cell death and prevention of tumor growth by ceramide
577		analogues in metastatic human colon cancer. Cancer Res *61*, 1233-1240.
578	25.	Ryoo, H.D., Gorenc, T., and Steller, H. (2004). Apoptotic cells can induce compensatory cell
579		proliferation through the JNK and the Wingless signaling pathways. Dev Cell *7*, 491-501.
580		10.1016/j.devcel.2004.08.019.
581	26.	Wandall, H.H., Pizette, S., Pedersen, J.W., Eichert, H., Levery, S.B., Mandel, U., Cohen, S.M.,
582		and Clausen, H. (2005). Egghead and brainiac are essential for glycosphingolipid biosynthesis
583		in vivo. J Biol Chem *280*, 4858-4863. 10.1074/jbc.C400571200.
584	27.	Muller, R., Altmann, F., Zhou, D., and Hennet, T. (2002). The Drosophila melanogaster brainiac
585		protein is a glycolipid-specific beta 1,3N-acetylglucosaminyltransferase. J Biol Chem *277*,
586		32417-32420. 10.1074/jbc.C200381200.
587	28.	Stolz, A., Haines, N., Pich, A., Irvine, K.D., Hokke, C.H., Deelder, A.M., Gerardy-Schahn, R.,
588		Wuhrer, M., and Bakker, H. (2008). Distinct contributions of beta 4GalNAcTA and beta
589		4GalNAcTB to Drosophila glycosphingolipid biosynthesis. Glycoconj J *25*, 167-175.
590		10.1007/s10719-007-9069-5.
591	29.	Chen, Y.W., Pedersen, J.W., Wandall, H.H., Levery, S.B., Pizette, S., Clausen, H., and Cohen, S.M.
592		(2007). Glycosphingolipids with extended sugar chain have specialized functions in
593		development and behavior of Drosophila. Dev Biol *306*, 736-749.
594		10.1016/j.ydbio.2007.04.013.
595	30.	Xu, N., Wang, S.Q., Tan, D., Gao, Y., Lin, G., and Xi, R. (2011). EGFR, Wingless and JAK/STAT
596		signaling cooperatively maintain Drosophila intestinal stem cells. Dev Biol *354*, 31-43.
597		S0012-1606(11)00182-5 [pii]10.1016/j.ydbio.2011.03.018 [doi].
598	31.	Jiang, H., Patel, P.H., Kohlmaier, A., Grenley, M.O., McEwen, D.G., and Edgar, B.A. (2009).
599		Cytokine/Jak/Stat signaling mediates regeneration and homeostasis in the Drosophila midgut.
600		Cell *137*, 1343-1355. S0092-8674(09)00571-6 [pii]10.1016/j.cell.2009.05.014 [doi].
601	32.	Ohlstein, B., and Spradling, A. (2007). Multipotent Drosophila intestinal stem cells specify
602		daughter cell fates by differential notch signaling. Science *315*, 988-992.
603	33.	Guo, Z., and Ohlstein, B. (2015). Bidirectional Notch signaling regulates Drosophila intestinal
604		stem cell multipotency. Science *350*, aab0988.
605	34.	Jiang, H., Grenley, M.O., Bravo, M.J., Blumhagen, R.Z., and Edgar, B.A. (2011).
606		EGFR/Ras/MAPK signaling mediates adult midgut epithelial homeostasis and regeneration in
607		Drosophila. Cell Stem Cell *8*, 84-95.
608	35.	Patel, P.H., Dutta, D., and Edgar, B.A. (2015). Niche appropriation by Drosophila intestinal
609		stem cell tumours. Nat Cell Biol *17*, 1182-1192. 10.1038/ncb3214.
610	36.	Beehler-Evans, R., and Micchelli, C.A. (2015). Generation of enteroendocrine cell diversity in
611		midgut stem cell lineages. Development *142*, 654-664. 10.1242/dev.114959.
612	37.	Bray, S.J., and Gomez-Lamarca, M. (2018). Notch after cleavage. Current opinion in cell
613		biology *51*, 103-109.

38. Yamamoto, S., Charng, W.L., and Bellen, H.J. (2010). Endocytosis and intracellular trafficking
of Notch and its ligands. Curr Top Dev Biol *92*, 165-200. 10.1016/S0070-2153(10)92005-X.

39. Simons, K., and Toomre, D. (2000). Lipid rafts and signal transduction. Nat Rev Mol Cell Biol *1*,
31-39. 10.1038/35036052.

40. Couturier, L., and Schweisguth, F. (2014). Antibody uptake assay and in vivo imaging to study
intracellular trafficking of Notch and Delta in Drosophila. Methods Mol Biol *1187*, 79-86.
10.1007/978-1-4939-1139-4_6.

41. Micchelli, C.A., Rulifson, E.J., and Blair, S.S. (1997). The function and regulation of cut
expression on the wing margin of Drosophila: Notch, Wingless and a dominant negative role
for Delta and Serrate. Development *124*, 1485-1495.

42. Deng, W.-M., Althauser, C., and Ruohola-Baker, H. (2001). Notch-Delta signaling induces a
transition from mitotic cell cycle to endocycle in Drosophila follicle cells.

43. López-Schier, H., and Johnston, D.S. (2001). Delta signaling from the germ line controls the
proliferation and differentiation of the somatic follicle cells during Drosophila oogenesis.
Genes & development *15*, 1393-1405.

44. van Es, J.H., van Gijn, M.E., Riccio, O., van den Born, M., Vooijs, M., Begthel, H., Cozijnsen, M.,
Robine, S., Winton, D.J., Radtke, F., and Clevers, H. (2005). Notch/gamma-secretase inhibition
turns proliferative cells in intestinal crypts and adenomas into goblet cells. Nature *435*,
959-963.

45. Riccio, O., van Gijn, M.E., Bezdek, A.C., Pellegrinet, L., van Es, J.H., Zimber-Strobl, U., Strobl,
L.J., Honjo, T., Clevers, H., and Radtke, F. (2008). Loss of intestinal crypt progenitor cells owing
to inactivation of both Notch1 and Notch2 is accompanied by derepression of CDK inhibitors
p27Kip1 and p57Kip2. EMBO Rep *9*, 377-383. 10.1038/embor.2008.7.

46. Fre, S., Huyghe, M., Mourikis, P., Robine, S., Louvard, D., and Artavanis-Tsakonas, S. (2005).
Notch signals control the fate of immature progenitor cells in the intestine. Nature *435*,
964-968.

47. Jennemann, R., Kaden, S., Sandhoff, R., Nordstrom, V., Wang, S., Volz, M., Robine, S., Amen,
N., Rothermel, U., Wiegandt, H., and Grone, H.J. (2012). Glycosphingolipids are essential for
intestinal endocytic function. J Biol Chem *287*, 32598-32616. 10.1074/jbc.M112.371005.

48. VanDussen, K.L., Carulli, A.J., Keeley, T.M., Patel, S.R., Puthoff, B.J., Magness, S.T., Tran, I.T.,
Maillard, I., Siebel, C., Kolterud, A., et al. (2012). Notch signaling modulates proliferation and
differentiation of intestinal crypt base columnar stem cells. Development *139*, 488-497.
10.1242/dev.070763.

49. Coskun, U., Grzybek, M., Drechsel, D., and Simons, K. (2011). Regulation of human EGF
receptor by lipids. Proc Natl Acad Sci U S A *108*, 9044-9048. 10.1073/pnas.1105666108.

50. Hamel, S., Fantini, J., and Schweisguth, F. (2010). Notch ligand activity is modulated by
glycosphingolipid membrane composition in Drosophila melanogaster. J Cell Biol *188*,
581-594. 10.1083/jcb.200907116.

51. Ideo, H., Seko, A., Ishizuka, I., and Yamashita, K. (2003). The N-terminal carbohydrate
recognition domain of galectin-8 recognizes specific glycosphingolipids with high affinity.
Glycobiology *13*, 713-723. 10.1093/glycob/cwg094.

52. Park, S.Y., Kwak, C.Y., Shayman, J.A., and Kim, J.H. (2012). Globoside promotes activation of
ERK by interaction with the epidermal growth factor receptor. Biochim Biophys Acta *1820*,
1141-1148. 10.1016/j.bbagen.2012.04.008.

53.   Suckling, R.J., Korona, B., Whiteman, P., Chillakuri, C., Holt, L., Handford, P.A., and Lea, S.M.
(2017). Structural and functional dissection of the interplay between lipid and Notch binding
by human Notch ligands. EMBO J *36*, 2204-2215. 10.15252/embj.201796632.
54.   Martins, T., Meng, Y., Korona, B., Suckling, R., Johnson, S., Handford, P.A., Lea, S.M., and Bray,
S.J. (2021). The conserved C2 phospholipid-binding domain in Delta contributes to robust
Notch signalling. EMBO Rep *22*, e52729. 10.15252/embr.202152729.
55.   Chillakuri, C.R., Sheppard, D., Ilagan, M.X., Holt, L.R., Abbott, F., Liang, S., Kopan, R., Handford,
P.A., and Lea, S.M. (2013). Structural analysis uncovers lipid-binding properties of Notch
ligands. Cell Rep *5*, 861-867. 10.1016/j.celrep.2013.10.029.
56.   Lin, G., Xu, N., and Xi, R. (2008). Paracrine Wingless signalling controls self-renewal of
Drosophila intestinal stem cells. Nature *455*, 1119-1123. nature07329
[pii]10.1038/nature07329 [doi].
57.   Zeng, X., Chauhan, C., and Hou, S.X. (2010). Characterization of midgut stem cell- and
enteroblast-specific Gal4 lines in drosophila. Genesis *48*, 607-611. 10.1002/dvg.20661 [doi].
58.   McGuire, S.E., Mao, Z., and Davis, R.L. (2004). Spatiotemporal gene expression targeting with
the TARGET and gene-switch systems in Drosophila. Sci STKE *2004*, pl6.
59.   el Marjou, F., Janssen, K.P., Chang, B.H., Li, M., Hindie, V., Chan, L., Louvard, D., Chambon, P.,
Metzger, D., and Robine, S. (2004). Tissue-specific and inducible Cre-mediated recombination
in the gut epithelium. Genesis *39*, 186-193. 10.1002/gene.20042.
60.   Lee, T., and Luo, L. (1999). Mosaic analysis with a repressible cell marker for studies of gene
function in neuronal morphogenesis. Neuron *22*, 451-461. S0896-6273(00)80701-1 [pii].
61.   Brand, A.H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell
fates and generating dominant phenotypes. Development *118*, 401-415.
681

 Figure legends

Figure 1. A mosaic genetic screen identifies *GlcT* as a tumor suppressor in adult *Drosophila*

midgut

(A) A diagram showing lineage hierarchy of intestinal stem cells (ISCs) in the *Drosophila*

intestine. Abbreviations: EB, enteroblast; EEP, enteroendocrine cell; EC, enterocyte; EE,

enteroendocrine cell. (B-D) Compared to control clones (green), the *EA30* and *E230* clones

induced at day 5 are larger and contain a higher number of *Pros⁺* cells within each clone. *GlcT*

null allele (*GlcT^{Δ8}*) mutant clones exhibit a similar phenotype. (E-F) Expression of *UAS-GlcT* in

EA30 or *E230* mutant clones rescues the tumor phenotype. (G-J) Depletion of *GlcT* in

progenitor cells using RNAi with *esg-GAL4^{ts}* leads to ISC proliferation (G-H, stained with

Phospho-Histone 3) and excessive EEC generation in the midgut (I-J). For G-J, flies were

shifted to 29°C for 14 days. Error bars represent the mean ± SEM, with p-values indicated in

the figure (two-tailed Student's t-test). Scale bars: 25 μm.

Figure 2. Genetic analysis of the components of the GSL synthesis pathway reveals specific

tumor-suppressive activity for *egh*, in addition to *GlcT*

(A) A diagram illustrating a portion of the metabolic pathways of glycosphingolipids (GSLs)

synthesized from ceramide, highlighting the key enzymes involved at each step. (B) *GlcT^{Δ8}*

mutant clones without or with co-expression of *p35* or *CDase* stained with anti-DCP-1 or

anti-Pros, as indicated. Clones were examined at day 7-10 after clone induction. (C-D)

Quantitative analysis of the percentage of *Pros⁺* cells in clones of specified genotypes. (E)

Genetic analysis of the key enzymes in the GSL synthesis pathway. Note that *egh* mutant

clones (*egh^A*, *egh^B*, or *egh^7*) exhibited a phenotype of increased EEs similar to that observed

in *GlcT* mutant clones, but *Brn^{228}*, *β4GalNAcTA-IR*, and *α4GT1-IR* clones did not. (F, G)

Quantitative analysis of cell number (F) and the percentage of *Pros⁺* cells (G) in clones of

specified genotypes. For E-G, clones were examined at day 7 after clone induction. Error bars

represent the mean ± SEM, with p-values indicated in the figure (two-tailed Student's t-test).

Scale bars: 25 μm.

Figure 3. The EE tumor phenotype is a result of MacCer deficiency

(A) A diagram showing the major compartments along the length of the *Drosophila* midgut, including the anterior (yellow), middle (Region 3, R3, green), and posterior midgut (blue). (B-C) Feeding *GlcT* mutant flies with LacCer alleviated the overgrowth and excessive EE phenotype, which was observed in clones of both the anterior and posterior midgut. Note that the suppressive effect was more pronounced in the anterior midgut compared to the posterior midgut. Error bars represent the mean ± SEM, with p-values indicated in the figure (two-tailed Student's t-test). Scale bar: 25 µm.

Figure 4. Loss of *GlcT* leads to reduced activation of Notch signaling in ISC progenies

(A-B) The expression of *NRE-lacZ*, the Notch activity reporter, in *GlcT* mutant clones. Note that the fluorescence intensity of LacZ was significantly reduced within *GlcT* mutant clones compared to the cells outside of the clone. (C) *GlcT* mutant clones in the posterior (R5) region of the *Drosophila* midgut stained with anti-Tk (left) or anti-AstC (right). The two EE subtypes at R5 region are normally present in a 1:1 ratio. Note that AstC$^+$ EEs, but not TK$^+$ EEs, were present in *GlcT* mutant clones. (D) The effect of forced activation of Notch in *GlcT* mutant clones by expressing N^{intra}. Error bars represent the mean ± SEM, with p-values indicated in the figure (two-tailed Student's t-test). Scale bars: 25 µm.

Figure 5. *GlcT* regulates the endocytic trafficking of Delta

(A-B) Anti-Pros staining in flies carrying *GlcT-IR* driven by Dl^{ts} and NRE^{ts} respectively. Flies were shifted to 29°C for 14 days before analysis. (C) In a 3-hour Dl antibody uptake assay performed in the *Drosophila* midgut, Dl in ISCs of *GlcT* mutant clones (green box) showed reduced membrane localization and increased intracellular fluorescence compared to that in ISCs of control clones (see the enlarged inset boxes). (D, F) ISCs in *esg^{ts}>GlcT-RNAi* flies exhibited increased accumulation of Dl in early endosomes (Rab5$^+$, with co-localization

indicated by yellow arrows). (E-F) The percentage of Dl^+ late endosomes ($Rab7^+$) was largely

unchanged. For D-F, flies were shifted to 29°C for 10 days before analysis. Co-localization

analysis was performed using the intensity-based quantification feature in Leica Application

Suite X (LAS X) software, with the Overlap Coefficient (ranging from 0 to 1) employed to

quantify co-localization (1 indicating perfect co-localization and 0 indicating no

co-localization). Error bars represent the mean ± SEM, with p-values indicated in the figure

(two-tailed Student's t-test). Scale bar: 25 μm, unless otherwise noted.

Figure 6. *GlcT* shows tissue specificity in regulating Notch signaling activity

(A) The expression of Cut in a *GlcT* mutant clone (green area) in wing disc spanning the

dorsal-ventral boundary. (B) The expression of EYA in *GlcT* mutant follicle cell clones

(non-green areas). Note that there was no obvious difference in the level of EYA expression

inside and outside of the clones. (C) Punctate accumulation of Dl was observed in *GlcT*

mutant germ cell clones (non-green area, indicated by arrows in the enlarged inset), but not

in the germline of a wild-type developing egg chamber (see enlarged inset with the dashed

yellow line). Scale bars: 25 μm.

Figure 7. Transient loss of *UGCG* in mouse small intestine causes reduced number of ISCs

and increased number of goblet cells

(A-B) The expression of the Notch target gene O*lfm4* was downregulated in *villin-Cre^{ERT2}*,

UGCG^{flox/flox} mice 48 hours post-induction (n=3). The *CKO* mice displayed shorter villi (above

the dotted line) and elongated crypts (below the dotted line). (C-D) The number of $Muc2^+$

goblet cells was significantly increased in the *CKO* mice. (E) A schematic model for the role of

MacCer in regulating the Notch signaling pathway. In signaling-sending cells, mutations in

GlcT result in the loss of the downstream metabolite MacCer, leading to a rapid endocytosis

of the membrane Dl and an accumulation of Dl in early endosomes. This causes a reduced

level of Notch signaling activation in signaling-receiving cells. Error bars represent the mean

± SEM, with p-values indicated in the figure (two-tailed Student's t-test). Scale bars: 100 μm.

Supplementary Figure 1. Cross scheme for the mosaic genetic screen on the 2R chromosome.

(A) A schematic illustrating the principles of FLPase-induced mitotic recombination for generating homozygous mutant cells in heterozygous animals. (B) The cross scheme and workflow for the forward genetic mosaic screen conducted in this study. Abbreviation: EMS, ethyl methanesulfonate.

Supplementary Figure 2. The EGFR/Ras/MAPK and JAK/STAT signaling activities in *GlcT* mutant clones.

GlcT mutant clones stained with pERK (red, upper panels), a direct indicator of EGFR/Ras/MAPK signaling activity, and STAT92E (red, lower panels), whose nuclear localization is indicative of JAK/STAT signaling activity. Note that pERK was not significantly increased in *GlcT* mutant cells compared to normal cells outside of the clones, and STAT92E activity was also not obviously altered in *GlcT* mutant cells compared to normal cells outside the clones. Scale bar: 25 μm.

Supplementary Figure 3. The Crypt-Villus morphology in *UGCG* CKO mice. The duodenal structure exhibits significant abnormalities in *CKO* mice

Small intestine (duodenum) from control and *villin-creERT2, UGCG^flox/flox* mice stained with K20, which marks enterocytes and goblet cells, and Ki67, which marks proliferative cells. Note that the mutant intestine has shortened villi and elongated crypts. Dotted lines indicate the boundary between the crypt and the villus. Samples were collected at 48h after tamoxifen injection. Scale bar: 100 μm.