Flower yeasts obfuscate intrinsic individual variation in nectar sugar concentration

in a bumble bee-pollinated plant

Carlos M. Herrera [1] and Conchita Alonso

Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas (CSIC), Avda.

Américo Vespucio 26, E-41092 Sevilla, Spain

[1] Author for correspondence: email herrera@ebd.csic.es, telephone +34 95 4466700

Open research statement:

Data and metadata are available at figshare (to be submitted)

Key words/phrases: extended phenotype, *Helleborus foetidus*, individual variation,

Metschnikowia reukaufii, nectar sugar, nectar yeasts, plant-fungi interactions, pollination

Abstract

Nectar-dwelling yeasts modulate the ecology and evolution of interactions between plants and pollinators. This paper evaluates the hypothesis that yeasts can reshape intrinsic individual differences in nectar traits. We address the question, Do local genotypes of the nectar yeast specialist *Metschnikowia reukaufii* have the ability of obfuscating intrinsic individual variation among *Helleborus foetidus* plants in nectar sugar concentration ? An array of paired plant–yeast genotypes mimicking a natural field situation was created in the laboratory by inoculating nectar from different plant individuals with different yeast genotypes following a factorial design. Chemical analyses of nectar sugars before and after exposure to yeasts were performed using ion-exchange high performance liquid chromatography. Plant individual, yeast genotype, and their interaction all had strong, significant effects on sucrose concentration of *H. foetidus* nectar. Metabolic activity of yeasts abolished 79% of intrinsic variance among plants in nectar sucrose concentration, considerably reducing the opportunity for selection by pollinators on this important nectar trait. We propose that pollinator selection on nectar traits of plant x yeast combinations will supersede "naïve" selection on plants' intrinsic nectar traits, and that nectar phenotypes found by pollinators will largely represent extended phenotypes of the yeast genotype(s) living in the nectar.

Introduction

Phenotypic differences among conspecifics are a central determinant of the opportunity for selection in natural populations. In the case of plants, however, certain traits can sometimes depend more on the expression of genes belonging to intimately associated fungi than to those of the plants themselves, exemplifying instances of fungal "extended phenotypes" (Dawkins 1982; see, e.g., Dawkins 2004, Hunter 2009, 2018, Hawkes et al. 2021, for discussion and reviews). When fungal modifications of plant phenotypes involve floral features (e.g., production of

pseudoflowers), there can be fitness consequences for individual plants via interference with

pollinator behavior and plant reproduction (Batra and Batra 1985, Jennersten 1988, Jennersten

and Kwak 1991, Roy 1993, 1996, Gange and Smith 2005, González-Mas et al. 2023). Fungi

known so far to account for modifications of floral phenotypes mostly are systemic endophytes,

generally parasites or pathogens bearing antagonistic relationships with their hosts. Nevertheless,

recognition in recent years of the influence of nectar-dwelling yeasts on the interaction between

plants and pollinators through their effects on floral nectar and the pollinators themselves

(Herrera et al. 2008, 2013, Schaeffer et al. 2014, 2017, 2019, Vanette and Fukami 2016, 2018,

Rering et al. 2018, Pozo et al. 2020, de Vega et al. 2022) motivates the hitherto unexplored

hypothesis that intraspecific variation in floral nectar traits might actually reflect, to some

variable and yet-unknown degree, variation in extended phenotypes of the nectar-dwelling yeasts.

Applicability of the extended phenotype notion to nectar yeasts requires at least that the

following three premises hold true: (1) yeast genotypes differ in physiological competence and

ability to transform floral nectar, e.g., via changes in sugar composition or concentration; (2) in

natural populations of a given plant, floral nectar of different individuals is inhabited by different

yeast genotypes, so that plants might present to pollinators nectars with different features *because*

they host different yeast genotypes; and (3) differences among coexisting plants in nectar traits of

flowers exposed to natural conditions (i.e., visited by pollinators and colonized by yeasts) depend

closely on the yeast genotypes living in their flowers. Premise 1 has been documented for the

cosmopolitan nectar yeast *Metschnikowia reukaufii* whenever metabolic differences among

genotypes have been looked for (Herrera 2014, Pozo et al. 2015, Dhami et al. 2018). Premise 2 is

supported by the finding that coexisting individuals of the bumble bee-pollinated herb *Helleborus*

foetidus host distinct *M. reukaufii* genotypes in their floral nectar (Herrera et al. 2014). This

widespread floricolous yeast (Brysch-Herzberg 2004) is therefore a promising candidate for

testing the notion that nectar yeasts can profoundly reshape intrinsic patterns of individual

variation in nectar traits in local plant populations.

This paper reports an experimental test of premise 3 above for the *H. foetidus–M. reukaufii*

system. An array of paired plant–yeast genotypes mimicking a natural field situation was created

in the laboratory by inoculating nectar from different plants with cells of different yeast

genotypes. Sugar analyses of nectar samples before and after a period of yeast growth will be

used to answer the following specific question: Do local genotypes of nectar yeasts collectively

have the transformative capacity for reshaping intrinsic individual differences in nectar traits ?

Our results revealed that the identity of the nectar yeast genotypes hosted by individual plants can

be more important than intrinsic plant traits in shaping the variation in nectar traits encountered

by foraging pollinators in natural plant populations, which in turn supports the view that

pollinators might actually be interacting with sets of extended phenotypes of the nectar yeasts

living in the flowers.

Material and methods

Study system and field sampling

This study focuses on the early-blooming perennial herb *Helleborus foetidus* (Ranunculaceae)

and the dominant yeast in its floral nectar, *Metschnikowia reukaufii* (Metschnikowiaceae,

Saccharomycetales). *Helleborus foetidus* inflorescences are produced in early winter and flowers

are pollinated by bumble bees (Herrera et al. 2001). Flowers produce a sucrose-dominated nectar

(Vesprini et al. 1999, Canto et al. 2011) that ordinarily harbors dense populations of *M. reukaufii*,

the colonizing inocula of which are brought to flowers by foraging bumble bees (Brysch-

Herzberg 2004, Canto et al. 2008). Metabolic activity of these yeasts induce substantial changes

in composition and concentration of nectar sugars (Herrera et al. 2008, Canto et al. 2008, 2011).

Field sampling was conducted during February-March 2015 in a large population of *H.*

foetidus in the Sierra de Cazorla, Jaén province, southeastern Spain. Floral nectar for the

experiment was obtained from 20 widely spaced *H. foetidus* plants whose inflorescences had

been previously bagged at the bud stage to exclude pollinators. For each individual plant, nectar

from many flowers was gathered into a single bulk sample and stored at -20ºC until used. Ten *M.*

reukaufii strains from the same population were randomly chosen for the experiment from a large

collection of local strains gathered from nectar of *H. foetidus* flowers exposed to pollinators.

Experimental and analytical procedures

Prior to the experiment, nectar samples were filter-sterilized using a polyvinylidene difluoride

0.22 µm filter. Genetic distinctness of the experimental *M. reukaufii* strains was verified by

fingerprinting them using polymorphic nuclear DNA microsatellite markers (Seino et al. 2013),

and will be referred to hereafter as "genotypes".

The experimental layout consisted of a two-way factorial design, with *M. reukaufii* genotype

($N = 10$ levels) and *H. foetidus* individual ($N = 20$ levels) as main factors, and two replicates per

each genotype x individual combination ($N = 381$ experimental units in total; a few plants had

insufficient nectar for replication). Each experimental unit consisted of a 0.2 mL PCR tube

containing 15 µL of virgin, sterile nectar plus 1 µL of *M. reukaufii* cell suspension taken from a

fresh culture on standard liquid medium. To assess differences among genotypes in their initial

cell densities in the experimental units, two replicated control tubes were also set per yeast

genotype at the time of inoculation, each consisting of 15 µL of 10% lactophenol cotton blue

fixing solution (LCB hereafter) plus 1 µL of *M. reukaufii* cell suspension. After inoculation, all

experimental tubes were held at 25ºC without agitation. Two aliquots were taken from each

experimental unit for chemical analyses at 24 h and 48 h after inoculation. These rather short

periods were chosen to preclude extensive sugar depletion of nectar (Herrera et al. 2008), which

would have biased comparisons.

Samples of virgin nectar from experimental plants ($N = 40$, two aliquots per plant) and all

nectar samples taken in the course of the experiment ($N = 762$) were analyzed using ion-exchange

high performance liquid chromatography. Analytical procedures and equipment were as

described in Herrera et al. (2006) and Canto et al. (2011). Separate estimates of the concentration

of the three sugars appearing in the samples (sucrose, fructose, glucose) were obtained by

integrating areas under chromatogram peaks, and total sugar concentration was computed by

summing partial figures. As previously found for *H. foetidus* nectar in the study area (Herrera et

al. 2006, Canto et al. 2011), sucrose was virtually the only sugar in all nectar samples. It

accounted on average for 99.2% (interquartile range = 99.4–100%) and 98.9% (interquartile

range = 100–100%) of total sugars in virgin (pre-exposure to yeasts) and experimental (post-

exposure) nectar samples, respectively. Only the results for sucrose will be considered hereafter.

Data analysis

Due to differences among source cultures, initial cell densities in experimental units differed

among yeast genotypes (Chi-square = 28.7, df = 9, $P = 0.0007$, Kruskal-Wallis rank sum test;

range of genotype means = 475–1399 cells/mm^3), and experimental plants differed in the sucrose

concentration of virgin nectar ($F_{19,20} = 73.8$, $P = 1.7E\text{-}14$; plant means range = 53–77% weight-

to-weight basis). To evaluate the robustness of experimental results to this initial heterogeneity

across experimental units, and to accurately dissect the role of yeast genotypes as agents of

variation in nectar sugar across plants, two separate statistical analyses were carried out. In the

first, sucrose concentration in the nectar of experimental units after exposure to yeasts was

regressed on initial yeast cell density, initial sucrose concentration, and their interaction. In the

second analysis, a linear model was fitted to the same response variable (post-exposure sucrose

concentration) using plant identity, yeast genotype and their interaction as predictors. Including

time of sample collection (24 or 48 h) as another predictor in the analyses had a negligible,

statistically nonsignificant effect on results, and the two groups of data were pooled for the

analyses. In addition to estimating main effects and interactions in the two linear models,

"relative importance" of each predictor was estimated by decomposition of the total variance (R^2)

accounted for by each model (Groemping 2007).

Statistical analyses were carried out using the R environment (R Core Team 2024). The raw

data used here are available in Herrera (2025). Linear models were fitted using the lm function in

stats package. Adequacy of fitted models was verified using the function check_model of the

performance package (Lüdecke et al. 2021). Relative importance of predictors was obtained with

function calc.relimp in package relaimpo (Groemping 2006). Adjusted predictions (model-based

estimates) of the response variable (sucrose concentration) and marginal means were estimated

from fitted models using the predict_response function in the ggeffects package (Lüdecke 2018).

The specific "blurring" effect of each yeast genotype on individual variation in nectar sucrose

concentration was inversely assessed with the correlation coefficient across plants between initial

sucrose concentration in virgin nectar and the model-adjusted, predicted values conditioned on

the corresponding yeast genotype.

Results

Initial heterogeneity across experimental units in yeast cell density and nectar sucrose

concentration had negligible explanatory effects on variation in sucrose concentration after

experimental nectar exposure to yeasts, as revealed by both the regression parameter estimates

and the proportion of variance explained by single effects and their interaction (Table 1).

Exposure to yeasts, therefore, nearly wiped out interplant differences in sucrose concentrations in

48 h. In contrast, plant and yeast identity, and their interaction, each had strong, statistically

significant effects on nectar sucrose concentration after exposure to yeasts (Table 1). The

proportion of variance explained by this model was remarkably high ($R^2 = 68.1\%$) given the large

number of samples. The estimated contribution of main effects (plant and yeast identity) and their

interaction to the variance explained by the model declined in the direction plant (32%) – plant x

yeast (20%) – yeast (15%) (Table 1). Model-adjusted, predicted values of nectar sucrose

concentration after exposure to yeasts varied among plants (range = 31–53%, conditioned on

strain 1_A_10.2) and yeast strains (range = 40–62%, conditioned on plant CV01), but the strong

plant x yeast interaction effect renders these figures only weakly informative (Appendix S1:

Figure S1).

The high explanatory value of the yeast x plant interaction was of particular significance in

relation to the hypothesis tested here. First, the relative effect of different yeast genotypes on

nectar sucrose concentration was remarkably different for different individual plants, as

illustrated by extensive criss-crossing of lines in the interaction plot of within-plant standardized

values (Figure 1). Second, as a consequence of the strong interaction, the set of paired

combinations plant x yeast genotype created a heterogeneous two-dimensional space of post-

exposure sucrose concentrations (Appendix S1: Figure S1). And third, yeast genotypes differed

widely in their capacity to blur individual variation in nectar sucrose concentration, as shown by

the broad range across plants of the correlation coefficient between mean sucrose concentration

before and after the experiment (r range = 0.142–0.736, mean = 0.454). The metabolic activity of

the yeast genotypes tested obliterated between 46–98% (mean = 79%) of intrinsic variance

among plants in nectar sucrose concentration over the experimental period (estimated as $1 - r^2$).

Discussion

Local *M. reukaufii* genotypes blurred intrinsic differences among individuals of *H. foetidus* in

nectar sucrose concentration, a trait potentially important for the relationship with pollinators.

Our results support premise 3 in the Introduction by showing that, after a relatively short

exposure to the metabolic activity of *M. reukaufii* genotypes, initial sucrose concentrations were

obfuscated, with plant and yeast identity, and their interaction acquiring overwhelming

explanatory power. These experimental results, however, may underestimate the actual impact of

yeasts on individual variation in sugar concentration in the field. On one side, initial yeast cell

densities in experimental units were lower than the densities commonly found in nectar of *H.*

foetidus flowers exposed to pollinators (Herrera et al. 2008, 2009). And on the other, length of

exposure to yeasts in this study was much shorter than natural lifetime of individual *H. foetidus*

flowers (up to 20 d; Herrera et al. 2001). Since sucrose concentration of *H. foetidus* nectar

declines as yeast cell density rises with flower age (Herrera et al. 2008), effects in the field could

perhaps be larger than those reported here.

Floral nectar is a major pollinator reward that strongly influences plant-pollinator

interactions, and nectar features have been mostly interpreted in the light of selection from

pollinators (Nicolson et al. 2007, Abrahamczyk et al. 2017). More recently, however, the

potential role of nectar microorganisms in nectar evolution has begun to be recognized, but our

understanding of their actual role is still in an incipient stage (Canto and Herrera 2012,

Parachnowitsch et al. 2019, Barberis et al. 2024). Results of this study have revealed an hitherto

unrecognized role of specialized nectar yeasts in nectar trait evolution: by reshaping intrinsic

individual differences among plants, *M. reukaufii* yeasts will reduce the opportunity for pollinator

selection on *H. foetidus* nectar sugar concentration. Such reduction will hinder selection on the

plants' intrinsic values of nectar sugar content, as their variation will have been "masked" by the

transformative action of the yeast genotypes on nectar. Such reduction in opportunity for

selection, however, will be in a sense compensated by new opportunities on an emergent axis.

The variation among yeast genotypes in sugar transformative capacity, plus the strong plant x

yeast genotype interaction effect on sugar concentration, will open opportunities for selection on

sugar concentration over the two-dimensional space defined by the combinations of plant x yeast

genotypes (Appendix S1: Figure S1). In this proposed scenario, pollinator selection on nectar

traits of plant x yeast combinations will supersede "naïve" pollinator selection on intrinsic nectar

traits of individual plants, since the plant nectar phenotypes encountered by pollinators will

largely be extended phenotypes of the yeast genotype(s) within the nectar. Nectar features could

thus be considered as extended phenotypes of yeasts insofar as they are "candidate adaptations

for the benefit of alleles responsible for variation in them" (Dawkins 2004), as discussed below.

Since initial conditions (sucrose concentration, yeast cell density) had a negligible effect on

our experiment results, it seems safe to conclude that variation among experimental units in

sucrose concentration after exposure to yeasts mainly arose from differences among yeast

genotypes in growth rate and metabolic competence to exploit not just sucrose, but also other

influential nectar constituents such as monosaccharides, proteins, amino acids and lipids which

also occur in the nectar of *H. foetidus* but were not measured here (Herrera 1989, Vesprini et al.

1999). The concentration of these other nectar constituents probably varied among plants, and in

our experiment their collective, unmeasured effects are subsumed under the "plant identity"

factor. Yeast genotypes with metabolic profiles variably suited to the chemical profile of a given

plant's nectar will probably grow and modify sucrose concentration at different rates (Herrera

2014). All else being equal, therefore, bumble bee discrimination for nectars with different

sucrose concentrations (Cnaani et al. 2006, Whitney et al. 2008, Pozo et al. 2020) will translate

into selection among plant-yeast combinations which have different sucrose concentrations (i.e.,

among different tiles in Appendix S1: Figure S1). As bumble bees simultaneously disperse yeast

propagules and pollen grains, any positive selection on a particular plant-yeast combination

arising from pollinator discrimination will automatically benefit both the plant and the yeast

genotype(s) involved, contribute to consolidate their association, increase its frequency, and

perhaps eventually become "stabilized" within *H. foetidus* populations over the blooming season.

In the long run, differential dispersal success of yeast genotypes associated with different plant

individuals, and the likely stabilization of plant-yeast genotypic pairs postulated here, could

eventually contribute to the genetic differentiation of *M. reukaufii* subpopulations found in

flowers of *H. foetidus* individuals growing side by side, as well as the remarkable genetic and

phenotypic diversity that characterize this yeast species (Herrera 2014, Herrera et al. 2014, Pozo

et al. 2015, Dhami et al. 2018).

Acknowledgments

We are grateful to Pilar Bazaga, Marina García and Esmeralda López-Perea for laboratory

assistance; Ricardo Pérez for the development and implementation of sugar analytical

procedures; and Consejería de Medio Ambiente, Junta de Andalucía por permission to work in

the Sierra de Cazorla and providing invaluable facilities there. Different stages of this work were

supported by grants CGL2010-15964 (Ministerio de Educación y Ciencia), P09-RNM-4517

(Junta de Andalucía), and PID2022-141530NB-C22 (Ministerio de Ciencia e Innovación).

Author contributions

250   C. M. H. conceptualized the study, performed field sampling, set up the experiment, analyzed the

251   data and led writing. C. A. set up and conducted sugar analytical work. Both authors contributed

252   to data curation, edited and revised manuscript drafts and approved the final version.

Conflict of interest statement

254   Authors declare no conflict of interest.

Data availability statement

256   Data and metadata are available at figshare (to be submitted)

ORCID

Carlos M. Herrera https://orcid.org/0000-0003-2452-3888

Conchita Alonso https://orcid.org/0000-0002-7418-3204

Literature cited

Abrahamczyk, S., M. Kessler, D. Hanley, D. N. Karger, M. P. J. Muller, A. C. Knauer, F. Keller,

M. Schwerdtfeger, and A. M. Humphreys. 2017. Pollinator adaptation and the evolution of

floral nectar sugar composition. Journal of Evolutionary Biology 30:112–127.

Barberis, M., M. Nepi, and M. Galloni. 2024. Floral nectar: fifty years of new ecological

perspectives beyond pollinator reward. Perspectives in Plant Ecology, Evolution and

Systematics 62:125764.

Batra, L. R., and S. W. T. Batra. 1985. Floral mimicry induced by mummy-berry fungus exploits

host's pollinators as vectors. Science 228:1011–1013.

Brysch-Herzberg, M. 2004. Ecology of yeasts in plant-bumblebee mutualism in Central Europe.

FEMS Microbiology Ecology 50:87–100.

Canto, A., and C. M. Herrera. 2012. Micro-organisms behind the pollination scenes: microbial

imprint on floral nectar sugar variation in a tropical plant community. Annals of Botany

110:1173–1183.

Canto, A., C. M. Herrera, M. Medrano, R. Pérez, and I. M. García. 2008. Pollinator foraging

modifies nectar sugar composition in *Helleborus foetidus* L. (Ranunculaceae): an experimental

test. American Journal of Botany 95:315–320.

Canto, A., C. M. Herrera, I. M. García, R. Pérez, and M. Vaz. 2011. Intraplant variation in nectar

traits in *Helleborus foetidus* (Ranunculaceae) as related to floral phase, environmental

conditions and pollinator exposure. Flora 206:668–675.

Cnaani, J., J. D. Thomson, and D. R. Papaj. 2006. Flower choice and learning in foraging

bumblebees: effects of variation in nectar volume and concentration. Ethology 112:278–285.

Dhami, M. K., T. Hartwig, A. D. Letten, M. Banf, and T. Fukami. 2018. Genomic diversity of a

nectar yeast clusters into metabolically, but not geographically, distinct lineages. Molecular

Ecology 27:2067–2076.

Dawkins, R. 1982. The extended phenotype: the long reach of the gene. Oxford University Press,

Oxford, UK.

Dawkins, R. 2004. Extended phenotype – But not too extended. A reply to Laland, Turner and

Jablonka. Biology and Philosophy 19:377–396.

de Vega, C., R. G. Albaladejo, S. Álvarez-Pérez, and C. M. Herrera. 2022. Contrasting effects of

nectar yeasts on the reproduction of mediterranean plant species. American Journal of Botany

109:393–405.

Gange, A. C., and A. K. Smith. 2005. Arbuscular mycorrhizal fungi influence visitation rates of

pollinating insects. Ecological Entomology 30:600–606.

González-Mas, N., M. Cuenca-Medina, H. García-Mozo, J. M. Muñoz-Redondo, J. M. Moreno-

Rojas, F. Padilla-Álvarez, I. M. Rodríguez-Gómez, and E. Quesada-Moraga. 2023.

Endophytic *Beauveria bassiana* modifies flowering phenology, floral volatile profile and

pollinator behaviour in melon. Entomologia Generalis 43:961–969.

Groemping, U. 2006. Relative importance for linear regression in R: the package relaimpo.

Journal of Statistical Software 17:1–27.

Groemping, U. 2007. Estimators of relative importance in linear regression based on variance

decomposition. American Statistician 61:139–147.

Hawkes, C. V., R. Kjøller, J. M. Raaijmakers, L. Riber, S. Christensen, S. Rasmussen, J. H.

Christensen, A. B. Dahl, J. C. Westergaard, and M. Nielsen. 2021. Extension of plant

phenotypes by the foliar microbiome. Annual Review of Plant Biology 72:823–846.

Herrera, C. M. 2014. Population growth of the floricolous yeast *Metschnikowia reukaufii*: effects

of nectar host, yeast genotype and host x genotype interaction. FEMS Microbiology Ecology

88:250–257.

Herrera, C. M. 2025. Effects of yeast genotype and plant identity on sucrose concentration

of *Helleborus foetidus* nectar. Figshare. Dataset. To be submitted

Herrera, C. M., A. M. Sánchez-Lafuente, M. Medrano, J. Guitián, X. Cerdá, and P. Rey. 2001.

Geographical variation in autonomous self-pollination levels unrelated to pollinator service in

Helleborus foetidus (Ranunculaceae). American Journal of Botany 88:1025–1032.

Herrera, C. M., R. Pérez, and C. Alonso. 2006. Extreme intraplant variation in nectar sugar

composition in an insect-pollinated perennial herb. American Journal of Botany 93:575–581.

Herrera, C. M., I. M. García, and R. Pérez. 2008. Invisible floral larcenies: microbial

communities degrade floral nectar of bumblebee-pollinated plants. Ecology 89:2369–2376.

Herrera, C. M., C. de Vega, A. Canto, and M. I. Pozo. 2009. Yeasts in floral nectar: a quantitative

survey. Annals of Botany 103:1415–1423.

Herrera, C. M., M. I. Pozo, and M. Medrano. 2013. Yeasts in nectar of an early-blooming herb:

sought by bumble bees, detrimental to plant fecundity. Ecology 94:273–279.

Herrera, C. M., M. I. Pozo, and P. Bazaga. 2014. Nonrandom genotype distribution among floral

hosts contributes to local and regional genetic diversity in the nectar living yeast

Metschnikowia reukaufii. FEMS Microbiology Ecology 87:568–575.

Herrera, J. 1989. Aminoácidos en el néctar de plantas del sur de España. Anales del Jardín

Botánico de Madrid 45:475–482.

Hunter, P. 2009. Extended phenotype redux. EMBO Reports 10:212–215.

Hunter, P. 2018. The revival of the extended phenotype. EMBO Reports 19:e46477.

Jennersten, O. 1988. Insect dispersal of fungal disease: effects of *Ustilago infection* on pollinator

attraction in *Viscaria vulgaris*. Oikos 51:163–170.

Jennersten, O., and M. M. Kwak. 1991. Competition for bumblebee visitation between

Melampyrum pratense and *Viscaria vulgaris* with healthy and *Ustilago*-infected flowers.

Oecologia 86:88–98.

Lüdecke, D. 2018. ggeffects: Tidy data frames of marginal effects from regression models.

Journal of Open Source Software 3:772.

Lüdecke, D., M. S. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021. performance:

An R package for assessment, comparison and testing of statistical models. Journal of Open

Source Software 6:3139.

Nicolson, S. W., M. Nepi, and W. Pacini, editors. 2007. Nectaries and nectar. Springer-Verlag,

Dordrecht, The Netherlands.

Parachnowitsch, A. L., J. S. Manson, and N. Sletvold. 2019. Evolutionary ecology of nectar.

Annals of Botany 123:247–261.

Pozo, M. I., C. M. Herrera, W. van den Ende, K. Verstrepen, B. Lievens, and H. Jacquemyn.

2015. The impact of nectar chemical features on phenotypic variation in two related nectar

yeasts. FEMS Microbiology Ecology 91:fiv055.

Pozo, M. I., G. van Kemenade, A. van Oystaeyen, T. Aledón-Catalá, A. Benavente, W. Van den

Ende, F. Wäckers, and H. Jacquemyn. 2020. The impact of yeast presence in nectar on bumble

bee behavior and fitness. Ecological Monographs 90:e01393.

R Core Team. 2024. R: A language and environment for statistical computing. Vienna, Austria: R

Foundation for Statistical Computing.

Rering, C. C., J. J. Beck, G. W. Hall, M. M. McCartney, and R. L. Vannette. 2018. Nectar-

inhabiting microorganisms influence nectar volatile composition and attractiveness to a

generalist pollinator. New Phytologist 220:750–759.

Roy, B. A. 1993. Floral mimicry by a plant pathogen. Nature 362:56–58.

Roy, B. A. 1996. A plan pathogen influences pollinator behavior and may influence reproduction

of nonhosts. Ecology 77:2445–2457.

Schaeffer, R. N., C. R. Phillips, M. C. Duryea, J. Andicoechea, and R. E. Irwin. 2014. Nectar

yeasts in the tall larkspur *Delphinium barbeyi* (Ranunculaceae) and effects on components of

pollinator foraging behavior. PLoS ONE 9:e108214.

Schaeffer, R. N., Y. Z. Mei, J. Andicoechea, J. S. Manson, and R. E. Irwin. 2017. Consequences

of a nectar yeast for pollinator preference and performance. Functional Ecology 31:613–621.

Schaeffer, R. N., C. C. Rering, I. Maalouf, J. J. Beck, and R. L. Vannette. 2019. Microbial

metabolites elicit distinct olfactory and gustatory preferences in bumblebees. Biology Letters

15:20190132.

Seino, M. M., C. de Vega, P. Bazaga, M. I. Pozo, and C. M. Herrera. 2013. Development and

characterization of microsatellite loci for the nectar-living yeast *Metschnikowia reukaufii*.

Pages 760-762 in Molecular Ecology Resources Primer Development Consortium. *Molecular*

Ecology Resources 13:760–762.

Vannette, R. L., and T. Fukami. 2016. Nectar microbes can reduce secondary metabolites in

nectar and alter effects on nectar consumption by pollinators. Ecology 97:1410–1419.

Vannette, R. L., and T. Fukami. 2018. Contrasting effects of yeasts and bacteria on floral nectar

traits. Annals of Botany 121:1343–1349.

Vesprini, J. L., M. Nepi, and E. Pacini. 1999. Nectary structure, nectar secretion patterns and

nectar composition in two *Helleborus* species. Plant Biology 1:560–568.

Whitney, H. M., A. Dyer, L. Chittka, S. A. Rands, and B. J. Glover. 2008. The interaction of

temperature and sucrose concentration on foraging preferences in bumblebees.

Naturwissenschaften 95:845–850.

Table 1. Summary of statistical analyses testing for effects on nectar sucrose concentration after experimental exposure of nectar from 20 different plants to 10 yeast genotypes using a factorial design. Separate models were fitted to the data ($N = 786$ sugar concentration measurements) in which the predictors were initial yeast cell density and sucrose concentration, and plant and yeast genotype identities, respectively.

Predictors in model	Effect	Parameter estimate (SE)	Test statistic	P-value	Variance explained (%)
Initial yeast and nectar features:			Student's t		
	Cell density (CD)	-0.0117 (0.009)	1.35	0.18	2.1
	Sucrose concentration (SC)	0.289 (0.13)	2.26	0.024	9.1
	CD x SC	0.0001 (0.001)	0.84	0.39	0.084
Plant and yeast identity:			F value (df)		
	Plant (P)		31.6 (19, 586)	< 2E-16	32.5
	Yeast genotype (YG)		31.7 (9, 586)	< 2E-16	15.3
	YG x P		2.2 (171,586)	7.9E-12	20.3

Caption to the figure

Figure 1. Interaction plot depicting differences in sucrose concentration after experimental exposure of nectar from different plants to different yeast genotypes. To emphasize the contrasting effect of the different yeast genotypes on different individual plants, and to facilitate comparisons across plants, model-predicted sucrose concentrations for each plant x yeast genotype combination were scaled and centered separately for each plant. See Appendix S1: Figure S1, for the distribution of model-predicted values of sucrose concentration over the two-dimensional space defined by the 20 plants and 10 yeast genotypes tested in the experiment.

Figure 1