Flowers as fungal extended phenotypes: nectar yeasts obfuscate among-plant differences

in nectar sugar concentration

Carlos M. Herrera [1] and Conchita Alonso

Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas (CSIC),

Avda. Américo Vespucio 26, E-41092 Sevilla, Spain

ORCID:

Carlos M. Herrera https://orcid.org/0000-0003-2452-3888

Conchita Alonso https://orcid.org/0000-0002-7418-3204

[1] Author for correspondence: email herrera@ebd.csic.es

Abstract

Nectar-dwelling yeasts modulate the ecology of interactions between flowers and pollinators.

We evaluate here the hypothesis that floral nectar traits, which are a central element in most

plant-pollinator relationships, can largely represent the extended fungal phenotypes of nectar-

dwelling yeasts. The following specific question is addressed, Do local genotypes of the

specialist nectar yeast *Metschnikowia reukaufii* have the ability to obfuscate intrinsic

individual variation among *Helleborus foetidus* plants in nectar sugar concentration ? An

array of paired plant-yeast genotypes mimicking a natural field situation was created in the

laboratory by inoculating field-collected nectar from different plant individuals with distinct

yeast genotypes following a factorial design. Chemical analyses of nectar sugars before and

after exposure to yeasts were performed using ion-exchange high performance liquid

chromatography. Plant individual, yeast genotype, and their interaction all had strong,

significant effects on sucrose concentration of *H. foetidus* nectar. Yeasts abolished 79% of

intrinsic variance among plants in nectar sucrose concentration and profoundly reshaped

patterns of intrinsic among-plant variation. Our results support the hypothesis that among-

plant variation in nectar sugar concentration found by pollinators in the field can sometimes

reflect the extended phenotypes of nectar yeasts more closely than intrinsic differences among

the plants themselves.

Introduction

Phenotypic differences among conspecifics are a central determinant of the opportunity for

selection in natural populations. In the case of plants, however, certain traits can sometimes

depend more on the expression of genes belonging to intimately associated fungi than to those

of the plants themselves, exemplifying instances of fungal "extended phenotypes" (Dawkins

1982; see, e.g., Dawkins 2004, Hunter 2009, 2018, Hawkes et al. 2021, for discussion and

reviews). When fungal modifications of plant phenotypes involve floral features there can be

fitness consequences for the individual plants via interference with pollinator behavior and

plant reproduction (Batra and Batra 1985, Jennersten 1988, Jennersten and Kwak 1991, Roy

1993, 1996, Gange and Smith 2005, González-Mas et al. 2023). Fungi known so far to

account for modifications of floral phenotypes mostly are systemic endophytes, generally

parasites or pathogens bearing antagonistic relationships with their hosts. Nevertheless,

recognition in recent years of the influence of nectar-dwelling yeasts on the interaction

between plants and pollinators through their effects on floral nectar and the pollinators

themselves (Herrera et al. 2008, 2013, Schaeffer et al. 2014, 2017, 2019, Vanette and Fukami

2016, 2018, Rering et al. 2018, Pozo et al. 2020, de Vega et al. 2022) motivates the hitherto

unexplored hypothesis that intraspecific variation in nectar traits naturally occurring in the

field might actually reflect the extended phenotypes of nectar-dwelling yeasts.

This paper reports results of an experimental test of this hypothesis for the interaction

between plants of *Helleborus foetidus* (Ranunculaceae) and the nectar yeast *Metschnikowia*

reukaufii (Metschnikowiaceae, Saccharomycetales). An array of paired plant-yeast genotypes

mimicking a natural field situation was created in the laboratory by inoculating nectar from

different plant individuals with cells of different yeast genotypes. Sugar analyses of nectar

samples before and after a period of yeast population growth will be used to answer the

following specific question: Do local genotypes of the yeast have the ability to obfuscate

intrinsic differences among plants in nectar sugar concentration ? Our results revealed that the

identity of the nectar yeast genotypes hosted by individual plants can become more important

than intrinsic plant traits in shaping the variation in nectar traits encountered by foraging

pollinators in natural plant populations, which in turn supports the hypothesis that pollinators

might actually be interacting with sets of extended phenotypes of the nectar yeasts living in

the flowers.

Material and methods

Study system and field sampling

This study focuses on the early-blooming perennial herb *Helleborus foetidus* and the dominant yeast in its floral nectar, *Metschnikowia reukaufii*. *Helleborus foetidus* inflorescences are produced in early winter and flowers are pollinated by bumble bees (Herrera et al. 2001). Flowers produce a sucrose-dominated nectar (Vesprini et al. 1999, Canto et al. 2011) that ordinarily harbors dense populations of *M. reukaufii*, the colonizing inocula of which are brought to flowers by foraging bumble bees (Brysch-Herzberg 2004, Canto et al. 2008). Metabolic activity of these yeasts induce substantial changes in composition and concentration of nectar sugars (Herrera et al. 2008, Canto et al. 2008, 2011).

Field sampling was conducted during February-March 2015 in a large population of *H. foetidus* in the Sierra de Cazorla, Jaén province, southeastern Spain. Floral nectar for the experiment was obtained from 20 widely spaced *H. foetidus* plants whose inflorescences had been previously bagged at the bud stage to exclude pollinators. For each individual plant, nectar from many flowers was gathered into a single bulk sample and stored at -20ºC until used. Ten *M. reukaufii* strains from the same population were randomly chosen for the experiment from a large collection of local strains gathered from nectar of *H. foetidus* flowers exposed to pollinators.

Experimental and analytical procedures

Prior to the experiment, nectar samples were filter-sterilized using a polyvinylidene difluoride 0.22 μm filter. Genetic distinctness of the experimental *M. reukaufii* strains was verified by fingerprinting them using polymorphic nuclear DNA microsatellite markers (Seino et al. 2013), and will be referred to hereafter as "genotypes".

The experimental layout consisted of a two-way factorial design, with *M. reukaufii* genotype ($N = 10$ levels) and *H. foetidus* individual ($N = 20$ levels) as main factors, and two

replicates per each genotype x individual combination ($N = 381$ experimental units in total; a

few plants had insufficient nectar for replication). Each experimental unit consisted of a

0.2 mL PCR tube containing 15 µL of virgin, sterile nectar plus 1 µL of *M. reukaufii* cell

suspension taken from a fresh culture on standard liquid medium. To assess differences

among yeast genotypes in initial cell density per experimental unit, two replicated control

tubes were also set per genotype at the time of inoculation, each consisting of 15 µL of 10%

lactophenol cotton blue fixing solution (LCB hereafter) plus 1 µL of *M. reukaufii* cell

suspension. After inoculation, all experimental tubes were held at 25ºC without agitation.

Aliquots for chemical analyses were taken from each experimental unit at 24 h and 48 h after

inoculation. These short periods were chosen to impede extensive sugar depletion of nectar

(Herrera et al. 2008), which would have biased comparisons.

Samples of virgin nectar from experimental plants ($N = 40$, two aliquots per plant) and all

nectar samples taken in the course of the experiment ($N = 762$) were analyzed using ion-

exchange high performance liquid chromatography. Analytical procedures and equipment

were as described in Herrera et al. (2006) and Canto et al. (2011). Separate estimates of the

concentration of the three sugars appearing in the samples (sucrose, fructose, glucose) were

obtained by integrating areas under chromatogram peaks, and total sugar concentration was

computed by summing partial figures. As previously found for *H. foetidus* nectar in the study

area (Herrera et al. 2006, Canto et al. 2011), sucrose was virtually the only sugar in all nectar

samples. It accounted on average for 99.2% (interquartile range = 99.4–100%) and 98.9%

(interquartile range = 100–100%) of total sugars in virgin (pre-exposure to yeasts) and

experimental (post-exposure) nectar samples, respectively. Only results for sucrose will be

considered hereafter.

Data analysis

Due to differences among source cultures, initial cell densities in experimental units differed

among yeast genotypes (Chi-square = 28.7, df = 9, P = 0.0007, Kruskal-Wallis rank sum test;

range of genotype means = 475–1399 cells/mm^3), and experimental plants differed in the

sucrose concentration of virgin nectar ($F_{19,20}$ = 73.8, P = 1.7E-14; plant means range = 53–

77% weight-to-weight basis). To evaluate robustness of experimental results to this initial

heterogeneity across experimental units, and to accurately dissect the role of yeast genotypes

as agents of variation in nectar sugar across plants, two separate statistical analyses were

carried out. In the first one, sucrose concentration in nectar of experimental units after

exposure to yeasts was regressed on initial yeast cell density, initial sucrose concentration,

and their interaction. In the second analysis, a linear model was fitted to post-exposure

sucrose concentration using plant identity, yeast genotype and their interaction as predictors.

Adding time of sample collection (24 h or 48 h) as a predictor had a negligible, statistically

nonsignificant effect on results, and the two data sets were pooled for the analyses. In addition

to estimating main effects and interactions in the two linear models, "relative importance" of

each predictor was estimated by decomposition of the total variance (R^2) accounted for by

each model (Groemping 2007).

Statistical analyses were carried out using the R environment (R Core Team 2024). The

raw data used here are available in Herrera (2025). Linear models were fitted using the lm

function in stats package. Adequacy of fitted models was verified using the function

check_model of the performance package (Lüdecke et al. 2021). Relative importance of

predictors was obtained with function calc.relimp in package relaimpo (Groemping 2006).

Marginal means and adjusted predictions (model-based estimates) of the response variable

(sucrose concentration) were estimated from fitted models using the predict_response

function in the ggeffects package (Lüdecke 2018). The average "blurring" effect of each yeast

genotype on intrinsic among-plant variation in nectar sucrose concentration was inversely

assessed with the correlation coefficient across plants between initial sucrose concentration in

virgin nectar and the model-adjusted, predicted values of post-exposure concentration

conditioned on the yeast genotype.

Results

Exposure to yeasts largely wiped out initial interplant differences in sucrose concentrations.

Initial heterogeneity across experimental units in yeast cell density and nectar sucrose

concentration had negligible explanatory effects on variation in sucrose concentration after

experimental nectar exposure to yeasts, as revealed by both the regression parameter estimates

and the proportion of variance explained by single effects and their interaction (Table 1).

Plant and yeast identity, and their interaction, each had strong, statistically significant

effects on nectar sucrose concentration after exposure to yeasts (Table 1). The proportion of

variance explained by this model was remarkably high ($R^2 = 68.1\%$) given the large number

of samples. Estimated contributions of main effects (plant and yeast identity) and their

interaction to the variance explained by the model declined in the direction plant (32%) –

plant x yeast (20%) – yeast (15%) (Table 1). Model-adjusted, predicted values of nectar

sucrose concentration after exposure to yeasts varied among plants (range = 31–53%,

conditioned on strain 1_A_10.2) and yeast strains (range = 40–62%, conditioned on plant

CV01), but the strong plant x yeast interaction effect renders these figures only weakly

informative (Figure 2).

The high explanatory value of the yeast x plant interaction deserves particular

consideration in the context of the hypothesis examined here. First, the comparative effect of

different yeast genotypes on nectar sucrose concentration was remarkably different for

different individual plants, as illustrated by extensive criss-crossing of lines in the interaction

plot of within-plant standardized values (Figure 1). Second, as a consequence of the strong

interaction the set of paired combinations plant x yeast genotype created a heterogeneous two-

dimensional space of post-exposure sucrose concentrations (Figure 2). And third, yeast

genotypes differed widely in their capacity to blur individual variation in nectar sucrose

concentration, as shown by the broad range across plants of the correlation coefficient

between mean sucrose concentration before and after the experiment (r range = 0.142–0.736,

mean = 0.454). The activity of yeast genotypes obliterated between 46–98% (mean = 79%) of

intrinsic variance among plants in nectar sucrose concentration (estimated as $1 - r^2$).

Discussion

Local *M. reukaufii* genotypes obfuscated intrinsic differences among individuals of *H.*

foetidus in nectar sucrose concentration, a trait potentially important for the relationship with

pollinators. After a relatively short exposure to the metabolic activity of *M. reukaufii*

genotypes, among-plant differences in initial sucrose concentrations were nearly wiped out,

while yeast identity and its interaction with plant identity acquired a major explanatory power.

It must be stressed, however, that these experimental results probably underestimate the actual

impact of yeasts on individual variation in sugar concentration in the field. On one side, initial

yeast cell densities in experimental units were lower than the densities commonly found in

nectar of *H. foetidus* flowers exposed to pollinators (Herrera et al. 2008, 2009). And on the

other, length of exposure to yeasts in this study was shorter than the natural lifetime of

individual *H. foetidus* flowers (up to 20 d; Herrera et al. 2001). Since sucrose concentration of

H. foetidus nectar declines as yeast cell density rises with flower age (Herrera et al. 2008),

yeast genotype effects in the field are probably larger than those reported here.

Applicability of the extended phenotype notion to nectar yeasts requires that the

following three premises hold true: (1) yeast genotypes differ in physiological competence

and ability to transform floral nectar, e.g., via changes in sugar composition or concentration;

(2) in natural populations of a given plant, floral nectar of different individuals is inhabited by

different yeast genotypes, so that plants might present to pollinators nectars with different

features *because* they host different yeast genotypes; and (3) differences among coexisting

plants in nectar traits of flowers exposed to natural conditions (i.e., visited by pollinators and

colonized by yeasts) depend closely on the yeast genotypes living in their flowers. Premise 1

has been previously documented for *M. reukaufii* whenever metabolic differences among

genotypes have been looked for (Herrera 2014, Pozo et al. 2015, Dhami et al. 2018), and

premise 2 was supported by the finding that coexisting individuals of *H. foetidus* host distinct

M. reukaufii genotypes in their floral nectar (Herrera et al. 2014). Premise 3 is supported by

the results of the present study. Past and present results, taken together, thus provide

compelling support for the hypothesis that natural variation in sugar concentration occurring

in populations of *H. foetidus* can reflect the extended phenotypes of the nectar-dwelling yeast

M. reukaufii. The potential implications of this finding for the interaction between *H. foetidus*

and its pollinators are briefly discussed below.

Floral nectar is a major pollinator reward that strongly influences plant-pollinator

interactions, and nectar features have been mostly interpreted in the light of selection from

pollinators (Nicolson et al. 2007, Abrahamczyk et al. 2017). More recently, the potential role

of nectar microorganisms in nectar evolution has begun to be acknowledged, although our

understanding of their actual role is still in an incipient stage (Parachnowitsch et al. 2019,

Barberis et al. 2024). Results of this study have revealed an hitherto unrecognized role of

specialized nectar yeasts in nectar trait evolution: by reshaping intrinsic individual differences

among plants, *M. reukaufii* yeasts will reduce the opportunity for pollinator selection on *H.*

foetidus nectar sugar concentration. Such reduction will hinder selection on the plants'

intrinsic values of nectar sugar content, insofar as that variation is overshadowed by the

transformative action of the yeast genotypes on nectar. Such reduction in opportunity for

selection, however, will be in a sense compensated by new opportunities on an emergent axis.

The variation among yeast genotypes in sugar transformative capacity, plus the strong plant x

yeast genotype interaction effect on sugar concentration, will create opportunities for

selection on sugar concentration over the two-dimensional space defined by combinations of

plant x yeast genotypes (Figure 2). In this proposed scenario, pollinator selection on nectar

traits of plant x yeast combinations will supersede "naive" pollinator selection on intrinsic

nectar traits of individual plants, since the plant nectar phenotypes encountered by pollinators

in the field will often be extended phenotypes of the yeast genotype(s) within the nectar.

Nectar features could thus be considered as extended phenotypes of yeasts insofar as they are

"candidate adaptations for the benefit of alleles responsible for variation in them" (Dawkins

2004), as discussed below.

Since initial conditions (sucrose concentration, yeast cell density) had a negligible effect

on our experiment results, it seems safe to conclude that variation among experimental units

in sucrose concentration after exposure to yeasts mainly arose from differences among yeast

genotypes in growth rate and metabolic competence to exploit not just sucrose, but also other

influential nectar constituents such as monosaccharides, proteins, amino acids and lipids

which are also present in the nectar of *H. foetidus* but were not measured here (Herrera 1989,

Vesprini et al. 1999). The concentration of these other nectar constituents most likely varied

among plants, and in our experiment their collective, unmeasured effects are subsumed under

the "plant identity" factor. Yeast genotypes with metabolic profiles variably suited to the

chemical profile of a given plant's nectar will probably grow and modify sucrose

concentration at different rates (Herrera 2014). All else being equal, therefore, bumble bee

discrimination for nectars with different sucrose concentrations (Cnaani et al. 2006, Whitney

et al. 2008, Pozo et al. 2020) will translate into selection among plant-yeast combinations

which have different sucrose concentrations (i.e., among different tiles in Figure 2). As

bumble bees disperse yeast propagules and pollen grains simultaneously, any positive

selection on a particular plant-yeast combination arising from pollinator discrimination will

automatically benefit both the plant and the yeast genotype(s) involved, contribute to

consolidate their association, increase its frequency, and perhaps eventually become

"stabilized" within *H. foetidus* populations over the blooming season. In the long run,

differential dispersal success of yeast genotypes associated with different plant individuals,

and the likely stabilization of plant-yeast genotypic pairs postulated here, could eventually

contribute to the genetic differentiation of *M. reukaufii* subpopulations found in flowers of *H.*

foetidus individuals growing side by side, as well as the remarkable genetic and phenotypic

diversity that characterize this yeast species (Herrera 2014, Herrera et al. 2014, Pozo et al.

2015, Dhami et al. 2018).

Acknowledgments

We are grateful to Pilar Bazaga, Marina García and Esmeralda López-Perea for laboratory

assistance; Ricardo Pérez for the development and implementation of sugar analytical

procedures; and Consejería de Medio Ambiente, Junta de Andalucía por permission to work

in the Sierra de Cazorla and providing invaluable facilities there. Different stages of this work

were supported by grants CGL2010-15964 (Ministerio de Educación y Ciencia), P09-RNM-

4517 (Junta de Andalucía), and PID2022-141530NB-C22 (Ministerio de Ciencia e

Innovación).

Data availability statement

Data, metadata and R script for analyses are available at figshare

(https://figshare.com/s/2e27d89d49dc46e4adb1) (Herrera 2025)

Literature cited

Abrahamczyk, S., M. Kessler, D. Hanley, D. N. Karger, M. P. J. Muller, A. C. Knauer, F.

Keller, M. Schwerdtfeger, and A. M. Humphreys. 2017. Pollinator adaptation and the

evolution of floral nectar sugar composition. Journal of Evolutionary Biology 30:112–127.

Barberis, M., M. Nepi, and M. Galloni. 2024. Floral nectar: fifty years of new ecological

perspectives beyond pollinator reward. Perspectives in Plant Ecology, Evolution and

Systematics 62:125764.

Batra, L. R., and S. W. T. Batra. 1985. Floral mimicry induced by mummy-berry fungus

exploits host's pollinators as vectors. Science 228:1011–1013.

Brysch-Herzberg, M. 2004. Ecology of yeasts in plant-bumblebee mutualism in Central

Europe. FEMS Microbiology Ecology 50:87–100.

Canto, A., C. M. Herrera, M. Medrano, R. Pérez, and I. M. García. 2008. Pollinator foraging

modifies nectar sugar composition in *Helleborus foetidus* L. (Ranunculaceae): an

experimental test. American Journal of Botany 95:315–320.

Canto, A., C. M. Herrera, I. M. García, R. Pérez, and M. Vaz. 2011. Intraplant variation in

nectar traits in *Helleborus foetidus* (Ranunculaceae) as related to floral phase,

environmental conditions and pollinator exposure. Flora 206:668–675.

Cnaani, J., J. D. Thomson, and D. R. Papaj. 2006. Flower choice and learning in foraging

bumblebees: effects of variation in nectar volume and concentration. Ethology 112:278–

285.

Dhami, M. K., T. Hartwig, A. D. Letten, M. Banf, and T. Fukami. 2018. Genomic diversity of

a nectar yeast clusters into metabolically, but not geographically, distinct lineages.

Molecular Ecology 27:2067–2076.

Dawkins, R. 1982. The extended phenotype: the long reach of the gene. Oxford University

Press, Oxford, UK.

Dawkins, R. 2004. Extended phenotype – But not too extended. A reply to Laland, Turner and

Jablonka. Biology and Philosophy 19:377–396.

de Vega, C., R. G. Albaladejo, S. Álvarez-Pérez, and C. M. Herrera. 2022. Contrasting effects

of nectar yeasts on the reproduction of mediterranean plant species. American Journal of

Botany 109:393–405.

Gange, A. C., and A. K. Smith. 2005. Arbuscular mycorrhizal fungi influence visitation rates

of pollinating insects. Ecological Entomology 30:600–606.

González-Mas, N., M. Cuenca-Medina, H. García-Mozo, J. M. Muñoz-Redondo, J. M.

Moreno-Rojas, F. Padilla-Álvarez, I. M. Rodríguez-Gómez, and E. Quesada-Moraga.

2023. Endophytic *Beauveria bassiana* modifies flowering phenology, floral volatile profile

and pollinator behaviour in melon. Entomologia Generalis 43:961–969.

Groemping, U. 2006. Relative importance for linear regression in R: the package relaimpo.

Journal of Statistical Software 17:1–27.

Groemping, U. 2007. Estimators of relative importance in linear regression based on variance

decomposition. American Statistician 61:139–147.

Hawkes, C. V., R. Kjøller, J. M. Raaijmakers, L. Riber, S. Christensen, S. Rasmussen, J. H.

Christensen, A. B. Dahl, J. C. Westergaard, and M. Nielsen. 2021. Extension of plant

phenotypes by the foliar microbiome. Annual Review of Plant Biology 72:823–846.

Herrera, C. M. 2014. Population growth of the floricolous yeast *Metschnikowia reukaufii*:

effects of nectar host, yeast genotype and host x genotype interaction. FEMS Microbiology

Ecology 88:250–257.

Herrera, C. M. 2025. Data: effects of yeast genotype and plant identity on sucrose concentration

of *Helleborus foetidus* nectar. Figshare. Dataset.

https://figshare.com/s/2e27d89d49dc46e4adb1

Herrera, C. M., A. M. Sánchez-Lafuente, M. Medrano, J. Guitián, X. Cerdá, and P. Rey.

2001. Geographical variation in autonomous self-pollination levels unrelated to pollinator

service in *Helleborus foetidus* (Ranunculaceae). American Journal of Botany 88:1025–

1032.

Herrera, C. M., R. Pérez, and C. Alonso. 2006. Extreme intraplant variation in nectar sugar

composition in an insect-pollinated perennial herb. American Journal of Botany 93:575–

581.

Herrera, C. M., I. M. García, and R. Pérez. 2008. Invisible floral larcenies: microbial

communities degrade floral nectar of bumblebee-pollinated plants. Ecology 89:2369–2376.

Herrera, C. M., C. de Vega, A. Canto, and M. I. Pozo. 2009. Yeasts in floral nectar: a

quantitative survey. Annals of Botany 103:1415–1423.

Herrera, C. M., M. I. Pozo, and M. Medrano. 2013. Yeasts in nectar of an early-blooming

herb: sought by bumble bees, detrimental to plant fecundity. Ecology 94:273–279.

Herrera, C. M., M. I. Pozo, and P. Bazaga. 2014. Nonrandom genotype distribution among

floral hosts contributes to local and regional genetic diversity in the nectar living yeast

Metschnikowia reukaufii. FEMS Microbiology Ecology 87:568–575.

Herrera, J. 1989. Aminoácidos en el néctar de plantas del sur de España. Anales del Jardín

Botánico de Madrid 45:475–482.

Hunter, P. 2009. Extended phenotype redux. EMBO Reports 10:212–215.

Hunter, P. 2018. The revival of the extended phenotype. EMBO Reports 19:e46477.

Jennersten, O. 1988. Insect dispersal of fungal disease: effects of *Ustilago infection* on

pollinator attraction in *Viscaria vulgaris*. Oikos 51:163–170.

Jennersten, O., and M. M. Kwak. 1991. Competition for bumblebee visitation between

Melampyrum pratense and *Viscaria vulgaris* with healthy and *Ustilago*-infected flowers.

Oecologia 86:88–98.

Lüdecke, D. 2018. ggeffects: Tidy data frames of marginal effects from regression models.

Journal of Open Source Software 3:772.

Lüdecke, D., M. S. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021.

performance: An R package for assessment, comparison and testing of statistical models.

Journal of Open Source Software 6:3139.

Nicolson, S. W., M. Nepi, and W. Pacini, editors. 2007. Nectaries and nectar. Springer-

Verlag, Dordrecht, The Netherlands.

Parachnowitsch, A. L., J. S. Manson, and N. Sletvold. 2019. Evolutionary ecology of nectar.

Annals of Botany 123:247–261.

Pozo, M. I., C. M. Herrera, W. van den Ende, K. Verstrepen, B. Lievens, and H. Jacquemyn.

2015. The impact of nectar chemical features on phenotypic variation in two related nectar

yeasts. FEMS Microbiology Ecology 91:fiv055.

Pozo, M. I., G. van Kemenade, A. van Oystaeyen, T. Aledón-Catalá, A. Benavente, W. Van

den Ende, F. Wäckers, and H. Jacquemyn. 2020. The impact of yeast presence in nectar on

bumble bee behavior and fitness. Ecological Monographs 90:e01393.

R Core Team. 2024. R: A language and environment for statistical computing. Vienna,

Austria: R Foundation for Statistical Computing.

Rering, C. C., J. J. Beck, G. W. Hall, M. M. McCartney, and R. L. Vannette. 2018. Nectar-

inhabiting microorganisms influence nectar volatile composition and attractiveness to a

generalist pollinator. New Phytologist 220:750–759.

Roy, B. A. 1993. Floral mimicry by a plant pathogen. Nature 362:56–58.

Roy, B. A. 1996. A plan pathogen influences pollinator behavior and may influence

reproduction of nonhosts. Ecology 77:2445–2457.

Schaeffer, R. N., C. R. Phillips, M. C. Duryea, J. Andicoechea, and R. E. Irwin. 2014. Nectar

yeasts in the tall larkspur *Delphinium barbeyi* (Ranunculaceae) and effects on components

of pollinator foraging behavior. PLoS ONE 9:e108214.

Schaeffer, R. N., Y. Z. Mei, J. Andicoechea, J. S. Manson, and R. E. Irwin. 2017.

Consequences of a nectar yeast for pollinator preference and performance. Functional

Ecology 31:613–621.

Schaeffer, R. N., C. C. Rering, I. Maalouf, J. J. Beck, and R. L. Vannette. 2019. Microbial

metabolites elicit distinct olfactory and gustatory preferences in bumblebees. Biology

Letters **15**:20190132.

Seino, M. M., C. de Vega, P. Bazaga, M. I. Pozo, and C. M. Herrera. 2013. Development and

characterization of microsatellite loci for the nectar-living yeast *Metschnikowia reukaufii*.

Pages 760-762 in Molecular Ecology Resources Primer Development Consortium.

Molecular Ecology Resources 13:760–762.

Vannette, R. L., and T. Fukami. 2016. Nectar microbes can reduce secondary metabolites in

nectar and alter effects on nectar consumption by pollinators. Ecology 97:1410–1419.

Vannette, R. L., and T. Fukami. 2018. Contrasting effects of yeasts and bacteria on floral

nectar traits. Annals of Botany 121:1343–1349.

Vesprini, J. L., M. Nepi, and E. Pacini. 1999. Nectary structure, nectar secretion patterns and

nectar composition in two *Helleborus* species. Plant Biology 1:560–568.

Whitney, H. M., A. Dyer, L. Chittka, S. A. Rands, and B. J. Glover. 2008. The interaction of

temperature and sucrose concentration on foraging preferences in bumblebees.

Naturwissenschaften 95:845–850.

Table 1. Summary of statistical analyses testing for effects on nectar sucrose concentration after experimental exposure of nectar from 20 different plants to 10 yeast genotypes using a factorial design. Separate models were fitted to the data ($N = 786$ sugar concentration measurements) in which the predictors were initial yeast cell density and sucrose concentration, and plant and yeast genotype identities, respectively.

Predictors in model	Effect	Parameter estimate (SE)	Test statistic	P-value	Variance explained (%)
Initial yeast and nectar features:			Student's t		
	Cell density (CD)	-0.0117 (0.009)	1.35	0.18	2.1
	Sucrose concentration (SC)	0.289 (0.13)	2.26	0.024	9.1
	CD x SC	0.0001 (0.001)	0.84	0.39	0.084
Plant and yeast identity:			F value (df)		
	Plant (P)		31.6 (19, 586)	< 2E-16	32.5
	Yeast genotype (YG)		31.7 (9, 586)	< 2E-16	15.3
	YG x P		2.2 (171,586)	7.9E-12	20.3

Caption to the figures

Figure 1. Interaction plot depicting differences in sucrose concentration after experimental exposure of nectar from different plants to different yeast genotypes. To emphasize the contrasting effect of the different yeast genotypes on different individual plants, and to facilitate comparisons across plants, model-predicted sucrose concentrations for each plant x yeast genotype combination were scaled and centered separately for each plant. See Figure 2 for the distribution of model-predicted values of sucrose concentration over the two-dimensional space defined by the 20 plants and 10 yeast genotypes tested in the experiment.

Figure 2. Model-adjusted, predicted sucrose concentration in nectar after exposure to yeasts for every plant identity x yeast strain combination.

Figure 1

Figure 2