Universal bacterial clade dynamics dominate under predation despite altered phenotypes and mutation targets

Dovydas Kičiatovas[1], Johannes Cairns[1,2,3], Veera Partanen[4], Julius Hoffmann[5], Lutz Becks[5,*], Teppo Hiltunen[2,*], Ville Mustonen[1,*]

[1] Department of Computer Science, Organismal and Evolutionary Biology Research Programme, University of Helsinki, Helsinki, Finland

[2] Department of Biology, University of Turku, Turku, Finland

[3] Turku Collegium for Science, Medicine and Technology, Turku, Finland

[4] Department of Microbiology, University of Helsinki, Helsinki, Finland

[5] Department of Biology, University of Konstanz, Konstanz, Germany

[*] Corresponding authors: Lutz Becks, lutz.becks@uni-konstanz.de; Teppo Hiltunen, teppo.hiltunen@utu.fi; Ville Mustonen, v.mustonen@helsinki.fi

Data availability

Raw sequence data (bam files) will be deposited in the EMBL ENA. Accession codes and links to be added upon acceptance of the manuscript. All code and pre-processed data needed to reproduce the downstream analyses and figures will be made available upon publication via GitHub: https://github.com/dovydask/clonal_structure.

Author contributions

J.C. and T.H. designed the long-term evolution experiment. J.C. and V.P. performed the long-term evolution experiment. J.H. performed phenotypic measurements. D.K. performed sequence data bioinformatics, downstream analyses and algorithm design. D.K., J.C., T.H., L.B. and V.M. were responsible for interpreting results. D.K. and J.C. wrote the original paper draft, with contributions from all authors. J.C. T.H., L.B. and V.M. supervised the work. All authors approved the final version of the paper.

Funding

This work was in part funded by the Research Council of Finland (Multidisciplinary Center of Excellence in Antimicrobial Resistance Research, grant no. 346126; grant nos. 330886 and 327741 to T.H; and grant no. 346128 to V.M.); Deutsche Forschungsgemeinschaft (BE 4135/12-1).

Conflict of interest statement

The authors declare no conflicts of interest.

Acknowledgments

We thank Liisa Ruusulehto, Paula Typpö, Roosa Jokela, Eeva Vakkari, Jutta Kasurinen, Iina Hepolehto, Tinja Hyvönen for their laboratory procedure work on the long-term experiment. We thank the WGGC Core Facility Cologne for the preparation and sequencing of our samples. We also thank Anthony Sun and Mikko Kivikoski for their helpful insights and comments on algorithmic and bioinformatic procedures.

Abstract

Recent studies have revealed bacterial genome-wide evolution to be complex and dynamic even in a constant environment. The evolution is characterized by the emergence of new clades competing or temporarily coexisting as each clade undergoes evolutionary change. Previous studies on predator-prey dynamics tracking simple ecological and phenotypic metrics have shown predation to fundamentally alter prey evolution, facilitating defense evolution followed by coevolution and frequency dependent selection between defended and undefended prey genotypes. Moreover, these dynamics can differ between prey species depending, for example, on mutation rate and the physiological and fitness effects of the mutations available. Here we sought to consolidate these fields by examining genome-wide evolution in five bacterial prey species separately subjected to long-term evolution under ciliate predation. For all species, we found mutational signals of prey adaptation, with genomic and phenotypic data demonstrating changes in composition between the experimental treatments. Mutational targets and dynamics also differed between prey species. Intriguingly, however, overall temporal molecular dynamics across the prey species were strikingly similar to those of bacteria evolving alone, with constant emergence, competition and quasi-stable coexistence of clades. This makes molecular evolution more interesting and less predictable than we might expect based on existing coevolutionary theories.

Keywords

Predator-prey dynamics, experimental evolution, coevolution.

Introduction

Understanding how organisms adapt and evolve is a fundamental challenge in biology that continues to intrigue researchers. While organisms seemingly respond to environmental pressures through predictable phenotypic changes (Stern 2013), the underlying genetic mechanisms reveal a far more complex and unpredictable reality (Tenaillon et al. 2012). At the phenotypic level, adaptation is often repeatable and predictable, both in controlled experiments (Wichman et al. 1999; Katz et al. 2021) and when comparing across natural populations (Papadopulos et al. 2021). This repeatability can allow us to identify key mechanisms driving adaptive changes, such as selection strength (Saxer, Doebeli, and Travisano 2010; Bernardes et al. 2021) and trade-offs between traits (Soudi et al. 2023; Maccagni and Willi 2022). However, at the nucleotide level, evolution is typically much less repeatable, even under strong selective pressures (Lässig, Mustonen, and Walczak 2017). This lack of repeatability arises from two main factors. First, different metabolic pathways can lead to the same phenotypic outcome, allowing for multiple routes of adaptation (Frickel, Theodosiou, and Becks 2017; Maharjan et al. 2015). Second, even in relatively simple environments and within a single, isolated population, clonal structure can undergo continuous and significant changes over time. These changes are driven by complex dynamics, such as competition and persistence among mutational or clonal cohorts, which can coexist for extended periods (Good et al. 2017; Lang et al. 2013). Clonal cohorts are groups of genetically identical individuals that arise from a common ancestor and share the same mutations. These cohorts play a crucial role in shaping the genetic composition of populations over time, as they compete, persist, or decline based on their fitness advantages or disadvantages, and chance. Thus, molecular evolution is shaped not only by direct selection and the emergence of new mutations but also by the complex interactions among clonal cohorts within populations over generations. Understanding these dynamics is crucial for uncovering how genetic changes respond to environmental pressures and why evolutionary outcomes can diverge, even under similar selective conditions.

Ecological interactions, those between individuals of different species, introduce additional layers of complexity to molecular evolution (Cairns et al. 2020b). For instance, species interactions can drive strong directional or stabilizing selection on populations. Strong selection may increase the efficiency of eliminating competing clonal lineages, simplifying clonal population dynamics (Corbett-Detig, Hartl, and Sackton 2015). On the other hand, species interactions can lead to coevolution and eco-evolutionary dynamics, where changes in population size both drive and result from evolutionary shifts occurring on similar timescales (Lion 2018). In these scenarios, selection pressures fluctuate continuously, varying in direction and/or strength. These fluctuations can promote the coexistence of clonal cohorts in a quasi-stable or frequency-dependent manner, making molecular evolution at least as dynamic as in single-species systems. For example, a study of microbial host-virus coevolution found that the host harbored significant genetic diversity, primarily driven by neutral molecular evolution rather than direct selection by the virus. The virus underwent fewer, strongly selected mutations, typically confined to specific genes (Retel et al. 2019). Furthermore, the complexity of species interactions can lead to conflicting selection pressures, potentially altering the trajectory of molecular evolution. This has been observed in tripartite interactions involving bacteria, phages, and plasmids, where the presence of multiple mobile genetic elements constrains the evolutionary responses typically seen in pairwise interactions (Harrison et al. 2017). Coevolution can also accelerate molecular evolution (Paterson et al. 2010), driving successive sweeps of beneficial mutations and, in some cases, leading to simpler evolutionary dynamics. These examples illustrates how co-evolutionary dynamics can lead to both complex or more simple patterns of molecular evolution. Since all populations are part of larger communities and interact with other species, these interactions play a fundamental role in shaping evolutionary dynamics. Therefore, it is crucial to study

their influence on molecular evolution. Understanding these dynamics is essential for predicting and managing evolutionary outcomes in natural and artificial systems, from ecosystem conservation to the development of sustainable disease control strategies (Lässig, Mustonen, and Nourmohammad 2023).

The patterns and rates of molecular evolution differ depending on the timescale observed. Experimental evolution studies, which track molecular evolution in replicated populations, have shown that, over short timescales, beneficial mutations accumulate rapidly in response to new selective pressures. These mutations are often large-effect and lead to significant fitness gains in a short period, allowing for an evaluation of predictability in evolutionary processes (Johnson et al. 2021). Long-term dynamics, however, are more likely governed by declining rates of adaptation, as beneficial mutations become rarer, while the overall rate of molecular evolution remains constant. Although much of our understanding of molecular evolution in the context of species interactions comes from short-term experiments, the more complex evolutionary dynamics are only observable over longer periods. Unfortunately, such long-term studies are currently lacking, leaving unanswered questions about how species interactions influence molecular evolution. Thus, the long-term evolutionary consequences of strong species interactions, particularly whether antagonistic relationships can leave a lasting molecular signature in prey, remain to be fully elucidated.

To that end, in this study, we investigated the long-term molecular and phenotypic evolution of five bacterial species (*Brevundimonas diminuta*, *Comamonas testosteroni*, *Pseudomonas fluorescens*, *Sphingomonas capsulata*, and *Serratia marcescens*) as they grew with and without a ciliate predator, *Tetrahymena thermophila*, over a period of 800 days (Fig. 1, first two panels; see Methods). Our aim was to quantitatively assess how the presence of the predator influenced molecular evolution in the bacterial prey (Fig. 1, third panel; see Methods). Specifically, we sought to determine whether the predator's presence led to measurable changes in long-term molecular evolution compared to predator-free populations. To achieve this, we tracked clonal diversity and phenotypic changes in replicated bacterial populations over time, identifying recurrent gene targets along the way.

Figure 1: Graphical abstract of this study. Each of the five prey species undergo two tracks of long-term evolution – evolving alone and coevolving with predator *T. thermophila*. With prey freeze-stored every four transfers (i.e., weeks), resulting genomic and phenotypic data are time-resolved and allow for clonal structure inferences.

Results

Growth phenotypes differ between prey populations coevolving with a predator and populations evolving alone

First, we investigated the effects of predation on the prey population growth dynamics. A previous study by Cairns *et al.* on the same predator-prey system (Cairns et al. 2020a) has shown that the ciliates evolve alongside the bacterial prey. Here, we sought to describe the phenotypic changes in these prey species over a longer period of time than previously studied. We analysed the growth curve data (i.e., growth phenotype) of sampled populations from each study population at the beginning and the end of the long-term experiment. The isolates were grown on three nutrient-rich experimental media – without ciliate (control), with ciliate and with added KCl salt without ciliate (see Methods). Adaptation to a particular stressor typically influences multiple bacterial phenotypes due to the pleiotropic effects of altering key structures or functions, as is the case with antibiotic cross resistance and collateral sensitivity (Colclough et al. 2019). Therefore, in addition to studying the phenotypes of bacteria evolved alone or with ciliate in the primary conditions of interest – experimental medium with and without ciliate predation – we also included a salt stress condition to further measure a fitness cost of prey of either treatment lineage growing without the predator.

For each growth curve, we computed yield Y, maximal per-capita growth rate ρ_{max}, time to the latter $t_{\rho_{max}}$ and the area under the curve (AUC). We contrasted these parameters between the coevolutionary histories and found that all of their means differ significantly when the bacterium is grown in a medium where the ciliate is present (Fig. 2A). Yield, AUC and ρ_{max} are higher, and the latter is attained sooner when the bacteria have coevolved together with the predator over the course of the long-term evolutionary experiment, meaning that in the presence of predation, they consistently grow better than bacterium that evolved alone. This is expected as bacteria evolve anti-predatory defenses in the presence of the ciliate, as shown before in a previously described ciliate-*P. fluorescens* system in Scheuerl *et al.* study (Scheuerl et al. 2019). However, there is no sufficiently strong statistical evidence that growth differs in either control or salt stress media. Nevertheless, it is noteworthy that the coevolved bacteria isolates do not display weaker growth capabilities in the control medium, compared to the bacteria that evolved alone. This could be explained by the expectation that coevolved prey populations would include both more reproductionally fit undefended phenotypes and ciliate-resistant but less fit defended phenotypes.

We did not observe a consistent direction of growth phenotype evolution with respect to time in the long-term experiment. Isolates sampled later in the experiment may grow better or worse in terms of area under the curve (AUC) than in the beginning without a clear pattern across replicates, though it clearly differs between species (Supplementary Fig. 4). Despite the overall signal separating coevolutionary histories in terms of differences in AUC in the presence of predation, each species has a unique phenotypic response to each growth medium, regardless of their coevolutionary history. Notably, all replicates of both treatments of *S. marcescens* do not appear to have any notable differences other than slight change in variance when grown in the presence of predation, possibly indicating pre-existing defenses to ciliate (Supplementary Fig. 4, bottom row). Isolates of *B. diminuta* and *S. capsulata* show the most stark difference in growth rates between the histories in the presence of predation. Overall, the variance between isolates' growth is lower in the presence of predation than in the other test conditions.

To determine whether predation exerts a sufficiently strong selective pressure in our experiment, we computed ratios of AUC between species-specific ancestral and (co-)evolved population growth curves (see Methods; Fig. 2B). We saw that in control and salt stress growth environments the AUC does not, on average, differ with respect to the ancestors (two-sided Welch's t-test p-value = 0.325

and p-value = 0.949 respectively), whereas there is a significant difference when isolates are grown in the presence of predator (p-value < 0.001, Fig. 2B, "Predator medium" panel). On average, populations that have evolved without the predator are growing worse compared to their ancestral populations when exposed to the predator (ratio < 1), possibly indicating that adaptation to the growth medium has made them less defended against the predator. On the other hand, coevolved populations in the presence of predators grow much better with respect to their ancestors (ratio > 1), indicating that indeed these populations have been forced to adapt to the stress of predation. However, it must be noted that there are clear differences in these ratios between each species and also isolate sampling timing, with some species clearly adapting to the predator over time, e.g., late experiment isolates in each of the *S. capsulata* populations (Supplementary Fig. 5).

To assess the predictability of the growth curves in terms of the species, coevolutionary history, replicate population within species and the timing of the isolate sampling, we constructed a label for each growth curve, comprised of these categorical variables. Using the growth curves, concatenated across the three types of media the isolates were grown in as our input, we trained an ensemble of 100 random forest models with random splits of 50% of the data, predicted the label of the remaining randomly split 50% of growth curves and averaged the prediction results across the random forest instances (see Methods; Fig. 3).

As expected, bacterial adaptation in the presence or absence of ciliate predation had a multidimensional effect on bacterial phenotypes. Moreover, this effect was specific to the bacterial species, and within species, to the coevolutionary history. However, the replicate populations are not as distinct within the pooled isolates of the same coevolutionary history, and neither across the time points within a replicate population. Nevertheless, we clearly see higher classification accuracy between the three coevolved populations in every species (Fig. 3, right hand panel) than in populations that evolved without the predator, suggesting that the phenotypic dynamics are more distinguishable between coevolved populations.

Figure 2: **A** Growth curve statistics: yield (optical density at the end of measurement), area under the curve (AUC), maximal per-capita growth rate ρ_{max} and time to maximal per-capita growth rate $t_{\rho_{max}}$. Statistically significant results from two-sided Welch's t-tests, comparing the means of all isolates between opposite coevolutionary histories when grown in the three different media (see Methods), are marked next to the relevant pair of boxes. The legend is shared with the lower panel. **B** Box plots of all growth phenotype AUC ratios between (co-)evolved and ancestral growth curves (species-specific growth curve AUC divided by corresponding species' ancestral growth curve AUC, see Methods). There is a clear difference in fitness (measured as AUC) between evolutionary treatments with respect to ancestral fitness, indicating that the predator exerted sufficient pressure to influence phenotypic changes.

Figure 3: Combined and averaged confusion matrix of 100 random forest classifier instances, trained on 50% of randomly sampled isolate growth profiles (combined isolate growth curves in all three growth media) and tested on the remaining 50% of the data (label proportions preserved in both training and testing sets). The X-axis represents predicted data labels and the Y-axis – true labels. The rows and columns of the confusion matrix are separated according to the alphabetic order of the species names, and each inner square of the confusion matrix, representing a species, is further subdivided into the two evolutionary histories, then three replicate populations and lastly, the time points. "ANC" denotes the growth profile of the ancestral strain of the respective species. The color gradient represents the average number of assigned samples from the test set, averaged over 100 random forest instances. While there is little confusion between the species and evolutionary histories, the growth profiles of replicate populations within species and their evolutionary history are more difficult to separate. However, coevolved species-specific replicates are distinguished with higher accuracy than when evolved alone (right panel).

Bacterial molecular evolutionary dynamics are similar regardless of the presence or absence of predator

Next, for each replicate of a given prey species, evolving with and without the predator present, we calculated the frequencies of single nucleotide polymorphisms (SNPs) and short insertions or deletions (indels) and their change over the course of the experiment. The final accumulated variant counts differ vastly between species and between replicate populations (Supplementary Fig. 1, Supplementary Table 1), indicating a large degree of variability in the accumulated mutation counts even within the replicates of the same species under same environmental conditions. Likewise, population 1 of *C. testosteroni* steadily acquires almost double the number of variants compared to the remaining populations of either treatment of the same species. However, at the 95% confidence level, none of the species has a significant two-sided Welch's t-test result when comparing the means of final accumulated variants counts between experimental groups – although it is worth noting that the sample sizes for comparison might be too low. The trajectories of most variants form clearly defined clusters, in line with the expectation that most mutations would hitchhike with beneficial mutations under selection (Lang and Desai 2014). Henceforth, we analyzed these clusters as variant cohorts (see Methods; Fig. 4A).

Several moderate to high-frequency variants in known mutator genes were observed together with a sharp increase in mutation counts right after the time of their detection, likely explaining why some study populations accrue much higher mutation counts than others within the same species and treatment. Variants in mutator gene *mutS* were detected in three species in both treatments: a stop-gain (appeared on day 700, mean frequency after appearance 0.32) and disruptive inframe deletion (appeared on day 476, mean frequency after appearance 0.36) in coevolved *P. fluorescens* populations 2 and 3, respectively; a missense variant in coevolved *S. marcescens* population 2 (appeared on day 679, mean frequency after appearance 0.75); disruptive (appeared on day 553, mean frequency after appearance 0.99) and conservative (appeared on day 553, mean frequency after appearance 0.96) inframe deletions in evolved-alone *C. testosteroni* populations 2 and 3, respectively; and a missense variant in evolved-alone *P. fluorescens* population 1 (appeared on day 412, mean frequency after appearance 0.64). Variants in *mutL* were detected in the following populations: a frameshift variant in coevolved *C. testosteroni* population 1 (appeared on day 448, mean frequency after appearance 0.35); a frameshift variant in evolved-alone *C. testosteroni* population 1 (appeared on day 146, mean frequency after appearance 0.66); and a stop-gain variant in evolved-alone *B. diminuta* population 1 (appeared on day 272, mean frequency after appearance 0.65). Additionally, in *C. testosteroni*, coevolved population 1 had a *uvrC* missense variant (appeared on day 504, mean frequency after appearance 0.13) and evolved-alone population 1 had two *uvrA* missense variants (appeared on days 553 and 776, mean frequencies after appearance 0.55 and 0.25).

Next, using the variant cohorts, we inferred the clonal structure within each population at every time point – treating the cohorts as genetic alleles, we developed a tree-based algorithm to assign each cohort to the role of either a new clone or a sub-clone of an existing clone (see Methods). Reconstructing clonal structures from variant frequency data is widely used in cancer research, where linking cancer evolution to clinical outcomes can inform therapies (Salcedo et al. 2020; Gillis and Roth 2020). Cancer clonal structure inference is difficult due to several factors, such as DNA copy number variation, genome diploidy and paucity of long term time-series data (Dentro et al. 2021), which are not present in our bacterial genomes, making this type of inference more readily viable. With the inferred allele hierarchy, we calculated the allele frequencies over time and visualize them as Muller plots and phylogenetic trees (Fig. 4A, B). Muller plots in this kind of population structure inference from variant data have been presented before (Frazão et al. 2022), however, to our knowledge, there is no tool that computes high-confidence population clonal structures automatically

₂₄₇ (nevertheless, we have manually checked all our visualised solutions as a form of quality control, see
₂₄₈ Methods).

₂₄₉ We observed a large degree of variability in clonal dynamics, even between the replicates of
₂₅₀ the same species (Fig. 4, "Coevolved with ciliate" and "Evolved alone" panels). Most prey species
₂₅₁ exhibit similar variant frequency dynamics, such as selective sweeps that occur throughout the entire
₂₅₂ duration of the experiment. We have also observed non-trivial dynamics of variant clusters being
₂₅₃ maintained at a high frequency over long periods of time without these mutations getting lost or
₂₅₄ fixed (Fig. 4, "Coevolved with ciliate" and "Evolved alone" panels), suggesting previously described
₂₅₅ state of quasi-stable coexistence in bacterial populations (Good et al. 2017). The latter is clearly
₂₅₆ visible in the population 1 of *B. diminuta* in clades AC and AD, both undergoing their own sub-
₂₅₇ clonal expansions at different points in time. Other species exhibit similar clade coexistence, albeit
₂₅₈ with more stark difference in the clade frequencies. Similarly, selective sweeps occur in several
₂₅₉ study replicates, with coevolved *S. capsulata* population 3 displaying textbook successive sweeps.
₂₆₀ Successive hard selective sweeps are only observed in four cases; interestingly, all three coevolved *S.*
₂₆₁ *capsulata* populations, grown with ciliate, are among them, and the remaining one is a replicate of
₂₆₂ *C. testosteroni* grown without ciliate (Fig. 4, "Coevolved with ciliate" and "Evolved alone" panels).
₂₆₃ Furthermore, many populations show a pattern of late large scale diversification events, where many
₂₆₄ different clades start to appear in a quick succession halfway into the experiment (more than 400
₂₆₅ days).

₂₆₆ The number of clades consistently in both treatments increases over the duration of the exper-
₂₆₇ iment to the maximum of nine clades (linear regression coefficient of determination $R^2 = 0.75$ and
₂₆₈ slope 0.0085 in coevolved populations, $R^2 = 0.69$ and slope 0.0081 in evolved-alone populations).
₂₆₉ However, interestingly, the number of clades present at any given time in the experiment, after
₂₇₀ approximately 250 days, consistently stays at 3-5 clades (Supplementary Fig. 2) – suggesting active
₂₇₁ turnover of clades within a population as time goes on. This is likely explained by clonal interference
₂₇₂ between clades – however, since we do not observe consistent and repeated selective sweeps across
₂₇₃ the whole population, clonal interference dynamics are likely moderated by frequency dependent
₂₇₄ selection dynamics that sustain clade coexistence for longer periods of time (Maddamsetti, Lenski,
₂₇₅ and Barrick 2015; Good et al. 2017).

₂₇₆ To quantitatively compare the inferred clonal structures between the experimental groups, in
₂₇₇ addition to Muller plots, we computed hierarchical representations in the form of strictly bifur-
₂₇₈ cating phylogenetic trees (Fig. 4B, Supplementary Fig. 7, Supplementary Fig. 8), where the total
₂₇₉ branch length represents the number of days a clone existed since its emergence until either the
₂₈₀ end of the studied experiment or the final point in time the clone retains a frequency above 0.1
₂₈₁ (provided no further sub-clones appear) – otherwise, the branch length is truncated. We computed
₂₈₂ the sum, mean and variance of branch lengths, the number of leaves (corresponds to number of
₂₈₃ clones plus the background clone), maximum and average depths of the leaves, as well as Sackin,
₂₈₄ Colless and B1 indices of tree balance (Shao and Sokal 1990). Using regression analysis, we tested
₂₈₅ these metrics for statistically significant differences between the experimental groups (with predator
₂₈₆ and without predator) in two ways: across all study replicates, using the species indicator as a co-
₂₈₇ variate (ANCOVA (Rutherford 2001)), and between species-specific experimental groups (ANOVA)
₂₈₈ – interestingly, none of these measures differ significantly between the treatments (95% CL).

₂₈₉ Going by species, only the experimental groups of *B. diminuta* are significantly differing in their
₂₉₀ trees' B1 indices (at 95%CL, p-value 0.004) – trees that have their leaves closer to the root on average
₂₉₁ are considered balanced, indicating that there is a greater imbalance observed in the structure of
₂₉₂ populations that evolved without ciliate. For complete table of within-species ANOVA results, see
₂₉₃ Supplementary Table 2.

₂₉₄ These results suggest that the presence of ciliate does not predictably affect the clonal structure

₁₀

of the studied prey populations, or, at the very least, the changes are subtle and remain largely hidden under shared evolutionary processes, such as adaptation to the growth medium. Instead, we observed the usual clonal interference and likely frequency dependent selection dynamics that universally characterize even simple microbial systems without predation (Maddamsetti, Lenski, and Barrick 2015).

Figure 4: **A** Clonal structure evolution in *B. diminuta* under predation (replicate population 1) in terms of variant frequency trajectories over the course of the long-term experiment. Trajectories are first clustered into cohorts and then assigned to a likely position in the clonal structure (see Methods) and represented as a Muller plot. Vertical dashed bars in the trajectory and the Muller plot mark the appearance of the clone (cohort trajectory exceeding the emergence threshold, see Methods). **B** A corresponding phylogenetic tree representation of the population structure of *B. diminuta* under predation (replicate population 1). The time of the appearance of each clone is color-coded to correspond to the clone information in the legend and the variant trajectories with shaded areas in the trajectory and Muller plots, respectively. The branch labels in the tree denote the time the clone remains above a survival threshold (provided no further sub-clones emerge; see Methods). A non-trivial long-term coexistence of the inferred clades is clearly seen in the Muller plot and the phylogenetic tree. **Evolved alone** and **Coevolved with ciliate** boxes: Muller plot showcase for the remaining 29 populations evolving under and without predation, respectively. There is significant variability both between species and between replicate populations of the same species. Long-term coexistence of clades is a recurring theme, along with examples of selective sweeps. Populations with truncated Muller plots represent populations where a portion of time series data was lost to contamination during laboratory procedures.

Mutation targets are recurrent both within and across species, and differ depending on the presence or absence of the predator

Lastly, we investigated the recurrence of non-synonymous variants identified by our genomic analyses to determine if there are patterns in recurrence and evidence of selection for mutations in specific genes. We defined the recurrence of a mutated gene as it being affected by at least one variant in at least two out of three replicates of a species, with and without the presence of the predator separately (Fig. 5, top panel).

We also investigated the appearance times of the recurrent gene mutations, i.e., the appearance of the clade they belong in (Fig. 5, bottom panel) and found that the overall distribution of the timing of appearance is significantly differing between the coevolutionary histories, i.e., populations evolving with the predator and populations evolving alone (two-sided Kolmogorov-Smirnov test p-value = 0.00003; Fig. 5, bottom left histogram). It appears that in the populations coevolving with the predator, more recurrently targeted genes are affected late in the experiment (> 600 days) than in populations evolving alone, moreover, only the coevolving populations show recurrently affected genes appearing within the first 100 days of the experiment.

In total, only eight genes – *ppk*, *sasA*, *glpK*, *rcsC*, *mutS* *dppA*, *rpoB* and *corA* – are recurrent in more than one of the species. The gene encoding the adaptive sensory kinase *sasA*, recurrently affected in all studied species except *S. marcescens*, is purported to be a core part of the circadian rhythm control mechanism in cyanobacteria (Iwasaki et al. 2000), though to our knowledge, its exact physiological role in other bacteria is not known. Polyphosphate kinase gene *ppk* is responsible for a variety of functions (Rao, Gómez-García, and Kornberg 2009) and is recurrently targeted in 3 species, though the recurrence is not limited to coevolved populations, unlike *sasA*. All remaining affected genes are limited to recurrence between at most two species.

To test whether the recurrence of any gene is statistically significant, we performed a regression analysis using ANCOVA on gene hit indicator data (1 if a non-synonymous variant has been detected in a gene at least once, 0 otherwise) between with- and no-predator groups using species indicator as a covariate. Nine genes – *barA*, *dppB*, *flhC*, *lptA*, *rcsC*, *relA*, *rpoB*, *rpoC* and *sasA* – have been identified as having a significantly differing recurrence between the experimental groups, i.e., growth with ciliate versus without ciliate (Supplementary Table 3).

The majority of remaining genes recurrent only in coevolved populations are responsible for various metabolic functions, cell wall formation and maintenance, though some are related to biofilm formation, which is expected given bacteria propensity to form biofilm in response to external stress, such as predation, in addition to simply being a viable strategy to adapt to the environment (Kimberly K Jefferson 2004). Genes *flhC* and *rpoC*, recurring only in coevolved *S. marcescens* populations, are associated with biofilm formation (Ferrières and Clarke 2003; Finn et al. 2000; Sule et al. 2011). Hybrid sensor kinase gene *rcsC* has been identified as an important part of *E. coli* biofilm formation (Ferrières and Clarke 2003) and is recurrently affected in all replicates of both *C. testosteroni* and *P. fluorescens*. In contrast, genes *rpoB*, *pilQ* and *relA* are associated with biofilms and motility (Mollet, Drancourt, and Raoult 1997; Sule et al. 2011) and are recurrent only in *C. testosteroni* populations grown without ciliate and are likely not related to defense against the predator. Gene *rpoB* is also recurrent in two out of three *B. diminuta* evolved-alone populations. Gene *cheB* (also known as *wspF*) is recurrent in all three coevolved *P. fluorescens* populations and has been previously reported to be also recurrently targeted in a dual-stressor (antibiotic and ciliate) system with *P. fluorescens* as prey (Hiltunen et al. 2018). Additionally, mutator gene *mutS* is recurrent in *P. fluorescens* populations. Many genes that recur in both experimental treatments are related to various metabolic functions and are likely responsible for adaptation to the growth medium rather than defense.

Figure 5: Top panel: non-synonymous variant hit recurrency across the experiments in known genes. Columns (i.e., genes) are hierarchically clustered according to the pairwise Hamming distance. Populations without ciliate (blue rows) have 17 recurrent genes containing variants that do not occur in populations grown with the ciliate (orange rows). Conversely, the set of genes that are hit only in populations with ciliate is larger. Bottom panel: variant timing of appearance (based on cohort appearance to which a variant belongs to). Darker circles indicate that there are multiple unique study replicates where a variant in a given gene appeared at the same time. Marginal counts of variant appearance timing are shown as a histogram.

Discussion

Our study builds upon the long-term experimental evolution methodology presented in Good *et al.* study (Good et al. 2017), where bacterial molecular evolution is shown to be complex and incessant over long time scales, even in a constant environment. With our experiment, we aimed to study whether a strong selective pressure, i.e., predation, would significantly change the clonal dynamics and the outcome of molecular adaptation in the prey bacterium. We observed clear signs of phenotypic adaptation, where coevolved bacterial prey populations grow significantly better in the presence of ciliate compared to control populations. Moreover, recurrently mutated gene sets are different between the experimental groups – both phenotypic and genomic data support treatment-specific evolution. Population sizes of predator and prey in coevolved populations, as expected, fluctuate in opposition to one another and are negatively correlated (linear regression $R^2 = -0.42$, slope -1.78, Supplementary Fig. 6). However, surprisingly, despite our observance of rich and non-trivial clonal structure dynamics, we cannot detect a strong signal of predation within the clonal structures of our study populations. Instead, we find that the molecular evolutionary dynamics are predominated by processes common to all environments. These processes feature constant dynamic emergence of adaptive clades, and coexistence and competitive dynamics between them (Good et al. 2017).

What we observed in this study may be a case of "emergent simplicity" – complex and un-

predictable genomic evolution is ultimately resulting in predictable phenotypic changes (Lässig, Mustonen, and Walczak 2017). There may be many metabolic pathways through which prey bacterium adapt to their immediate environment (both the growth medium and the ciliate), changing their mode of growth in a predictable manner, but on a genomic level, prey population clonal dynamics may be largely driven by the aforementioned universal evolutionary processes, much like in previous studies on pairwise bacterial community evolution (Meroz et al. 2024). Moreover, it has been suggested that the repeatability of evolution may not necessarily be mainly driven by strong selective pressure (Bailey et al. 2017). Partly repeatable evolution in terms of clade dynamics has been described in influenza (Strelkowa and Lässig 2012; Łuksza and Lässig 2014), where strong clonal interference drives successful clades to constantly emerge victorious over the other clades in the population – that is predominantly not the case in our clade trees, and instead we observed constant competition and coexistence of clades. An interesting thing to note is that we observe rapid changes in clonal diversity that occur rather late – in some cases, even in the final 100 days in our 800 day-long experiment (Fig. 5) – so it would be interesting to see whether there are any significant changes in clonal structure over much longer timescales, though that can be, unfortunately, time and cost prohibitive.

There might be several explanations for the lack of clearly observable predator signal in clade dynamics in our study: 1) our data resolution might be insufficient to observe subtle changes in between genomic sampling (every four weeks in our study); 2) the predator in our study, *Tetrahymena Thermophila*, is a generalist predator, and therefore may not exert a strong enough selective pressure to influence clade dynamics; and 3) frequency dependent selection, which prevents a quick competitive exclusion of the prey clades (Maddamsetti, Lenski, and Barrick 2015), unlike in influenza evolution. Moreover, within each population, there is a constant maintenance of several clades coexisting at the same time above threshold frequencies (Supplementary Fig. 2).

Also, we recognize that our statistical power (three replicates per species, per treatment) might be a limiting factor in detecting subtle clonal dynamics. To fully disentangle the clonal dynamics of coevolving prey, we believe that a larger number of phenotypic measurements (than 10 clones per sampling time point in our study) would be needed from every sampling time point to account for the ever-increasing clonal diversity of the populations – growth profiles of the limited number of isolates taken from a time point where one clone dominates might be hiding subtle effects of clones that are present at lower frequencies. Additionally, since the first time point of isolate sampling represents populations that have already (co-)evolved for at least a month, we cannot compare early prey growth dynamics between the treatments. Furthermore, sampling the populations for genomic data every four weeks might be too coarse to account for rapid eco-evolutionary dynamics, such as overall prey and predator population size fluctuations during one-week transfer cycles (see Methods), though it is not clear whether increasing the overall time resolution with respect to the genomic data sampling alone would result in different conclusions.

For future studies, we believe that long-term coevolution experiments need carefully designed linking between genomic and phenotypic data tracks in order to disentangle the impact of strong environmental pressure (i.e., predation) from other evolutionary processes. Ignoring the identity of mutated genes, clonal dynamics, represented by variant cohort dynamics of SNPs and indels, seems quickly to become so complex that differentiating predation pressure from other evolutionary processes molding it remains an outstanding challenge. Furthermore, it would be interesting to study the coevolution of the predator as well and contrast the evolution of prey defense to evolution of predator offense – for that, one would need to produce equally rich genomic data from the predator and investigate clonal dynamics and gene recurrence in a similar fashion, though it is expected that the prey would experience greater evolutionary changes than the predator. Nevertheless, studying predator coevolution in terms of predator population clade dynamics might yield interesting novel

insights.

While it is still surprising that in this study we do not observe differences in clade dynamics between our evolutionary treatments, given the strong phenotypic and genomic effects from predation, we believe that the predator signals, while being detectable, remain masked within the clade dynamics by pervasive prey clade interactions, stable coexistence and frequency dependent selection that maintains both undefended and defended types within the prey populations. We believe that our study further reinforces the non-triviality of clonal dynamics even in the presence of a strong selective pressure despite easily detectable phenotypic differences and also provides new methodology to represent and investigate the clonal structure of a population solely through genomic time series data.

Materials and Methods

Experimental setup

The focal predator of this experiment is a ciliate *Tetrahymena thermophila*, preying upon these five bacterial species: *Brevundimonas diminuta*, *Comamonas testosteroni*, *Pseudomonas fluorescens*, *Sphingomonas capsulata* and *Serratia marcescens*. *Brevundimonas diminuta* has been previously studied and is found in lake habitats shared with *Tetrahymena* (Becks et al. 2005), *Comamonas testosteroni* is known to have specific defenses against protozoan grazing (Matz and Kjelleberg 2005). The remaining three – *Pseudomonas fluorescens*, *Sphingomonas capsulata* and *Serratia marcescens* – are model species commonly used in prey–predator studies. They are a subset of the same organisms studied in the Cairns *et al.* 2020 study (Cairns et al. 2020a); therefore, a detailed description of these species is omitted in this article.

For each prey species, three replicate populations were prepared and grown in 5% King's Broth (KB) medium with and without the presence of the ciliate. In total, 30 bacterial populations have been evolved over the course of 800 days, with 1% of the population being transferred to a fresh medium every 7 days. A sample from each prey population was taken and freeze-stored every 4 transfers (i.e., 28 days), and predator cells were made axenic 1–2 times per year by transferring coevolving cultures to 50 ml of 20% R2A (Reasoner's 2A agar) medium, then to antibiotic cocktail in 6 ml PPY (Proteose Peptone Yeast Extract Medium), and finally to 50 ml PPY for growth. Axenic predator cells were stored in liquid nitrogen.

Optical densities (prey) and cell counts (predator) were recorded at every transfer to obtain a time series of population growth over the course of the experiment.

Genomic data and variant calling

Reference genome sequences of all species in this study were obtained from ancestral (i.e., unevolved) samples in the PacBio HiFi long-read sequencing laboratory (URL: https://www.pacb.com). Genetic variants of each single-species bacterial population sample were obtained via bulk sequencing using short-read next–generation sequencing (NGS) equipment Illumina NovaSeq 6000. After assessing the read quality metrics of each prey population sample, genetic variants were obtained by following this read processing, variant calling and annotation workflow: Illumina DNA adapters trimmed with *cutadapt* (version v3.5; NBIS) (Martin 2011), reads aligned to reference sequences with *BWA* (version 0.7.17; Li and Durbin) (Li and Durbin 2009), read duplicates marked with *Picard* (version 2.27.5; Broad Institute, URL: https://broadinstitute.github.io/picard/), variants called in a multi-sample mode and subsequently filtered with *GATK Mutect2* (version 4.3.0.0; Broad Institute) (Benjamin et al. 2019), and finally annotated with *snpEff* (version 5.1d; Pablo Cingolani) (Cingolani et al.

2012). Sequence data for populations coevolved with the predator extends beyond 800 days (up to approximately 1400 days) and all samples have been used to call the variants, however, because populations evolving alone were limited to a 800 day period, the variant data corresponding to samples taken later than 800 days into the experiment were not included in any downstream analyses of this study (see Supplementary Information for sample curation and tool usage details).

Clonal structure inference from mutation cohorts

Variant trajectories have been assigned to variant cohorts (clusters) using read counts of reference and alternate alleles with PyClone-VI (Gillis and Roth 2020). Though any other similar clustering software would likely be sufficient for this task, we chose PyClone-VI for its acknowledged status in bioinformatics research and ease of use. Since this software is originally intended for human cancer genomics and the genomes under our study are haploid, we set the required parameters for major and minor copy numbers to one and zero respectively for every variant within the genomic time series data of any given population from our experiments. For each replicate separately, using 15 for an initial guess for the number of clusters with 100 restarts of the inference algorithm, we repeated the inference ten times, each with a different random seed (seed values 1, 2, ..., 10), and selected the model with the highest final evidence lower bound (ELBO) value, as described in the PyClone-VI manuscript (Gillis and Roth 2020). The final number of inferred clusters varied from four to nine – however, we note that repeated inferences with different seed values may yield slightly different variant trajectory clusters and their numbers.

Given the computed variant cohorts of a prey population, we infer its clonal structure by following two clonality rules: 1) for two competing clones to co-exist, the sum of their frequencies cannot exceed 1 at any time point, and 2) the sum of the frequency of every sub-clone cannot exceed the frequency of the clone they belong to. The hierarchy of the clonal structure is represented by a tree, where each node represents a mutation cohort. As the cohorts "emerge" over time, i.e., their frequency exceeds a chosen threshold at a given time point (0.1 by default; some replicates needed higher thresholds, e.g., 0.15, to resolve obviously wrong hierarchical assignments due to multiple cohorts emerging at the same time with a similar frequency, close to the threshold), they are added as new nodes to the growing trees. After the first cohort has emerged and has been placed under the root node of the tree, every subsequent emerging cohort creates new trees, each representing one of the possible placements of the new node within each tree constructed before. Therefore, the total number of such possible trees that include every mutational cohort is the factorial of the number of inferred cohorts. The order of the cohort emergence is known, except in cases where two or more cohorts exceed the emergence threshold at the same time, doubling the number of possible trees with each additional emerging cohort at such a time point – because of that, we specify the order manually to prevent possibly degenerate cases due to incorrect clonality role assignment and to limit computational cost.

Next, having a set of complete clonal hierarchy trees for a given population, we use the mutational cohort frequencies Q to calculate the allele frequencies F that correspond to the inferred clonal structure at a given time t. Given a clone c and using $x \subset c$ to denote that clone x is a sub-clone of c, we define the allele fraction as:

$$F_c = Q_c - \sum_{x \subset c} Q_x$$

Since the originally computed mutation cohort frequencies are inherently noisy because of the variant trajectory variation within trajectory clusters, we use the newly calculated allele frequencies to marginalize the cohort frequencies, i.e., adjust their values per time point so that they fully agree with the two clonality rules. That is done by iteratively subtracting a small value (0.1% of the

absolute F_c value) from the frequencies of alleles under consideration until their respective sum is either less than 1 or less than the frequency of the clone they are sub-clones of, respective to the placement of the considered allele in the previously inferred tree structure. This way, we evaluate the consistency of the inferred clonal hierarchy both quantitatively and visually.

Lastly, we select a tree with the lowest total marginalization cost to be the representative clonal structure of a given replicate. Since the marginalization of every possible tree is computationally time-prohibitive, we first select a subset of most likely tree configurations by implementing an arbitrary scoring function which corresponds to the two clonality rules; when considering a node that represents a cohort that emerged at time t, the score is the number of future time points $t_f > t$ where neither of the rules are violated, allowing up to 0.01 of absolute error to account for the potential noise in the frequency data. While the minimum score usually represents low-cost solutions, they may not be the lowest possible ones, and an objective solution other than the marginalization is not known. To help mitigate this, we marginalize 50 best scoring clonal structures and choose the one with the minimal cost as our representative structure. Inferred clonal structures presented in this article are intended as examples of automated, yet realistic, solutions and not objectively best ones – it would likely be possible to find marginally lower-cost solutions.

We note that multiple rule-consistent and high scoring solutions are sometimes possible, especially when low-frequency cohorts emerge late in a population with multiple competing clones present at sufficiently high frequencies. Additionally, some cohorts never reach our emergence threshold, and we remove these cohorts altogether. Nevertheless, this method could be extended to potentially handle every possible configuration of the clonal structure tree in a more efficient way.

Phenotypic data

From each bacterial population in our study, in addition to four ancestral (day 0) isolates, ten isolates were taken at two time points in the experiment. The first sampling, early in the experiment, is done at days 135 (*S. marcescens*) and 64 (rest of the study species) from the evolved-alone populations; and at day 43 (all study species) from the coevolved populations. The second sampling, late in the experiment, is done at days 723 (*S. marcescens*) and 652 (rest of the study species) from the evolved-alone populations; and at day 631 (all study species) from the coevolved populations. In total, there were $5 \times 2 \times 3 \times 2 \times 10 = 600$ isolate growth curves (5 species, 2 evolutionary treatments, 3 replicate populations, 2 time points, 10 isolates).

The isolates were then grown in three kinds of media: 5% King's Broth (KB) medium (without the presence of ciliate), predator medium (with ciliate) and salt stress medium (potassium chloride KCl; without ciliate). Growth curves were extracted by measuring the optical density of the grown bacterial colony every 20 minutes using the Agilent LogPhase 600 nm device. For each isolate, optical density measurements were replicated three times in total. Due to some inherent variability in these technical replicate measurements, we computed their median at each time point and used the resulting curve as our representative growth curve of that isolate.

Additionally, there were four ancestral strain isolates taken for each of the five species and for each of the three growth mediums, totalling $5 \times 3 \times 4 = 60$ ancestral growth curves. To compute ratios of area under the curve (AUC) metric between isolates of either evolutionary treatment versus their ancestral counterparts, we computed the median AUC of the four technical replicates of ancestral growth per species, per growth medium (control, salt stress and predator mediums). Then, per each species separately, isolates of populations of both evolutionary treatments were divided by the corresponding species median AUC in the relevant growth medium.

Random forest classifier

To assess the classification accuracy in classifying phenotypic growth profiles (growth curves from all three growth media combined) according to species, evolutionary history, replicate and sampling time, we used scikit-learn (Pedregosa et al. 2011) (version 1.4.2) Python (version 3.10.14) implementation of the random forest classifier to train 100 random forest models and averaged the resulting confusion matrices representing the test dataset over all model instances. The final averaged confusion matrix (Fig. 3) is an element-wise sum of all produced confusion matrices, divided by the number of model instances, i.e., 100. Before learning each model, phenotypic growth profile data was randomly split in half for training and testing datasets in a stratified fashion, i.e., keeping the proportion of each unique label equal across both datasets. For result repeatability, each train-test split and random forest model instance were generated with a Numpy (Harris et al. 2020) (version 1.26.4) random seed $s = 0, \ldots, 99$ for each of the 100 runs.

References

Bailey, Susan F. et al. (2017). "What drives parallel evolution?" In: *BioEssays* 39.1. ISSN: 0265-9247. DOI: 10.1002/bies.201600176.

Becks, Lutz et al. (June 2005). "Experimental demonstration of chaos in a microbial food web". In: *Nature* 435.7046, pp. 1226–1229. ISSN: 00280836. DOI: 10.1038/nature03627.

Benjamin, David et al. (2019). "Calling Somatic SNVs and Indels with Mutect2". In: *bioRxiv*.

Bernardes, Joana P. et al. (2021). "The evolution of convex trade-offs enables the transition towards multicellularity". In: *Nature Communications* 12.1. ISSN: 20411723. DOI: 10.1038/s41467-021-24503-z.

Cairns, Johannes et al. (June 2020a). "Evolution in interacting species alters predator life-history traits, behaviour and morphology in experimental microbial communities". In: *Proceedings of the Royal Society B: Biological Sciences* 287.1928. ISSN: 14712954. DOI: 10.1098/rspb.2020.0652.

Cairns, Johannes et al. (2020b). "Repeatable ecological dynamics govern the response of experimental communities to antibiotic pulse perturbation". In: *Nature Ecology and Evolution* 4.10. ISSN: 2397334X. DOI: 10.1038/s41559-020-1272-9.

Cingolani, Pablo et al. (2012). "A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff". In: *Fly* 6.2. ISSN: 1933-6934. DOI: 10.4161/fly.19695.

Colclough, Abigail et al. (2019). "Patterns of cross-resistance and collateral sensitivity between clinical antibiotics and natural antimicrobials". In: *Evolutionary Applications* 12.5. ISSN: 17524571. DOI: 10.1111/eva.12762.

Corbett-Detig, Russell B., Daniel L. Hartl, and Timothy B. Sackton (2015). "Natural Selection Constrains Neutral Diversity across A Wide Range of Species". In: *PLoS Biology* 13.4. ISSN: 15457885. DOI: 10.1371/journal.pbio.1002112.

Dentro, Stefan C. et al. (2021). "Characterizing genetic intra-tumor heterogeneity across 2,658 human cancer genomes". In: *Cell* 184.8. ISSN: 10974172. DOI: 10.1016/j.cell.2021.03.009.

Ferrières, Lionel and David J. Clarke (2003). "The RcsC sensor kinase is required for normal biofilm formation in Escherichia coli K-12 and controls the expression of a regulon in response to growth on a solid surface". In: *Molecular Microbiology* 50.5. ISSN: 0950382X. DOI: 10.1046/j.1365-2958.2003.03815.x.

Finn, Robert D. et al. (2000). "Escherichia coli RNA polymerase core and holoenzyme structures". In: *EMBO Journal* 19.24. ISSN: 02614189. DOI: 10.1093/emboj/19.24.6833.

Frazão, N. et al. (Dec. 2022). "Two modes of evolution shape bacterial strain diversity in the mammalian gut for thousands of generations". In: *Nature Communications* 13.1. ISSN: 20411723. DOI: 10.1038/s41467-022-33412-8.

Frickel, Jens, Loukas Theodosiou, and Lutz Becks (Oct. 2017). "Rapid evolution of hosts begets species diversity at the cost of intraspecific diversity". In: *Proceedings of the National Academy of Sciences of the United States of America* 114.42, pp. 11193–11198. ISSN: 10916490. DOI: 10.1073/pnas.1701845114.

Gillis, Sierra and Andrew Roth (Dec. 2020). "PyClone-VI: scalable inference of clonal population structures using whole genome data". In: *BMC Bioinformatics* 21.1, p. 571. ISSN: 1471-2105. DOI: 10.1186/s12859-020-03919-2. URL: https://bmcbioinformatics.biomedcentral.com/articles/10.1186/s12859-020-03919-2.

Good, Benjamin H. et al. (Nov. 2017). "The dynamics of molecular evolution over 60,000 generations". In: *Nature* 551.7678, pp. 45–50. ISSN: 0028-0836. DOI: 10.1038/nature24287. URL: https://www.nature.com/articles/nature24287.

Harris, Charles R. et al. (2020). *Array programming with NumPy.* DOI: 10.1038/s41586-020-2649-2.

Harrison, Ellie et al. (2017). "Conflicting selection alters the trajectory of molecular evolution in a tripartite bacteria–plasmid–phage interaction". In: *Molecular Ecology* 26.10. ISSN: 1365294X. DOI: 10.1111/mec.14080.

Hiltunen, Teppo et al. (Dec. 2018). "Dual-stressor selection alters eco-evolutionary dynamics in experimental communities". In: *Nature Ecology and Evolution* 2.12, pp. 1974–1981. ISSN: 2397334X. DOI: 10.1038/s41559-018-0701-5.

Iwasaki, Hideo et al. (2000). "A KaiC-interacting sensory histidine kinase, SasA, necessary to sustain robust circadian oscillation in cyanobacteria". In: *Cell* 101.2. ISSN: 00928674. DOI: 10.1016/S0092-8674(00)80832-6.

Johnson, Milo S. et al. (2021). "Phenotypic and molecular evolution across 10,000 generations in laboratory budding yeast populations". In: *eLife* 10. ISSN: 2050084X. DOI: 10.7554/eLife.63910.

Katz, Sophia et al. (2021). "Dynamics of Adaptation during Three Years of Evolution under Long-Term Stationary Phase". In: *Molecular Biology and Evolution* 38.7. ISSN: 15371719. DOI: 10.1093/molbev/msab067.

Kimberly K Jefferson (July 2004). "What drives bacteria to produce a biofilm?" In: *FEMS Microbiology Letters* 236.2, pp. 163–173. ISSN: 03781097. DOI: 10.1016/j.femsle.2004.06.005. URL: http://doi.wiley.com/10.1016/j.femsle.2004.06.005.

Lang, Gregory I. and Michael M. Desai (2014). "The spectrum of adaptive mutations in experimental evolution". In: *Genomics* 104.6. ISSN: 10898646. DOI: 10.1016/j.ygeno.2014.09.011.

Lang, Gregory I. et al. (2013). "Pervasive genetic hitchhiking and clonal interference in forty evolving yeast populations". In: *Nature* 500.7464. ISSN: 00280836. DOI: 10.1038/nature12344.

Lässig, Michael, Ville Mustonen, and Armita Nourmohammad (2023). *Steering and controlling evolution — from bioengineering to fighting pathogens.* DOI: 10.1038/s41576-023-00623-8.

Lässig, Michael, Ville Mustonen, and Aleksandra M. Walczak (2017). "Predicting evolution". In: *Nature Ecology and Evolution* 1.3. ISSN: 2397334X. DOI: 10.1038/s41559-017-0077.

Li, Heng and Richard Durbin (2009). "Fast and accurate short read alignment with Burrows-Wheeler transform". In: *Bioinformatics* 25.14. ISSN: 13674803. DOI: 10.1093/bioinformatics/btp324.

Lion, Sébastien (2018). "Theoretical approaches in evolutionary ecology: Environmental feedback as a unifying perspective". In: *American Naturalist* 191.1. ISSN: 15375323. DOI: 10.1086/694865.

Łuksza, Marta and Michael Lässig (2014). "A predictive fitness model for influenza". In: *Nature* 507.7490. ISSN: 14764687. DOI: 10.1038/nature13087.

Maccagni, Alessio and Yvonne Willi (2022). "Trait divergence and trade-offs among Brassicaceae species differing in elevational distribution". In: *Evolution* 76.9. ISSN: 15585646. DOI: 10.1111/evo.14554.

Maddamsetti, Rohan, Richard E. Lenski, and Jeffrey E. Barrick (2015). "Adaptation, clonal interference, and frequency-dependent interactions in a long-term evolution experiment with escherichia coli". In: *Genetics* 200.2. ISSN: 19432631. DOI: 10.1534/genetics.115.176677.

Maharjan, Ram P. et al. (2015). "Simple phenotypic sweeps hide complex genetic changes in populations". In: *Genome Biology and Evolution* 7.2. ISSN: 17596653. DOI: 10.1093/gbe/evv004.

Martin, Marcel (2011). "Cutadapt removes adapter sequences from high-throughput sequencing reads". In: *EMBnet.journal* 17.1. DOI: 10.14806/ej.17.1.200.

Matz, Carsten and Staffan Kjelleberg (2005). "Off the hook - How bacteria survive protozoan grazing". In: *Trends in Microbiology* 13.7. ISSN: 0966842X. DOI: 10.1016/j.tim.2005.05.009.

Meroz, Nittay et al. (Oct. 2024). "Evolution in microbial microcosms is highly parallel, regardless of the presence of interacting species". In: *Cell Systems* 15.10, pp. 930–940. ISSN: 24054712. DOI: 10.1016/j.cels.2024.09.007.

Mollet, Christophe, Michel Drancourt, and Didier Raoult (1997). "rpoB sequence analysis as a novel basis for bacterial identification". In: *Molecular Microbiology* 26.5. ISSN: 0950382X. DOI: 10.1046/j.1365-2958.1997.6382009.x.

Papadopulos, Alexander S.T. et al. (2021). "Rapid Parallel Adaptation to Anthropogenic Heavy Metal Pollution". In: *Molecular Biology and Evolution* 38.9. ISSN: 15371719. DOI: 10.1093/molbev/msab141.

Paterson, Steve et al. (2010). "Antagonistic coevolution accelerates molecular evolution". In: *Nature* 464.7286. ISSN: 00280836. DOI: 10.1038/nature08798.

Pedregosa, Fabian et al. (2011). "Scikit-learn: Machine learning in Python". In: *Journal of Machine Learning Research* 12. ISSN: 15324435.

Rao, Narayana N., María R. Gómez-García, and Arthur Kornberg (2009). *Inorganic polyphosphate: Essential for growth and survival.* DOI: 10.1146/annurev.biochem.77.083007.093039.

Retel, Cas et al. (2019). "The feedback between selection and demography shapes genomic diversity during coevolution". In: *Science Advances* 5.10. ISSN: 23752548. DOI: 10.1126/sciadv.aax0530.

Rutherford, Andrew (2001). *Introducing Anova and Ancova: A GLM Approach.*

Salcedo, Adriana et al. (2020). "A community effort to create standards for evaluating tumor subclonal reconstruction". In: *Nature Biotechnology* 38.1. ISSN: 15461696. DOI: 10.1038/s41587-019-0364-z.

Saxer, Gerda, Michael Doebeli, and Michael Travisano (2010). "The repeatability of adaptive radiation during long-term experimental evolution of Escherichia coli in a multiple nutrient environment". In: *PLoS ONE* 5.12. ISSN: 19326203. DOI: 10.1371/journal.pone.0014184.

Scheuerl, Thomas et al. (2019). "Predator coevolution and prey trait variability determine species coexistence". In: *Proceedings of the Royal Society B: Biological Sciences* 286.1902. ISSN: 14712954. DOI: 10.1098/rspb.2019.0245.

Shao, Kwang Tsao and Robert R. Sokal (1990). "Tree balance". In: *Systematic Zoology* 39.3. ISSN: 1063-5157. DOI: 10.2307/2992186.

Soudi, Shaghayegh et al. (2023). "Repeatability of adaptation in sunflowers reveals that genomic regions harbouring inversions also drive adaptation in species lacking an inversion". In: *eLife* 12. ISSN: 2050084X. DOI: 10.7554/eLife.88604.

Stern, David L. (2013). *The genetic causes of convergent evolution.* DOI: 10.1038/nrg3483.

Strelkowa, Natalja and Michael Lässig (2012). "Clonal interference in the evolution of influenza". In: *Genetics* 192.2. ISSN: 00166731. DOI: 10.1534/genetics.112.143396.

Sule, Preeti et al. (2011). "Regulation of cell division, biofilm formation, and virulence by FlhC in Escherichia coli O157:H7 grown on meat". In: *Applied and Environmental Microbiology* 77.11. ISSN: 00992240. DOI: 10.1128/AEM.00069-11.

Tenaillon, Olivier et al. (2012). "The molecular diversity of adaptive convergence". In: *Science* 335.6067. ISSN: 10959203. DOI: 10.1126/science.1212986.

Wichman, H. A. et al. (1999). "Different trajectories of parallel evolution during viral adaptation". In: *Science* 285.5426. ISSN: 00368075. DOI: 10.1126/science.285.5426.422.