1

Programmed *in vivo* excision of prophage Φ13 is an essential timing device for the *Staphylococcus aureus* infectious process

Olivier Poupel,[1] Gérald Kenanian,[1] Lhousseine Touqui,[2] Charlotte Abrial,[2] Grégory

Jouvion,[3,4] Tarek Msadek[1,6*], Sarah Dubrac[1,5,6*]

[1] Institut Pasteur, Université Paris Cité, CNRS UMR2001, Department of Microbiology,

Biology of Gram-Positive Pathogens Unit, F-75015 Paris, France

[2] Institut Pasteur, Université Paris Cité, Department of Global Health, Cystic Fibrosis

and Bronchial Diseases Unit, F-75015 Paris, France.

[3] Institut Pasteur, Université Paris Cité, Department of Infection and Epidemiology,

Experimental Neuropathology Unit, F-75015 Paris, France

[4] Present address: Ecole Nationale Vétérinaire d'Alfort, Veterinary Histology and

Pathology Unit, Maisons-Alfort, France

[5] Present address: Institut Pasteur, Université Paris Cité, CNRS UMR6047, Department

of Microbiology, Stress Adaptation and Metabolism in Enterobacteria Unit, F-75015

Paris, France

24 [6] These authors jointly supervised this work : Tarek Msadek, Sarah Dubrac.

25 *Address correspondence to Tarek Msadek, tmsadek@pasteur.fr, or Sarah Dubrac

26 sarah.dubrac@pasteur.fr

ABSTRACT Mobile genetic elements play an essential part in the infectious process of major pathogens, yet the role of prophage dynamics in *Staphylococcus aureus* remains poorly understood. Here we studied the impact of the Φ13 *hlb*-converting prophage, whose integration inactivates the *hlb* β-toxin gene, on staphylococcal pathogenesis. We showed that prophage Φ13 is lost in approximately half the bacterial population during the course of infection. Inactivation of the Φ13 *int* recombinase gene, essential for insertion/excision, locked the prophage in the bacterial chromosome, leading to a significant loss of virulence in a murine systemic infection model. In contrast, the non-lysogen strain (ΔΦ13), where the *hlb* beta-hemolysin gene is reconstituted, displayed strongly increased virulence. Accordingly, histopathological analyses revealed more severe nephritis in mice infected with bacteria lacking prophage Φ13 (ΔΦ13), in contrast to infection with the parental strain and particularly with the Δ*int* mutant, which led to less severe renal lesions. Cytokine induction in a human neutrophil model showed significantly increased IL-6 expression following infection with the beta-hemolysin producing strain (ΔΦ13). Our results indicate that programmed *in vivo* excision of the Φ13 prophage is essential for progression of the *Staphylococcus aureus* infectious process, since early excision rapidly kills the host whereas the absence of excision significantly lowers staphylococcal virulence.

IMPORTANCE This study highlights prophage Φ13 excision as a critical factor in *S. aureus* pathogenesis, influencing infection outcomes by balancing rapid host killing with reduced bacterial virulence. This mechanism may represent a bet-hedging strategy in genetic regulation, resulting in a mixed bacterial population capable of rapidly switching between two processes: bacterial colonization and host damage. Unraveling this dynamic opens new possibilities for developing targeted therapies to disrupt or modulate prophage activity, offering a novel approach to mitigating *S. aureus* infections.

INTRODUCTION

Staphylococcus aureus, a major human pathogen, causes severe invasive infectious diseases targeting every type of organ, such as pneumonia, endocarditis, osteomyelitis or septicemia. Yet it is also a commensal, asymptomatically colonizing up to 30% of the human population, with the transition between the two lifestyles poorly understood (1). Mobile genetic elements make up approximately 20% of its genome, leading to variations in gene combinations that can strongly influence *S. aureus* virulence (2). Among these, temperate bacteriophages play an important role in genetic variability. Φ13, a member of the Sa3int Siphoviridae *S. aureus* bacteriophage family, is responsible for a dual phenotypic conversion (3, 4) and is by far the most widespread among human clinical isolates (5). Its chromosomal attachment site (*attB*) lies within the coding sequence of the hemolysin B gene, *hlb*, encoding the neutral sphingomyelinase C β-toxin, such that it acts as a phage regulatory switch (6). Φ13 lysogeny leads to disruption of *hlb* and loss of β-hemolytic activity, with excision reconstituting an intact *hlb* gene (5, 7).

Phage Φ13 also carries the *iec* immune evasion cluster including several genes involved in escape from the host innate immune response: *scn*, *chp* and *sak* (8, 9). Natural mobility has been shown for the Φ13-related ΦSa3mw phage in strain MW2 during murine skin colonization and rabbit invasive infection (10, 11).

We studied the effect of phage Φ13 on *S. aureus* virulence during systemic infection. We show that timing of Φ13 prophage excision is essential for pathogenesis, with early β-hemolysin production leading to a strong increase in virulence and elevated levels of the pro-inflammatory cytokine IL-6.

RESULTS

Mobility dynamics of *Staphylococcus aureus* prophages

S. aureus strain HG001, a derivative of NCTC 8325 (12), carries prophages Φ11, Φ12 and Φ13 (7, 13). Specific oligonucleotide primer pairs were designed to detect the presence or absence of each prophage (Supplementary Table 2 and Supplementary Fig. 1A). Using chromosomal DNA from a closely related strain cured of all phages, SH1000, an amplicon corresponding to the native bacterial locus without Φ13 (oligonucleotide pair OMA7/OMA9, hybridizing on either side of the *attB* site) was detected, but no amplification occurred with oligonucleotides specific for Φ13 insertion (OMA7/OMA8) (Supplementary Fig. 1B). With the HG001 Φ13 lysogen strain, both types of amplicon were generated by PCR, suggesting spontaneous Φ13 excision within the bacterial population (Supplementary Fig. 1B). We quantified relative prophage levels within the bacterial population by qRT-PCR during growth under optimal laboratory conditions (TSB at 37°C with aeration). Our results indicate that prophage Φ11 is highly mobile with approximately 10% of cells lacking the phage whereas Φ12 was much more stable with less than 0.01% of non-lysogen cells (Fig. 1A). Interestingly, the Φ13 prophage was lost in about 0.5% of the population under these growth conditions (Fig. 1A).

Excision of prophage Φ13 is increased *in vivo*

As shown above, prophage Φ13 is spontaneously excised during growth in laboratory conditions. Since this phage carries genes encoding the SAK staphylokinase virulence factor, as well as SCIN and CHIP, involved in escape from the innate immune response (14), we examined Φ13 excision frequency during the infectious process.

We generated a strain carrying the *aad*(9) spectinomycin resistance gene inserted within the Φ13 genome (see Materials and Methods, and Supplementary Table 1 strain ST1387). Four days following sub-lethal (2×10^7 c.f.u.) intravenous inoculation with *S. aureus* strain ST1387, mice were sacrificed and kidneys collected for c.f.u. counts. Out of 25 harvested kidneys, 18 were colonized by a mixed bacterial population with respect to Φ13 lysogeny (Fig.1B). In some mice a single kidney was colonized with a

mixed population with the other only by Φ13 lysogens, suggesting that loss of Φ13 is

a *post facto* event to kidney colonization. The average loss of Φ13 was around 50% in

the 18 kidneys colonized with a mixed bacterial population (Fig. 1C), a significant

increase over the spontaneous excision rate of 0.5% observed during growth *in vitro*.

These results suggest that Φ13 excision can be induced and/or the non-lysogen

population favored during the infectious process under conditions such as those

encountered during kidney colonization.

Excision of prophage Φ13 is essential for the *S. aureus* infectious process

Increased prophage Φ13 excision during infection suggests it may impact virulence.

The Φ13 SAOUHSC-*02293* gene, referred to henceforth as *int*, encodes a protein highly

similar to tyrosine recombinase integrases such as Int of phage lambda (13, 15). We

deleted the entire coding sequence of the Φ13 *int* gene in *S. aureus* strain HG001 to lock

the prophage within the bacterial chromosome. We also constructed a strain where the

entire Φ13 genome is deleted, restoring an intact and functional *hlb* gene (see Materials

and Methods). These mutations had no effect on bacterial growth *in vitro*

(Supplementary Fig. 2). β-hemolytic activity following Φ13 excision was detected on

sheep blood agar plates as a distal cold-inducible halo of partial hemolysis (indicated

by the white arrow in Fig. 2A).

As shown in Fig. 2B, the frequency of Φ13 excision in the Δ*int* mutant is lowered

more than 1000-fold as compared to that of the wild type strain (0.6%), confirming the

essential role of Int in Φ13 excision. Full complementation of the Δ*int* mutant was

obtained by expressing the *int* gene from the Pprot constitutive promoter in the

pMK4Pprot-*int* plasmid (Fig. 2B).

We next examined the effect on *S. aureus* virulence of either the absence of

prophage Φ13 (HG001ΔΦ13 strain, intact *hlb* gene) or the lack of Φ13 excision

(HG001Δ*int* strain, *hlb* gene disrupted). As shown in Fig. 3A, using a systemic model

of murine infection (5 X 10^7 c.f.u. intravenous injection), the absence of Φ13 with a

resulting intact *hlb* gene led to acute virulence with the rapid death of infected animals

(50% mortality 24 h post-infection for the ΔΦ13 strain vs. six days for the wild type

strain), and more than 90% mortality at the end of the assay (ten days post-infection).

Conversely, when Φ13 excision is abolished (Δ*int* strain), *S. aureus* virulence is strongly

reduced compared to the wild type strain (31% mortality at the end of the assay, *vs*
63%). These results indicate that Φ13 excision and its timing are crucial parameters for
the *S. aureus* infectious process.

Φ13 dynamics are involved in organ damage during infection

To understand the effect of Φ13 dynamics on *S. aureus* virulence, we measured
bacterial colonization, and evaluated the kidney lesions of infected mice by
histopathological analysis. Mice were inoculated intravenously with sub-lethal
concentrations of *S. aureus* cells (approximately 2×10^7 c.f.u.). At six days post-infection,
mice were sacrificed and one kidney was fixed for histopathological analysis and the
other homogenized for c.f.u. counts. Kidney bacterial loads for the parental HG001
strain and the HG001ΔΦ13 and HG001Δ*int* mutants were not significantly different
(approximately 5×10^7 c.f.u. per kidney; Supplementary Fig. 3). However,
histopathological analysis revealed nephritis (very probably embolic nephritis) with
significantly varying degrees of severity between mice infected with the parental
HG001 strain or the ΔΦ13 and Δ*int* mutants. At six days post-infection, mice inoculated
with the wild type strain generally displayed mild to moderate nephritis (5/6 mice) or
multifocal inflammatory infiltrates (macrophages and other mononucleated cells) in
the renal cortex (1/6 mouse) (Fig. 3B1 and 3B2). In contrast, the ΔΦ13 strain leads to
more severe nephritis (5/6 mice), with neutrophilic inflammation forming multifocal
abscesses, extension to tubules and interstitium, and ischemic necrosis of the papillae
(Fig. 3B3 and 3B4). Mice infected with the Δ*int* strain displayed less severe lesions, with
mononuclear cell infiltrates mostly in the cortex (4/6 mice) or mild to moderate (1/6
mouse) or marked (1/6 mouse) nephritis (Fig. 3B5 and 3B6).

These results show that Φ13 dynamics and correlated β-hemolysin production
are not essential for host colonization but instead are key factors in host organ damage
during infection.

Φ13 excision triggers host inflammatory response in infected THP-1 cells

Given the major impact of Φ13 excision on virulence and inflammation, we tested its
effect on induction of the cellular inflammatory response using human monocytic
THP-1 cells. We compared the bacterial internalization capacity of THP-1 cells, the

intracellular viability of bacterial cells and the induction of inflammatory markers
following infection with the parental HG001 strain and the ΔΦ13 and Δ*int* mutants.
THP-1 differentiated macrophages were very efficient for *S. aureus* phagocytosis
regardless of prophage content, with nearly 100% of bacteria internalized with a MOI
(Multiplicity of Infection) of 10 after 2 hours (Supplementary Fig. 4A). Intracellular
survival was not significantly different for the three strains with approximately 10%
survival after 24 h post infection and less than 5% after 48 h (Supplementary Fig. 4B).
To measure induction of the inflammatory response, duplicate aliquots of infected
THP-1 cells were frozen 24 h post-infection, treated to recover cellular RNA and
induction of inflammatory cytokine genes measured by qRT-PCR. We measured
induction of three pro-inflammatory cytokines previously shown to be induced by
S. aureus infection, TNFα, IL-1β and IL-6 (16). Although there was no significant
difference in induction levels of TNFα and Pro-IL-1β for the three strains, IL-6 gene
expression was strongly increased when THP-1 cells were infected with the ΔΦ13
strain as compared to the HG001 strain and the Δ*int* mutant (Fig. 4). Thus, lysogeny
by Φ13 and the concomitant loss of β-hemolytic activity appear to down-regulate the
inflammatory response through lower induction of the pro-inflammatory cytokine IL-
6.
197
198   **Impact of *hlb* on *S. aureus* virulence**
199
Taken together the results shown above suggest a major role of β-hemolysin
production in the increased virulence of the ΔΦ13 strain. To test this hypothesis, we
compared the virulence of the ΔΦ13 strain, which produces β-hemolysin, with the
virulence of the ΔΦ13Δ*hlb* strain in which the *hlb* gene has also been deleted. As shown
in Fig. 5A, 44% of mice infected by the ΔΦ13Δ*hlb* strain survived at nine days post-
infection, whereas survival of mice infected by the strain producing β-hemolysin
(ΔΦ13) was only 14%, indicating that it is the major factor responsible for the effect of
Φ13 excision on *S. aureus* virulence. To test the contribution of Φ13 in virulence, we
constructed a strain where both Φ13 and an intact *hlb* gene are present (Δ*int hlb*
complemented strain; see Materials and Methods, and Supplementary Table 1). As
shown in Fig. 5B, mouse mortality caused by the ΔΦ13 strain producing β-hemolysin
but lacking the Φ13 prophage was essentially the same as that following infection by

the Δ*int hlb* complemented strain (with both the Φ13 prophage and a functional *hlb*

gene). This result confirms the major role of Φ13 dynamics in *S. aureus* virulence by

controlling timing of β-hemolysin production during the infectious process.

DISCUSSION

Staphylococcus aureus is a highly successful human pathogen with a sophisticated arsenal of virulence factors whose production is tightly controlled by multiple regulatory networks, such as two-component systems (17). Further fine-tuning is provided by mobile genetic elements such as pathogenicity islands (SaPI), plasmids, transposons, staphylococcal cassette chromosomes (SCC) and bacteriophages conferring additional virulence factors or antibiotic resistance genes (18). Prophages are widely distributed amongst *S. aureus* strains. Among these, the Φ13 *S. aureus* prophage is inserted within the *hlb* gene encoding β-hemolysin and carries at least three genes involved in host interactions and escape from the immune response, known as the immune evasion cluster (IEC): *sak*, encoding staphylokinase, involved in converting host plasminogen to active plasmin, *scn*, encoding a staphylococcal complement inhibitor, and *chp*, encoding a neutrophil and monocyte chemotaxis inhibitory protein (8). The Φ13 *hlb*-converting prophage family is present in approximately 90% of *S. aureus* clinical isolates (8).

β-hemolysin is a sphingomyelinase that can lyse proliferating human lymphocytes, contributing to the host immune evasion (19). Independently of its pore forming activity, β -hemolysin has a biofilm ligase effect by forming a nucleoprotein matrix both in vitro and during infectious endocarditis (20, 21). It has also been shown that Hlb favors mouse skin colonization (11). We noted that *S. aureus* cells isolated from infected mice frequently displayed β-hemolytic activity due to *in vivo* excision of the Φ13 prophage. This prompted us to examine the effect of phage mobility on host pathogen interactions. Our results indicate that mobility of this prophage is required for full virulence of *S. aureus* as part of the time-course of the infectious process. We identified and inactivated the potential integrase gene of this prophage, effectively locking the prophage within the *S. aureus* genome by preventing its excision, and showed that this leads to strongly decreased virulence, with 70% of the mice still surviving after 10 days. Conversely, a strain lacking the Φ13 prophage exhibited greater virulence than the wild type strain, particularly evident in the rapid mortality of infected mice, with 50% succumbing within the first day of infection, compared to six days for the wild-type strain. These Φ13-associated virulence traits are not correlated to differences in the strain's ability to colonize the host, as bacterial loads in the infected kidneys were similar regardless of the strain's lysogenic status. However,

kidney damage following infection was strongly associated with the presence of Φ13,
showing a clear gradient where a lower proportion of lysogenic bacteria resulted in
more severe tissue lesions. In a human monocyte (THP-1) infection model used to
assess inflammatory response, we found that the non-lysogenic strain significantly
increased the production of the pro-inflammatory cytokine IL-6. IL-6 triggers the
release of acute-phase response proteins into the bloodstream, leading to a systemic
inflammatory response (22). It is also considered a key biomarker for predicting poor
outcomes in sepsis patients and is a promising target for inhibitors to help mitigate the
adverse effects of inflammation (16, 23). The induction of IL-6 by the non-lysogenic
strain may be linked to its heightened virulence and the significant neutrophil
recruitment observed at the inflammation site, as evidenced by the histopathological
analysis of infected kidneys.

The exact trigger causing excision of Φ13 during infection remains to be
established. Interestingly, our observations show that after intravenous infection,
kidneys of the same mouse can be colonized differently, one kidney by a lysogenic
population, while the other harbors a mixed population. This suggests that Φ13
excision is likely induced in the kidneys following colonization rather than during the
initial spread of bacteria through the bloodstream. Interestingly, a previous study
using a rabbit infection model found that bacteria producing β-hemolysin emerged
during organ colonization, but this was not observed in the bacterial population
recovered from the bloodstream (10). Stress conditions activating the SOS response are
known to induce prophage excision. However, despite many attempts, we were not
able to induce Φ13 excision from the bacterial chromosome through treatment with
Mitomycin C as SOS-inducing agent. Similar results were previously described with a
Newman strain carrying the closely related *hlb*-converting prophage, ΦNM3 (24).
More recently, it has been demonstrated that exposure to oxidative stress (such as
hydrogen peroxide) and biofilm growth conditions promote the excision of ΦSA3mw
(25). This may correlate with our in vivo observations, i.e. that phage induction occurs
after organ colonization, under biofilm development conditions.

Epidemiological studies have shown that *hlb*-converting prophages are more
commonly found in strains colonizing humans (from nasal swabs) than in invasive
strains (isolated from blood) (3). Additionally, Φ13-derivative carriers are significantly
more frequent in human *S. aureus* strains than in animal strains (26). These *hlb*-
converting phages not only inactivate *hlb* but also carry virulence genes, including an

immune evasion cluster (IEC) (8). The Sak, SCIN, and CHIPS determinants encoded
by this cluster play crucial roles in modulating neutrophil chemotaxis, bacterial
phagocytosis, and killing (14). However, their activity is highly specific, primarily
targeting the human immune system (27-29). This specificity of the Φ13-derivative IEC
in counteracting the human immune response likely contributes to the greater
prevalence of *hlb*-converting prophages in human *S. aureus* strains.

Interestingly, we demonstrated using an infection mouse model, that the *hlb* gene
is the major determinant responsible for the increased virulence observed in the
absence of the Φ13 phage. When Hlb is produced, no effect of Φ13 alone was observed
(Fig. 5B), likely because the strain is already highly virulent due to constant Hlb
production. However, in the absence of *hlb*, the presence of Φ13 led to lowered mouse
mortality (Fig. 3A and 5A). While the immune evasion factors encoded by Φ13 have
been described as human-specific, it cannot be excluded that they may also exhibit
some activity during mouse colonization. Alternatively, Φ13 may encode
uncharacterized proteins that could modulate *S. aureus* virulence.

Excision of Φ13 and the chromosomal rearrangement reconstituting an intact *hlb*
gene can be viewed as a timing device, ensuring that the bacteria first colonize the host,
benefiting from the immune evasion cluster proteins encoded by the prophage, with
production of β-hemolysin and host tissue damage occurring once the bacteria have
reached a threshold level, in agreement with the temporal control of *Staphylococcus*
aureus virulence factors (30). One of the first examples of prophage excision as a timing
device for a developmental process is the *skin* element whose excision leads to a
chromosomal rearrangement reconstituting the gene encoding the Sigma K sigma
factor, essential for *Bacillus subtilis* sporulation (31, 32).

As demonstrated in this study and supported by the literature, the
heterogeneity within *S. aureus* populations with respect to *hlb*-converting prophages
appears to be a common trait. This could be seen as a bet-hedging strategy, allowing
the bacteria to retain the ability to produce both immune evasion proteins (in the
lysogenic population) and the potent Hlb cytotoxin (in the non-lysogenic cells).
Additionally, since both β-hemolysin and the immune evasion cluster proteins
encoded by the prophage are secreted, it is plausible that functional complementation
may occur within a mixed population of lysogenic and non-lysogenic bacteria. Similar
dual strategies have been observed before, where bistability leads to cell-to-cell
variation in gene expression (33, 34). This mixed population with distinct genetic

content would be well-equipped both to resist host defenses during colonization and
to cause cellular damage.

MATERIALS AND METHODS

Bacterial strains and growth media

Escherichia coli K12 strain DH5α™ (Invitrogen) was used for cloning experiments. *Staphylococcus aureus* strain HG001, derived from strain NCTC 8325 by repairing the mutated *rsbU* gene (12), was used for genetic and functional studies. Plasmids were first transformed into the restriction deficient *S. aureus* strain RN4220 before introduction into the HG001 or derivative strains. *E. coli* strains were grown in LB medium with ampicillin (100 μg/mL) added when required. *S. aureus* strains and plasmids used in this study are listed in Supplementary Table 1. *S. aureus* strains were grown in Trypticase Soy Broth (TSB; Difco) supplemented with chloramphenicol (10 μg/mL) or erythromycin (1 μg/mL) when required. *E. coli* and *S. aureus* strains were transformed by electroporation using standard protocols (35) and transformants were selected on LB or Trypticase Soy Agar (TSA; Difco) plates, respectively, with the appropriate antibiotics.

DNA manipulations

Oligonucleotides used in this study were synthesized by Eurofins Genomics and are listed in Supplementary Table 2. *S. aureus* chromosomal DNA was isolated using the MasterPure™ Gram-positive DNA purification Kit (Epicentre Biotechnologies). Plasmid DNA was isolated using a QIAprep Spin Miniprep kit (Qiagen) and PCR fragments were purified using the Qiaquick PCR purification kit (Qiagen). T4 DNA ligase and restriction enzymes, PCR reagents and Q5 high-fidelity DNA polymerase (New England Biolabs) were used according to the manufacturer's recommendations. Nucleotide sequencing of plasmid constructs was carried out by Eurofins Genomics .

Plasmids and mutant strain construction

Construction of the ΔΦ13 mutant strain was carried out by cloning an 1800 bp DNA fragment centered around the *attB* site, generated by PCR using oligonucleotides OSA418 and OSA419 and chromosomal DNA from the SH1000 strain (a NCTC8325-4 derivative cured for prophages), between the *Bam*HI and *Eco*RI sites of the pMAD vector (36), to give plasmid pMAD-Φ13. For deletion of the Φ13 integrase (*int*; *saouhsc-*

02239), DNA fragments corresponding to the *int* upstream and downstream regions were generated using oligonucleotides OSA411/OSA412 (900 bp) and OSA413/OSA414 (820 bp) cloned between the pMAD *Bam*HI and *Nco*I sites. The *aad*(9) spectinomycin resistance gene (37) was inserted in the center of the 23 nt intergenic region between the convergent Φ13 *saouhsc-02232* and *saouhsc-02233* genes, to avoid any polar effects on transcription. DNA fragments corresponding to the regions upstream and downstream from the insertion site were generated by PCR using primers OSA508/OSA509 and OSA510/OSA511, respectively, and for the *aad*(9) cassette primers OSA299/OSA470 and plasmid pIC333 (38) as the DNA matrix. The DNA fragments were cloned into pMAD using the *Bam*HI/*Sal*I sites for the upstream fragment, *Sal*I/*Nco*I for the *aad*(9) cassette, and the *Nco*I/*Bgl*II sites for the downstream fragment, generating plasmid pMAD-Φ13*aad*(9). For deletion of *hlb*, we generated upstream and downstream fragments by PCR using oligonucleotides OSA442/OSA443 (884 bp) and OSA444/OSA445 (889 bp) respectively. These fragments were introduced into pMAD between the *Bam*HI and *Eco*RI sites, thus generating the pMAD-*hlb* plasmid. To introduce a functional *hlb* allele into the Φ13 lysogenic strain, a pMAD-*hlb*comp plasmid was constructed to place the intact *hlb* gene upstream from the Φ13 genome. A PCR fragment including the *hlb* upstream region and the *hlb* intact gene (1760 bp) was generated using the OSA418 and OSA573 primers and the SH1000 genomic DNA as matrix and it was cloned into pMAD using the *Bam*H*I and *Sal*I restriction sites. A second PCR fragment corresponding to the part of Φ13 adjacent to *att*L was generated with the OSA574-575 primers and cloned between the *Sal*I and *Bgl*II restriction sites of pMAD.

Plasmids were introduced by electroporation into *S. aureus* strain RN4220, and transformants selected at 30°C on TSA plates containing erythromycin and 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-Gal) (100 µg/mL). Plasmids were then introduced into strain HG001 and integration and excision performed as previously described (36), yielding strains HG001ΔΦ13, HG001Δ*int*, and HG001Φ13-aad(9) and into strain HG001ΔΦ13 for pMAD-*hlb* and HG001Δ*int* for pMAD-*hlb*comp to give strains HG001Δ*int hlb*comp, HG001ΔΦ13Δ*hlb*.

Plasmid pMK4Pprot-*int* was used for complementation of the Δ*int* mutant and constructed by cloning the entire *int* coding sequence (amplicon OSA446/OSA447) under the control of the constitutive P*prot* promoter (39).

Prophage content measurement by quantitative real time PCRs (qRT-PCRs)

Strains were grown overnight in TSB at 37°C, with aeration. Cells were pelleted by centrifugation (2 min, 20,800 *g*) from 2 mL culture samples and genomic DNA extracted using the MasterPure™ Gram Positive DNA Purification Kit (Tebu-Bio).

Oligonucleotides were designed with the BEACON Designer 4.02 software (Premier Biosoft International, Palo Alto, CA) to synthesize 100-200 bp amplicons (see Supplementary Table 2). Quantitative real-time PCRs (qRT-PCRs) and critical threshold cycles (CT) were performed and determined using the SsoFast™ EvaGreen Supermix (Bio-Rad, Hercules, CA). All assays were performed using quadruplicate technical replicates, and repeated with three independent biological samples, with one representative experiment shown. Prophage loss frequency was calculated using the formula: $1/(1+2^{\Delta Ct})$ derived from the ΔCt method (40).

Hemolysis detection

Bacteria were grown overnight in TSB at 37°C with aeration. Culture aliquots (3 μl) were spotted on Columbia plates containing 5% sheep blood (COS, Biomerieux Craponne). Plates were incubated for 24h at 37°C and transferred to 4°C for an additional 24h to for detection of β-hemolysis (41).

THP-1 infection experiments

THP-1 monocytes were maintained as previously described (42). Differentiation and adherence were obtained by incubation with Phorbol 12-Myristate 13-Acetate (PMA, 50 ng/mL) during 4 days at 37°C in a 5%CO_2 atmosphere. Bacterial uptake experiments and survival assays were carried out as previously described (43). Briefly, bacteria were grown in TSB until OD_{600nm} = 1. Cultures were washed in PBS and adjusted to the desired inoculum in RPMI 1640 medium (Gibco), and c.f.u. counts determined by plating serial dilutions on TSA plates. Differentiated THP-1 macrophages were incubated with bacteria (MOI = 10) in RPMI 1640 at 37°C with 5% CO2 for 1 h to allow bacterial phagocytosis. They were then washed with RPMI and incubated in RPMI-10% Fetal Calf Serum with streptomycin (100 μg/mL), penicillin (100 U/mL), and gentamycin 40 μg/mL to eliminate extracellular bacteria. At the

indicated times, infected macrophages were washed once with RPMI and lysed by
incubation in ice-cold water for 15 min and c.f.u. counts were determined by plating
serial dilutions on TSA plates. For cytokine gene expression studies by qRT-PCR, THP-
1 cells were frozen at -20°C 24h post-infection. Total RNA was extracted using a
RNeasy mini kit (Qiagen, CA, USA). Cytokine gene expression was analysed by qRT-
PCR using the StepOnePlus real-time PCR system (Applied Biosystems). Primers used
for qRT-PCR were TNFα, IL6, pro-IL1beta and HPRT as the reference gene. The qRT-
PCRs were performed by Taqman and results presented as relative quantification of
gene expression using the $2^{-\Delta\Delta CT}$ method.
431

432 **Animal experiments**
433

Seven-week-old female *RjOrl:SWISS* mice (Janvier Labs) were inoculated
intravenously with the specified *S. aureus* strains. To analyse Φ13 excision during the
infectious process, mice were infected with a sublethal dose (2×10^7 c.f.u.) of the Φ13-
aad(9) strain. Mice were sacrificed four days post-infection and kidneys harvested and
homogenized for c.f.u. determination following dilution and plating on TSA and TSA
with spectinomycin (100 μg/mL). The rate of prophage excision was calculated as
[(c.f.u. on TSA) - (c.f.u. on TSA+spectinomycin)]/(c.f.u. on TSA).

Strain HG001 and the Δ*int*, ΔΦ*13*, ΔΦ*13*Δ*hlb* and Δ*int hlb*comp mutants were used
for survival studies. Groups of seven mice were injected in the caudal lateral vein with
5×10^7 c.f.u. per mouse in 0.2 mL. Survival was monitored daily for 9 days following
infection and three independent experiments were carried out. To measure kidney
colonization and histopathological analysis, mice were infected intravenously with a
sublethal dose (1×10^7 c.f.u.) of the HG001 parental and Δ*int* and ΔΦ*13* mutant strains
(six mice for each strain). At six days post-infection, kidneys were collected and for
each mouse, bacterial c.f.u. counts were carried out using one kidney and
histopathological observations were done on the other kidney. For histopathology,
kidneys were removed and immediately fixed in 10% neutral buffered formalin. After
48 h of fixation, the kidneys were longitudinally sectioned through the hilus in order
to obtain the renal cortex, medulla, crest, and pelvis on the same section. These
samples were embedded in paraffin, and 4-μm sections were cut and stained with
hematoxylin and eosin (lesion description) and Gram (bacterial detection).
455

Ethics statement

Animal experiments were carried out at the Institut Pasteur according to European

Union guidelines for handling of laboratory animals

(http://ec.europa.eu/environment/chemicals/lab_animals/index_en.htm).

Animals were monitored daily with a particular attention for any signs of distress.

Protocols used were approved by the Institut Pasteur ethics committee (dap160057).

ACKNOWLEDGMENTS

We are grateful to Patrick Trieu-Cuot, in whose lab this work was carried out, and Shaynoor Dramsi for helpful discussion. All authors have read and agreed to the submitted version of the manuscript.

FUNDING

This work was supported by funds from Institut Pasteur, CNRS, and Vaincre La Mucoviscidose Association (Grant «Phage induction, a novel essential factor for *Staphylococcus aureus* infections?»).

AUTHOR CONTRIBUTIONS

Conceptualization: Sarah Dubrac, Lhousseine Touqui, Tarek Msadek.

Data curation: Olivier Poupel, Sarah Dubrac, Lhousseine Touqui, Tarek Msadek.

Formal analysis: Sarah Dubrac, Grégory Jouvion, Lhousseine Touqui, Tarek Msadek.

Funding acquisition: Sarah Dubrac, Lhousseine Touqui, Tarek Msadek.

Investigation: Olivier Poupel, Gérard Kénanian, Charlotte Abrial, Sarah Dubrac.

Methodology: Olivier Poupel, Sarah Dubrac.

Project administration: Tarek Msadek, Sarah Dubrac.

Resources: Tarek Msadek, Sarah Dubrac.

Supervision: Tarek Msadek, Sarah Dubrac.

Validation: Tarek Msadek, Sarah Dubrac.

Writing – original draft: Tarek Msadek, Sarah Dubrac.

Writing – review & editing: Tarek Msadek, Sarah Dubrac.

ADDITIONAL FILES

The following material is available online.

Supplemental Material

Supplemental Figures and Tables. Fig. S1–S4 and Tables S1–S2.

REFERENCES

1. Piewngam P, Otto M. 2024. *Staphylococcus aureus* colonisation and strategies for decolonisation. Lancet Microbe 5:e606-e618.

2. Fitzgerald JR, Sturdevant DE, Mackie SM, Gill SR, Musser JM. 2001. Evolutionary genomics of *Staphylococcus aureus*: insights into the origin of methicillin-resistant strains and the toxic shock syndrome epidemic. Proc. Nat. Acad. Sci. USA: 98:8821-6.

3. Goerke C, Pantucek R, Holtfreter S, Schulte B, Zink M, Grumann D, Broker BM, Doskar J, Wolz C. 2009. Diversity of prophages in dominant *Staphylococcus aureus* clonal lineages. J. Bacteriol. 191:3462-8.

4. Xia G, Wolz C. 2014. Phages of *Staphylococcus aureus* and their impact on host evolution. Infect. Genet. Evol. 21:593-601.

5. Goerke C, Wirtz C, Fluckiger U, Wolz C. 2006. Extensive phage dynamics in *Staphylococcus aureus* contributes to adaptation to the human host during infection. Mol. Microbiol.61:1673-85.

6. Feiner R, Argov T, Rabinovich L, Sigal N, Borovok I, Herskovits AA. 2015. A new perspective on lysogeny: prophages as active regulatory switches of bacteria. Nat. Rev. Microbiol. 13:641-50.

7. Iandolo JJ, Worrell V, Groicher KH, Qian Y, Tian R, Kenton S, Dorman A, Ji H, Lin S, Loh P, Qi S, Zhu H, Roe BA. 2002. Comparative analysis of the genomes of the temperate bacteriophages phi 11, phi 12 and phi 13 of *Staphylococcus aureus* 8325. Gene 289:109-18.

8. van Wamel WJ, Rooijakkers SH, Ruyken M, van Kessel KP, van Strijp JA. 2006. The innate immune modulators staphylococcal complement inhibitor and chemotaxis inhibitory protein of *Staphylococcus aureus* are located on beta-hemolysin-converting bacteriophages. J. Bacteriol. 188:1310-5.

9. Rohmer C, Wolz C. 2021. The role of hlb-converting bacteriophages in *Staphylococcus aureus* host adaption. Microb. Physiol. 31:109-122.

10. Salgado-Pabon W, Herrera A, Vu BG, Stach CS, Merriman JA, Spaulding AR, Schlievert PM. 2014. *Staphylococcus aureus* beta-toxin production is common in strains with the beta-toxin gene inactivated by bacteriophage. J. Infect. Dis. 146:784-92.

11. Katayama Y, Baba T, Sekine M, Fukuda M, Hiramatsu K. 2013. Beta-hemolysin promotes skin colonization by *Staphylococcus aureus*. Journal of Bacteriology 195:1194-203.

12.   Herbert S, Ziebandt AK, Ohlsen K, Schafer T, Hecker M, Albrecht D, Novick R,
Gotz F. 2010. Repair of global regulators in *Staphylococcus aureus* 8325 and
comparative analysis with other clinical isolates. Infect. Immun. 78:2877-89.

13.   Gillaspy AF, Worrell V, Orvis J, Roe BA, Dyer DW, Iandolo JJ. 2006. The
Staphylococcus aureus NCTC 8325 Genome, p 381-412. *In* Fischetti VA, Novick
RP, Ferretti JJ, Portnoy DA, Rood JI (ed), Gram-Positive Pathogens, 2nd ed.
ASM Press, Washington, DC.

14.   Rooijakkers SH, van Kessel KP, van Strijp JA. 2005. Staphylococcal innate
immune evasion. Trends Microbiol. 13:596-601.

15.   Nunes-Duby SE, Kwon HJ, Tirumalai RS, Ellenberger T, Landy A. 1998.
Similarities and differences among 105 members of the Int family of site-specific
recombinases. Nucleic Acids Res. 26:391-406.

16.   van den Berg S, Laman JD, Boon L, ten Kate MT, de Knegt GJ, Verdijk RM,
Verbrugh HA, Nouwen JL, Bakker-Woudenberg IA. 2013. Distinctive cytokines
as biomarkers predicting fatal outcome of severe *Staphylococcus aureus*
bacteremia in mice. PLoS One 8:e59107.

17.   Haag AF, Bagnoli F. 2017. The role of two-component signal transduction
systems in *Staphylococcus aureus* virulence regulation. Curr. Top. Microbiol.
Immunol. 409:145-198.

18.   Lindsay JA. 2010. Genomic variation and evolution of *Staphylococcus aureus*. Int
J. Med. Microbiol. 300:98-103.

19.   Huseby M, Shi K, Brown CK, Digre J, Mengistu F, Seo KS, Bohach GA,
Schlievert PM, Ohlendorf DH, Earhart CA. 2007. Structure and biological
activities of beta toxin from *Staphylococcus aureus*. J. Bacteriol. 189:8719-26.

20.   Herrera A, Vu BG, Stach CS, Merriman JA, Horswill AR, Salgado-Pabon W,
Schlievert PM. 2016. *Staphylococcus aureus* beta-toxin mutants are defective in
biofilm ligase and sphingomyelinase activity, and causation of infective
endocarditis and sepsis. Biochemistry 55:2510-7.

21.   Huseby MJ, Kruse AC, Digre J, Kohler PL, Vocke JA, Mann EE, Bayles KW,
Bohach GA, Schlievert PM, Ohlendorf DH, Earhart CA. 2010. Beta toxin
catalyzes formation of nucleoprotein matrix in staphylococcal biofilms. Proc.
Nat. Acad. Sci. USA: 107:14407-12.

22.   Tanaka T, Narazaki M, Kishimoto T. 2014. IL-6 in inflammation, immunity, and
disease. Cold Spring Harb. Perspect. Biol. 6:a016295.

23.   Cao YZ, Tu YY, Chen X, Wang BL, Zhong YX, Liu MH. 2012. Protective effect
570   of Ulinastatin against murine models of sepsis: inhibition of TNF-alpha and IL-
571   6 and augmentation of IL-10 and IL-13. Exp. Toxicol. Pathol. 64:543-7.

572   24.   Bae T, Baba T, Hiramatsu K, Schneewind O. 2006. Prophages of *Staphylococcus*
573   *aureus* Newman and their contribution to virulence. Mol. Microbiol. 62:1035-47.

574   25.   Tran PM, Feiss M, Kinney KJ, Salgado-Pabon W. 2019. øSa3mw prophage as a
575   molecular regulatory switch of *Staphylococcus aureus* β -toxin production. J.
576   Bacteriol. 201:e00766-18

577   26.   Verkaik NJ, Benard M, Boelens HA, de Vogel CP, Nouwen JL, Verbrugh HA,
578   Melles DC, van Belkum A, van Wamel WJ. 2011. Immune evasion cluster-
579   positive bacteriophages are highly prevalent among human *Staphylococcus*
580   *aureus* strains, but they are not essential in the first stages of nasal colonization.
581   Clin. Microbiol. Infect. 17:343-8.

582   27.   de Haas CJ, Veldkamp KE, Peschel A, Weerkamp F, Van Wamel WJ, Heezius
583   EC, Poppelier MJ, Van Kessel KP, van Strijp JA. 2004. Chemotaxis inhibitory
584   protein of *Staphylococcus aureus*, a bacterial antiinflammatory agent. J. Exp. Med.
585   199:687-95.

586   28.   Kwiecinski J, Josefsson E, Mitchell J, Higgins J, Magnusson M, Foster T, Jin T,
587   Bokarewa M. 2010. Activation of plasminogen by staphylokinase reduces the
588   severity of *Staphylococcus aureus* systemic infection. J. Infect. Dis. 202:1041-9.

589   29.   Rooijakkers SH, Ruyken M, Roos A, Daha MR, Presanis JS, Sim RB, van Wamel
590   WJ, van Kessel KP, van Strijp JA. 2005. Immune evasion by a staphylococcal
591   complement inhibitor that acts on C3 convertases. Nat. Immunol. 6:920-7.

592   30.   Novick RP. 2003. Autoinduction and signal transduction in the regulation of
593   staphylococcal virulence. Mol. Microbiol. 48:1429-49.

594   31.   Stragier P, Kunkel B, Kroos L, Losick R. 1989. Chromosomal rearrangement
595   generating a composite gene for a developmental transcription factor. Science
596   243:507-12.

597   32.   Kunkel B, Losick R, Stragier P. 1990. The *Bacillus subtilis* gene for the
598   development transcription factor sigma K is generated by excision of a
599   dispensable DNA element containing a sporulation recombinase gene. Genes
600   Dev. 4:525-35.

601   33.   Dubnau D, Losick R. 2006. Bistability in bacteria. Mol. Microbiol. 61:564-72.

602   34.   Veening JW, Smits WK, Kuipers OP. 2008. Bistability, epigenetics, and bet-
603   hedging in bacteria. Annu. Rev. Microbiol. 62:193-210.

604   35.   Sambrook J, Fritsch EF, Maniatis T. 1989. Molecular cloning : a laboratory
605   manual, second edition. Cold Spring Harbor Laboratory, Cold Spring Harbor,
606   N. Y.

607   36.   Arnaud M, Chastanet A, Debarbouille M. 2004. New vector for efficient allelic
608   replacement in naturally nontransformable, low-GC-content, gram-positive
609   bacteria. Appl. Environ. Microbiol. 70:6887-91.

610   37.   Murphy E. 1985. Nucleotide sequence of a spectinomycin adenyltransferase
611   AAD(9) determinant from *Staphylococcus aureus* and its relationship to AAD(3")
612   (9). Mol. Gen. Genet. 200:33-39.

613   38.   Steinmetz M, Richter R. 1994. Easy cloning of mini-Tn*10* insertions from the
614   *Bacillus subtilis* chromosome. J. Bacteriol. 176:1761-3.

615   39.   Archambaud C, Gouin E, Pizarro-Cerda J, Cossart P, Dussurget O. 2005.
616   Translation elongation factor EF-Tu is a target for Stp, a serine-threonine
617   phosphatase involved in virulence of *Listeria monocytogenes*. Mol. Microbiol.
618   56:383-96.

619   40.   Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using
620   real-time quantitative PCR and the 2(-Delta Delta C(T)) method. Methods
621   25:402-8.

622   41.   Haque RU, Baldwin JN. 1964. Types of hemolysins produced by *Staphylococcus
623   aureus*, as determined by the replica plating technique. J. Bacteriol. 88:1442-7.

624   42.   Stokes RW, Doxsee D. 1999. The receptor-mediated uptake, survival, replication,
625   and drug sensitivity of *Mycobacterium tuberculosis* within the macrophage-like
626   cell line THP-1: a comparison with human monocyte-derived macrophages.
627   Cell. Immunol. 197:1-9.

628   43.   Soutourina O, Dubrac S, Poupel O, Msadek T, Martin-Verstraete I. 2010. The
629   pleiotropic CymR regulator of *Staphylococcus aureus* plays an important role in
630   virulence and stress response. PLoS Pathog. 6:e1000894.

 FIGURE LEGENDS

 Figure 1: Prophage Φ13 in vitro and in vivo. A) Prophage excision frequencies in

 strain HG001. qRT-PCR quantification of Φ11, 12, and 13 prophage content in *S. aureus*

 strain HG001 grown in TSB at 37°C with aeration. Specific primer pairs are described

 in Supplementary Table 3. The results are the mean of 4 technical replicates and are

 representative of 3 independent biological replicates.

 B) Quantification of Φ13 lysogens in bacterial populations from infected kidneys. Mice

 were infected from the caudal lateral vein with a sub-lethal dose (2 X 10^7 c.f.u.) of *S.*

 aureus strain HG001 Φ13::*aad*(9). Mice were sacrificed four days post-infection and

 kidneys harvested for c.f.u. counts on TSA and TSA with spectinomycin. 25 infected

 kidneys were examined and the proportion of homogeneous lysogen population (less

 than 0.5% non-lysogen) and heterogeneous bacterial population with a mix of lysogen

 and non-lysogen bacteria is shown.

 C) Proportion of non-lysogen bacteria per kidney in organs infected with a mixed

 population of lysogenic and non-lysogenic bacteria.

 Figure 2: Characterization of the *S. aureus* Δ*int* and ΔΦ13 mutants. Strains were

 grown overnight at 37°C with aeration. A) 3 μl of the bacterial cultures was spotted on

 sheep blood agar plates and incubated consecutively for 24 h at 37°C and 24 h at 4°C

 to detect hemolytic activities. The white arrow indicates the β-hemolytic halo

 characteristic of bacteria lacking prophage Φ13. B) Prophage Φ13 excision frequency

 detected by qRT-PCR using oligonucleotide pairs OMA7-OMA8 and OMA7-OMA9

 specific for the presence or absence of Φ13, respectively.

 Figure 3: Impact of prophage Φ13 and *hlb* on *S. aureus* virulence. A) Kaplan-Meier

 survival curves of *RjOrl:SWISS* mice infected intravenously with *S. aureus* HG001 wild

 type strain and the Δ*int* and ΔΦ13 mutants (5 X 10^7 c.f.u./injection). A total of 14 mice

 were used in each group in two independent experiments. *, *P*<0.05 and **, *P*<0.01

 (Wilcoxon test). For each strain, the genotype is indicated followed by the relevant

 genetic context (presence or absence of prophage Φ13 and a functional *hlb* gene). B)

 Histopathological kidney analyses 6 days post-infection by intravenous inoculation

 with sub-lethal bacterial loads (2 X 10^7 c.f.u./injection) of the *S. aureus* HG001 wild

type (1 and 2), Δ*int* (3 and 4) and ΔΦ13 (5 and 6) strains. Most (5/6) mice infected with
the wild type strain displayed mild to moderate nephritis, with neutrophilic
infiltrates/abscesses (black arrowheads) of varying sizes in the cortex (1), medulla, and
papilla (2). Infection with strain ΔΦ13 led to more severe nephritis (5/6 mice), with
multifocal abscesses (3 and 4, black arrowheads), extension to tubules and interstitium,
and ischemic necrosis of the papilla (4, black arrow). Most mice (4/6) infected with the
Δ*int* strain displayed less severe lesions, with mononuclear cell infiltrates most often
in the cortex (5, not seen at a low magnification) but also in the papilla (6, arrowhead).
673

Figure 4: Impact of Φ13 on cytokine induction by THP-1 neutrophils. PMA-activated
THP-1 cells were infected with the HG001 wild type strain or the ΔΦ13 and Δ*int*
mutants with a MOI=10. After 24 H of infection, cytokine gene expression was
determined by qRT-PCR performed with cDNA synthesized from RNA of THP-1 cells
infected by the indicated strains (dark grey) or non-infected control (light grey).
Expression levels are given in arbitrary units normalized against that of HPRT. Each
experiment was carried out 6 times in two independent series (means +/- SD). ***:
P<0.005 (Student's *t*-test).
682

Figure 5: Impact of *hlb* on *S. aureus* virulence. Kaplan-Meier survival curves of
RjOrl:SWISS mice infected intravenously with *S. aureus* HG001 wild type strain and
the Δ*int hlb*$_{comp}$, ΔΦ13 and ΔΦ13Δ*hlb* mutants (5 X 10^7 c.f.u./injection). A total of 14 mice
were used in each group in two independent experiments. *, *P*<0.05 and **, *P*<0.01
(Wilcoxon test). For each strain, the genotype is indicated followed by the relevant
genetic context (presence or absence of prophage Φ13 and a functional *hlb* gene).
A) Role of *hlb* in virulence by comparing infection with the HG001 strain and the ΔΦ13
and ΔΦ13Δ*hlb* derivatives.
B) Role of Φ13 in virulence by comparing infection with the HG001 strain and the
Δ*int hlb*$_{comp}$ (dashed line) and ΔΦ13 derivatives.

Figure 1

A.

B.

Colonization by Φ13 lysogenized *S. aureus*

Colonization by *S. aureus* with or without Φ13

C.

Figure 2

A.

B.

Figure 3

A.

B.

WT	ΔΦ13	Δint

Figure 4

Figure 5

A.

B.