A systematic analysis of in-source fragments in LC-MS metabolomics

Yuanye Chi[1], Joshua M. Mitchell[1] and Shuzhao Li[1,2*]

[1]The Jackson Laboratory for Genomic Medicine, 10 Discovery Drive, Farmington, CT 06032,

USA

[2]University of Connecticut School of Medicine, Farmington, CT 06032, USA

*Corresponding author, E-mail: shuzhao.li@jax.org

The "dark matter" of metabolomics refers to the large number of unidentified features in

metabolomic studies, mostly from mass spectrometry (MS) based analysis (deSilva2015;

David2021; Giera2024). The topic is pertinent to the analytical coverage of small molecules in

biomedical research (Kind2009; Uppal2016), approaches to metabolite annotation

(Domingo2018; Chaleckis2019; Metz2025), mapping reaction pathways (Zamboni2015) and the

promise of applying metabolomics and exposomics to precision medicine (Wishart2016;

Vermeulen2020). The number of unidentified features is not a direct account of number of

compounds, as a metabolite can have isotopologues, adducts and fragments that are measured

in the same data (Mahieu2017; Wang2019; Li2023a). Giera et al (2024) recently reported that

in-source fragments (ISFs) accounted for over 70% of MS/MS features in METLIN, one of the

leading spectral databases, suggesting that ISFs could be a significant portion of the "dark

matter". Since the reference spectra in METLIN are based on chemical standards, we examine

here the LC-MS (liquid chromatography coupled mass spectrometry) metabolomics from

biological samples, which are the most relevant in biomedical investigations.

LC-MS metabolomics data from 45 studies were retrieved from public repositories (orbital mass

analyzer, electron spray ionization (ESI) in positive or negative mode, **Table S1**). For

consistence of data analysis, only studies of human plasma or serum samples are included, and

large datasets are down selected to about 100 samples, because low-frequency features

cumulate by sample numbers. We use the asari software (Li2023b) for processing raw data to

features of unique m/z (mass-to-charge ratio) and retention time (RT). Asari is designed with

explicit data models that report all features above a quality threshold (signal-to-noise ratio, or

SNR, of 2, peak shape of 0.5). The median number of reported features in these studies is

around 50,000, of which about a quarter to a third are considered high-quality features (SNR >

5, peak shape > 0.9). Because all organic molecular species contain both ^{12}C and ^{13}C (carbon

13, often under detection limit for low-abundance compounds) natural isotopologues, the

presence of $^{13}C/^{12}C$ pair indicates high-confidence detection of a compound. The numbers of

$^{13}C/^{12}C$ pairs are typically between 2,000 to 3,000 in these metabolomic datasets (**Figure 1a**).

Signal intensity in a typical study follows a power law distribution (**Figure 1b**).

Typically, ESI is used in MS to avoid breaking up the molecules. In contrast, MS/MS techniques

fragment molecules intentionally by a collision energy, and the fragments serve as a fingerprint

of the molecule. If ISFs of a molecule are present in LC-MS data, they must a) have the same

retention time as the intact molecules, and b) have m/z values that are similar to those in the

corresponding MS/MS spectra. In Giera et al (2024), MS/MS data at zero collision energy were

used to mimic MS data. We take a similar strategy here to compare an MS/MS database

(MoNA2024) to the LC-MS metabolomics data, requiring both the precursor and at least one

MS/MS fragment to be observed in MS data (**Figure 1c**). We parsed out 13,672 compounds

from MoNA MS/MS spectra. They typically match to 5~10,000 features by precursor m/z values

(**Figure 1d**). Of these matched precursor ions, about 7% have at least one matched MS/MS

fragment in the elution window for positive ESI data, about 5% for negative ESI (**Figure 1e**).

The elution window here is defined as two standard deviation of RT differences of all $^{13}C/^{12}C$

pairs. These results indicate that ISFs impact less than 10% of LC-MS features. The results are

expected to include false positives (thus overestimation), as a) MS/MS fragments generated by

higher collision energies are not expected to be in the LC-MS data; and b) matched features are

not necessarily generated by ISF.

A limitation of the above approach is that MoNA database does not fully cover the LC-MS

metabolomics data, though the number of matched MS/MS fragments relative to the matched

precursor ions should indicate the prevalence of ISFs. A more systematic analysis can be

performed using the khipu algorithm, which constructs ion patterns de novo into empirical

compounds, in a process called pre-annotation (Li2023a, Mitchell2024). In short, a khipu aligns

coeluting ions on a grid (a tree in computer term) of isotopologues and modifications, where

modifications include adducts and fragments, as exemplified in **Figure 2a**. This approach

depends on the list of mass differences underlying the ion patterns. We systematically calculate

all common mass differences in these LC-MS datasets (**Table S2-5**). Not surprisingly, the m/z

difference between $^{13}C/^{12}C$ isotopologues is by far the most frequent observation in both ESI+

and ESI- data. The top frequent mass differences, excluding known isotopic and adduct

patterns, can be considered as candidates for ISFs (also referred as neutral losses, **Figure**

2b,c). Examining the RT shift of these candidate ISFs shows that the majority have identical RT

as their molecular ions, confirming coelution in chromatography (**Figure 2d**).

To assess the impact of ISFs on full metabolomic profiles, we compare the pre-annotations

without and with consideration of ISFs. Without ISFs, the khipu algorithm is applied to the LC-

MS datasets using a conservative set of isotopologues and adducts (Methods), to produce a set

of khipus and singletons per dataset. A singleton is a feature that has no known relationship to

other features. The numbers of khipus and singletons can be impacted by ISFs, but ISFs do not

impact relationships from isotopologues and adducts. Next, to consider ISFs, the candidate

ISFs (from **Figure 2b,c**) are searched between khipus, and between singletons and khipus. The

comparison between two steps reveals that in the ESI+ data, about 500 khipus and 700

singletons can be explained by the candidate ISFs (**Figure 2e**), which correspond to about 8%

of khipus and 1% of all features, respectively (**Figure 2f**). In the ESI- data, the candidate ISFs

explain about 5% of khipus and 1% of extra features (**Figure 2e,f**). While these results do not

rule out additional ISFs among singleton features, singletons are of lower intensity and expected

to contain even less percentage of fragments. The RT shift distributions in **Figure 2d** bear two

shoulder peaks, which suggest that a subset of matched ISFs is not truly coeluting and thus not

fragments but different compounds. This implies that the results in **Figure 2e,f** are an

overestimation.

The khipu approach allows a full calculation of pre-annotation of all datasets now. A

comprehensive set of isotopologues, adducts and ISFs are used in a new round of analysis

(**Table S6**). The results show that about 16,000 features are explained by khipus in ESI+ or

ESI- data, which correspond to 5~6,000 unique empirical compounds (**Figure 3a**). These

numbers are in line with high-quality features in **Figure 1a**, suggesting that we are close to a full

explanation of them. To test the dependency of pre-annotation on ion abundance, we split the

features in each dataset by intensity into quartiles and visualize the pre-annotation for each

quartile. The results show that about 60% of features in the top quartiles are explained by

khipus in ESI+ or ESI- data, while the pre-annotated percentage is smaller in lower quartiles

(**Figure 3b**). Plotting the result using full quantile range returns a similar conclusion, where pre-

annotation reaches 80% at the top end of quantiles (**Figure S1a**). Within the top quartile, the

singletons have significantly lower intensity than M0 features (**Figure S1b**), corroborating this

observation that pre-annotation is bound by abundance. To test the impact of peak alignment on

the results, we selected one random sample (thus avoiding peak alignment) from each dataset

to repeat the full data processing and pre-annotation, which returned similar results (not shown).

In summary, we have employed two approaches to search ISFs in a sizable collection of LC-MS

datasets. Both resulted in less than 10% of features being explained by ISFs. Additional

analysis using 16 studies on time-of-flight platforms returned similar results (**Tables S1**, **S7**). If

we assume ESI always fragments a small percentage of all analytes, our results are consistent

with that by Giera et al (2024): in the analysis of authentic chemical standards, most ISFs are

detected by mass spectrometer because there is little background and in the analysis of

biological samples, the ISFs are outcompeted by more abundant molecular species in the

complex matrix.

Overall, these results represent a broad survey of the current metabolomics data landscape: the

khipu-based pre-annotation already explains the majority of abundant features. The less

abundant features should not be considered as mystery - their analysis is just limited by

concentration and ionization efficiency. Therefore, the "dark matter" of metabolomics is largely

explainable. This means that the overwhelming majority of LC-MS metabolomic features from

human blood samples are real compounds (not necessarily of biological origin). The differences

between studies are often affected by experimental methods and conditions, which should be a

focus of future computational data processing and annotation.

Data and Code Availability:

All asari processed data are available at: https://zenodo.org/records/14541717. The lists of mass patterns for isotopologues, adducts and ISFs are included in the mass2chem software package, freely available at https://github.com/shuzhao-li-lab/mass2chem. All data analysis in this work is provided as Jupyter notebooks at https://github.com/shuzhao-li-lab/in_source_fragments_serum.

Author Contributions

Project design and writing: SL. Data collection and processing: YC. Data analysis: SL, YC. Software development: JM, YC, SL. All authors participated in and approved the final manuscript.

Acknowledgements

We thank Dr. Shujian Zheng for discussion and review of this work. This work was in part supported by NIH grant R01 AI149746 (NIAID) and ARPA-H award D24AC00345.

Supplemental Information

Table S1. List of datasets used in this work, include 45 Orbitrap studies and 16 TOF studies.

Table S2. Most frequent mass differences in Orbitrap ESI+ data.

Table S3. Most frequent mass differences in Orbitrap ESI- data.

Table S4. Most frequent mass differences in TOF ESI+ data.

Table S5. Most frequent mass differences in TOF ESI- data.

Table S6. Isotopologues, adducts and ISFs used for extended khipu search in Figure 3.

Table S7. Results of features explained in TOF LC-MS metabolomics data.

Figure S1. Dependency of pre-annotation on feature abundance.

References

Chaleckis, R., Meister, I., Zhang, P. and Wheelock, C.E., 2019. Challenges, progress and promises of metabolite annotation for LC–MS-based metabolomics. *Current opinion in biotechnology*, *55*, pp.44-50.

David, A., Chaker, J., Price, E.J., Bessonneau, V., Chetwynd, A.J., Vitale, C.M., Klánová, J., Walker, D.I., Antignac, J.P., Barouki, R. and Miller, G.W., 2021. Towards a comprehensive characterisation of the human internal chemical exposome: Challenges and perspectives. *Environment International*, *156*, p.106630.

Domingo-Almenara, X., Montenegro-Burke, J.R., Benton, H.P. and Siuzdak, G., 2018. Annotation: a computational solution for streamlining metabolomics analysis. *Analytical chemistry*, *90*(1), p.480.

Giera, M., Aisporna, A., Uritboonthai, W. and Siuzdak, G., 2024. The hidden impact of in-source fragmentation in metabolic and chemical mass spectrometry data interpretation. *Nature metabolism*, pp.1-2.

Huber, F., van der Hooft, J. J. J., Spaaks, J. H., Diblen, F., Verhoeven, S., Geng, C., Meijer, C., Rogers, S., Belloum, A., Spreeuw, H., Villanueva Castilla, E. M., Ashouritaklimi, K., de Jonge, N., Hecht, H., Skoryk, M., Ahmad, Z., Aleksander Piciga, Rutz, A., & Pollmann, J. (2024). matchms (0.27.0). Zenodo. https://doi.org/10.5281/zenodo.12706344

Kind, T., Scholz, M. and Fiehn, O., 2009. How large is the metabolome? A critical analysis of data exchange practices in chemistry. *PloS one*, *4*(5), p.e5440.

Li, S., Siddiqa, A., Thapa, M., Chi, Y. and Zheng, S., 2023b. Trackable and scalable LC-MS metabolomics data processing using asari. *Nature Communications*, *14*(1), p.4113.

Li, S. and Zheng, S., 2023a. Generalized tree structure to annotate untargeted metabolomics and stable isotope tracing data. *Analytical chemistry*, *95*(15), pp.6212-6217.

Mahieu, N. G.; Patti, G. J., Systems-Level Annotation of a Metabolomics Data Set Reduces 25 000 Features to Fewer than 1000 Unique Metabolites. *Anal Chem* **2017,** *89* (19), 10397-10406.

Mitchell, J.M., Chi, Y., Thapa, M., Pang, Z., Xia, J. and Li, S., 2024. Common data models to streamline metabolomics processing and annotation, and implementation in a Python pipeline. *PLOS Computational Biology*, *20*(6), p.e1011912.

MassBank of North America (MoNA) [retrieved Feb 8, 2024]; Available from: https://mona.fiehnlab.ucdavis.edu/.

Uppal, K., Walker, D.I., Liu, K., Li, S., Go, Y.M. and Jones, D.P., 2016. Computational metabolomics: a framework for the million metabolome. *Chemical research in toxicology*, *29*(12), pp.1956-1975.

Vermeulen, R., Schymanski, E.L., Barabási, A.L. and Miller, G.W., 2020. The exposome and health: Where chemistry meets biology. *Science*, *367*(6476), pp.392-396.

200
201 Wang, L.; Xing, X.; Chen, L.; Yang, L.; Su, X.; Rabitz, H.; Lu, W.; Rabinowitz, J. D., Peak
202 Annotation and Verification Engine for Untargeted LC-MS Metabolomics. *Anal Chem* **2019,** *91*
203 (3), 1838-1846.
204
205 Wishart, D.S., 2016. Emerging applications of metabolomics in drug discovery and precision
206 medicine. *Nature reviews Drug discovery*, *15*(7), pp.473-484.
207
208 Zamboni, N., Saghatelian, A. and Patti, G.J., 2015. Defining the metabolome: size, flux, and
209 regulation. *Molecular cell*, *58*(4), pp.699-706.
210
211

Figures

Figure 1. Search of in-source fragments in 45 LC-MS studies from Orbitrap platforms using MS/MS spectra.

a. Across the 22 studies using positive ionization, the median number of features is 50,771, the median number was 18,117, and the number of matched $^{13}C/^{12}C$ pairs was 2,176. Similar numbers were observed across the 23 negative ionization studies with 48,212 features, 11,783 good features, and 3,008 $^{13}C/^{12}C$ pairs. Good features are defined as having SNR > 5 and peak shape > 0.9 when fitting to a gaussian curve.

b. Example distribution of feature intensity (dataset ST002200_RPpos_17min_B3_ppm5_3422144, total 50,647 features). Quartiles are marked by dashed lines.

c. Example of matching MS/MS and MS spectra for searching ISFs. Top: MS/MS from MoNA. Bottom: sample CHCL3_20min_B from dataset MTBLS1465_HILICpos__ppm5_3505731, MS scan number 1408.

d. Absolute numbers of features matched to MoNA MS/MS spectra across the 45 LC-MS studies, using positive and negative ionization respectively. In each boxplot, 1st column is the number of features matching a precursor m/z value (P), the 2nd column the subset of P with match to at least one MS/MS fragment (Y), and the 3rd column number of all matched MS/MS fragments per study.

e. Conversion of **d** to percentages: P over total features (T), Y over T and Y over P.

233   Figure 1

234
235

Figure 2: Most frequent neutral losses account for a small subset of LC-MS features.

a. Example of khipu pre-annotation of isotopologues, adducts and fragments in a tree structure,
with m/z values labeled for each ion. The mass spectrum is shown on the right. Data from
sample 214 in study ST002112_RPpos_B3_ppm5_356194, RT around 619.49 second. This
compound matches to glycochenodeoxycholic acid.

b, **c**. Most frequence delta m/z values in positive and negative ionization datasets, respectively.
These exclude obvious isotopologues and adducts, thus as candidates for fragments (neutral
losses).

d. Example distribution of retention time shift (in seconds) associated with each delta m/z in **b**.

e, **f**. The numbers and percentages of khipus and additional features explained by candidate
fragments, in positive and negative ionization datasets, respectively.

Figure 2

248
249

Figure 3: Khipu based pre-annotation explains most of the abundant features.

a. Numbers of pre-annotated features cross studies, median value 16,574 and 16,296 for positive and negative ionization datasets, respectively. They correspond to 5,539 and 5,937 unique khipus, respectively.

b. Percentage distribution of pre-annotated features in each intensity quartile (exemplified in **Figure 1b**) cross studies, shown as violin plots, for positive and negative ionization datasets respectively. The pie charts show median percentage of contributions from M0, other isotopologues, in-source fragments and singletons. Adducts are included in khipus, and singletons are the unexplained features.

261   Figure 3

262
263

Figure S1: Dependency of pre-annotation on feature abundance.

a. Percentage of pre-annotated features versus feature quantile by peak area (1%-99% with

stepwise as 1%) from 45 Orbitrap studies, for positive and negative ionization datasets

respectively.

b. Example of abundance difference between [M0+H]+ features and singletons from study

ST001237_HILICpos_B2_ppm5_3524314, p-value < 1E-25 by Student t-test.

Methods

Data Retrieval and Processing

A total of 61 public metabolomics studies were downloaded from Metabolomics Workbench (metabolomicsworkbench.org) and MetaboLights (www.ebi.ac.uk/metabolights/). All studies were based on human serum or plasma samples, totaling 3482 samples. Of the 61 total studies, 45 were collected using orbitrap-type mass analyzers and 16 using time-of-flight mass analyzers. The breakdown by chromatography and ionization type are noted in Table S1 but all permutations of HILIC, RP, ESI+ and ESI- were represented in all analyzer subsets. Large studies were down selected to between 100~120 samples.

The raw files were converted to centroided mzML files using ThermoRawFileParser v1.3.1 for Orbitrap data or msConvert v3 for ToF data. Data preprocessing was performed using Asari (v 1.13.1), yielding feature tables with m/z, retention time, peak shape, signal-to-noise ratio, chromatographic selectivity, detection count and intensity values per study. For all analyses, the full feature table was used which enforces the default feature quality filters (signal-to-noise ratio, or SNR, of 2, and a peak shape > 0.5 defined as goodness of fit to a gaussian model). Orbitrap studies were processed using the autoheight option enabled, while ToF data was processed using a mass accuracy of 25 ppm, a min_peak_height of 1000, a cal_min_peak_height as 3e4, and a min_intensity_threshold of 500. Mass calibration was performed according to the default parameters in Asari.

The MoNA MS/MS library were downloaded from https://mona.fiehnlab.ucdavis.edu/ (Feb 8, 2024) and the resulting spectra were deduplicated as follows using utilities from MatchMS (Huber2024). First all spectra for the same precursor inchikey were identified to yield a spectral cluster. Within each spectral cluster, in the second step, the pairwise cosine similarity is calculated for all pairs. The sum of the cosine similarity score weighted by the number of matched peaks for all such comparisons per spectrum is then calculated. Lastly, the spectrum with the largest sum score for that inchikey is then selected as the exemplar for that inchikey and all others are discarded. All collision energy spectra were considered during deduplication with the assumption that the most common fragments should be frequently shared across collision energies.

Search of m/z patterns

The isotopic pairs, $^{13}C/^{12}C$ features, are defined by annotation of matched retention time, mass

distance of 1.003355, and intensity of ^{13}C feature under 50% of that of ^{12}C feature. The RT

differences of all isotopic pairs in each dataset were calculated, and their standard deviation

was used to define the RT coelution window per study (one standard deviation on either side of

the molecular ion).

To compare MS/MS data to LC-MS data, our deduplicated MoNA MS/MS library is used (13973

precursors in positive and 9184 in negative). A 5-ppm mass tolerance was used for both

precursor and MS/MS peak matching with features in retention time window. Only MS/MS

peaks whose normalized intensity was higher than 0.1 was considered. To compare common

mass difference to LC-MS data, in-source fragment candidates are manually selected from

common mass differences, and the minimum of 5 ppm of m/z value or 0.0005 as absolute value

was used for mass delta matching. Results here are searched on the top 5 MS/MS fragments,

while using top 10 fragments led only to minor increase of matches (not shown).

Calculation of common mass differences

For the molecular ion in each khipu, the m/z differences to all other features in the elution

window were calculated and counted in a histogram (bin size 0.0001 for Orbitrap data, 0.0005

for TOF data). The histogram was smoothed, and peak values were selected by a threshold

(100 for Orbitrap data, 20 for TOF data). The top 20 mass differences (deltas) were selected per

ionization mode to be used as candidate in-source fragments.

Pre-annotation using Khipu

Khipu performs pre-annotation by assigning co-eluting features to adduct and isotopologue

relations using a generic tree structure based on *a priori* mass delta patterns (Li2023a). The

minimum of 5 ppm of m/z value or 0.0005 as absolute value was used for mass delta matching.

The conservative parameters used in Figure 2 are as follows: in positive ionization mode,

isotope patterns 13C/12C at m/z 1.0034, 13C/12C × 2 at m/z 2.0067, and 37Cl/35Cl at m/z

1.9970; adducts Na/H at m/z 21.9819, ACN at m/z 41.0265, NaCOOH at m/z 67.9874, K/H at

m/z 37.9559, and CH3OH at m/z 32.0262. In negative ionization mode, isotope patterns

13C/12C at m/z 1.0034, 13C/12C × 2 at m/z 2.0067, 37Cl/35Cl at m/z 1.9970, and 32S/34S at

m/z 1.9958; adducts Na/H at m/z 21.9819, NaCOOH at m/z 67.9874, and NaCH2COOH at m/z

82.0030. The parameters for comprehensive khipu analysis in Figure 3 are given in Table S6.

339