# Towards a comprehensive view of the pocketome universe – biological implications and algorithmic challenges.

Hanne Zillmer[1] and Dirk Walther[1*]

[1]Max Planck Institute of Molecular Plant Physiology

Am Mühlenberg 1

14476 Potsdam-Golm

Germany

*Corresponding author

Tel.: +49-331-5678 108

Email: walther@mpimp-golm.mpg.de

# 1 Abstract

With the availability of reliably predicted 3D-structures for essentially all known proteins, characterizing the entirety of compound-binding sites (binding pockets on proteins) has become a possibility. The aim of this study was to identify and analyze all compound-binding sites, i.e. the pocketomes, of eleven different species from different kingdoms of life to discern evolutionary trends as well as to arrive at a global cross-species view of the pocketome universe. Computational binding site prediction was performed on all protein structures in each species as available from the AlphaFold database. The resulting set of potential binding sites was inspected for overlaps with known pockets and annotated with regard to the protein domains in which they are located. 2D-projection plots of all pockets embedded in a 128-dimensional feature space, and characterizing them with regard to selected physicochemical properties, provide informative, global pocketome maps that unveil differentiating features between pockets. Our study revealed a sub-linear scaling law of the number of unique binding sites relative to the number of unique protein structures per species. Thus, as proteome size increased during evolution and therefore potentially diversified, the number of distinct binding sites, reflecting potentially diversifying functions, grew less than proportionally. We discuss the biological significance of this finding as well as identify critical and unmet algorithmic challenges.

**Authors' summary**

The function of proteins is governed by specific interactions with other molecules, notably small

molecules (compounds, such as metabolites). The precise nature of the protein-compound

interaction, and thus, the associated function, is determined by the stereochemical and

physicochemical properties of the sites at which the interaction occurs (binding pockets). Thus,

novel functions (binding of novel compounds) generally require the emergence of new binding

sites. With the recent breakthroughs in protein structure prediction, the complete set of protein

structures has become available. This allowed us to apply computational binding site predictions

and to investigate the entirety of all pockets (the "pocketome") across eleven different species

from all kingdoms of life, and to study the relationship between emergence of novel bindings sites

with increasing sizes of proteomes, the set of all protein structures in a species. We found that

there is a sub-linear relationship between unique pockets and unique protein structures,

suggesting that during evolution, functional diversity shows evidence of saturation, which is in line

with other reports to that effect, but is approached here from the perspective of compound-binding

specificities. Our study constitutes the first large-scale investigation of the pocketomes based on

the now available high-confidence protein structures.

# Introduction

To perform their biochemical function, proteins interact with various entities such as small molecules, referred to generally as compounds or specifically in biological systems as metabolites. These interactions occur at cavities on the protein surface known as "binding pockets" or "compound binding sites". Binding pockets have been the subject of both experimental and bioinformatic research for many years, with the aim to assess similarities between them, to identify potential ligands for specific binding sites, or all potential target proteins for a given compound. A binding pocket is typically defined as all residues within a certain distance to the ligand's heavy atoms in a given protein-ligand complex [1]. As this definition requires detailed 3D-structural information of protein-compound complexes, i.e. the position and conformation of the ligand needs to be known, computational approaches have been developed to predict binding pockets from protein 3D-structures alone, i.e. in the absence of an actually bound ligand. Some of these approaches are geometry-based [2,3], while others are energy-based [4,5] or machine-learning-based [6]. With the recent advances in protein structure prediction and the release of the AlphaFold database, these approaches allow to predict and analyze potential binding sites for whole proteomes of the covered 48 different and evolutionarily diverse species, such as *Arabidopsis thaliana* or human. Thus, a comprehensive and systematic view of compound binding sites on protein surfaces has now become possible.

In this study, we aimed to systematically characterize, cluster, and compare complete sets of protein-compound binding sites. As in the case of small compound ligands, binding sites are frequently referred to as binding pockets, we essentially took aim at describing whole "pocketomes" of different species. By taking a cross-species view and including species of different evolutionary history (bacteria, fungi, plants, mammals), we wished to understand how the pocketomes evolved along with different metabolic needs and organismic capabilities. In particular, as species differ markedly with regard to their proteomes, notably by the number of encoded proteins, we wished to discern the corresponding scaling law that relates the number of unique pockets, reflecting on the potential to bind a corresponding number of unique compounds, to the number of proteins encoded in the respective species. Such an analysis may shed light on

the trends of evolution with regard of "investment", meaning whether binding capacities scale

proportionally with proteome size, or emerged at increased or decreased rates, with all possible

outcomes being relevant for an understanding of selection processes in evolution.

Each species has a unique set of proteins and, thus, binding pockets that is involved in

metabolic interactions as well as signaling processes. These binding pockets collectively define

the species' pocketome. Thus far, only a few studies on pocketomes have been performed. These

include studies on the pocketome of certain enzyme classes, such as human kinases [7] and G-

protein-coupled receptors [8], the pocketome of known 3D structures [9–11], and the species-

specific pocketomes, such as *Mycobacterium tuberculosis* [12] and human [13,14]. Most of these

studies focused on applications in the field of biomedical research and the druggability of binding

sites or discussed differences between drug-protein vs. metabolite-protein binding events [15,16].

An inter-species perspective that exploits the new scale at which structural data are available and

that aims at deducing evolutionary trends has not been presented yet.

To computationally compare and cluster binding sites, they have to be represented in a

machine-readable format, i.e. in a numeric format. Binding site descriptions have been developed

at different levels of detail, ranging from coarse-grained representations where each amino acid

residue is represented by a single atom, e.g., the $C_\alpha$ atom, to full-atom representations where all

atoms of the binding site are used [17]. Other approaches represent binding pockets not based

on residue positions but rather as a point cloud [3]. In addition to 3D coordinates, binding pocket

representations often include additional information, such as the type of residues, thereby aiming

to capture the physicochemical properties. Other methods, such as DeeplyTough [18], employ

deep learning techniques to learn a representation for a protein binding pocket as a numeric

vector of a chosen dimensionality (referred to as "embedding").

Applying established computational methods that allow for an analysis at different levels

of structural detail, our study revealed a sub-linear scaling law of the number of unique binding

sites relative to the number of unique protein structures per species. Thus, larger proteomes

harbor less than proportionally more different binding sites than species with smaller proteomes.

We also identified and discuss challenges encountered during the process of performing a global,

multi-species comparative computational pocketome analysis, such as the speed and specificity

of binding site comparison algorithms, and accounting for co-factors, structural flexibility, and

protein-protein interactions.

# Materials and Methods

**Dataset**

Predicted 3D-protein structures of 11 different species covering all kingdoms of life (excluding

*Archaea*) were downloaded from the AlphaFold (AF) database v4 (https://alphafold.ebi.ac.uk/)

(Table 1) [19]. All predicted structures of all species were considered for compound binding site

predictions. Approximately 34% of the investigated protein structures contain at least one

transmembrane segment. To ensure a high quality of proteins with alpha-helical transmembrane

regions, evaluation files from the TmAlphaFold Transmembrane Protein Structure Database

(https://tmalphafold.ttk.hu/) were downloaded [20]. All proteins passing less than six quality

checks (as reported by the database) were excluded from further analysis, as they are classified

as poor (if five quality checks are passed) and failed (if less than five quality checks are passed)

models. In addition, all proteins in the AF database that are marked as fragments in UniProt and

all proteins with fewer than 100 amino acid residues were excluded from further analysis. With

regard to the latter, P2Rank has been reported to underpredict on proteins shorter than 100 amino

acid residues [21].

Additional information, such as known binding sites and domains, was obtained from

UniProt [22] and InterPro [23], respectively. The assigned FoldSeek (FS) cluster for each protein

was obtained from https://cluster.foldseek.com/ [24].

All ligands found in known (annotated) pockets were mapped to their respective compound

classes using ChEBI ontologies [25]. Only pockets with four or more residues were kept. Inter-

chain pockets were split using an all-against-all distance matrix of $C_\alpha$ atoms and distance

threshold of 12 Å. This distance matrix was interpreted as an adjacency matrix, and distinct

components were determined using scipy [26]. The ChEBI ID for each ligand was used as input

for a recursive search on ChEBI ontologies. If a ChEBI ID could not be mapped to any of the

compound classes, it was assigned as "other compound". An overview of all compound classes

considered in this study is listed in Table 2.

**Compound binding site/ Pocket detection**

Potential binding sites were predicted for each investigated protein using P2Rank in conservation

mode [6]. P2Rank is a state-of-the-art, machine-learning-based program that was trained to

predict potential binding sites via the solvent-accessible surface (SAS) of a protein. After

generating a regularly distributed set of points lying on the SAS, feature descriptors are

calculated , and a ligandability score is predicted for each of the points. Points with a high

ligandability are then clustered, and the respective amino acid residues form the predicted binding

sites. These pockets are ranked by the cumulative ligandability scores.

In this study, P2Rank was used with its standard settings, as those do not make use of

the Debye-Waller B-factor as a feature. For AF-predicted structures, the predicted local distance

difference test (pLDDT) is provided in place of the B-factor, which indicates the confidence of the

predicted position for each residue and should not be mistaken as B-factors.  Pockets with a mean

pLDDT < 70 or a mean predicted aligned error (PAE) > 10 Å were excluded from further analysis,

as they are found either in low-quality or intrinsically disordered regions. Both thresholds were set

based on visual inspection of the obtained distributions (Fig 2).

The overlap between computationally predicted and the respective known (annotated)

pockets from the UniProt database was determined for each protein. If there was any overlap

between the amino acid residues of a known and a predicted pocket, the respective known pocket

was marked as "found" and the predicted pocket was mapped to the respective compound class

corresponding to the ligand that is known to bind to that pocket.

Each compound class was tested for enrichment/ depletion in found pockets compared to

all other compound classes using Fisher's exact test. Odds ratios were computed to signify the

fold increase or decrease of detecting sites for the respective class from a 2x2 contingency table

of observations (found/not-found x is-class/other-class). Fisher exact test p-values for each

compound class were corrected for multiple testing using the Benjamini-Yekutieli procedure to

control the false discovery rate.

**Pocket comparison and dimensionality reduction**

Many different binding site comparison tools have been published in recent years. A subset of them has been compared on the ProSPECCTs datasets in [27] and [28]. In this study, we decided to use two different binding site comparison tools, ProBis [29] and DeeplyTough [18].

Other binding site comparison tools were not suitable for this study for several reasons. Tools, such as SiteMine [28] or IChem [30], depend on internal binding site prediction tools, thus making it difficult to compare their performance to other binding site descriptors for predicted binding sites as the predicted sites potentially differ between different prediction tools. In addition, most of them were implemented to compare one pocket to a set of known pockets and not to perform all-against-all comparison. This results in a high average pairwise runtime, ranging between 8 and 28 s, e.g., for SiteHopper [31] and SMAP [32], making them unsuitable for large-scale comparisons [28]. Lastly, the descriptors for some tools, e.g., KRIPO [33] are based on pharmacophore features that are defined for a protein-ligand complex. This makes them unsuitable for predicted binding sites, where the ligand is not known, which is the case for most pockets.

ProBiS [29] compares protein binding sites by detecting common surface structural patches. The solvent accessible surface atoms within the binding site are converted to overlapping subgraphs, where each vertex represents one functional group with physicochemical properties, such as hydrogen bond donor or hydrogen bond acceptor. The subgraphs are then compared by constructing a modular product graph containing only vertices with matching physicochemical properties. For each product graph, a maximum clique is identified that corresponds to the substructure common to the two compared binding sites. This approach allows ProBiS to detect local structure matches of the two compared binding sites and outputs easily interpretable alignments to the user. In this study, ProBiS was used at the level of within-species pocket comparisons. As the identification of maximum cliques is computationally expensive, ProBiS is not suitable for inter-species comparisons (11 different species) of all predicted binding sites across all species.

To enable a fast comparison of different pockets via a simple Euclidean distance metric and visualizations using established dimensionality reduction methods, DeeplyTough [18] was

employed for the large-scale, inter-species comparisons. DeeplyTough consists of a steerable

CNN that is trained to separate dissimilar pockets by a certain margin during the training process.

In the first step, a 24 x 24 x 24 Å grid is centered on the defined binding site and each binding site

is converted to a 4D tensor with eight channels each one indicating the presence or absence of

an atom or pharmacophoric features. These 4D tensors serve as input to a steerable CNN with

six convolutional layers. To achieve rotational invariance, the authors used data augmentation

and introduced an additional stability loss. This loss drives the descriptors of the original binding

site and its randomly rotated copy towards being the same. The output of DeeplyTough consists

of 128-dimensional embedding vectors that can easily be compared, clustered, and projected to

lower numbers of dimensions.

To plot the resulting 128-dimensional embedding vectors in 2D, different dimensionality

reduction techniques were employed. These methods include principal component analysis

(PCA), independent component analysis (ICA), uniform manifold approximation (UMAP) and t-

distributed stochastic neighbor embedding (tSNE). While for PCA, ICA, and tSNE the

implementation in the scikit-learn python package was used [34], the python package umap-learn

was used for UMAP [35]. For UMAP, n_neighbors was set to 10, 20, 50, 100, and 200 to focus

on local neighborhoods (smaller values for n_neightbors) and more global relationships (larger

values of n_neighbors) and min_dist was set to 0 to enable clustering. For tSNE, the following

parameters were used: n_iter = 5000, random_state=1, learning_rate="auto",

init="random". Perplexity was set to 10, 20, 30, 40, 50 and 100 to test the influence of local and

global shapes within the dataset, and set to 50 for the main results set with results of the other

settings available in the supporting information.

**Calculation of pocket descriptors**

As DeeplyTough does not provide any explicitly interpretable features, several simple pocket

descriptors (i.e. properties) were calculated for each pocket to identify the most important features

that DeeplyTough is learning. These descriptors include physicochemical properties, such as

aromaticity, hydrophobicity, net charge, solvent accessible surface area (SASA), and the

structure prediction quality parameters: predicted aligned error (PAE) and predicted local distance

difference test (pLDDT).

The aromaticity and hydrophobicity of each predicted pocket was calculated using the

ProteinAnalysis module of biopython [36]. This module calculates the aromaticity and

hydrophobicity of an amino acid sequence according to [37] and [38], respectively. The net charge

of each pocket was estimated using the amino acid residues lining each pocket and the charge

of the respective residue type at pH 7.0. All neutral residue types were assigned a charge of 0,

and all negative residue types (Glu, Asp) were assigned a charge of -1. All positively charged

residue types (Arg, Lys) were assigned a charge of +1. Histidine was assigned a charge of +0.5.

The SASA of each pocket was calculated using the Shrake and Rupley algorithm

implemented in biopython [36,39]. The sum of all SASA values for each residue in a pocket was

obtained and then divided by the maximal solvent accessible surface area for the respective

amino acid residue type as defined by [40] to obtain the relative SASA for each pocket.

To check how confident AF is about the relative and local positions of all residues within

one binding site, we calculated the mean PAE and the mean pLDDT for each pocket. The PAE

and pLDDT values for each protein of interest were obtained from the AF database, with the

pLDDT values extracted directly from the corresponding protein structure files [19].

**Clustering and analysis**

As DeeplyTough's embedding space for all binding sites was found to be too uniformly populated

to obtain a meaningful clustering, only ProBiS alignment scores were used to estimate the number

of unique pockets per species, as ProBiS operates at an increased level of structural resolution

compared to DeeplyTough. The pairwise binding site score matrix for each species was clustered

in two ways. In a simple approach, the similarity matrix for each species was clustered by treating

it as an adjacency matrix and applying different thresholds for setting edges. Clusters were then

detected by determining the number of components detected per graph. In addition, using the

igraph package (R 4.1.2), pocket clusters were created for both weighted, undirected graphs

based on the computed ProBiS similariy scores (normalized by the maximum thereof obtained in

a given species), and unweighted, undirected graphs by setting an edge between pockets (nodes)

if the max-normalized ProBiS score was greater than 0.1 (determined based on reviewing the

score distributions across species), and now edge otherwise. The latter allowed for randomizing

the graph under preservation of the degree distribution using the igraph "rewire" function with

"keeping_degseq" set active (niter=10). Furthermore, completely random, unweighted, undirected

networks (same number of nodes and edges) were also created and assessed. Clusterings were

obtained using the "cluster_leiden" function with the resolution_parameter set to 0.01 (different

thresholds were also tested, but were determined not to qualitatively affect the results. Respective

results are provided on the github page (see below).

When performing a cross-species analysis of binding site similarity, to estimate the

kingdom or species diversity of binding sites mapping to same region in tSNE projection plots, we

computed the entropy, $S$, on binned tSNE regions, with

241                            $$S = -\sum_{i=1}^{N} p_i \ log(p_i),$$                Eq. (1)

where $p_i$ is the fractional frequency of pockets associated with a given kingdom or species relative

to all pockets mapping into the same region (bin) of tSNE space, and $N$ is the number considered

kingdoms or species, respectively.

**Visualizations**

All protein structure visualizations were created using ChimeraX v. 1.7 [41]. All other plots were

created with the matplotlib [42] and seaborn library [43]. The Plotly library [44] was used to create

interactive plots of the embedding vectors produced by DeeplyTough.

# Results

**Pocket identification with P2Rank**

A set of 223,091 predicted protein structures found in the AF database for eleven species,

including bacteria, plants, unicellular fungi, an insect, and two mammal species (Table 1), was

subjected to binding site predictions using P2Rank. Testing various thresholds, we determined

the recall, i.e., the detection of known binding sites, was relatively stable across the P2Rank score

threshold range from 0.2-0.7, with 59% and 45%, respectively. To balance the number of true

positive and potential false positive binding sites, i.e., sites that are not actually involved in any

ligand binding, and to arrive at a manageable set size (176,156 pockets for a threshold of 0.2), a

threshold of 0.5 was applied to the probability score as predicted by P2Rank for all pockets. This

resulted in a set of 97,515 potential binding sites for the eleven different species used in this

study. A summary of the known and predicted binding sites of each species is provided in Table

1. The number of predicted pockets is strongly positively correlated with the number of predicted

protein structures (Pearson correlation coefficient, r=0.98). The largest number of pockets was

predicted for SOYBN, which also has the largest proteome. The smallest number of pockets is

predicted for ECOLI with only 3,809 predicted protein structures. Note that at this stage, no protein

or pocket clustering was performed. Thus, this correlation pertains to the raw counts.

**Table 1**. **Summary of proteins and pockets for all eleven species used in this study.**

| Species | Abbr. | Kingdom | UniProt acc. | # pred. struct. | # known pockets | # known pockets (exp.) | # pred. pockets |
|---|---|---|---|---|---|---|---|
| *Escherichia coli* | ECOLI | *Bacteria* | 625 | 3,813 | 1,014 | 224 | 2,648 |
| *Saccharomyces cerrevisiae* | YEAST | *Fungi* | 2311 | 5,623 | 1,207 | 109 | 3,000 |
| *Candida albicans* | CANAL | *Fungi* | 559 | 5,622 | 400 | 11 | 3,124 |
| *Arabidopsis thaliana* | ARATH | *Plantae* | 6548 | 24,778 | 4,668 | 169 | 11,075 |
| *Oryza sativa* | ORYSJ | *Plantae* | 59680 | 30,708 | 1,925 | 25 | 10,618 |
| *Zea mays* | MAIZE | *Plantae* | 7305 | 35,165 | 681 | 9 | 13,177 |
| *Glycine max* | SOYBN | *Plantae* | 8827 | 47,539 | 1,220 | 0 | 20,915 |
| *Drosophila melanogaster* | DROME | *Animalia* | 803 | 12,389 | 1,110 | 19 | 5,867 |
| *Caenorhabditis elegans* | CAEEL | *Animalia* | 1940 | 17,819 | 1,091 | 10 | 8,850 |

| *Mus musculus* | MOUSE | *Animalia* | 589 | 20,367 | 4,439 | 146 | 10,133 |
| *Homo sapiens* | HUMAN | *Animalia* | 5640 | 19,268 | 4,451 | 975 | 8,108 |

"# pred. structures" corresponds to the number of protein structures in the AF database without fragments and small proteins. "# known pockets" corresponds to the number of pockets found in UniProt and the "# pred. pockets" to the number of pockets predicted by P2Rank. Note the pocket sets (known/ predicted) are not mutually exclusive.

First, we wished to verify that the quality of the protein structures is comparable across

the species used in this study. The distribution of PAE (AF-predicted aligned error) values

associated with all pockets of the 11 species and the applied threshold value of 10 Å is visualized

in Figure 1A. The median PAE ranges between 2.9 and 4.3 Å for all species.  The median pLDDT

ranges between 87.4 and 92.7 for all species. As expected, pLDDT  and PAE medians show a

negative correlation. High pLDDT values (ECOLI) are matched by low PAE, whereas low pLDDT

(ORYSJ) are mirrored by a high PAE median. The largest variation in both the PAE- and the

pLDDT-values was observed for MAIZE, ORYSJ, and SOYBN. These species have the largest

proteomes and the most predicted pockets. The least variation was observed for ECOLI,

suggesting a lower fraction of less-defined, disordered regions in ECOLI.  Notwithstanding the

species-related differences, we considered all AF-predicted protein structure sets suitable for a

systematic cross-species comparison. Based upon visual inspection of the PAE and pLDDT

distribution plots, we set the thresholds for including pockets to PAE < 10 Å and pLDDT > 70.

**Fig 1. Structural confidence of pocket conformations as judged by AF-Predicted Aligned**
**Error (PAE) and predicted local distance difference test (pLDDT).**

(A) Violin plots for PAE values per-species. PAE-values correspond to the average of the pairwise

all-against-all residue PAE matrix. All pockets above the red dotted line were not considered for

further analysis. The median for all PAE distributions varies between 2.9 Å and 4.3 Å. MAIZE

shows the largest variation in PAE values. (B) Violin plots for pLDDT values per-species. pLDDT-

values correspond to the average of the respective pLDDT-values found for all residues within a

pocket. All pockets below the red dotted line were not considered for further analysis. The median

for all pLDDT distributions varies between 87.4 and 92.7. MAIZE shows the largest variation in

pLDDT values as well.

If available for a given protein, the domain information was mapped onto the respective,

predicted binding pockets. While 34,101 (35 %) pockets could not be assigned to any domain,

most of the pockets that were assigned to a single domain are annotated as protein kinase domains (InterPro: IPR000719, 4,007 pockets) and GPCR, rhodopsin-like, 7TM domain (InterPro: IPR017452, 2,083 pockets). A total of 38,929 pockets predicted by P2Rank for all species are inter-domain binding sites that are assigned to two or more different domain types. Most of these pockets are at the interface between two different domains, but 9,979 pockets are found at the interface of more than two domains. Another subset includes pockets that were found at a "domain-no_domain" interface (9,979 pockets). A small subset of 583 pockets was found in domains of unknown function (DUF).

Figure 2 shows the number of residues per binding site, i.e. residues lining the respective pocket reflecting on binding site size, for known and predicted pockets. Predicted pockets are, on average, significantly larger (median number of residues=24) than known binding sites (median=6). While known pockets are derived from experimentally solved structures with oftentimes manually annotated binding site residues, P2Rank predicts binding sites by clustering potentially druggable surface atoms. Therefore, residues not directly involved in the binding of any ligand are potentially included in the definition of predicted pockets as well. In addition, P2Rank does not include any additional pocket segmentation, resulting in large pockets if several ligands are binding to a similar region, as is, for example, the case, for ATP-dependent reactions.

**Fig 2. Pocket size statistic as gauged by the number of residues lining a given pocket in known and predicted pockets.**

On average, predicted pockets consist of more residues than experimentally annotated (known) pockets. While the median number of residues for known pockets is 6, the median is 24 for predicted pockets.

Exploring the reliability of P2Rank pocket predictions by checking the overlap of predicted sites with known sites, designating them as "found" if at least one residue-based overlap was detected, and "not found" otherwise, we observed that the most "found" as well as "not found" pockets were annotated as binding "nucleobases/ nucleosides/ nucleotides". An 8.55-fold enrichment (p-value << 0.05, Fisher's exact test) for known pockets that are denoted as "found" was observed for this ligand class as compared to all other ligand classes. About 15 % of the "nucleobases/ nucleosides/ nucleotides" pockets that were not identified were inter-chain pockets.

For inorganic ions, the second largest compound class (Fig 3), we observed a significant depletion

with p-value << 0.05 and odds ratio = 0.05 (Fisher's exact test) for known pockets that are denoted

as "found" as compared to all other ligand classes. The binding sites of inorganic ions are often

formed by a few amino acid residues, via charge-driven complex formation or covalently, thereby

exhibiting less of the hallmark features of compound binding sites (surface invaginations). This

makes it difficult to predict their location if they are bound to a site that is distant from any other

binding site. Thus, inorganic ions are proportionally overrepresented in the "not found" set. For

other ligand classes, there are significantly fewer binding sites annotated in UniProt. These

classes include heme, lipids, carbohydrates and derivatives, and amino acids and derivatives. A

significant enrichment for pockets that were "found" was observed for "Glycans", "Carbohydrates

& derivatives", "Amino acids & derivatives", "Lipids" and "Monosaccharides & derivatives" while a

significant depletion of pockets that were "found" was observed for "Hetero nuclear clusters". For

"Hemes", all pockets were found. For nucleic acids, none of the pockets could be identified by

P2Rank using a threshold of 0.5 for the probability. "Oligo- and polypeptides" were found at the

expected average rate, odds-ratio=0.79, p=0.387. A summary of the respective statistics of the

Fisher's exact test and the number of pockets denoted as "found" and "not found" for all ligand

classes can be found in Table 2.

**Fig 2. Ligand annotation of pockets.**

Known (i.e., annotated) pockets identified by P2Rank used as the binding site predictor and with

correspondingly available ligand annotation were classified as "found" (n=11,662, 52.1 %), and

"not found" (n=10,709, 47.9 %), otherwise. The total number of occurrences for each compound

class is denoted in brackets in the legend.

**Table 2. Summary table for the compound classes of the ligands binding to known pockets.**

| Compound class | ChEBI ID | # Pockets ("Found") | # Pockets ("Not found") | Odds ratio | $p_{adj}$-value |
|---|---|---|---|---|---|
| Nucleobases, nucleosides & nucleotides | CHEBI:18282, CHEBI:33838, CHEBI:36976 | 8,092 | 2,244 | 8.55 | <<0.001 |
| Inorganic ions | CHEBI:36914 | 1,191 | 7,265 | 0.05 | <<0.001 |
| Other compounds | - | 1,114 | 296 | 3.72 | <<0.001 |
| Carbohydrates & derivatives | CHEBI:16646, CHEBI:63299 | 548 | 103 | 5.08 | <<0.001 |
| Amino acids & derivatives | CHEBI:33709, CHEBI:35238, CHEBI:83821 | 260 | 135 | 1.79 | <<0.001 |
| Lipids | CHEBI:18059 | 173 | 94 | 1.7 | <0.001 |
| Hetero nuclear clusters | CHEBI:33733 | 84 | 433 | 0.17 | <<0.001 |
| Oligo- & polypeptides | CHEBI:25676, CHEBI:15841 | 79 | 92 | 0.79 | 0.387 |
| Monosaccharides & derivatives | CHEBI:35381, CHEBI:63367 | 71 | 34 | 1.92 | 0.006 |
| Glycans | CHEBI:37163, CHEBI:167559 | 73 | 4 | 8.52 | <<0.001 |
| Hemes | CHEBI:30413 | 13 | 2 | 5.97 | 0.003 |
| Nucleic acids | CHEBI:16991, CHEBI:33697 | 0 | 7 | 0.0 | 0.002 |
| No known compounds | - | - | - | - | - |

For all compound classes considered in this study, the respective ChEBI IDs, the number of pockets identified ("found") and not identified ("not found") by P2Rank and the corresponding statistic of the Fisher's exact test adjusted p-value (see Materials and Methods) are listed. "No known compounds" is listed for reasons of completeness of considered categories only.

Several factors may explain why known pockets might not be predicted by P2Rank. One

reason is that flexible loops flanking the binding site do not have a stable conformation. These

flexible loops are often missing in experimental structures. In AF, they are assigned to a single

conformation with a mostly low pLDDT values. An example for this case is shown in Fig 4A. While

the two flexible regions are missing completely in the experimental structure of tubulin

polyglutamylase TTLL7 (PDB ID: 4YLR), the ADP binding site is not accessible in the AF

predicted structure (UniProt ID: Q6ZT98). Other known binding sites, such as the LacNAc binding

site on galectin-3, only consist of few residues that form a flat binding surface (see Fig 4B).

**Fig 3. Examples of binding sites that could not be identified by P2Rank due to flexible regions and flat surfaces.**

(A) Experimental (PDB ID: 4YLR, orange) and AF structure for Q6ZT98. In 4YLR, ADP (beige) is bound as a ligand. This ADP binding site is blocked in the AF structure (green) by two flexible regions (residues: 161-172, 235-269) that are not present in the experimental structure. Therefore, P2Rank cannot predict the ADP binding sites. (B) Galectin-3 (PDB ID: 1KJL, orange) binds LacNAc (beige). The binding site is found on a flat surface and consists of only few AAs.

**Per-species binding site comparison (ProBiS)**

As we wished to discern the underlying scaling law linking the pocketomes of different species to their respective proteomes viewed from an evolutionary perspective, we performed a per-species comparison of all predicted binding sites using ProBiS to determine the number of unique binding sites present in each species. This graph-based comparison algorithm results in a sparse similarity matrix, where zeros denote pocket pairs that could not be aligned and thus are considered dissimilar. The higher the alignment scores, the more similar two pockets are. Pockets that could not be aligned to any other pocket of the respective species were termed "singletons". The number of singletons and the number of communities as detected by the Leiden algorithm are listed in Table 3. The fraction of "number of singletons" over the "number of predicted" pockets is smaller for species with larger proteomes and more predicted pockets than for species with smaller proteomes, such as ECOLI, YEAST, and CANAL. For all three species, about 40% of all pockets were singletons. The smallest fraction of singletons over the number of pockets (0.079) was detected for SOYBN. The inverse relationship (Pearson correlation r= -0.84, p=0.001) between proteome size and fraction of singletons is, to some degree, to be expected, given that the pocket space crowds up more with more proteins and their associated pockets. It would, however, be different if more proteins meant altogether new and different pockets, a question we will turn to below.

As singletons might potentially be falsely predicted binding sites that are not actually involved in any ligand binding, as it may seem unlikely that proteomes harbor sites that occur only once and thus for a single-case ligand binding event, we compared various properties of the

pockets for singletons and non-singletons, such as P2Rank probability, PAE, pLDDT, number of

residues, solvent-accessible surface area (SASA), aromaticity, net-charge, and hydrophobicity

Fig 5). All of the tested properties show significant differences between singletons and non-

singletons with p-value < 0.01 (t-test), but the actual effect sizes, calculated using Cohen's d,

differ markedly across the various properties. The largest effect size was observed for the P2Rank

probability of the pockets, i.e., the confidence of the pocket prediction. The median probability of

the predicted pockets drops from 0.83 for the non-singletons to 0.67 for singletons with an

associated effect size of 0.738 (Fig 5G). The effect size for all other properties were found in the

range 0.053 to 0.468 (see Fig 5). Taken together, singletons differ markedly from non-singleton

pockets, which may indicate that they are either special because of their respective particular

binding modes and ligands outside the P2Rank training sets and score-tuning approaches, or that

they may be falsely predicted pockets.

**Fig 5. Annotation confidence of pockets for non-singletons and singletons.**

Violin plots of various properties contrasted for singletons and non-singletons (A) hydrophobicity,

effect size (Cohen's d) between the two groups is 0.334. (B) aromaticity, d=0.207. (C) net charge

d= 0.053. (D) solvent accessible surface area (SASA),d=0.230. (E) pLDDT,d=0.369. (F) PAE,

d=0.223. (G) P2rank, d=0.738. (H) number of residues, d=0.468.

**Table 3. Pocket clustering statistics.**

| Species | # Communities | # Singletons | # Singletons/ # Pockets | Largest community |
|---------|---------------|--------------|-------------------------|-------------------|
| ECOLI | 273 | 1,076 | 0.41 | 185 |
| YEAST | 335 | 1,182 | 0.39 | 133 |
| CANAL | 300 | 1,293 | 0.41 | 124 |
| ARATH | 823 | 1,532 | 0.14 | 893 |
| ORYSJ | 780 | 1,711 | 0.16 | 924 |
| MAIZE | 966 | 1,879 | 0.14 | 1,086 |
| SOYBN | 1,376 | 1,792 | 0.09 | 1,868 |
| DROME | 514 | 1,688 | 0.29 | 328 |
| CAEEL | 617 | 2,225 | 0.25 | 406 |
| MOUSE | 708 | 1,805 | 0.18 | 1,068 |
| HUMAN | 682 | 1,705 | 0.21 | 516 |

Communities, i.e. sets of similar pockets, were found by applying the Leiden algorithm on unweighted graphs (binary similarity matrix based on threshold-based discretization of the ProBiS scores, see Materials and Methods). Singletons are lone pockets, not similar to any other pocket in a given species. While the number of singletons is getting larger, the number of singletons per pocket is getting lower with larger proteomes. The highest fraction is observed for CANAL, ECOLI, and YEAST. These species have the lowest number of communities. The lowest fraction is observed for ARATH and SOYBN. "Largest community" indicates the number of member pockets in the largest found community.

Next, we addressed the question as to whether expanding proteomes during the course

of evolution are commensurate with proportionally increased pocketome sizes. While a

proportional increase may be the naive expectation, alternatively, relatively more or fewer pockets

may have emerged during evolution than is expected purely based on proportionality. That is, we

wished to understand whether evolution "invests" in new proteins to expand the set of diverse

compounds that can be processed (via binding to proteins), or whether functions other than

compound binding are invested in. In case of the latter, with more unique protein folds, fewer than

proportionally more binding pockets would be found with increased proteome sizes. By contrast,

evolution could also proceed "economically" by reusing the same fold but evolving new and

additional binding sites on them via selected point mutations. Under that scenario, more than

proportionally more pockets per protein fold would be seen.

To explore this relation between unique binding sites and unique folds across species, we

plotted the number of unique binding sites (communities + singletons) against the number of

FoldSeek (FS) clusters (https://cluster.foldseek.com/) [24] and the number of clusters

(communities only) against the number of FS clusters. We decided to gauge the proteome size

as the number of FS clusters and not by the number of encoded proteins to avoid wrongly

associating different sizes of protein families (paralogs) with protein "novelty". This approach also

accounts for the effect of gene duplication in some plants, such as SOYBN, where 73.6 % of the

genes are duplicated according to its BUSCO score. Note that unlike reported above, where we

stated a high linear correlation of numbers of proteins and pockets, here we performed a

clustering of both entities to really capture unique proteins and unique pockets.

As shown in Figure 6A, there is an inverse association between the number of FoldSeek

clusters, $N_{FS}$, in the different species and the number of pockets clusters per FS cluster, $N_{Pocket}$

$_{clusters}/N_{FS}$, where $N_{Pocket\ clusters}$ is the number of pocket clusters in a given species. Compared to

ORYSJ, which has the lowest ratio of pocket clusters to FS clusters, the ratio between the number

of pocket clusters and FS clusters is about four times higher for ECOLI, which has only about

1/7th the number of FS clusters compared to ORYSJ. Note that under pure proportionality, we

would expect this relationship to be a constant (as we normalize the number of pocket clusters

by the number of FS clusters for better visual clarity). Regressing log($N_{Pocket\ clusters}$) as a linear

function of log($N_{FS}$), a strong linear relationship (Pearson correlation coefficient $r$= 0.942) was

obtained, suggesting a power law relationship with proportionality of $N_p \sim N_{FS}^a$, and $a$= 0.398 (slope

of the regression line, unweighted graph), i.e. substantially smaller than one. A smaller than one

exponent $a$ suggests a less than linear (sub-linear) increase of the number of unique pocket

clusters with increasing number of FS clusters, i.e. larger and more divers proteomes do not

harbor a proportionally greater diversity of binding sites. For the original ProBiS-score-weighted

graph, $a$=0.44 was obtained.

**Fig 4. Scaling law of the number of unique pockets vs. proteome size.**

(A) Number of unique pockets per species ($N_{Pocket\ clusters}$) normalized by the number of FoldSeek

(FS) clusters vs. the number of FS clusters ($N_{FS\ clusters}$). On a log-log scale for log($N_{Pocket\ clusters}$) ~

log($N_{FS\ clusters}$), a linear regression yielded: slope = 0.398, intercept = 4.015, Pearson r = 0.942.

(B) Number of pocket communities per species, $N_{Communities}$, i.e. without considering singleton pockets, normalized by the number of FoldSeek (FS) clusters vs. the number of FS clusters ($N_{FS clusters}$). On a log-log scale for $\log(N_{Communities}) \sim \log(N_{FS clusters})$, a linear regression yielded: slope = 0.73, intercept = -0.364, Pearson r = 0.952). Note that the reported slope values correspond to the power law exponent, $a$, discussed in the main text. The reported results relate to the unweighted (binarized) network (see Materials and Methods).

To obtain a null-model, we randomized the unweighted graphs (binary adjacency matrix), preserving the degree distribution (the actual network was found to follow a power-law degree distribution (see below). From five performed randomization runs, an average scaling law coefficient, $a=0.882+/-0.001$ (s.d.), for the randomized network was obtained. i.e., substantially larger than the one obtained for the actual network ($a=0.398$). Thus, compared to random expectation, evolution shows evidence of a pocketome diversity plateau. Despite more and different protein structures appearing, not proportionally and less than randomly expected different binding sites appeared in evolution. Note, under a completely random graph model (randomizations performed without degree-preservation, but preserving node and edge counts only, randomly assigned), we obtained a very loosely defined (r= -0.42+/-0.01 (s.d.) negative coefficient (-0.98~/-0.01 (s.d.), which, however, is a) dictated by outliers (mouse clusters into one cluster), and b) is not the proper reference network, as the actual network is a power-law node degree network. Furthermore, for the original ProBiS-score-weighted network, degree-preserving randomizations are not possible; they require unweighted graphs (binary adjacency matrix). For completely randomized ProBiS-score input matrices (kept symmetric), 0.45+/-0.001 (s.d.) was obtained.

As reported above, singletons pockets, i.e., pockets not found similar to other pockets within a given species, differ markedly from non-singletons with regard to their properties. To assess the effect of singletons, which may be false-positive pockets, on the scaling law statistic, we performed analyses ignoring singletons. Excluding singletons, we determined a power law exponent of $a= 0.73$ for the actual network, and $a=0.925+/-0.002$ for degree-preserved random networks.. Thus, even without considering singletons, actual pocket communities grow more slowly than randomized versions and are distinctly lower than expected for proportionality, albeit

the difference is smaller for communities compared to the overall clustering statistic, i.e., including

singletons.

Regarding the sizes of the communities, the Leiden algorithm detects a few large

communities for each species and many smaller communities that only consist of two or three

different pockets (Fig 7). Overall, linear relationships are observed in the log-log plot of cluster-

frequency and cluster size, suggesting an overall power-law relationship for the cluster-size

distribution, which is oftentimes observed for such data types in biological systems [45]. The

largest communities are found in SOYBN and MAIZE, whereas in ECOLI, the distribution is shifted

to smaller clusters. Thus, as expected, the size distribution of clusters correlates with set size

(number of pockets, Table 1).

**Fig 5. Cluster size frequency distribution for the eleven species used in this study.**

The log(Cluster size) plotted against the log($N_{Clusters}$). The clustering for all species reveals that
there are many clusters that only contain few pockets and only few clusters with many pockets,
and overall, a power-law degree distribution is observed. The largest clusters are found in
SOYBN, MAIZE, MOUSE.

As ATP is the most frequent known ligand (n = 6,327 for ChEBI:30616) and protein kinase

is the most frequent domain annotated for the predicted pockets, it is expected that the respective

largest communities are enriched for protein kinase domain annotations. This is the case for

ARATH, ORYSJ, MAIZE, SOYBN, and CAEEL. For these species, 81.03 % (CAEEL) to 96.41 %

(SOYBN) of the pockets in the largest cluster are found at least partially in a protein kinase

domain. For YEAST, CANAL, DROME, and HUMAN, the protein kinase domain dominates in the

second largest cluster with 127, 105, 253, and 457 members, respectively. The two largest

communities in MOUSE contain mainly pockets that are found at least partially in the "GPCR,

rhodopsin-like, 7TM" domain. The fourth largest community in MOUSE is enriched for pockets

found in the "Protein kinase" domain. The paucity of protein kinase annotations in ECOLI is

expected, as it does not have the eukaryotic protein kinase domain. The largest proportion of

pockets is found at least partially in the "ABC transporter-like, ATP-binding" domain (74 of 185

pockets in the largest community) for ECOLI.

**Large-scale, cross-species binding site comparison (DeeplyTough)**

Thus far, we clustered pockets and proteins per species. Evidently, the pocket datasets could

also be inspected across all species. As the computational resources needed for an all-against-

all comparison using ProBiS are prohibitively high, we decided to use the embedding vectors

obtained by DeeplyTough for a large-scale, inter-species comparison. By using DeeplyTough, a

method to describe pockets as numeric vectors of length 128, we were able to perform an all-

against-all pocket comparison of all 97,515 pockets across all eleven species used in this study,

arriving at a global view of the pocketome "universe". We present this view primarily by using

tSNE (perplexity = 50 (~number of neighbors considered during embedding), tSNE maps using

different settings are provided in the supplement) as an appropriate projection method, but

included projections for other perplexity values, UMAPs with varying number of neighbors, PCA

(principal component analysis), and ICA (independent component analysis) as well as part of

supporting material (see S3 - S9 Fig).

**Fig 6. tSNE plots for all predicted pockets of all eleven species colored by different properties.**

Perplexity set to 50. tSNE plots obtained for other perplexity settings are provided in the
supporting information.

Figure 8 shows all pockets projected into the first two tSNE components colored by the

pocket properties: hydrophobicity, aromaticity, net charge, solvent accessible surface area

(SASA), pLDDT. PAE, number of residues, and probability. Each point in the plot represents one

unique pocket, allowing us to gauge the most important properties by which the pockets

segregate. Most strongly, pockets differ by hydrophobicity, to a lesser degree by aromaticity,

charge, and the quality parameters pLDDT, PAE, and pocket probability. Size-related properties

(SASA and number of residues) do not cause a notable segregation of pockets.

To examine whether particular compound classes correspond to binding sites displaying

characteristic properties, we examined the pocketome projection with regard to their annotation,

i.e. which compounds are known to bind to it, provided this information is available (Fig 9). The

most abundant ligands found in known pockets are "nucleobases, nucleosides, and nucleotides",

such as ADP and ATP. The predicted pockets that overlap with these pockets are mostly found

on the right side of the tSNE plot. As evident from Figure 8, these pockets are hydrophilic to

neutral on the hydrophobicity scale and are predicted with moderate to high probability. In

addition, they contain few aromatic amino acid residues. As mentioned above, P2Rank does not

perform any sub-sampling of the predicted pockets. Binding sites that contain several ligands are

therefore predicted as one large pocket instead of several separate pockets.

**Fig 7. tSNE plot of all pockets of all eleven species colored by different ligand classes.**

The most abundant ligand class are nucleobases/nucleosides/nucleotides which are mostly

found on the right side of the tSNE plot. While glycan binding sites cluster well, the binding sites

for other ligand classes are found in different regions of the tSNE plot. An interactive plot for

further inspection is made available as well (see Data availability).

Glycan binding sites are rather hydrophilic, contain moderate amounts of aromatic amino

acid residues, and are all found in the Glycoside hydrolase family 16 domain (InterPro:

IPR000757). They are found on the left side of the tSNE plot and cluster well. One reason for that

might be their unique shape as they are formed by a set of anti-parallel β-sheets and their

connecting loops (example shown in Fig 10). In the vicinity of these pockets, there are several

other, not annotated pockets that are found on the glycoside hydrolase family 16 domain. This

suggests that more glycan binding sites could be potentially identified with this approach.

For other ligands, such as lipids, carbohydrates, and amino acid derivatives, the predicted

binding sites that overlap with known binding sites of those compounds are not clustering strongly.

Their associated binding sites are found in several parts of the tSNE plot.

**Fig 8. Glycan binding site of the protein Q8LC45.**

The binding site residues are displayed as sticks, other parts of the protein as cartoon. The

binding site is formed by several anti-parallel β-sheets. This protein visualization was created

using ChimeraX (v. 1.7).

In the left part of the tSNE map, there are two regions where only few pockets could be

annotated. The first region, marked in orange in Fig S14, contains about 12,000 pockets. The

median probability for these pockets is significantly lower than for the rest of the pockets (0.65 vs.

0.83), and about one third of these pockets are at least partially found in a transmembrane

domain. Regarding the domains that these pockets are assigned to, about 47 % of these pockets

could not be assigned to any domain. The rest of the pockets are assigned to various other

domains with "GPCR, rhodopsin-like, 7TM" domain being the most common domain. The second region corresponds to the cluster to the left in the tSNE map. This region contains about 3,650, mainly hydrophobic pockets (median hydrophobicity = 1.858) that were predicted by P2Rank with a slightly higher median probability than the rest of the pockets (0.842 vs. 0.806). As noted for the first described region, about half of the pockets found in this region could not be assigned to any domain, while the other pockets are assigned to various other domains.

As we wished to discern kingdom- and species-specific differences between the pocketomes, we colored the tSNE projection by kingdom of life and species. The kingdom-specific projection map shown in Figure 11 does not reveal any strong kingdom-specific differences for *bacteria*, *fungi*, *plantae*, and *animalia*, with pockets from all kingdoms occupying all of the tSNE space (Fig 11A) and the binned entropy heatmap capturing diversity revealing a rather uniform kingdom representation. The only region that contains less binding sites found on bacterial proteins, is the upper right part of the tSNE plot (Fig 11A and 11B). This corresponds to the region where most of the pockets are found in the protein kinase domain. This eukaryotic domain is not found in any bacterial protein. Similarly, no species-specific binding sites are revealed in the respective tSNE plot shown in S15 Fig.

Taken together, these results suggest that either compound binding is not very specifically associated with pocket descriptors, or, and we consider this to be more likely, the binding site definition of DeeplyTough's embeddings may be too coarse-grained to capture local similarities between different binding sites or differences, respectively. The binding sites are rather separated by their overall physicochemical features and high-level domain information. This leads to an embedding space that is populated too uniformly to allow for meaningful fine-grained clustering.

**Fig 9. tSNE 2D-projection plots colored by the respective kingdoms of life.**

(A) All pockets in their respective tSNE embedding space, (B) Binned entropy, S (see Materials and Methods), heatmap plot, capturing for every square bin the kingdom diversity with low S values indicating low diversity and high values corresponding to respectively increased diversity. All grid cells containing less than 20 pockets are colored black, as no robust entropy estimate can be computed. An interactive plot (also for species-level annotations) for further inspection is made available as well (see Data availability).

# Discussion

In this study, we attempted to generate a global view of the pocketome universe. Inspecting the eleven species for which the AlphaFold database contains a comprehensive set of predicted protein structures and applying established binding site (pocket) prediction algorithms, we performed a systematic analysis of binding site properties, ligand sets, clustering, and their scaling with proteome sizes.

Our analysis suggests that during the course of evolution, with species evolving larger proteomes and more differentiated molecular functions, unlike perhaps expected, the increase in proteome size and complexity does not commensurate with a proportional increase of binding site "innovation". Instead, fewer than proportionally expected unique binding sites seem to have emerged in evolution (Fig 6A). This implies that, rather than compound binding sites, other functions may have emerged and expanded, such as possibly transcriptional and translational regulation and signaling. However, it needs to be cautioned that this conclusion is based on the approach pursued here, with all the caveats pertaining to the used tools and settings (detailed below).

How evolutionary increases in molecular repertoire (genome-size, number of genes etc.) relate to increases in functional diversity has been studied before, in particular in recent years, with close-to-comprehensive coverage of sequence-and-structure space, whether there is evidence of a saturation of sequence-structure-function space exploration by evolution [46,47]. Notably, Nimwegen [46] reported on scaling laws relating number of genes with the size of various functional categories (number of genes devoted to the respective functions) based on gene-ontology, found for all but one sub-linear scaling laws, with the notable exception of "signal transduction", for which higher than linear (exponent 1.55-1.90) were reported, in line with our hypothesis stated in the previous paragraph. With regard to structural space and its association with function, it was repeatedly reported that both same-structure-different-function and different-structure-same-function can be observed [48,49]. Our results suggest that at large scale, across kingdoms, the functional space (with pocket diversity taken as a proxy of functional capabilities) seems more constrained than structure space and that the predominating mode of evolution is

not the invention of novel function using the same fold, as this would have resulted in a greater

than linear scaling law, but rather that functional novelty requires disproportionally more novel

structures. Interestingly, a recent survey of microbial protein structures and exploiting the recent

leap in structure information coverage suggested that structure space is already largely saturated

[47] which, in light of our findings, may mean that functional diversity is also saturated, and has

been saturated already earlier (in evolution) than the structural space.

While the pocketome universe exhibits a partitioning into pockets of particular

physicochemical properties (Fig 8) and compound call binding (Fig 9), we did not identify any

such segregation with regard to species or kingdom (Fig 11 A, B), except for the missing protein

kinase domain in bacteria. This may either result from technical limitations to discern any present

species/kingdom-specific pockets (discussed below), or may suggest that species/kingdoms are

not characterized by specific binding sites. However, as, for example, plants do contain

specialized metabolic reactions (secondary metabolism), it seems surprising that this should be

the case. Yet, as those specialized compounds are consumed and degraded by other species

as food, perhaps it is a wrong assumption to expect unique pockets in plants, as in species that

feed on plants, similar binding sites need to exist to process the compounds.

One source of uncertainty possibly affecting our analyses concerns the reliability of

predicted pockets. As P2Rank [6] does not perform any sub-sampling, the predicted pockets are,

on average, larger than known pockets (Fig 2) and are likely to merge several distinct pockets,

i.e., for pockets that are found in protein kinase domains. Introducing a sub-sampling algorithm

as a post-processing step of P2Rank, could improve the precision of the binding site comparison

as well as the quality of the P2Rank predictions in general.

One option to potentially refine predicted binding sites may be to exploit domain

information. In our study we identified a subset of 32,098 pockets that are found at a domain -

no_domain interface, i.e. sites for which some residues belong to an annotated domain, while

others carry no domain annotation information. Those pockets may be less well defined

structurally than pockets fully embedded in specific domains of known function and may, hence,

require a focused analysis.

Inorganic ions are the second largest fraction of known pockets that are not correctly

identified via P2Rank (Fig 3). P2Rank may often fail to correctly identify such sites as they tend

to involve a smaller number of residues during binding. P2Rank prediction might be influenced by

missing ions or cofactors as it depends on physicochemical properties. Employing AlphaFill to

implant the most likely ions or cofactors into the predicted structures could potentially replace

binding site prediction for this compound class [50].

Another type of ligand for which P2Rank struggles to identify the binding sites are

carbohydrates, likely due to their complexity in shape, flexibility, and size. This problem has been

recently addressed by the development of several specialized prediction tools, such as

DeepGlycanSite [51] and PeSTo-Carbs [52]. Specialized binding site prediction tools, such as

PocketMiner [53], have also been developed for the detection of cryptic pockets. Cryptic pockets

are pockets that are only apparent following a conformational change or upon ligand binding.

Complementing the current data set of predicted pockets with binding sites predicted by

specialized tools, is a promising route towards investigating the full space of protein binding sites.

Our study yet again reveals that binding site comparison remains a challenging task,

although it has been a subject of research and tool development for many years. While in proteins

there are secondary structural elements that allow for a relatively fast and relatively well-defined

comparison between proteins, the comparison of protein binding sites is more complex, as they

generally do not have a continuous sequence. For a meaningful spatial comparison, rotational

and translational invariance is required for binding site comparison tools, such as implemented

by DeeplyTough [18]. DeeplyTough compares binding sites in an alignment-free manner by

reducing the representation of each pocket to a feature vector. While this captures the overall

shape, domain information and physicochemical features of a pocket and allows for fast

comparison (simple vector comparison metrics), it is limited in the detection of similar sub-shapes

of two binding pockets. Although DeeplyTough was trained to separate dissimilar binding sites by

a certain margin with a contrastive loss, the embedding space of DeeplyTough is densely and

uniformly populated, making it difficult to obtain a clustering for inter-species comparison as true

positives (= similar pockets) as well as false positives (= dissimilar binding sites) are close in the

vector space. One reason for that might be pocket types encountered in our study, but that were

not found in the limited training data set used by the DeeplyTough developers.

In contrast to alignment-free methods, alignment-based methods, such as ProBiS [29],

allow insights into which amino acid residues or surface atoms of the compared pockets are

similar and contribute positively to the comparison score. This allows for the easy interpretation

of why two binding sites are considered similar or different and, importantly, allows for a sensitive

identification of local sub-structural matches. However, these approaches are computationally too

expensive at the moment to allow large-scale, inter-species studies on tens of thousands of

binding sites as attempted here [28]. Hence, the development of a fast and reliable alignment-

based or alignment-free comparison method is still needed to improve inter-species binding site

comparison and clustering at scale.

On the other hand, the search space of potentially similar binding sites could be reduced

by additional filtering to make alignment-based methods feasible for an all-against-all inter-

species comparison. Binding sites that are known to be dissimilar could be pre-filtered by their

amino acid composition or their overall physicochemical properties. Binding sites that are known

to be similar could be pre-clustered based on the domain to which they belong. This would reduce

the number of pairwise comparisons needed, especially for redundant domains, such as the

protein kinase domain.

Many of the recently published binding site comparison tools have been evaluated on the

ProSPECCTs datasets [27]. ProSPECCTs is an ensemble of seven different data sets of pairs of

binding sites with known similarities. In contrast to the aim of this study, the ProSPECCTs

datasets do not contain all-against-all comparisons of the respective binding sites. Therefore, the

actual performance of the binding sites comparison tools might be overestimated. In addition, the

datasets were derived from known protein-ligand complexes. Reim et al. [28] investigated the

performance of their tool, SiteMine, for predicted pockets and observed that it is performing worse

for some datasets. As predicted pockets consist, on average, of more residues, they might lose

some specificity and overall physicochemical properties represented unspecifically or even

incorrectly. This might lead to pockets that are more similar than they are in reality. Hence,

complicating the binding sites comparison at a large scale. DeeplyTough might lose additional

specificity because it is using all surrounding residues of the center of mass of the pocket for

embedding, and the usage of predicted pockets instead of known pockets, as they contain, on

average, more residues than known pockets.

In addition, using predicted, monomeric protein structures from AlphaFold database

results in different challenges that need to be addressed in future research. These challenges

include protein-protein interactions, post-translational modifications, and conformational changes

that occur upon binding. Currently, the AlphaFold database only includes monomeric structures.

However, studies for, e.g., HUMAN and YEAST, suggest that most proteins are at least part of

one protein complex [54,55]. Structural predictions might differ between monomer and multimer

predictions. Therefore, it would be desirable to investigate potential binding sites from a multimer

perspective. Although multimer prediction is challenging due to often unknown stoichiometries

and the exploding combinatorics with higher number of chains, reliable prediction of structures for

dimers has been achieved by new models, such as AlphaFold-Multimer [56]. The challenge to

identify potential interaction partner(s) for each protein of interest through, e.g., protein-protein

interaction prediction, remains.

The AF database contains one predicted structure per protein. However, there are proteins

that change their conformation upon binding or have an ensemble of different conformations.

These conformational changes are currently not taken into consideration. In addition, studies

showed that AF tends to predict the "mean" structure in cases where the apo- and holo-structure

of a protein differ [57]. This might influence the prediction and classification of so-called cryptic

pockets that are only found in holo-structures. This problem can be tackled by established

methods, such as normal mode analysis (NMA), as well as DL-based methods that have been

developed recently to overcome this issue [58,59].

Importantly, the AF database does not consider any post-translational modifications which

might influence the structural conformation of proteins. Prominently, protein phosphorylation has

been discussed as a modulator of compound binding [60]. A recent, PDB-wide study [61] could

show that for phosphorylation, these conformational changes are rather small, with only ~30% of

phosphorylation events causing an RMSD $\geq 2$ Å as compared to the non-phosphorylated

structure. However, this number might be too conservative as the PDB database is biased

towards less flexible conformations as conformational flexible structures are harder to crystallize.

In the same study, the authors could identify a small subset of structures for which

phosphorylation is mechanically coupled to functional sites. Extending their work to AF predicted

structures, might be a promising avenue of future research.

Another caveat concerns the notion of binding specificity. We approached our study by

positing that novel ligands require novel binding sites. However, it is known that binding sites as

well as ligands can be promiscuous [15], thereby a one-to-one relationship will not always be true.

As a switch from selective to promiscuous binding may involve only small structural changes,

functional diversity may grow even in the absence of easily noticeable structural changes.

Lastly, it is known that a substantial portion of proteins and regions within proteins are

structurally disordered under physiological conditions [62]. However, it is also known or surmised

that these proteins or protein regions do engage in (specific) molecular interactions, including

binding to small molecules [63,64]. While structure-based binding site detection in disordered

regions are challenging, attempts have been made to apply molecular dynamics simulation to

interactions of compounds with disordered protein regions [65], or to predict sequence regions

that are likely to interact with other molecules, for example, proteins [66]. Future computational

predictions of small molecules with disordered regions of proteins will critically depend on the

availability of large-enough experimental datasets as generated by specific NMR protocols [67]

and others.

Although some of the challenges mentioned above, such as the post-translational

modification, are addressed by AlphaFold3 [68], structures produced by AF3 are not yet available

at such a large scale as they can be found in the AF database. Hence, they cannot yet be used

for a proteome-wide comparison of binding pockets.

**Conclusions**

To our knowledge, we present the first large-scale intra- and inter-species study of compound

binding sites on protein surfaces using AlphaFold structures to arrive at a map of the pocketome

universe. The binding site comparisons conducted for different species using ProBiS reveal that

species with larger and more diverse proteomes are not commensurate with proportionally larger

and more diverse pocketomes; instead, a sub-linear relationship between the number of unique

pockets vs. unique structures was observed. Our study also documents the huge potential of the

new quality and quantity of structural data availability, as well as the power and shortcomings of

existing pocket detection and comparison methods.

**Acknowledgements**

We wish to thank Zoran Nikoloski, Ariane Nunes Alves, and Alisdair Fernie for helpful discussions.

**Data availability**

The software code developed and/ or used to conduct this study, along with corresponding input

files, is available at https://github.com/zillmerhanne/PocketomeUniverse. In addition, this

repository contains all pocket descriptors and embeddings produced by DeeplyTough, as well as

interactive Plotly plots for the tSNE projections of the DeeplyTough embeddings, which allow

filtering for specific compound classes or species.

**Authors' contributions**

The study was conceived and designed by both authors, all computations were performed by

H.Z., except for the final version of the Leiden clustering computations and random controls.

Both authors reviewed and interpreted the results and wrote the manuscript.

**Competing Interests**

The authors have declared that no competing interests exist.

# References

1.     Eguida M, Rognan D. Estimating the Similarity between Protein Pockets. Int J Mol Sci.
2022;23. doi:10.3390/ijms232012462

2.     Le Guilloux V, Schmidtke P, Tuffery P. Fpocket: An open source platform for ligand
pocket detection. BMC Bioinformatics. 2009;10: 1–11. doi:10.1186/1471-2105-10-168

3.     Desaphy J, Azdimousa K, Kellenberger E, Rognan D. Comparison and druggability
prediction of protein-ligand binding sites from pharmacophore-annotated cavity shapes. J Chem
Inf Model. 2012;52: 2287–2299. doi:10.1021/ci300184x

4.      An J, Totrov M, Abagyan R. Pocketome via Comprehensive Identification and
Classification of Ligand Binding Envelopes. Mol Cell Proteomics. 2005;4: 752–761.
doi:10.1074/MCP.M400159-MCP200

5.      Halgren TA. Identifying and characterizing binding sites and assessing druggability. J
Chem Inf Model. 2009;49: 377–389. doi:10.1021/ci800324m

6.      Krivák R, Hoksza D. P2Rank: machine learning based tool for rapid and accurate
prediction of ligand binding sites from protein structure. J Cheminformatics. 2018;10: 39.
doi:10.1186/s13321-018-0285-8

7.      Volkamer A, Eid S, Turk S, Jaeger S, Rippmann F, Fulle S. Pocketome of human
kinases: Prioritizing the ATP binding sites of (Yet) untapped protein kinases for drug discovery.
J Chem Inf Model. 2015;55: 538–549. doi:10.1021/ci500624s

8.      Hedderich JB, Persechino M, Becker K, Heydenreich FM, Gutermuth T, Bouvier M, et al.
The pocketome of G-protein-coupled receptors reveals previously untargeted allosteric sites.
Nat Commun 2022 131. 2022;13: 1–12. doi:10.1038/s41467-022-29609-6

9.      Kufareva I, Ilatovskiy AV, Abagyan R. Pocketome: an encyclopedia of small-molecule
binding sites in 4D. Nucleic Acids Res. 2012;40: D535. doi:10.1093/NAR/GKR825

10.     Bhagavat R, Sankar S, Srinivasan N, Chandra N. An Augmented Pocketome: Detection
and Analysis of Small-Molecule Binding Pockets in Proteins of Known 3D Structure. Structure.
2018;26: 499-512.e2. doi:10.1016/j.str.2018.02.001

11.     Gao M, Skolnick J. A Comprehensive Survey of Small-Molecule Binding Pockets in
Proteins. PLOS Comput Biol. 2013;9: e1003302. doi:10.1371/journal.pcbi.1003302

12.     Anand P, Chandra N. Characterizing the pocketome of Mycobacterium tuberculosis and
application in rationalizing polypharmacological target selection. Sci Rep 2014 41. 2014;4: 1–
17. doi:10.1038/srep06356

13.     Konc J, Janežič D. ProBiS-Fold Approach for Annotation of Human Structures from the
AlphaFold Database with No Corresponding Structure in the PDB to Discover New Druggable
Binding Sites. J Chem Inf Model. 2022;62: 5821–5829. doi:10.1021/acs.jcim.2c00947

14.     Pallante L, Cannariato M, Androutsos L, Zizzi EA, Bompotas A, Hada X, et al.
VirtuousPocketome: a computational tool for screening protein–ligand complexes to identify
similar binding sites. Sci Rep 2024 141. 2024;14: 1–14. doi:10.1038/s41598-024-56893-7

15.     Korkuć P, Walther D. Physicochemical characteristics of structurally determined
metabolite-protein and drug-protein binding events with respect to binding specificity. Front Mol
Biosci. 2015;2. doi:10.3389/fmolb.2015.00051

16.     Walther D. Specifics of Metabolite-Protein Interactions and Their Computational Analysis
and Prediction. In: Skirycz A, Luzarowski M, Ewald JC, editors. Cell-Wide Identification of
Metabolite-Protein Interactions. New York, NY: Springer US; 2023. pp. 179–197.
doi:10.1007/978-1-0716-2624-5_12

17.     Binkowski TA, Joachimiak A. Protein functional surfaces: Global shape matching and
local spatial alignments of ligand binding sites. BMC Struct Biol. 2008;8: 1–23.
doi:10.1186/1472-6807-8-45

18.    Simonovsky M, Meyers J. DeeplyTough: Learning Structural Comparison of Protein
Binding Sites. J Chem Inf Model. 2020;60: 2356–2366. doi:10.1021/acs.jcim.9b00554

19.    Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold
Protein Structure Database: massively expanding the structural coverage of protein-sequence
space with high-accuracy models. Nucleic Acids Res. 2022;50: D439–D444.
doi:10.1093/NAR/GKAB1061

20.    Dobson L, Szekeres LI, Gerdán C, Langó T, Zeke A, Tusnády GE. TmAlphaFold
database: membrane localization and evaluation of AlphaFold2 predicted alpha-helical
transmembrane protein structures. Nucleic Acids Res. 2023;51: D517–D522.
doi:10.1093/nar/gkac928

21.    Utgés JS, Barton GJ. Comparative evaluation of methods for the prediction of protein–
ligand binding sites. J Cheminformatics. 2024;16: 126. doi:10.1186/s13321-024-00923-z

22.    Bateman A, Martin MJ, Orchard S, Magrane M, Ahmad S, Alpi E, et al. UniProt: the
Universal Protein Knowledgebase in 2023. Nucleic Acids Res. 2023;51: D523–D531.
doi:10.1093/NAR/GKAC1052

23.    Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, et al.
InterPro in 2022. Nucleic Acids Res. 2023;51: D418–D427. doi:10.1093/nar/gkac993

24.    Barrio-Hernandez I, Yeo J, Jänes J, Mirdita M, Gilchrist CLM, Wein T, et al. Clustering
predicted structures at the scale of the known protein universe. Nature. 2023;622: 637–645.
doi:10.1038/s41586-023-06510-w

25.    Hastings J, De Matos P, Dekker A, Ennis M, Harsha B, Kale N, et al. The ChEBI
reference database and ontology for biologically relevant chemistry: enhancements for 2013.
Nucleic Acids Res. 2012;41: D456–D463. doi:10.1093/nar/gks1146

26.    Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al.
SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat Methods. 2020;17:
261–272. doi:10.1038/s41592-019-0686-2

27.    Ehrt C, Brinkjost T, Koch O. A benchmark driven guide to binding site comparison: An
exhaustive evaluation using tailor-made data sets (ProSPECCTs). PLOS Comput Biol. 2018;14:
e1006483. doi:10.1371/JOURNAL.PCBI.1006483

28.    Reim T, Ehrt C, Graef J, Günther S, Meents A, Rarey M. SiteMine: Large-scale binding
site similarity searching in protein structure databases. Arch Pharm (Weinheim). 2024;357:
2300661. doi:10.1002/ARDP.202300661

29.    Konc J, Janežič D. ProBiS algorithm for detection of structurally similar protein binding
sites by local structural alignment. Bioinformatics. 2010;26: 1160–1168.
doi:10.1093/BIOINFORMATICS/BTQ100

30.    Da Silva F, Desaphy J, Rognan D. IChem: A Versatile Toolkit for Detecting, Comparing,
and Predicting Protein–Ligand Interactions. ChemMedChem. 2018;13: 507–510.
doi:10.1002/cmdc.201700505

31.    Batista J, Hawkins PC, Tolbert R, Geballe MT. SiteHopper - a unique tool for binding site
comparison. J Cheminformatics. 2014;6: P57, 1758-2946-6-S1-P57. doi:10.1186/1758-2946-6-
S1-P57

32.    Ren J, Xie L, Li WW, Bourne PE. SMAP-WS: a parallel web service for structural
proteome-wide ligand-binding site comparison. Nucleic Acids Res. 2010;38: W441–W444.
doi:10.1093/nar/gkq400

33.    Wood DJ, Vlieg JD, Wagener M, Ritschel T. Pharmacophore Fingerprint-Based
Approach to Binding Site Subpocket Similarity and Its Application to Bioisostere Replacement. J
Chem Inf Model. 2012;52: 2031–2043. doi:10.1021/ci3000776

34.    Pedregosa F, Michel V, Grisel O, Blondel M, Prettenhofer P, Weiss R, et al. Scikit-learn:
Machine Learning in Python. J Mach Learn Res. 2011;12: 2825–2830.

35.    McInnes L, Healy J, Melville J. UMAP: Uniform Manifold Approximation and Projection
for Dimension Reduction. arXiv; 2020. Available: http://arxiv.org/abs/1802.03426

36.    Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, et al. Biopython: freely
available Python tools for computational molecular biology and bioinformatics. Bioinformatics.
2009;25: 1422–1423. doi:10.1093/BIOINFORMATICS/BTP163

37.    Lobry JR, Gautier C. Hydrophobicity, expressivity and aromaticity are the major trends of
amino-acid usage in 999 Escherichia coli chromosome-encoded genes. Nucleic Acids Res.
1994;22: 3174–3180. doi:10.1093/nar/22.15.3174

38.    Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a
protein. J Mol Biol. 1982;157: 105–132. doi:10.1016/0022-2836(82)90515-0

39.    Shrake A, Rupley JA. Environment and exposure to solvent of protein atoms. Lysozyme
and insulin. J Mol Biol. 1973;79: 351–371. doi:10.1016/0022-2836(73)90011-9

40.    Tien MZ, Meyer AG, Sydykova DK, Spielman SJ, Wilke CO. Maximum Allowed Solvent
Accessibilites of Residues in Proteins. Porollo A, editor. PLoS ONE. 2013;8: e80635.
doi:10.1371/journal.pone.0080635

41.    Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, et al. UCSF
ChimeraX: Structure visualization for researchers, educators, and developers. Protein Sci Publ
Protein Soc. 2021;30: 70–82. doi:10.1002/pro.3943

42.    J. D. Hunter. Matplotlib: A 2D Graphics Environment. Comput Sci Eng. 2007;9: 90–95.
doi:10.1109/MCSE.2007.55

43.    Waskom ML. seaborn: statistical data visualization. J Open Source Softw. 2021;6: 3021.
doi:10.21105/joss.03021

44.    Inc PT. Collaborative data science. Montreal, QC: Plotly Technologies Inc.; 2015.
Available: https://plot.ly

45.    Koonin EV, Wolf YI, Karev GP. Power Laws, Scale-Free Networks and Genome Biology.
Boston, MA: Springer US; 2006. doi:10.1007/0-387-33916-7

46.    van Nimwegen E. Scaling laws in the functional content of genomes. Trends Genet.
2003;19: 479–484. doi:10.1016/S0168-9525(03)00203-8

47.    Koehler Leman J, Szczerbiak P, Renfrew PD, Gligorijevic V, Berenberg D, Vatanen T, et
al. Sequence-structure-function relationships in the microbial protein universe. Nat Commun.
2023;14: 2351. doi:10.1038/s41467-023-37896-w

48.    Osadchy M, Kolodny R. Maps of protein structure space reveal a fundamental
relationship between protein structure and function. Proc Natl Acad Sci. 2011;108: 12301–
12306. doi:10.1073/pnas.1102727108

49.    Hegyi H, Gerstein M. The relationship between protein structure and function: a
comprehensive survey with application to the yeast genome. J Mol Biol. 1999;288: 147–164.
doi:10.1006/jmbi.1999.2661

50.    Hekkelman ML, De Vries I, Joosten RP, Perrakis A. Nature Methods nature methods
AlphaFill: enriching AlphaFold models with ligands and cofactors. 2023. doi:10.1038/s41592-
022-01685-y

51.    He X, Zhao L, Tian Y, Li R, Chu Q, Gu Z, et al. Highly accurate carbohydrate-binding site
prediction with DeepGlycanSite. Nat Commun. 2024;15: 5163. doi:10.1038/s41467-024-49516-
2

52.    Bibekar P, Krapp L, Peraro MD. PeSTo-Carbs: Geometric Deep Learning for Prediction
of Protein–Carbohydrate Binding Interfaces. J Chem Theory Comput. 2024;20: 2985–2991.
doi:10.1021/acs.jctc.3c01145

53.    Meller A, Ward M, Borowsky J, Kshirsagar M, Lotthammer JM, Oviedo F, et al.
Predicting locations of cryptic pockets from single protein structures using the PocketMiner
graph neural network. Nat Commun. 2023;14: 1177. doi:10.1038/s41467-023-36699-3

54.    Gavin A-C, Bösche M, Krause R, Grandi P, Marzioch M, Bauer A, et al. Functional
organization of the yeast proteome by systematic analysis of protein complexes. Nature.
2002;415: 141–147. doi:10.1038/415141a

55.    Huttlin EL, Bruckner RJ, Paulo JA, Cannon JR, Ting L, Baltier K, et al. Architecture of the
human interactome defines protein communities and disease networks. Nature. 2017;545: 505–
509. doi:10.1038/nature22366

56.    Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, et al. Protein complex
prediction with AlphaFold-Multimer. Bioinformatics; 2021. doi:10.1101/2021.10.04.463034

57.    Saldaño T, Escobedo N, Marchetti J, Zea DJ, Mac Donagh J, Velez Rueda AJ, et al.
Impact of protein conformational diversity on AlphaFold predictions. Valencia A, editor.
Bioinformatics. 2022;38: 2742–2748. doi:10.1093/bioinformatics/btac202

58.    Wayment-Steele HK, Ojoawo A, Otten R, Apitz JM, Pitsawong W, Hömberger M, et al.
Predicting multiple conformations via sequence clustering and AlphaFold2. Nat 2023 6257996.
2023;625: 832–839. doi:10.1038/s41586-023-06832-9

59.    Monteiro Da Silva G, Cui JY, Dalgarno DC, Lisi GP, Rubenstein BM. High-throughput
prediction of protein conformational distributions with subsampled AlphaFold2. Nat Commun.
2024;15: 2464. doi:10.1038/s41467-024-46715-9

60. Korkuć P, Walther D. Spatial proximity statistics suggest a regulatory role of protein
phosphorylation on compound binding. Proteins Struct Funct Bioinforma. 2016;84: 565–579.
doi:10.1002/prot.25001

61. Correa Marrero M, Mello VH, Sartori P, Beltrao P. Global comparative structural analysis
of responses to protein phosphorylation. Bioinformatics; 2024. doi:10.1101/2024.10.18.617420

62. Wright PE, Dyson HJ. Intrinsically disordered proteins in cellular signalling and
regulation. Nat Rev Mol Cell Biol. 2015;16: 18–29. doi:10.1038/nrm3920

63. Uversky VN. Intrinsic Disorder-based Protein Interactions and their Modulators. Curr
Pharm Des. 2013;19: 4191–4213. doi:10.2174/13816128113199230005

64. Mollica L, Bessa LM, Hanoulle X, Jensen MR, Blackledge M, Schneider R. Binding
Mechanisms of Intrinsically Disordered Proteins: Theory, Simulation, and Experiment. Front Mol
Biosci. 2016;3. doi:10.3389/fmolb.2016.00052

65. Herrera-Nieto P, Pérez A, De Fabritiis G. Small Molecule Modulation of Intrinsically
Disordered Proteins Using Molecular Dynamics Simulations. J Chem Inf Model. 2020;60: 5003–
5010. doi:10.1021/acs.jcim.0c00381

66. Mészáros B, Simon I, Dosztányi Z. Prediction of Protein Binding Regions in Disordered
Proteins. Casadio R, editor. PLoS Comput Biol. 2009;5: e1000376.
doi:10.1371/journal.pcbi.1000376

67. Brutscher B, Felli IC, Gil-Caballero S, Hošek T, Kümmerle R, Piai A, et al. NMR Methods
for the Study of Instrinsically Disordered Proteins Structure, Dynamics, and Interactions:
General Overview and Practical Guidelines. In: Felli IC, Pierattelli R, editors. Intrinsically
Disordered Proteins Studied by NMR Spectroscopy. Cham: Springer International Publishing;
2015. pp. 49–122. doi:10.1007/978-3-319-20164-1_3

68. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, et al. Accurate structure
prediction of biomolecular interactions with AlphaFold 3. Nat 2024. 2024; 1–3.
doi:10.1038/s41586-024-07487-w

# Supporting Information

**960** **S1 Fig. Violin plots for all species used in the study for the pLDDT of proteins with and**
**961** **without binding sites.**

**962** The highest median pLDDT for proteins with and without binding sites was observed for ECOLI,
**963** with 93.24 and 90.07, respectively. The lowest median pLDDT of proteins with binding sites was
**964** observed for MAIZE with 79.41 and for ORYSJ of proteins without binding sites with 67.05. The
**965** median pLDDT values of proteins for the other species range from 80.95 to 85.40 for proteins
**966** with binding site and from 68.48 to 75.75 for proteins without any binding site predicted by
**967** P2Rank.

**968** **S2 Fig. tSNE plots (perplexity = 10) for all predicted pockets of all 11 species colored by**
**969** **different properties.**

**970** **S3 Fig. tSNE plots (perplexity = 20) for all predicted pockets of all 11 species colored by**
**971** **different properties.**

**972** **S4 Fig. tSNE plots (perplexity = 30) for all predicted pockets of all 11 species colored by**
**973** **different properties.**

**974** **S5 Fig. tSNE plots (perplexity = 40) for all predicted pockets of all 11 species colored by**
**975** **different properties.**

**976** **S6 Fig. tSNE plots (perplexity = 100) for all predicted pockets of all 11 species colored by**
**977** **different properties.**

**978** **S7 Fig. UMAP plots (n_neighbors = 10) for all predicted pockets of all 11 species colored**
**979** **by different properties.**

**980** **S8 Fig. UMAP plots (n_neighbors = 20) for all predicted pockets of all 11 species colored**
**981** **by different properties.**

**982** **S9 Fig. UMAP plots (n_neighbors = 50) for all predicted pockets of all 11 species colored**
**983** **by different properties.**

**984** **S10 Fig. UMAP plots (n_neighbors = 100) for all predicted pockets of all 11 species colored**
**985** **by different properties.**

**986** **S11 Fig. UMAP plots (n_neighbors = 200) for all predicted pockets of all 11 species colored**
**987** **by different properties.**

**988** **S12 Fig. PCA plots for all predicted pockets of all 11 species colored by different**
**989** **properties.**

**S13 Fig. ICA plots for all predicted pockets of all 11 species colored by different properties.**

**S14 Fig. tSNE plot with the first low probability region highlighted in orange.**

Pockets that are found in the first low probability region are highlighted in orange. All other pockets
are colored in orange

**S15 Fig. tSNE plot colored by species and binned entropy.**

(A) tSNE projection colored by different species. All pockets in their respective tSNE embedding
space, (B) Binned entropy, S (see Materials and Methods), heatmap plot, capturing for every
square bin the species diversity with low entropy, S, values indicating low diversity and high
values corresponding to respectively increased diversity. All grid cells containing less than 20
pockets are colored black.

**S1 Table.** Additional information for all predicted pockets, including the number of FoldSeek (FS)
clusters per species, the ratio of pockets per protein, the number of pockets before filtering, and
the number of pockets that did not pass the respective quality check.

**S2 Table.** Number of unique pockets, singletons, and real clusters with threshold 5 applied to the
ProBiS alignment score.

**Found**

- 10.21%
- 9.55%
- 4.70%
- 2.23%
- 1.48%
- 0.72%
- 0.68%
- 0.61%
- 0.32%
- 0.11%
- 0.00%
- 69.39%

**Not found**

- 20.95%
- 4.04%
- 2.76%
- 1.26%
- 0.96%
- 0.88%
- 0.86%
- 0.32%
- 0.07%
- 0.04%
- 0.02%
- 67.84%

Legend:
- Nucleobase/nucleoside/nucleotide (10336)
- Inorganic ion (8456)
- Other compound (1410)
- Carbohydrate & derivatives (651)
- AA & derivatives (395)
- Lipid (267)
- Hetero nuclear cluster (517)
- Oligo- & polypeptide (171)
- Monosaccharide & derivatives (105)
- Glycan (41)
- Heme (15)
- Nucleic acid (7)