Consumer resilience suppresses the recovery of overgrazed ecosystems.

Abstract

1. Many heterotroph species perish when faced with severe food limitation, others can

persist, adapt, and thrive. Sea urchins are emblematic of this paradox: they can overgraze

kelp forests to form barren habitats, but can then survive for decades in these nutritionally

depauperate seascapes. Understanding the mechanisms enabling persistence under

starvation, and rapid recovery when food returns, provides insight into how consumer

resilience shapes ecosystem dynamics.

2. We quantified how food abundance, quality, deprivation, and reintroduction influence

bioenergetic performance in the red sea urchin (*Mesocentrotus franciscanus*), integrating

field observations of kelp forest and barren populations with a controlled feeding

experiment. We measured respiration, feeding rates, gonadal growth, and fatty acid

biomarkers to test how habitat history and diet jointly govern metabolic plasticity and

nutrient assimilation.

3. Resting metabolic rates (RMR) were nearly twofold higher in kelp forest urchins than

barrens conspecifics, yet feeding rates were equivalent across habitats, indicating that

metabolic depression does not constrain food intake. Reciprocal shifts emerged in the

experiment: starvation reduced RMR and lipid reserves in kelp forest urchins, while

feeding elevated both traits in barrens urchins to levels comparable with kelp forest

conspecifics. These results demonstrate rapid physiological compensation in response to

both food deprivation and reintroduction.

4. Diet quality strongly modulated performance. Urchins fed nutritionally poor

monospecific diets consumed more biomass and calories than those on diverse,

polyunsaturated fatty acid (PUFA)-rich diets, but did so with markedly lower efficiency

of conversion to gonadal tissue. Fatty acid assimilation revealed that starvation elevated

bacterial and biofilm biomarkers in tissues, whereas algal diets enriched essential PUFA

profiles, particularly when diets were diverse. These results highlight that both quantity

and quality of food influence consumer recovery trajectories, with nutritional geometry

shaping efficiency of energy and nutrient use.

5.   Together, our findings show that *M. franciscanus* exhibits pronounced metabolic

resilience, allowing persistence in barren habitats and rapid reactivation of grazing and

reproduction when food becomes available. This work links nutritional ecology to

ecosystem feedbacks by showing how compensatory feeding and metabolic flexibility

enable consumers to maintain pressure on primary producers, thereby influencing the

stability, hysteresis, and recovery of degraded ecosystems.

1. Introduction

Variation in food availability can directly shape the fitness of consumers and associated

food web dynamics. Such variation results from, for example, bottom-up changes in primary

production (Edwards et al., 2020), shifts in availability of allochthonous food resources (Leal et

al., 2022), and/or strong consumer pressure (Gangal et al., 2021). Declines in the quantity or

quality of available food can reduce rates of reproduction, juvenile survival, growth, and adult

survival in many taxa (Okamoto et al., 2012, Mduma et al., 1999, Olsen et al., 2011). A diverse

range of animals across trophic levels, taxonomic groups, and ecosystems (Randall, 1961,

Silliman and Zieman, 2001, Mysterud, 2006, Kayal et al., 2012, Ling et al., 2015) are known to

periodically graze down and deplete resource productivity leading to barren landscapes of

nutritionally poor, scarce, or low productivity food resources. While many fields of study have

focused on how these trophic interactions shape population dynamics, less is known about the

mechanistic basis of consumer resilience to nutritional deficits. If consumers are to avoid death

by starvation, they must emigrate, switch their diet to alternative foods, and/or cope (McCue et

al., 2017). Among those that cope, it remains unclear how metabolic plasticity and resilience

interact with changes in the abundance and quality of food to shape energetic traits and fitness.

Recent research in nutritional ecology has shown that consumers may regulate intake not

simply to maximize calories, but to balance multiple nutrients simultaneously (Simpson et al.,

2004, Raubenheimer et al., 2009). Across diverse taxa, from beetles to primates, individuals can

adjust their intake of macronutrients to fulfill physiological requirements for individual

micronutrients, even at a cost to growth or survival (Jensen et al., 2012, Harrison et al., 2014,

Solon-Biet et al., 2014, Tait et al., 2014). Mobile consumer species may respond to reductions in

food density or productivity by expanding or shifting their geographic range (Brown and Kotler,

2004) or reducing activity (Storey, 2015). Consumers may also expand their dietary breadth

(Tinker et al., 2012) capitalize on exogenous food subsidies (Britton-Simmons et al., 2009), or

engage in a last-gasp reproductive bout before dying (Kirkwood and Rose, 1991). In addition to

these behavioral shifts, a wide range of eukaryotes avoid minimize the effects of starvation or

malnutrition by mobilizing energy reserves for basic functions and/or prioritizing maintenance

at the cost of structural growth and/or reproduction (Vilela et al., 2008, Lesage et al., 2001,

Chippindale et al., 1993, Ellers, 1995). While heterotrophs can conserve energy and reduce

metabolic demands under food deprivation (Storey, 2015, Spindel et al., 2021), the effects of

nutritional limitation and resurgence on such metabolic flexibility and the attendant

consequences for performance remain poorly studied (McCue et al., 2017). Metabolic rate

correlates strongly with specific physiological processes like respiration rate (Glazier, 2015), but

it is less clear how metabolic demands feed-back to affect maximum potential food intake and

investment in reproduction. For example, while resting metabolic rate (RMR) reflects baseline

energetic maintenance, food intake must supply not only energy for homeostasis but also

essential nutrients for the growth of reproductive tissues (Sterner and Elser, 2002). A

comprehensive understanding of an organism's metabolic flexibility under food limitation

therefore requires integration of energetics with nutrient balance. Beyond the capacity of

consumers to modulate metabolic demands under scarcity, the quality and composition of

available resources emerge as equally critical determinants of performance and recovery.

Shifts in both resource quantity and quality affect consumer energetics and performance

even when caloric intake is sufficient. Macronutrient deficiencies can alter feeding behavior,

growth, and longevity in a wide range of organisms (Solon-Biet et al., 2014, Cruz-Rivera and

Hay, 2000). Some micronutrients are "essential" in the diets of many consumers (Parrish, 2009).

For example, highly unsaturated fatty acids (HUFAs), such as docosahexaenoic acid (DHA), can

be critical biomolecules for successful reproduction, growth, and survival (Fuiman and Perez,

2015, Ruiz et al., 2021). Yet many consumers lack the capacity to synthesize these essential

nutrients (Twining et al., 2021). When quality of available food is low, consumers may

compensate by increasing consumption to meet metabolic and nutritive demands. This

phenomenon is a central prediction of nutritional geometry and has been documented across both

terrestrial and marine systems (Raubenheimer et al., 2009, Raubenheimer and Simpson, 1993,

Cruz-Rivera and Hay, 2000, Berner et al., 2005, Fink and Von Elert, 2006, Jochum et al., 2017).

This framework suggests that the balance of nutrients, not energy alone, governs performance

and recovery (Raubenheimer and Simpson, 1993, Simpson et al., 2004, Raubenheimer et al.,

2009, Simpson and Raubenheimer, 2012).

Sea urchins provide a powerful model to investigate metabolic flexibility under food

limitation. They are dominant herbivores that often graze down highly productive and biodiverse

algal assemblages (Schiel and Foster, 2015) leaving behind low-productivity barrens which can

persist for decades (Filbee-Dexter and Scheibling, 2014). Survival rates of some sea urchins in

barrens have been hypothesized to be high as a result of phenotypic plasticity (Edwards and

Ebert, 1991). Recent observations demonstrated a 50% decline in RMR between sea urchins

occupying food-rich versus food-poor habitats (Spindel et al., 2021). Understanding how such

metabolic depression in food-poor habitats affects the capacity of sea urchins to quickly resume

consumption and reproduction when food becomes available provides a direct test of theories of

metabolic flexibility and nutritional ecology. Moreover, the timescale over which animals

respond to resurgent food availability and how different food qualities influence the recovery of

reproductive capacity, energy stores, and RMR remains poorly understood (McCue et al., 2017).

We focused on the barren-forming red sea urchin, *Mesocentrotus franciscanus*, because

of its important roles in structuring nearshore algal ecosystems, commercial fisheries, and

indigenous culture (Rogers-Bennett and Okamoto, 2020). We addressed two objectives: (1)

quantifying feeding rates and metabolic responses following starvation as a function of food

quality, and (2) characterizing how habitat and diet affect assimilated FA biomarker profiles as a

step toward linking experimental nutrition with wild dietary dynamics. Based on large

differences in RMR between barren and kelp forest urchins (Spindel et al., 2021), we

hypothesized that (a) RMR would be highly flexible as a function of declines and resurgences in

food availability, (b) these metabolic shifts would be accompanied by shifts in energy stores and

reproductive capacity, and (c) calorically rich but nutritionally poor diets would induce

compensatory feeding and yield less efficient recovery than nutritionally-rich diets.

2. Materials and Methods

We evaluated the effects of habitat of origin (kelp forest vs. barrens) and experimental diet on both bioenergetics (i.e., metabolism, grazing rate, growth, and gonad production) and FA dietary biomarkers of *M. franciscanus*. Field assays provided natural ecological context for our laboratory experiment and included measurements of RMR, body size specific gonad mass, and FA biomarker profiles in each habitat. We then subjected individuals from neighboring barrens and kelp forest habitats to a six-week aquarium experiment where we controlled their diets and measured bioenergetic responses and FA biomarker profile outcomes.

Field Collections

We collected individuals using SCUBA from Surge Narrows, British Columbia, (50.22°N, 125.16 °W) from a depth of 2-3 m and 10-12 m relative to mean tide from kelp forest and nearby barrens habitat, respectively (n = 30 per habitat, 60 total; mean test diameter = 69.4 mm ± 1.02 mm SE) in August of 2019. We immediately transported collected individuals in insulated coolers to the Hakai Institute Quadra Island Ecological Observatory (HIQIEO) where we transferred them to flow-through sea tables and allowed them to recover for a period of 48 hours. We collected additional individuals from similar depths for sacrificial sampling to compare habitat-specific gonadal mass allometry and fatty acid composition at Surge Narrows with additional study sites in Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve, and Haida Heritage Site on Haida Gwaii, including Faraday (52.61° N, 131.49° W) and Murchison (52.60° N, 131.45° W) Islands.

Aquarium Setup

We used outdoor sea tables supplied with flow-through treated seawater for the laboratory experiment. Seawater was pumped from a depth of 20 m, filtered at 100 µm a. UV

sterilized,. We subdivided each sea table into multiple 17L aquaria (n = 60 aquaria) using plastic

bins, each supplied with independent flow-through seawater (flow = 2.20 Lhr^{-1}; turnover = 3.12

d^{-1}) via irrigation plumbing (**figure S1**). Temperature, salinity, and pH were recorded daily

(temperature = 13□ ± 1.01□ SE, salinity = 28.85 ppt ± 0.35 ppt SE, pH = 7.61 ± 0.13 SE). We

haphazardly assigned individual urchins to solitary chambers in each aquarium following the

recovery period

Acclimation and Experimental Treatments

We minimized relocation stress by allowing the urchins to acclimate to aquarium

conditions for 48 hours prior to initiating treatments. We assigned experimental subjects to one

of three diet treatments: 1) Monospecific diet (*Nereocystis luetkeana*), *ad libitum*; 2) Diverse

diet (*N. luetkeana, Ulva sp., Chondracanthus corymbiferus,* and *Dilsea californica*), *ad libitum*;

and 3) Starvation, resulting in a fully-crossed factorial design (n = 10 urchins per habitat-diet

treatment combination). These algal taxa have similar caloric content, C:N ratios, and protein

content but substantially different concentrations of nutritionally valuable fatty acids (electronic

supplemental material; **table S11**). We collected fresh macroalgae for diet treatments from a

depth of 3-6 m at nearshore sites around Quadra Island (*N. luetkeana* at Quathiaski Cove

50.04˚N, 125.22 ˚W, *Ulva sp., C. corymbiferus,* and *D. californica* at Wa Wa Kie Beach

50.04˚N, 125.17 ˚W) and transferred collected algae to flow-through sea tables at the HIQIEO.

We prevented decomposition of macroalgal tissue by collecting fresh macroalgae immediately

prior to and periodically throughout the experiment, only using algae collected within one week.

To standardize composition of fresh macroalgal diets we only used thallus tissue which was free

of reproductive patches, edges prone to decomposition, and conspicuous epibionts. We

maintained experimental treatments for thirty-three days to allow for controlled diets to be

assimilated into new urchin biomass.

Energetics

Respirometry

We quantified respiration rate as a proxy for metabolic rate, focusing on resting

metabolic rates (RMR). We selected RMR to avoid the confounding effects of specific dynamic

action (Lighton, 2018); to control for this we starved focal subjects for 48 hours prior to

respirometry then immediately resumed prescribed feeding after respirometry sessions. We

quantified respiration rates using custom-built sealed acrylic chambers (Spindel et al., 2021). To

account for the displacement volume of the urchins when calculating respiration rates, we

subtracted internal test volume of the urchins (V – calculated from test diameter (D) and height

(H) assuming oblate spheroid geometry ($V = 4/3\pi\ D^2H$)) from the total chamber system volume.

We accounted for background oxygen dynamics using an identical empty respiration chamber in

the same environmental conditions in parallel with every set of three respiration chambers

containing urchins. We conducted reproducible and transparent quality control on oxygen time

series data using the R package respR (Harianto et al., 2019).

We standardized RMR with total metabolically active biomass which we measured as

ash-free dry mass (AFDM). AFDM represents non-skeletal soft tissue as the difference between

dry mass and post-combustion ash mass. We first cracked the test of the urchins and discarded

the coelomic fluid, then dried the carcasses for 24 hours at 60 ☐ in a drying oven, then weighed

the dried carcasses. We measured post-combustion ash mass by combusting dried carcasses for

six hours at 450 ☐ in a muffle furnace, then weighed the ashes of each carcass (scale resolution

=0.01 mg). To contextualize measurements of RMR made on experimental subjects maintained

in the laboratory with natural variation, we also sampled the wild population at the beginning and

end of the experiment from the same collection site.

Feeding Rate and Assimilation Efficiency

We measured food consumption and assimilation efficiency using repeated trials

spanning from day 7-14 and 23-33 in shaded sea tables to limit macroalgal growth. For each

trial, we measured the wet mass of food at the beginning and endpoints, as well as the mass of

feces at the endpoints. We standardized wet weight measurements of algae by spinning thallus

tissue in a colander for 30 s prior to recording a mass measurement. We approximated algal dry

weights for each of the four species of algae by generating species specific linear regressions

between wet and dry weights (electronic supplemental material; **figure S3**). We calculated mean

feeding rate as the difference in dry weight of algae from the beginning and end of each feeding

cycle divided by the amount of time in the feeding cycle (electronic supplemental material;

method S2).

Gonadal Mass and Food Conversion Efficiency

To quantify shifts in gonadal mass in experimental subjects, we calculated the difference

between an estimated initial and measured final state. To estimate initial gonadal mass, we used

body size and gonadal mass measurements from representative wild individuals in barrens and

kelp forest habitats at Surge Narrows to model the effects of habitat and body size on gonadal

dry mass. We measured test volume and gonadal dry mass after heating samples at 60 ▢ for 24 h

in a drying oven to correct for variation in water content. We sampled a representative size

gradient of benthic stage individuals spanning small through large body sizes at Surge Narrows

in the barrens and kelp forest habitats at the beginning of the experiment and combined these

measurements with identical measurements taken three months prior (Spindel et al., 2021) (kelp

forest habitat: n = 49 individuals, minimum test volume = 26 mL, maximum test volume = 668

mL; barrens habitat: n = 59, minimum test volume = 18 mL, maximum test volume = 452 mL).

We quantified food conversion efficiency as a ratio between increase in dry gonadal

biomass and total dry food consumption between the beginning and end of the experiment (Brett

and Groves, 1979). To infer the beginning habitat specific dry gonadal biomass of experimental

subjects for a given body size, we used a model fit to data collected from wild sacrificial samples

in the beginning of the experiment collected from the same source populations as the

experimental subjects. We used an inverse problem approach to estimate food conversion

efficiency, combining error in measurement models estimating treatment specific feeding rates

and gonad mass. To estimate treatment specific feeding rates, we fit an error in measurement

model to data from replicate feeding trials, combined with an error in measurement model of

gonad mass fit to data from wild individuals at the beginning of the experiment and experimental

subjects at the end of the experiment.

Total Lipid Content

We evaluated changes in biomass-specific energy reserves by comparing the lipid content

of gonadal tissue in wild individuals of comparable size to experimental subjects at the beginning

of the experiment with the lipid content of gonadal tissue of experimental subjects at the end of

the experiment. To prevent degradation of lipids, we immediately flash-froze all harvested tissue

samples using liquid nitrogen, then transferred frozen samples to -80 □ freezers for storage and

later lipid extraction and FA composition profiling (see below). To quantify lipid content of

gonadal tissue, we extracted lipids from gonadal tissue biopsies (Schram et al., 2018): tissue

samples were lyophilized for 48 h then homogenized. Following homogenization, lyophilized

tissue samples were digested in 2 mL chloroform, then sonicated, vortexed, and centrifuged

(3000 rpm for 5 m) twice in a 4:2:1 chloroform:methanol:0.9% NaCl solution. After each of

these cycles, the lower chloroform layer containing organic compounds was pipetted out and

pooled. Subsamples were collected from this pooled solution for gravimetric analysis of total

lipid content (Kainz et al., 2017).

Fatty Acid Composition

To quantify shifts in FA composition associated with nutritional history and diet, we

compared the FA composition of lipids extracted from tissue biopsies collected from individuals

in their natural habitat and experimental subjects fed controlled diets of known FA composition.

We isolated FAs from the lipid solution extracted from tissue samples. To generate a FA-based

dietary resource library, we extracted and characterized FAs from thallus tissue samples

harvested from the four species of algae supplied in our diet treatments, avoiding epibiota from

algal thallus tissue prior to storage. To quantify shifts in FA composition of gonadal tissues, we

estimated the state of gonadal tissue FA composition at the beginning of the experiment using

gonadal tissue biopsies from wild individuals of comparable size to experimental subjects

collected at the same time and from the same place as the experimental subjects (Foster et al.,

2015) and compared this estimated initial state against biopsies of gonadal tissue collected from

dissected experimental subjects at the end of the experiment.

To quantify the FA composition of tissue samples, we isolated and analyzed FAs from

extracted lipids. Briefly, total lipids were extracted using a modified Folch method and

derivatized into fatty acid methyl esters (FAME) using methods described in (Schram et al.,

2018) and Thomas et al. (2020). Samples were added to chloroform:methanol, sonicated,

vortexed, and centrifuged to separate the organic phase, which was removed and evaporated to

dryness under nitrogen gas. The extraction process was repeated after resuspending dried lipid in

2 mL of chloroform. To produce FAME, the dried lipid extracts were resuspended in a 1:2 ratio

of toluene and 1% sulfuric acid in methanol and incubated at 90°C for 90 minutes. Following

transesterification, the samples were allowed to cool, and the acid was neutralized and hexane

was added and the samples were vortexed and centrifuged to capture the FAME. Dried FAME

were resuspended in 1.5 mL hexane and stored at -20°C. FAME were analyzed with a gas

chromatograph-mass spectrometer (Shimadzu GCMS model QP-2020) fitted with a DB-23

column (30 × 0.25 mm × 0.15 μm, Agilent, Santa Clara, CA, USA), using helium as the

carrier gas. The heating program described in Taipale et al. (2016) was utilized to ensure

sufficient separation between chromatogram peaks. Fatty acids were identified using external

standards (GLC 566C, Nu-Chek Prep, Elysian, MN, USA) and mass spectrometry.

Replication Statement

In this study, our inferences are drawn at the individual urchin level, as we tested how

habitat of origin and diet treatments influence physiological performance and nutritional

assimilation in *M. franciscanus*. Treatments were applied at two main scales: (i) habitat of origin

(urchins collected from kelp forest vs. barren habitats) and (ii) dietary treatments (starvation,

monospecific diet, or diverse algal diet) applied to individual urchins housed in independent

flow-through chambers. Replication was based on multiple individuals per treatment, while

repeated measurements (feeding rate, assimilation efficiency, fatty acid composition) were taken

from the same individuals to improve precision but not as independent replicates. Chamber

identity was modeled as a group-level effect to account for potential non-independence.

Table 1. Replication schema for laboratory and field-based analyses.

Scale of inference	Scale at which the factor of interest is applied	Number of replicates at the appropriate scale

Individual urchins	Habitat of origin (kelp forest vs. barren)	30 kelp forest urchins, 30 barren urchins in manipulative experiment. An additional 24 individuals from kelp forests and 13 individuals from barrens were sampled at the beginning of the experiment, but not subjected to experimental treatments (total n = 97).
Individual urchins	Diet treatments (applied to individuals in independent chambers: starvation, monospecific diet, diverse algal diet)	10 individuals per diet x habitat combination (total n = 60). An additional 24 individuals from kelp forests and 13 individuals from barrens were sampled at the beginning of the experiment, but not subjected to experimental treatments.
Individual urchins	Time (repeated measures: start vs 33 days)	Not independent replicates; subsamples for within-individual precision (total n = 60).

Habitat (gonad FA profiling in wild populations)	Depth: 2-3 m and 10-12 m relative to mean tide from kelp forest and nearby barrens habitat, respectively	3 islands, 2 depths each (2-3 m = kelp forest, 10-12 m = barrens), 10-15 urchins per combination (total n = 76).

Data Analyses

Univariate

We estimated the population level effects of body size, habitat of origin, and diet on

bioenergetic and FA outcomes using Bayesian generalized linear models. RMR models also

included a group-level effect of respiration chamber and tube feet FA models included a group-

level effect of individual to account for repeated measures. We used a Gamma likelihood and log

link function to model these continuous positive outcomes. We accounted for uncertainty in

model structures by approximate leave-one-out cross validation (loo) and stacking weights

(Vehtari et al., 2016, Yao et al., 2018), a method designed to maximize predictive accuracy. To

evaluate goodness of fit for our models, we conducted graphical posterior predictive checks,

ensuring that model simulations did not systematically differ from empirical data (Gelman et al.,

2013) and estimated Bayesian R^2 values(Gelman et al., 2019). We evaluated the sensitivity of

parameter estimates and model fits to choices of priors by systematically altering priors and

refitting models. Subsequently, we tested hypotheses using Bayes Factors (BF) (Kass and

Raftery, 1995) computed from marginal likelihoods using the R package bridgesampling

(Gronau et al., 2020). We contrasted estimated marginal means (EMMs) to explore differences

between levels of factors while accounting for interactions and adjustments of covariates,

expressed in terms of highest posterior density (HPD) intervals the R package emmeans (Lenth

et al., 2024). We estimated model posteriors using Stan (Team, 2022) via the R (Team, 2024)

package brms (Bürkner, 2017) with 16,000 iterations across four chains, discarding the first half

of the iterations per chain as a warm-up, resulting in a posterior sample of 32,000 iterations for

each outcome. To ensure convergence, we verified that Rhat (the potential scale-reduction

factor) was no higher than 1.01 and the minimum effective sample size (n_{eff}) was greater than

1,000 (Gelman et al., 2013).

We estimated starting body size and habitat specific gonadal biomass of experimental

subjects using a model fit to data collected from sacrificing individuals at the beginning of the

experiment from the same source populations as the experimental subjects. To evaluate gonad

production during the experiment, we combined this estimated initial state with measurements of

per capita food consumption and final gonadal biomass. Because both consumption and gonadal

measurements were subject to observational error, we implemented a probabilistic framework

that explicitly propagated this uncertainty. In this framework, gonad production was modeled as

a function of body size, habitat of origin, and diet, while incorporating error terms for both

feeding rate and gonad mass to reflect their respective measurement uncertainty (electronic

supplemental material; **method S2**).

Multivariate

We visualized multivariate patterns associated with habitat of origin and diet using non-

metric multidimensional scaling ordination (NMDS) and tested for significance of predictors

using a multivariate analysis of variance. We accounted for the compositionality of fatty acid

data using Aitchison dissimilarity, which implements a centered log ratio transformation

(Aitchison, 1982). NMDS were fit using the package vegan (Oksanen et al., 2007) in R (Team,

2024). Ordination stress and the number of dimensions are reported within each NMDS plot. We

fit correlation vectors for constituent fatty acids, such that vector lengths were scaled by

correlation coefficients with respect to NMDS ordination scores for each data point. Fitted fatty

acid vector directions indicated the ordination space toward which the focal fatty acid changed

most rapidly and had the greatest correlation with the ordination configuration (Oksanen et al.,

2023). We identified fatty acids that contributed most to the observed multivariate separation by

habitat using a SIMPER analysis (Clarke, 2006). To assess significance of multivariate additive

and interaction effects of habitat of origin and diet while accounting for heteroskedasticity and

unequal sample sizes, we ran W_d^* (Hamidi et al., 2019) tests.

3. Results

Energetics

Resting metabolic rate

Resting metabolic rate (RMR) scaled positively with body size and was strongly

influenced by both habitat of origin and diet (Figure 1). Wild kelp forest urchins had nearly twice

the RMR of barrens conspecifics (ratio = 1.978, 95% HPD: 1.613-2.380). Experimental

manipulations produced reciprocal shifts: kelp forest urchins depressed their RMR after 33 days

of starvation (ratio = 0.560, 95% HPD: 0.404–0.731) to levels similar to barrens individuals,

while barrens urchins increased their RMR when fed to levels comparable with kelp forest

conspecifics. After 33 days, habitat differences were no longer detectable: kelp vs. barrens

comparisons overlapped zero both under starvation (ratio = 1.135, 95% HPD: 0.749-1.525) and

feeding (monospecific ratio = 1.058, 95% HPD: 0.691-1.430; diverse ratio = 0.985, 95% HPD:

0.661-1.342). Feeding produced especially strong effects in barrens individuals: RMR increased

more than threefold relative to wild (monospecific ratio = 3.182, 95% HPD: 2.355-4.073; diverse

ratio = 3.280, 95% HPD: 2.199-4.431). In contrast, kelp forest urchins responded strongly to

starvation but not feeding, with depressed RMR under starvation (ratio = 0.560, 95% HPD:

0.404-0.731) and high values under either diet (monospecific ratio = 3.060, 95% HPD: 2.040-

4.177). Effects of monospecific versus diverse algal diets were largely indistinguishable (kelp

ratio = 0.872, 95% HPD: 0.586-1.179; barrens ratio = 0.937, 95% HPD: 0.610-1.272). Full

statistical comparisons are presented in the electronic supplementary material (Table S1, Figure

S4).

Feeding rate

Diet quality was the strongest predictor of consumption (Figure 2). At the start of the

experiment, feeding rates did not differ between habitats or diets (ratio = 1.017, 95% HPD:

0.784-1.240). By the end, individuals fed a diverse diet consumed less total biomass and calories

than those on a monospecific diet, despite similar per capita intake earlier (ratio = 0.553, 95%

HPD: 0.425-0.690). Conversely, polyunsaturated fatty acid (PUFA) intake was initially higher

under diverse diets but converged across treatments after 33 days. Statistical model selection

details are provided in Tables S2–S4.

Assimilation efficiency

Assimilation efficiency was generally stable across treatments. Neither body size, habitat

of origin, nor diet produced consistent differences, although there was a slight tendency for

efficiency to increase over the course of the experiment (odds ratio = 1.16, 95% HPD 0.86-1.45).

These findings suggest that differences in performance between diets were driven primarily by

consumption and conversion efficiency, not assimilation itself (Table S5).

Gonad mass and lipid content

For a given body size, kelp forest urchins had larger gonads and greater lipid content than

barrens conspecifics in the wild (gonad mass: ratio = 1.89, 95% HPD: 1.34–2.46; lipids: ratio =

1.37, 95% HPD: 1.03–1.74). Starvation reduced both metrics regardless of habitat (kelp gonad

mass: ratio = 0.66, 95% HPD: 0.40–0.84; kelp lipids: ratio = 0.734, 95% HPD: 0.55–0.93;

barrens gonad mass: ratio = 0.66, 95% HPD: 0.45–0.91; barrens lipids: ratio = 0.57, 95% HPD:

0.37–0.77). Feeding restored gonads and lipids in both habitats. In barrens urchins, feeding

increased gonadal biomass and lipids relative to wild conspecifics (monospecific: gonad mass

ratio = 1.47, 95% HPD: 1.00–1.97, lipids ratio = 1.53, 95% HPD: 1.09–2.01; diverse: gonad

mass ratio = 1.38, 95% HPD: 0.95–1.87, lipids ratio = 1.63, 95% HPD: 1.07–2.23). In contrast,

fed kelp forest urchins resembled wild conspecifics, suggesting they already had high baseline

energy reserves (Figure 3a). Model comparisons supporting these results are reported in Tables

S6–S7.

Food conversion efficiency

Urchins fed diverse diets converted food to gonadal tissue almost three times more

efficiently than those fed monospecific diets, regardless of habitat (ratio = 2.81, 95% HPD: 1.91–

3.81; Figure 3b). This demonstrates that diet quality strongly influences the efficiency of

reproductive recovery (Table S8).

Fatty acid composition

Urchins from Faraday, Murchison, and Quadra Islands exhibited distinct fatty acid

profiles by habitat of origin, and these patterns were sensitive to diet treatments and starvation.

Parallel responses were observed in both gonads (figure 4 & 5; electronic supplemental material,

figure S7) and tube feet (electronic supplemental material, figure S8). SIMPER analysis

identified specific fatty acids associated with habitat and diet (figure 4). Among these, kelp

biomarkers (oleic, linoleic, stearidonic, and α-linolenic acids) and biofilm biomarkers (odd-chain

fatty acids and the diatom marker palmitoleic acid) were particularly diagnostic. In the field, kelp

forest urchins consistently had higher kelp biomarker and lower biofilm biomarker

concentrations than neighboring barrens urchins. In the laboratory, controlled macroalgal diets

increased kelp biomarker concentrations across habitats. Biofilm biomarkers were similar

between starved urchins and barrens urchins in the field, but for kelp forest urchins, starvation

more than doubled biofilm biomarker concentrations (ratio = 2.50, 95% HPD = 1.66–3.37).

In gonad tissue, fatty acid profiles differed significantly by habitat (W*d = 11.18, p =

0.001, nrep = 999), diet (W*_d = 10.70, p = 0.001, nrep = 999), and their interaction (W*d =

18.00, p = 0.001, nrep = 999). Kelp biomarker concentrations were 27.5% higher in urchins fed a

monospecific diet (ratio = 1.28, 95% HPD = 1.07–1.48) and 45.5% higher in those fed a diverse

diet (ratio = 1.46, 95% HPD = 1.22–1.70) relative to wild individuals. For biofilm biomarkers,

credible intervals for differences between barrens urchins in the field and experimentally starved

urchins overlapped zero. Additional details of fatty acid assimilation and trophic modification

are provided in the electronic supplemental material.

Discussion

Our study demonstrates that *M. franciscanus* exhibits striking physiological plasticity in

response to variation in food quantity and quality. RMR depressed under starvation but

rebounded rapidly when food became available, while feeding rates and tissue composition were

strongly shaped by diet quality. Individuals fed diverse algal diets converted food to gonadal

tissue more efficiently than those on monospecific diets, and FA profiles confirmed assimilation

of complementary nutrients. Together, these results provide empirical evidence for principles

long emphasized in nutritional ecology: consumers respond not only to the quantity of food but

also to its nutrient balance (Raubenheimer and Simpson, 1993, Simpson et al., 2004,

Raubenheimer et al., 2023).

Classic models of consumer resource dynamics predict that overgrazing leads to

consumer population decline, potentially allowing resources to recover. However, the ability of

M. franciscanus to enter hypometabolic states and then rapidly restore RMR and gonadal mass

when food reappears suggests an alternative: consumers can persist in food-poor environments

and resume grazing with little delay once resources return. This echoes prior work showing

metabolic depression and recovery in sea urchins (Dolinar and Edwards, 2021, Okamoto, 2014)

and underscores the role of compensatory physiological responses in stabilizing consumer

populations.

Compensatory feeding emerged as a central mechanism in our study. Urchins maintained

similar caloric intake across treatments but required higher consumption on monospecific, lower-

quality diets to achieve reproductive recovery. This pattern parallels findings across diverse taxa

where consumers regulate intake of multiple nutrients, not just energy (Simpson and

Raubenheimer, 2012, Ruiz et al., 2021, Jensen et al., 2012). The efficiency gains observed in

urchins fed diverse diets are consistent with nutritional geometry theory, which predicts that

complementary nutrients reduce the need for compensatory overconsumption (Simpson et al.,

2004, Raubenheimer and Simpson, 2018). In this way, diet quality mediates the balance between

energetic costs and reproductive performance.

Our FA analyses reinforce these conclusions by linking metabolic performance to

assimilated nutrient profiles. Wild kelp forest urchins had higher concentrations of kelp-derived

FA biomarkers, while barrens and starved individuals showed enrichment of bacterial and

biofilm markers. Feeding on diverse macroalgal diets shifted FA profiles toward kelp

biomarkers, confirming direct assimilation of high-value nutrients. These results extend prior

work tracing FA assimilation in echinoids (Schram et al., 2018, Raymond et al., 2014) and link

starvation to elevated microbial biomarkers. Our FA data show that experimentally starved kelp

forest urchins and wild barrens urchins elevate bacterial/biofilm biomarkers (odd-chain and

branched FAs; palmitoleic acid 16:1n-7) relative to kelp forest conspecifics, consistent with a

greater reliance on microbial substrates in food-poor habitats. This finding aligns with prior FA-

based reports suggesting the occurrence of benthic diatoms in the diets of urchins found in

barrens (Kelly et al., 2008). Parallel microbiome studies show that urchin gut communities shift

with habitat; notably, some taxa abundant in *M. franciscanus* guts, such as *Achromobacter*, are

rare or absent in their available foods (Miller et al., 2021). This genus can produce protein-rich

biomass, illustrating a plausible route by which microbial associates contribute nitrogen- and

amino acid-rich nutrition when macrophyte foods are limiting (Ehsani et al., 2019). In addition,

Campylobacteraceae dominate the guts of *M. franciscanus*, and predictive metagenomics

suggest these bacteria metabolize lipids, amino acids, and carbohydrates, potentially enhancing

host persistence in food-poor habitats (Hakim et al., 2019, Hakim et al., 2016). Classic

experiments corroborate nutritional provisioning by gut microbes: suppressing gut bacteria

reduced incorporation of essential amino acids into gonads (Fong and Mann, 1980), and

nitrogen-fixing *Vibrio spp.* have been isolated from urchin guts (Guerinot and Patriquin, 1981).

Finally, the presence of odd- and branched-chain FAs in urchin tissues, particularly in detritus-

feeding contexts, has long been attributed to microbial sources, reinforcing our interpretation of

FA shifts under starvation and in barrens as signals of microbial and/or detrital reliance

(Zhukova, 2023, Hayashi and Takagi, 1977).

If consumers are able to depress energetic demands in the absence of food, but rebound

when food becomes abundant, such phenomena are likely to inhibit such cyclic dynamics and

facilitate persistent barren states. Our results demonstrate metabolic resilience of the dominant

nearshore grazer, *M. franciscanus* with strong consequences for grazing effects in nearshore

ecosystems. While starvation induces metabolic depression, individuals presented with food can

quickly recover to feed at consistent rates regardless of whether it has a history of metabolic

depression (**figures 1 and 2**). In this study, emaciated individuals were capable of nearly

complete recovery of gonadal mass in just over one month when supplied ample food. Moreover,

recovery was more efficient with a diverse diet (**figure 3**), supporting the hypothesis that

consumers balance the intake of complementary nutrients (Raubenheimer et al., 2023) rather

than selecting singular substitutable food resources based on energy density (Houston and

McNamara, 2014).

The persistence of barren habitats has often been attributed to grazer abundance (Ling et

al., 2015, Filbee-Dexter and Scheibling, 2014), yet our results suggest that grazer nutritional

ecology also contributes. Metabolic depression prolongs survival during scarcity, while

compensatory feeding on low-quality foods can maintain energy balance once early colonizing

algae or microbial biofilms appear. This dynamic creates hysteresis wherein nutritionally poor

but abundant foods can sustain populations at high density, suppressing recovery of more

diverse, nutrient-rich macroalgal assemblages. Comparable dynamics have been documented in

other systems where food quality determines consumer efficiency and population impacts

(Marzetz et al., 2017, Foster et al., 2015). Our findings therefore broaden the mechanistic basis

for barren stability from consumer density alone to include nutrient-specific feedbacks in grazer-

producer interactions.

Although our study was framed in nutritional ecology, the findings have implications for

management. The remarkable resilience of *M. franciscanus* suggests that barren formation

reduces the resilience of kelp forests by maintaining high consumer capacity to exploit resurgent

algae. This aligns with observations that urchin overgrazing exacerbates climate-driven kelp

declines (Smale, 2020, Ling et al., 2015). Restoration strategies such as predator recovery

(Watson and Estes, 2011, Burt et al., 2018), targeted urchin removals (Miller et al., 2022), and/or

commercial urchin ranching (Powell et al., 2020, Angwin et al., 2022) may therefore be

necessary to reset the system by reducing consumer density and competition. Importantly, our

results suggest that interventions which shift the nutritional landscape (e.g., enhancing the

diversity of macroalgal recruitment) could also reduce compensatory feeding pressure and aid

kelp recovery.

Ethics

This work did not require ethical approval from a human subject or animal welfare committee.

Data, code, and materials

Datasets, code, and electronic supplementary material will be made available to reviewers upon

request, and pending acceptance will be archived on an open-access repository preferred by the

journal.

Declaration of AI use

We have not used AI-assisted technologies in creating this article.

Conflict of interest declaration

We declare we have no competing interests.

Funding

This study was funded by an FSU APACT grant to DKO, Parks Canada Conservation and

Restoration (CoRe) project (no. 1808) funding to Gwaii Haanas National Park Reserve, National

Marine Conservation Area Reserve, and Haida Heritage Site with LL as technical lead and DKO

via contribution agreement with FSU, NSF grant 2023649 to DKO, the William R. and Lenore

Mote Eminent Scholar in Marine Biology Endowment at FSU to NBS, a Professional

Association of Diving Instructors (PADI) Foundation Research Grant to NBS, an Academy of

Underwater Arts and Sciences Zale Parry Scholarship to NBS, and the Tula Foundation.

Acknowledgements

We thank Markus Thompson of Thalassia for assistance with specimen collection and

underwater photography at Quadra Island, Brenna Collicutt of the Hakai Institute for technical

support; Eric Peterson and the Hakai Institute for providing laboratory facilities at Quadra Island

and for technical support; Reyn Yoshioka for assistance with fatty acid analysis at the Oregon

Institute of Marine Biology.

Figure 1. Shifts in resting metabolic rate associated with habitat of origin and diet. Smaller faded

symbols represent measured data points, larger solid symbols represent modeled mean values,

and thin and thick vertical bars represent 95% and 80% credible intervals, respectively. Lettered

labels next to modeled means indicate statistically homogenous groups.

Figure 2. Effects of diet on per capita consumption as a function of time, after controlling for

habitat of origin, in terms of (a) algal biomass, (b) calories, and (c) nutritionally valuable

polyunsaturated fatty acids (PUFA – fatty acids with two or more double bonds). Smaller faded

symbols represent measured data points, larger solid symbols represent modeled mean values,

and thin and thick vertical bars represent 95% and 80% credible intervals, respectively. Lettered

labels next to modeled means indicate statistically homogenous groups.

Figure 3. Effects of habitat and diet on (a) gonad mass and (b) food conversion efficiency.

Smaller faded symbols represent measured data points, larger solid symbols represent modeled

mean values, and thin and thick vertical bars represent 95% and 80% credible intervals,

respectively. Lettered labels next to modeled means indicate statistically homogenous groups.

Figure 4. Shifts in gonadal fatty acid (FA) composition as a function of habitat of origin and

diet. (a & b) Icons represent individual profiles and ellipses indicate 95% confidence intervals.

(c) Arrows indicate correlation vectors for constituent fatty acids, with FAs representing kelp

biomarkers, biofilm biomarkers, and diatoms colored in gold, red, and aquamarine, respectively.

536   **Figure 5** Effects of habitat of origin and diet on gonadal concentrations of (*a*) kelp biomarkers,

537   and (*b*) biofilm biomarkers. Smaller faded symbols represent measured data points, larger solid

538   symbols represent modeled mean values, and thin and thick vertical bars represent 95% and 80%

539   credible intervals, respectively. Lettered labels next to modeled means indicate statistically

540   homogenous groups.

References

Aitchison, J. (1982) The Statistical-Analysis of Compositional Data. *Journal of the Royal Statistical Society*
Series B-Statistical Methodology, **44,** 139-177.
Angwin, R. E., Hentschel, B. T. & Anderson, T. W. (2022) Gonad enhancement of the purple sea urchin,
Strongylocentrotus purpuratus, collected from barren grounds and fed prepared diets and kelp.
Aquaculture International, **30,** 1353-1367.
Berner, D., Blanckenhorn, W. U. & Korner, C. (2005) Grasshoppers cope with low host plant quality by
compensatory feeding and food selection: N limitation challenged. *Oikos,* **111,** 525-533.
Brett, J. & Groves, T. (1979) Physiological energetics. *Fish physiology,* **8,** 280-352.
Britton-Simmons, K. H., Foley, G. & Okamoto, D. (2009) Spatial subsidy in the subtidal zone: utilization of
drift algae by a deep subtidal sea urchin. *Aquatic Biology,* **5,** 233-243.
Brown, J. S. & Kotler, B. P. (2004) Hazardous duty pay and the foraging cost of predation. *Ecology*
Letters, **7,** 999-1014.
Bürkner, P.-C. (2017) Advanced Bayesian multilevel modeling with the R package brms. *arXiv preprint*
arXiv:1705.11123.
Burt, J. M., Tinker, M. T., Okamoto, D. K., Demes, K. W., Holmes, K. & Salomon, A. K. (2018) Sudden
collapse of a mesopredator reveals its complementary role in mediating rocky reef regime shifts.
Proc Biol Sci, **285,** 20180553.
Chippindale, A. K., Leroi, A. M., Kim, S. B. & Rose, M. R. (1993) Phenotypic plasticity and selection in
Drosophila life-history evolution. I. Nutrition and the cost of reproduction. *Journal of*
evolutionary biology, **6,** 171-193.
Clarke, K. R. (2006) Non-parametric multivariate analyses of changes in community structure. *Australian*
Journal of Ecology, **18,** 117-143.
Cruz-Rivera, E. & Hay, M. E. (2000) Can Quantity Replace Quality? Food Choice, Compensatory Feeding,
and Fitness of Marine Mesograzers. *Ecology,* **81,** 201-219.
Dolinar, D. & Edwards, M. (2021) The metabolic depression and revival of purple urchins
(Strongylocentrotus purpuratus) in response to macroalgal availability. *Journal of Experimental*
Marine Biology and Ecology, **545,** 151646.
Edwards, M., Konar, B., Kim, J. H., Gabara, S., Sullaway, G., McHugh, T., Spector, M. & Small, S. (2020)
Marine deforestation leads to widespread loss of ecosystem function. *PLoS One,* **15,** e0226173.
Edwards, P. B. & Ebert, T. A. (1991) Plastic responses to limited food availability and spine damage in the
sea urchin Strongylocentrotus purpuratus (Stimpson). *Journal of Experimental Marine Biology*
and Ecology, **145,** 205-220.
Ehsani, E., Dumolin, C., Arends, J. B. A., Kerckhof, F. M., Hu, X., Vandamme, P. & Boon, N. (2019)
Enriched hydrogen-oxidizing microbiomes show a high diversity of co-existing hydrogen-
oxidizing bacteria. *Appl Microbiol Biotechnol,* **103,** 8241-8253.
Ellers, J. (1995) Fat and Eggs: an Alternative Method To Measure the Trade-Off Between Survival and
Reproduction in Insect Parasitoids. *Netherlands Journal of Zoology,* **46,** 227-235.
Filbee-Dexter, K. & Scheibling, R. E. (2014) Sea urchin barrens as alternative stable states of collapsed
kelp ecosystems. *Marine Ecology Progress Series,* **495,** 1-25.
Fink, P. & Von Elert, E. (2006) Physiological responses to stoichiometric constraints: nutrient limitation
and compensatory feeding in a freshwater snail. *Oikos,* **115,** 484-494.
Fong, W. & Mann, K. H. (1980) Role of Gut Flora in the Transfer of Amino Acids Through a Marine Food
Chain. *Canadian Journal of Fisheries and Aquatic Sciences,* **37,** 88-96.
Foster, M. C., Byrnes, J. E. & Reed, D. C. (2015) Effects of five southern California macroalgal diets on
consumption, growth, and gonad weight, in the purple sea urchin Strongylocentrotus
purpuratus. *PeerJ,* **3,** e719.

Fuiman, L. A. & Perez, K. O. (2015) Metabolic programming mediated by an essential fatty acid alters
body composition and survival skills of a marine fish. *Proc Biol Sci,* **282,** 20151414.
Gangal, M., Gafoor, A. B., D'Souza, E., Kelkar, N., Karkarey, R., Marbà, N., Arthur, R. & Alcoverro, T.
(2021) Sequential overgrazing by green turtles causes archipelago-wide functional extinctions of
seagrass meadows. *Biological Conservation,* **260,** 109195.
Gelman, A., Carlin, J. B., Stern, H. S., Dunson, D. B., Vehtari, A. & Rubin, D. B. (2013) *Bayesian data*
analysis. CRC press.
Gelman, A., Goodrich, B., Gabry, J. & Vehtari, A. (2019) R-squared for Bayesian Regression Models.
American Statistician, **73,** 307-309.
Glazier, D. S. (2015) Is metabolic rate a universal 'pacemaker' for biological processes? *Biol Rev Camb*
Philos Soc, **90,** 377-407.
Gronau, Q. F., Singmann, H. & Wagenmakers, E. J. (2020) bridgesampling: An R Package for Estimating
Normalizing Constants. *Journal of Statistical Software,* **92,** 1-29.
Guerinot, M. L. & Patriquin, D. G. (1981) N2-fixing vibrios isolated from the gastrointestinal tract of sea
urchins. *Canadian Journal of Microbiology,* **27,** 311-317.
Hakim, J. A., Koo, H., Kumar, R., Lefkowitz, E. J., Morrow, C. D., Powell, M. L., Watts, S. A. & Bej, A. K.
(2016) The gut microbiome of the sea urchin, Lytechinus variegatus, from its natural habitat
demonstrates selective attributes of microbial taxa and predictive metabolic profiles. *FEMS*
Microbiol Ecol, **92,** fiw146.
Hakim, J. A., Schram, J. B., Galloway, A. W. E., Morrow, C. D., Crowley, M. R., Watts, S. A. & Bej, A. K.
(2019) The Purple Sea Urchin Strongylocentrotus purpuratus Demonstrates a
Compartmentalization of Gut Bacterial Microbiota, Predictive Functional Attributes, and
Taxonomic Co-Occurrence. *Microorganisms,* **7,** 35.
Hamidi, B., Wallace, K., Vasu, C. & Alekseyenko, A. V. (2019) W(*)(d) -test: robust distance-based
multivariate analysis of variance. *Microbiome,* **7,** 51.
Harianto, J., Carey, N. & Byrne, M. (2019) respR-An R package for the manipulation and analysis of
respirometry data. *Methods in Ecology and Evolution,* **10,** 912-920.
Harrison, S. J., Raubenheimer, D., Simpson, S. J., Godin, J. G. & Bertram, S. M. (2014) Towards a synthesis
of frameworks in nutritional ecology: interacting effects of protein, carbohydrate and
phosphorus on field cricket fitness. *Proc Biol Sci,* **281,** 20140539.
Hayashi, K. & Takagi, T. (1977) Occurrence of branched-and odd-chain fatty acids in a mud-feeding sea
urchin, Strongylocentrotus franciscanus. 北海道大學水産學部研究彙報, **28,** 40-46.
Houston, A. I. & McNamara, J. M. (2014) Foraging currencies, metabolism and behavioural routines. *J*
Anim Ecol, **83,** 30-40.
Jensen, K., Mayntz, D., Toft, S., Clissold, F. J., Hunt, J., Raubenheimer, D. & Simpson, S. J. (2012) Optimal
foraging for specific nutrients in predatory beetles. *Proc Biol Sci,* **279,** 2212-8.
Jochum, M., Barnes, A. D., Ott, D., Lang, B., Klarner, B., Farajallah, A., Scheu, S. & Brose, U. (2017)
Decreasing Stoichiometric Resource Quality Drives Compensatory Feeding across Trophic Levels
in Tropical Litter Invertebrate Communities. *American Naturalist,* **190,** 131-143.
Kainz, M. J., Hager, H. H., Rasconi, S., Kahilainen, K. K., Amundsen, P. A. & Hayden, B. (2017)
Polyunsaturated fatty acids in fishes increase with total lipids irrespective of feeding sources and
trophic position. *Ecosphere,* **8,** e01753.
Kass, R. E. & Raftery, A. E. (1995) Bayes Factors. *Journal of the American Statistical Association,* **90,** 773-
795.
Kayal, M., Vercelloni, J., Lison de Loma, T., Bosserelle, P., Chancerelle, Y., Geoffroy, S., Stievenart, C.,
Michonneau, F., Penin, L., Planes, S. & Adjeroud, M. (2012) Predator crown-of-thorns starfish
(Acanthaster planci) outbreak, mass mortality of corals, and cascading effects on reef fish and
benthic communities. *PLoS One,* **7,** e47363.

Kelly, J. R., Scheibling, R. E., Iverson, S. J. & Gagnon, P. (2008) Fatty acid profiles in the gonads of the sea
urchin Strongylocentrotus droebachiensis on natural algal diets. *Marine Ecology Progress Series,*
373, 1-9.
Kirkwood, T. B. & Rose, M. R. (1991) Evolution of senescence: late survival sacrificed for reproduction.
Philos Trans R Soc Lond B Biol Sci, **332,** 15-24.
Leal, I., Tremblay, R. & Flores, A. A. V. (2022) Allochthonous subsidies drive early recruitment of a
subtropical foundation species. *Oikos,* **2022,** e08991.
Lenth, R., Singmann, H., Love, J., Buerkner, P. & Herve, M. (2024) emmeans: Estimated Marginal Means,
aka Least-Squares Means.
Lesage, L., Crete, M., Huot, J. & Ouellet, J. P. (2001) Evidence for a trade-off between growth and body
reserves in northern white-tailed deer. *Oecologia,* **126,** 30-41.
Lighton, J. R. (2018) *Measuring metabolic rates: a manual for scientists.* Oxford University Press.
Ling, S. D., Scheibling, R. E., Rassweiler, A., Johnson, C. R., Shears, N., Connell, S. D., Salomon, A. K.,
Norderhaug, K. M., Pérez-Matus, A., Hernández, J. C., Clemente, S., Blamey, L. K., Hereu, B.,
Ballesteros, E., Sala, E., Garrabou, J., Cebrian, E., Zabala, M., Fujita, D. & Johnson, L. E. (2015)
Global regime shift dynamics of catastrophic sea urchin overgrazing. *Philosophical Transactions*
of the Royal Society B-Biological Sciences, **370,** 20130269.
Marzetz, V., Koussoroplis, A. M., Martin-Creuzburg, D., Striebel, M. & Wacker, A. (2017) Linking primary
producer diversity and food quality effects on herbivores: A biochemical perspective. *Sci Rep,* **7,**
11035.
McCue, M. D., Terblanche, J. S. & Benoit, J. B. (2017) Learning to starve: impacts of food limitation
beyond the stress period. *J Exp Biol,* **220,** 4330-4338.
Mduma, S. A. R., Sinclair, A. R. E. & Hilborn, R. (1999) Food regulates the Serengeti wildebeest: a 40-year
record. *Journal of animal ecology,* **68,** 1101-1122.
Miller, K. I., Blain, C. O. & Shears, N. T. (2022) Sea Urchin Removal as a Tool for Macroalgal Restoration:
A Review on Removing "the Spiny Enemies". *Frontiers in Marine Science,* **9**.
Miller, P. M., Lamy, T., Page, H. M. & Miller, R. J. (2021) Sea urchin microbiomes vary with habitat and
resource availability. *Limnology and Oceanography Letters,* **6,** 119-126.
Mysterud, A. (2006) The concept of overgrazing and its role in management of large herbivores. *Wildlife*
Biology, **12,** 129-141.
Okamoto, D. K. (2014) *The role of fluctuating food supply on recruitment, survival and population*
dynamics in the sea. UC Santa Barbara.
Okamoto, D. K., Schmitt, R. J., Holbrook, S. J. & Reed, D. C. (2012) Fluctuations in food supply drive
recruitment variation in a marine fish. *Proc Biol Sci,* **279,** 4542-50.
Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'hara, R., Simpson, G. L., Solymos, P.,
Stevens, M. H. H. & Wagner, H. (2023) Package 'vegan'. *Community ecology package, version*
2.6-5.
Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M. H. H., Oksanen, M. J. & Suggests, M. (2007)
The vegan package. *Community ecology package,* **10,** 631-637.
Olsen, E. M., Ottersen, G., Llope, M., Chan, K. S., Beaugrand, G. & Stenseth, N. C. (2011) Spawning stock
and recruitment in North Sea cod shaped by food and climate. *Proc Biol Sci,* **278,** 504-10.
Parrish, C. C. (2009) Essential fatty acids in aquatic food webs. *Lipids in Aquatic Ecosystems* (eds M.
Kainz, M. T. Brett & M. T. Arts), pp. 309-326. Springer New York, New York, NY.
Powell, M. L., Marsh, A. G. & Watts, S. A. (2020) Biochemical and energy requirements of gonad
development in regular sea urchins. *Developments in Aquaculture and Fisheries Science* pp. 51-
64. Elsevier.
Randall, J. E. (1961) Overgrazing of Algae by Herbivorous Marine Fishes. *Ecology,* **42,** 812-812.

Raubenheimer, D., Hou, R., Dong, Y., Ren, C. & Cui, Z. (2023) Towards an integrated understanding of
dietary phenotypes. *Philos Trans R Soc Lond B Biol Sci,* **378,** 20220545.
Raubenheimer, D. & Simpson, S. J. (1993) The geometry of compensatory feeding in the locust. *Animal*
Behaviour, **45,** 953-964.
Raubenheimer, D. & Simpson, S. J. (2018) Nutritional ecology and foraging theory. *Curr Opin Insect Sci,*
27, 38-45.
Raubenheimer, D., Simpson, S. J. & Mayntz, D. (2009) Nutrition, ecology and nutritional ecology: toward
an integrated framework. *Functional Ecology,* **23,** 4-16.
Raymond, W. W., Lowe, A. T. & Galloway, A. W. E. (2014) Degradation state of algal diets affects fatty
acid composition but not size of red urchin gonads. *Marine Ecology Progress Series,* **509,** 213-
225.
Rogers-Bennett, L. & Okamoto, D. (2020) Mesocentrotus franciscanus and Strongylocentrotus
purpuratus. *Sea Urchins: Biology and Ecology* (ed J. M. Lawrence), pp. 593-608. Elsevier.
Ruiz, T., Koussoroplis, A. M., Danger, M., Aguer, J. P., Morel-Desrosiers, N. & Bec, A. (2021) Quantifying
the energetic cost of food quality constraints on resting metabolism to integrate nutritional and
metabolic ecology. *Ecology Letters,* **24,** 2339-2349.
Schiel, D. R. & Foster, M. S. (2015) *The biology and ecology of giant kelp forests.* Univ of California Press.
Schram, J. B., Kobelt, J. N., Dethier, M. N. & Galloway, A. W. E. (2018) Trophic Transfer of Macroalgal
Fatty Acids in Two Urchin Species: Digestion, Egestion, and Tissue Building. *Frontiers in Ecology*
and Evolution, **6,** 83.
Silliman, B. R. & Zieman, J. C. (2001) Top-down control of

production by periwinkle grazing in a Virginia salt marsh. *Ecology,* **82,** 2830-2845.
Simpson, S. J. & Raubenheimer, D. (2012) The nature of nutrition. *The Nature of Nutrition.* Princeton
university press.
Simpson, S. J., Sibly, R. M., Lee, K. P., Behmer, S. T. & Raubenheimer, D. (2004) Optimal foraging when
regulating intake of multiple nutrients. *Animal Behaviour,* **68,** 1299-1311.
Smale, D. A. (2020) Impacts of ocean warming on kelp forest ecosystems. *New Phytologist,* **225,** 1447-
1454.
Solon-Biet, S. M., McMahon, A. C., Ballard, J. W., Ruohonen, K., Wu, L. E., Cogger, V. C., Warren, A.,
Huang, X., Pichaud, N., Melvin, R. G., Gokarn, R., Khalil, M., Turner, N., Cooney, G. J., Sinclair, D.
A., Raubenheimer, D., Le Couteur, D. G. & Simpson, S. J. (2014) The ratio of macronutrients, not
caloric intake, dictates cardiometabolic health, aging, and longevity in ad libitum-fed mice. *Cell*
Metab, **19,** 418-30.
Spindel, N. B., Lee, L. C. & Okamoto, D. K. (2021) Metabolic depression in sea urchin barrens associated
with food deprivation. *Ecology,* **102,** e03463.
Sterner, R. W. & Elser, J. J. (2002) *Ecological stoichiometry: the biology of elements from molecules to the*
biosphere. Princeton university press.
Storey, K. B. (2015) Regulation of hypometabolism: insights into epigenetic controls. *J Exp Biol,* **218,** 150-
9.
Tait, A. H., Raubenheimer, D., Stockin, K. A., Merriman, M. & Machovsky-Capuska, G. E. (2014)
Nutritional geometry and macronutrient variation in the diets of gannets: the challenges in
marine field studies. *Marine Biology,* **161,** 2791-2801.
Team, R. C. (2024) R: A language and environment for statistical computing. *R Foundation for Statistical*
Computing.
Team, S. D. (2022) Stan Modeling Language Users Guide and Reference Manual, 2.29.

Tinker, M. T., Guimaraes, P. R., Jr., Novak, M., Marquitti, F. M., Bodkin, J. L., Staedler, M., Bentall, G. &
Estes, J. A. (2012) Structure and mechanism of diet specialisation: testing models of individual
variation in resource use with sea otters. *Ecology Letters,* **15,** 475-83.
Twining, C. W., Bernhardt, J. R., Derry, A. M., Hudson, C. M., Ishikawa, A., Kabeya, N., Kainz, M. J., Kitano,
J., Kowarik, C., Ladd, S. N., Leal, M. C., Scharnweber, K., Shipley, J. R. & Matthews, B. (2021) The
evolutionary ecology of fatty-acid variation: Implications for consumer adaptation and
diversification. *Ecology Letters,* **24,** 1709-1731.
Vehtari, A., Gelman, A. & Gabry, J. (2016) Practical Bayesian model evaluation using leave-one-out cross-
validation and WAIC. *Statistics and Computing,* **27,** 1413-1432.
Vilela, A., Cariaga, R., González-Paleo, L. & Ravetta, D. (2008) Trade-offs between reproductive allocation
and storage in species of

L. (Onagraceae) native to Argentina. *Acta Oecologica-International Journal of Ecology,* **33,** 85-92.
Watson, J. & Estes, J. A. (2011) Stability, resilience, and phase shifts in rocky subtidal communities along
the west coast of Vancouver Island, Canada. *Ecological Monographs,* **81,** 215-239.
Yao, Y. L., Vehtari, A., Simpson, D., Gelman, A., Clarke, B., Li, M., Grünwald, P., de Heide, R., Dawid, A. P.,
Yoo, W. W., Winkler, R. L., Jose, V. R. R., Lichtendahl, K. C., Grushka-Cockayne, Y., McAlinn, K.,
Aastveit, K. A., West, M., Shin, M., Zhou, T. J., Hoogerheide, L., van Dijk, H. K., Bakka, H. C.,
Castro-Camilo, D., Franco-Villoria, M., Freni-Sterrantino, A., Huser, R., Opitz, T., Rue, H., Ferreira,
M. A. R., Pericchi, L., Franck, C. T., Belitser, E., Nurushev, N., Iacopini, M. & Tonellato, S. (2018)
Using Stacking to Average Bayesian Predictive Distributions (with Discussion). *Bayesian Analysis,*
13, 917-1003.
Zhukova, N. V. (2023) Fatty Acids of Echinoderms: Diversity, Current Applications and Future
Opportunities. *Marine Drugs,* **21,** 21.